


713

625

BOOK NO.

083

K



FRONTISPIECE.

XIII. B. 37.

2 THE ENTERTAINING NATURALIST:

BEING POPULAR DESCRIPTIONS, TALES, AND ANECDOTES

OF MORE THAN

83/

FIVE HUNDRED ANIMALS,

COMPREHENDING ALL THE

Quadrupeds, Birds, Fishes, Reptiles, Insects, &c.

OF WHICH A KNOWLEDGE IS INDISPENSABLE IN

POLITE EDUCATION.

WITH INDEXES OF SCIENTIFIC AND POPULAR NAMES, AN EXPLANATION OF TERMS, AND AN APPENDIX OF FABULOUS ANIMALS.

ILLUSTRATED BY NEARLY

FIVE HUNDRED ENGRAVINGS ON WOOD BY BEWICK, HARVEY, LANDELLS, THOMPSON, WHIMPER, SLY, AND OTHERS.

A New Edition, revised, enlarged, and completed to the present state of Zoological Science,

BY MRS. (LOUDON.)



LONDON:

HENRY G BOHN, YORK STREET, COVENT GARDEN.

1850.

590
Lou

TRANSFER
Imperial Library

12

1903

LONDON:
JOSEPH BICKERBY, PRINTER,
SHERBOURN LANE.

Acc. no. - 156
18/11/06

Acc. no. 156
dt. 18-11-06

P R E F A C E.

THE care of preparing a new edition of this long, established work having been confided to me, I have endeavoured to render it as instructive as I could without destroying its popular and entertaining character. For this purpose I have added scientific names to the popular ones at the head of each description, and I have placed all the animals of each genus and all the genera in each order together, that the child who reads the book may have the animals most nearly allied to each other associated together in his mind. For the same reason, I have used, in most cases, the Linnæan names, as Linnæus placed a great many animals in one genus, which are now divided into several. At the same time that the child may learn both the Linnæan and the modern name of each animal, I have given both in the Table of Contents. I have not arranged the orders quite scientifically; as I could not persuade myself either to displace the Lion from the situation he has held so long at the com-

mencement of the book, or to remove the Whales and other Cetacea from the place they have near the fishes. I have therefore placed the Monkeys and their allies at the end of the terrestrial Mammalia, instead of at the beginning; and commenced the third book with the Cetacea.

In the present edition, the whole of the descriptions of the animals have been corrected, so as to bring the whole down to the present state of Zoological knowledge; and in the Introduction a Sketch of the Modern System of Arrangement of the Animal Kingdom has been added to that of Linnæus.

J. W. L.

Bayswater, December 20th, 1849.

XIII U 37

INTRODUCTION.

ZOOLOGY is that branch of Natural History which treats of animals, and embraces not only their structure and functions, their habits, instincts, and utility, but their names and systematic arrangement.

Various systems have been proposed by different naturalists for the scientific arrangement of the animal kingdom, but that of Cuvier, with some modifications by Lamarck and Latreille, is now thought the best, and a sketch of it will be found under the head of the Modern System in this Introduction. As, however, the System of Linnæus was formerly in general use, and is still often referred to, it has been thought advisable to give a sketch of it first; that the reader may be aware of the difference between the old system and the new one.

LINNÆAN SYSTEM.

ACCORDING to the system of Linnæus, the objects comprehended within the animal kingdom were divided into six classes: Mammalia or Mammiferous Animals, Birds, Amphibia, or Amphibious Animals, Fishes, Insects, and Worms, which were thus distinguished;

CLASSES.

Body	{	With vertebræ	Hot Blood.....	{ Viviparous	I. MAMMALIA.
			Cold red Blood.....	{ Oviparous	II. BIRDS.
{	Without vertebræ	Cold white Blood	{ With lungs	III. AMPHIBIA.	
			{ With gills	IV. FISHES.	
			{ Having antennæ	V. INSECTS.	
			{ Having tentacula	VI. WORMS.	

ORDERS OF MAMMALIA.

The first class, or Mammalia, consists of such animals as produce living offspring, and nourish their young ones with milk supplied from their own bodies; and it comprises both the quadrupeds and the cetacea.

*This class was divided by Linnæus into seven Orders: viz. *primates*, *bruta*, *feræ*, *glivæ*, *pecora*, *belluæ*, and *cetacea* (this order was called Cete by Linnæus) or whales. The characteristics of these were founded, for the most part, on the number and arrangement of the teeth; and

on the form and construction of the feet, or of those parts in the seals, manati, and cetacea, which supply the place of feet.

- I. PRIMATES.—Having the upper front teeth, generally four in number, wedge-shaped, and parallel; and two teats situated on the breast, as the apes and monkeys.
- II. BRUTA.—Having no front teeth in either jaw; and the feet armed with strong hoof-like nails, as the elephant.
- III. FERÆ.—Having in general six front teeth in each jaw; a single canine tooth on each side in both jaws; and the grinders with conic projections, as the dogs and cats.
- IV. GLIRES.—Having in each jaw two long projecting front teeth, which stand close together; and no canine teeth in either jaw, as the rats and mice.
- V. PECORA.—Having no front teeth in the upper jaw; six or eight in the lower jaw, situated at a considerable distance from the grinders; and the feet with hoofs, as cattle and sheep.
- VI. BELLUÆ.—Having blunt wedge-shaped front teeth in both jaws; and the feet with hoofs, as horses.
- VII. CETACEA.—Having spiracles, or breathing-holes on the head; fins instead of fore-feet; and a tail flattened horizontally, instead of hind feet. This order consists of the narvals, whales, cachalots, and dolphins.

ORDERS OF BIRDS.

The second class, or Birds, comprises all such animals as have their bodies clad with feathers. Their jaws are elongated, and covered externally with a horny substance, called a bill or beak, which is divided into two parts called mandibles. Their eyes are furnished with a thin, whitish, and somewhat transparent membrane, that can at pleasure be drawn over the whole external surface like a curtain. Birds respire by air vessels, which are extended through their body, and which, in the abdominal cavity, adhere to the under surface of the bones. Their organs of motion are two wings and two legs; and they are destitute of external ears, lips, and many other parts which are important to quadrupeds. That part of zoology which treats of Birds is called Ornithology.

Linnæus divided this class into six orders.

I. Land Birds.

- I. RAPACIOUS BIRDS (*Accipitres*).—Having the upper mandible hooked, and an angular projection on each side near the point, as the eagles, hawks, and owls.
- II. PIES (*Pice*).—Having their bills sharp at the edge, somewhat compressed at the sides, and convex on the top, as the crow.
- III. PASSERINE BIRDS (*Passeres*).—Having the bill conical and pointed, and the nostrils oval, open, and naked, as the sparrow and linnet.

IV. GALLINACEOUS BIRDS (*Gallinæ*).—Having the upper mandible arched, and covering the lower one at the edge, and the nostrils arched over with a cartilaginous membrane, as the common poultry.

2. Water Birds.

V. WADERS (*Grallæ*).—Having a roundish bill, a fleshy tongue, and the legs naked above the knees, as the herons, plovers, and snipes.

VI. SWIMMERS (*Anseres*).—Having their bills broad at the top, and covered with a soft skin, and the feet webbed; as ducks and geese.

ORDERS OF AMPHIBIA.

Under the third class, or Amphibia, Linnæus arranged such animals as have a cold, and, generally, naked body, a lurid colour, and nauseous smell. They respire chiefly by lungs, but they have the power of suspending respiration for a long time. They are extremely tenacious of life, and can repair certain part of their bodies which have been lost. They are also able to endure hunger, sometimes even for months, without injury.

The bodies of some of them, as the turtles and tortoises, are protected by a hard and horny shield or covering; those of others are clad with scales, as the serpents, and some of the lizards; whilst others, as the frogs, toads, and most of the water-lizards, are entirely naked, or have their skin covered with warts. Many of the species shed their skins at certain times of the year. Several of them are furnished with a poison, which they eject into wounds that are made by their teeth. They chiefly live in retired, watery, and marshy places; and, for the most part, feed on other animals, though some of them eat water-plants, and many feed on garbage and filth. None of these species chew their food; they swallow it whole, and digest it very slowly.

The offspring of all these animals are produced from eggs, which, after they have been deposited by the parent animals in a proper place, are hatched by the heat of the sun. The eggs of some of the species are covered with a shell; those of others have a soft and tough skin or covering, not much unlike wet parchment; and the eggs of several are perfectly gelatinous. In those few that produce their offspring alive, as the vipers, and some other serpents, the eggs are regularly formed, but are hatched within the bodies of the females.

This class Linnæus divided into two Orders.

I. REPTILES.—Having four legs, and walking with a crawling pace, as the tortoises, toads, and lizards.

II. SERPENTS.—Having no legs, but crawling on the body.

ORDERS OF FISHES.

Fishes constituted Linnæus's fourth class of animals. They are all inhabitants of the water, in which they move by certain organs called

fins. Those situated on the back, are called dorsal fins; those on the sides, behind the gills, pectoral fins; those below the body, near the head, are ventral; those behind the vent are anal; and that which forms the tail is called the caudal fin. Fishes breathe by gills, which, in most species, are situated at the sides of the head. In some of the flat-fish, however, as the skate and the thornback, they are on the under-part of the body. Fishes rise and sink in the water, generally by a kind of bladder in the interior of the body, called an air-bladder. Some of them, as the skate and other flat-fish, do not possess this organ, and consequently are seldom found but at the bottom of the sea. The bodies of these animals are usually covered with scales, which keep them from injury by the pressure of the water.

The fishes were divided by Linnæus into six Orders.

- I. **APODAL.**—Having bony gills; and no ventral fins, as the eel.
- II. **JUGULAR.**—Having bony gills; and the ventral fins situated in front of the pectoral fins, as the cod, haddock, and whiting.
- III. **THORACIC.**—Having bony gills; and the ventral fins situated directly under the pectoral fins, as the perch and mackerel.
- IV. **ABDOMINAL.**—Having bony gills; and the ventral fins on the lower part of the body below the pectoral fins, as the salmon, herrings, and carp.
- V. **BRANCHIOSTEGOUS.**—Having gills destitute of bony rays.
- VI. **CHONDROPTERYGEOUS.**—Having cartilaginous fins, as the sturgeons, sharks, and skate.

ORDERS OF INSECTS.

The fifth class of Linnæus comprised the Insects; and the branch of zoology which treats of them is called Entomology. Nearly all insects go through certain great changes at different periods of their existence. From the egg is hatched the larva, which is a grub or caterpillar, and destitute of wings; this afterwards changes to a pupa, or carysalis, wholly covered with a hard shell, or strong skin, from which the perfect or winged insect bursts forth. Spiders and their allies, which were included by Linnæus in the insects, issue from the egg in nearly a perfect state.

Linnæus divided his insects into seven Orders.

- I. **COLEOPTEROUS.**—Having elytra, or crustaceous cases covering the wings; and which, when closed, form a longitudinal division along the middle of the back, as the cockchafer.
- II. **HEMIPTEROUS.**—Having four wings, the upper ones partly crustaceous, and partly membranous; not divided straight down the middle of the back, but crossed, or incumbent on each other, as the cock-roach.
- III. **LEPIDOPTEROUS.**—Having four wings covered with fine scales almost like powder, as the butterflies and moths.
- IV. **NEUROPTEROUS.**—Having four membranous and semi-transparent wings, veined like net-work; and the tail without a sting, as the dragon-fly and ephemera.

- V. **HYMENOPTEROUS.**—Having four membranous and semi-transparent wings, veined like net-work; and the tail armed with a sting, as the wasp and bee.
- VI. **DIPTEROUS.**—Having only two wings, as the common house-flies.
- VII. **APTEROUS.**—Having no wings, as the spiders.

ORDERS OF VERMES, OR WORMS.

The sixth and last Linnæan class consisted of Worms, or Vermes. These are slow of motion, and have soft and fleshy bodies. Some of them have hard internal parts, and others have crustaceous coverings. In some of the species, eyes and ears are very perceptible, whilst others appear to enjoy only the senses of taste and touch. Many have no distinct head, and most of them are destitute of feet. They are, in general, so tenacious of life, that parts which have been destroyed will be reproduced. These animals are principally distinguished from those of the other classes by having tentacula, or feelers, and are divided by Linnæus into five Orders.

- I. **INTESTINAL.**—Are simple and naked, without limbs; some of them live within other animals, as the ascarides and tape-worms; others in water, as the leeches; and a few in the earth, as the earth-worm.
- II. **MOLLUSCOUS.**—Are simple animals, without shells, and furnished with limbs, as the cuttle-fish, medusæ, star-fish, and sea-urchin.
- III. **TESTACEOUS.**—Are animals similar to the last, but covered with shells, as oysters, cockles, snails, and limpets.
- IV. **ZOOPHYTES.**—Are composite animals, and appear to hold a rank between animals and vegetables; though they are in fact true animals, and possess sensation and voluntary motion. In many instances a great number of them inhabit the same stone, but some are soft, naked, and separate. The coral, sponge, and polypes are instances of this order.
- V. **ANIMALCULES.**—Are destitute of tentacula or feelers, and are generally so minute as to be invisible to the naked eye. They are chiefly found in different infusions of animal and vegetable substances.

MODERN SYSTEM.

It will be found by reading the following sketch of the Modern System, that the greatest change has taken place in the latter two classes. The others remain nearly the same in effect, though their distinctions are different, and the classes are not arranged in the same order.

ACCORDING to Cuvier, all animals are arranged in four great divisions, which are subdivided into classes and orders, as follows:—

DIVISIONS.	CLASSES.	NUMBER OF ORDERS.
I. VERTEBRATA. Four Classes. Twenty-seven Orders.	1 Mammalia	Eight.
	2 Aves	Six.
	3 Reptilia	Four.
	4 Pisces	Nine.

DIVISIONS.	CLASSES.	NUMBER OF ORDERS.
II. MOLLUSCA.	1 Cephalopoda	One.
Six Classes. Fifteen Orders.	2 Pteropoda	One.
	3 Gasteropoda	Nine.
	4 Acephala	Two.
	5 Brachiopoda	One.
	6 Cirrhopodes	One.
III. ARTICULATA.	1 Annelides	Three.
Four Classes. Twenty- four Orders.	2 Crustacea	Seven.
	3 Arachnides	Two.
	4 Insecta	Twelve.
	IV. RADIATA.	1 Echinodermata
Five Classes. Eleven Orders.	2 Entozoa	Two.
	3 Acalephæ	Two.
	4 Polyphi	Three.
	5 Infusoria	Two.

THE VERTEBRATED ANIMALS

Have a backbone divided into vertebræ or joints, whence they take their name. They have also separate senses for hearing, seeing, tasting, smelling, and feeling; a distinct head, with a mouth opening by two horizontal jaws; a muscular heart, and red blood. The four classes of Vertebrata and their orders are as follow.

I. THE MAMMALIA are all furnished with mammæ, or teats, through which they give milk to their young, which they bring forth alive. They have warm blood, which circulates from the heart through the lungs, and returns to the heart before it passes through the body. Their skins are naked, or covered with wool or hair, and their mouths are generally furnished with teeth. There are eight orders which are thus distinguished:—

Unguiculata; that is, having nails or claws.	Having hands.	I. Bimana.
		II. Quadrumana.
Ungulata; that is, without nails or claws.	Quadrupeds, or animals with four feet.	III. Cheiroptera.
		IV. Carnivora.
	Having fins.	V. Marsupialia.
		VI. Glires, or Rodentia.
		VII. Edentata.
		VIII. Pachydermata.
		IX. Ruminantia
		X. Cetacea.
		XI. Monothrema.

SECTION I.—*Unguiculated Animals, or Mammalia having nails or claws.*

- I. *Bimana*, or two handed. This order contains only the human species.
- II. *Quadrumana*, or four handed. This order contains the apes, baboons, and monkeys, and the lemurs.
- III. *Cheiroptera*, the bat family
- IV. *Carnivora*, or beasts of prey. This order is divided into the following three tribes:—

1. *The Insectivora*, consisting of those animals which live upon insects, as the hedgehog, the shrew, and the mole.
2. *The Carnivora proper*, consisting chiefly of the cat family, including lions, tigers, and their allies; the bear family, including the badger, the coati-mondi, the racoon, &c.; the dog family, including the wolf and the fox; the weazel family; the civet-cats; and the hyæna.
3. *The Amphibia*, consisting of the seals, and other allied animals.
- V. *The Marsupialia*, including the opossums, and the kangaroos.
- VI. *Rodentia*, or gnawing animals. The principal of these are the squirrel family, mice and rats, hares and rabbits, the beaver, the porcupine, and the guinea-pig.
- VII. *Edentata*, or toothless animals, that is, without front-teeth. The principal of these are the sloths, the armadillos, the ant-eaters, and the water-mole, or ornithorhynchus.

SECTION II.—*Mammalia without nails or claws.*

- VIII. *Pachydermata*, or thick-skinned animals. The principal of these are the elephant, the hippopotamus, the rhinoceros; the horse family, including the ass, the mule, the zebra, and the quagga; the wild boar family, and the tapir.
- IX. *Ruminantia*, or ruminating animals, the principal of which are the camel family, the deer family, the giraffe, the antelope family, the goat family, the sheep family, and the ox family.
- X. *Cetacea*, or sea mammalia, the principal of which are the whale family, the dolphin family, the manati, the porpoise family, and the narwal, or sea-unicorn.
- XI. *Monothrema*, containing the Echidna and Ornithorhynchus of Australia.

THE AVES, OR BIRDS,

Lay eggs from which their young are hatched by what is called incubation. Their skins are covered with feathers; and their jaws are horny, without teeth. Their blood is warm, and circulates like that of the mammalia. The six orders of Aves are as follows:—

1. *Raptores*, or birds of prey. These birds are distinguished by a very strong and sharp bill more or less curved, but always hooked at the extremity of the upper mandible, which is covered at the base with a kind of skin called the cere. The nostrils are usually open. The legs are very strong, and generally only partially covered with feathers; the feet are large, and the toes, which are four in number, are armed with very strong, sharp, curved, prehensile claws. The principal raptorial birds are the vultures, including the condor; the falcon family, including the eagles, hawks, kites, and buzzards; and the owls.
2. *Incessores*, or perching birds. These birds have all feet formed for perching, the hind toe springing from the same place as the other toes, which gives them great power of grasping. Their legs are of moderate length, and their claws not sharply curved. This order includes the thrushes, nightingales, and all the finest songsters of our groves, with the robin-redbreast, the sparrow, and other birds seen about dwellings, the swallows, the larks,

the crow family, the kingfishers, the birds of paradise, and the humming birds.

3. *Scansores*, or climbers. These birds have two toes before and two behind, or rather they have the power of throwing one of the fore toes back at pleasure. This construction gives them such great power of climbing, that they can ascend the perpendicular trunk of a tree. The principal birds in this order are the parrot family and the woodpeckers.
4. *Rasores*, or gallinaceous birds. These birds have the head small in proportion to the body. The bill is generally short, with the upper mandible somewhat curved, and furnished with a cere. The nostrils have usually a protecting fleshy membrane. The tarsus, or lower part of the leg, is long and bare, and there are four toes, those in front being united by a slight membrane, while that behind is generally higher up the leg, and smaller than the others. This order comprises nearly all the birds used as food, and includes the peacock, the turkey, the common cock and hen, the partridge, the pheasant, and the pigeon family.
5. *Grallatores*, or Waders. These birds are characterized by their long and slender legs, and by the thighs being more or less bare. There are three anterior toes, more or less united at the base by a membrane, or rudimentary web. The hind toe is wanting in one division of the order. This order contains the ostrich family, the bustards and plovers; the cranes, herons, and storks; and the snipes and woodcocks.
6. *Palmipedes*, or web-footed birds. These birds have the legs and feet short, and placed behind, with their fore toes united by a thick, and strong membrane. The neck is much longer than the legs, and their bodies are covered with a dense layer of down, beneath the outer plumage, which is close, and imbued with an oily fluid that repels the water. The principal birds in this order are the coots and grebes, the auks and penguins, the petrels, the pelican and cormorant, and the swans, ducks, and geese.

THE REPTILIA,

Or Reptiles, having neither hair, wool, nor feathers, and their bodies are either naked, or covered with scales. Some lay eggs, and some bring forth their young alive. Some have gills, and others lungs, but the latter have only a portion of the blood passing through them; and thus the blood of reptiles is cold, as it is respiration which gives the blood heat. The senses of reptiles are dull, and their movements are either slow or laborious. The following are the four orders into which this class is divided:—

1. *Chelonian Reptiles*. These animals have a heart with two auricles, and a body with four legs. The body is enclosed in an upper buckler, called the corapace, and an under one, called the plastron. They have lungs which are much expanded; but they have no teeth, though they have hard horny jaws. The females lay eggs covered with a hard shell. The principal animals belonging to this division are the tortoises, which live on land, and the turtles, which are always found in or near water.

2. *The Saurian Reptiles.* These animals have also a heart with two auricles, expanded lungs, and generally four legs, but some have only two. Their bodies are covered with scales, and their mouths filled with teeth. This order includes all the crocodiles and lizards. The crocodiles have broad flat tongues, attached throughout to the jaws, and the lizards have long narrow tongues, which they can extend to a great distance from the mouth.
3. *The Ophidian Reptiles,* are the snakes and serpents. The body is covered with scales, but it is destitute of feet. The heart has two auricles, and the lungs are generally well developed. Serpents are frequently furnished with poison-bags at the base of their teeth.
4. *The Batrachian Reptiles* include the frogs and toads. The heart has only one auricle, and the body is naked. The greater part of these reptiles undergo a transition from a fish-like tadpole furnished with gills, to a four-legged animal with lungs. Others never lose their gills, though they acquire lungs, and of this kind are the siren and the proteus.

THE PISCES,

Or Fishes, are defined by Cuvier to be vertebrated animals with red blood, breathing through the medium of water by means of their branchiæ or gills. To this definition may be added, that fishes have no neck, and that the body generally tapers from the head to the tail; that most of the species are furnished with air-bladders which enable them to swim; and that their bodies are generally covered with scales. The heart has only one auricle, and the blood is cold. The gills require to be kept moist to enable the fish to breathe, and as soon as they become dry, the fish dies. Thus fishes with large gills die almost as soon as they are taken out of the water; while those with very small gills, like the eel, live a long time. Fishes have no feet, but are furnished with fins. The scientific knowledge of Fishes is called Ichthyology. Fishes are first divided into two great series, viz. the Bony Fishes, and the Cartilaginous Fishes, and these are again subdivided into nine orders, as follows;—

OSSEOUS OR BONY FISHES.

1. *Acanthopterygii,* or fishes with hard fins.
2. *Malacopterygii abdominales,* or fishes with soft abdominal fins.
3. *Malacopterygii sub-brachiati,* or fishes with soft fins under the gills.
4. *Malacopterygii apodes,* or fishes without ventral fins.
5. *Lopobranchii,* or fishes with tufted branchiæ.
6. *Plectognathii,* or fishes with the upper jaw fixed.

CHONDROPTERYGIC, OR CARTILAGINOUS FISHES.

7. *Cyclostomi,* or fishes with jaws fixed in an immoveable ring, and with holes for the gills.
8. *Selachii,* or fishes with moveable jaws and holes for the gills.
9. *Sturiones,* with the branchiæ in the usual form.

Of the bony fishes the *Acanthopterygii*, or fishes with hard spiny fins, are divided into fifteen families, the principal of which are the perch family, the mailed cheek fishes, including the gurnards, the flying fish of the Mediterranean, and the sticklebacks, or jack banticles; the mackerel family, including the tunny, bonito, and sword-fish; the pilot-fish, the dolphin of the Mediterranean, so celebrated for the beauty of its dying tints, and the john doree. Among the *Malacopterygii abdominales*, or soft-finned fishes, that have their ventral fins suspended from the abdomen, the most interesting are the carp family, the pike family, the flying-fish of the ocean, the salmon family, and the herr.g family, including the sprat, pilehard, and anchovy.

The *Malacopterygii sub-brachiati* are soft-finned fishes, with the ventral fins beneath the pectorals; the principal of which are the cod family, including the haddock, whiting, and ling; the flat-fish family, including soles, turbot, plaice, and flounders; and the suckers or Limp-fish.

The *Malacopterygii apodes* are confined to the eel family.

The *Lopobranchii* include the pipe-fish and other fishes of similar form.

The *Plectognathi* comprise the very singular forms of the balloon-fish, the sun-fish, and other similar fishes.

The *Chondopterygii*, or *Cartilaginous fishes*, are divided into three orders, viz. the *Sturiones*, or sturgeon family; the *Selachi*, or sharks and rays, including the torpedo; and the *Cyclostomi*; or lamp-prey family.

THE MOLLUSCOUS ANIMALS

Have no bones except their shells. Their sense of feeling appears to be very acute, but the organs for the other senses are either wanting or very imperfect. The blood is cold and white, and the heart often consists of only one ventricle; a few of them have imperfect lungs, but the greater number breathe through gills. They have all the power of remaining a long time in a state of rest, and their movements are either slow or violently laborious. Some of them appear incapable of locomotion. They produce their young from eggs, but some lay their eggs on a part of their own body, where the young are hatched. The following are Cuvier's six classes:—

1. *Cephalopoda*, or *Head-footed Mollusca*. These animals are furnished with long fleshy arms or feet, proceeding from the head, which is not distinct from the body, and on which they crawl. There is only one order, which includes the cuttle-fish, nautilus, and belemnites.
2. *Pteropoda*, or *Wing-footed Mollusca*. These animals have two membranous feet or arms, like wings, proceeding from the neck. There is only one order, which contains six genera, the best known of which, is the *Hyalea*, the shell of which is commonly called Venus's chariot.
3. *Gasteropoda*, or *Body-footed Mollusca*. All these animals crawl with the flat part of the body, which acts as a kind of sucker. There are nine orders; but the common snail will give an idea of the habits of the class.

4. *Acephala* or *Headless Mollusca*. These animals have two shells united by a hinge, and no distinct head, like the oyster. There are two orders.
5. *Brachiopoda* or *arm-footed Mollusca*. These animals have two shells not united by a hinge. There are two orders.
6. *Cirrhopoda*, or *fringed-footed Mollusca*. These animals are soft and headless, like the oyster, but they are furnished with fringe or hair—like tentacula; as for example, the lepas or barnacle.

It must be observed, that the system of Cuvier is not generally adopted in classing shells, as his classes are formed from the animals alone, and we have many shells, the animals of which are unknown; and that of Lamarck is preferred, as he founds his classes partly on the animals, and partly on the shells. Lamarck divides the species into three classes, viz. *Cirrhopoda*, *Conchifera*, and *Mollusca*. The first is the same as the sixth class of Cuvier; the *Conchifera* are cephalous, or headless, and two-shelled or bivalves, like the oyster; and the *Mollusca* are univalves and cephalous, that is, they have only one shell, and a kind of head, like the snail.

THE ARTICULATED ANIMALS

Have no back-bone. The covering of the body is sometimes hard and sometimes soft, but it is always divided into segments by a number of transverse incisions. The limbs, when the body is provided with any, are jointed; and they can be separated from the body without any serious injury being sustained by the animal, new limbs being shortly after formed to replace them. The senses of tasting and seeing are more perfect than those of the *Mollusca*, though that of feeling seems much less acute. In other respects the four classes differ considerably from each other.

- I. *The Annelida*, or *Red-Blooded Worms*, have no heart, properly so called, but have sometimes one or more fleshy ventricles. They breathe through branchiæ. Their bodies are soft, and more or less elongated, being divided into numerous rings or segments. The head, which is at one extremity of the body, can scarcely be distinguished from the tail, except by having a mouth. These animals have no feet, properly so called, but they are furnished with little fleshy projections, bearing tufts of hairs or bristles, which enable them to move. They are generally of carnivorous habits, and live principally on the blood of other creatures. They lay eggs, but the young are frequently hatched before exclusion, and hence these creatures are said to be oviparous. Their study is called Helminthology. As examples of the three orders of this class may be mentioned the serpulæ or worm-like animals, often found on shells, the common earth-worm, and the leech family.
- II. *The Crustacea* comprise the shell-fish commonly called crabs, lobsters, shrimps, and prawns. They have a distinct head, furnished with antennæ, eyes, and mouth; and their bodies are covered with a crust or shell, divided into segments by transverse incisions, the segments being united by a strong membrane. The crust turns red by boiling, whatever colour it may have borne while the animals were in a living state. Once a year these animals moult, throwing off their old crust or shell, and

forming a new one, the animal remaining in a naked, and greatly weakened state, during the intermediate time. The Crustacea swim with great ease, but on land their motions are generally cramped and awkward; and they are confined to crawling, or leaping by means of the tail. When a limb is injured they possess the extraordinary power of throwing it off, and forming a new one. Another singular property which they possess is that of emitting light, which as it is well known the shell of a lobster, or crab, will do when beginning to decay; there is indeed a smaller kind of crab, called *Cancer fulgens*, which produces a luminous effect when alive. The Crustacea lay eggs, and the young of some of the species undergo a transformation before they attain their full size. The Crustacea were divided into two sections and seven orders by Cuvier, which are as follow:

SECTION I.—*Malacostraca*.

Shell solid, legs ten or fourteen, foot-jaws six or ten, mandibles two, maxillæ four; mouth with a labrum.

Sub-section 1. *Podophthalma*, eyes on foot-stalks.

ORDER 1. *Decapoda*, legs ten.

Sub-order 1. *Brachyura*, the crabs.

Sub-order 2. *Macroura*, the lobsters.

ORDER 2. *Stomapoda*, legs more than ten.

Sub-section 2. *Edriophthalma*, eyes not on foot-stalk.

ORDER 3. *Amphipoda*, body compressed; mandibles palpigerous.

ORDER 4. *Læmodipoda*, abdomen rudimental, with only the rudiments of one or two pairs of appendages.

ORDER 5. *Isopoda*, body depressed; abdominal appendages flat; mandibles not palpigerous.

SECTION II.—*Entomostraca*.

Shell not solid; legs variable in number; mouth variable.

ORDER 6. *Branchiopoda*. Integuments horny, branchiæ feathery, forming part of the feet.

ORDER 7. *Pacilopoda*, mouth suctorial.

Sub-order 1. *Xiphosura*, or king-crabs.

Sub-order 2. *Siphonostoma*, or fish parasites.

Lamarck divides the Crustacea into only the two following orders, viz.

1. *The Homobranchia*, which have the body covered with an entire shell, and fringed appendages in the tail to retain the eggs. This order contains the crab family, which have an orbicular body and an imperfectly developed tail; and the lobster family, including the shrimp and the prawn, which have a cylindrical body and a fully-developed tail.

2. *The Heterobranchia*, which consist chiefly of the wood-lice, the brine-worm, and the ship-worm. The curious fossil animals, called trilobites, belong to this division.

III. *The Arachnida* are defined by Lamarck to be oviparous animals, provided with six or more articulated legs, not subject to meta-

morphoses, and never acquiring any new kinds of organs. Their respiration is either by means of air-sacks, which serve for lungs, or of a kind of tube with circular openings for the admission of air. There is a rudimentary heart and circulation in most of the species. There are two orders; those with lungs, and those without.

ORDER I. Pulmonaria. The Arachnides comprised in this division have air-sacks, which serve for lungs, a heart with distinct vessels, and from six to eight simple eyes. There are two distinct families: viz. *Aranides*, comprising all the spiders and spinners; and *Pedipalpi*, comprising the tarentula and scorpions.

ORDER II. Trachearia. These Arachnides are distinguished by their respiratory organs, which consist of radiated or branched tracheæ, receiving air by two circular openings. They have no circulation, and their eyes vary from two to four. The principal animals belonging to this division are the long-legged spiders (*Phalangium*), the book-worms, and the mites (*Acarus*), including the gardener's pest, the little red spider (*Acarus telarius*), the cheese mite (*Acarus Siro*), and the harvest bug (*Acarus or Lept. autumnalis*).

IV. The Insecta form the fourth and last class of articulated animals, and they derive their name from the Latin word *insectum*, which signifies cut into, in allusion to the distinct divisions of head, thorax, and abdomen in the true insects: and in contradistinction to the Annelides, the bodies of which present no such divisions. The true insects are defined as animals without vertebrae, possessing six feet, with a distinct head furnished with antennæ, and breathing through stigmatic openings, which lead to interior tracheæ. The Myriapoda have, however, more feet. The following are the twelve orders into which this class is divided.

SECTION I.—*Insects undergoing Metamorphoses.*

1. *Coleoptera*, (from two Greek words signifying sheathed wings). These are the beetles, which are all furnished with membranous wings, with which they fly, and which are protected by bony upper wings, or wing-cases, called elytra. They are all masticators, and are all provided with mandibles or projecting jaws, and maxillæ.
2. *Orthoptera*, or straight-winged insects. This order comprises the crickets, grasshoppers, locusts, and similar insects. They have their upper wings of the consistence of parchment, and have mandibles and maxillæ.
3. *Hemiptera*, or half-winged insects, have frequently half the upper wing membranous, like the under ones, while the other half is leathery. To this division belong the bugs, the water-scorpions, the cicadæ or froghoppers, and the aphides. These insects have neither mandibles nor maxillæ, but in their place have a sheath and sucker.
4. *Neuroptera*, or nerved-winged insects, such as the dragon-flies, have both pairs of wings membranous, naked, and finely reticulated. The mouth is adapted for mastication, and furnished with mandibles and maxillæ.
5. *Hymenoptera*, membranous winged insects, such as bees, wasps,

ichneumon flies, &c. All the four wings are membranous, but they have fewer nervures, and are not reticulated like those of the preceding order. The mouth is furnished with mandibles and maxillæ, and the abdomen is terminated either by an ovipositor or a sting.

6. *Lepidoptera*, or scaly-winged insects.* These are the butterflies and moths, which are characterised by the farinaceous or scaly aspect of their wings, and the tubular or thread-like extension of the parts of the mouth.
7. *Strepsiptera* or *Rhipitera*, with twisted wings. These creatures resemble the ichneumon, in laying their eggs in the bodies of other insects, though they generally attack wasps and bees. The principal genera are *Xenos* and *Stylops*.
8. *Diptera*, or double-winged insects, including the flies. The mouth is furnished with a proboscis, and there are two small wings called *halteres* placed behind the true wings, which act as balancers.
9. *Suctoria*, or sucking insects, such as the flea, which have no wings, but are furnished with an apparatus for sucking blood.

SECTION II.—*Insects not undergoing Metamorphoses.*

10. *Thysanoura*, or spring-tail insects. These creatures are of small size, and without wings; they are found in crevices of wood-work, or under stones. The principal genera are *Lepisma* and *Podura*.
11. *Parasita*, or parasitical insects, such as the louse. They are also without wings.
12. *Myriapoda*. This order is made a separate class by many naturalists, as the creatures contained in it are distinguished from the true insects by the great number of their feet; by the want of distinct divisions into thorax and abdomen; and by the great number of segments into which the body is divided. The principal insects in this order are included in the Linnæan genera *Julus* and *Scolopendra*, commonly called centipedes.

The term larva is applied to the young of all insects included in the first seven orders, when first hatched. The different kinds have, however, other names; that is to say, the larva of a butterfly, or moth, is called a caterpillar; that of a beetle, a grub; and that of a fly, a maggot. The larva changes its skin several times, and at last goes into the pupa state, when it is called a chrysalis, an aurelia, or a nymph. Sometimes the pupa is wrapped up in a loose outer covering called a cocoon. From the pupa in time bursts forth the imago, or perfect insect. The Apterous, or wingless true insects, and the Myriapoda, which are also without wings, do not undergo any metamorphosis. Entomology is a very difficult study, from the great number of systems that have been broached by modern naturalists, and from the changes continually making in the names of the insects.

THE RADIATED ANIMALS.

Are so called because their organs of locomotion, and even their internal viscera are generally arranged in a circle round a centre, so as to

give a radiated appearance to the whole body. The animals included in this class are the very lowest in the scale; they have scarcely any external senses; their movements are slow, and almost their only sign of life is a craving for food. Some of them, however, have a distinct mouth and alimentary canal, with an anal orifice; others have a bag-like stomach with a kind of mouth, through which they both take their food and reject their excrements; while others have no mouth, and only appear to absorb nourishment through pores. In the like manner, though some are oviparous, others may be propagated by division like plants. There are five classes, which are as follow.

- I. *Echinodermata*, or sea-urchins. These animals have a leathery or crustaceous skin or shell, commonly covered with numerous tubercles. The mouth is in the centre of the animal, and is armed with five or more pieces of bone, which serve as teeth; the stomach is a loose bag; the organs for respiration are vascular; and the animals are oviparous. They are furnished with tentacular tubes, which serve as arms or feet, and which they can push out and draw back at pleasure; and they have yellowish or orange-coloured blood, which appears to circulate. Cuvier divides this class into those with feet, and those without; but Lamarck, whose arrangement is now generally followed, divides them into three orders, viz.
 1. *The Fistuloides* or *Holothurida*, which have cylindrical bodies, leathery skins, and mouths surrounded by tentacula. These creatures live in the sea, or in the sands on the sea-shore; and the trepang, or eatable worm of the Chinese, is one of them.
 2. *The Echnides*. These are the sea-urchins, properly so called, and the shells, when the animals are out of them, are called sea-eggs. The Echnides live in the sea, and they have long arms to secure their prey, and two series of bone-like teeth to crush it. They lay eggs, and the roe, or imperfect eggs, occupy a large portion of the space within the shell when the animal is still alive.
 3. *The Stellerides*, or *Asterias*, are the star-fish. The mouth in these creatures is in the middle of the lower surface, and it has a membranous lip, capable of great dilation, but furnished with angular projections for capturing and crushing its prey. The skin is soft but leathery, and it is covered on the back with spongy tubercles or scales. The rays are hollow beneath, and furnished with tentacula, by the aid of which the star-fish manages to crawl backwards, forwards, or sideways, as the case may be, any of the rays serving as a leader. These animals are found on the sea-shore, forming large beds, which are washed over by the sea. *The Crinoidea*, or stone-lilies, of which such curious fossil specimens have been found, are nearly allied to the star-fish.
- II. *The Intestina*, or *Entozoa*. The intestinal worms were divided into two kinds by Cuvier, viz. *the Cavitaires*, including the worms of children, the bots in horses, and other cylindrical worms; and the *Parenchymateux*, or flat worms; such as the fluke in sheep and the tape-worm in human beings.
- III. *Acalephæ*, or *Sea-Jellies*. These creatures are of a soft and jelly-like substance, with a thin skin, and an unarmed mouth. There are two kinds, the *Medusides*, which have round bodies, and the

Acalephes, the bodies of which are of an irregular shape. The Medusae are very numerous, and produce that beautiful phosphorescent light noticed by voyagers in the Australian seas. The most interesting of the Acalephes is the Portuguese man-of-war, or *Physalia*; and some naturalists include in it the actinia, or sea-anemone.

IV. *Polypes* or *Anthozoa*, according to Cuvier, were arranged in three orders, viz. the Polypes, the Lithophytes, and the Sponges; but to these Lamarck has added the Molluscous Polypes, calling the others the Radiated Polypes. The molluscous polypus is a very small animal. The mouth is wide and circular, and it is surrounded by a fringe of tentacula. This kind of polypi always dwells in a calcareous cell, which is so connected with its body that it cannot leave it. The crust, or shell, is called the polypidom, and as a number of polypi dwell together their polypidom soon becomes of a great size; particularly as when the first set of polypi die, their cells remain to be increased by the next generation. The sea-mat, or floster, is a specimen of these animals. The radiated polypi are divided into three orders, which are thus named and characterised.

1. *Helianthoida*. Some naturalists include in this order the actinia, or sea-anemone; but all agree that it contains the animals of the madrepores, sea-mushrooms, and brainstones, which live in communities, and possess the power of secreting calcareous matters, which they emit to form these stony substances.
2. *Asteroida*. Some of the animals belonging to this division are called sea-pens, and others form some of the different kinds of coral, particularly that used for necklaces, &c.
3. *Hydroïda*. This order includes the fresh-water polypi, which, it is well known, by the experiments that have been tried, may be cut in pieces and even turned inside out without destroying life. The *Lithophytes* are divided into two families, viz. those covered with a thick jointed crust, which are the coral insects, or *Coralines* of Linnæus; and those called by De Blainville *Fucoïdeæ*, which closely resemble sea-weeds. The *Sponges* differ so much from the other zoophytes, that they have been formed by De Blainville into a separate class, under the name of *Amorphozoa*. The sponge, in its native state, appears to be filled with a kind of jelly, but when examined with a microscope, this jelly will be discovered to be full of transparent spherical granules, which are living animal matter. Sometimes sandy or calcareous particles are found in sponge, as though it were on the point of being changed into a polypidom.

V. *The Infusoria* or *Animalcula*, are so small as to be invisible to the naked eye, and they are all inhabitants of liquids. Cuvier arranged them in two orders, one of which he called *Les Rotiferes*, and the other *Les Infusories homogenes*, but by Lamarck's system, which is now generally followed, the first of these divisions is now included in the polypi; and the latter is divided into two orders, one with external organs and the other without. The great number of these animals and their singular forms must be present to the mind of every one who has seen a drop of stagnant water magnified in a powerful microscope.

EXPLANATION
OF
TERMS USED IN NATURAL HISTORY.

<i>Amphibious.</i>	Capable of living both on the land and in the water.
<i>Animalcules.</i>	Small animals, visible only with the assistance of the microscope.
<i>Annulated.</i>	Marked with rings.
<i>Antennæ.</i>	The horns or feelers of insects.
<i>Antlers.</i>	Horns overhanging the brows.
<i>Aquatic.</i>	Living or growing in the water.
<i>Bifid.</i>	Divided into two parts.
<i>Bimaculated.</i>	With two spots, or two series of spots.
<i>Bivalve.</i>	With two shells or openings.
<i>Callosity.</i>	A hard lump, an excrescence.
<i>Canine.</i>	Of the dog kind.
<i>Carnivorous.</i>	Feeding on flesh.
<i>Cere.</i>	A skin over the bill of birds; sometimes moveable, as in parrots.
<i>Cetaceous.</i>	Of the whale kind.
<i>Cinereous.</i>	Of the colour of ashes.
<i>Columbine.</i>	Of the dove or pigeon kind.
<i>Cordiform.</i>	Heart-shaped.
<i>Crustaceous.</i>	Covered with a shell or crust; as lobsters, crabs, &c.
<i>Digitated.</i>	Having the feet divided into toes or fingers; as in cats and dogs.
<i>Dorsal.</i>	Belonging to the back.
<i>Elytra.</i>	The wing-cases of insects of the beetle-tribe.
<i>Exsanguineous.</i>	Without red blood, as worms.
<i>Entomology.</i>	A description of insects.
<i>Feelers.</i>	The antennæ or horns of insects.
<i>Feline.</i>	Belonging to the cat kind.
<i>Ferruginous.</i>	Of an iron or rust colour.
<i>Frugivorous.</i>	Feeding on seeds.
<i>Furcated.</i>	Forked
<i>Gallinaceous.</i>	Belonging to the hen kind.
<i>Gestation.</i>	The time of going with young.
<i>Granivorous.</i>	Feeding on grain.
<i>Gregarious.</i>	Associating together.
<i>Herbivorous.</i>	Feeding on grass.
<i>Hoof.</i>	The horny substance that protects the external part of the foot in some quadrupeds.

<i>Ichthyology.</i>	A description of fishes.
<i>Imbricated.</i>	Tiled, or lying over each other
<i>Incubation.</i>	The act of hatching eggs.
<i>Insectivorous.</i>	Feeding on insects.
<i>Laminated.</i>	Covered with, or divided into plates or scales.
<i>Larva.</i>	The young of insects.
<i>Lateral.</i>	Belonging to the side, placed sideways.
<i>Mandibles.</i>	Upper and lower, the two divisions of a bird's beak, or projecting jaws of an insect.
<i>Migratory.</i>	Coming and going at certain seasons.
<i>Multivalve.</i>	With many shells or openings.
<i>Nictitating.</i>	Winking; applied to a membrane with which birds cover their eyes at pleasure.
<i>Obscured.</i>	Of a darkish colour.
<i>Olfactory.</i>	Relating to smell.
<i>Opperculum.</i>	A shield or cover.
<i>Ornithology.</i>	A description of birds.
<i>Oviparous.</i>	That lays eggs.
<i>Parturition.</i>	The act of bringing forth young.
<i>Passerine.</i>	Belonging to the sparrow tribe.
<i>Pectoral.</i>	Belonging to the breast.
<i>Pendulous.</i>	Hanging down.
<i>Piscivorous.</i>	Feeding on fishes.
<i>Predaceous.</i>	Formed to pursue prey.
<i>Prehensile.</i>	Capable of laying hold of.
<i>Proboscis.</i>	The flexible trunk of the elephant, bee, &c.
<i>Quadrifid.</i>	Divided into four parts.
<i>Quadruped.</i>	Four-footed.
<i>Reptiles.</i>	Animals of the serpent tribe, with legs.
<i>Ruminating.</i>	Chewing the cud.
<i>Scarrous.</i>	Rough.
<i>Scapulars.</i>	Shoulders.
<i>Semilunar.</i>	In the form of a half-moon.
<i>Setaceous.</i>	Having bristles or strong hairs.
<i>Spiral.</i>	Winding like a screw.
<i>Striated.</i>	Streaked or striped.
<i>Subulated.</i>	Formed like an awl.
<i>Tentacula.</i>	The feelers of snails and other mollusca.
<i>Testaceous.</i>	Covered with a shell, as oysters.
<i>Trifurcated.</i>	Three-forked.
<i>Truncated.</i>	Appearing as if cut off.
<i>Umbrageous.</i>	Spreading; thick, tufted.
<i>Univalve.</i>	With one shell or opening.
<i>Ventral.</i>	Belonging to the belly.
<i>Vertebated.</i>	Having a jointed spine-bone.
<i>Vivaporous.</i>	Bringing forth the young alive.
<i>Webbed.</i>	Connected with a membrane, as the claws of aquatic birds.
<i>Zoologists.</i>	Writers on animated nature.
<i>Zoology.</i>	The history of animated nature.
<i>Zoophyte.</i>	An animal plant, or sensitive vegetable.

TABLE OF CONTENTS.

BOOK I.—QUADRUPEDS, OR FOUR-FOOTED BEASTS.

POPULAR AND SCIENTIFIC NAMES OF THE ANIMALS DESCRIBED.

* * Where no synonyme is given, the Linnæan name is the only one in use; and when the synonymes are seldom used, they are marked thus *. When no Linnæan name is given, the animal was not described by Linnæus

1. Carnivorous, or Flesh-eating Animals.

English Name.	Linnæan Name.	Synonymes.	Page.
LION . . .	Felis Leo . . .	*Leo vulgaris.— <i>Leach</i>	1
LIONESS . . .	Ibid . . .		6
TIGER . . .	Felis Tigris . . .		8
LEOPARD . . .	Felis Leopardus . . .	Felis Pardus.— <i>Cuv.</i>	10
PANTHER . . .	Felis Pardus . . .	Felis Pardus var.	11
OUNCE . . .			12
HUNTING LEOPARD, OR CHEETAH . . .	Felis jubata . . .	Cynaclurus jubata.— <i>Wag.</i>	13
JAGUAR . . .	Felis onca . . .		14
PUMA . . .	Felis concolor . . .	{ Felis Puma.— <i>Trail</i> *Leo Americanus.— <i>Her.</i> *Puma concolor.— <i>Jard.</i>	15
CARACAL LYNX . . .	Felis caraca . . .	*Lynchus vulgaris.— <i>Gray.</i>	16
CAT . . .	Felis Catus . . .	Felis Catus var. domestica	17
DOGS . . .	{ Canis familiaris and var . . .		19-31
FOXHOUND . . .			19
BLOODHOUND . . .			21
POINTER . . .			22
MASTIFF . . .			23
BULLDOG . . .			24
SPANIEL . . .			27
WATER SPANIEL . . .			28
NEWFOUNDLAND DOG . . .			29
GREYHOUND . . .			31
FOX . . .	Canis Vulpes . . .	Vulpes vulgaris.— <i>Briss.</i>	32
WOLF . . .	Canis Lupus . . .	*Lupus vulgaris	34
JACKAL . . .	Canis aureus . . .		36

English Name.	Linnean Name.	Synonymes.	Page.
HYÆNA . . .	Canis Hyæna . . .	Hyæna vulgaris.—Cuv. . .	37
BLACK BEAR . . .	Ursus Americanus	38
BROWN BEAR . . .	Ursus Arctos	38
POLAR BEAR	Ursus maritimus.—Gmel. . .	41
RACCOON . . .	Ursus Lotor . . .	Procyon Lotor.—Cuv. . .	42
BADGER . . .	Ursus Meles . . .	Meles vulgaris.—Des. . .	44
COATI-MONDI . . .	Viverra Nasua . . .	Nasua nasica.—F. Cuv. . .	45
CIVET CAT . . .	Viverra Civetta	46
ICHNEUMON, or EGYPTIAN MANG- GOUSTE . . .	Viverra Pharaonis . . .	{ Herpestes Pharaonis.— Illig. Mangusta Pha- raonis.—Cuv. }	47
WEASEL . . .	Mustela vulgaris	49
FERRET . . .	Mustela furo . . .	*Viverra furo.—Shaw. . .	51
GENET . . .	Viverra Genetta . . .	Genetta vulgaris.—Cuv. . .	52
POLICAT . . .	Mustela putorius . . .	Putorius vulgaris.—Cuv. . .	53
SABLE . . .	{ Mustela or Martes Zibellina . . . }	54
MARTEN . . .	Mustela Martes . . .	Martes fagorum.—Ray. . .	54
OTTER . . .	Mustela Lutra . . .	Lutra vulgaris.—Erzl. . .	55
SEAL . . .	{ Phoca vitulina . . . Trichechus Rosma- rus . . . }	{ *Phoca variegata.—Niel Calocephalus vitulinus.— Cuv. }	57
WALRUS	59

SECTION II.—Insect-eating Animals.

HEDGEHOG . . .	Erinaceus Europæus	61
MOLE . . .	Talpa Europea . . .	Talpa vulgaris.—Briss. . .	63
SHREW . . .	Sorex araneus	65
WATER SHREW . . .	Sorex Fodiens	66

SECTION III.—Chiropterous Animals.

BAT . . .	Vespertilio noctula	67
-----------	---------------------------	-----------	----

SECTION IV.—Marsupialia, or Pouch-bearing Animals.

KANGAROO	{ Macropus major.—Shaw and Cuv. *Halmaturus Illig. and *Kangurus. —Desm. }	68
OPOSSUM . . .	Didelphis Virginiana	70

SECTION V.—Rodentia, or Gnawing Animals.

BEAVER . . .	Castor Fiber	71
HARE . . .	Lepus timidus	73
RABBIT (Wild) . . .	Lepus cuniculus	76
RABBIT (Domestic)	77
SQUIRREL . . .	Sciurus vulgaris	78
AMERICAN FLY- ING SQUIRREL . . .	Pteromys volucella	78
MARMOT, or AL- PINE RAT . . .	Mus alpinus . . .	{ Arctomys Marmotta.— Gmel. }	79

English Name.	Linnæan Name.	Synonymes.	Page.
GUINEAPIG . . .	Mus porcellus	{ <i>Cavia cobaya.</i> — <i>Pall.</i> <i>Cavia</i> <i>aperea.</i> — <i>Erxl.</i> <i>Hydro-</i> <i>chærus aperea.</i> — <i>F. Cuv.</i> }	80
MOUSE . . .	Mus musculus	81
RAT . . .	Mus decumanus	82
WATER RAT . . .	Mus amphibius	{ <i>Mus aquaticus.</i> — <i>Briss.</i> <i>Lemnus aquaticus.</i> — <i>F.</i> <i>Cuv.</i> — <i>Arvicola amphi-</i> <i>bia.</i> — <i>Desm.</i> and <i>Jenyns.</i> <i>Arvicola aquatica.</i> — <i>Hem.</i> <i>Fiber zibethicus.</i> — <i>Des.</i> <i>Ondrata zibethicus.</i> — <i>Lacep.</i> }	84
MUSK RAT	{ <i>Myoxis muscardinus.</i> — <i>Gmel.</i> <i>Myoxis glis</i> <i>Dipas Jerboa.</i> — <i>Gmel.</i> <i>Mus sagitta.</i> — <i>Pall.</i> <i>Chincella lanigera</i> . . . }	85
DORMOUSE . . .	Mus avellanarius	86
JERBOA . . .	Dipas jaculus	86
CHINCHILLA	88
PORCUPINE . . .	Hystrix cristata	89

SECTION VI.—EDENTATA, or Toothless Animals.

SLOTH . . .	Bradypus tridactylus	90
ARMADILLO . . .	Dasypus sexcinctus	92
ANT EATER . . .	{ <i>Myrmecophaga ju-</i> <i>bata</i> }	93
DUCK-BILLED PLATYPUS	{ <i>Ornithorhynchus para-</i> <i>doxus.</i> — <i>Blum.</i> <i>Platypus</i> <i>anatinus.</i> — <i>Shaw.</i> . . . }	94
AUSTRALIAN HEDGEHOG . . .	Echidna setosa	95

SECTION VII.—Pachydermata, or thick-skinned Animals.

ELEPHANT . . .	Elephas Indicus	96
HIPPOTAMUS, or RIVERHORSE }	Hippopotamus am- phibius }	99
RHINOCEROS . . .	Rhinoceros Indicus	100
HOG (<i>Domestic</i>) . . .	Sus scrofa	101
WILD BOAR . . .	Sus scrofa	<i>Sus aper.</i> — <i>Briss.</i>	103
TAPIR . . .	Tapir Americanus	105
HORSE . . .	Equus caballus	106
ASS . . .	Equus Asinus	<i>Asinus vulgaris.</i> — <i>Gray.</i>	109
WILD A	112
MULE	113
ZEBRA . . .	Equus Zebra	114

SECTION VIII.—Ruminating Animals.

BULL . . .	{ <i>Bos Taurus,</i> var. do- <i>mesticus</i> }	116
COW	118
WILD BULL . . .	{ <i>Bos Taurus,</i> var. <i>Scoticus</i> }	119
BUFFALO . . .	<i>Bos Bubalus</i>	<i>Bubalus Caffer</i>	121

English Name.	Linneæan Name.	Synonymes.	Page.
BISON	Bos Bonasus	123
BRAHMIN BULL, } OF ZEBU	Bos Taurus, var. } Indicus	125
SHEEP	Ovis Aries	*Capra ovis.—Blum.	126
RAM	128
WALLACHIAN RAM	Ovis Musmon	128
ARGALI, OF WILD } SHEEP OF ASIA }	Ovis Ammon	129
GOAT	Capra Hircus	129
IBEX, OF BOQUE- } TIN &	Capra Ibex	130
ANTELOPE	Antilope cervicapra	131
GAZELLE	Antilope Dorcas	132
SHAMOI	Antilope rupicapra	133
NYL GHAU	Antilope picta	134
GNU	Antilope Gnu.—Gmel.	135
STAG	Cervus Elephas	136
ROEBUCK	Cervus capreolus	138
FALLOW DEER	Cervus Dama	139
ELK	Cervus Alces	140
REINDEER	{ Cervus Tarantus, or } Rangifer	141
AXIS	Cervus Axis	143
WAPITI	{ Cervus canadensis.—Gmel. } *Cervus strongyloceros.— } Schres	144
GIRAFFE	Camelopardalis Giraffa	145
CAMEL	Camelus Bactrianus	148
DROMEDARY	Camelus Dromedarius	150
LLAMA	Camelus Llama	Auchenia glama.—Illig.	152

SECTION IX.—Quadrupana, or four-handed Animals.

OURANGOUTANG, } OF CHIMPANSEE }	Simia troglodytes	{ Troglodytes niger.—Geoff. } and Cuv.	153
BARBARY APE	Simia inuus	Inuus sylvanus.—Cuv.	154
BABOON	Simia mormon	{ Cynocephalus Mormon.— } Desm. and Cuv.	155
PROBOSCIS MON- } KEY	Nasalis larvatus—Geoff.	156
DIANA MONKEY	Simia Diana	{ Cereopithecus Diana.— } Geoff.	156
CAPUCHIN MON- } KEY	Simia capucina	Cebus capucinus.—Des.	158
SPIDER MONKEY	Ateles arachnoides.—Geoff.	158
OUISTITI, OR } MARMOZET	Simia Jacchus	Jacchus vulgaris.—Geoff.	159
MARIKINA	Ateles hypoxanthus.—Kuhl.	159
LEMUR & MON- } GOOS	Lemur Mongoz	{ Lemur nigrifrons and al- } bifrons.—Geoff.	160

BOOK II.—INHABITANTS OF THE AIR.

SECTION I.—RAPTORES.—Diurnal Birds of Prey.

English Name.	Linnean Name.	Synonyms.	Page
GOLDEN EAGLE	Falco chrysaetos	Aquila chrysaetos	161
SEA EAGLE	Falco albicilla	Halæetus albicilla.— <i>Sav.</i>	164
BALD EAGLE	Falco leucocephalus	Halæetus leucocephalus. — <i>Sav.</i>	165
BLACK EAGLE	Falco melaneotos		167
CONDOR	Vultur Gryphus	Sarcorhampus Gryphus.— <i>Dum.</i>	169
VULTURE	Vultur Papa	Sarcorhampus Papa.— <i>Dum.</i>	170
BUZZARD	Falco Buteo	Buteo vulgaris.— <i>Bech.</i>	172
HONEY BUZZARD			174
GOSHAWK	Falco palumbarius	Astur palumbarius	175
SPARROW-HAWK	Falco Nisus	Accipiter Nisus.— <i>Ray.</i> Nisus communis	176
KITE	Falco Milvus	Milvus regalis.— <i>Cuv.</i>	178
FALCON	Falco communis	Falco peregrinus and Fal- co islandicus	179
MERLIN	Falco æsalon		182
HEN HARRIER	Circus cyaneus		183
OSPREY, or FISH- ING HAWK	Falco halæetus	Pandion halæetus	184

SECTION II.—Nocturnal Birds of Prey.

HORNED OWL	Strix Otus	Bubo Virginarius	185
HARFANG, OF SNOWY OWL	Strix nyctea		186
BARN OWL	Strix flammea		187

SECTION III.—Essencers, or Perching Birds.

BUTCHER-BIRD, or SHRIKE	Lanius excubitor		188
LITTLE BUTCHER BIRD	Lanius collurio		189
WATER OUZEL, or DIPPER	Sturnus Cinclus	Turdus Cinclus.— <i>Lath.</i> Merula aquatica.— <i>Briss.</i> Cinclus aquaticus.— <i>Bech.</i>	190
BLACKBIRD	Turdus Merula		191
THRUSH	Turdus musicus		192
MISSIL THRUSH	Turdus viscivorus		192
REDWING	Turdus iliacus		193
FIELDFARE	Turdus pilaris		194
RING OUZEL	Turdus torquatus		195
REDBREAST	Motacilla rubecula	Sylvia rubecula.— <i>Lath.</i> Rubecula familiaris Ficedula.— <i>Bech.</i>	195
NIGHTINGALE	Motacilla lusciniæ, Philomela lusciniæ	Sylvia lusciniæ.— <i>Lath.</i> Curruca lusciniæ.— <i>Bech.</i>	198
BLACKCAP	Motacilla atricapilla	Sylvia.— <i>Lath.</i> and Cur- ruca.— <i>Bech.</i>	201

English Name.	Linnean Name.	Synonyms.	Page.
WREN	{ Motacilla Troglodytes . . . }	{ Sylvia. — <i>Lath.</i> Troglodytes Europæus.— <i>Cuv.</i> Troglodytes vulgaris.— <i>Flem.</i> }	202
WILLOW WREN.	Motacilla trochilus	{ Sylvia trochilus.— <i>Lath.</i> Regulus trochilus.— <i>Cuv.</i> }	203
WATER WAGTAIL	Motacilla alba	Motacilla boarula	204
YELLOW SHEPHERDESS	Budytes flava	Motacilla flava	205
SWALLOW	Hirundo rustica		206
MARYIN	Hirundo urbana		209
SWIFT	Cypselus apus		211
GOATSUCKER	{ Caprimulgus Europæus . . . }		212
SKYLARK	Alauda arvensis		213
WOODLARK	Alauda arborea		215
TITMOUSE	Parus major		216
YELLOW HAMMER	Emberiza citrinella	Emberiza flava	217
ORPHEAN	Emberiza hortulana		217
WHEATEAR	Motacilla Ænanthe	{ Sylvia Ænanthe.— <i>Lath.</i> Saxicola Ænantæ.— <i>Bech.</i> }	218
WHINCHAT			218
SPARROW	Fringilla domestica	{ *Pyrgita domestica.— <i>Cuv.</i> Passer domesticus.— <i>Ray.</i> Fringilla Linota.— <i>Gmel.</i> Linaria Linota.— <i>Cuv.</i> }	220
LINNET	Fringilla cannabina	*Serinus canarius.— <i>Briss.</i>	221
CANARY BIRD	Fringilla Canaria		222
CHAFFINCH	Fringilla coelebs		224
BULLFINCH	Loxia pyrrhula	Pyrrhula vulgaris.— <i>Tem.</i> Carduelis communis.— <i>Cuv.</i> ; Carduelis elegans.— <i>Steph.</i>	226
GOLDFINCH	Fringilla carduelis		227
STARLING	Sturnus vulgaris	Sturnus varius.— <i>Mey.</i>	229
RAVEN	Corvus corax		230
CROW	Corvus corone		233
ROOK	Corvus frugilegus		234
JACKDAW	Corvus monedula		236
MAGPIE	Corvus pica	Caudata Pica	237
CROSSBILL	Loxia		239
JAY	Corvus glandarius	{ Garrulus glandarius.— <i>Briss.</i> and <i>Cuv.</i> }	240
CHOUGH	Corvus graculus	{ Pyrrhocorax graculus.— <i>Tem.</i> Ptilonorhynchus Holosericeus.— <i>Kuhl.</i> Kitta.— <i>Lesson.</i> Grauculus.— <i>Cuv.</i> }	241
SATIN BOWER BIRD			242
BIRD OF PARADISE	Paradisea apoda		244
NUTHATCH	Sitta Europæa		246
WRYNECK	Yunx torquilla		247
CREEPER	Certhia familiaris		247
HUMMING-BIRD	Trochilus colubris		248
HOPOE	Upupa epops		249

SECTION IV.—Scansores, or Climbers.

English Name	Linnean Name.	Synonymes.	Page.
CUCKOO . . .	Cuculus canorus . . .	Cuculus hepaticus . . .	251
ROLLER . . .	Coracias garrula	255
KINGFISHER . . .	Alcedo ispida	255
WOODPECKER . . .	Picus viridus	257
TOUCAN . . .	Ramphastos tucanus	259
PARROT . . .	Psittacus erythacus	260
MACAW . . .	Psittacus aracanga . . .	{ Macrocerus aracanga— Viell. }	262
PAROQUET . . .	{ Psittacus NovaHol- landiæ . . . }	Palæornis torquatus . . .	263
RING PAROQUET . . .	Psittacus Alexandri . . .	Palæornis Alexandri . . .	263
COCKATOO . . .	Psittacus galeritus . . .	Plyctolophus galeritus . . .	264

SECTION V.—Gallinaceous Birds.

PEACOCK . . .	Pavo cristatus	265
TURKEY . . .	{ Meleagris Gallo- Pavo . . . }	267
GUINEA FOWL, . . .	Numida Meleagris	269
JUNGLE FOWL, or BRUSH TUR- KEY	Megapodius tumulus . . .	271
LYRE BIRD	Menura superba . . .	274
PHEASANT . . .	Phasianus Colchicus	277
PARTRIDGE . . .	Tetrao Perdix . . .	Perdix cinerea.—Lath. . .	279
RED-LEGGED PARTRIDGE . . .	Perdix Rufus	279
QUAIL . . .	Tetrao Coturnix . . .	{ Coturnix major.—Briss. Coturnix vulgaris.— Flem. Coturnix Euro- pæus.—Wils. Perdix Coturnix.—Lath. . . }	282
AMERICAN QUAIL . . .	Ortyx borealis	283
GROUSE, or MOOR FOWL . . .	Tetrao Scoticus . . .	{ Lagopus Scoticus.—Lath.; *Bonasa Scotica.—Briss. }	283
BLACK COCK . . .	Tetrao Tetrix . . .	*Uriogallis minor.—Ray. Lagopus vulgaris.—Wils.	285
PTARMIGAN . . .	Tetrao Lagopus . . .	{ Tetrao rupestris.—Gmel. Gallus domesticus.—Wils. }	285
COCK . . .	Phasianus Gallus . . .	Gallus Sonati . . .	286
BANKIVA COCK	288
JAGO COCK	289
SPANISH COCK	289
BANTAM FOWL	289
DODO . . .	Didus ineptus	290
RINGDOVE, or CU- SHAT . . .	Columba palumbus	292
STOCKDOVE . . .	Columba Ænas	293
PIGEON . . .	Columba livia	295
TURTLEDOVE . . .	Columba Turtur	297

English Name.	Linnean Name.	Synonymes.	Page.
BISON	Bos Bonasus	123
BRAHMIN BULL, } or ZEBU }	Bos Taurus, var. } Indicus }	125
SHEEP	Ovis Aries	*Capra ovis.—Blum.	126
RAM	128
WALLACHIAN RAM	Ovis Musmon	128
ARGALI, or WILD } SHEEP OF ASIA }	Ovis Ammon	129
GOET	Capra Hircus	129
IBEX, or BOQUE- } TIN T }	Cap. a Ibex	130
ANTELOPE	Antilope cervicapra	131
GAZELLE	Antilope Dorcas	132
SHAMUIS	Antilope rupicapra	133
NYL GHAU	Antilope picta	134
GTU	Antilope Gnu.—Gmel.	135
STAG	Cervus Elephas	136
ROEBUCK	Cervus capreolus	138
FALCOW DEER	Cervus Dama	139
ELK	Cervus Alces	140
REINDEER	{ Cervus Tarantus, or } Rangifer }	141
AXIS	Cervus Axis	143
WAPITI	{ Cervus canadensis.—Gmel. } *Cervus strongyloceros.— } Schres }	144
GIRAFFE	Camelopardalis Giraffa	145
CAMEL	Camelus Bactrianus	148
DROMEDARY	Camelus Dromedarius	150
LLAMA	Camelus Llama	Auchenia glama.—Illig.	152

SECTION IX.—Quadrupana, or four-handed Animals.

OURANGOUTANG, } or CHIMPANSEE }	Simia troglodytes	{ Troglodytes niger.—Geoff. } and Cuv. }	153
BARBARY APE	Simia inuus	Inuus sylvanus.—Cuv.	154
BABOON	Simia mormon	{ Cynocephalus Mormon.— } Desm. and Cuv. }	155
PROBOSCIS MON- } KEY }	Nasalis larvatus—Geoff.	156
DIANA MONKEY	Simia Diana	{ Cercopithecus Diana.— } Geoff. }	156
CAPUCHIN MON- } KEY }	Simia capucina	Cebus capucinus.—Des.	158
SPIDER MONKEY	Ateles arachnoides.—Geoff.	158
QUISTITI, or } MARMOZET }	Simia Jacchus	Jacchus vulgaris.—Geoff.	159
MARIKINA	Ateles hypoxanthus.—Kuhl.	159
LEMUR & MON- } GOOS }	Lemur Mongoz	{ Lemur nigrifrons and al. } bifrons.—Geoff. }	160

BOOK II.—INHABITANTS OF THE AIR.

SECTION I.—RAPTORES.—Diurnal Birds of Prey.

English Name.	Linnean Name.	Synonymes.	Page
GOLDEN EAGLE	Falco chrysaetos	Aquila chrysaetos	161
SEA EAGLE	Falco albicilla	Haliaeetus albicilla.—Sav.	164
BALD EAGLE	Falco leucocephalus	{ Haliaeetus leucocephalus. } —Sav.	165
BLACK EAGLE	Falco melaneotos		167
CONDOR	Vultur Gryphus	{ Sarcorhampus Gryphus.— Dum.	169
VULTURE	Vultur Papa	Sarcorhampus Papa.—Dum.	170
BUZZARD	Falco Buteo	Buteo vulgaris.—Bech.	172
HONEY BUZZARD			174
GOSHAWK	Falco palumbarius	Astur palumbarius	175
SPARROW-HAWK	Falco Nisus	{ Accipiter Nisus.—Ray. Nisus communis	176
KITE	Falco Milvus	Milvus regalis.—Cuv.	178
FALCON	Falco communis	{ Falco peregrinus and Fal- co islandicus	179
MERLIN	Falco aesalon		182
HEN HARRIER	Circus cyaneus		183
OSPREY, OF FISH- ING HAWK	Falco haliaeetus	Pandion haliaeetus	184

SECTION II.—Nocturnal Birds of Prey.

HORNED OWL	Strix Otus	Bubo Virginianus	185
HARFANG, OF SNOWY OWL	Strix nyctea		186
BARN OWL	Strix flammea		187

SECTION III.—Essessores, or Perching Birds.

BUTCHER-BIRD, OF SHRIKE	Lanius excubitor		188
LITTLE BUTCHER BIRD	Lanius collurio		189
WATER OUZEL, OF DIPPER	Sturnus Cinclus	{ Turdus Cinclus.—Lath. Merula aquatica.—Briss. Cinclus aquaticus.—Bech.	190
BLACKBIRD	Turdus Merula		191
THRUSH	Turdus musicus		192
MISSEL THRUSH	Turdus viscivorus		192
REDWING	Turdus iliacus		193
FIELDFARE	Turdus pilaris		194
RING OUZEL	Turdus torquatus		195
REDBREAST	Motacilla rubecula	{ Sylvia rubecula.—Lath. Rubecula familiaris Ficedula.—Bech.	195
NIGHTINGALE	{ Motacilla luscini- Philomela luscini- nia	{ Sylvia luscini- Curreca luscini- —Bech.	198
BLACKCAP	Motacilla atricapilla	{ Sylvia.—Lath. and Cur- ruca.—Bech.	201

English Name.	Linnean Name.	Synonymes.	Page.
WREN	{ Motacilla Troglodytes	{ Sylvania. — <i>Lath.</i> Troglodytes Europæus. — <i>Cuv.</i> Troglodytes vulgaris. — <i>Flem.</i>	202
WILLOW WREN.	Motacilla trochilus	{ Sylvania trochilus. — <i>Lath.</i> Regulus trochilus — <i>Cuv.</i>	203
WATER WAGTAIL	Motacilla alba	Motacilla boarula	204
YELLOW SHEPHERDESS	Budytes flava	Motacilla flava	205
SWALLOW	Hirundo rustica		206
MARTIN	Hirundo urbica		209
SWIFT	Cypselus apus		211
GOATSUCKER	{ Caprimulgus Europæus		212
SKYLARK	Alauda arvensis		213
WOODLARK	Alauda arborea		215
TITMOUSE	Parus major		216
YELLOW HAMMER	Emberiza citrinella	Emberiza flava	217
ORPHEAN	Emberiza hortulana		217
WHEATEAR	Motacilla Ænanthe	{ Sylvania Ænanthe. — <i>Lath.</i> Saxicola Ænantiae. — <i>Bech.</i>	218
WHINCHAT			218
SPARROW	Fringilla domestica	{ *Pyrgita domestica. — <i>Cuv.</i> Passer domesticus. — <i>Ray.</i>	220
LINNET	Fringilla cannabina	{ Fringilla Linota. — <i>Gmel.</i> Linaria Linota. — <i>Cuv.</i>	221
CANARY BIRD	Fringilla Canaria	*Serinus canarius. — <i>Briss.</i>	222
CHAFFINCH	Fringilla œolebs		224
BULLFINCH	Loxia pyrrhula	Pyrrhula vulgaris. — <i>Tem.</i>	226
GOLDFINCH	Fringilla carduelis	{ Carduelis communis. — <i>Cuv.</i> ; Carduelis elegans. — <i>Steph.</i>	227
STARLING	Sturnus vulgaris	Sturnus varius. — <i>Mey.</i>	229
RAVEN	Corvus corax		230
CROW	Corvus corone		233
ROOK	Corvus frugilegus		234
JACKDAW	Corvus monedula		236
MAGPIE	Corvus pica	Caudata Pica	237
CROSSBILL	Loxia		239
JAY	Corvus glandarius	{ Garrulus glandarius. — <i>Briss.</i> and <i>Cuv.</i>	240
CHOUGH	Corvus graculus	{ Pyrrhcorax graculus. — <i>Tem.</i>	241
SATIN BOWER BIRD		{ Ptilonorhynchus Holoseiceus. — <i>Kuhl.</i> Kitta. — <i>Lesson.</i> Grauculus — <i>Cuv.</i>	242
BIRD OF PARADISE	Paradisea apoda		244
NUTHATCH	Sitta Europæa		246
WRYNECK	Yunx torquilla		247
CREEPER	Certhia familiaris		247
HUMMING-BIRD	Trochilus colubris		248
HOOPOE	Upupa epops		249

SECTION IV.—Scansores, or Climbers.

English Name	Linnean Name.	Synonyms.	Page.
CUCKOO . . .	Cuculus canorus . . .	Cuculus hepaticus . . .	251
ROLLER . . .	Coracias garrula	255
KINGFISHER . . .	Alcedo ispada	257
WOODPECKER . . .	Picus viridus	259
TOUCAN . . .	Ramphastos tucanus	260
PARROT . . .	Psittacus erythacus	262
MACAW . . .	Psittacus aracanga . . .	Macrocerus aracanga— Viell.	262
PAROQUET . . .	{ Psittacus Nova Hol- landia . . .	Palæornis torquatus . . .	263
RING PAROQUET . . .	Psittacus Alexandri . . .	Palæornis Alexandri . . .	263
COCKATOO . . .	Psittacus galeritus . . .	Plectolophus galeritus . . .	264

SECTION V.—Gallinaceous Birds.

PEACOCK . . .	Pavo cristatus	265
TURKEY . . .	{ Meleagris Gallo- Pavo	267
GUINEA FOWL, JUNGLE FOWL, or BRUSH TUR- KEY . . .	Numida Meleagris	269
LYRE BIRD	Megapodius tumulus . . .	271
PHEASANT . . .	Phasianus Colchicus . . .	Menura superba . . .	274
PARTRIDGE . . .	Tetrao Perdix . . .	Perdix cinerea.—Lath. . .	277
RED-LEGGED PARTRIDGE . . .	Perdix Rufus	279
QUAIL . . .	Tetrao Coturnix . . .	{ Coturnix major.—Briss. Coturnix vulgaris.— Flem. Coturnix Euro- pæus.—Wils. Perdix Coturnix.—Lath.	282
AMERICAN QUAIL GROUSE, or MOOR FOWL . . .	Ortyx borealis . . .	Lagopus Scoticus.—Lath. . .	283
BLACK COCK . . .	Tetrao Scoticus . . .	*Bonasa Scotica.—Briss. . .	283
PTARMIGAN . . .	Tetrao Tetrix . . .	*Uriogallis minor.—Ray. . .	285
COCK . . .	Tetrao Lagopus . . .	Lagopus vulgaris.—Wils. . .	285
BANKIVA COCK . . .	Phasianus Gallus . . .	Tetrao rupestris.—Gmel. . .	286
JAGO COCK	Gallus domesticus.—Wils. . .	286
SPANISH COCK	Gallus Sonati . . .	288
BANTAM FOWL	289
DODO . . .	Didus ineptus	289
RINGDOVE, or CU- SHAT . . .	Columba palumbus	290
STOCKDOVE . . .	Columba Ænas	293
PIGEON . . .	Columba livia	295
TURTLEDOVE . . .	Columba Turtur	297

SECTION VI.—Grallatores, or Waders.

English Name.	Linnean Name	Synonymes.	Page.
OSTRICH . . .	Struthio Camelus . . .		299
CASSOWARY . . .	Struthio Casuarius . . .	{ Casuarius galeatus.— <i>Viel.</i> Casuarius Emeu . . . }	302
EMEU . . .	{ Struthio Novæ Hol- landæ . . . }	Dromaius ater.— <i>Viel.</i> . . .	304
CRANE . . .	Ardea Grus . . .	Grus cinerea.— <i>Cuv.</i> . . .	305
BALEARIC CRANE	Ardea pavonina . . .	{ Anthropoides pavonina.— <i>Viel.</i> . . . }	307
NUMIDIAN CRANE	Anthropoides Virgo . . .		307
STORK . . .	Ardea Ciconia . . .	Ciconia alba.— <i>Cuv.</i> . . .	308
ADJUTANT . . .	Ardea dubia . . .	Ciconia argala.— <i>Cuv.</i> . . .	310
HERON . . .	Ardea major . . .	Ardea Herodias . . .	311
BITTERN . . .	Ardea stellaris . . .	Botaurus stellaris.— <i>Briss</i>	313
SPOONBILL . . .	Platalea leucorodia . . .		315
CURLEW . . .	Scolopax arquata . . .	Numenius arquata.— <i>Cuv.</i>	316
REDHANK . . .	Scolopax calidris . . .	Totanus calidris.— <i>Bech.</i>	317
GODWIT . . .	Scolopax Egocephala	Limosa melanura.— <i>Tem.</i>	318
SNIPE . . .	Scolopax Gallinago . . .		318
WOODCOCK . . .	Scolopax rusticola . . .		320
LAPWING, OF PE- WIT . . .	Tringa vanellus . . .	Vanellus cristatus.— <i>Mey.</i>	321
RUFF and REEVE	Tringa pugnax . . .	Machetes Pugnax . . .	323
KNOT . . .	Tringa Canutus . . .	Tringa cinerea.— <i>Gmel.</i>	324
GREY PLOVER . . .	Tringa squatarola . . .		325
DOTTREL . . .	Charadrius Morinellus . . .		326
WATER HEN . . .	Fulica chloropus . . .	Gallinula chloropus . . .	327
CORNCRAKE, OR LAND RAIL . . .	Ortygometra crex . . .		328
COOT . . .	Fulica atra . . .		329

SECTION VII.—Palmipedes, or Web-footed Birds.

PELICAN . . .	Pelicanus onocrotalus . . .		330
CORMORANT . . .	Pelicanus Carbo . . .	{ Corvus aquaticus.— <i>Ray</i> Carbo Cormoranus.— <i>Mey.</i> Phalacrocorax Carbo.— <i>Briss.</i> . . . }	332
CRESTED CORMO- RANT	Phalacrocorax cristatus . . .	332
SOLOON GOOSE, OR GANNET . . .	Pelicanus Bassanus . . .	{ Pelicanus maculatus.— <i>Gmel.</i> —Anser bassanus. <i>Ray.</i> —Sula alba.— <i>Mey.</i> Sula Bassana.— <i>Bris.</i> . . . }	333
WILD SWAN . . .	Anas Cygnus . . .	Cygnus ferus.— <i>Ray.</i> . . .	335
TAME SWAN . . .	Anas olor . . .	Cygnus olor.— <i>Ray.</i> . . .	335
GOOSE . . .	Anas anser . . .	{ Anser palustris.— <i>Flem.</i> Anser ferus.— <i>Wils.</i> An- ser sylvestris.— <i>Briss.</i> . . . }	337
DUCK . . .	Anas Boschas . . .	Anas fera.— <i>Briss.</i> . . .	339
WIGEON . . .	Anas Penelope . . .	{ Mareca fistularis.— <i>Step.</i> Anatra Mangiana.— <i>Stor.</i> . . . }	341

English Name.	Linnaean Name.	Synonymes.	Page.
TEAL	Anas Crecca	Querquedula Crecca.—Step.	342
STORMY PETREL, OR MOTHER CAR- RY'S CHICK- ENS	} Procellaria pelagica	{ Thalassidroma pelagica.— Vigors }	} 343
AUK, OR NORTH ERN PENGUIN			

BOOK III.

INHABITANTS OF THE WATER.

SECTION I.—Cetacea, or Sea Mammalia.

WHITE WHALE	Delphinus leucas	{ Beluga leucas.—Gray.— Beluga artica.—Less. Delphinapterus Beluga. —Lacep. }	346
COMMON, OR GREENLAND WHALE	} Balæna mysticetus	349
SPERMACETI WHALE, OF CA- CHALOT			
DOLPHIN	Delphinus Delphis	354
PORPOISE	Delphinus Phocæna	Phocæna vulgaris	356
GRAMPUS	Delphinus orca	357
SEA UNICORN	Monodon monoceros	358

SECTION II.—Cartilaginous fishes.

SHARK	Squalus Carcharias	359
DOG-FISH	361
SKATE, OR MAID	Raja batis	362
TORPEDO	Raja Torpedo	Torpedo vulgaris	364
HAG-FISH	{ Gastrobranchus glutinosa }	365
PILOT FISH	Gasterosteus ductor	Naucrates ductor.—Cuv.	366
REMORA, OR SUCK- ING FISH	Echeneis Remora.—Tur.	367
SEA WOLF	Anarrhichas lupus	368
HORNED SILURE	369
FATHER LASHER	Cottus scorpius	Cottus bubalis	370
MONK FISH, OR ANGEL FISH	Squatina Angelus	370
SAW FISH	Pristis antiquorum	371
SWORD FISH	Xiphias gladius	372
FLYING SCORPION	373
BUTTERFLY FISH	374
BULL HEAD	Cottus	374
TOAD FISH	375

English Name.	Linnean Name.	Synonymes.	Page.
SUN FISH	Orthogoriscus	375
TRUMPET FISH	Centricus Scolopax	376
LUMP-SUCKER	Cylopterus lumpus	377
OCCELLATED-SUCKER	Cylopterus ocellatus	378
ANGLER	Lophius piscatorius	379
FOUR-HORNED TRUNK FISH	Ostracion quadricornis	380
GLOBE FISH	Tetraodon Hispidus	380
SEA HORSE	Hippocampus v. virostris	381
STURGEON	Accipenser Sturio	382
FLYING FISH	Exocoetus volitans	384
GURNARD	Trigla cuculus	382
JOHN DORY	Zeus faber	382
ZEUS OPAH	385
COD FISH	Gadus Morrhua	Morrhua vulgaris.—Cuv.	388
HADDOCK	Gadus Æglefinus	Morrhua Æglefinus.—Cuv.	389
WALTING	Gadus Merlangus	Merlangus vulgaris.—Cuv.	391
LING	Gadus molva	Lota molva.—Cuv. Asellus —Will. Molva vulgaris.—Flem.	391
HAKE	Gadus Merlucius	Merlucius vulgaris.—Cuv.	392
SMELT	Salmo eperlanus	Osmerus eperlanus.—Flem. Eperlanus Ronderlettii.—Will.	392
MACKEREL	Scomber Scomber	Scomber scombrus.—Cuv.	393
GAR FISH	Esox Belone	Scomber vulgaris.—Flem.	394
HERRING	Clupea Harengus	Belone vulgaris.—Cuv.	396
SPRAT	Clupea Sprattus	396
PILCHARD	Clupea pilchardus	Clupea Pilchardus.—Bloch.	397
ANCHOVY	Clupea encrasicolus	Engraulis encrasicolus.—Flem. Engraulis vulgaris.—Cuv.	398
WHITE BAIT	Clupea latu	Clupea alba.—Yarrell Clupea alosa.—Don.	399
TURBOT	Pleuronectes maximus	Rhombus maximus.—Cuv.	399
PLAICE	Pleuronectes platessa	Platessa vulgaris.—Flem.	400
FLOUNDER	Pleuronectes fesus	Platessa fesus.—Flem. Pleuronectes fluviatilis.—Will.	401
SOLE	Pleuronectes solea	Solea vulgaris.—Cuv.	402
SALMON	Salmo solar	403
SALMON PINK	405
SALMON TROUT	Salmo trutta	405
TROUT	Salmo fario	406
GRAYLING	Salmo thymallus	Thymallus vulgaris.—Cuv.	407
CHAR	Salmo umbla	Salmo alpinus.—Pen.	408
PIKE	Esox lucius	409
PERCH	Perca fluviatilis	411
POPE, OR RUFFE	412
CARP	Cyprinus carpio	413

English Name.	Linnean Name.	Synonymes	Page.
TENCH . . .	Cyprinus tinca . . .	Tinca vulgaris.—Cuv. . .	414
GOLD FISH . . .	Cyprinus auratus	415
GUDGEON . . .	Cyprinus gobio . . .	Gobio fluviatilis.—Will. . .	415
CHUB . . .	Cyprinus cephalus . . .	Leuciscus cephalus.—Flem. . .	416
BARBEL . . .	Cyprinus barbatus	417
DACE . . .	Cyprinus leuciscus . . .	Leuciscus vulgaris.—Cuv. . .	417
ROACH . . .	Cyprinus rutilus . . .	Leuciscus rutilus.—Cuv. . .	418
BLEAK . . .	Cyprinus alburnus	418
BREAM . . .	Cyprinus brama . . .	Abramis brama.—Cuv. . .	419
MINNOW . . .	Cyprinus phoxinus . . .	Leuciscus phoxinus.—Cuv. . .	420
BULLHEAD . . .	Cottus Gobio	420
ELECTRICAL EEL . . .	Gymnotus electricus	421
EEL . . .	Muraena Anguilla . . .	Anguilla vulgaris.—Thun. . .	423
CONGER EEL	423
LAMPREY . . .	Petromyzon marinus	424

BOOK IV.

REPTILES.

SECTION I.—Serpents, or Ophidian Reptiles.

VIPER, or ADDER . . .	Coluber Berus . . .	{ Vipera Berus.—Daud. . . } 427
HORNED VIPER . . .	Coluber cerastes . . .	{ Pelias Berus.—Merr. . . } 429
ASP . . .	Coluber Hajé . . .	Vipera cerastes 430
SNAKE . . .	Coluber natrix . . .	Coluber aspes 432
RATTLE SNAKE . . .	Crotalus horridus . . .	Natrix torquata.—Ray. . . 433
AMPHISBENA . . .	Amphisbæna fuliginosa 434
BOA . . .	Boa constrictor 435
ANACONDA . . .	Boa murina 436

SECTION II.—Batrachian Animals.

FROG . . .	Rana temporaria 436
TOAD . . .	Rana Bufo . . .	Bufo vulgaris.—Laur. . . 438

SECTION III.—Saurian Reptiles.

LIZARD . . .	Lacerta vivipara . . .	{ Lacerta agilis.—Briss. . . } 440
CHAMELEON . . .	Lacerta Chamælon . . .	{ Zootoca vivipara.—Wag. . . } 441
CROCODILE . . .	Lacerta Crocodilus . . .	Chamæleo vulgaris.—Cuv. . . 443
ALLIGATOR, or CAYMAN . . .	Lacerta Alligator . . .	Crocodilus vulgaris.—Cuv. . . 441

SECTION IV.—Chelonian Reptiles.

TORTOISE . . .	Testudo Græca 446
TURTLE . . .	Testudo Mydas . . .	Chelonia Mydas.—Briss. . . 447

BOOK V.

MOLLUSCOUS ANIMALS.

SECTION I.—Bivalves, or those having two Shells.

English Name.	Linnaean Name.	Synonymes.	Page.
PEARL OYSTER . . .	{ Mytilus Margaritifera . . . }	Avicula margaritifera.— <i>Lam.</i>	449
OYSTER . . .	Ostrea edulis	450
COCKLE . . .	Cardium edule . . .	Cardium fimbria . . .	451
MUSCLE . . .	Mytilus edulis	452

SECTION II.—Univalves.

ADMIRAL . . .	Conus ammiralis	452
SNIFE SHELL . . .	Murex haustellus	453
LINNET . . .	Patella vulgata	453
SNAIL . . .	Helix aspersa	454
GREY SLUG . . .	Limax cinereus	456
BLACK SLUG	456
CUTTLEFISH . . .	Sepia octopodia	457
NAUTILUS	457

BOOK VI.

ARTICULATED ANIMALS.

SECTION I.—Annelida, or ringed Animals.

EARTHWORMS . . .	Lumbricus terrestris	458
LEECH . . .	Hirudo medicinalis . . .	Sanguisuga officinalis . . .	459

SECTION II.—Crustacea.

LOBSTER . . .	Cancer gammarus . . .	Astacus marinus.— <i>Leach.</i>	461
CRAYFISH . . .	Cancer astacus . . .	{ Astacus fluviatilis.— <i>Des.</i> . . . Potamobius.— <i>Leach.</i> . . . }	462
CRAB . . .	Cancer Pagurus	462
LANE CRAB	463
SOLDIER CRAB	Pagurus Miles.— <i>Oliv.</i>	464
SHRIMP . . .	Cancer crangon . . .	Crangon vulgaris.— <i>Fab.</i>	465
PRAWN . . .	Cancer serrator . . .	Palæmon serratus.— <i>Pen.</i>	465

SECTION III.—Arachnida.

SPIDER . . .	Aranea domestica	467
TARANTULA . . .	Phalangium hirsutum . . .	Lycosa tarantula . . .	468
CHEESE MITE . . .	Acarus domesticus	469

SECTION IV.—*INSECTS.*

ORDER I.—COLEOPTERA, OR BEETLES.

English Name.	Linnean Name.	Synonymes.	Page.
COCK CHAFER . . .	Scarabæus Melolontha	Melolontha vulgaris . . .	472
DOR BEETLE . . .	Scarabæus stercorarius	Geotrupes stercorarius . . .	473
STAG BEETLE . . .	Lucanus Cervus . . .		474
ELEPHANT BEETLE . . .	Scarabæus elephantis . . .	Dynastes elephas . . .	475
MUSK BEETLE, OR GOAT CHAFER . . .	Cerambyx moschatus . . .		476
GROUND BEETLE . . .	Carabus clathratus . . .		476
GLOW WORM . . .	Lampyris noctiluca . . .		477
DEATH WATCH . . .	Ptinus pulsator . . .	Anobium tessellatum . . .	478
SPANISH FLY . . .	Cantharis vesicatoria . . .		478
LADY BIRD . . .	Coccinella septempunctata . . .		479

ORDER II.—ORTHOPTERA.

EARWIG . . .	Forficula auricularia . . .		479
WALKING LEAF . . .	Mantis sp. . .	Phyllium siccifolium.— <i>Ill.</i>	480
GRASSHOPPER . . .	Gryllus viridissimus . . .	Tetrix viridissima.— <i>Lat.</i>	481
LOCUST . . .	Gryllus migratorius . . .	Acrydium migratorium.— <i>Geoff.</i>	482
MOLE CRICKET . . .	Gryllus Gryllotalpa . . .	Gryllotalpa vulgaris.— <i>Lat.</i>	484
CRICKET . . .	Gryllus domesticus . . .		485

ORDER III.—HEMIPTERA.

LANTHERN FLY . . .	Fulgora lanternaria . . .		486
COCHINEAL INSECT . . .	Coccus cacti . . .		486
GREEN FLY . . .	Aphis rosæ . . .		488

ORDER IV.—NEUROPTERA.

ANT LION . . .	{ Myrmeleon formicarium . . . }		489
DRAGON FLY . . .	Libellula grandis . . .	*Æshna.— <i>Fab.</i>	491

ORDER V.—HYMENOPTERA.

BEE . . .	Apis mellifica . . .		492
WASP . . .	Vespa vulgaris . . .		494
ICHNEUMON . . .	{ Ichneumon mani- festator . . . }		495
ANT . . .	Formica rufa . . .		496

ORDER VI.—LEPIDOPTERA, MOTHS AND BUTTERFLIES.

MOTH, with its CHRYSLIS and CATERPILLARS . . .			498
--	--	--	-----

English Name.	Linnean Name.	Synonymes	Page.
TORTOISE-SHELL BUTTERFLY . . .	Papilio Vanessa	Vanessa urtica	499
CABBAGE BUTTERFLY . . .	Papilio Brassicæ	{ Pieris Brassicæ.— <i>Lat.</i> Pontia Brassicæ.— <i>Fab.</i>	500
MAGPIE MOTH . . .	Phalæna grossulari- ata	{ Abraxas grossulariata.— <i>Leach.</i>	
WINTER MOTH . . .	Phalæna brumata	Hibernia brumata.— <i>Lat.</i>	502
BLACK ARCHES . . .	Bombyx monacha	{ Liparis monacha.— <i>Och.</i> Psilura monacha.— <i>Steph.</i>	503
SILKWORM . . .	Bombyx mori		
CLOTHES MOTH . . .	Tinea vestianella		504
BOOKWORM . . .	{ Crambus pingui- nella		505

ORDER VII.—DIPTERA.

HOUSE FLY . . .	Musca domestica		507
GNAT . . .	Culex pipiens		507

ORDER VIII.—SUCTORIA.

FLEA . . .	Pulex irritans		508
------------	----------------	--	-----

BOOK VII.

RADIATA.

STAR FISH . . .	Asterias glacialis		509
SEA URCHIN . . .	Echinus		510
ZOOPHYTES . . .			511
CORALS . . .			512
POLYPPES (fresh water) . . .			519
— (sea) . . .			521

APPENDIX.

FABULOUS ANIMALS.

SPHINX . . .	523	KRAKEN . . .	530
DRAGON . . .	525	DOLPHIN . . .	530
WIVERN, or WOLVERINE . . .	526	UNICORN . . .	531
COCKATRICE, or BASILISK . . .	527	PEGASUS . . .	532
GRIFFIN . . .	528	CENTAUR . . .	532
PHENIX . . .	529	SATYR . . .	533
MERMAID, or SIREN . . .	529		

The Entertaining Naturalist.


BOOK E.

I. QUADRUPEDS, OR FOUR-FOOTED BEASTS.

§ I. Carnivorous, or Flesh-eating Animals.



590
Lou



THE LION. (*Felis Leo.*)

“What if the Lion in his rage I meet?
Oft in the dust I view his printed feet:
And, fearful oft when Day's declining light
Yields her pale empire to the mourner Night,
By hunger roused, he scours the groaning plain,
Gaunt wolves and sullen tigers in his train:
Before them Death, with shrieks, directs their way,
Fills the wild yell, and leads them to their prey.”—COLLINS.

THE LION is called the king of beasts, not only from his grave and majestic appearance, but from his prodigious

strength. Zoologists describe him as an animal of the cat kind, distinguished from the other species of the genus by the uniformity of his colour, the mane which decorates the male, and a tuft of hair at the tip of the tail, which conceals a small prickle or claw.

Lions were formerly found in every part of the old world; but they are now confined to Africa, and some parts of Asia. The African Lion stands four or five feet high, and his body is from seven to nine feet long. The mane is thick, and somewhat curly; and the colour varies in different parts of Africa, but it is generally of a clear dark brown, deepening in some cases almost into black. The Asiatic Lions are much smaller than those of Africa, and their colour paler. The Bengal Lion is of a light brown, with a long flowing mane; the Persian Lion is of a sort of cream-colour, with a short thick mane; and the Lion of Guzerat is of a reddish brown, without any mane.

All the varieties agree in their habits; they lie hid in jungles in the long grass, and when roused either walk quietly and majestically away, or turn and look steadily at their pursuers. Their roar is terrific; and in a wild state, the animal generally roars with his mouth close to the ground, which produces a low rumbling noise, like that of an earthquake. The effect is described by those who have heard it, as making the stoutest heart quail; and the feebler animals, when they hear it, fly in dismay, often in their terror falling in the way of their enemy, instead of avoiding him. Serpents, and some of the larger animals, will, however, fight with Lions, and occasionally kill them; and Lions, when pursued by man, are sometimes hunted with dogs, but are oftener shot, or speared. Those which are exhibited in menageries have generally been caught in pits. The pit is dug where traces have been discovered of a Lion's path; and it is then covered with sticks and turf. He is deceived by the appearance of solidity presented by the turf, and attempts to walk over it; but the moment he sets his foot upon the covering of the trap, it breaks beneath his weight, and he falls into the pit. He is then kept without

food for several days, shaking the ground with his roaring, and fatiguing himself by vainly attempting to escape; till, at last, he becomes exhausted, and so tame as to permit his captors to put ropes round him, and drag him out. He is then put into a cage, and removed in a kind of waggon, wherever his captors may wish to take him.

The generosity of the Lion has been much extolled; but the tales related of it appear to have had no other foundation than the fact, that, like many other beasts, when gorged with food, he will not attack a man. As the Lion belongs to the cat tribe, his eyes are incapable of bearing a strong light; it is therefore generally in the night that he prowls about for prey, and when the sun shines in his face, he becomes confused and almost blinded. Lion hunters are aware of this fact. In the day-time they always consider themselves safe, so long as they have the sun on their backs. In the night a fire has nearly the same effect; and travellers in Africa and the deserts of Arabia can always protect themselves from Lions and Tigers, by making a large fire near their sleeping-place. The strength of the African species is so great that he has been known to carry away a young heifer, and leap a ditch with it in his mouth. The power that man may acquire over this animal has been often shown in the exhibitions of Van Amburgh, Carter, and others; but the attachment which Lions sometimes form for their keepers, was never more strongly exemplified than in the following anecdote.

M. Felix, the keeper of the animals in Paris, some years ago brought two Lions, a male and female, to the national menagerie. About the beginning of the following June he was taken ill, and could no longer attend them; and another person was under the necessity of performing this duty. The male, sad and solitary, remained from that moment constantly seated at the end of his cage, and refused to take food from the stranger, whose presence was hateful to him, and whom he often menaced by bellowing. The company even of the female seemed now to displease him, and he paid no attention to her. The uneasiness of the animal afforded a belief that

he was really ill; but no one dared to approach him. At length Felix recovered, and, with an intention to surprise the Lion, crawled softly to the cage, and showed his face between the bars: the Lion, in a moment, made a bound, leaped against the bars, patted him with his paws, licked his hands and face, and trembled with pleasure. The female also ran to him; but the Lion drove her back, and seemed angry; and, fearful lest she should snatch any favours from Felix, a quarrel was about to take place; but Felix entered the cage to pacify them. He caressed them by turns; and was afterwards frequently seen between them. He had so great a command over these animals, that whenever he wished them to separate and retire to their cages, he had only to give the order: when he wished them to lie down, and show strangers their paws or throats, they would throw themselves on their backs on the least sign, hold up their paws one after another, open their jaws, and, as a recompense, obtain the favour of licking his hand.

The Lion, like all animals of the cat kind, does not devour his prey the moment he has seized it. When those in cages are fed, they generally hide their food under them for a minute or two before they eat it. Thus an instance is known of a man, who was struck down by a Lion, having time to draw his hunting-knife and stab the ferocious beast, who was growling over him, to the heart, before it had seriously injured him. The Lion also resembles a cat in his mode of stealing after, and watching his prey, a long time before seizing it.


Dr. Sparrman mentions a singular instance of the animal's habits in this respect. A Hottentot perceiving that he was followed by a Lion, and concluding that the creature only waited the approach of night to make him his prey, began to consider what was the best mode of providing for his safety, and at length adopted the following:—Observing a piece of broken ground with a precipitate descent on one side, he sat down by the edge of it; and found, to his great joy, that the Lion also made a halt, and kept at a distance behind him. As soon as it grew dark the man, sliding gently forward, let

himself down a little below the edge of the steep; and held up his cloak and hat on his stick, at the same time gently moving them backward and forward. The Lion, after a while, came creeping towards the object; and, mistaking the cloak for the man himself, made a spring at it, and fell headlong down the precipice.

Lions have been sometimes known to attain a great age, as Pompey, a large male Lion that died, in 1760, in the Tower of London, which was upwards of seventy years old. The usual period, however, seldom exceeds twenty years. The Lion is generally represented as the companion of Britannia, as a national symbol of strength, courage, and generosity. In ancient gems, paintings, and statuary, his skin is the attribute of Hercules. In Scriptural compositions, he is painted at the side of the evangelist St. Mark; and holds the fifth place among the signs of the zodiac, answering to the months of July and August.

In the various sculptured Lions discovered by Mr. Layard at Nineveh in 1848, the claw in the Lion's tail is distinctly marked, and is represented as being of large size. It is, however, really a very small, dark, horny prickle at the tip of the fleshy part of the tail, and entirely hidden by the hair.





THE LIONESS AND CUBS.

THE LIONESS is in all her dimensions about one-third less than the male, and has no mane. She has generally from two to four cubs at a time, which are born blind, like kittens, which they greatly resemble, though they are as large as a pug-dog, when born. When quite young, they are striped and spotted, but these marks soon disappear; they also at first mew like a cat, and do not begin to roar till they are about eighteen months old. About the same time, the mane begins to appear on the males, and soon after the tuft of hair on the tail, though the animal is generally five or six years before it attains its full size.

The Lioness, though naturally less strong, less courageous, and less mischievous than the Lion, becomes terrible as soon as she has young ones to provide for. The ferocity of her disposition then appears with tenfold

vigour; and woe be to the wretched intruder, whether man or beast, who should unwarily approach the precincts of her sanctuary. She makes incursions for food for her young with even more intrepidity than the Lion himself; throws herself indiscriminately among men and other animals; destroys without distinction; loads herself with the spoil, and brings it home reeking to her cubs. She usually brings forth her young in the most retired and inaccessible places; and when she fears the discovery of her retreat, often hides her track, by running back over the ground, or by brushing it out with her tail. She sometimes also, when her apprehensions are great, transports her young from one place to another, like a cat; and if obstructed, defends them with determined courage, and fights to the last.

Mr. Fennel, in his *History of Quadrupeds*, relates an interesting anecdote of a Lioness kept at the Tower in 1773. This creature had become "greatly attached to a little dog, which was her constant companion. When the Lioness was about to whelp, the dog was removed; but shortly after her accouchement had taken place, the dog contrived to enter the den, and approached the Lioness with his usual fondness. She, alarmed for her cubs, immediately seized him, and seemed about to kill him; but, as if suddenly recollecting their former friendship, she carried him to the door of her den, and allowed him to escape unhurt." Mr. Fennel also tells us, that the first Lioness ever brought to England, died in the Tower in 1773, after having attained a great age.

Another Lioness, which was kept at the Tower in 1806, became extremely attached to a little dog, and whenever he attempted to pass through the bars of the den, would draw him back by the hinder parts, and place her paw gently upon his body, as if entreating him not to leave her.



THE TIGER.—(*Felis Tigris*.)


THOUGH very inferior to the lion in majesty of appearance and deportment, this ferocious animal nearly equals him in size and strength. The Tiger is another of the feline species, and may be compared to an enormous cat, the whiskers and the tail being exactly similar; and both the Tiger and the lion resemble the cat in the form of their feet, and the power they possess of drawing in their claws. The Tiger, however, bears the strongest resemblance, and when pleased, purrs and curves up his back as he rubs himself against the nearest object. When enraged, he growls rather than roars; and springs up to a great height before he pounces on his prey.

The Tiger has a smaller and rounder head than the lion; he has no mane; his tail is without any tuft at the extremity, and his body much more slender and flexible. His colour is yellowish on the back and sides, becoming white beneath, with numerous lines of a very dark rich brown, or glossy black, sloping from the centre of the back down the sides, and over the head, and continued down the tail in the form of rings. Tigers are only found

wild in Asia; but they are very abundant, and very destructive in the East Indies, as from their enormous strength they can carry off a bullock with the greatest ease.

The attack of one of these animals upon Mr. Monro, son of Sir Hector Monro, was attended with the most tragical consequences. "We went," says an eye-witness, "on shore on Sangar Island, to shoot deer, of which we saw innumerable tracks, as well as of Tigers. We continued our diversion till near three o'clock, when sitting down by the side of a jungle to refresh ourselves, a roar like thunder was heard, and an immense Tiger seized our unfortunate friend, and rushed again into the jungle, dragging him through the thickest bushes and trees, everything giving way to his monstrous strength. All we could do was to fire on the Tiger; and our shots took effect, as in a few moments our unfortunate friend came up to us bathed in blood. Every medical assistance was vain, and he expired in the space of twenty-four hours, having received such deep wounds from the teeth and claws of the animal as rendered his recovery hopeless. A large fire, consisting of ten or twelve whole trees, was blazing near us at the time this accident took place; and ten or more of the natives were with us. The human mind can scarcely form any idea of this scene of horror."

Tiger-hunting, though very dangerous, is a very favourite sport in India. The hunters are mounted in carriages called howdahs, on the backs of elephants, well armed. The first indication is generally given by the elephants, who scent their enemy at some distance, and commencing a peculiar kind of snorting, become greatly agitated. As soon as the motion of the Tiger through the jungle is perceived, the nearest elephant is halted, and the hunter fires instantly. Should the Tiger be wounded, he will, in all probability, spring up with a hideous roar, and rush at the nearest elephant, his mouth open, his tail erect, or lashing his sides, and his whole fur bristled up. Sometimes, however, he endeavours to sneak away, artfully diminishing his size by drawing in his breath and creeping along the ground, and often with such success as to enable him to escape to ravines where it would be madness to attempt pursuit.



THE LEOPARD (*Felis Leopardus*)

Differs from the tiger in being smaller, and in having the skin spotted instead of striped. His length from nose to tail is about four feet, the colour of the body is a more lively yellow, and the spots of his skin are smaller and closer than those of the panther; they are composed of four or five black dots arranged in a circle, and not imperfectly representing the print left by the animal's foot upon the sand. There are three varieties of the Leopard; viz., the Panther, the Ocelot, and the Ounce.

Like all animals of the cat tribe, Leopards are a compound of ferocity and cunning; they prey upon the smaller animals, such as antelopes, sheep, and monkeys; and are enabled to secure their food with great success, from the extraordinary flexibility of their bodies. Kolben informs us, that in the year 1708, two of these animals, a male and female, with three young ones, broke

into a sheepfold at the Cape of Good Hope. They killed nearly a hundred sheep, and regaled themselves with the blood, after which, they tore a carcass into three pieces, one of which they gave to each of their offspring; they then took each a whole sheep, and, thus laden, began to retire; but having been observed, they were waylaid on their return, and the female and young ones killed, while the male effected his escape. They appear afraid of man, and never attack him unless driven by hunger, when they spring upon him from behind. The Leopard is sometimes called the tree-tiger.



THE PANTHER (*Felis Pardus*)


Is in shape somewhat like the leopard, of which he is generally considered only a variety. His hair is short, sleek, and glossy, and of a bright yellow, beautifully marked with round, annular black spots. He is about the size of a large mastiff-dog, with legs not quite so long; his voice is strong and hoarse, and he growls continually.

He is fierce, greedy of blood, swift in the chase, and catches his prey by leaping from some lurking-place, where he squats himself in the manner and shape of a cat, which he resembles in many points. Like all the tribe, his tongue, in licking, grates like a file.

Some years ago, Mrs. Bowditch brought a tame Panther over with her from Africa. This animal was called Sai. One day, at Cape Coast Castle, he found the servant appointed to attend on him sitting asleep, resting his back against a door; Sai instantly lifted up his paw, and gave the sleeper a tap on the side of the cheek, which knocked him over, and when the man awaked, he found Sai wagging his tail, and seeming to enjoy the fun. Another day, when a woman was scrubbing the floor, he jumped on her back; and when the woman screamed with fright, he sprang off, and began rolling over and over like a kitten. When put on board ship, he was first confined in a cage; and the greatest pleasure he had was when Mrs. Bowditch gave him a little twisted cup or cornet of stiff paper with some lavender-water in it, and with this he was so delighted, that he would roll himself over and over, and rub his paws against his face. At first he used to put his claws out when he attempted to snatch anything; but as Mrs. Bowditch would never give him any lavender-water when this was the case, he soon learnt to keep his claws in. This poor Panther died soon after it reached England.

THE OUNCE

DIFFERS from the leopard principally in the shape of its spots, which are sometimes scalloped and sometimes ring-shaped, and are very irregularly disposed. This animal is a native of the Old World. The *ocelot*, or tiger-cat, is, on the contrary, only found in South America and Mexico, and is described by Buffon as the most beautiful of all the animals of the cat tribe. Its colour is grey, tinged with fawn, and it is marked with long compressed rings of black, each enclosing a spot of yellow. The movements of this animal are peculiarly graceful, and it is easily tamed.



THE CHEETAH, OR HUNTING LEOPARD.


(*Felis Jubata.*)

THE Hunting Leopard seems to form the connecting link between the cat and the dog-tribes; as it has the long tail and flexible body of the cat, with the sharp nose and elongated limbs of the dog. Its claws also resemble those of a dog, and cannot be drawn into the foot, as they can in all animals of the cat kind. The Cheetah is easily tamed, and Cuvier describes one which was accustomed to go at large in a park, and associated with the children and domestic animals, purring like a cat when pleased, and mewling when he wished to call attention to his wants. In the East the Cheetah is used in hunting, and is carried in a carriage, or chained on a pad behind the saddle of a horseman, with a hood over his eyes: when a herd of antelopes is found, the hood is taken off the Cheetah, who is let loose, and as soon as he sees the antelopes, steals cautiously along, till he comes within reach, when he springs suddenly upon them; making several bounds with the greatest rapidity, till he has killed his victim, when he begins instantly to suck its blood. The keeper then approaches, and throwing the Cheetah some pieces of raw meat, contrives to hoodwink and chain him again to his pad behind the saddle, on which he crouches like a dog. If the Cheetah is not successful in catching an antelope before the herd takes flight, he never pursues them, but returns to his keeper with a discontented and sullen air.



THE JAGUAR. (*Felis Onca.*)

THE Jaguar is a native of the New World, and is sometimes called the American Tiger. He is generally larger and stronger than the leopard, which he resembles in colour; but the black ring-like marks have always a spot in the centre, which is not the case with those of the leopard. The tail is also much shorter, and the head larger and rounder. The Jaguar has great strength, and will kill a horse or an antelope, and carry it off. He is, however, a cowardly animal, always springing upon his prey from behind, and attacking in preference the hindmost of a herd. He fastens upon its neck, placing one paw upon the head, which he twists round with the other, and thus instantly deprives it of life. His principal haunt is the long grass on the banks of a river, where he subsists chiefly on turtles; turning them on their backs, and then insinuating his paw between the shells so as to scoop out the flesh. No instance is yet known of a Jaguar being tamed.



THE PUMA. (*Felis concolor.*)

THE Puma, or American Lion, is smaller than the jaguar, and has a shrill hissing cry, very different from that of other animals of the cat kind. The fur is of a silvery fawn-colour, nearly white below, but becoming black at the head; the animal has no mane, and its tail is without any tuft at the tip. The cubs are spotted when young. The habits of the Puma are somewhat peculiar; when attacked, he climbs the nearest tree for safety, and there is generally shot by his hunters. When hunted with dogs, however, and cut off from all retreat, he stands at bay and fights furiously. The flesh is eaten by the Indians, and is said to be much prized by them. The Puma flies from the sight of man, and seldom attacks any animal larger than a sheep; but when he can surprise a flock of sheep, he kills as many as he can, only sucking the blood of each. He never devours the whole of his prey at once, carefully covering with leaves what he cannot eat; but if these should be removed, he will not touch the food again. It was formerly supposed that the Puma could not be tamed; but this is incorrect, as the late Edmund Kean, the tragedian, had one which followed him about like a dog, and was often permitted to come, at perfect liberty, into the drawing-room when it was full of company.



THE CARACAL LYNX. (*Felis Caracal.*)

THERE are several kinds of Lynx, but the Caracal is generally supposed to be the Lynx of the ancients, which was so celebrated for the keenness of its sight. The name of Caracal is derived from two Turkish words, signifying black-ears, and the animal is, in fact, remarkable for the blackness of the tips of its ears. He is somewhat larger and stronger than the fox; his body of a reddish brown, becoming white below, and the tail rather short, being only about eight or nine inches in length. This animal is said to follow the lion's footsteps, and to feed on what the lion leaves, like the jackal. The Caracal is both irritable and sulky in confinement, and is very seldom tamed; indeed, on the slightest irritation, it expresses its anger by a sort of snarl, like what is called swearing in a cat, but much louder, and sometimes ending in a scream.

When left to its own resources for support, it preys upon hares, rabbits, and birds; and will pursue the latter, of which it is immoderately fond, with remarkable activity, to the tops of the tallest trees.



THE CAT. (*Felis Catus*, var. *domestica*.)

“ Grimalkin, to domestic vermin sworn
 An everlasting foe, with watchful eye
 Lies nightly brooding o'er a chinkey gap,
 Protending her fell claws, to thoughtless mice
 Sure ruin.”

JOHN PHILIPS.

It was formerly supposed that the common domestic Cat was nothing more than the wild Cat of the woods, rendered tame by education. This opinion is, however, now doubted, on the ground that the tail of the wild Cat is thick and bushy, like that of a fox, while that of the domestic Cat tapers to the point. The Cat of the Egyptians, of which so many mummies have been found, differed still more in this respect, as its tail was long and slender, ending in a kind of tuft. There are four or five distinct varieties of the domestic Cat: the tabby, the tortoise-shell, the Chartreuse, and the Angora. Of these the tabby bears most resemblance to the wild Cat, and the black Cats are from this breed: the tortoise-shell is said to have been brought from Spain, the females of this race being generally of a pure tortoise-shell, and the males buff, with stripes of a darker hue. All the white, and whitish Cats are descended from the Chartreuse breed; they have all a blue tinge in their fur, and reddish eyelids: the tailless cats of Cornwall and the Isle of Man belong to this race. The Angoras are quite distinct, and are well known by their long silky hair. Cats are fond of warmth, and are generally affected by changes in

the weather. They are very affectionate, purring at the sight of those who are kind to them; and will curve up their backs and rub themselves against a door when it is opened for them, as if to thank the kind friend who has done them this service, before they take advantage of it. The female Cat has generally five or six kittens at a time, which she carries about in her mouth, and hides, when she thinks them in danger. When a Cat is enraged, its hair stands erect, and its tail swells to an enormous size. Cats fight savagely, and often tear the skin off each other's necks: when two are about to fight, they stand for some time looking at each other, growling, and then dart at each other with the greatest fury, yelling with rage.

Most cats are good mousers, and some bring everything they kill to their master or mistress, displaying their mice and rats with as much pride as a sportsman would his game. They are very fond of catmint and valerian, rolling themselves in a kind of ecstasy when they smell the latter plant. They are very cleanly, often sitting stroking their faces with their paws, as if washing themselves.

In the eye of the Cat, the pupil is perpendicularly oval, extending from above downwards, and when contracted appears like a straight line. This conformation is suited to the habits of these animals, for they are not content with prowling along the ground, but occasionally spring to great heights, their heads being directed upwards, and their eyes placed in front and more nearly parallel.





[Chasseur and Cuba Bloodhounds.]

THE BLOODHOUND. (*Canis sanguinarius.*)

“—Conscious of the recent stains, his heart
Beats quick ; his snuffling nose, his active tail,
Attest his joy : then with deep opening mouth,
That makes the welkin tremble, he proclaims
Th’ audacious felon.—”

THE BLOODHOUND is taller than the old English hound, most beautifully formed, and superior to every other kind in activity, speed, and sagacity. It is commonly of a reddish or brown colour, with long ears. It seldom barks, except in the chase ; and never leaves its game until it has caught and killed it.

Bloodhounds were formerly used in certain districts lying between England and Scotland, which were much infested by robbers and murderers; and a tax was laid upon the inhabitants for keeping and maintaining a certain number of them. But as the arm of justice is now extended over every part of the country, and there is no secret recesses where villany may lie concealed, these services are no longer necessary. In former times these Dogs were used to hunt renegade negroes and others in the Spanish West Indies, and many surprising anecdotes are told of their wonderful sagacity and power of scent.

In Dallas' "History of the Maroons," an anecdote is given of the extent of their accomplishments in this way, which seems truly marvellous. A ship, attached to a fleet under convoy to England, was manned chiefly by Spanish sailors, who, as they passed Cuba, took the opportunity of running the vessel on shore, when they murdered the officers, and other Englishmen on board, and carried off all the available plunder into the mountains of the interior. The place was wild and unfrequented, and they fully expected to elude all pursuit. The moment, however, the news reached Havanna, a detachment of twelve chasseurs, with their Dogs, was sent off. The result was, that in a few days the whole of the murderers were brought in and executed, not a man having been injured by the Dogs in the capture.

The old English Hound, the original stock of this island, and used by the ancient Britons in the chase, is a most valuable Dog; though the breed has been gradually declining, and the size studiously diminished by a mixture of other kinds, in order to increase their speed. It seems to have been accurately described by Shakspeare in the following lines:—

“ My hounds are bred out of the Spartan kind,
 So flew'd, so sanded; and their heads are hung
 With ears that sweep away the morning dew;
 Crook-kneed and dew-lapp'd, like Thessalian bulls;
 Slow in pursuit; but match'd in mouth like bells;
 Each under each.”



THE FOXHOUND.

THIS most valuable of all the Dogs of the chase, is smaller than the staghound, its average height being from twenty to twenty-two inches. No country in Europe can boast of Foxhounds equal in fleetness, strength, and perseverance to those of Britain, where the utmost attention is paid to their breeding, education, and food. The climate also seems congenial to their nature, for when taken to France or Spain, and other southern countries of Europe, they quickly degenerate, and lose all the admirable qualities they possess in this country.


Our predilection for fox-hunting appears to have descended from our forefathers, and to have gone on increasing in ardour. Certainly no other country can boast of such splendid establishments for this valuable breed: the Duke of Richmond's Kennel at Goodwood cost no less than £19,000.





THE POINTER (*Canis familiaris avicularis*)

Is about the size of a bulldog, and spotted like a spaniel. In disposition it is docile, and when trained, is of the greatest service to the sportsman who delights in shooting. It is astonishing to see to what a degree of obedience these animals may be brought. Their sight is equally acute with their scent, and they are enabled to perceive at a distance the smallest sign from their master. So admirably have they been trained, that their acquired propensities seem as inherent as a natural instinct, and appear to be transmitted from parent to progeny. When they scent their game, they fix themselves like statues, in the very attitude in which they happen to be at the moment. If one of their fore feet is not on the ground when they first scent, it remains suspended, lest, by putting it to the ground, the game might be too soon alarmed by the noise. In this position they remain, until the sportsman comes near enough, and is prepared to take his shot; when he gives the word, and the Dog immediately springs the game. This attitude has often been selected by the artist.



THE MASTIFF (*Canis molossus*)

Is the largest of the whole species: he is a strong and fierce animal, with short pendent ears and a large head, large and thick lips hanging on each side, and a noble countenance; he is a faithful guardian, and a powerful defender of the house.

A curious account is given, by Stow, of an engagement between three Mastiffs and a lion, in the presence of James the First. "One of the Dogs, being put into the den, was soon disabled by the lion, which took him by the head and neck, and dragged him about: another Dog was then let loose, and served in the same manner: but the third, being put in, immediately seized the lion by the lip, and held him for a considerable time; till, being severely torn by his claws, the Dog was obliged to quit his hold; and the lion, greatly exhausted in the conflict, refused to renew the engagement; but, taking a sudden leap over the Dogs, fled into the interior part of the den. Two of the Dogs soon died of their wounds; the last survived, and was taken great care of by the king's son, who said, 'He that had fought with the king of the beasts should never after fight with any inferior creature.'"

The following anecdote will show that the Mastiff, conscious of its superior strength, knows how to chastise the impertinence of an inferior:—A large Dog of this kind, belonging to a gentleman near Newcastle, being frequently molested by a mongrel, and teased by its continual barking, at last took it up in his mouth, by the back, and, with great composure, dropped it over the quay into the river, without doing any further injury to an enemy so much its inferior.



THE BULLDOG (*Canis domesticus molossus*)

Is much less than the mastiff, but the fiercest of all the Dog kind, and is probably the most courageous creature in the world. His short neck adds to his strength. Those of a brindled colour are accounted the best of the kind: they will run at and seize the fiercest bull without barking, making directly at his head, sometimes catch hold of his nose, pin the animal to the ground, and make him roar in a most tremendous manner, nor can they without difficulty be made to quit their hold. Whenever a Bulldog attacks in any of the extremities of the body, it is invariably considered a mark of his degeneracy from the original purity of blood.

Some years since, at a bull-baiting in the north of England, when this barbarous custom was very common, a young man, confident of the spirit of his Dog, laid a wager that he would, at separate times, cut off all the animal's feet, and that he would continue to attack the bull after each amputation. The experiment was tried, and the brutal wretch won his wager.



THE SHEPHERD'S DOG. (*Canis familiaris.*)

To no animal is mankind so much indebted for its services and affection as to the Dog. Among all the various orders of brute creatures, none have hitherto been found so entirely adapted to our use, and even to our protection, as this. There are many countries, both of the old and new continent, in which, if man were deprived of this faithful ally, he would unsuccessfully resist the foes that surround him, seeking opportunities to encroach upon his property, destroy his labour, and attack his person. His own vigilance, in many situations, could not secure him, on the one hand, against their rapacity, nor, on the other, against their speed. The Dog, more tractable than any other animal, conforms himself to the movements and habits of his master. His diligence, his ardour, and his obedience are inexhaustible; and his disposition is so friendly, that, unlike every other animal, he seems to remember only the benefits he receives: he soon forgets our blows; and instead of discovering resentment while we chastise him, exposes himself to torture, and even licks the hand from which it proceeds.


Dogs, even of the dullest kind, seek the company of other animals; and by instinct take to the care of flocks and herds.

The *Shepherd's Dog* may be considered the primitive stock, from whence those of Siberia, Lapland, Iceland, the Cape of Good Hope, Madagascar, and indeed all others,

are derived. He makes the stem of the genealogical tree, which has branched out into every part of the world. This animal still continues nearly in its original state among the poor in temperate climates: being transported into the colder regions, it becomes smaller, and covered with a shaggy coat. Whatever differences there may be among the Dogs of these cold countries, they are not very considerable, as they all have straight ears, long and thick hair, a savage aspect, and do not bark either so often or so loud as Dogs of the more cultivated kind. The Shepherd's Dog, transported into temperate climates, and among people entirely civilized, such as into England, France, and Germany, will be divested of his savage air, his pricked ears, his rough, long, and thick hair, though he will still retain his large skull, abundant brain, and consequent great sagacity.

Many interesting anecdotes are told of the shepherd's tyke or colley, as this kind of Dog is frequently called, particularly of its sagacity in rescuing sheep from snow-drifts. When sheep are missing in a snow-storm, as is frequently the case in Scotland and the North of England, the shepherd arms himself with a spade, and watching the motions of his faithful Dog, digs into the snow wherever the Dog begins to scratch it away, and is thus sure to find his lost sheep.

This valuable boon to the shepherd is the least voracious of his genus, and endures fatigue and hunger with patience.





THE SPANIEL. (*Canis avicularius.*)

OF this elegant animal, said to be of Spanish extraction, there are several varieties in this country; but it is more than probable that the English Spaniel, the most common and useful breed, is indigenous. It has received from nature a very keen smell, good understanding, and uncommon docility, and is employed in setting for partridges, pheasants, quails, &c. His steadiness in the field, his caution in approaching game, his patience in keeping the bird at bay till the fowler discharges his piece, are objects worthy of admiration. Many sportsmen prefer him to the pointer; and if water is plentiful he is more useful, for his feet are much better defended against the sharp cutting of the heath than those of the pointer, as he has a great deal of hair growing between the toes and round the ball of the feet, of which the pointer is almost destitute. He also ranges much faster, and can endure more fatigue.

“When milder autumn summer’s heat succeeds,
 And in the new-shorn field the partridge feeds,
 Before his lord the ready spaniel bounds;
 Panting with hope, he tries the furrow’d grounds;
 But when the tainted gales the game betray,
 Couch’d close he lies and meditates the prey:
 Secure they trust th’ unfaithful field beset,
 Till hovering o’er them sweeps the swelling net.”

POPE’S WINDSOR FOREST.



THE WATER-SPANIEL

Is excellent for hunting otters, wild ducks, and other game whose retreat is among the rushes and reeds which cover the banks of rivers, the fens, and the ponds. He is very sagacious, and perhaps the most docile and tractable of all the canine tribe. The Newfoundland Dogs are of this class, and the largest of the family.

The *Water-Spaniel* will fetch and carry whatever he is bid, and often dive to the bottom of deep water in search of a piece of money, which he brings up in his mouth, and lays at the feet of whoever sent him: the best breed has black curly hair and long ears.

The beautiful breed of Spaniels known as King Charles's, are highly prized for their diminutive size and length of ears. They are found of all colours, but those which are black, with tanned cheeks and legs, are considered the purest breed.

They derive their name from King Charles the Second, who, as Evelyn tells us, "took great delight in having a number of little spaniels follow him and lie in his bed-chamber."





THE NEWFOUNDLAND DOG. (*Canis sensilis.*)

THIS animal was originally brought into Europe from Newfoundland, whence it derives its name, and where it is extremely useful to the settlers, almost supplying the place of a horse. There are several varieties, differing slightly in size and appearance, but the full size is about six feet and a half from the nose to the tip of the tail, the length of which is two feet. He is noble in appearance, and covered with long shaggy hair of a black and white colour, in which the latter generally predominates.

The Newfoundland Dog is affectionate, sagacious, and docile beyond all others; and being web-footed is excellently adapted for the water; and there are innumerable instances of his rescuing man from a watery grave.

The anecdotes which illustrate the affection and sagacity of this animal would fill a volume, but we select one relating to the water, as that appears his noblest scene of action.

Some time ago a young woman was nursing an infant on one of the quays on the Liffey, when it made a sudden spring from her arms and fell into the water. The screaming nurse and anxious spectators saw the child sink, as they thought to rise no more; when at the very instant a Newfoundland Dog, which was accidentally passing, rushed to the spot, and at the sight of the child, who at that moment reappeared, sprang into the water. The child again sunk, and the faithful animal was seen anxiously swimming round the spot. Once more the

child rose, and the Dog gently, but firmly, seized him and bore him to land. Meanwhile a gentleman arrived who appeared to take much interest in the affair, and on the person who had the child turning to show it him, he recognized the well-known features of his own son. A mixed sensation of horror, joy, and surprise struck him mute. When he recovered himself he lavished a thousand caresses on the faithful animal, and offered his master five hundred guineas for him; but the latter felt too much affection for the noble animal to part with him on any consideration whatever.

We also subjoin another equally interesting.


A native of Germany, fond of travelling, was pursuing his course through Holland, accompanied by a large Newfoundland Dog. Walking one evening on a high bank, which formed one side of a dike, or canal, so common in that country, his foot slipped, and he was precipitated into the water, and being unable to swim he soon became senseless. When he recovered his recollection he found himself in a cottage on the other side of the dike, surrounded by peasants, who had been using means to restore suspended animation. The account given by them was, that one of them, returning home from his labour, observed at a considerable distance a large Dog in the water swimming, and dragging the body of a man into a small creek on the opposite side to which the men were.

The Dog having shaken himself, began industriously to lick the hands and face of his master, while the rustic hastened across; and, having obtained assistance, the body was conveyed to a neighbouring house, where the usual means of resuscitation soon restored him to sense and recollection. Two very considerable bruises, with the marks of teeth, appeared, one on his shoulder and the other on the nape of his neck; whence it was presumed that the faithful animal first seized his master by the shoulder, and swam with him in this manner some time; but that his sagacity had prompted him to let go this hold, and shift his grasp to the neck, by which he had been enabled to support the head out of water. It was in the latter position that the peasant observed the Dog making his way along the dike, which it appeared he had done for the distance of nearly a quarter of a mile.



THE GREYHOUND (*Canis familiaris graius*)

Is well known, and was formerly held in such estimation, that he was the especial companion of a gentleman, who, in ancient times, was distinguished by his horse, his hawk, and his Greyhound, and it was penal for any person of inferior rank to keep one. He is the fleetest of all Dogs, and can outrun every animal of the chase. He has a long body, and is of an elegant shape; his head is neat and sharp, with a full eye, a good mouth, sharp and very white teeth; his tail is long, and curls round above his hind part. There are several varieties; as the Italian Greyhound, the Oriental Greyhound, and the Irish Greyhound, or Wolf-dog. They are used for coursing; that is, hunting by sight instead of scent; and are principally employed in chasing hares. Daniel, in his *Rural Sports*, tells us, that a brace of Greyhounds have been known to course a hare four miles in twelve minutes; turning it several times, till the poor creature dropped at last quite dead from fatigue.



THE FOX. (*Canis Vulpes.*)

THIS well-known animal, which is found in most countries, varies much in size and colour. In the North of Europe some are white, some red, and others black; but the general colour is a yellow-brown. His abode is generally on the skirt of a wood, as near a farm-yard as possible, in a hole, of which some other animal has either been dispossessed or voluntarily deserted. Thence he issues at night, and cautiously approaching the poultry, kills all that he can find, conveying them one by one to different hiding places, which he visits when hungry. He will continue his depredations till day-break, or until he is alarmed, often depopulating a whole poultry-yard in one night. When, however, his choice food, the chicken, is not accessible, he devours animal food of every description; and if his habitation be near the water he will even content himself with shell-fish. In France and Italy he does much

damage to the vineyards, being very fond of grapes, and spoiling many for the sake of one bunch.


His name has passed into a proverb for cunning and deceitfulness; and, unlike the dog tribe to which he belongs, he is totally unsusceptible of any sentiment of gratitude.

His bite is tenacious and dangerous, as the severest blows cannot make him quit his hold; his eye is most significant, and expressive of almost every passion. He generally lives about twelve or fifteen years

The female produces but once a year, and seldom has more than four or five cubs at a litter. The first year the young is called a Cub, the second year a Fox, and the third year an Old Fox. The tail is very bushy, and is called the brush.

In this country he is hunted with horses and hounds, and no animal affords greater diversion and occupation to the sportsman. When pursued he usually makes for his hole; but should his retreat be cut off, his stratagems and shifts to escape are singularly acute. He seeks woody and uneven parts of the country, preferring the paths the most embarrassed by thorns and briars, and running in a straight line before the hounds, at no great distance from them; and, when overtaken, he turns on his assailants, and fighting with obstinate despair, dies in silence.





THE WOLF, (*Canis Lupus*,)

When hungry, is an undaunted and most ferocious inhabitant of the woods, but a coward when the stimulus of appetite is no longer in action. He delights to roam in mountainous countries, and is a great enemy to sheep and goats; the watchfulness of dogs can hardly prevent his depredations, and he often dares to visit the haunts of men, howling at the gates of cities and towns. His head and neck are of a cinereous colour, and the rest of a pale yellowish brown. He commonly lives to the age of fifteen or twenty years. He possesses a most exquisite power of smelling his prey at a great distance. Wolves are found nearly everywhere, except in the British islands, where this noxious race has been entirely extirpated. King Edgar first attempted to effect this by remitting the punishment of certain crimes on producing a number of Wolves' tongues; and in Wales, the tax of gold and silver was commuted for an annual tribute of Wolves' heads. In the reign of Athelstan, Wolves abounded so

much in Yorkshire, that a retreat was built at Flixton, to defend passengers from their attacks. They infested Ireland many centuries after their extinction in England: the last presentment for killing Wolves was made in the county of Cork about the year 1710. They abound in the immense forests of Germany, and they are also found in considerable numbers in the South of France. Everywhere that they are wild, so great is the general detestation of this destructive creature, that all other animals endeavour to avoid it. In a state of captivity, however, the Wolf is said to be remarkably anxious to attract the attention of man, and to rub itself against the bars of its cage when noticed. The Wolf is sometimes affected with madness, in symptoms and consequences exactly similar to that which affects the dog; but this disease, as it generally happens in the depth of winter, cannot be attributed to the great heat of the dog-days. In the northern parts of the world, wolves are said, frequently, in the spring, to get upon the fields of ice adjoining the sea, for the purpose of preying upon the young seals, which they there find asleep; but vast pieces of the ice occasionally detaching themselves from the mass, they are carried with them to a great distance from the land, where they perish amidst the most hideous and dreadful howling. The language of the poet is beautifully descriptive of this creature's insatiable fury:—

“ By wintry famine roused, from all the tract
Of horrid mountains, which the shining Alps,
And wavy Apennine, and Pyrenees,
Branch out, stupendous, into distant lands,
Cruel as death! and hungry as the grave!
Burning for blood! bony, and gaunt, and grim!
Assembling Wolves, in raging troops, descend;
And, pouring o'er the country, bear along,
Keen as the north wind sweeps the glossy snow:
All is their prize.”



THE JACKAL, (*Canis aureus*,)

Commonly called *the lion's provider*, is not much larger than the fox, which he resembles in the appearance of the fore part of his body. His skin is of a bright yellow colour. The Jackals often unite to attack their prey, and make a most hideous noise, which, rousing the king of the forest from his slumbers, brings him to the place of food and plunder: at his arrival, the petty thieves, awed by the greater strength of their new messmate, retire to a distance; and hence the fabulous story of their attendance on the lion, to provide for his food.—These animals are always seen in large flocks of forty or fifty; and hunt, like hounds in full cry, from evening till morning. In the absence of other food, they drag the dead out of their tombs, and feed greedily on putrid corpses; but notwithstanding their natural ferocity, it is said that when taken young they may be easily tamed, and, like dogs, they love to be fondled, wag their tails, and show a considerable degree of attachment to their masters. They are common in many parts of the East: and as they act as scavengers, the people do not annoy them in their nocturnal visits.



THE HYÆNA. (*Canis Hyæna.*)

THIS animal was long supposed to be the most savage and untractable of all quadrupeds; but it is now found he may be tamed. He is covered with long, coarse, and rough ash-coloured hair, marked with long black stripes, from the back downwards; the tail is very hairy. His teeth and jaws are so constructed, as to enable him to crush the largest bones with ease; and his tongue is as rough as a coarse file. Like the jackal, he attacks the flocks and herds, caring little for the watchfulness or strength of dogs; and when pressed with hunger, comes and howls at the gates of towns, and violates the repositories of the dead, tearing up the bodies from the graves, and devouring them. He is now only found wild in Asia and Africa, but is supposed to have formerly inhabited Europe. When receiving his food, the eyes of this fierce animal glisten, the bristles of his back stand erect, he grins fearfully and utters a snarling growl.



AMERICAN BLACK BEAR. (*Ursus Americanus*.)

THIS animal inhabits the Northern districts of America, where it is found in considerable numbers. It is somewhat smaller than the Brown or European Bear; its colour of an uniform and glossy black. Its food consists chiefly of fruits, the young shoots and roots of vegetables and grain. In quest of these it occasionally emigrates from the northern to the more southern regions. Their retreats, during the period of gestation, are so impenetrable, that although immense numbers of Bears are annually killed in America, a female is rarely found among them. In autumn, when they are become exceedingly large by feeding on acorns and other arborescent food, their flesh is extremely delicate, the hams in particular are highly esteemed, and the fat is remarkably white and sweet. At this time and during the winter they are hunted, and killed in great numbers by the American Indians.

THE BROWN EUROPEAN BEAR, (*Ursus Arctos*,) is a native of the North of Europe, and also of the Alps

of Switzerland, and the Pyrenees. He is a great sleeper, and passes the whole winter in his den, without any particular food: but if we consider his being at rest, losing little by perspiration, and never retiring to his winter-quarters before he is properly fattened, his abstinence will cease to be wonderful. When tamed, this animal appears mild and obedient to his master; he may be taught to walk upright, to dance, to lay hold of a pole with his paws, and perform various tricks to entertain the multitude, who are highly pleased to see the awkward movements of this rugged creature, which it seems to suit to the sound of an instrument, or to the voice of its leader. The discipline Bears undergo in teaching them to dance is so severe that they never forget it; and an amusing story is told of a gentleman who was pursued by a Bear, and who, when in despair he turned and raised his stick against his assailant, was astonished to see the Bear rear itself on its hind legs, and begin to dance. It had escaped from captivity, and had been taught to dance when a stick was held up by its keeper. But to give the Bear this kind of education, it must be taken when young, and accustomed early to restraint and discipline, as an old Bear will not suffer constraint without discovering the most furious resentment: neither the voice nor the menaces of his keeper have any effect upon him; he growls equally at the hand that is held out to feed, and that which is raised to correct him. The female Bears bring forth two or three young, and are very careful of their offspring. The fat of the Bear is reckoned very useful in rheumatic complaints, and for anointing the hair; his fur affords comfort to the inhabitants of cold climates, and ornaments to those of warm. It was anciently supposed that the young Bear, when first brought forth, was merely an unformed mass, till its mother licked it into shape; and hence the expression "he wants licking into shape" was frequently employed by the old dramatists, when speaking of an awkward, clownish man.

The Brown Bear was at one time common in the British islands. "Many years ago it has been swept away so completely, that we found it imported for baiting,

a sport in which our nobility, as well as the commonalty, of the olden time—nay, even royalty itself—delighted. A bear-bait was one of the recreations offered to Elizabeth at Kenilworth, and in the Earl of Northumberland's Household Book, we read of twenty shillings for his bearward. In Southwark there was a regular bear-garden, that disputed popularity with the Globe and Swan theatres on the same side of the water. Now, however, so much do tastes alter, (in this instance certainly for the better) such barbarous sports are banished from the metropolis."

The Bear is a flat-footed animal, and can stand easily upon its broad hind feet, but is extremely awkward and sluggish in its movements. He possesses, however, the faculty of climbing to an extraordinary degree; and in his native country ascends frequently lofty trees in pursuit of honey, of which he is excessively fond. Bears swim well, and will cross not only broad rivers, but sometimes even an arm of the sea.





THE POLAR, OR GREAT WHITE BEAR.

(*Ursus maritimus.*)

THE Polar Bear is generally from six to eight feet long. The fur is long and white, with a tinge of yellow, which becomes darker as the animal advances in age; the ears are small and round, and the head long. It inhabits Greenland and Lapland, and is found as far north as eighty degrees. It walks heavily, and is very clumsy in all its motions; its senses of hearing and seeing appear very dull, but its smell is very acute; and it does not appear destitute of some degree of understanding, or at least of cunning. Captain King, who visited the shores of the Arctic Ocean in 1835, relates a curious instance of the cunning of this animal "On one occasion a Polar Bear was seen to swim cautiously to a large piece of ice, on which two female walrus were lying asleep with their cubs. The Bear crept up some hammocks behind them, and with his fore feet loosened a large block of ice, which, with the help of his nose and paws, he rolled and carried till it was immediately over the heads of the sleepers, when he let it fall on one of the old animals, which was instantly killed. The other walrus, with its cubs, rolled into the water, but the young one of the murdered female remained by its dam, and on this help-

less creature the Bear rushed, thus killing two animals at once."

The ferocity of this kind of Bear is equal to its cunning. A few years since, the crew of a boat belonging to a ship in the whale-fishery, shot at a Bear at a short distance and wounded it. The animal immediately set up the most dreadful yells, and ran along the ice towards the boat. Before it reached it, a second shot was fired, and hit it. This served to increase its fury. It presently swam to the boat; and in attempting to get on board, placed its fore foot upon the gunwale; but one of the crew having a hatchet, cut it off. The animal still, however, continued to swim after them till they arrived at the ship, and several shots were fired at it, which also took effect; but on reaching the ship it immediately ascended the deck, and the crew having fled into the shrouds, it was pursuing them thither, when a shot from one of them laid it dead on the deck.



THE RACCOON. (*Procyon*, or *Ursus lotor*.)

THIS animal is a native of America, of the bear tribe: in Jamaica they are very numerous, and do incredible mischief to the plantations of sugar-cane and Indian corn, especially to the latter while it is young. The Raccoon is less than the fox in size, and has a sharp-pointed nose. His fore legs are shorter than the others. The colour of his body is grey, with two broad rings of black round the eyes, and a dusky line running down the middle of the face. In the wild state the Raccoon is savage and sanguinary, committing great destruction among both wild and domesticated birds, without consuming any part of them

except the head, or the blood which flows from their wounds. It is a good climber, the form of its claws enabling it to adhere to the branches of trees with great tenacity. Raccoons are easily domesticated, and then become very amusing and harmless animals. They are as mischievous as a monkey, seldom at rest, and extremely sensible of ill treatment, which they never forgive. They have great antipathy to sharp and harsh sounds, such as the bark of a dog, and the cry of a child. They eat of everything that is given them, and, like the cat, are good providers, hunting after eggs, fruit, corn, insects, snails, and worms; and generally dip their food in water before devouring it. A peculiarity which few other animals are found to possess is, that they drink as well by lapping like the dog, as by sucking like the horse. These animals are hunted for the sake of their fur, which is used by the hatters, and is considered next in value to that of the beaver; it is used also in linings for garments. The skins, when properly dressed, are made into gloves and upper-leathers for shoes. The negroes frequently eat the flesh of the Raccoon, and are very fond of it, though it has a very disagreeable and rank smell. The American hunters pique themselves on their skill in shooting Raccoons; which, from the extraordinary vigilance and cunning of the animals, is by no means an easy task.

When eating they support themselves on their hind feet, and carry their food to the mouth with their fore paws. Some of them are very fond of oysters and other shell-fish, and show great dexterity in keeping the shells open, while they extract the contents. Their most remarkable peculiarity, however, is that already mentioned, of dipping their food in water when there is any within their reach; though when there is not, they seem quite contented to eat it dry.

THE BADGER. (*Ursus Meles.*)

THIS animal inhabits most parts of the world. The length of the body is about two feet six inches from the nose to the insertion of the tail, which is short, and black like the throat, breast, and belly; the hair of the other part of the body is long and rough, of a yellowish white at the roots, black in the middle, and greyish at the point: the toes are much enveloped in the skin, and the long claws of the fore feet enable the animal to dig with great effect: under the tail there is a receptacle, in which is secreted a white fetid substance, that constantly exudes through the orifice, and thus gives the body a most unpleasant smell. Being a solitary animal, it digs a hole for itself, at the bottom of which it remains in perfect security: it feeds upon young rabbits, birds and their eggs, and honey. The cruel sport of Badger-hunting with dogs is still practised; and the motions of the animal when attacked are so quick, that a dog is often desperately wounded in the first assault. The Badger, when nearly overpowered, falls upon his back, combats with desperate resolution, and seldom dies unrevenged on his enemies. It is indeed not easy to overcome him; his skin is so thick that it resists the impression of the teeth, and so loose, that even when a dog has caught hold of it, the Badger is enabled to turn round easily and bite his assailants in the most tender parts. The female has generally three or four young ones at a time.

THE COATI-MONDI. (*Viverra nasua.*)

THIS creature is a native of Brazil, not unlike the Raccoon in the general form of the body, and, like that animal, frequently sits up on the hinder legs, and in this position, with both paws carries its food to its mouth. Even in a state of tameness, it will pursue poultry, and destroy every living thing that it has strength to conquer. When it sleeps it rolls itself into a ball, and remains immoveable for fifteen hours together. Its eyes are small, but full of life; and, when domesticated, it is very playful and amusing. A great peculiarity belonging to this animal is the length of its snout, which resembles in some particulars the trunk of the elephant, as it is moveable in every direction. The ears are round, and like those of a rat; the fore feet have five toes each. The hair on the back is short and rough and of a blackish hue; the tail marked with rings of black, like the wild cat; the rest of the body is a mixture of black and red. This animal is very apt to eat its own tail, which is very long; but this strange appetite is not peculiar to the Coati alone; the mococo, or honey-bear, and some of the monkey tribe do the same, and seem to feel no pain in wounding a part of the body so remote from the centre of circulation.




THE CIVET CAT (*Viverra Civetta*)

Is found throughout India, Guinea, and the Philippine Islands, and is famous for producing the perfume called *civet*. He is kept for the sake of this perfume, and fed with a kind of soup made of millet, or rice, with a little fish or flesh boiled with it in water. The civet obtained from the male is esteemed the best. It is found in a large double glandular receptacle, situated at a little distance beneath the tail. When a sufficient time for the secretion has been allowed, one of these animals is put into a long wooden cage, so narrow that it cannot turn itself round. The cage being opened by a door behind, a small spoon is introduced through the orifice of the pouch, which is carefully scraped; this is done twice or thrice a week, and the animal is said always to produce the most civet after being irritated. The Civet, although a native of the warmest climates, is yet found to live in temperate, and even cold countries, provided it be defended carefully from the injuries of the air.

In general appearance this animal reminds one of the fox, which it also resembles in its predatory habits. It is bred in Holland, for the sake of its perfume, which is sold

as an article of commerce.* The civet of Amsterdam is considered purer than any other, as people of other countries adulterate it with gums, and other ingredients, to increase its weight. In a wild state, the Civet lives entirely on birds and small quadrupeds; and at any time a small quantity of salt is said to poison it.



THE ICHNEUMON, OR EGYPTIAN MANGOUSTE,
OR PHARAOH'S RAT. (*Herpestes Pharaonis.*)

THIS animal bears a close resemblance to the weasel tribe, both in form and habits. From the tip of the nose to the root of the tail, it is about eighteen inches in length. At the base, the tail is very thick, tapering gradually towards the point, which is slightly tufted. It has a long active body, short legs, lively and piercing eyes, and a pointed nose; the hair is rough and bristly, of a pale reddish grey. In a wild state the Ichneumon usually resides on the banks of rivers; and swims and dives like the otter, being able to keep under water for a great length of time.

* Important medical virtues were formerly attributed to the civet, but it now no longer forms an article in the *Materia Medica*, and even as a perfume it has been laid aside.




THE CIVET CAT (*Viverra Civetta*)

Is found throughout India, Guinea, and the Philippine Islands, and is famous for producing the perfume called *civet*. He is kept for the sake of this perfume, and fed with a kind of soup made of millet, or rice, with a little fish or flesh boiled with it in water. The civet obtained from the male is esteemed the best. It is found in a large double glandular receptacle, situated at a little distance beneath the tail. When a sufficient time for the secretion has been allowed, one of these animals is put into a long wooden cage, so narrow that it cannot turn itself round. The cage being opened by a door behind, a small spoon is introduced through the orifice of the pouch, which is carefully scraped; this is done twice or thrice a week, and the animal is said always to produce the most civet after being irritated. The Civet, although a native of the warmest climates, is yet found to live in temperate, and even cold countries, provided it be defended carefully from the injuries of the air.

In general appearance this animal reminds one of the fox, which it also resembles in its predatory habits. It is bred in Holland, for the sake of its perfume, which is sold

as an article of commerce.* The civet of Amsterdam is considered purer than any other, as people of other countries adulterate it with gums, and other ingredients, to increase its weight. In a wild state, the Civet lives entirely on birds and small quadrupeds; and at any time a small quantity of salt is said to poison it.



THE ICHNEUMON, OR EGYPTIAN MANGOUSTE,
OR PHARAOH'S RAT. (*Herpestes Pharaonis.*)

THIS animal bears a close resemblance to the weasel tribe, both in form and habits. From the tip of the nose to the root of the tail, it is about eighteen inches in length. At the base, the tail is very thick, tapering gradually towards the point, which is slightly tufted. It has a long active body, short legs, lively and piercing eyes, and a pointed nose; the hair is rough and bristly, of a pale reddish grey. In a wild state the Ichneumon usually resides on the banks of rivers; and swims and dives like the otter, being able to keep under water for a great length of time.

* Important medical virtues were formerly attributed to the civet, but it now no longer forms an article in the *Materia Medica*, and even as a perfume it has been laid aside.

The Ichneumon is celebrated in the mythology of ancient Egypt, where it has long been domesticated, and where it was ranked amongst its divinities, on account of its great utility in destroying serpents, snakes, rats, mice, and other vermin: it is also fond of crocodiles' eggs, which it digs out of the sand where they have been deposited. It is a very fierce, though small animal, and will fight with dogs, foxes, and even jackals, with great fury. It will not breed in confinement, but may be easily tamed when taken young.

The following particulars are related by M. D'Obsonville, in his *Essays on the Nature of various foreign Animals*:—"I had an Ichneumon very young, which I brought up. I fed it at first with milk, and afterward with baked meat mixed with rice. It soon became even tamer than a cat; for it came when called, and followed me, though at liberty, in the country. One day I brought this animal a small water-serpent alive, being desirous to know how far his instinct would carry him against a being with which he was as yet totally unacquainted. His first emotion seemed to be astonishment mixed with anger, for his hair became erect; but in an instant he slipped behind the reptile, and with remarkable swiftness and agility leaped upon its head, seized it, and crushed it between its teeth. This essay, and new food, seemed to have awakened in him his innate and destructive voracity, which till then had given way to the gentleness he had acquired from education. I had about my house several curious kinds of fowls, among which he had been brought up, and which, till then, he had suffered to go and come unmolested and unregarded: but a few days after, when he found himself alone, he strangled them every one, ate a little, and, as it appeared, drank the blood of two."

The mode in which the Ichneumon seizes a serpent is thus described by Lucan in his *Pharsalia*:—

"Thus oft the Ichneumon, on the banks of Nile,
Invades the deadly aspic by a wile;
While artfully his slender tail is played,
The serpent darts upon the dancing shade,
Then turning on the foe with swift surprise,
Full on the throat the nimble traitor flies,
And in his grasp the panting serpent dies."



THE WEASEL. (*Mustela vulgaris*.)

THE animals belonging to this genus, notwithstanding their small size, are all carnivorous, and from their slender and lengthened bodies, short legs, and the very free motion in every direction, permitted by the loose articulations of the spine, are well formed for pursuing their prey into the deepest recesses. Constituted by nature to subsist on animals, many of which have great strength and courage, they possess an undaunted and ferocious disposition. The Weasel has a long and thin body; its length, with its tail, is only seven inches, and its height not more than an inch and a half. There are some of this species of different colours, as red, brown, black, and some all white. There are two sorts, the domestic Weasel, living in houses, barns, &c.; and the wild, living in the woods and mountains. In the northern parts of Europe they are very numerous. Mice of every description, the field and the water-vole, rats, moles, and small birds are their ordinary food, and occasionally

rabbits and partridges. When driven by hunger, it will boldly attack the poultry-yard. The Weasel, when it enters a hen-roost, never meddles with the cocks or old hens, but makes choice of the pullets and young chickens; these it kills with a single stroke on the head, and carries away one after the other. It sucks the eggs with avidity, making a small hole at one end, through which it draws out the yolk. In winter it resides in granaries and hay-lofts, and in summer chooses the low lands about the mills and streams, where it hides among the bushes, and in the hollows of old trees.

It was formerly supposed that the Weasel was untamable; but Buffon, in a supplementary volume, corrects this error, and from a letter of a female correspondent, shows that it may be rendered as familiar as a cat or a lapdog. It frequently eat from his correspondent's hand, and seemed fonder of milk and fresh meat than of any other food. "If I present my hands," says this lady, "at the distance of three feet, it jumps into them without ever missing. It shows a great deal of address and cunning, in order to accomplish its ends, and seems to disobey certain prohibitions merely through caprice. During all its actions it seems solicitous to divert and be noticed, looking at every jump and at every turn to see whether it be observed or not. If no notice be taken of its gambols, it ceases them immediately, and betakes itself to sleep; and when awaked from the soundest sleep, it instantly resumes its gaiety, and frolics about in as sprightly a manner as before. It never shows any ill humour, unless when confined or too much teased, in which case it expresses its displeasure by a sort of murmur, very different from that which it utters when pleased."

Weasels and ferrets are used by rat-catchers to drive the rats out of their holes; and they kill a great many, the habit of the Weasel being to kill its prey by biting the head, so that the teeth penetrate the brain, and then to throw the body aside, or hide it till a future period.



THE FERRET (*Mustela furo*)

Is a small, yet bold animal, and an enemy to all others but those of his own kind. He is made somewhat like the weasel, and breeds in all countries of Europe. His eyes are remarkably fiery. The tame ones live on milk, bread, barley, &c.; the wild ones on the blood of rabbits, hares, &c. He is much used to drive rabbits from their holes, and for this purpose is always muzzled, for he is such an inveterate enemy to the rabbit, that if a dead one be presented to a young Ferret, he instantly bites it with an appearance of rapacity; or if it be living, the Ferret seizes it by the neck, winds himself round it, and continues to suck its blood till he be satiated; indeed, his appetite for blood is so strong, that he has been known to attack and kill children in the cradle. He is very soon irritated; and his bite is very difficult to be cured.

Our figure is full large, as the length of the animal is usually about thirteen inches, exclusive of the tail, which is about five.



THE GENET. (*Viverra Genetta.*)

- THIS animal is about the size of a small cat. The skin is spotted and beautiful, of a reddish grey colour. The spots on the sides are round and distinct, those on the back almost close; its tail is long, and marked with seven or eight rings of black. From an orifice beneath its tail, it yields a kind of perfume, which smells faintly of musk. This little animal is meek and gentle, except when provoked, and is easily domesticated. In Constantinople it strays from house to house like our cat, and keeps whatever house it is in perfectly free from mice and rats, which cannot endure its smell. It is found wild in various parts of Europe, and also at the Cape of Good Hope. Its fur is beautiful and soft, and valuable as an article of commerce. The eyes of the Genet contract when exposed to the light, like those of the cat; and it can draw in its claws in nearly the same manner. The favourite animal called *Delun-drung* seems to be an intermediate genus between the cat and the Genet.

The Genet nearly resembles the marten in its habits and disposition, but can be tamed with much greater facility. "Its nature," says Goldsmith, "is mild and gentle, its colour various and glossy, its fur valuable; and, upon the whole, it seems to be one of those animals that, with proper care, might be propagated among us, and become one of the most serviceable of our domestics."



THE POLECAT. (*Mustela putorius.*)

THE strong and disagreeable smell of this animal is proverbial; its skin is stiff, hard, and rugged, and when well prepared, is very desirable as clothing. It is about seventeen inches in length, exclusive of the tail, which is about six inches. The breast, tail, and legs are of a blackish colour, but the belly and sides yellowish. It conceals itself in secret corners at tops of houses, and is a disastrous pest to the poultry-yard. Some of these animals frequent the woods and destroy a great quantity of birds; and some, forsaking the haunts of man, retire to the rocks and crevices of the cliffs on the sea-shore, preferring a meagre and scanty diet with security, to the daintiness of chicken-flesh and eggs, attended with trouble and fear. Rabbits seem to be their favourite prey, and a single Polecat is often sufficient to destroy a whole warren; for with that insatiable thirst for blood which is natural to all the weasel tribe, it kills much more than it can devour; and twenty rabbits have been found dead, which one Polecat had destroyed by a wound hardly perceptible. The *Polecat* is the same with the *Fitchet* or *Foumart*, the hair of which is made into fine brushes and pencils for the use of painters. This small animal is fierce and bold. When attacked by a dog, it will defend itself with great spirit, attack him in turn, fastening upon the nose of its enemy with so keen a bite, as frequently to oblige him to desist. When heated or enraged, the smell it emits is absolutely intolerable.



THE SABLE. (*Mustela*, or *Martes Zibellina*.)

THIS animal is a native of North America, Siberia, Kamtschatka, and Asiatic Russia, and it frequents the banks of rivers, and the thickest parts of the woods. It lives in holes under the ground, and especially under the roots of trees; but sometimes makes its nest, like the squirrel, in the hollows of trees. The skin of the Sable is more valuable than that of any other animal of equal size. One of these skins, not more than four inches broad, has sometimes been valued at as high a rate as fifteen pounds; but the general price is from one to ten pounds, according to the quality. The Sable's fur is different from all others, its peculiarity being that the hair turns with equal ease either way; on which account fur dealers sometimes blow the fur of any article they may be selling, to show that it is really Sable. The tails are sold by the hundred, at from four to eight pounds.

The common, or Beech Marten, (*Mustela Martes*), like the Sable, boasts the honour of adorning with his fur the rich and the beautiful; as princes, ladies, and opulent people of all nations pride themselves in wearing his spoils. He is about as big as a cat, but the body is much longer proportionately, and the legs shorter. His skin is of a light brown, with white under the throat, and yellow on the back. The fur of the Marten fetches a good price, and is much used in European countries, though very far inferior to that of the Sable: the best, which is called Stone Marten fur by the furriers, is imported from Sweden and Russia. The Pine, or Yellow-breasted Marten, is another species, the fur of which is nearly equal to that of the sable, though it is much cheaper.



THE OTTER. (*Mustela Lutra.*)

“Forth from his den the Otter drew,—
 Grayling and trout their tyrant knew,
 As between reed and sedge he peers,
 With fierce round snout and sharpened ears,
 Or, prowling by the moonbeam cool,
 Watches the stream or swims the pool.”

SCOTT.

As the Otter lives principally on fish, the formation of his body is such as will enable him to swim with the greatest facility. His body is flattened horizontally; his tail is flat and broad; his legs are short, and his toes webbed. His teeth are very strong and sharp; and his body, besides its fur, has an outer covering of coarse shining hair. The Otter is a perfect epicure in his food; he seldom eats an entire fish, but beginning at the head, eats that, and about half the body, always rejecting the tail. When the rivers and ponds are frozen so that the Otter can get no fish, he will visit the neighbouring farm-yards, where he will attack the poultry, sucking-pigs, and even lambs. An Otter may be tamed, and taught to catch fish enough to sustain not only himself, but a whole family. Goldsmith states that he saw an Otter go to a gentleman's pond at the word of command, drive the fish into a corner, and seize upon the largest of the whole, bring it off, and give it to his master.


Bewick, in his *History of Quadrupeds*, states, that a person of the name of Collins, who lived at Kilmerston, near Wooler, in Northumberland, had a tame Otter, which followed him wherever he went. He frequently

took it to fish in the river; and, when satiated, it never failed to return to him. One day, in the absence of Collins, the Otter, being taken out to fish by his son, instead of returning as usual, refused to come at the accustomed call, and was lost. The father tried every means in his power to recover the animal; and, after several days' search, being near the place where his son had lost it, and calling it by name, to his inexpressible joy it came creeping to his feet, and showed many marks of affection and attachment.

The female Otter produces four or five young ones at a birth, and these in the spring of the year. Where there have been ponds near a gentleman's house, instances have occurred of their littering in cellars or drains. The male utters no noise when taken, but the females sometimes emit a shrill squeak.

Otters are generally caught in traps placed near their landing-places, and carefully concealed in the sand. When hunted by dogs, the old ones defend themselves with great obstinacy. They bite severely, and do not readily quit their hold. Otter-hunting is a favourite sport in many parts of Great Britain; particularly in the midland counties of England, and in Wales.





THE SEAL. (*Phoca vitulina.*)

THE amphibious flesh-eating animals, though nearly allied to the otter in their habits, are very different in the construction of their bodies. Their feet are so short and so enveloped in skin, that they are of scarcely any use in assisting the animal on dry land; so that the Seal's progress on solid ground is only effected by a sort of half tumbling, jumping, and shuffling motion, excessively ridiculous to a looker-on. The feet, however, which are furnished with strong claws, are of use in enabling the animal to climb out of the water over a rocky shore. For swimming, the Seal is admirably adapted; its long flexible body is shaped like that of a fish, tapering to the tail; and it is furnished with strong webs between the toes, so as to make the fore feet act as oars, and the hind feet, which the animal generally drags behind it like a tail, to serve as a rudder. The Seal lives generally in the water, and feeds entirely on fish; only coming to shore occa-

sionally to bask on the sands, and to lie there to suckle its young. The usual length of a Seal is five or six feet. The head is large and round; the neck small and short; and on each side of the mouth there are several strong bristles. From the shoulders the body tapers to the tail, which is very short. The eyes are large: there are no external ears; and the tongue is cleft or forked at the end. The body is covered with short thick-set hair, which in the common species is generally grey, but sometimes brown or blackish. There are, however, several species; and one of them, which is called the sea-leopard, has the fur spotted with white or yellow.

Seals are hunted by the Greenlanders for the sake of their oil, and also for their skins, which are used for making waistcoats and other articles of clothing, and are much prized by the fishermen for their great warmth. As Seals cannot breathe below water, they are obliged to swim with their heads as much above the waves as a dog would, and thus are easily seen by the fishermen. When attacked, they fight with great fury; but when taken young, are capable of being tamed; they will follow their master like a dog, and come to him when called by the name given to them. Some years ago a young Seal was thus domesticated. It was taken at a little distance from the sea, and was generally kept in a vessel full of salt water: but sometimes it was allowed to crawl about the house, and even to approach the fire. Its natural food was regularly procured for it; and it was carried to the sea every day, and thrown in from a boat. It used to swim after the boat, and always allowed itself to be taken back. It lived thus for several weeks, and probably would have lived much longer, had it not been sometimes too roughly handled. The females in this climate bring forth in winter, and rear their young upon some sand-bank, rock, or desolate island, at some distance from the main land. When they suckle their young, they sit up on their hinder legs, while the little Seals, which are at first white, with woolly hair, cling to the teats, of which there are four in number. In this manner the young continue in the place where they are brought forth

for twelve or fifteen days; after which the dam brings them down to the water, and accustoms them to swim and get their food by their own industry.

In Newfoundland the Seal-fishery forms an important source of wealth, and numerous ships are sent out every season among the ice in search of Seals. One ship has been known to catch five thousand Seals, but about half that number is the usual quantity taken. As soon as the Seal is killed, it is skinned, and the pelt, as the skin and blubber together is called, being preserved, the body of the Seal is either eaten by the sailors, or left on the ice for the polar bears.

The aboriginal inhabitants of the northern regions have several strange superstitions about Seals. They believe that Seals delight in thunder-storms; and say, that during these times they will sit on the rocks, and contemplate, with apparent pleasure and gratification, the convulsion of the elements. The Icelanders, in particular, are said to believe that these animals are the offspring of Pharaoh and his host, who were converted into Seals when they were overwhelmed in the *Red Sea*.



THE WALRUS, MORSE, OR SEA-COW.

(*Trichecus Rosmarus.*)

THIS very curious animal is nearly allied to the Seal, but is of much greater size, being frequently eighteen feet in length, and from ten to twelve feet in girth. The head is round, the eyes are small and brilliant, and the upper

lip, which is enormously thick, is covered with pellucid bristles, as large as a straw. The nostrils are very large, and there are no external ears. The most remarkable part of the Walrus is, however, his two large tusks in the upper jaw; they are inverted, the points nearly uniting, and sometimes exceed twenty-four inches in length! the use which the animal makes of them is not easily explained, unless they help him to climb up the rocks and mountains of ice among which he takes up his abode, as the parrot employs his beak to get upon his perch. The tusks of the Walrus are equal for durability and whiteness to those of the elephant, and, as they keep their colour much longer, are preferred by dentists to any other substance for making artificial teeth.

The Walrus is common in some of the northern seas, and will sometimes attack a boat full of men. They are gregarious animals, usually found in herds, sleeping and snoring on the icy shores; but when alarmed, they precipitate themselves into the water with great bustle and trepidation, and swim with such rapidity, that it is difficult to overtake them with a boat. One of their number always keeps watch while the others sleep. They feed on shellfish and sea-weeds, and yield an oil equal in goodness to that of the whale. The white bear is their greatest enemy. In the combats between these animals, the Walrus is said to be generally victorious, on account of the desperate wounds it inflicts with its tusks. The females have only one young one at a time, which, when born, resembles a good-sized pig.






SECTION II.—Insect-eating Animals.

THE HEDGEHOG. (*Erinaceus Europæus*.)

THIS animal is something like a porcupine in miniature, and is covered all over with strong and sharp spines or prickles, which he erects when irritated. He is no more than ten inches in length; and, by rolling himself up in the shape of a ball, he can resist the attack of his strongest enemies, among which are the cat, the weasel, the marten, and the dog. His common food consists of worms, slugs, and snails; and thus, far from being a noxious animal in a garden, he is a very useful one, as he feeds upon all the insects he can find. Hedgehogs inhabit most parts of Europe. Notwithstanding its formidable appearance, it is one of the most harmless animals in the world. While other creatures trust to their force, their cunning, or their swiftness, this quadruped, destitute of all, has but one expedient for safety, and from this alone it generally finds protection. The instant it perceives an enemy, it withdraws all its vulnerable parts, rolls itself into a ball, and presents nothing to view but a round mass of spines, impervious on every side. When the Hedgehog is thus rolled up, the cat, the weasel, the ferret, and the marten, after wounding themselves with the prickles, quickly decline the combat; and the dog himself generally spends his time in empty menaces rather than in effectual efforts, while the little

animal waits patiently till its enemy, by retiring, affords an opportunity for retreat.

The female produces from two to four young ones at a birth. When first born they are blind, and their spines white and soft, but they become hard in a few days. The Hedgehog is said to suck the milk from cows; but this is impossible, as the mouth of the Hedgehog would not admit the teat of the cow. The Hedgehog, however, sometimes destroys eggs, and has been known to attack frogs, mice, and even toads when pressed by hunger; it will also occasionally eat the tuberous roots of plants, boring under the root, so as to devour it, and yet leave the stem and leaves untouched. The Hedgehog makes himself a nest of leaves and soft wool for the winter, in the hollow trunk of an old tree, or in a hole in a rock or bank; and here, having coiled himself up, he passes the winter in one long unbroken sleep. Hedgehogs may easily be tamed, and are sometimes kept in the kitchens in London houses to destroy the black-beetles. The flesh of the Hedgehog is sometimes eaten; especially by gipsies, who appear to consider it a delicacy. It is said to be well-tasted, and to have abundance of yellow fat.






THE MOLE. (*Talpa Europaea.*)

THE Mole is a curious, awkwardly-shaped animal, with a long flexible snout, very small eyes, and four hand-like feet, armed with very strong claws, with which it scrapes its way through the ground, when it is forming the subterranean passages in which it takes up its abode. The Mole, though it is supposed not to possess the advantage of sight, has the senses of hearing and feeling in great perfection; and its fur, which is short and thick, is set erect from its skin, so as not to impede its progress whether it goes forward or backwards along its runs. These runs are very curiously constructed: they cross each other at different points, but all lead to a nest in the centre, which the Mole makes his castle, or place of abode. The passages are made by the Mole in his search after the earth-worms and grubs, on which he lives; and the molehills are formed by the earth he scrapes out of his runs. These molehills do a great deal of mischief to grass lands, as they render the ground very difficult to mow; and on this account mole-catchers are employed to fix traps in the ground, so that when the mole

is running through one of his passages, he passes through the trap, which instantly springs up out of the ground with the poor Mole in it. The female Mole makes her nest at a distance from the male's castle. She has young only once a year, but she has four or five at a time.

The following curious fact respecting a Mole is related by Mr. Bruce. "In visiting the Loch of Clunie, I observed in it a small island, at the distance of a hundred and eighty yards from the land. Upon this island Lord Airley, the proprietor, had a castle and small shrubbery. I observed frequently the appearance of fresh molehills; but for some time took it to be the water mouse, and one day I asked the gardener if it was so. He replied it was the Mole, and that he had caught one or two lately: but that five or six years ago he had caught two in traps, and for two years after this he had observed none. But about four years since, coming ashore one summer's evening in the dusk, he and Lord Airley's butler saw, at a small distance upon the smooth water, an animal paddling to and not far distant from the island; they soon closed with the feeble passenger, and found it to be the common Mole, led by a most astonishing instinct from the nearest point of land, (the castle-hill,) to take possession of this island. It was at this time for about the space of two years, quite free from any subterraneous inhabitant; but the Mole has, for more than a year past, made its appearance again."

The Mole is very pugnacious, and sometimes two of the males will fight furiously till one of them is killed.



THE SHREW. (*Sorex araneus.*)

THIS curious little animal closely resembles a mouse, except in its snout, which is long and pointed, to enable it to grub in the ground for its food, which consists of earth-worms, and the grubs of beetles. Like the mole and the hedgehog, it is soon famished for want of food; but the hedgehog has the advantage of being able to sleep for several months at a time, which the Shrew does not possess, and, consequently, great numbers of the latter animal die every year in autumn, when their food becomes scarce. The Shrew, like the mole, is very fond of fighting; and when two are seen together, they are generally engaged in a furious battle. Like the hedgehog, it has been much scandalized by false reports, as will be seen by the following extract from that most amusing and interesting work, *White's Selborne*: "At the south corner of the area, near the church, there stood, about twenty years ago, a very old, grotesque, hollow pollard-ash, which for ages had been looked upon with no small veneration as a shrew-ash. Now a shrew-ash is an ash whose twigs and branches, when applied to the limbs of cattle, will immediately relieve the pains which a beast

suffers from the running of a Shrew-mouse over the part affected; for it is supposed that a Shrew-mouse is of so baneful and deleterious a nature, that whenever it creeps over a beast, be it a horse, or cow, or sheep, the suffering animal is afflicted with cruel anguish, and threatened with the loss of the use of the limb. Against this accident, to which they were continually liable, our provident forefathers always kept a shrew-ash at hand, which, when once medicated, would maintain its virtue for ever. A shrew-ash was made thus:—into the body of the tree a deep hole was bored with an auger, and a poor devoted Shrew-mouse was thrust in alive, and plugged in." The cruelty of this, and many other practices of our ancestors, ought to make us thankful that we live in more enlightened days.

The body of the Shrew exhales a rank musky odour, which renders the animal so offensive to cats, that though they will readily kill them, they will not eat their flesh. This noisome odour probably gave rise to the notion that the Shrew-mouse is a venomous animal, and its bite dangerous to cattle, particularly horses. It is, however; neither venomous nor capable of biting, as its mouth is not sufficiently wide to seize the double thickness of the skin, which is absolutely necessary in order to bite.

The female Shrew makes her nest in a bank, or if on the ground, she covers it at the top, always entering on the side; and she has generally from five to seven young ones at a time.

The water Shrew (*Sorex fodiens*,) is a beautiful little creature, with somewhat differently formed feet and tail, to enable it to paddle through the water, in which it dives and swims with great agility. When floating "on the calm surface of a quiet brook," or diving after its food, its black velvety coat becomes silvered over with the innumerable bubbles of air that cover it when submerged; though when it rises again, the fur is observed to be perfectly dry, repelling the water as completely as the feathers of a water-fowl.


SECTION III.—Chiropterous Animals.

THE BAT. (*Vespertilio noctula.*)

THE Bat has the body of a mouse, and the wings of a bird. It has an enormous mouth, and very large ears, which are of a kind of membrane, thin and almost transparent. The pinions of its wings are furnished with hooks, by which it hangs to trees or the crevices in old walls during the day, a great number of them together, as they only fly at night. The wings of the Bat are very large; those of the Great Bat measuring fifteen inches across. It feeds on insects of various kinds, particularly on cockchafers and other winged beetles, part of which, however, it always throws away. A female Bat that was caught, and kept in a cage, ate meat when it was given to her in little bits, and lapped water like a cat. She was very particular in keeping herself clean, using her hind feet like a comb, and parting her fur so as to make a straight line down the back. Her wings she cleaned by thrusting her nose into the folds, and shaking them. She had a young one born in the cage. It was blind, and quite destitute of hair, and its mother wrapped it in the membrane of her wing, pressing it so closely to her breast, that no one could see her suckle it. The next day the poor mother died, and

the little one was found alive, hanging to her breast. It was fed with milk from a sponge, but only lived about a week.

SECTION IV.—The Marsupialia, or Pouch-bearing Animals.



THE KANGAROO. (*Macropus major*.)

THIS remarkable animal was first discovered by the celebrated Captain Cook, in New Holland; and as it was the only quadruped discovered on the island by the first settlers, they attempted to hunt it with greyhounds. The astonishing leaps it took, however, quite puzzled the colonists, who found it extremely difficult to catch. At first it was supposed that there was only one kind of Kangaroo, but now thirty or forty have been discovered, some of them not larger than a rat, and others as big as a calf. Kangaroos live in herds; one, older and larger than the rest, appearing to act as a kind of king. The ears of the Kangaroo are large, and in almost constant motion; it has a hare-lip, and a very small head. The fore legs, or rather paws, are short and weak, with five toes, each ending in a strong curved claw. The hind legs, on the contrary, are very large

and strong, but the feet have only four toes, and much weaker claws. The tail is very long and tapering; but is so thick and strong near the body, that it forms a kind of third hind leg, and wonderfully assists the animal in making its extraordinary leaps, which are often from twenty to thirty feet in length, and six or eight feet high. When the animal is attacked, it uses its tail as a powerful instrument of defence, and also scratches violently with its hind feet. It generally sits upright, but brings its fore feet to the ground when it is grazing. It lives entirely on vegetable substances. The most curious part of the Kangaroo is the pouch which the female has in front for carrying her young. It is just below her breasts, and the young ones sit there to suck; and even when they are old enough to leave the pouch, take refuge in it whenever they are alarmed.

The Kangaroo is easily tamed, and there are many in a tame state in England. In Australia, Kangaroo beef, as it is called, is eaten, and found very nourishing; but is hard and coarse. The female has generally two young ones at a time, which do not attain their full growth until they are a year old.

When a large Kangaroo is pursued by dogs, it generally takes refuge in a pond, where, from the great length of its hind legs and tail, it can stand with its body half out of the water, while the dogs are obliged to swim. Thus, the Kangaroo has a decided advantage; for, as each dog approaches him, he seizes it with his fore paws, and holds it under water, shaking it furiously till the dog is almost suffocated, and very glad to sneak off as soon as the Kangaroo lets him go.

The female, when pursued and hard-pressed by the dogs, will, while making her bounds, put her fore paws into her pouch, take a young one from it, and throw it as far out of sight as she possibly can. But for this manœuvre, her own life, and that of her young one would be sacrificed; whereas, she frequently contrives to escape, and returns afterwards to seek for her offspring.



THE VIRGINIAN OPOSSUM.

(*Didelphis Virginiana.*)


THIS creature, which is a native of North America, is about the size of a cat, and its fur is of a dingy white, except the legs, which are brown, and the nose and ears, which are yellowish. There is also a brownish circle round each eye, and the ears are nearly black at the base.

The Opossum generally lives in trees, suspending itself by the tail, by means of which it swings from branch to branch. In this manner it catches the insects and small birds, on which it generally feeds; but sometimes it descends from the tree, and invades poultry-yards, where it devours the eggs, and sometimes the young fowls. It resembles the kangaroo in its pouch for carrying its young, but in no other particular, as it walks on four feet, and its legs are uniform in length; and it has a long flexible tail, which is of no use to it either in leaping, or as a weapon of defence. The tail is, however, of singular use to the young, as when they get too large to be carried in the pouch, they fly to their mother when alarmed, and twisting their long slender

tails round hers, leap upon her back. The female Opossum may be sometimes seen thus carrying four or five at once.

The Opossum may be easily tamed, but is an unpleasant inmate, from its awkward figure and stupidity, and its very disagreeable smell. The American Indians spin its hair and dye it red, and then weave it into girdles and other articles of clothing. The flesh of these animals is white and well tasted, and is preferred by the Indians to pork: that of the young ones eats very much like the sucking-pig.

SECTION V.—Rodentia, or Gnawing Animals.



THE BEAVER. (*Castor Fiber.*)

THE Beaver is about the size of the badger; his head short, his ears round and small, his two fore teeth long, sharp, and strong, and well calculated for the part which Nature has allotted him: the tail is of an oval form, and covered with a scaly skin.

Beavers are natives of North America, and more particularly the north of Canada. Their houses are constructed with earth, stones, and sticks, neatly arranged and

worked together by their paws. The walls are about two feet thick, and are surmounted by a kind of dome, which generally rises about four feet above them. The entrance is on one side, always at least three feet below the surface of the water, so as to prevent it being frozen up. The number of Beavers in each house is from two to four old ones, and about twice as many young. When Beavers form a new settlement, they build their houses in the summer; and then lay in their winter provisions, which consist principally of bark and the tender branches of trees, cut into certain lengths, and piled in heaps on the outside of their habitation, and always under the water; though sometimes the heap is so large as to rise above the surface. One of these loads will occasionally contain more than a cart-load of bark, young wood, and the roots of the water-lily.

Beavers are hunted for the sake of their skins, which are covered with long hairs, and a short thick fur beneath, which is used in making hats, after the long hairs have been destroyed.

A great many stories have long been believed respecting the Beaver, on the authority of a French gentleman who had resided a long time in North America; but it is now ascertained that the greater part of them are false. The house of the Beaver is not divided into rooms, but consists of only one apartment; and the animals do not use their tails either as a trowel or a sledge, but only as an assistance in swimming. Some years ago a Beaver was brought to this country from America, that had been quite tamed by the sailors, and was called Bunney. When he arrived in England, he was made quite a pet of, and used to lie on the hearth-rug in his master's library. One day he found out the housemaid's closet, and his building propensities began immediately to display themselves. He seized a large sweeping brush, and dragged it along with his teeth to a room where he found the door open; he afterwards laid hold of a warming-pan in the same manner; and having laid the handles across, he filled up the walls of the angle made by the brushes with the wall,

with hand-brushes, baskets, boots, books, towels, and anything he could lay hold of. As his walls grew high, he would often sit, propped up by his tail, (with which he supported himself admirably,) to look at what he had done; and if the disposition of any of his building-materials did not satisfy him, he would pull part of his work down, and lay it again more evenly. It was astonishing how well he managed to arrange the incongruous materials he had chosen, and how cleverly he contrived to remove them, sometimes carrying them between his right fore paw and his chin, sometimes dragging them with his teeth, and sometimes pushing them along with his chin. When he had built his walls, he made himself a nest in the centre, and sat up in it, combing his hair with the nails of his hind feet.



THE HARE. (*Lepus timidus*.)

THIS small quadruped is well known at our tables as affording a favourite food, notwithstanding the dark colour of its flesh. Its swiftness cannot save it from the search of its enemies, among whom man is the most inveterate. Unarmed and fearful, the Hare appears almost to sleep


with open eyes, so easily is it alarmed. Its hind legs are longer than its fore ones, to enable it to run up hills; its eyes are so prominently placed, that they can encompass at once the whole horizon of the plain where it has chosen its form, for so its seat or bed is called; and its ears so long, that the least noise cannot escape it. It seldom outlives its seventh year, and breeds plentifully. Naturally wild and timorous, the Hare may, however, be occasionally tamed. The following is from the entertaining account given by Cowper, of three Hares that he brought up tame in his house; the names he gave them were Puss, Tiney, and Bess. Tiney was a reserved and surly Hare; Bess, who was a Hare of great humour and drollery, died young. "Puss grew presently familiar, would leap into my lap, raise himself upon his hinder feet, and bite the hair from my temples. He would suffer me to take him up and carry him about in my arms, and has more than once fallen fast asleep upon my knee. He was ill three days, during which time I nursed him, kept him apart from his fellows that they might not molest him, (for, like many other wild animals, they persecute one of their own species that is sick,) and by constant care, and trying him with a variety of herbs, restored him to perfect health. No creature could be more grateful than my patient after his recovery, a sentiment which he most significantly expressed by licking my hand, first the back of it, then the palm, then every finger separately, then between all the fingers, as if anxious to leave no part of it unsaluted; a ceremony which he never performed but once again upon a similar occasion.

"Finding him extremely tractable, I made it my custom to carry him always after breakfast into the garden, where he hid himself generally under the leaves of a cucumber vine, sleeping or chewing the cud, till evening; in the leaves also of that vine he found a favourite repast. I had not long habituated him to this taste of liberty, before he began to be impatient for the return of the time when he might enjoy it. He would invite me to the garden by drumming upon my knee, and by a look of such expression as it was not possible to misinterpret. If

this rhetoric did not immediately succeed, he would take the skirt of my coat between his teeth, and pull at it with all his force. Thus Puss might be said to be perfectly tamed, the shyness of his nature was done away, and, on the whole, it was visible, by many symptoms, which I have not room to enumerate, that he was happier in human society than when shut up with his natural companions."

Hares are included in the list of animals called game, and are hunted with greyhounds, which is called coursing; and also by packs of dogs called harriers and beagles. There are white Hares in the northern regions, the change in colour being the effect of cold.






THE RABBIT. (*Lepus cuniculus.*)

THIS animal, in a wild state, resembles the hare in all its principal characters, but is distinguished from it by its smaller size, the comparative shortness of the head and hinder legs, the gray colour of the body, the absence of the black tip to the ears, and the brown colour of the upper part of the tail. Its habits, however, are very different, as being from its organization unable to outstrip its enemies in the chase, it seeks its safety and shelter by burrowing in the ground; and instead of leading a solitary life, its manners are eminently social. Its flesh is white and good, though not so much prized as that of the hare.

The female begins to breed when she is about twelve months old, and bears at least seven times a year, generally eight at each time; now supposing this to happen regularly, a couple of Rabbits at the end of four years might see a progeny of almost a million and a half! Fortunately their destruction by various enemies is in proportion to their fecundity, or we might justly apprehend being overstocked by them. The young are born blind, and almost destitute of hair; while those of the hare can see, and are covered with hair.



THE DOMESTIC RABBIT.

THE Domestic Rabbit is larger than the wild species, owing to its taking more nourishment, and less exercise; (our example, however, is drawn disproportionately large.) Like pigeons, they have their regular fanciers, and are bred of various colours; grey, reddish-brown, black more or less mixed with white, or perfectly white. The ears are considered to constitute a principal feature of their beauty, and the animal is most valued when both ears hang down by the side of the head; the animal is then called a double lop; when only one ear drops, it is called a single or horn lop, and when both stretch out horizontally, an oar-lop. Unlike the wild Rabbit, they do not burrow in the ground.



THE SQUIRREL. (*Scirus vulgaris.*)

ELEGANCE of shape, spiritedness, and agility to leap from bough to bough in the forest, are the principal characteristics of this pretty animal. The Squirrel is of a deep reddish brown colour; his breast and belly white. He is lively, sagacious, docile, and nimble: he lives upon nuts, and has been seen so tame as to dive into the pocket of his mistress, and search after an almond or a lump of sugar. In the woods, he leaps from tree to tree with surprising agility, living a most frolicsome life, surrounded with abundance, and having but few enemies. His time, however, is not entirely devoted to idle enjoyment, for in the luxuriant season of autumn, he gathers provisions for the approaching winter, as if conscious that the forest would then be stripped of its fruits and foliage. His tail serves him as a parasol to defend him from the rays of the sun, as a parachute to secure him from dangerous falls when leaping from tree to tree, and as a sail in crossing the water, which he sometimes does in Lapland on a bit of ice or bark inverted in the manner of a boat.

The American Flying Squirrel (*Pteromys volucella*) has a large membrane proceeding from the fore feet to the hind legs, which answers the same purpose as the Squirrel's tail, and enables him to give surprising leaps that almost resemble flying. In the act of leaping, the loose skin is stretched out by the feet; whereby the surface of

the body is augmented, the animal becomes lighter in proportion to its bulk, its fall is retarded, and it appears to sail or fly from one place to another. Where numbers of them are seen at a time leaping, they appear like leaves blown off by the wind.



THE MARMOT, OR ALPINE RAT.

(*Arctomys Marmotta.*)

THIS is a harmless inoffensive animal, and seems to bear enmity to no creature but the dog. He is caught in Savoy, and carried about in several countries for the amusement of the mob. When taken young, he is easily tamed, and possesses great muscular power and agility. He will often walk on his hinder legs, and uses his fore paws to feed himself, like the squirrel. The Marmot makes his hole very deep, and in the form of the letter Y, one of the branches serving as an avenue to the innermost apartment, and the other sloping downwards, as a kind of sink or drain; in this safe retreat he sleeps throughout the winter, and if discovered may be killed without appearing to undergo any great pain. These animals produce but once a year, and bring forth three or four at a time. They grow very fast, and the extent of their lives is not above nine or ten years. They are about the size of a hare or cat, but much more corpulent. When a number of Marmots are feeding together, one of them stands sentinel upon a wall; and on the first appearance of a man, a dog, an eagle, or any dangerous animal, utters a loud and shrill cry, as a signal for immediate retreat. The Marmot inhabits the highest regions of the Alps; and is likewise found in Poland, Ukraine, and Chinese Tartary.



THE GUINEAPIG. (*Cavia Cobaya.*)

THIS animal is generally white, variegated with red and black. It is a native of the Brazils, but now domesticated in most parts of Europe, and is about the size of a large rat, though more stoutly made and without any tail; and its legs and neck are so short, that the former are scarcely seen, and the latter seems stuck upon its shoulders. Guinea-pigs, though they have a disagreeable smell, are extremely cleanly, and the male and female may be often seen alternately employed in smoothing each other's skins, disposing their hair, and improving its gloss. They sleep like the hare with their eyes half open, and continue watchful if they apprehend any danger. They are very fond of dark retreats; previously to their quitting which, they look round, and seem to listen attentively; then, if the road be clear, they sally forth in quest of food; but run back on the slightest alarm. They utter a sound like the snore of a young pig. The female begins to produce young, when only two months old, and as she does so every two or three months, and has sometimes as many as twelve at a time, a thousand might be raised from a single pair in the course of a year. They are naturally gentle and tame; as incapable of mischief as they seem to be of good, although rats are said to avoid their locality. The upper lip is only half divided; it has two cutting teeth in each jaw, and large and broad ears. They feed on bread, grain, vegetables, and tea-leaves.



THE MOUSE. (*Mus musculus.*)

THIS is a lively, active animal, and the most timid in nature, except the hare, and a few other defenceless species. Although timid, he eats in the trap as soon as he is caught; yet he never can be thoroughly tamed, nor does he betray any affection for his assiduous keeper. He is beset by a number of enemies, among which are the cat, the hawk, and owl, the snake and weasel, and the rat himself, though not unlike the mouse in his habits and shape. The mouse is one of the most prolific of animals, sometimes producing seventeen at a birth; but it is supposed that the life of this small inmate of our habitations does not extend much further than three years. This creature is known all over the world, and breeds wherever it finds food and tranquillity. There are Mice of various colours, but the most common kind is of a dark, cinereous hue: white Mice are not uncommon, particularly in Savoy and some parts of France.

A remarkable instance of sagacity in a long-tailed Field Mouse, occurred to the Rev. Mr. White, as his people were pulling off the lining of a hot-bed, in order to add some fresh dung. From the side of this bed something leaped with great agility, that made a most grotesque appearance, and was not caught without much difficulty. It proved to be a large Field Mouse, with three or four

young ones clinging to her teats by their mouths and feet. It was amazing that the various and rapid motions of the dam did not oblige her litter to quit their hold, especially when it appeared that they were so young as to be both naked and blind. Mr. White appears to be the first to describe and accurately examine that diminutive creature the Harvest Mouse, the least of all the British quadrupeds. He measured some of them, and found that from the nose to the tail they were two inches and a quarter long. Two of them in a scale only weighed down one copper halfpenny, about the third of an ounce avoirdupoise! Their nest is a great curiosity.




THE RAT. (*Mus decumanus*.)

THE Rat is about four times as large as the mouse, but of a dusky colour, with white under the body; his head is longer, his neck shorter, and his eyes comparatively larger. These animals are so attached to our dwellings, that it is almost impossible to destroy the breed, when

they have once taken a liking to any particular place. Their produce is enormous; as they have from ten to twenty young ones at a litter, and this thrice a year. Thus their increase is such, that it is possible for a single pair (supposing food to be sufficiently plentiful, and that they had no enemies to lessen their numbers) to amount, at the end of two years, to upwards of a million; but an insatiable appetite impels them to destroy each other; the weaker always fall a prey to the stronger; and the large male Rat, which usually lives by itself, is dreaded by those of its own species as their most formidable enemy. The Rat is a bold and fierce little animal, and when closely pursued, will turn and fasten on its assailant. Its bite is keen, and the wound it inflicts is painful and difficult to heal, owing to the form of its teeth, which are long, sharp, and of an irregular form.

It digs with great facility and vigour, making its way with rapidity beneath the floors of our houses, between the stones and bricks of walls, and often excavating the foundations of a dwelling to a dangerous extent. There are many instances of their totally undermining the most solid mason-work, or burrowing through dams which had for ages served to confine the waters of rivers and canals.

A gentleman, some time ago, travelling through Mecklenburgh, was witness to a very singular circumstance respecting one of these animals, in the post-house at New Hargarell. After dinner, the landlord placed on the floor a large dish of soup, and gave a loud whistle. Immediately there came into the room a mastiff, an Angora cat, an old raven, and a large Rat with a bell about its neck. They all four went to the dish, and without disturbing each other, fed together; after which, the dog, cat, and Rat lay before the fire, while the raven hopped about the room. The landlord, after accounting for the familiarity which existed among these animals, informed his guest that the Rat was the most useful of the four; for that the noise he made had completely freed the house from the Rats and mice with which it had been before infested.



THE WATER RAT (*Arvicola amphibius*)

DIFFERS very little from the *Land Rat*, and inhabits the banks of rivers and ponds, where he digs holes always above the water-mark, and feeds on fish, frogs, and insects.

Water Rats live near rivers and ponds in summer, but when winter approaches frequent the farm-house, burrow in the corn-fields, infest the mows and stacks, attack the poultry-yard, and commit vast depredations.

This animal is nearly as large as the brown Rat, but has a larger head, a blunter nose, and smaller eyes; its ears are very short, and almost hid in the fur, and the tip of its tail is whitish; the cutting-teeth are of a deep yellow colour in front, very strong, and much resembling those of the beaver. Its head and back are covered with long black hair, and its belly with iron gray. Tail more than half the length of the body, covered with hairs. Fur thick and shining; of a rich reddish brown, mixed with gray above, yellowish gray beneath. The female produces a brood of five or six young ones once (and sometimes twice) a year.



THE MUSK RAT (*Fiber zibethicus*)

Is a native of Canada, and resembles the beaver in many of his habits. He has a fine musky scent, and makes his holes in marshes, and by the waterside, with two or three ways to get in or go out, and several distinct apartments: he is said to contrive one entrance to his hole always below the water, that he may not be frozen out by the ice. He lives upon small fish and water insects, as well as vegetables. This animal is called the Musquash in America, and its fur is used, like that of the beaver, in the manufacture of hats, four or five hundred thousand skins being said to be sent to Europe every year for that purpose. Musk Rats are always seen in pairs; and though watchful, are not timid, as they will often approach quite close to a boat or other vessel. In spring they feed on pieces of wood, which they peel carefully; and they are particularly fond of the roots of the sweet flag (*Acorus Calamus*). In Canada this animal is called the Ondrata.



THE DORMOUSE, OR SLEEPER. (*Myoxis glis*.)

THESE animals build their nests either in the hollow parts of trees, or near the bottom of thick shrubs, and line them

upon its legs, would give us the nearest resemblance to the figure of a Jerboa when pursued. It uses, however, all its four feet upon ordinary occasions, and it is only when pursued that it presses its fore feet close to its body, and leaps on its hind ones. The ancients called it the two-footed rat. This creature is about the size of a rat; the head resembles that of a rabbit with long whiskers; the tail is ten inches long, and terminated by a tuft of black hair. The fur of the body is tawny, except the breast and throat, and part of the belly, which are white. The Jerboa is very active and lively, and jumps and springs, when pursued, six or seven feet from the ground, with the assistance of its tail; but if this useful member be in any manner injured, the activity of the Jerboa is proportionately diminished; and one which had been accidentally deprived of its tail, was found unable to leap at all. The Jerboa is supposed to be both the coney and the mouse of the Holy Scriptures. It burrows like the rabbit, and feeds like the squirrel: it is a native of Egypt, and all the adjacent countries.



most industriously with moss, soft lichens, and dead leaves. Conscious of the length of time they have to pass in their solitary cells, Dormice are very particular in the choice of the materials they employ to build and furnish them; and generally lay up a store of food, consisting of nuts, beans, and acorns; and on the approach of cold weather, roll themselves in balls, their tail curled over their head between the ears, and in a state of apparent lethargy, pass the greatest part of the winter, till the warmth of the sun, pervading the whole atmosphere, kindles their congealed blood, and calls them back again to the enjoyment of life. Except in the time of breeding and bringing up its young, the Dormouse is generally found alone in its cell. This animal is remarkable for the very small degree of heat its body possesses during its torpid state, when it appears actually frozen with the cold, and it may be tossed or rolled about without being roused, though it may be instantly revived by the application of gentle heat, such as that of the hands. If a torpid Dormouse, however, be placed before a large fire the sudden change will kill it.

The American Dormouse, or Ground Squirrel, is a very beautiful animal, striped down the back, and resembling the squirrel in its habits, except that instead of living in trees, it burrows in the ground.




THE JERBOA. (*Dipos Jerboa.*)

THE principal peculiarity of this animal consists in its having very short fore legs, and very long hinder ones: a bird divested of its feathers, and wings, and jumping

upon its legs, would give us the nearest resemblance to the figure of a Jerboa when pursued. It uses, however, all its four feet upon ordinary occasions, and it is only when pursued that it presses its fore feet close to its body, and leaps on its hind ones. The ancients called it the two-footed rat. This creature is about the size of a rat; the head resembles that of a rabbit with long whiskers; the tail is ten inches long, and terminated by a tuft of black hair. The fur of the body is tawny, except the breast and throat, and part of the belly, which are white. The Jerboa is very active and lively, and jumps and springs, when pursued, six or seven feet from the ground, with the assistance of its tail; but if this useful member be in any manner injured, the activity of the Jerboa is proportionately diminished; and one which had been accidentally deprived of its tail, was found unable to leap at all. The Jerboa is supposed to be both the coney and the mouse of the Holy Scriptures. It burrows like the rabbit, and feeds like the squirrel: it is a native of Egypt, and all the adjacent countries.





THE CHINCHILLA. (*Chincella lanigera.*)

THE Chinchilla is a native of America, and its coat produces the beautiful fur known by its name. This animal was but little known to naturalists until Captain Beechey brought home with him a specimen from the north-west coast of America, and presented it to the Zoological Society.

The length of the body of this little animal is about nine inches, and its tail nearly five; its limbs are comparatively short, the hind legs being much the longest. The fur is of a remarkably close and fine texture, somewhat crisped, and entangled together; of a grayish or ash colour above, and paler beneath. It is used for muffs, tippets, and linings of cloaks, and is perhaps prettier than the Sable, although less durable, and less valuable in commerce, excepting when fashion rules. The form of the head resembles that of the rabbit; the eyes are full, large, and black; and the ears broad, naked, round at the tips, and nearly as long as the head. The whiskers are plentiful and strong, the longest being twice as long as the head, some of them black, others white. Four short toes, with an appearance of a thumb, terminate the fore feet; the hinder have the same number of toes, but have less the appearance of hands: on all the claws are short, and nearly hidden by tufts of bristly hairs. The tail is about half the length of the body, of equal thickness throughout, and covered with long bushy hairs. It resembles in some degree the jerboa, and takes its food, like that animal, in its fore paws, sitting on its haunches. The temper of the Chinchilla is mild and tractable. It dwells in burrows under ground, and produces young twice a year, bringing forth five or six at a time. It feeds upon the roots of bulbous plants.

THE PORCUPINE. (*Hystrix cristata.*)

WHEN full grown this animal measures about two feet and a half in length, and his body is covered with hair and sharp quills, from ten to fourteen inches long, and bent backwards. When he is irritated, they stand erect; but the story that the Porcupine can shoot them at his enemies, is only one of the many fables formerly related as facts in Natural History. The female has only one young one at a time. It is reported to live from twelve to fifteen years. The Porcupine is dull, fretful, and inoffensive; it feeds upon fruits, roots, and vegetables; and inhabits the south of Europe, India, Persia, and almost every part of Africa; particularly Barbary.

Mr. Bewick, in his History of Quadrupeds, asserts, that, upon the smallest irritation, this animal erects its quills, and shakes them with great violence, directing them to the quarter whence it is in danger of being attacked. "We have observed," says he, "on an occasion of this sort, when the animal was moulting or casting its quills, that they would fly out to the distance of a few yards, with such force as to bend the points of them against the board where they struck: and it is not improbable that a circumstance of this kind may have given rise to an opinion of its power to use them in a more effectual manner."

SECTION VI.—Edentata, or toothless Animals.


THE SLOTH. (*Bradypus tridactylus*.)

THIS animal, which is sometimes also called Ai, in reference to a noise it makes when caught, is most curiously formed. The arms or fore legs are nearly twice as long as the hind legs: and consequently when the animal attempts to walk or rather crawl, it can only do so on its hind feet, and the elbows of its fore legs. The claws also are larger than the foot, and bent inwardly, so as to prevent the animal from placing the ball of its foot on the ground. From these peculiarities in its construction the progress of the Sloth on land is extremely slow and laborious; but he is not intended to be a terrestrial animal. It lives in trees, always hanging below the branch, with its back to the ground; and for a life of this kind, its long arms and hooked claws are admirably adapted. Mr. Waterton, whose long residence in the wilds of South America, and whose habits of close observation, render him an excellent authority, observes, that when the Sloth travels from branch to branch of the tree which

it, inhabits, particularly in windy weather, it moves with such rapidity as to make it quite a misnomer to call it a Sloth. "The Sloth," says Mr. Waterton, "in its wild state, spends its whole life in the trees, and never leaves them, but through force or accident; and what is more extraordinary, not *upon* the branches, like the squirrel and monkey, *but under them*. He moves suspended from the branch, he rests suspended from the branch, and he sleeps suspended from the branch. Hence his seemingly bungled composition is at once accounted for; and in lieu of the Sloth leading a painful life, and entailing a melancholy existence upon its progeny, it is but fair to conclude, that it enjoys life just as much as any other animal, and that its extraordinary formation and singular habits are but further proofs to engage us to admire the wonderful works of Omnipotence."

The common Sloth has always three toes; but there is another kind, called the Unan, which has only two toes, and much shorter fore legs.

The female Sloth has only one young one at a time, which hangs to her breast, and makes a kind of cradle of her body, during her journeys from branch to branch; in fact, it appears never to quit her, till it is able to provide for itself. When hanging from the branch, she hides her young one in her thick, matted hair, which resembles in texture and appearance dry withered grass, and, indeed, is so like the rough bark and moss on old trees, as to render the animal scarcely distinguishable. When the Sloth has got possession of a tree, it will not descend while a leaf or bud is remaining; and it is asserted, that in order to obviate the necessity of a slow and laborious descent, it suffers itself to fall to the ground; the toughness of its skin and the thickness of its hair securing it from any unpleasant consequences.



THE ARMADILLO. (*Dasyus sexcinctus.*)

NATURE seems to have been singularly careful in the preservation of this animal, for she has surrounded it with a strong coat of armour to protect it from its enemies. When closely pursued, it withdraws its head under the shell, and assumes the shape of a ball; and if near a precipice, rolls from one rock to another, and escapes without receiving any injury. The shell, which covers the whole of the body, is composed of several scales, very hard, and of a square shape, united by a kind of cartilaginous substance, which gives flexibility to the whole. The Armadillo lives principally on roots, carrion, and ants; and in a wild state resides in subterranean burrows, like the rabbit. It is a native of South America. There are several species differing chiefly in the number of their bands. When naturalists wish to obtain a specimen of the Armadillo in its native country, they are obliged to employ an Indian to dig one out of its hole; and as the holes are almost innumerable, only a few of them containing Armadillos, the Indians try them first by putting a stick down, when, if a number of musquitos rise, the Indians know the hole contains an Armadillo, as, if there were none, there would be no musquitos.



THE GREAT ANT-EATER. (*Myrmecophaga jubata*.)

THE body of the Great Ant-eater is covered with exceedingly coarse and shaggy hair. Its head is very long and slender, and the mouth but just large enough to admit its tongue, which is cylindrical, nearly two feet in length, and lies folded double within it. The tail is of enormous size, and covered with long black hair, somewhat like the tail of a horse. The whole length of the animal, from the end of the snout to the tip of the tail, is sometimes eight or nine feet. Its food consists principally of ants, which it obtains in the following manner:—When it comes to an ant-hill, it scratches it up with its long claws, and then unfolds its slender tongue, which much resembles an enormously long worm. This being covered with a glutinous matter or saliva, the ants adhere to it in great numbers: these it swallows alive, repeating the operation till no more are to be caught.

He also tears up the nests of wood-lice, which it in like manner discovers; and frequently climbs trees in quest of these insects, and of wild bees or their honey. But should it meet with little success in its pursuit of food, it is able to fast for a considerable time without inconvenience. The motions of the Ant-eater are in general very slow. It swims, however, over great rivers with ease; and on these occasions, its tail is always

thrown over its back. With this extraordinary member, when asleep, or during heavy showers of rain, the animal also covers its back; but at other times he trails it along the ground. The Ant-eater is a native of South America.



THE DUCK-BILLED PLATYPUS, OR WATER MOLE. (*Ornithorhynchus paradoxus*.)


THIS extraordinary creature has the bill and webbed feet of a duck, united to the body of a mole. It is a native of Australia, where it is found on the banks of rivers, in the sides of which it burrows and forms its nest. It feeds on aquatic insects and small molluscous animals, always, however, rejecting the shells of the latter, after crushing them in its mouth, so as to extract the body. A number of these animals are always found together, but it is very difficult to watch their habits, as their sense of hearing is so acute, that they disappear at the slightest noise, plunging into the water, in which they swim so low, that they only look like a mass of weeds floating on the surface.

These curious animals produce their young in eggs, which are hatched the very moment they are laid; and

they nourish them with milk, which oozes through the fur from the chest of the female, though she has no teats. When the animal feeds, he plunges his beak into the mud, just like a duck; and appears to be equally at home on land and in water. Two young ones that were kept for some time at Sidney, by Mr. Bennet, were very fond of rolling themselves up like a hedgehog, in the form of balls. They often slept in this position, and "awful little growls" issued from them when disturbed. They were fed with worms and bread and milk; but captivity did not seem to agree with them, and they soon died. They dressed their fur by combing it with their feet, and pecking at it with their beaks, seeming to take great delight in keeping it smooth and clean.

The shape of this animal is so extraordinary, that when a specimen was first sent to Europe it was supposed to have been manufactured by fixing the beak of a duck into the head of some small quadruped, with the intention to deceive. Subsequent experience has proved, beyond the possibility of a doubt, the existence of the animal, without in the smallest degree diminishing the wonder excited by its first appearance, as it partakes in almost equal parts of the nature of quadrupeds, birds, and reptiles.

The Australian hedgehog (*Echidna setosa*) has, like the Duck-billed Platypus, a spur on its hind legs, which contains a poisonous venom, similar to the tooth of a serpent. It has a long very slender muzzle, resembling the bill of a snipe, at the end of which is a very small mouth, containing a long tongue, which the creature can extend at pleasure. The body is short and rounded: it is covered with strong sharp spines mixed with hair; and its tail is so short that it was at first doubted whether it had one.



SECTION VII.—*Pachydermata*, or thick-skinned
Animals.

THE ELEPHANT. (*Elephas Indicus*.)

PROVIDENCE, always impartial in the distribution of its gifts, has given this bulky quadruped a quick instinct nearly approaching to reason, in compensation for the uncouthness of his body. The Ceylon Elephant is about thirteen feet high, and seven broad, and is much the largest of all living animals. His skin is in general a mouse colour, but is sometimes white and sometimes black. His eyes are rather small for the size of his head, and his ears, which are very expanded and of a peculiar shape, have the flaps hanging down, instead of standing up, as in most quadrupeds. The Elephant is a gregarious animal in his wild state, and when domesticated is susceptible of attachment and gratitude, as well as of anger and revenge. Several anecdotes are related of his quick ap-

prehension, and particularly of his vindictive treatment of those who have either scoffed at or abused him. To disappoint him is dangerous, as he seldom fails to be revenged. The following instance is given as a fact, and deserves to be recorded:—An Elephant, disappointed of his reward, out of revenge, killed his governor. The poor man's wife, who beheld the dreadful scene, took her two children and thrust them towards the enraged animal, saying, "Since you have slain my husband, take my life also, as well as those of my children!" The Elephant instantly stopped, relented, and, as if stung with remorse, took the eldest boy in his trunk, placed him on his neck, adopted him for his governor, and would never afterwards allow any other person to mount him.

The Elephant's mouth is armed with broad and strong grinding teeth, and two large tusks, which measure sometimes nine or ten feet, and from which the finest ivory is produced. The ivory from the tusks of the female is thought the best, as the tooth, being smaller, admits less porosity in the cellular part of the mass.


Becoming tame under the mild treatment of a good master, the Elephant is not only a most useful servant, for the purposes of state or war, but is also of great assistance in taming the wild ones that have been recently caught. Hindostan superstition has paid great honours to the white race of this quadruped; and the island of Ceylon is supposed to breed the finest of the kind. This immense beast, by the wisdom of Providence, has not been placed among the carnivorous animals: and vegetable food being much more abundant than animal, he is destined to live on grass and the tender shoots of trees. This noble creature bears in state on his back the potentates of the East, and seems to delight in pompous pageantry: in war he carries a tower filled with archers; and in peace lends his assistance in domestic operations. The female is said to go a year with young, and to bring forth one at a time. The Elephant lives a hundred and twenty or a hundred and thirty years, though they have been known to live to the great age of four hundred. When Alexander the Great had

conquered Porus, King of India, he took a large Elephant which had fought very valiantly for the king, and naming him Ajax, dedicated him to the sun, and then let him loose with this inscription:—"Alexander, the son of Jupiter, hath dedicated Ajax to the sun." This Elephant was found with this inscription 350 years after.

The greatest wonder the Elephant presents to the admiration of the intelligent observer of nature is his *proboscis*, or trunk, composed of elastic rings to the extent of six or eight feet, and so flexible that he uses it as dexterously as a man does his hand. It was erroneously said, that the Elephant could receive nourishment through his trunk; this sort of pipe is nothing but a prolongation of the snout, for the purpose of breathing, into which the animal can, by the strength of his lungs, draw up a great quantity of water or other liquid, which he spouts again, or brings back to his mouth by inverting and shortening his proboscis for this purpose.

Captain Marryat, in his very entertaining work called *Masterman Ready*, relates a curious instance of the sagacity of an Elephant in India, which had fallen into a deep tank. The tank was so deep that it was impossible to hoist the Elephant up, but when the people threw down several bundles of faggots, the sagacious animal laid one bundle above another, always standing on each tier as he arranged it, till at last he raised the pile high enough to allow him to walk out of the tank. But instances of the sagacity of this noble creature might be cited ad infinitum. In the East, where they are made available in the service of man, they will load a boat with singular dexterity, carefully keeping every article dry, and disposing and balancing the cargo with the utmost precision.

Its strength is proportionate to its bulk: it will carry three or four thousand pounds weight on its back, and upwards of a thousand pounds on its tusks.



THE HIPPOPOTAMUS, OR RIVER-HORSE.

(*Hippopotamus amphibius.*)

THIS animal lives as well on land as in water, and yields in size to none but the elephant: he weighs sometimes more than fifteen hundred pounds. His skin is very sleek, and covered with short and soft hair, of a mouse colour. The head is flattish on the top, about four feet long and nine in circumference; the lips are large, the jaws open about two feet wide, and the cutting-teeth, of which it has four in each jaw, are about a foot long; he has broad ears, and large eyes, a thick neck, and a short tail, tapering like that of a hog. He grazes and eats the leaves and young branches of trees on shore, but retires to the water if pursued, and will sink down to the bottom, where he can remain five or six minutes at a time. When he rises to the surface and remains with his head out of the water, he makes a bellowing noise which may be heard at a great distance. The female brings forth her


young upon land, and it is supposed that she seldom produces more than one at a time. The calf at the instant that it comes into the world, flies to the water for shelter, if pursued; a circumstance which has been noticed as a remarkable instance of pure instinct. The Hippopotamus is supposed to be the Behemoth of the Scripture. See Job, chap. xl.



THE RHINOCEROS, (*Rhinoceros Indicus*.)

So called because of the horn on his nose, is bred in India and Africa, is of a dark slate-colour, and nearly as large as the elephant; as he measures about twelve feet in length, and has short legs. His skin, which is not penetrable by any ordinary weapon, is folded upon his body, in the manner represented in the figure above; his eyes are small and half closed, and the horn on his nose is attached to the skin only. In confinement he often wears it to a mere stump, by rubbing it against his crib. He is perfectly indocile and untractable; a natural enemy to the elephant, to whom he often gives battle, and is

said never to go out of his way, but to endeavour to destroy whatever obstacles present themselves, rather than turn about. He lives on the coarsest vegetables, and frequents the banks of rivers, and marshy grounds; his hoofs are divided into four, and he grunts like a hog, which he resembles in many other particulars. The female produces but one at a time, and during the first month her young are not bigger than a large dog. The Rhinoceros is supposed by some to be the Unicorn of holy writ, and possesses all the properties ascribed to that animal,—rage, untameableness, great swiftness, and immense strength. It was known to the Romans in very early times. Augustus introduced one into the shows, on his triumph over Cleopatra. The African Rhinoceros has two horns.




THE COMMON OR DOMESTIC HOG (*Sus scrofa*)

DIFFERS chiefly from the wild animal in having smaller tusks, and large and pendant ears. Of all domestic

quadrupeds, this is the most filthy and impure. Its form is clumsy and unsightly, and its appetite gluttonous and excessive. Nature, however, has fitted its stomach to receive nutriment from a variety of things that would be otherwise wasted, as the refuse of the field, the garden, and the kitchen afford it a luxurious repast. The Hog is naturally stupid, inactive, and drowsy; much inclined to increase in fat, which is disposed in a different manner from that of other animals, forming a thick, distinct, and regular layer between the flesh and skin. Their flesh, Linnæus observes, is a wholesome food for those that use much exercise, but improper for such as lead a sedentary life. It is of great importance to this country, as a naval and commercial nation, for it salts better than any other flesh, and is capable of being longer preserved.

The domestic Sow brings forth twice a year, producing from ten to twenty at a litter. She goes four months with young, and brings forth in the fifth. At that time she must be carefully watched, to prevent her from devouring her young. Still greater attention is necessary to keep off the male, as he would destroy the whole litter. Jews and Mahomettans not only abstain from the flesh of swine from a religious principle, but consider themselves defiled by even touching it.




THE WILD BOAR (*Sus scrofa*)

INHABITS, for the most part, marshes and woods, and is of a black or brown colour: his flesh is very tender and good for food. The Wild Boar has tusks, which are sometimes nearly a foot in length, and have often proved dangerous to men as well as to dogs in the chase. His life is confined to about thirty years; his food consists of vegetables, but when pressed by hunger, he devours animal flesh. This creature is strong and fierce, and undauntedly turns against his pursuers. To hunt him is one of the principal amusements of the grandees in those countries where he is to be found. The dogs provided for this sport are of the slow heavy kind. Those used for hunting the stag, or the roe-buck, would be very improper, as they would too soon come up with their prey, and instead of a chase, would only furnish an engagement. Small mastiffs are therefore chosen; nor do the hunters much regard the goodness of their nose, as the Wild Boar leaves so strong a scent that it is impossible for them to mistake its course. They never hunt any but the largest and the oldest, which are known by their tusks. When the boar is *reared*, as is the expression for driving him from his covert, he goes slowly and sullenly forward, without any indication of fear, not very far before his pursuers. At the end of every half-mile, or thereabouts, he turns round, stops till the hounds come up, and offers to attack them. These, on the other hand, knowing their danger, keep off and bay him at a distance. After they have for a while gazed upon each other, with mutual animosity, the Boar again slowly goes on his course,

and the dogs renew the pursuit. In this manner the charge is sustained, and the chase continues, till the Boar is quite tired, and refuses to go any further. The dogs then attempt to close in upon him from behind; those which are young, fierce, and unaccustomed to the chase, are generally the foremost, and often lose their lives by their ardour. Those which are older and better trained are content to wait until the hunters come up, who despatch him with their spears.

In former times, the Wild Boar was a native of Britain, as appears from the laws of the Welsh prince, Howell the Good, who permitted his grand huntsman to chase that animal from the middle of November to the beginning of December; and in the reign of William the Conqueror, those who were convicted of killing the Wild Boars, in any of the royal forests, were punished with the loss of their eyes. Our domestic pigs are descended from the wild race; but the tame Boar has two tusks, smaller than those of the wild ones, and the sow has none.



THE TAPIR. (*Tapir Americanus*.)

THIS animal bears considerable resemblance to the wild boar, but is without tusks, and has its snout prolonged into a small fleshy proboscis or trunk. This trunk, however, has not the flexibility of that of the elephant, and is incapable of holding anything. The colour of the Tapir is of a deep brown, and the male has a small mane on the upper part of his neck. It stands about three feet and a half high, and measures nearly six feet in length. It lies in thickets, the thorny branches of which cannot affect it from the thickness of its skin, while they lacerate the skins of its pursuers. Its favourite food is the water-melon. It is generally found alone, and always roams for prey at night; and it is easily tamed if taken young. It possesses the same power of remaining under water as the hippopotamus, and when it enters a pond, can descend to the bottom, and remain there five or six minutes.



THE HORSE. (*Equus caballus.*)

THE noblest conquest that man ever made over the brute creation was the taming of the Horse, and adapting him to his service. He lessens the labours of man and adds to his pleasures; shares, with equal docility and cheerfulness, the fatigues of hunting or the dangers of war; and draws with appropriate strength, rapidity, or grace, the heavy ploughs and carts of the husbandman, the light vehicles of the fashionable, and the stately carriages of the aristocratic.

The Horse is now bred in most parts of the world: those of Arabia, Turkey, and Persia are accounted better proportioned than many others; but the English Race-Horse may justly claim the precedence over all the other European breeds, and is not inferior to any in strength and symmetry.

The beautiful Horses produced in Arabia are in general of a brown colour; their mane and tail are very short, with the hair black and tufted. The Arabs for the most part use the Mares in their ordinary excursions; experience having taught them that they are less vicious than the


males, and more capable of sustaining abstinence and fatigue. As the Arabs have no other residence than a tent, this also serves for a stable; the husband, the wife, the child, the mare, and the foal lie together indiscriminately, and the younger branches of the family may be often seen embracing the neck, or reposing on the body of the Mare, without any idea of fear or danger.


Of the remarkable attachment which the Arabs have to these animals, St. Pierre has given an affecting instance in his *Studies of Nature*.—"The whole stock of a poor Arabian of the desert consisted of a beautiful Mare: this the French consul at Said offered to purchase, with an intention to send her to Louis XIV. The Arab, pressed by want, hesitated a long time, but at length consented, on condition of receiving a very considerable sum of money, which he named. The consul wrote to France for permission to close the bargain; and, having obtained it, sent the information to the Arab. The man, so indigent as to possess only a miserable covering for his body, arrived with his magnificent courser: he dismounted, and first looking at the gold, then steadfastly at his Mare, heaved a sigh. 'To whom is it,' exclaimed he, 'that I am going to yield thee up? To Europeans? who will tie thee close, who will beat thee, who will render thee miserable! Return with me, my beauty, my jewel! and rejoice the hearts of my children:' as he pronounced the last words, he sprung upon her back, and was out of sight almost in a moment."

The intelligence of the Horse is next to that of the elephant, and he obeys his rider with so much punctuality and understanding, that the Americans, who had never seen a man on horseback, thought, at first, that the Spaniards were a kind of centaurs, half men and half horses. The Horse, in a domestic state, seldom lives longer than twenty years; but it is supposed that in a wild state he attains a much greater age. The Mare is as elegant in her shape as the Horse; and her young is called a foal. The age of the Horse is known from his teeth; and his colour, which varies from black to white, and from the darkest brown to a light hazel tint, has been reckoned a criterion by which to judge of his strength.

The Horse feeds upon grass, either fresh or dry, and corn: he is liable to many diseases, and often dies suddenly. In the state of nature, he is a gregarious animal, and even when domesticated, his debased situation of slavery has not entirely destroyed his love of society and friendship; for Horses have been known to pine at the loss of their masters, their stable fellows, and even at the death of a dog which had been bred near the manger. Virgil, in his beautiful description of this noble animal, seems to have imitated Job:

“The fiery courser, when he hears from far
The sprightly trumpets, and the shouts of war,
Pricks up his ears, and trembling with delight,
Shifts place, and paws, and hopes the promised fight.
On his right shoulder his thick mane reclined,
Ruffles at speed, and dances in the wind.
His horny hoofs are jetty black and round,
His chine is double; starting with a bound,
He turns the turf and shakes the solid ground.
Fire from his eyes, clouds from his nostrils flow;
He bears his rider headlong on the foe.”





THE ASS. (*Equus Asinus.*)

THE ASS is a beast of burden, and extremely serviceable to man. Of greater strength than most animals of his size, he bears fatigue with patience, and hunger with apparent cheerfulness. A bundle of dried herbs, or a thistle on the road, is sufficient for his daily meal, and he is content with the clear and pure water of a neighbouring brook (in the choice of which he is particularly nice) in the absence of better fare. It is probable that the Ass was originally a native of Arabia, and other parts of the East: the deserts of Libya and Numidia, and many parts of the Archipelago, contain vast herds of wild Asses, which run with such amazing swiftness, that even the fleetest horses of the country can hardly overtake them. At present perhaps the best breed in Europe is the Spanish; and very valuable Asses are still to be had in the southern continent, where, during the existence of the Spanish dominion, the breed was very carefully attended to. In the time of Elizabeth, we are informed, there

were no Asses in this country. Our treatment of this very useful animal is both wanton and cruel, and most ungrateful, considering the great services he renders us at so little expense. The ears of the Ass are of an uncommon length; and he is of a greyish or dun colour, with a black cross on his back and shoulders. When very young, the Ass is sprightly, and even tolerably handsome; but he soon loses these qualifications, either by age or ill treatment, and becomes slow, sullen, and headstrong. The female is passionately fond of her young one; and it is said she will even cross fire and water to protect or rejoin it. The Ass is also sometimes greatly attached to his owner, whom he scents at a distance, and plainly distinguishes from others in a crowd.


The female goes with young eleven months, and seldom produces more than one foal at a time: the teeth follow the same order of appearance and renewal as those of the horse. Asses' milk has long been celebrated for its sanative qualities; invalids suffering from debility of the digestive and assimilative functions make use of it with great advantage; and to those also who are consumptive it is very generally recommended.

An old man who, a few years ago, sold vegetables in London, used in his employment an Ass, which conveyed his baskets from door to door. Frequently he gave the poor industrious creature a handful of hay, or some pieces of bread, or greens, by way of refreshment or reward. The old man had no need of any goad for the animal, and seldom, indeed, had he to lift up his hand to drive it on. His kind treatment was one day remarked to him, and he was asked if his beast was apt to be stubborn? "Ah! master," replied he, "it is of no use to be cruel, and as for stubbornness, I cannot complain; for he is ready to do anything, and go anywhere. I bred him myself. He is sometimes skittish and playful, and once ran away from me; you will hardly believe it, but there were more than fifty people after him, attempting in vain to stop him; yet he turned back of himself, and he never stopped till he ran his head kindly into my bosom."

The ancients had a great regard for this animal. The


Romans had a breed which they held in such high estimation, that Pliny mentions one of the males selling for a price greater than three thousand pounds of our money; and he says that in Celtiberia, a province in Spain, a she Ass had colts that were bought for nearly the same sum. The Ass lives nearly to the same age as the horse. From the general resemblance between the Ass and the horse, it might naturally be supposed that they were closely allied, and that one had degenerated; they are, however, perfectly distinct. There is that inseparable barrier placed between them which nature provides for the protection and preservation of her productions; their mutual offspring, the mule, being incapable of reproducing its kind.





THE WILD ASS. (*Equus hemionus*.)

THE Wild Ass, or *Koulan*, as it is called by the Persians, is a native of Asia, and it is scarcely possible to imagine a greater contrast than is presented by this noble and spirited animal to the common domestic Ass. The Wild Ass in Persia and Arabia is as large as a horse, and as fleet and as full of fire. It is used only for riding by the Persian nobles, and is preferred by them to the horse. It is indeed, in its wild state, a noble animal, with fine slender limbs and a glossy skin, carrying its head high, and moving in a very swift and graceful manner. The colour of its body is a silvery grey; the upper part of the face, the sides of the neck and body being of a flaxen hue, and a broad black stripe running down the back from the mane to the tail and across the shoulders, as in the common Ass. The inhabitants of the East have a legend that the Ass was the first animal ever subjected to the dominion of man. Its flesh is held in great esteem by the Tartars and Persians, who hunt it in preference to all kinds of game.



THE MULE.

THIS useful and hardy animal is the offspring of the horse and the ass, and partakes of the good qualities of both. The common Mule is very healthy, and will live above thirty years. The size and strength of our breed have been much improved by the importation of Spanish male asses; and it is much to be wished that the useful qualities of this animal were more attended to; for, by proper care in its breaking, its natural obstinacy would in a great measure be corrected; and it might be formed with success for the saddle, the draught, or the burden. People of the first quality are drawn by Mules in Spain, where fifty and sixty guineas is no uncommon price for them; nor is it surprising, when we consider how far they excel the horse in travelling in a mountainous country, the Mule being able to tread securely where the former can hardly stand. It is much less dainty in its food than the horse, and not so liable to disease; and has been known to go a distance of eighty or a hundred miles in one day, with a heavy weight on its back, without much fatigue.



THE ZEBRA. (*Equus Zebra.*)


THIS is the most elegantly marked quadruped in nature. He is striped all over with the most pleasing regularity; in size he resembles the mule, being smaller than the horse, and larger than the ass. The hair of his skin is uncommonly smooth, and he looks at a distance like an animal that some fanciful hand has surrounded with ribbons of white or buff, and jet black. He is a native of southern Africa—chiefly of the Cape of Good Hope; but he is also found in the kingdom of Angola; and, as we are assured by Lopez, in several provinces of Barbary. In these boundless forests the Zebra has nothing to restrain his liberty. He is too shy to be caught in traps, and therefore seldom taken alive. Were the Zebra inured to our climate, there is little doubt but he might be soon domesticated. The black cross which the ass bears on his back and shoulders, proves the affinity

between these two animals. The Zebra feeds in the same manner as the horse, ass, and mule; and seems to delight in having clean straw and dried leaves to sleep upon. His voice can hardly be described; it is thought by some persons to have a distant resemblance to the sound of a post-horn, and is more frequently exerted when the animal is alone than at other times. In former times, Zebras were often sent as presents to the oriental princes. A governor of Batavia is said to have given one to the emperor of Japan, for which he received, as an equivalent, a present to the value of sixty thousand crowns; and Teller informs us, that the Great Mogul gave two thousand ducats for one of these animals. It is usual with the African ambassadors to the court of Constantinople, to bring Zebras with them as presents for the Grand Signior. In a wild state they live in herds, and can only be tamed when taken young, or bred in captivity. Four of them, bred in the Zoological Gardens, Regent's Park, may often be seen drawing a kind of chaise-cart belonging to the gardens.

Instinct having taught these beautiful animals that in union consists their strength, they combine in a compact body when menaced by an attack either from man or beast; and if overtaken by the foe, they unite for mutual defence, with their heads together in a close circular band, presenting their heels to the enemy, and dealing out kicks in equal force and abundance. Beset on all sides or partially crippled, they rear on their hinder legs, fly at their adversary with jaws distended, and use both teeth and heels with the greatest freedom.

The *Quagga* is also a native of Southern Africa. It is more wild than the Zebra, and less beautifully marked; the stripes, indeed, do not extend over the whole body, but only over the head and neck. The colour is a blackish brown, banded with white. The *Quagga* is less than the Zebra, and not so elegantly formed, the hind quarters being higher than the shoulders. The ears are also much shorter. The *Quagga* bears the reputation of being naturally vicious, and so treacherous that it is said that, like a cat, it will bite the hand that feeds and caresses it.

SECTION VIII.—Ruminating Animals.

THE BULL. (*Bos Taurus.*)

THERE are, perhaps, no animals more generally useful to mankind than the race of oxen, in all their states of existence. They are called ruminating animals; that is, after they have eaten their food, they possess the power of returning it from the first stomach into the mouth, to be again masticated before it is finally digested. This is called chewing the cud; and as the animal generally lies down, and looks very thoughtful while the operation is performing, it is said to be ruminating.


The Bull is a very fierce creature, and when enraged, runs about, tossing up his tail, and roaring most fearfully. When attacked by men or dogs, he tears up the ground with his feet, and then gallops after his assailants endeavouring to toss them with his horns; and very often pursues in this manner any one he sees, particularly if they appear frightened. When in danger of being attacked by a Bull, the best course is to stand still, and open an umbrella, or flap a shawl, or something of that kind, in the Bull's face; as with all his fierceness he is a great coward, and only pursues those who fly from him.

The Ox, or Bullock, is used in some parts of the

country for drawing carts and waggons, and ploughing; and its flesh is called beef. The skin is tanned and made into leather; the hair is mixed with mortar; the bones are used for knife-handles, chess-men, counters, and other things, as a substitute for ivory; from its horns are made combs, and various other articles; the fat is used in making candles; the blood in refining sugar; and, in short, every part has some important use.

The common charge of stupidity urged against the Ox is wholly unfounded, as the following anecdote recorded by Mr. Bell will show. A cow, feeding in a pasture, the gate of which was open, was much annoyed by a mischievous boy, who amused himself by throwing stones at her. The peaceful animal, after enduring this patiently for some time, went up to him, and hooking the end of her horn into his clothes, carried him out of the field and laid him down in the road. She then returned calmly to her pasture, leaving him quit for a severe fright and a torn garment.






THE COW.

THE Cow is the female of the ox tribe, and her young is called a calf. A young Cow, when under two years old, is called a heifer. The Cow is as useful to mankind as the ox, except in ploughing and drawing; but to make amends, she supplies us with milk, from which butter and cheese are made. The Cow gives from six to twenty quarts of milk in a day; and the faculty of giving it in such abundance, and with so much ease, is a striking peculiarity, for this animal differs in this part of its organization from most others, having a large udder, and longer and thicker teats, than the largest animal we know of; it has likewise four teats, whilst all other animals of the same nature have but two; it also yields the milk freely to the hand, whilst all other animals, at least those that do not ruminate in the same manner, refuse it, unless their young, or some adopted animal, be allowed to partake it. The age of the Cow is known by her horns; at four a ring is formed at their roots, and every succeeding year another ring is added. Thus, by allowing three years before their appearance, and then reckoning the number of rings, the creature's age may be exactly known. Calves, when quite young, are helpless creatures, from the great length and weakness of their legs. Sometimes they are killed when young, and their flesh is then called veal. The stomach of the calf, when it is killed, is taken


out, and cleaned and salted; it is then hung up to dry, and is called rennet. In making cheese, a bit of rennet is soaked in water, which when poured into milk, turns it to curd. The curd is then separated from the whey, and put into a press, when it becomes cheese.



THE WILD BULL. (*Bos scoticus*.)

IN the Duke of Hamilton's park in Scotland, Lord Tankerville's at Chillingham, in Northumberland, and some other places, there is a breed of wild cattle, probably the last remains of those which at one period overran the island. The colour is invariably white, with muzzle and ears black, or very dark red.

At the first appearance of any person near them, these animals set off at full gallop; and at the distance of two or three hundred yards, wheel round and come boldly up again, tossing their heads in a menacing manner. On a sudden they make a full stop at the distance of forty or fifty yards, and look wildly at the object of their surprise; but on the least motion they all turn round, and gallop off again with equal speed, but not to the same distance, forming a smaller circle; and again returning with a bolder and more threatening aspect than before, they approach much nearer, when they make another stand, and again gallop off. This they do several times, shortening



THE COW.

THE Cow is the female of the ox tribe, and her young is called a calf. A young Cow, when under two years old, is called a heifer. The Cow is as useful to mankind as the ox, except in ploughing and drawing; but to make amends, she supplies us with milk, from which butter and cheese are made. The Cow gives from six to twenty quarts of milk in a day; and the faculty of giving it in such abundance, and with so much ease, is a striking peculiarity, for this animal differs in this part of its organization from most others, having a large udder, and longer and thicker teats, than the largest animal we know of; it has likewise four teats, whilst all other animals of the same nature have but two; it also yields the milk freely to the hand, whilst all other animals, at least those that do not ruminate in the same manner, refuse it, unless their young, or some adopted animal, be allowed to partake it. The age of the Cow is known by her horns; at four a ring is formed at their roots, and every succeeding year another ring is added. Thus, by allowing three years before their appearance, and then reckoning the number of rings, the creature's age may be exactly known.

Calves, when quite young, are helpless creatures, from the great length and weakness of their legs. Sometimes they are killed when young, and their flesh is then called veal. The stomach of the calf, when it is killed, is taken

their distance, and advancing nearer till they come within a few yards, when most persons consider it prudent to leave them, not choosing to provoke them further, as it is probable that in a few turns more they would make an attack.

The mode of killing these animals, as was practised a few years ago, was the only remnant of the ancient mode of hunting that existed in this country. On notice being given that a Wild Bull would be killed on a certain day, the inhabitants of the neighbourhood assembled, sometimes to the number of a hundred horsemen, and four or five hundred foot, all armed with guns or other weapons. Those on foot stood upon the walls, or climbed into trees, while the horsemen separated a Bull from the rest of the herd, and chased him until he stood at bay, when they dismounted and fired. At some of these huntings, twenty or thirty shots have been discharged before the animal was subdued. On such occasions the bleeding victim grew desperately furious from the smarting of his wounds, and the shouts of savage joy echoing from every side.

When the Cows calve, they hide their young ones for a week or ten days in some sequestered retreat, and go to suckle them two or three times in a day. If any person comes near one of the calves, it crouches close upon the ground, and endeavours to hide itself, a proof of the native wildness of the animals. In one instance where a calf was disturbed, it pawed the ground like an old Bull, and attempted to butt with its head, till it fell from weakness. It had done enough, however, to raise an alarm, and the whole herd came to its rescue, compelling the intruder to decamp; for the dams will allow no one to touch their young without attacking him with impetuosity. In the Duke of Hamilton's park, in the summer of 1841, a calf, which was disturbed by the passing of a carriage near it, bellowed so fearfully, as to rouse the whole herd, though they were at a considerable distance.



THE AFRICAN BUFFALO. (*Bubalus Caffer.*)

IN its general form the Buffalo has a great resemblance to the ox; but it differs from that animal in its horns, and in some particulars of its internal structure. It is larger than the ox; the head is also bigger in proportion, the forehead higher, and the muzzle longer. The horns are large, and of a compressed form, with the exterior edge sharp; they are straight for a considerable length from their base, and then bend slightly upward. The general colour of the animal is blackish, except the forehead and the tip of the tail, which are of a dusky white. The hunch is not, as many have supposed it, a large fleshy lump, but is occasioned by the bones that form the withers being continued to a greater length than in most other animals. The Buffalo is a native of most parts of the torrid zone, and of almost all warm climates; always dwelling in moist and marshy places, where he delights to roll in the mire. In a wild state the Buffalo is exceedingly fierce; but in some of the tropical countries he is

perfectly domestic, and very useful for many purposes, being an animal of patience and great strength. When employed in the labours of agriculture, he has a brass ring put through his nose, by which means he is led at pleasure. Buffaloes are common in the Pontine Marshes near Rome, where they were brought from India in the sixth century. In India they constitute the riches and food of the poor, who employ them in their fields, and make butter and cheese from their milk. They are much valued for their hides; of which, in several countries, and especially in England, military belts, boots, and other implements of war are made. There are various species of Buffaloes, of which the Cape Buffalo, from South Africa, is the best known, and most valuable.

Buffaloes, in their native country, fight so fiercely with each other, that African travellers have remarked that they are seldom found without torn ears, and scars of various kinds on the neck and body. And they are no less treacherous than ferocious, lurking among the trees in concealment, until some unfortunate passenger passes. The animal will then suddenly rush upon him, and there is little chance of the victim escaping unless a tree be at hand. The furious beast, not contented with throwing him down and killing him, stands over him for a long time, trampling on and tearing the body to pieces, he then strips off the skin with his rough and prickly tongue. Even after all this he repeatedly returns to the body to gratify afresh his savage disposition.



THE BISON. (*Bos Bonasus*.)


THERE are two kinds of Bison ; one a native of Europe, and the other of America. The European Bison, or *Bonasmus*, is as large as a bull or ox ; maned about the back and neck, like a lion ; and his hair hanging down under his chin, or nether jaw, like a large beard. The fore parts of his body are thick and strong, but the hinder parts are comparatively slender. He has a little ridge along his face from his forehead down to his nose, which is very hairy ; his horns are large, very sharp, and turning towards his back, like those of a wild goat. The American Bison attains a size far superior to that of the largest breeds of our common oxen ; and is met with throughout nearly the whole of the uninhabited parts of North America, from Hudson's Bay to Louisiana and the frontiers of Mexico. Captains Lewis and Clarke, and Dr. James, bear frequent testimony to the almost incredible numbers in which these animals assemble on the banks of the Missouri. "Such was their multitude," say the first-named travellers, "that, although the river, including an island over which they passed, was a mile in

breadth, the herd stretched, as thick as they could swim, completely from one side to the other." And again they say: "If it be not impossible to calculate the moving multitude which darkened the whole plains, we are convinced that twenty thousand would be no exaggerated number." Dr. James tells us that, "in the middle of the day, countless thousands of them were seen coming in from every quarter to the stagnant pools;" their paths, as he informs us elsewhere, being "as frequent, and almost as conspicuous, as the roads in the most populous parts of the United States."

These wild cattle defend themselves from the wolves in the most admirable manner. When they hear their savage enemies approaching, they form themselves adroitly into a circle. The weakest are left in the middle, whilst the strongest are on the outside, and present to their foes an impenetrable phalanx of horns. The vignette is an illustration of this subject.

Exciting stories of the Buffalo hunt, both American and African, will be seen in Catlin's North American Indians, and Harris's wild Animals and Sports of Southern Africa.





THE ZEBU, OR BRAHMIN BULL. (*Bos indicus*.)

PENNANT describes the Zebu, or Indian Ox, as sometimes surpassing in size the largest of the European breeds, and the hunch on his shoulders as weighing frequently fifty pounds. There are many varieties, with and without horns, differing in size from that above named down to the dimensions of an ordinary hog. They are spread over the whole of Southern Asia, and from Abyssinia to the Cape of Good Hope. In all these countries the Zebu supplies the place of the Ox, both as a beast of burden and as an article of food. By the Hindoos they are treated with great veneration, and it is held sinful to deprive them of life, or eat their flesh. A select number are exempted from all labour, and allowed to wander about, and subsist on the voluntary and pious contributions of the devotees of their faith.

Emboldened by the toleration they experience, they make free with every vegetable to which they take a fancy, no one daring to resist or drive them away; often they lie down in the street; no one must disturb them; every one must give place to the sacred Ox of Brahma: thus they are frequently nuisances, which superstition alone would endure.



THE SHEEP. (*Ovis Aries.*)

THE Sheep has been so long subjected to the empire of man, that it is not known with certainty from what race our domestic species has been derived. It is supposed, however, to be from the Mouflon, or Musmon, of Sardinia and Crete. This animal is one of the most useful ever bestowed on us by a bountiful Providence; and in patriarchal times, the number of Sheep constituted the riches of kings and princes. It is universally known, its flesh being one of the chief kinds of human food, and its wool being of great use for clothing. Although of a moderate size, and well covered, it does not live more than nine or ten years. The Ewe has one or two young at a time, and the young one, which is called a lamb, has always been an emblem of innocence.

In its domestic state, it is too well known to require a detail of its peculiar habits, or of the methods which have been adopted to improve the breed. No country produces finer Sheep than England, either with larger fleeces, or better adapted for the business of clothing. Those of Spain have confessedly finer wool, some of which we generally require to work up with our own; but the weight of a Spanish fleece is much inferior to one of Lincoln or Tees Water. Merino, or Spanish Sheep, have of late years been introduced with some success into our

English pastures, and the wool of the hybrids, raised between the Merino Sheep and the South Down Sheep, is thought nearly equal to that of Spain.

In stormy weather, these animals generally hide themselves in caves from the fury of the elements; but if such retreats are not to be found, they collect themselves together, and during a fall of snow, place their heads near each other, with their muzzles inclined to the ground. In this situation they sometimes remain, till hunger compels them to gnaw each other's wool, which forms into hard balls in the stomach, and destroys them. But in general, they are sought out and extricated soon after the storm has subsided.

“The Sheep,” Mr. Bell observes, “is one of the most interesting of all animals as regards its historical relations with man. It was the subject of the first sacrifices, and was used in its typical character as an offering of atonement; and the relation which existed between the patriarchal shepherds and their flocks, was of so intimate and even affectionate a nature, as to have afforded the subject of many beautiful passages in the Holy Scriptures.”





THE RAM

Is the male sheep, and is so strong and fierce that he will boldly attack a dog, and often come off victorious: he has even been known, regardless of danger, to engage a bull; and his forehead being much harder than that of any other animal, he seldom fails to conquer. He overcomes the bull who, by lowering his head, receives the stroke of the Ram between his eyes, which usually brings him to the ground.



THE WALLACHIAN RAM.

THE singular conformation of the horns, which adorn the head of this breed of Sheep, has induced us to insert a

figure of the animal in this work, though it is only a variety of the common species. The horns of the Ewe are twisted also, but not so much as those of the Ram, which form, near the head, a spiral line. The wool is much longer than that of the common Sheep, and resembles the hair of the goat. A fine Ram of this species was presented some years since to the Zoological Gardens in the Regent's park, by Dr. Bowring. It is there called the Parnassian Sheep, having been brought from Mount Parnassus.

THE ARGALI, OR WILD SHEEP OF ASIA, in figure somewhat resembles a ram, but his wool is rather like the hair of a goat. His horns are large and bent backwards, and his tail is short. He is of the size of a small deer, active, swift, wild, and found in flocks in the rocky, dry deserts of Asia, Kamschatka, Barbary, and Corsica. His flesh and fat are delicious. He is called also the Siberian Sheep or Goat, and is considered by some to be the parent stock of the domestic Sheep.



THE GOAT. (*Capra hircus.*)

THE Goat, next to the cow and the sheep, has been always reckoned, especially in ancient and patriarchal times, the most useful domestic animal. Its milk is sweet, nourishing, and medicinal, and better adapted for persons of weak digestion than that of the cow, as it is not so apt to curdle on the stomach. The female has generally two

young ones at a time, which are called kids. This animal is admirably adapted for living in wild places; it delights in climbing precipices, and is often seen reposing in peaceful security on rocks overhanging the sea. Nature indeed has in some measure fitted it for traversing these eminences; the hoof being hollow underneath, with sharp edges, so that it can walk as securely on the ridge of a house as on the level ground. The flesh of the Goat is seldom eaten; but that of the kid is esteemed a very delicate food, and is frequently eaten on the Continent. In the East, the long soft hair of the Goat is used in making the beautiful Cashmere shawls; and of the skin is manufactured morocco leather. The skin of the kid is well known for its use in making gloves.



THE IBEX, OR BOQUETIN,

Is a Wild Goat, which inhabits the Pyrennean-mountains, the Alps, and the highest mountains of Greece. He is of an admirable swiftness; his head is armed with

two long, knotted horns, inclining backwards; his hair is rough, and of a deep brown colour. The male only has a beard, and the female is less than the male. This animal skips from rock to rock, and often, when pursued, leaps down enormous precipices, and is said to bend his head between his fore legs while springing, so as to break his fall, by alighting partly on his horns. The Ibex has been known to turn on the incautious huntsman, and tumble him down the precipice, unless he has time to lie down, and let the animal pass over him.



THE ANTELOPE. (*Antilope cervicapra*.)

THESE beautiful inhabitants of the temperate regions of Africa possess swiftness and elegance of shape in an eminent degree. They are timid, inoffensive, and gregarious. The males have horns like those of the goat, and never shed them; they are smooth, long, and twisted spirally or annulated. The general colour of the hair is brown, and, in some species, a beautiful yellow. The

eyes are exceedingly bright, and have often been compared to those of a beautiful nymph by Persian and other poets. Enjoying perfect liberty, they range in herds, through the deserts of Arabia, and bound from rock to rock with wonderful agility. Their long and slender legs are peculiarly suited to their habits and manners of life, and are, in some of the species, so slender and brittle as to snap with a very trifling blow. The Arabs, taking advantage of this circumstance, catch them by throwing sticks at them, by which their legs are broken.



THE GAZELLE. (*Antilope Dorcas.*)

“The wild Gazelle, on Judah’s hills,
 Exulting yet may bound,
 And drink from all the living rills
 That gush on holy ground.
 Its airy step and glorious eye
 May glance in tameless transport by.”—BYRON.

The Gazelle is the most elegant of antelopes. The Ara-

bian poets have applied their choicest epithets to the beauty of this animal, and their descriptions have been adopted into our own poetry. Byron, in speaking of the dark eyes of an eastern beauty, says,

“Go look on those of the Gazelle.”

When the Persian describes his mistress, she is “an antelope in beauty,”—“his Gazelle employs all his soul;” and thus, in their figurative language, perfect beauty and Gazelle beauty are synonymous. These animals are spread, in innumerable herds, from Arabia to the river Senegal in Africa. Lions and panthers feed upon them; and man chases them with the dog, the cheetah, and the falcon. The height of the Gazelle is about twenty inches, the skin beautifully sleek, its body extremely graceful, its head unusually light, its ears flexible, its eyes most brilliant and glancing, and its legs as slender as a reed.

THE CHAMOIS. (*Antilope Rupicapra.*)

THE Chamois is about three feet in length and two in height; its horns six or seven inches long, its ears small, and its head resembling that of the goat. The body is covered with long brown hair, the hue of which varies with the season.

The flesh is considered a savoury food, and the skin is wrought into a soft pliable leather, well known in domestic economy.

The Chamois is found only in mountainous regions, where they herd together on lofty and almost inaccessible cliffs and precipices. They are so acute and shy, that it is only by the greatest patience and skill that the hunter can approach near enough to shoot them; and they are so swift, and leap with such extraordinary sureness of foot, that to overtake them is impossible.

“————— But beasts have reason too;
And that we know, we men that hunt the Chamois,
They never turn to feed—sagacious creatures—
Till they have placed a sentinel a-head,
Who pricks his ears whenever we approach,
And gives alarm with clear and piercing pipe.”

SCHILLER'S WILLIAM TELL.



THE NYL GHAU, OR BLUE OX. (*Antelope picta*.)

THIS is a kind of antelope, found in India. In the wild state, these animals are said to be ferocious, but they may be domesticated, and in that condition, give frequent tokens of familiarity, and even of gratitude, to those under whose care they are placed. The female, or doe, is much smaller than the male, and of a yellowish colour, by which she is easily distinguished from the buck, who is of a grey tint.

Its manner of fighting is very particular, and is thus described:—Two of the males, at Lord Clive's, being put into an enclosure, were observed, while they were at some distance from each other, to prepare for the attack, by falling down upon their knees; they then shuffled towards each other, still keeping upon their knees; and, at the distance of a few yards, they made a spring, and darted against each other with great force.

The following anecdote will serve to show that these animals are sometimes fierce and vicious, and not to be

depended upon :— A labouring man, without knowing that the animal was near him, went up to the outside of the enclosure : the Nyl Ghau, with the quickness of lightning, darted against the woodwork with such violence, that he dashed it to pieces, and broke one of his horns close to the root. The death of the animal, soon after, was supposed to be owing to the injury he sustained by the blow.



THE GNU. (*Antelope Gnu.*)

THIS very singular animal may be called a horned horse ; as it has the shape and mane of a horse, with the addition of a formidable pair of horns, a kind of beard below the chin, and a fringe of hair below the body, along the breastbone. Its habits also agree in many respects with those of the wild horse. The Gnus live together in herds, and when alarmed, fling up their heels, and plunge and rear, tossing their heads and tails, before they gallop off ; which they do, the whole herd following their leader singly, like a troop of soldiers. The Gnu inhabits the sandy deserts of South Africa ; and its flesh, which is said to resemble beef, is sometimes eaten by the colonists near the Cape of Good Hope. When caught young, the Gnu may be tamed, but its disposition is always uncertain, and when offended, it throws itself on its knees, like the nyl ghau, and then springing up, butts furiously with its horns.



THE STAG. (*Cervus Elephas.*)

THIS animal is the male of the red Deer, and is generally famed for long life, though upon no certain authority. Naturalists agree, however, upon this point, that his life may exceed forty years: but that his existence, as it has been asserted, reaches to three centuries, is too absurd to be believed. His horns are at first very small, but gradually increase in size, as they are yearly shed and renewed, till the stag has completed his fifth year, when they become very large and branching, and remain so during the remainder of his life. The Stag is the tallest of the deer kind, and is called a Hart after he has completed his fifth year; the female, called the hind, is without horns. Every year, in the month of April, when the Stag has lost his horns, he appears conscious of his temporary weakness, and hides himself till his new ones have grown and are hardened. This is generally in about ten weeks, even when the Stag is full grown; his horns at this age weigh between twenty and thirty pounds. Little need be said of the pleasure taken in hunting the Stag, the Hart, and the Roe-buck, it being a matter well known in this country, and in all parts of Europe. The following fact, recorded in

history, will serve to show that the Stag is possessed of an extraordinary share of courage, when his personal safety is concerned:—In the reign of George the Second, William, Duke of Cumberland, caused a tiger and a Stag to be enclosed in the same area; and the Stag made so bold a defence, that the tiger was at length obliged to give up. The flesh of the Stag is accounted excellent food, and his horns are useful to cutlers; even their shavings are used to make ammonia, so much esteemed in medicine, under the name of *hartshorn*. The swiftness of the Stag has become proverbial, and the diversion of hunting this creature has, for ages, been looked upon as a royal amusement. In the time of William Rufus and Henry the First, it was less criminal to destroy a human being than a full-grown Stag. This animal, when fatigued in the chase, often throws himself into a pond of water, or crosses a river; and, when caught, sheds tears like a child.

“To the which place a poor sequestered Stag,
That from the hunter's aim had ta'en a hurt,
Did come to languish; and indeed, my lord,
The wretched animal heaved forth such groans
That their discharge did stretch his leathern coat
Almost to bursting; and the big round tears
Coursed one another down his innocent nose
In piteous chase.”


SHAKESPEARE.






THE ROEBUCK (*Cervus capreolus*)

Is one of the least of the Deer kind known in these climates, being not above three feet in length, and two in height, and seldom lives more than fifteen years. His horns are about nine inches long, round, and divided into three small branches, and his colour is of a brown shade on the back, his face partly black and partly ash-colour, the chest and belly yellow, and the rump white; his tail is short. The Roe buck is more graceful, more active, more cunning, and comparatively swifter than the stag; his flesh is much esteemed. He is very delicate in the choice of his food, and requires a larger tract of country, suited to the wildness of his nature, which can never be thoroughly subdued. No arts can teach him to be familiar with his keeper, nor in any degree attached to him. These animals are easily terrified; and in their attempts to escape will run with such force against the walls of their inclosure, as sometimes to disable themselves: they are also subject to capricious fits of fierceness; and, on these occasions, will strike furiously with their horns and feet at the object of their dislike. The only parts of Great Britain where they are found, are the Highlands of Scotland.

THE FALLOW DEER. (*Cervus dama.*)

THESE are the Deer now usually kept in our parks. The beautifully spotted kind are said to have been brought from Bengal, and the very deep brown from Norway by King James I. Their horns are broad and flat; the male is called a buck, the female a doe, and the young one a fawn. The buck casts his horns every spring, and they increase in size annually till he has attained his fifth year. The venison of this Deer is very far superior to that of the red deer, which is coarse and tough. The buck-skin and doe-skin are well known, as furnishing a peculiarly soft and warm leather, which is used for gloves, gaiters, &c. The horns are used for the handles of knives, &c., like those of the stag; and the refuse is, in the like manner, used in the manufacture of ammonia. The buck stands about three feet high, and measures about five feet in length; the doe is somewhat smaller. The tail is much longer than either that of the stag or the roebuck, being nearly seven inches and a half long.



THE ELK (*Cervus Alces*)

Is the largest of all the Deer kind. The antlers, at first simple, and then divided into narrow slips, assume in the fifth year the form of a triangular blade, denticulated on the external edge and very thick at the base, and increase with age, till they weigh fifty or sixty pounds, and have fourteen branches to each horn. The Elk lives in forests, feeding upon branches and sprouts of trees, and inhabits Europe, Asia, and America; in the last-named country he is known by the name of the moose-deer. There is very little difference between the European Elk and the American moose-deer, though they are larger in the New World than with us, owing perhaps to the extensive forests in which they range. In all places, however, they are timorous and gentle; content with their pasture, and never willing to disturb any other animal. The pace of the Elk is a high, shambling trot, but it runs with great swiftness. Formerly these animals were made use of in Sweden to draw sledges, but their swiftness gave criminals such means of escape, that this employment of them was prohibited under great penalties. The female is less than the male, and has no horns.



THE REIN-DEER (*Cervus Tarantus, or rangifer*)

Is found in most of the northern regions of Europe, Asia, and America, and its general height is about four feet and a half. The colour is brown above and white beneath; but as the animal advances in age it often becomes of a grayish white. The hoofs are long, large, and black. Both sexes are furnished with horns, but those of the male are much the largest. To the Laplanders this animal supplies the place of the horse, the cow, the goat, and the sheep; and is their only wealth. The milk affords them cheese; the flesh, food; the skin, clothing; of the tendons they make bowstrings, and when split, thread; of the horns, glue; and of the bones, spoons. During the winter, the Rein-deer supplies the want of a horse, and draws sledges with amazing swiftness over the frozen lakes and rivers, or over the snow, which at that time covers the whole country. Innumerable are the uses, the comforts, and advantages which the poor inhabitants of this dreary climate derive from this animal. We cannot sum them up better than in the beautiful language of the poet:—

“ Their Rein-deer form their riches. These their tents,
Their robes, their beds, and all their homely wealth

Supply, their wholesome fare, and cheerful cups :
Obsequious at their call, the docile tribe
Yield to the sled their necks, and whirl them swift
O'er hill and dale, heaped into one expanse
Of marbled snow, as far as eye can sweep,
With a blue crest of ice unbounded glazed."

The mode of hunting the wild Rein-deer by the Laplanders, the Esquimaux, and the Indians of North America, has been accurately described by late travellers. Captain Franklin gives the following interesting account of the mode practised by the Dog-rib Indians, to kill these animals. "The hunters go in pairs, the foremost man carrying in one hand the horns and part of the skin of the head of a Deer, and in the other a small bundle of twigs, against which he, from time to time, rubs the horns, imitating the gestures peculiar to the animal. His comrade follows, treading exactly in his footsteps, and holding the guns of both in a horizontal position, so that the muzzles project under the arms of him who carries the head. Both hunters have a fillet of white skin round their foreheads, and the foremost has a strip of the same round his wrists. They approach the herd by degrees, raising their legs very slowly, but setting them down somewhat suddenly, after the manner of a Deer, and always taking care to lift their right or left feet simultaneously. If any of the herd leave off feeding to gaze upon this extraordinary phenomenon, it instantly stops, and the head begins to play its part, by licking its shoulders, and performing other necessary movements. In this way the hunters attain the very centre of the herd without exciting suspicion, and have leisure to single out the fattest. The hindmost man then pushes forward his comrade's gun, the head is dropped, and they both fire nearly at the same instant. The Deer scamper off, the hunters trot after them; in a short time the poor animals halt, to ascertain the cause of their terror; their foes stop at the same moment, and having loaded as they ran, greet the gazers with a second fatal discharge. The consternation of the Deer increases; they run to and fro in the utmost confusion; and sometimes a great part of the herd is destroyed within the space of a few hundred yards."

THE AXIS, (*Cervus Axis*,)

A VERY beautiful species of the deer found in Africa and the East Indies, of a light red colour, though some of the kind are of a deeper red. It is about the size of a fallow deer, and often variegated with beautiful spots of bright white. The horns are slender and triple forked. The Axis is a timid and harmless creature, more ornamental to the landscape, where it skips and plays in a wild state, than useful to man. It is extremely docile, and possesses the sense of smelling to an exquisite degree. Though it is a native of the banks of the Ganges, it appears to bear the climates of Europe without injury.

The Great Axis, which is a native of Borneo and Ceylon, is about the height of a horse, and of a reddish-brown colour. The horns are bifurcated, thick, strong, and rugged; about two feet nine inches long, and two feet four inches between the tips.

The Lesser Axis is a gregarious animal, inhabiting Java, Ceylon, Borneo, and some other oriental islands. It is hunted with ardour, the sport affording the highest diversion, and the flesh being esteemed excellent.



THE WAPITI (*Cervus Canadensis*)

Is a native of Canada and other northern parts of America, and is one of the most gigantic of the deer tribe, growing to the height of our tallest oxen, and uniting great activity to strength of body and limbs. His horns, which he sheds annually, are very large, branching in serpentine curves, and measuring, from tip to tip, upwards of six feet. These animals make a shrill noise, resembling the braying of an ass, and are supposed to be the most stupid of the deer kind. The flesh is coarse, and little esteemed, but the hide, when made into leather, is said not to become hard in drying after being wetted, a quality which entitles it to a preference over almost every other kind. There are several of these splendid animals in the collection of the Zoological Society, in the Regent's Park, where they continue to form objects of singular interest and attraction. The male is, however, very fierce, always endeavouring to attack those who approach him; and on one occasion seriously injured one of the visitors to the gardens.



THE GIRAFFE, OR .CAMELOPARD.

(*Camelopardalis Giraffa.*)

THIS most remarkable ruminant, which in its general structure nearly approaches the deer, has points of affinity, also, with the antelopes and camels, besides very striking peculiarities of its own.

The head is the most beautiful part of the animal: it is small, and the eyes are large, brilliant, and very full. Between the eyes, and above the nose, is a swelling very prominent and well-defined. This prominence is not a fleshy excrescence, but an enlargement of the bony substance; and it seems to be similar to the two little lumps, or horns with which the top of the head is armed, and which, being about three inches in length, spring on each

side of the head, just above the ears, and are covered with a thick tuft of stiff upright hairs. The neck is remarkably elongated, and it is furnished with a very short, stiff mane, which stands out erect from the skin. The height of a full-grown Giraffe in a wild state is said to be twenty feet, measuring from the hoofs to the tip of the ears; but none of those in England exceed fourteen feet. At first sight, the fore legs appear much longer than the hind ones; but the fact is, that the legs are of the same length, and it is only the height of the withers that occasions the apparent disproportion. The vertebræ of the back are slightly curved. The colour of the Giraffe is a light fawn, marked with spots only a few shades darker. The legs are very slender; and notwithstanding the length of the neck, it manifests great difficulty in taking anything from the ground. To do this it puts out first one foot, then the other; repeating the same process several times; and it is only after several of these experiments that it at length bends its neck, and applies its lips and tongue to the object in question. It willingly accepts fruit and branches of a tree when offered to it; and seizes the foliage in a very singular manner, thrusting forth a long, reddish, and very narrow tongue, which it rolls round whatever it wishes to secure. Indeed, the tongue is a most remarkable organ in this animal, and we have been witness of some amusing exploits with it. In the Zoological Gardens at Regent's Park, many a fair lady has been robbed of the artificial flowers which adorned her bonnet, by the nimble, filching tongue of the rare object of her admiration.

The Giraffe is a native of Africa; and it was for a long time known only by the descriptions of travellers. It was first sent to Europe in 1829; but since that time many have been introduced, and two young ones have been born in the Zoological Gardens in the Regent's Park.

Vaillant, in his entertaining Travels in Africa, gives an animated account of a Giraffe hunt:—"After several hours' fatigue, we discovered, at the turn of a hill, seven Giraffes, which my pack instantly pursued. Six of them went off together; but the seventh, cut off by my dogs,

took another way. I followed it at full speed, but, in spite of the efforts of my horse, she got so much ahead of me, that in turning a little hill, I lost sight of her altogether. My dogs, however, were not so easily put out. They were soon so close upon her that she was obliged to stop to defend herself. From the place where I was, I heard them give tongue with all their might; and, as their voices appeared all to come from the same spot, I conjectured that they had got the animal in a corner, and I again pushed forward. I had scarcely got round the hill, when I perceived her surrounded by the dogs, and endeavouring to drive them away by heavy kicks. In a moment I was on my feet, and a shot from my carbine brought her to the earth. Enchanted with my victory, I returned to call my people about me, that they might assist in skinning and cutting up the animal. On my return I found her standing under a large ebony-tree, assailed by my dogs. She had staggered to this place, and fell dead at the moment I was about to take a second shot."

The horns of the Giraffe, small as they are, and muffled with skin and hair, are by no means the insignificant weapons they seem. We have seen them wielded by the males against each other with fearful and reckless force; and we know that they are the natural arms of the Giraffe, most dreaded by the keeper of the present living Giraffes in the Zoological Gardens, because they are most commonly and suddenly put in use. The Giraffe does not butt by depressing and suddenly elevating the head, like the deer, ox, or sheep; but strikes the callous obtuse extremities of the horns, against the object of his attack, with a sidelong sweep of the neck.

The Giraffe has a peculiarly awkward manner of trotting, as it appears to move both the legs on one side at the same time, though this is not really the case. The female Giraffe in the Regent's Park was a very bad mother to her first young one, as she would not let it suck, and beat it away whenever it approached. The poor thing was fed with cow's milk, but it soon died. The present young one has been more kindly treated, and has in consequence thriven well.



THE CAMEL. (*Camelus bactrianus*.)

“ In silent horror, o’er the boundless waste,
 The driver Hassan with his Camels passed :
 One cruise of water on his back he bore,
 And his light scrip contained a scanty store :
 A fan of painted feathers in his hand,
 To guard his shaded face from scorching sand ;
 The sultry sun had gained the middle sky,
 And not a tree, and not a herb was nigh :
 The beasts with pain their dusty way pursue,
 Shrill roar’d the winds, and dreary was the view !”

COLLINS.

THE Bactrian Camel is a native of Asia, and is generally of a brown or ash colour. His height is about six feet. He is one of the most useful quadrupeds in oriental countries ; his docility and strength, his endurance of hunger and thirst, and his swiftness, on the sands of Arabia, make him a most valuable acquisition to the inhabitants of those desert places. The principal characteristics of the Camel

are these:—He has two large and hard bunches on his back, and is destitute of horns; the upper lip is divided like that of the hare; and the hoofs small and undeciduous. But the peculiar and distinguishing characteristic of the Camel is, its faculty of abstaining from water for a greater length of time than any other animal; for which nature has made a wonderful provision, in giving it, besides the four stomachs which it has in common with all ruminating animals, a fifth bag, serving as a reservoir for water, where it remains without corrupting or mixing with the other aliments. By this singular structure it can take a prodigious quantity of water at one draught, and is enabled to pass as much as fifteen days without drinking again. But besides this reservoir of water the animal is said in cases of emergency to draw sustenance from the humps on his back, which are of a fatty substance; thus, after long privation, they become absorbed. A large Camel is capable of carrying a thousand, and sometimes even twelve hundred weight, and, like the elephant, is tame, tractable, and strong; but like him he has his periodical fits of rage, and at these times has been known to take up a man in his teeth, throw him on the ground, and trample him under his feet. Like the horse, he gives security to his rider; and, like the cow, he furnishes his owner with meat for his table, and the female with milk for his drink. The flesh of the young Camel is esteemed a delicacy, and the milk of the female, diluted in water, is the common drink of the Arabians. The hair or fleece, which falls off entirely in the spring, is superior to that of any other domestic animal, and is made into very fine stuffs, for clothes, coverings, tents, and other furniture. The female goes one year with young, and produces but one at a time. The Camel kneels to receive his burthen, and it is said that he refuses to rise, if his master impose upon him a weight above his strength. He has callosities on his knees, and on his breast, which prevent him from being hurt by kneeling to take up his load; and sleeps with his knees bent under him, and his breast on the ground. He arrives at maturity in about five years, and the duration of his life is from forty to fifty years.



THE DROMEDARY, (*Camelus Dromedarius*.)

ANOTHER species of Camel, of less stature than the former, but much swifter, and having but one hard bunch on his back, is domesticated throughout Africa, as well as in Asia. It is said that a Dromedary can travel one hundred miles a day, and carry fifteen hundred weight. Attempts have been made to introduce the Camel and Dromedary into our West India islands, but they have not succeeded; they have, however, been comparatively naturalized near Pisa in Italy. The Camels used as beasts of burden in Egypt are all Dromedaries; and the first experiment which an European makes in bestriding one is generally a service of some little danger, from the peculiarity of the animal's movement in rising. Denon, the French traveller, has described this with his usual vivacity: "During the French invasion of Egypt, a part of Dessaix's division," to which the scientific traveller was attached, "was sent with Camels to a distant post across the desert. The Camel, slow as he generally is in his actions, lifts up his hind legs very briskly at the instant the rider is in the

saddle; the man is thus thrown forward; a similar movement of the fore legs throws him backward; each motion is repeated; and it is not till the fourth movement, when the Dromedary is fairly on his feet, that the rider can recover his balance. None of us could resist the first impulse, and thus nobody could laugh at his companions." Macfarlane, in his work on Constantinople, tells us that upon his first Camel adventure he was so unprepared for the probable effect of the creature's rising behind, that he was thrown over his head, to the infinite amusement of the Turks, who laughed heartily at his inexperience.

Though the name of Dromedary is very generally applied to all the one-humped camels, both in common parlance and books on Natural History, it is said that the true Dromedary (*El Herie*) is merely a peculiarly swift camel, which may have either one or two humps, like the common kind. The name of Dromedary, indeed, appears to be applied in the East to all the higher bred camels, the genealogy of which is kept by the Arabs as carefully as that of their horses.

Possessing strength and activity, surpassing that of most beasts of burthen, docile, patient of hunger and thirst, and contented with small quantities of the coarsest provender, the camel is one of the most valuable gifts of Providence. There is nothing, however, in the exterior appearance of the animal to indicate the existence of any of its excellent qualities. In form and proportions it is very opposite to our usual ideas of perfection and beauty. A stout body, having the back disfigured by one or two humps; limbs long, slender, and seemingly too weak to support the trunk; a long, thin, crooked neck, surmounted by a heavily-proportioned head, are all ill-suited to produce favourable impressions. Nevertheless, there is no creature more excellently adapted to its situation, nor is there one in which more of creative wisdom is displayed in the peculiarities of its organization. To the Arabs, and other wanderers of the desert, the camel is at once, wealth, subsistence, and protection



THE LLAMA, OR CAMEL OF AMERICA,

(*Auchenia glama*,)

Is a mild, timorous creature, not above four feet and a half in height, and usually of a brown colour. It bears in form a general resemblance to the camel; but, instead of a protuberance on the back, it has one on the breast. Llamas are used as beasts of burden by the South Americans, and are so capriciously vindictive, that, if their drivers strike them they immediately squat down, and nothing but caresses can induce them to rise again. They have been known to kill themselves by striking their heads against the ground in their rage, when by blows they have been urged forward against their will. They express their anger by spitting at their adversary. The *Alpacas* are much smaller than the Llamas, and of different colours in a domestic state. They are used for the same purposes, and differ little in habits and nature. The wool of both these animals is made use of for several purposes, and is a principal ingredient in the composition of hats, in several parts of the new and old continent; and the flesh of the young Llamas is, in their native country, considered a great delicacy, and is as good as that of the fat sheep of Castile. In Peru, where the animals are found, there are public shambles for the sale of their flesh.


SECTION IX.—Quadrupana, or Four-handed Animals.



THE OURANG OUTANG, OR CHIMPANSEE.

(*Simia troglodytes.*)

ANIMALS of the Monkey tribe are furnished with hands instead of paws; their ears, eyes, eyelids, lips, and breasts resemble those of the human species. For greater facility of description, the animals of this extensive tribe are usually arranged in the three divisions of Apes, Baboons, and Monkeys. Apes are destitute of tails, and the chief of this kind is the Ourang Outang, or Wild Man of the Woods: he is found in the interior parts of Africa, and Madagascar. He is a solitary animal, and avoids mankind. The largest are said to be six feet high, very active, strong, and intrepid, capable of overcoming the strongest man: they are likewise exceedingly swift, and cannot easily be taken alive. When young, however, the Ourang Outang is capable of being tamed: one of them, shown in London some years ago, was taught to sit at table, make use of a spoon or fork in eating, and drink wine out of a glass. It was mild and affectionate, much attached to its keeper, and obedient to his commands.



THE BARBARY APE.

THE animal next to the ourang outang, and to be placed in the same class, is the Margot, or Barbary Ape. Caubasson relates a laughable anecdote of one of these animals, which he brought up tame, and which became so attached to him as to be desirous of accompanying him wherever he went: when, therefore, he had to perform divine service, he was under the necessity of shutting him up. One day, however, the animal escaped, and followed the father to church, where, silently mounting on the top of the sounding-board, above the pulpit, he lay perfectly quiet till the sermon began. He then crept to the edge, and overlooking the preacher imitated his gestures in so grotesque a manner, that the whole congregation were unavoidably excited to laughter. Caubasson, surprised and displeased at this ill-timed levity, reprov'd his auditors for their inattention; and on the obvious failure of his reproof, he, in the warmth of zeal, redoubled his actions and his vociferations. These the Ape so exactly imitated that all respect for their pastor was swallowed up in the scene before them, and they burst into a loud and continued roar of laughter. A friend of the preacher at length stepped up to him, and on pointing out the cause of this improper conduct, it was with the utmost difficulty he could command a serious countenance while he ordered the Ape to be taken away.



THE BABOON, (*Cynocephalus*,)

A GENUS of *Quadrumania*, which forms the last link in the chain that unites the *Simiæ* with quadrupeds: it comprises a large, fierce, and formidable race of animals, who, though they in a slight degree partake of the human conformation, as seen in the Ourang Outang, &c., are in their habits, propensities, and dispositions the very reverse of gentleness, and docility.

This creature is commonly covered with black hair, a little intermixed with yellow. He has a canine face, the lower parts of which are of a bright vermillion; the snout resembles that of a hog, and the nails are flat, but sharp and very strong. It is said that he follows goats and sheep in order to drink their milk; he partakes of human dexterity in getting the kernels out of nuts, and loves to be covered with garments; he stands upright, and imitates with ease many human actions. The cunning of these animals is well exemplified in this mode of plunder. They form long lines, extending from their retreat to the object in view, and then pitch the produce of their theft from hand to hand till it is secure. Baboons differ from the apes on the one hand, and the monkeys on the other, by having short tails. There is a large specimen in the Surrey Zoological Gardens, known as "Old Peter."



THE PROBOSCIS. THE DIANA MONKEY.

(*Nasalis larratus.*)

(*Cercopithecus Diana.*)


THE Proboscis Monkey is so called from its long projecting and disproportionate nose; it is an inhabitant of the island of Borneo, where it lives in troops, on trees in the vicinity of its rivers. It is of a savage disposition.— The Diana Monkey is called after the goddess of that name, from the crescent of white hair which ornaments its brow. It is very playful, and one of the most graceful of the tribe; and is found in the hottest parts of Africa. Monkeys are less in stature, but more numerous than the apes and baboons. They live almost entirely in trees. Their natural food is vegetable, fruit of all sorts, corn, and even grass; but when domesticated, they learn to eat almost anything that is served on our tables.

There are few persons that are not acquainted with the various mimicries of these animals, and their capricious feats of activity: anecdotes of this kind are too numerous for selection; we shall therefore content ourselves by giving the following:—Captain Stedman, while hunting among the woods of Surinam for provisions, says, that he shot at two of these animals: but that the destruction of one of them was attended with such circumstances, as to ever afterwards deter him from going a monkey hunting. “Seeing me nearly on the bank of the river, in the canoe,” says he, “the creature made a halt from

skipping after his companions, and being perched on a branch that overhung the water, examined me with the strongest marks of curiosity; while he chattered prodigiously, and kept shaking the boughs on which he rested, with incredible strength and agility. At this time I laid my piece to my shoulder and brought him down from the tree: but may I never again be witness to such a scene! The miserable animal was not dead, but mortally wounded. I seized him by the tail, and taking him in both my hands, to end his torment, swung him round, and hit his head against the side of the canoe; but the poor creature still continued to live, and looked at me in the most affecting manner that can be conceived. I therefore knew no other means of ending his murder than to hold him under water till he was drowned: but even in doing this, my heart sickened; for his little dying eyes still continued to follow me with seeming reproach, till their light gradually forsook them, and the wretched animal expired."

The manner in which some of the Monkey tribe capture shell-fish, is remarkably indicative of their cunning and ingenuity. The oysters of the tropical climates, being larger than ours, the Monkeys, when they reach the sea-side, pick up stones, and thrust them between the opening shells, which being thus prevented from closing, the cunning animals eat the fish at their ease. In order to attract crabs, they put their tails before the holes in which they have taken refuge; and when the creatures have fastened on the lure, the Monkeys suddenly withdraw their tails, and thus drag their prey on shore.

The Monkey generally brings forth one at a time, and sometimes two. They are rarely found to breed when brought over into Europe; but those that do, exhibit a very striking picture of parental affection. The male and female are never tired of fondling their young one. They instruct it with no little assiduity; and often severely correct it, if stubborn, or disinclined to profit by their example. They hand it from one to the other, and when the male has done showing his regard, the female takes her turn in the work of affection.



THE CAPUCHIN AND SPIDER MONKEYS


ARE both natives of South America ; they live in large troops, feeding on roots, fruits, and insects, and are much more gentle than those of the old world. Of the *Capuchin* there are many varieties, differing from each other in colour only ; they are very lively, active, and amusing, and about a foot long. The *Spider Monkey*, like the *Capuchin*, has a long prehensile tail, which it uses like a fifth hand, and even employs it in preference to either its hands or feet. Nature seems by this addition to have more than recompensed them for the want of a thumb, for by it, when they are unable to leap from one tree to another, on account of the distance, they form a kind of chain, with their young upon their backs, hanging down by each other's tails. One of them holding the branch above, and the rest swing to and fro like a pendulum, until the undermost is enabled to catch hold ; the first then lets go his hold, and thus comes undermost in his turn ; in this way they can travel a great distance without ever touching the ground. Curious illustrations of this are daily seen at the Zoological Gardens, where there are several of these Monkeys.



THE OUISTITI AND MARIKINA MONKEYS.

THE OUISTITI, or MARMOZET, inhabits the Brazils, and is of small size, not measuring more than seven inches, though his tail is near eleven; he weighs about six ounces, and, like others of his species, lives entirely on vegetables. His face is almost naked, of a swarthy flesh colour, with a white spot above the nose: the tail is full of hair, and annulated with ash colour and black rings alternately; his nails are sharp, and his fingers like those of a squirrel.

The MARIKINA is a beautiful little animal, not above nine inches long, and is sometimes called the Lion Monkey; his hair is long, soft, and glossy; his head is round, his face brown, and his ears hid under the long hairs which surround his face, and which are of a bright red, while those on his body and tail are of beautiful pale yellow, or gold colour. He is very playful, and of a seemingly robust temperament, for we have seen one which lived five or six years in Paris, without any other particular care, than keeping it during the winter in a chamber in which there was a fire every day.



THE LEMUR AND THE MONGOOS

(*Lemur nigrifrons* and *Lemur albifrons*)

MAY be considered as the connecting link between the Monkeys and the genuine quadruped. Their habits are nocturnal, whence they have been called Lemurs, or ghosts. They pass a considerable portion of the day in sleep, rolled up like a ball, with the large tail passed between the hind legs, and twisted round the neck. They live in troops, more or less numerous, like the apes and monkeys, on trees, and climb with great quickness, and leap with so much force, as frequently to rise ten feet at a single bound. They feed on fruits, roots, &c., and carry their food to their mouth with their hands, like the apes; their voice, when not alarmed, is a quick grunt. Their nocturnal and unobtrusive habits may probably account in some degree for the rarity of their appearance. They are all inhabitants of Madagascar, but are widely spread, having been also found in Bengal, and other parts of Hindostan, in Ceylon, and Java. The above specimens are from the Zoological Gardens, and are the White-fronted, and the Black and White Lemurs.

BOOK II.

INHABITANTS OF THE AIR.

§ 1. RAPTORES. *Diurnal Birds of Prey.*



THE GOLDEN EAGLE. (*Aquila chrysaelos.*)

“ But who the various nations can declare,
That plough with busy wing the peopled air?
These cleave the crumbling bark for insect food,
Those dip the crooked beak in kindred blood :
Some haunt the rushy moor, the lonely woods ;
Some bathe their silver plumage in the floods ;
Some fly to man, his household gods implore,
And gather round his hospitable door,
Wait the known call, and find protection there
From all the lesser tyrants of the air.
The tawny Eagle seats his callow brood
High on the cliff, and feasts his young with blood.”

BARRAULD.

THE *Golden Eagle* is the largest and the most powerful of all those birds that have received the name of Eagle.

It weighs above twelve pounds. Its length, from the point of the beak to the end of the tail, is about three feet nine inches; the breadth, when the wings are extended, is eight spans. The beak is horny, crooked, and very strong. The feathers of the neck are of a rusty colour, and the rest nearly black with lighter spots. The feet are feathered down to the claws, which have a wonderful grasp; the leg is yellowish, and the four talons are crooked and strong. As in all birds of prey, the female is the larger, and more powerful.

Eagles are remarkable for their longevity, and their faculty of sustaining a long abstinence from food. Of all birds the Eagle flies highest; and from thence the ancients have given it the epithet of the *bird of heaven*.

“ Bird of the broad and sweeping wing,
 Thy home is high in heaven,
 Where wide the storms their banners fling,
 And the tempest's clouds are driven.
 Thy throne is on the mountain top,
 Thy fields the boundless air;
 And hoary peaks, that proudly prop
 The skies, thy dwellings are.”

This formidable bird may be considered among its own species what the lion is among quadrupeds; and in many respects they have a strong similitude to each other. Solitary, like the lion, he keeps the wilds to himself alone; it is as extraordinary to see two pairs of Eagles in the same mountain, as two lions in the same plain.

The Eagle is found in Great Britain and Ireland, in Germany, and nearly all parts of Europe. It is carnivorous, and, when unable to obtain the flesh of larger animals, feeds on serpents and lizards. The story of the Eagle, brought to the ground after a severe conflict with a cat, which it had seized and taken up into the air with its talons, is very remarkable; Mr. Barlow, who was an eye-witness of the fact, made a drawing of it, which he afterwards engraved. Two instances are said to have occurred in Scotland, of the Eagle having flown

away with infants to its nest: but in both cases it is added that the children were recovered, without being materially injured. This bird has been often tamed, but in this situation it still preserves an innate love of liberty. The nest of the Eagle is composed of strong sticks, covered with rushes, and generally built on the point of an inaccessible rock, whence it darts upon its prey with the rapidity of lightning. The period of incubation is said to be thirty days; and when the young are hatched, both the male and female exert all their industry to provide for their wants. In the county of Kerry, a peasant is said once to have formed the resolution of plundering an Eagle's nest, built upon a small island in the beautiful lake of Killarney. He accordingly swam to the island while the parents were away; and, after robbing the nest of the young, was preparing to swim back, with the Eaglets tied in a string; but while he was yet up to the chin in the water, the old Eagles returned, and, missing their family, fell upon the invader with such fury, that, in spite of all his resistance, they dispatched him with their beaks and talons.





THE SEA EAGLE. (*Haliaeetus albicilla.*)

THIS bird, known also as the White-tailed Eagle, from the inside feathers of its tail being white, differs from the golden eagle in the greater length of its beak, in its sluggish and cowardly habits, and in its coarser taste. It is a native of Great Britain, where it inhabits the high rocks and cliffs that overhang the sea, and whence it pounces on the birds, fish, or seals that it can procure for its prey. It is much smaller than the golden eagle, seldom exceeding twenty-eight inches in length; and in young birds the tail feathers are brown.



THE WHITE-HEADED, OR BALD EAGLE.

(*Haliaetus leucocephalus.*)

THIS bird is about three feet long, and seven feet broad, measuring to the tips of the extended wings. The bill resembles that of the golden eagle, and from the chin hang some small hairy feathers like a beard. As it is found alike in the frigid and the torrid zone, it is provided for enduring rapid changes of temperament, and its whole body is clothed under the feathers with a kind of down, white and soft like that of the swan. This bird builds its nest on some romantic cliff by the sea-shore, or on the bank of some river or lake, and feeds almost entirely upon fish.

It is generally regarded by the Anglo-Americans with peculiar respect, as the chosen emblem of their native land. The great cataract of Niagara is mentioned as one of its favourite places of resort, not merely as a fishing station, where it is enabled to satiate its hunger upon its most congenial food, but also in consequence of the vast quantity of four-footed beasts, which, unwarily venturing

into the stream above, are borne away by the torrent, and precipitated down those tremendous falls.

“ High o'er the watery uproar silent seen,
Sailing sedate in majesty serene,
Now 'midst the pillar'd spray sublimely lost,
And now emerging, down the rapids toss'd,
Glides the Bald Eagle, gazing calm and slow
O'er all the horrors of the scene below ;
Intent alone to sate himself with blood,
From the torn victims of the raging flood.”

The number of birds of prey of various kinds, which assemble at the foot of the rocks to glut themselves upon the banquet thus provided for them, is said to be incredibly great, but they are all compelled to give place to the Eagle when he deigns to feed on dead animals; and the crow and the vulture submit without a struggle to the exercise of that tyranny, which they know it would be in vain to resist. “ We have ourselves,” says Wilson, “ seen the Bald Eagle, while seated on the dead carcase of a horse, keep a whole flock of vultures at a respectful distance, until he had fully sated his own appetite:” and he adds another instance, in which many thousands of tree squirrels having been drowned, in one of their migrations, in attempting to pass the Ohio, and having furnished for some length of time a rich banquet to the vultures, the sudden appearance among them of the Bald Eagle at once put a stop to their festivities, and drove them to a distance from their prey, of which the Eagle kept sole possession for several successive days.

These Eagles sometimes hunt in pairs in a manner which shows their great sagacity. Aware that water-fowl have the power of eluding their grasp by diving, they hover at a distance from each other over their prey. One of them then darts towards it with great swiftness, but the water-fowl easily avoids the first attack by diving. The pursuer then rises into the air and his mate resumes the attack just as the fowl is emerging to breathe, and compels it to plunge again. The Eagles continue alternately to proceed in this manner till their victim is so exhausted that it falls an easy prey.




THE BLACK EAGLE.

SOME ornithologists suppose this to be merely the golden eagle in its young state, but others make it a distinct species. It is about twice as large as the raven. The parts about the beak and the eye are bare of feathers, and somewhat reddish; the head, neck, and breast black; in the middle of the back, between the shoulders, there is a large white spot, dashed with red; a black streak sweeps along the feathers, and is followed by a white one; the remaining part of the wing, to the tip, is of a dark ash-colour. This bird has beautiful hazel eyes, full of animation: his legs are feathered down a little below the tarsal joint, the

naked part being red; his talons are very long. He is found in France, Germany, Poland, and delights in Alpine mountains, where he makes the vales and woods resound with his incessant screamings when in search of prey.

The Abbé Spallanzani had an eagle of this species, so powerful as to be able to kill dogs that were much larger than itself. When a dog was placed before it, the bird would ruffle up the feathers on its head and neck, cast a dreadful look at its victim, take a short flight, and immediately alight on its back. It held the head firmly with one foot, and thus secured the dog from biting, and with the other grasped one of his flanks, at the same time driving its talons into the body; and in this attitude it continued, till the dog expired with fruitless outcries and efforts.

The eyes of Eagles are celebrated for their brilliancy and strength, which has given rise to the popular opinion that they can gaze on the sun without shrinking; though this, from the overhanging eyebrow of the Eagle, would be an extremely difficult feat for the bird to perform. The eyes of all birds are curiously constructed, so as to enable them to see both distant objects and near ones with equal facility; and for this purpose they are furnished with a membrane placed near the edge of the crystalline lens of the eye, by which it can be moved at pleasure. The orbit of the eye is formed of about twelve or sixteen bony plates, which slide over each other when necessary. Birds are also furnished with an additional eyelid, of extremely thin texture, with which they occasionally appear to shade their eyes.

THE CONDOR. (*Vultur Gryphus.*)

THIS bird measures three or four feet long, and its wings, when expanded, from ten to twelve feet. Its bill and talons are exceedingly large and strong; and its courage is equal to its strength. The throat is naked, and of a red colour. The upper parts in some individuals (for they differ greatly in colour) are variegated with black, grey, and white; and the body is scarlet. Round the neck it has a white ruff of loose hairy feathers. The feathers on the back are generally quite black, and perfectly bright. These enormous birds, which are inhabitants of South America, breed among the highest and most inaccessible rocks. The female makes no nest, but lays two white eggs, somewhat bigger than those of a turkey, on the bare rock. Some writers have affirmed that a Condor can carry off a sheep in its claws, and others, that it has carried off children in the same manner; but these tales are manifestly absurd, as the Condor's feet and talons are not fitted for carrying any great weight. Both the talons

and the bill are indeed of extraordinary strength, but they are intended for tearing objects to pieces; and consequently we find that the Condor feeds chiefly on dead or dying cattle, or horses, which he tears to pieces, and devours where they lie. When the Condor is gorged, the hunters attack him, but his strength and fierceness are so great, that one of Sir Francis Head's companions, who attempted to seize a gorged Condor, said he never had "such a battle in his life;" though he had been a Cornish miner, and was reckoned an excellent wrestler in his own country.



THE VULTURE. (*Vultur Papa.*)

THE first rank in the description of birds has been given to the eagle, not on account of its size, but because it is nobler in its habits and more delicate in its appetites. But it belongs to the falcon tribe, and should be placed after the Vultures. The eagle, unless pressed by famine, will not stoop to carrion; and only devours what he has earned by his own pursuit. The Vulture, on the contrary, is disgustingly voracious; and seldom attacks living ani-

imals when it can be supplied with dead. The eagle meets and singly opposes his enemy; the Vulture, if it expects resistance, calls in the aid of its kind, and overpowers its prey by combination. Putrefaction, instead of deterring, only serves to allure it. The Vulture seems among birds what the jackal and hyæna are among quadrupeds, who prey upon carcasses, and root up the dead.

Vultures may be easily distinguished from eagles by the nakedness of their heads and necks, which are without feathers, and only covered with a very slight down, or a few scattered hairs; their eyes are more prominent; those of the eagle being buried more in the socket, and shaded by an overhanging eyebrow. Their claws are shorter and less hooked. The inside of the wing is covered with a thick down, which is different in them from all other birds of prey. Their attitude is not so upright as that of the eagle, and their flight is more difficult and heavy.

In this description we may include the Golden, the Ash-coloured, and the Brown Vulture, which are inhabitants of Europe; the Spotted and the Black Vulture of Egypt; the Bearded Vulture, the Brazilian Vulture, and the King of the Vultures, of South America. They all agree in their nature, being equally indolent, rapacious, and unclean. The Condor also belongs to the Vulture tribe. The King of the Vultures is the species represented in the plate. The head and neck of this bird are without feathers; the body above, reddish buff, beneath, yellowish white; quills greenish black; tail black; craw pendulous, and orange-coloured. It is about the size of a turkey; and is chiefly remarkable for the odd formation of the skin of the head and neck; this skin, which is of an orange colour, arises from the base of the bill, whence it stretches on each side to the head; the eyes are surrounded by a red skin, and the iris has the colour and lustre of pearl. Upon the naked part of the neck is a collar formed by soft longish feathers. Into this collar the bird sometimes withdraws its whole neck, and sometimes a part of its head, so that it looks as if it had hidden its neck in its body.



THE BUZZARD. (*Falco Buteo.*)

“ The noble Buzzard ever pleased me best ;
Of small renown, 't is true ; for, not to lie,
We call him but a Hawk by courtesy.”

HIND AND PANTHER.

THIS is a rapacious bird, of the long-winged hawk kind, and the most common of all in England. It is of a sluggish indolent nature, often remaining perched on the same bough for the greater part of the day ; as if, indifferent either to the allurements of food or of pleasure, it were doomed, like some of the human species, to pass its allotted span of life in passive contemplation. It feeds on mice, rabbits, frogs, and often on all sorts of carrion. Too idle to build itself a nest, it generally seizes upon the old habitation of a crow, which it lines afresh with wool and other soft materials. In general this bird, whose colour varies considerably, is brown, varied with yellow specks ; at a certain age its head becomes entirely grey. The female generally lays two or three eggs, which are mostly white, though sometimes spotted with yellow. Its length is usually twenty-two inches, and its breadth upwards of fifty.

The following anecdote related by Buffon, will show that the Buzzard may be so far tamed as to be rendered a faithful domestic. A Buzzard which had been caught in

a snare, was brought to a gentleman, who undertook to tame it. It was at first wild and ferocious, but he succeeded, by depriving it of food and constraining it to come and eat out of his hand. By pursuing this plan, he brought it to be very familiar; and, after having shut it up about six weeks, he began to allow it a little liberty, taking the precaution, however, to tie both pinions of its wings. In this condition it walked out into his garden, and returned when called to be fed; after some time, thinking he might trust to its fidelity, he removed the ligatures, and fastened a small bell above its talon, and also attached to its breast a bit of copper with his name engraved on it. He then gave it entire liberty, which it soon abused; for it took wing and flew into the forest of Belesme. The bird was given up for lost; but four hours afterwards, it rushed into the gentleman's hall, pursued by five other Buzzards, which had constrained it to seek its former asylum. After this adventure it preserved its fidelity, coming every night to sleep under the window. It soon became familiar, attended constantly at dinner, sat on a corner of the table, and often caressed its master with its head and bill, emitting a weak, sharp cry, which, however, it sometimes softened. It had a singular propensity of seizing from the head and flying away with the red caps of the peasants; and so alert was it in whipping them off, that they found their heads bare without knowing what was become of their caps: it even treated the wigs of the old men in the same way, hiding its booty in the tallest trees.

Wilson says that one he shot in the wing lived with him several weeks; but refused to eat. It amused itself by hopping from one end of the room to the other, and sitting for hours at the window, looking down on the passengers below. At first, he put himself in an attitude of defence when approached; but after some time became quite familiar, permitting himself to be handled. Though he lived so long without food, his stomach was found on dissection to be enveloped in solid fat of nearly an inch in thickness.



THE HONEY-BUZZARD. (*Falco-pivorus*, Penn.)

THIS Buzzard feeds upon honey and upon the larvæ of bees and wasps. It is also said to eat lizards, frogs, and snails. Buffon says that in winter, when fat, it is good eating, a very rare circumstance with birds of this genus. It seldom flies excepting from one bush to another; but when on the ground it runs with great rapidity, like a domestic fowl.

Willoughby observes, that it builds its nest with twigs, on which it lays wool to receive its eggs. He saw one that took possession of an old kite's nest to breed in, and that fed its young with the nymphæ of wasps, for in the nest were found the combs of wasps' nests, and in the stomachs of the young fragments of wasp-maggots. In the nest were two young ones, covered with white down, spotted with black. In the crop of one of them were two lizards entire, with their heads lying towards the mouth, as if they sought to creep out.


It would be highly interesting could we discover the manner in which this bird conducts its attack on a wasps' nest. The close feathering round the base of the bill is, no doubt, a protection against the stings of the insects which they attack.



THE GOSHAWK (*Falco*, or *Astur palumbarius*)

BREEDS in lofty trees in Scotland, and destroys a great quantity of small game, which he seizes with his sharp and crooked talons, and carries to his nest. He is of the falcon tribe; and somewhat larger than the common buzzard, his bill is blue, and he has a white stripe over each eye, and also a large white spot on each side of the neck. The general colour of the plumage is deep brown; the breast and belly white, transversely streaked with black; and the legs yellow. Buffon, who brought up two young Goshawks, a male and a female, makes the following observations: "The Goshawk, before it has shed its feathers, that is in the first year, is marked on the breast and belly with longitudinal brown spots; but after it has had two moultings they disappear, and their place is occupied by transverse bars, which continue during the rest of its life." He further observes, "that though the male was much smaller than the female, it was fiercer and more vicious." The Goshawk is found in France and Germany; it is not very common in England, but is more so in Scotland. In former times, the custom of carrying a Hawk or Falcon on the hand was confined to men of high distinction; so that it was a saying among the Welsh, "You may know a gentleman by his Hawk, horse, and greyhound." Even the ladies in those times were partakers of this gallant sport, and have been represented in pictures with Hawks on their hands. At present, hawking is almost entirely laid aside in this country,

as the expense which attended it being very considerable, confined it to princes and men of the highest rank. In the time of James the First, Sir Thomas Monson is said to have given a thousand pounds for a cast of Hawks. In the reign of Edward the Third it was made felony to steal a Hawk; to take its eggs, even in a person's own grounds, was punishable with imprisonment for a year and a day, together with a fine at the king's pleasure. Such was the delight our ancestors took in this royal sport, and such were the means by which they endeavoured to secure it. The Falcons, or Hawks, chiefly used in these kingdoms, were the Goshawk, the Falcon, the Peregrine Falcon, Iceland Falcon, and the Ger Falcon. The game usually pursued were cranes, wild geese, pheasants, and partridges. The Duke of St. Albans is still hereditary grand falconer of England, but the office is not now exercised, except for the Duke's own amusement.



THE SPARROWHAWK. (*Falco nisus*.)

THE Sparrowhawk is a bold-spirited bird, the length of the male is twelve inches, that of the female fifteen; the beak

is short, crooked, and of a bluish tint, but very black towards the tip; the tongue black, and a little cleft; the eyes of a middling size. The crown of the head is of a dark brown: above the eyes, in the hinder part of the head, there are sometimes white feathers; the roots of the feathers of the head and neck are white; the rest of the upper side, back, shoulders, wings, and neck, of a dark brown. The wings, when closed, scarcely reach to the middle of the tail; the thighs are strong and fleshy, the legs long, slender, and yellow; the toes also long, and the talons black. The female, which is, as in other birds of prey, much larger than the male, lays about five eggs, spotted near the blunt end with blood-red specks. When wild, they feed only upon birds, and possess a boldness and courage above their size; but in a domestic state they do not refuse raw flesh and mice. They can be made obedient and docile, and readily trained to hunt quails and partridges.






THE KITE. (*Falco Milvus.*)

THIS bird, though it belongs to the falcon tribe, is called ignoble, because it is never used in hawking. It is easily distinguished from all the other birds of prey by its forked tail, and the slow and circular eddies it describes in the air, whenever it spies from the regions of the clouds a young duck or a chicken which has strayed too far from the brood. When this is the case, the Kite, pouncing on it with the rapidity of a dart, seizes it in its talons, and carries it off to its nest. It is, however, a great coward, and if the hen flies at it, which she always does if she sees it, it will drop the chicken and fly off. It is larger than the common buzzard; and though it weighs somewhat less than three pounds, the extent of its wings is more than five feet. The head and back are of a pale ash-colour, varied with longitudinal lines across the shafts

of the feathers; the neck is reddish; the lesser rows of the wing feathers are party-coloured, of black, red, and white; the feathers covering the inside of the wings are red, with black spots in the middle. The eyes are large, the legs and feet yellow, the talons black. It is a handsome bird, and seems almost always on the wing. It rests itself on the air, and does not appear to make the smallest effort in flying, but rather to glide along with the gentlest breeze.



THE FALCON. (*Falco communis*.)

THE Falcon is a predaceous bird, of which there are several species. Of these the *Gerfalcon* is the largest, and is found in the northern parts of Europe; and, next to the eagle, is the most formidable, active, and intrepid of all voracious birds, and the most esteemed for falconry. The bill is crooked and yellow; the irides of the eye dusky; and the whole plumage of a whitish hue, marked with dark lines on the breast, and dusky spots on the back.

The *Peregrine Falcon*, which is the most common kind, is as large as the moor buzzard. The bill is blue at the base, and black at the point; the head, back, scapulars, and coverts of the wings are barred with deep black and blue; the throat, neck, and upper part of the breast are white, tinged with yellow; the bottom of the breast,

belly, and thighs are of a greyish white; and the tail is black and blue. Wilson enumerates no less than ten varieties, dependent chiefly upon age, sex, and country. It is found, more or less abundantly, throughout the whole of Europe, principally in the mountain districts in North and South America, and in New Holland, dwelling in the clefts of rocks, especially such as are exposed to the mid-day sun. It breeds upon the cliffs in several parts of England, but appears to be more common in Scotland and Wales. Its food consists principally of small birds; but it scruples not to attack the larger species, and sometimes gives battle even to the kite. Falcons rarely take their prey upon the ground, like the more ignoble birds of the class to which they belong; but pounce upon it from aloft, in a directly perpendicular descent as it flies through the air, bear it downwards by the united impulse of the strength and rapidity of their attack, and sticking their talons into its flesh, carry it off in triumph to the place of their retreat. Like most predatory animals, they are stimulated to action by the pressure of hunger alone, and remain inactive and almost motionless while the process of digestion is going on, until the renewed cravings of their appetite stimulate them to further exertion. In different stages of its growth, the Peregrine Falcon has been known by various English names. Its proper appellation among falconers is the Slight Falcon, the term Falcon Gentle being equally applicable to all the species when rendered manageable. In the immature state, this Falcon is also called a Red Hawk, from the prevailing colour of its plumage. The male is called a Tiercel, to distinguish it from the female, which, in the Falcon tribe, is commonly one-third larger than the male.

In China there is said to be a variety, which is mottled with brown and yellow, and used by the emperor of China in his sporting excursions, when he is usually attended by his great falconer, and a thousand of inferior rank. Every bird has a silver plate fastened to its foot, with the name of the falconer who has the charge of it, that, in case it should be lost, it may be restored to the proper person;

but if it should not be found, the name is delivered to another officer, called the guardian of lost birds, who, to make his situation known, erects his standard in a conspicuous place among the army of hunters.

In Syria, there is a variety of the Gentle Falcon, which the inhabitants call Shaheen, and which is of so fierce and courageous a disposition, that it will attack any bird, however large or powerful, which presents itself. "Were there not," says Dr. Russel, in his Account of Aleppo, "several gentlemen now in England to bear witness to the fact, I should hardly venture to assert that, with this bird, which is about the size of a pigeon, the inhabitants sometimes take large eagles. This Hawk was in former times taught to seize the eagle under the pinion, and thus depriving him of the use of one wing, both birds fell to the ground together; but the present mode is to teach the Hawk to fix on the back, between the wings, which has the same effect, only that as the bird tumbles down more slowly, the falconer has more time to come to his Hawk's assistance; but in either case, if he be not very expeditious, the Falcon is inevitably destroyed. I never saw the Shaheen fly at eagles, that sport having been disused before my time; but I have often seen him take herons and storks. The Hawk, when thrown off, flies for some time in a horizontal line, not six feet from the ground; then mounting perpendicularly, with astonishing swiftness, he seizes his prey under the wing, and both together come tumbling to the ground."

Audubon states, that he has seen this bird come at the report of a gun, and carry off a teal, not thirty paces distant from the sportsman who had killed it, with a daring assurance as surprising as unexpected; and adds, that this conduct is a notorious characteristic of the species.



THE MERLIN (*Falco aesalon*)

Is the least of the Falcon tribe, and, as its name implies, is not very different in size from the blackbird; the word Merlin signifying in French a small *merle*, or blackbird. Though small, the Merlin is not inferior in courage to any of the other hawks; it is noted for its boldness and spirit, often attacking and killing at one stroke a full-grown partridge or a quail; but it differs from the Falcons and all the other rapacious kinds, in the male and female being of equal size. The back of this bird is party-coloured, of dark blue and brown; the quill-feathers of the wings black, with rusty spots; the tail is about five inches long, of a dark brown or blackish colour, with transverse white bars: the breast is of a yellowish white, with streaks of rusty brown pointing downwards; the legs are long, slender, and yellow; the talons black. The head is encircled with a row of yellowish feathers, not unlike a coronet. In the male the feathers on the rump, next the tail, are bluer; a mark by which the falconers easily discern the sex of the bird. The Merlin does not breed here, but visits us in October: it flies low, and

with great celerity and ease. In the days of falconry, the Merlin was considered the lady's hawk.


"In ancient days—in ancient days,
When ladies took a strange delight
In hawks and hounds and sporting ways,
A Merlin was a pleasant sight.

'T was gentle when, in trappings gay,
Upon its lady's wrist it stood ;
Till its hood was raised and it saw its prey,
When its eye betrayed the bird of blood."



THE HEN HARRIER (*Circus cyaneus*)

Is seen about forests, heaths, and other retired places, especially in the neighbourhood of marshy grounds, where it destroys vast numbers of snipes, woodcocks, and wild ducks. It is about seventeen inches long, and three feet wide; its bill is black, and cere yellow. The upper part of its body is of a bluish grey; and the back of the head, breast belly, and thighs are white. The legs are long, slender, and yellow; and the claws black.



THE OSPREY, OR FISHING HAWK.

(*Falco, or Pandion Haliaetus.*)

“ True to the season, o'er our sea-beat shore,
 The sailing Osprey high is seen to soar
 With broad unmoving wing ; and circling slow,
 Marks each loose straggler in the deep below ;
 Sweeps down like lightning, plunges with a roar,
 And bears his struggling victim to the shore.”

THIS bird is always found on the sea-shore, or near rivers or lakes, as it feeds entirely on fish. It is common in Great Britain, and also in America, where large colonies of it are found, the birds living together like rooks. “ When looking out for its prey,” says Dr. Richardson, “ it sails with great care and elegance, in undulating and curved lines, at a considerable height above the water, till it perceives its prey, when it pounces down upon it. It seizes the fish with its claws, sometimes scarcely appearing to dip its feet in the water, and at others plunging entirely under the surface with force sufficient to throw up a considerable spray. It emerges again, however, so speedily, as to render it evident that it does not attack fish swimming at any great depth.”

The Osprey builds a large nest either on trees or rocks, and lays two or three eggs, which have a reddish tinge, and are spotted with brown at the larger end. The old birds feed the young ones even after they have left the nest, and only rear one brood in the year.

SECTION II.—Nocturnal Birds of Prey.



THE HORNERD OWL (*Bubo Virginianus*)

Is one of the largest of the Owls, and has two long tufts growing from the top of its head, above its ears, and composed of six feathers, which it can raise or lay down at pleasure. The eyes are large, and encircled with an orange-coloured iris; the ears are large and deep, and

the beak black; the breast, belly, and thighs are of a dull yellow, marked with brown streaks; the back, coverts of the wings, and quill-feathers are brown and yellow; and the tail is marked with dusky and red bars. It inhabits the north and west of England, and Wales. The conformation of the organ of sight in the Owl, is so peculiar and so much in its nature resembling that of the feline kind, that it can see much better at dusk than by daylight. The Barn Owl sees in a greater degree of darkness than the others; and, on the contrary, the Horned Owl is enabled to pursue his prey by day, though with difficulty. Owls are sometimes tamed by persons in the country, who carefully rear them in a domestic state, from their propensity to chase and devour mice and other vermin, of which they clear the houses with as much address as cats. The Owl is a solitary bird, and is said to retire into holes in towers and old walls in the winter, and pass that season in sleep.

“The solitary bird of night,
Through the pale shade now wings his flight,
And quits the time-shook tower;
Where, shelter'd from the blaze of day,
In philosophic gloom he lay,
Beneath his ivy bower.”

CARTER.

The Harfang, or great Snowy Owl, (*Surnia nyctea*) is another species which takes its prey occasionally by daylight. It is seldom seen in England, but frequently visits North Britain, particularly the Orkney and Shetland Islands. It is the only kind of Owl that is known to feed on fish, into which it strikes its talons while in the water, and carries them off to its nest. These Owls are very common in the northern parts of North America, and are eaten not only by the Indians, but by the Europeans engaged in the fur-trade.



THE WHITE, BARN, OR SCHREECH OWL.

(*Srix flammea.*)

“—— from yonder ivy-mantled tower
 The moping Owl does to the moon complain
 Of such as, wandering near her secret bower,
 Molest her ancient solitary reign.”

GRAY.

THIS bird is about the size of a pigeon. Its beak, hooked at the end, is more than an inch and a half long. There is a circle or wreath of white, soft, and downy feathers, encompassed with yellow ones, beginning from the nostrils on each side, passing round the eye and under the chin, somewhat resembling the hood that women used to wear; so that the eyes appear to be sunk in the middle of the feathers. The breast and the feathers of the inside of the wings are white, and marked with a few dark spots; the upper parts of the body are of a fine pale yellow colour, variegated with white spots. The legs are

covered with a thick down to the feet, but the toes have only thin-set hairs around them.

In ancient mythology, this bird was consecrated to Minerva, the goddess of wisdom; in allusion to the lucubrations of wise men, who study in retirement and during the night.

“ Now the Hermit Owlet peeps
From the barn, or twisted brake;
And the blue mist slowly creeps,
Curling on the silver lake.” CUNNINGHAM.

SECTION III.—Insectores, or Perching Birds.



THE BUTCHER-BIRD, OR SHRIKE.

(*Lanius excubitor.*)

THE Great Butcher-bird, or Shrike, is about as large as a thrush; its bill is black, an inch long, and hooked at the end. It is only an occasional visitor to this country, where it is generally found between autumn and spring. “The Shrike,” says Mr. Yarrell, “feeds on mice, shrews, small birds, frogs, lizards, and large insects. After having killed its prey, it fixes the body in a forked branch, or upon a sharp thorn, the more readily to tear off small pieces from it. It is from their habit of killing and hanging up their meat, that the Shrikes are called Butcher-birds.” The head, back, and rump are ash-coloured; the chin and lower part of the body white; the breast

and throat varied with dark lines, crossing each other; the tips of the feathers of the wings are, for the most part, white; it has a black spot by the eye; the utmost feathers of the male are all over white; the two middlemost have only their tips white, the rest of the feathers being black, as well as the legs and feet. It builds its nest among thorny shrubs and dwarf trees, and furnishes it with moss, wool, and downy herbs, where the female lays five or six eggs. A peculiarity belonging to the birds of this kind is, that they do not, like most other birds, expel the young ones from the nest, as soon as they can provide for themselves, but the whole brood live together in one family. The Butcher-bird will chase all the small birds upon the wing, and will sometimes venture to attack partridges, and even young hares. Thrushes and blackbirds are frequently their prey: the Shrike fixes on them with its talons, splits the skull with its bill, and feeds on them at leisure. On this account Linnæus classed the Shrikes with the birds of prey; but modern naturalists have placed them with the insect-eaters, as insects are their principal food. It is easy to distinguish these birds at a distance, not only from their going in companies, but also from their manner of flying, which is always up and down, seldom in a direct line, or obliquely.

The Little Butcher-bird, called in Yorkshire, *Flusher*, is about the size of a lark, with a large head. About the nostrils and corners of the mouth it has black hairs or bristles; and round the eyes a large black longitudinal spot; the back and upper side of the wings are of a rusty colour; the head and rump cinereous; the throat and breast white, spotted with red. It builds its nest of grass; and the female lays six eggs, nearly all white, except at the blunt end, which is encircled with brown or dark red marks. The female is somewhat larger than the male; the head is of a rust colour, mixed with grey; the breast, belly, and sides of a dirty white; the tail deep brown; the exterior web of the outer feathers white. Its manners are similar to those of the large Butcher-bird, like which, it imitates the notes of other birds, in order the more surely to decoy them. It frequently preys on young birds, which

it takes in the nest; it likewise feeds on grasshoppers, beetles, and other insects. During the period of incubation, the female soon discovers herself at the approach of any person by her loud and violent outcries.



THE WATER OUZEL, OR DIPPER,
(*Cinclus aquaticus*,)

Is found in most parts of this island, and is about the size of the common blackbird. It feeds upon aquatic insects, and small fish, such as sand-eels and minnows. The head and upper side of the neck are of a kind of umber colour, and sometimes black with a shade of red; the back and coverings of the wings are a mixture of black and ash-colour, the throat and breast perfectly white.

The Dipper is said to walk along the bottom of a lake or river, as easily as on land; but this is far from being the case, as though it readily plunges into the water, it appears to tumble about in a very extraordinary manner, with its head downwards. Even on land the bird walks awkwardly, as its feet are best adapted for the slippery stones on which it passes the greater part of its life, watching for the insects which it picks up on the edge of the water. When disturbed, it usually flirts up its tail, and makes a chirping noise. Its song in spring is said to be very pretty. In some places this bird is supposed to be migratory.



THE BLACKBIRD. (*Turdus Merula.*)

“The smiling morn, the breathing spring,
 Invite the tuneful birds to sing;
 And, while they warble from each spray,
 Love melts the universal lay.” MALLETT.

THIS well-known songster does not soar up to the clouds, like the lark, to make his voice resound through the air; but keeps to the shady groves, which he fills with his melodious notes. Early at dawn, and late at dusk, he continues his pleasing melody; and when incarcerated in the narrow space of a cage, still cheerful and merry, he strives to repay the kindness of his keeper by singing to him his natural strains; and beguiles his irksome hours of captivity, by studying and imitating his master's whistle. Blackbirds build their nest with great art, making the outside of moss and slender twigs, cemented together and lined with clay, and covering the clay with soft materials, as hair, wool, and straw. The female lays four or five eggs, of a bluish green colour, spotted all over with brown. The bill is yellow, but in the female the upper part and point are blackish; the inside of the mouth, and the circumference of the eyelids, are yellow. The name of this bird is sufficiently expressive of the general colour of his body. He feeds on berries, fruit, insects, &c. The species found in Surinam and South America is not of so deep a black; and the throat and part of the breast are of a crimson colour.



THE MISSEL THRUSH. (*Turdus viscivorus.*)

THE Missel Thrush, so called from its feeding on the berries of the misletoe, differs but little from the Song Thrush, except in size. He is larger than the fieldfare, while the Thrustle is smaller. The female lays five or six bluish eggs, with a tint of green, and marked with dusky spots.

The Song Thrush or Thrustle is one of the best songsters of the evening hymn in the grove. His voice is loud and sweet; the melody of his song is varied, and, although not so deep in the general diapason of the woodland concert as that of the blackbird, yet it fills up agreeably, and hursts through the inferior warblings of smaller performers. His breast is of a yellowish white, spotted with black or brown dashes, like ermine spots.

The term *Merle* for the Blackbird, and *Mavis* for the Thrush, are used chiefly by the poets.

“Merry is it in the good green wood,
 When the Mavis and Merle are singing,
 When the deer sweeps by, and the hounds are in cry,
 And the hunter's horn is ringing.”

SCOTT.

“Take thy delight in yonder goodly tree,
 Where the sweet Merle and warbling Mavis be.”

DRAYTON.



THE REDWING (*Turdus iliacus*)

Is rather less than the song thrush; but the upper part of the body is of the same colour; the breast not so much spotted; the coverings of the feathers of the under side of the wings, which in the thrush are yellow, are of orange colour in this bird; by which marks it is generally distinguished. The body is white, the throat and breast yellowish, marked with dusky spots. It is migratory in this island, builds its nest in hedges, and lays six bluish eggs. Like the fieldfare, it leaves us in spring, for which reason its song is quite unknown to us; but it is said to be very pleasing. It is delicate eating; and the Romans held it in such estimation, that they kept thousands of them together in aviaries, and fed them on a sort of paste made of bruised figs and flour, to improve the delicacy and flavour of their flesh. Under this management these birds fattened, to the great profit of their proprietors, who sold them to Roman epicures for three denarii, or about two shillings sterling each, which at that early period was a large price.



THE FIELDFARE (*Turdus pilaris*)


Is a well-known bird in this country. Fieldfares fly in flocks, together with the redwing and starlings, and change their haunts according to the season of the year. They abide with us in winter, and disappear in spring, so punctually, that after that time not one is to be seen. The flesh is esteemed a great delicacy, and is highly prized in Germany, where it is known as the *kramms vogel*, and is sold in the markets of Westphalia by the dozen. Their favourite food is the juniper-berry, whence its German name. The head is ash-coloured, and spotted with black: the back and coverts of the wings of a deep chesnut colour; the rump cinereous; and the tail black, except the lower part of the two middle feathers, which are ash-coloured, and the upper sides of the exterior feathers, which are white. They collect in large flocks; and it is supposed they keep watch, like the crow, to mark and announce the approach of danger. On any person approaching a tree that is covered with them, they continue fearless, till one at the extremity of the bush, rising on its wings, gives a loud and peculiar note of alarm. They then all fly away, except one which continues till the person approaches still nearer, to certify, as it were, the reality of the danger, and afterwards he also flies off, repeating the note of alarm.

Mr. Knapp, in his "Journal of a Naturalist," says, that in the county of Gloucestershire the extensive lowlands of the river Severn, in open weather, are visited by prodigious flocks of these birds.



THE RING OUZEL. (*Turdus torquatus.*)

THE Ring Ouzel differs from the fieldfare and redwing, to which it is nearly allied, in being a summer visitor to the British islands, instead of a winter one. It is found only in the wildest and most mountainous districts; particularly among the Welsh mountains and on Dartmoor, in Devonshire, where it has been known to breed.



THE ROBIN, OR REDBREAST.
(*Rubecula familiaris.*)

“ The Redbreast oft, at evening hours,
Shall kindly lend his little aid,

With hoary moss, and gathering flowers,
To deck the ground where thou art laid."

COLLINS.

THE Redbreast, or *Robin*, as he is popularly called, seems always to have enjoyed the protection of man, more than any other bird. The prettiness of his shape, the beauty of his plumage, the quickness of his motions, his familiarity with us in winter, and, above all, the melody and sweetness of his voice, claim our admiration, and have insured him that security which he enjoys among us; though the aid of fable has also been called in, to guard him from the assaults of thoughtless boys.

" Little bird with bosom red,
Welcome to my humble shed!
Courtly domes of high degree
Have no room for thee and me;
Pride and pleasure's fickle throng
Nothing mind an idle song.
Daily near my table steal,
While I pick my scanty meal;
Doubt not, little though there be,
But I'll cast a crumb for thee;
Well rewarded if I spy
Pleasure in thy glancing eye;
And see thee, when thou 'st eat thy fill,
Plume thy breast, and wipe thy bill."

LANGHORNE.

In the winter season, impelled by the potent stimulus of hunger, the Redbreast frequents our barns, gardens, and houses, and often alights, on a sudden, on the rustic floor; where, with his broad eye incessantly open, and looking askew upon the company, he picks up eagerly the crumbs of bread that fall from the table, and then flies off to the neighbouring bush, where, by his warbling strains, he expresses his gratitude for the liberty he has been allowed. He is found in most parts of Europe, but nowhere so commonly as in Great Britain. His bill is dusky; his forehead, chin, throat, and breast are of a deep orange-colour, inclining to vermilion; the back of his head, neck, back, and tail are of ash-colour, tinged

with green; the wings are somewhat darker, the edges inclining to yellow; the legs and feet are the colour of the bill. The female generally builds her nest in the crevice of some mossy bank, near places which human beings frequent, or in some part of a human dwelling. Robins have been known to build in a sawpit, where men worked every day, and in various other equally extraordinary places. The elegant poet of *The Seasons* gives us a very exact and animated description of this bird in the following lines:

“————— Half afraid, he first
Against the window beats: then, brisk alights
On the warm hearth; then, hopping on the floor,
Eyes all the smiling family askance,
And pecks, and starts, and wonders where he is,
Till, more familiar grown, the table crumbs
Attract his slender feet.”

An old Latin proverb tells us, that two Robin Redbreasts will not feed on the same tree; it is certain that the Redbreast is a most pugnacious bird, and that he does not live in much harmony and friendship with those of his own kind and sex. The male may be known from the female by the colour of his legs, which are blacker.

The Redbreast attends the gardener when digging his borders; and will, with great familiarity and tameness, pick out the worms almost close to his spade.





THE NIGHTINGALE. (*Philomela luscina.*)

“ Sweet bird, that shunn’st the noise of folly,
 Most musical, most melancholy !
 Thee, chantress, oft, the woods among,
 I woo to hear thy even song.” MILTON.

THE Nightingale has little to boast of in respect to plumage, which is of a pale tawny colour on the head and back, dashed with a slight shade of olive; the breast and upper part of the belly incline to a greyish tint, and the lower part of the belly is almost white; the exterior web of the quill feathers is of a reddish brown; the tail of a dull red; the legs and feet ash-coloured; the irides hazel; and the eyes large, bright, and staring. But it is hardly possible to give an idea of the extraordinary power which this small bird possesses in its throat, as to extension of sound, sweetness of tone, and versatility of notes. Its song is composed of several musical passages, each of which does not continue more than the third part of a minute; but they are so varied, the passing from one tone to another is so fanciful and so rapid, and the melody so sweet and so mellow, that the most consummate musician is pleasingly led to a deep sense of admiration at hearing it. Sometimes joyful and merry, it runs down the diapason with the velocity of lightning, touching the

treble and the bass nearly at the same instant; at other times, mournful and plaintive, the unfortunate *Philomela* draws heavily her lengthened notes, and breathes a delightful melancholy around. These have the appearance of sorrowful sighs; the other modulations resemble the laughter of the happy. Solitary on the twig of a small tree, and cautiously at a certain distance from the nest, where the pledges of his love are treasured under the fostering breast of his mate, the male fills constantly the silent woods with his harmonious strains; and during the whole night entertains and repays his female for the irksome duties of incubation. The Nightingale not only sings at intervals during the day, but waits till the blackbird and the thrush have uttered their evening call, even till the stock and ring doves have, by their soft murmurings lulled each other to rest, and then pours forth his full tide of melody.

“————— Listening *Philomela* deigns
To let them joy, and purposes, in thought
Elate, to make her night excel their day.”

THOMSON.

It is a great subject of astonishment that so small a bird should be endowed with such potent lungs. If the evening is calm, it is supposed that its song may be heard above half-a-mile. This bird, the ornament and charm of our spring and early summer evenings, as it arrives in April, and continues singing till June, disappears on a sudden about September or October, when it leaves us to pass the winter in the North of Africa and Syria. Its visits to this country are limited to certain counties, mostly in the south and east; as, though it is plentiful in the neighbourhood of London, and along the south coast in Sussex, Hampshire, and Dorsetshire, it is not found in either Cornwall or Wales. As soon as the young are hatched, the song of the male bird ceases, and he only utters a harsh croak, by way of giving alarm when any one approaches the nest. Nightingales are sometimes reared up, and doomed to the prison of a cage; in this

state they sing ten months in the year, though in their wild life they sing only as many weeks. Bingley says that a caged Nightingale sings much more sweetly than those which we hear abroad in the spring.

The Nightingale is the most celebrated of all the feathered race for its song. The poets have in all ages made it the theme of their verses; some of these we cannot resist giving :

“ The Nightingale, as soon as April bringeth
 Up her rested sense a perfect waking,
 Which late bare earth, proud of new clothing, springeth,
 Sings out her woes ———.”—SIR PHILIP SIDNEY.

“ ————Beast and bird,
 They to their grassy couch, these to their nests,
 Were slunk; all but the wakeful Nightingale;
 She all night long her amorous descant sung.”
 MILTON.

“ And in the violet-embroidered vale,
 Where the lovelorn Nightingale
 Nightly to thee her sad song mourneth well.”
 MILTON.

“ O Nightingale, that on yon bloomy spray
 Warblest at eve, when all the woods are still,
 Thou with fresh hope the lover's heart dost fill,
 While the jolly hours lead on propitious May,
 Thy liquid notes that close the eye of day,
 First heard before the shallow cuckoo's bill,
 Portend success in love. Oh, if Jove's will
 Have link'd that amorous power to thy soft lay,
 Now timely sing, ere the rude bird of hate
 Foretell my hopeless doom in some grove nigh;
 As thou from year to year hast sung too late
 For my relief, yet hadst no reason why:
 Whether the muse, or love, call thee his mate,
 Both them I serve, and of their train am I.”
 MILTON.

“ ——— Now is the pleasant time,
 The cool, the silent, save where silence yields
 To the *night-warbling bird*, that, now awake,
 Tunes sweetest his love-laboured song.”
 MILTON.


“ How all things listen while thy muse complains,
Such silence waits on Philomela's strains,
In some still evening, when the whispering breeze
Pants on the leaves, and dies upon the trees.”

POPE.

“ There's a bower of roses by Bendemeer's stream,
And the Nightingale sings round it all the year long ;
In the days of my childhood, 't was like a sweet dream
To sit in the roses, and hear the bird's song.

“ That bower and its music I never forget,
But oft when alone, in the bloom of the year,
I think, Is the Nightingale singing there yet ?
Are the roses still bright by the calm Bendemeer ?”

MOORE.




THE BLACK-CAP (*Sylvia atricapilla*)

Is a very small warbler, not weighing above half-an-ounce. The top of the head is black, whence he takes his name ; the neck ash-coloured, the back a dark green, the wings of a dusky colour, with green edges ; the tail nearly the same ; the nether part of the neck, throat, and upper part of the breast of a pale ash-colour ; the lower part of the belly white tinged with yellow.

The Black-cap visits us about the middle of April, and retires in September ; it frequents gardens, and builds its nest near the ground. The female lays five eggs of a pale

reddish brown, sprinkled with spots of a darker colour. This bird sings sweetly, and so like the nightingale, that in Norfolk it is called the mock nightingale. White observes, that it has usually a full, sweet, deep, loud, and wild pipe, yet the strain is of short continuance, and its motions desultory; but when it sits calmly, and earnestly engages in song, it pours forth very sweet but inward melody; and expresses a great variety of modulations, superior perhaps to any of our warblers, the nightingale excepted; and while it sings, its throat is greatly distended.



THE WREN. (*Troglodytes vulgaris*.)

“Fast by my couch, congenial guest,
The Wren has wove her mossy nest;
From busy scenes and brighter skies
To lurk with innocence she flies;
Her hopes in safe repose to dwell,
Nor aught suspects the sylvan cell.”

T. WARTON.

THE Wren is a very small bird; indeed, one species, the golden crested, is the smallest in Europe; but as if nature had intended to compensate the want of size and bulk in the individuals, by multiplying them to a greater extent, this little bird is one of the most prolific of the feathered tribe, its nest containing often upwards of eighteen eggs of a whitish colour, and not much bigger than a pea. The male and female enter by a hole con-


trived in the middle of the nest, and which, by its situation and size, is accessible only to themselves. The Wren weighs no more than three drachms. Its notes are very sweet, and rival those of the robin redbreast, in the middle of winter, when the coldness of the weather has condemned the other songsters to silence. Like the redbreast it frequently approaches the habitation of man, enlivening the rustic garden with its song during the greater part of the year. It begins to make a nest early in the spring, but frequently deserts it before it is lined, and searches for a more secure place. The Wren does not, as is usual with most other birds, begin to build the bottom of the nest first. When against a tree, its primary operation is to trace upon the bark the outline, and thus to fasten it with equal strength to all parts. It then, in succession, closes the sides and top, leaving only a small hole for entrance.

THE WILLOW WREN. (*Sylvia trochilus*.)

The Willow Wren is somewhat larger than the common Wren. The upper parts of the body are of a pale olive-green; the under parts are pale yellow, and a streak of yellow passes over the eyes. The wings and tail are brown, edged with yellowish green; and the legs are inclined to yellow. This bird is migratory, visiting us usually about the middle of April, and taking its departure towards the end of September. The female constructs her nest in holes at the roots of trees, in hollows of dry banks, and other similar places. It is round, and not unlike the nest of the Wren. The eggs are dusky white, marked with reddish spots; and are five in number. A Willow Wren had built in a bank of one of the fields of Mr. White near Selborne. This bird, a friend and himself observed as she sat in her nest, but were particularly careful not to disturb her, though she eyed them with some degree of jealousy. Some days afterwards, as they passed the same way, they were desirous of remarking how the brood went on; but no nest could be found, till Mr. White happened to take up a large bundle of

long green moss, which had been thrown, as it were, carelessly over the nest, in order to mislead the eye of any impertinent intruder.

The Willow Wren may justly be termed the nightingale of the northern snowy countries of Europe. It settles on the most lofty branches of the birch-trees, and makes the air resound with its bold and melodious song.



THE WATER WAGTAIL. (*Motacilla boarula.*)

THERE is not a brook purling along two flowery banks, not a rivulet winding through the green meadow, which is not frequented by this beautifully coloured and elegantly shaped little creature. We even see them in the streets of country towns, following with a quick pace, the half-drowned fly or moth, which the road-side streamlet carries away. Next to the robin redbreast and the sparrow, they are the boldest in approaching our habitations. The Wagtails are much in motion; seldom perch, and perpetually flirt their long and slender tails, (whence they derive their name,) principally after picking up some food from the ground, as if that tail were a kind of lever, or counterpoise, used to balance the body on the legs. They


are observed to frequent, more commonly, those streams where women come to wash their linen; probably not ignorant that the soap, the froth of which floats upon the water, attracts those insects which are most acceptable to them.

There are two species of Wagtails, the common kind, which has somewhat of a proud conceited air, and which is common in Britain only in summer; and what is called the Winter Wagtail, which is never found after February or March. The first, which is a very lively little bird, and seems always in a bustle, is black, softening into ash-colour and white; it is also bold, and will take the food thrown to it with as much confidence as a robin redbreast. The Winter Wagtail, on the contrary, is retiring in its habits, and much slower in its motions; its breast is yellow, and its wings greyish.

The Yellow Shepherdess (*Budytes flava*) was once included in the Wagtails. The male is olive-green on the back, and yellow on the lower part of the body, but the breast of the female is nearly white. These birds do not frequent the banks of rivers, but are generally found walking among the grass of meadows, and following sheep. They are rare in England; and are only seen here in spring; on which account they are sometimes called spring Wagtails.

White says, that "while the cows are feeding in the moist, low pastures, broods of Wagtails, white and grey, run round them, close up to their noses, and under their very bellies, availing themselves of the flies that settle on their legs, and probably finding worms and larvæ that are roused by the trampling of their feet. Nature is such an economist that the most incongruous animals can avail themselves of each other."

"Interest makes strange friendships!"



THE SWALLOW. (*Hirundo rustica*.)

“ From the low-roof'd cottage ridge,
 See the chattering Swallow spring ;
 Darting through the one-arch'd bridge,
 Quick she dips her dappled wing.”

CUNNINGHAM.

SWALLOWS are easily distinguished from all other birds, not only by their general structure, but by their twittering note and mode of flying, or rather darting from place to place.

They appear in Britain in April, and build in some out-house, or, in part of a human dwelling, where they lay their eggs and hatch their young. About August they disappear, and do not return till the following spring. Swallows kept in a cage moult about Christmas, and seldom live till spring.

There are several species of the Swallow : the general characters of which are a small beak, but large, wide mouth, for the purpose of swallowing flying insects, their natural food ; and long forked tail and extensive wings, to

enable them to pursue their prey. The common House Swallow builds under the eaves of houses, or in chimneys, near their top; the martin also builds under eaves, and very commonly against the upper corner or side of our very windows, and seems not afraid at the sight of man, yet it cannot be tamed, or even kept long in a cage. The nature of the Swallow's nest is worthy of our serious observation: how the mud is extracted from the sea-shores, rivers, or other watery places; how masoned and formed into a solid building, strong enough to support a whole family, and to face the "pelting storm," are wonders which ought to raise our mind to Him who bestowed that instinct upon them.

It is related that a pair of Swallows built their nest for two successive years on the handle of a pair of garden shears, that were stuck up against the boards in an out-house; and, therefore, must have had their nest spoiled whenever the implement was wanted. And what is still more strange, a bird of the same species built its nest on the wings and body of an owl that happened to hang dead and dry from the rafter of a barn, and so loose as to be moved by every gust of wind. This owl, with the nest on its wings, and with eggs in the nest was taken to the museum of Sir Ashton Lever as a curiosity. That gentleman, struck with the singularity of the sight, furnished the person who brought it with a large shell, desiring him to fix it just where the owl had hung. The man did so; and in the following year a pair of Swallows, probably the same, built their nest in the shell, and laid eggs.

Modern poets have not been unmindful of the Swallows; and our immortal Shakspeare mentions the martin, in *Macbeth*, in the following manner:

" This guest of summer,
The temple-haunting Martlet, does approve,
By his loved masonry, that Heaven's breath
Smells wooingly here. No jutting frieze,
Buttress, or coignes of 'vantage, but this bird
Hath made his pendant bed and procreant cradle:
Where they most breed and haunt, I have observed,
The air is delicate."

"The Swallow," writes Sir Humphrey Davy, "is one of my favourite birds, and a rival of the nightingale, for he cheers my sense of seeing as much as the other does my sense of hearing. He is the glad prophet of the year, the harbinger of the best season—he lives a life of enjoyment amongst the loveliest forms of nature—winter is unknown to him; and he leaves the green meadows of England in autumn for the myrrh and orange groves of Italy, and for the palms of Africa; he has always objects of pursuit, and his success is secure. Even the beings selected for his prey are poetical, beautiful, and transient. The ephemeræ are saved by his means from a slow and lingering death in the evening, and killed in a moment when they have known nothing but pleasure. He is the constant destroyer of insects, the friend of man, and may be regarded as a sacred bird. His instinct, which gives him his appointed season, and teaches him when and where to move, may be regarded as flowing from a divine source; and he belongs to the oracles of nature, which speak the awful and intelligible language of a present Deity."

The chimney Swallow is on the head, neck, back, and rump of a shining black colour, with purple gloss and sometimes with a blue shade; the throat and neck are of the same colour; the breast and belly are white, with a dash of red. The tail is forked, and consists of twelve feathers. The wings are of the same colour with the back. Swallows feed upon flies, worms, and insects; and generally hunt their prey on the wing.

"Away! away! thou summer bird;
For Autumn's moaning voice is heard,
In cadence wild, and deepening swell,
Of winter's stern approach to tell."



HOUSE MARTIN, OR WINDOW SWALLOW.

(Hirundo urbana.)

THE Martin is something less than the swallow, with a comparatively large head, and a wide mouth; the colour of the upper parts a bluish black, the rump and all the under parts of the body white, the bill black; its legs covered with short white down.

These birds begin to appear about the middle of April, and for some time pay no attention to the business of nidification, but sport and play about, either to recruit themselves from the fatigue of the journey, or else that their blood may recover its true tone and texture, after having been so long benumbed by the severities of the winter.

Should the weather prove favourable, it begins to build early in May, placing its nest generally beneath the eaves of a house, often against a perpendicular wall: without any projecting ledge to support any part of the nest, its utmost efforts are necessary to get the first foundation firmly fixed, so as to carry the superstructure safely. On this occasion, it not only clings with its claws, but partly

“The Swallow,” writes Sir Humphrey Davy, “is one of my favourite birds, and a rival of the nightingale, for he cheers my sense of seeing as much as the other does my sense of hearing. He is the glad prophet of the year, the harbinger of the best season—he lives a life of enjoyment amongst the loveliest forms of nature—winter is unknown to him; and he leaves the green meadows of England in autumn for the myrrh and orange groves of Italy, and for the palms of Africa; he has always objects of pursuit, and his success is secure. Even the beings selected for his prey are poetical, beautiful, and transient. The ephemerae are saved by his means from a slow and lingering death in the evening, and killed in a moment when they have known nothing but pleasure. He is the constant destroyer of insects, the friend of man, and may be regarded as a sacred bird. His instinct, which gives him his appointed season, and teaches him when and where to move, may be regarded as flowing from a divine source; and he belongs to the oracles of nature, which speak the awful and intelligible language of a present Deity.”

The chimney Swallow is on the head, neck, back, and rump of a shining black colour, with purple gloss and sometimes with a blue shade; the throat and neck are of the same colour; the breast and belly are white, with a dash of red. The tail is forked, and consists of twelve feathers. The wings are of the same colour with the back. Swallows feed upon flies, worms, and insects; and generally hunt their prey on the wing.

“Away! away! thou summer bird;
For Autumn’s moaning voice is heard,
In cadence wild, and deepening swell,
Of winter’s stern approach to tell.”



HOUSE MARTIN, OR WINDOW SWALLOW.

(*Hirundo urbica.*)

THE Martin is something less than the swallow, with a comparatively large head, and a wide mouth; the colour of the upper parts a bluish black, the rump and all the under parts of the body white, the bill black; its legs covered with short white down.

These birds begin to appear about the middle of April, and for some time pay no attention to the business of nidification, but sport and play about, either to recruit themselves from the fatigue of the journey, or else that their blood may recover its true tone and texture, after having been so long benumbed by the severities of the winter.

Should the weather prove favourable, it begins to build early in May, placing its nest generally beneath the eaves of a house, often against a perpendicular wall: without any projecting ledge to support any part of the nest, its utmost efforts are necessary to get the first foundation firmly fixed, so as to carry the superstructure safely. On this occasion, it not only clings with its claws, but partly

supports itself by strongly inclining its tail against the wall, making that a fulcrum; and thus fixed, it plasters the materials into the face of the brick or stone. But that this work may not, while soft, sink by its own weight, the provident architect has the prudence and forbearance not to proceed too fast; but by building only in the morning, and dedicating the rest of the day to food and amusement, he gives it sufficient time to dry and harden. By this method, in about ten days, the nest is formed, strong, compact, and warm, and perfectly fitted for all the purposes for which it is intended. But nothing is more common than for the house-sparrow, as soon as the shell is finished, to seize on it, eject the owner, and line it according to its own peculiar manner. The Martin will return for several seasons to the same nest, where it happens to be well sheltered and secured from the injuries of the weather. They breed the latest of all our swallows, never being without unfledged young ones even so late as Michaelmas.

The first hatch consists of five eggs, which are white, inclining to dusky at the thicker end; the second, of three or four; and of a third, of only two or three. While the young birds are confined to the nest the parents feed them, adhering by the claws to the outside; but as soon as they are able to fly they receive their nourishment on the wing, by a quick and almost imperceptible motion.

“ Welcome, welcome, feathered stranger,
Now the sun bids Nature smile;
Safe arrived and free from danger,
Welcome to our blooming isle.”

FRANKLIN.



THE SWIFT, OR BLACK MARTIN,
(*Cypselus apus*,)

ARRIVES in England later, and takes its departure earlier than any of its congeners. The Swift is the largest of the swallow species, and the most rapid in its flight. Its nest, which is generally built in the crevices of old towers and steeples, is constructed of dried grass, feathers, thread, and similar materials. These the bird collects whilst on the wing, picking them up with great dexterity. They seldom alight upon the ground, and if by accident they fall upon a level surface, they recover themselves with difficulty, owing to the shortness of their legs, and the length of their wings. During the heat of the day they remain within their holes, and at morning and evening sally out in quest of food. They may then be seen in flocks, whirling round some lofty edifice, or describing in mid-air, an endless series of circles upon circles. Swifts fly higher, and wheel with bolder wing than the swallows, with whom they never intermingle.



THE GOATSUCKER. (*Caprimulgus Europæus*.)

THIS curious bird, called also the Nightjar, and the Fern Owl, comes to this country from Africa about the middle of May and usually leaves by the end of August. These birds are generally found in low bushes, or amongst tufts of large ferns; and generally fly at night, hence their name of Fern Owl. The beak is furnished with bristles, and the middle toe of each foot has a claw toothed like a comb, with which the bird clears and smooths his bristles, when necessary. The female lays her eggs upon the ground, without any nest, and lays only two. The name of Goatsucker originated in an absurd idea that this bird sucked the goat's milk, from its habit of lying on the ground near cows or she goats, and catching the flies that torment them by fixing on their udders.

Mr. Waterton, who is certainly the closest observer of nature who ever wrote on Natural History, states, in one of his very interesting works, that he has frequently seen the Goatsuckers catching insects in this manner, and thus proving themselves the best of friends to the animals they are accused of annoying.



THE SKYLARK. (*Alda arvensis.*)

“Go, tuneful bird, that gladd’st the skies,
To Daphne’s window speed thy way ;
And there on quivering pinions rise,
And there thy vocal art display.” SHENSTONE.

THE Skylark is distinguished from most other birds, by the long spur on his back toe, the earthy colour of his feathers, and by singing as he mounts in the air. The common Skylark is not much bigger than the house-sparrow. These birds generally make their nest in meadows among the high grass, and the tint of their plumage resembles so much that of the ground, that the body of the bird is hardly distinguishable as it hops along.

“The daisied lea he loves, where tufts of grass
Luxuriant crown the ridge : there, with his mate,
He finds their lonely house, of withered herbs,
And coarsest spear-grass ; next the inner work,
With finer, and still finer fibres lays,
Rounding it curious with his speckled breast.”
GRAHAME.

Larks breed thrice a year, in May, July, and August, rearing their young in a short space of time. They are caught in great quantities in winter, and are considered choice and delicate food. It is a melancholy observation, that man should feed upon, and indulge his sense of taste with those very birds who have so often delighted his sense of hearing with their songs, when they

usher to the gladdened creation the return of their best friend, the sun. The instinctive warmth of attachment which the female Skylark bears towards her own species, even when not her nestling, is remarkable. "In the month of May," says Buffon, "a young hen bird was brought to me, which was not able to feed without assistance. I caused her to be reared; and she was hardly fledged, when I received from another place, a nest of three or four unfledged larks. She took a strong liking to these new comers, which were but little younger than herself; she tended them night and day, cherished them beneath her wings, and fed them with her bill. Nothing could interrupt her tender offices. If the young ones were torn from her she flew to them as soon as they were liberated, and would not think of effecting her own escape, which she might have done a hundred times. Her affection grew upon her; she neglected food and drink; she at length required the same support as her adopted offspring, and expired at last, consumed with maternal solicitude. None of the young ones long survived her. They died one after another; so essential were her cares, which were equally tender and judicious."

The Lark mounts almost perpendicularly, and by successive springs, into the air, where it hovers at a vast height. Its descent is in an oblique direction, unless threatened by some ravenous bird of prey, or attracted by its mate, when it drops to the ground like a stone. On its first leaving the earth, its notes are feeble and interrupted; but, as it rises, they gradually swell to their full tone. As the Lark's flight is always at sun-rise, there is something in the scenery that renders its song peculiarly delightful: the opening morning, the landscape just gilded by the rays of the returning sun, and the beauty of the surrounding objects, all contribute to heighten our relish for its pleasing melody.

"———Up springs the Lark,
 Shrill-voiced and loud, the messenger of morn;
 Ere yet the shadows fly, he, mounted, sings
 Amid the dawning clouds, and from their haunts
 Calls up the tuneful nations." THOMPSON.

“ Alas ! it's no thy neebor sweet,
 The bonnie Lark, companion meet !
 Bending thee 'mang the dewy weat !
 Wi' speckled breast,
 When upwards springing, blythe to greet
 The purpling east.”

BURNS.

“ Early, cheerful, mounting lark,
 Light's gentle usher, morning's clerk,
 In merry notes delighting.”

SIR JOHN DAVIS.



THE WOODLARK. (*Alauda arborea*.)

THIS species is smaller than the skylark, and its voice deeper ; it has also a circle of white feathers encompassing the head, from eye to eye, like a crown or wreath, and the utmost feather of the wing is much shorter than the second, whereas in the common lark they are nearly equal. This bird sometimes emulates the nightingale ; for which, when pouring forth his sweet melody in the grove, during a silent night, he is often mistaken. These birds sit and perch upon trees, unlike the common lark, which always keeps to the ground. They build their nest at the foot of a bush, near the bottom of a hedge, or in high dry grass. The number of eggs is about four ; they are of a pale bloom colour, beautifully mottled, and clouded with red and yellow. Like the skylark, they assemble in large flocks during frosty weather.

Their usual food consists of small beetles, caterpillars, and other insects, as well as the seeds of numerous kinds of wild plants.

“ Bright o’er the green hills rose the morning ray,
The Woodlark’s song resounded on the plain,
Fair nature felt the warm embrace of day,
And smiled through all her animated reign.” LANGBOURN.



THE TITMOUSE, OR TOM-TIT. (*Parus major*.)

THIS small bird presents us with the curious anomaly of a tongue split at the end into four distinct filaments. The head is curiously coloured; the beak is generally black, with strong hairs at its base. The head and throat are of the same colour, the body of a yellowish green; the back and coverts of the wings green, and the rump inclining to blue; the tail is black, but touched with blue at its exterior edges. This bird, and all the species related to it, live on insects, as well as on seeds. When kept in a cage, it is really amusing to see with what quickness the Titmouse darts at any fly or moth which comes imprudently within its reach. If this kind of food be deficient, as generally happens in winter, it feeds upon several kinds of seed, and particularly that of the sun-

flower, which it dexterously holds upright between its claws and strikes powerfully with its sharp little bill, till the black covering splits, and yields its white contents to the persevering bird. They build their nest with great ingenuity; and the female lays seven or eight eggs.



THE YELLOWHAMMER, OR YELLOW
BUNTING. (*Emberiza citrinella*.)

THIS bird is somewhat larger than the sparrow; and has a pretty note, not unlike that of the linnet. Its head is of a greenish yellow, spotted with brown; the throat and belly are yellow; the breast and sides, under the wings, mingled with red. These birds build their nests on the ground, near some bush, where the female lays five or six eggs. The Yellowhammer may be sometimes seen perched on the finger of some poor man or woman in the streets of London, in a state of complete tameness; but this is the transitory effect of intoxication, and soon after the bird is bought and brought home, it dies, overcome by the power of the laudanum that has been given it.

This bird feeds on seeds and various sorts of insects, and is common in every lane, on every hedge throughout the country, flitting before the traveller, and about the bushes. Happily for him, we have not yet acquired the taste of the natives of Italy, where the Yellowhammer falls a daily victim to the delicacy of the table, and where its flesh is esteemed very delicious eating. There he is often fattened, for the purpose of gratifying the palate of epicures.

The Ortolan, (*Emberiza hortulana*,) which is another species of the same genus, is common in the central and

southern provinces of Europe, where it is thought exquisitely flavoured as an article of food. When first taken it is frequently very lean, but if supplied with abundance of food, it is said to be so greedy, that it will eat till it dies of repletion.



THE WHEATEAR, AND WHIN CHAT.

(*Saxicola ænanthe.*)


THE Wheatear is one of our earliest visitants, and may be found in every part of Britain. In the North, it generally frequents heaps of stones, ruins, or the dry stone walls of burial-grounds, and though it is a very handsome bird, and in the early season sings sweetly, its haunts have obtained it a bad name. The common alarm-note resembles the sound made in breaking stones with a hammer, and as it utters that note from the top of the heap which haply covers the bones of one who perished by the storm, or his own hand, popular fancy has not unnaturally associated the Wheatear with the supersti-

tion that belongs to the place of graves. Beneath that heap of stones, or in some neighbouring fallow, its nest may be discovered, formed of moss and dried grass, lined with hair, feathers, or wool, and containing five or six eggs of a delicate bluish white. These birds congregate on the southern downs about the middle of July; they are caught in vast numbers, in horse-hair nooses, which are set between two pieces of turf turned against each other

The Whin Chat is a beautiful bird, compact in form, with a rich and elegant plumage. Its song, which is peculiarly soft and sweet, may be heard in spring on the bushy margins and gorse of extensive heaths. Its nest, constructed in thick tufts of grass, and under bushes, is most carefully concealed. It is usually approached by a labyrinth to which the rising of the bird affords no clue, and it may long be sought in vain, though perhaps not more than a yard distant all the time. The eggs are bluish green, without any spots, and are never more than six in number.

The following lines, addressed to the English Ortolan, or Wheatear, by Mrs. Charlotte Smith, allude to the foolish timidity of that bird.

“ To take you, shepherd boys prepare
 The hollow turf, the wiry snare,
 Of those weak terrors well aware,
 That bid you vainly dread
 The shadows floating o'er downs,
 Or murmuring gale, that round the stones
 Of some old beacon, as it moans,
 Scarce moves a thistle's head.
 And if a cloud obscure the sun,
 With faint and fluttering heart you run
 Into the pitfall you should shun,
 And only leave when dead.”




THE SPARROW. (*Pyrgita domestica.*)

THIS bird is, next to the robin redbreast, the boldest of the small feathered tribe which frequent our barns and houses: he is a courageous little creature, and fights undauntedly against birds ten times bigger than himself. Sparrows are accused of destroying a great quantity of corn, and in several counties the landlord or farmer puts a price on a Sparrow's head; but the farmer is the person most injured by the plan, as the good Sparrows, in ridding land of caterpillars, do more than compensate for the loss of the grain they destroy. Mr. Bradley, in his *Treatise on Husbandry and Gardening*, shows by a calculation, that a pair of Sparrows, during the time they have their young ones to feed, destroy on an average, every week, three thousand three hundred and sixty caterpillars.

This bird is easily tamed, and will hop about the house, and on the table, with great familiarity. It will feed on anything, and is particularly fond of meat cut into small pieces. The song of the Sparrow, if we can so call it

chirping, is far from agreeable: this arises, however, not from want of powers, but from its attending solely to the note of the parent bird. A Sparrow, when fledged, was taken from the nest and educated under a linnet: it also heard by accident, a goldfinch; and its song was in consequence a mixture of the two. The male is particularly distinguished by a jet-black spot under the bill upon a whitish ground. Sparrows are found nearly in every country of the world.




THE LINNET (*Fringilla linota*)

Is about the size of the goldfinch; and compensates, by an extremely melodious voice, the want of variety in its plumage, which, except in the red-breasted species, is nearly all of one colour. Its musical talents are, like those of many other birds, repaid with captivity; for it is kept in cages on account of its singing.

The Red-breasted Linnet, or Redpole, generally builds on the sea-coast in this country, and on the Continent in vineyards; but that livery of nature, the crimson scarf, that grows so beautifully under his neck, disappears as soon as he is domesticated. This bird is one of the first whose appearance announces the spring.

The Green Linnet is rather larger than the house sparrow. Its head and back are of a yellowish green, the edges of the feathers grayish; the rump and breast more yellow. The plumage of the female is much less vivid, inclining to brown. Its song is trifling, but in confinement it becomes tame and docile; and will catch the notes of other birds.

southern provinces of Europe, where it is thought exquisitely flavoured as an article of food. When first taken it is frequently very lean, but if supplied with abundance of food, it is said to be so greedy, that it will eat till it dies of repletion.



THE WHEATEAR, AND WHIN CHAT.

(*Saxicola ananthe.*)

THE Wheatear is one of our earliest visitants, and may be found in every part of Britain. In the North, it generally frequents heaps of stones, ruins, or the dry stone walls of burial-grounds, and though it is a very handsome bird, and in the early season sings sweetly, its haunts have obtained it a bad name. The common alarm-note resembles the sound made in breaking stones with a hammer, and as it utters that note from the top of the heap which haply covers the bones of one who perished by the storm, or his own hand, popular fancy has not unnaturally associated the Wheatear with the supersti-

tion that belongs to the place of graves. Beneath that heap of stones, or in some neighbouring fallow, its nest may be discovered, formed of moss and dried grass, lined with hair, feathers, or wool, and containing five or six eggs of a delicate bluish white. These birds congregate on the southern downs about the middle of July; they are caught in vast numbers, in horse-hair nooses, which are set between two pieces of turf turned against each other.

The Whin Chat is a beautiful bird, compact in form, with a rich and elegant plumage. Its song, which is peculiarly soft and sweet, may be heard in spring on the bushy margins and gorse of extensive heaths. Its nest, constructed in thick tufts of grass, and under bushes, is most carefully concealed. It is usually approached by a labyrinth to which the rising of the bird affords no clue, and it may long be sought in vain, though perhaps not more than a yard distant all the time. The eggs are bluish green, without any spots, and are never more than six in number.

The following lines, addressed to the English Ortolan, or Wheatear, by Mrs. Charlotte Smith, allude to the foolish timidity of that bird.

“ To take you, shepherd boys prepare
The hollow turf, the wiry snare,
Of those weak terrors well aware,
That bid you vainly dread
The shadows floating o'er downs,
Or murmuring gale, that round the stones
Of some old beacon, as it moans,
Scarce moves a thistle's head.
And if a cloud obscure the sun,
With faint and fluttering heart you run
Into the pitfall you should shun,
And only leave when dead.”



THE CANARY-BIRD. (*Carduelis canaria.*)


As his name imports, this bird is a native of the Canary Islands; where, in his wild state, he has a dusky gray plumage, and a much stronger voice than when in a cage. In our northern countries his feathers undergo a great alteration; and the bird becomes entirely white or yellow. This effect of cold, upon animals of all kinds, is general and progressive, according to the distance of the climate from the equator. Of this bird, Buffon says, "that if the nightingale is the chantress of the woods, the Canary is the musician of the chamber; the first owes all to nature, the second something to art. With less strength of organ, less compass of voice, and less variety of note, the Canary has a better ear, greater facility of imitation, and a more retentive memory; and as the difference of genius, especially among the lower animals, depends in a great measure on the perfection of their senses, the Canary, whose organ of hearing is more susceptible of receiving and retaining foreign impressions, becomes more social, tame, and familiar; is capable of gratitude and even attachment; its caresses are endearing, its little humours innocent, and its anger neither hurts nor offends. Its education is easy; we rear it with pleasure, because we are able to instruct it. It leaves the melody of its own natural note, to listen to the melody of our voices and instruments. It accompanies us, and repays the pleasure it receives with interest, while the nightingale, more proud of its talent, seems desirous of preserving it in all its purity, at least it appears to attach

very little value to ours, and it is with great difficulty that it can be taught any of our airs. The Canary can whistle; the nightingale despises our airs, and never fails to return to its own wild wood notes. Its pipe is a masterpiece of nature, which human art can neither alter nor improve; while that of the Canary is a model of more pliant materials, which we can mould at pleasure; and therefore it contributes in a much greater degree to the pleasures of society. It sings at all seasons, cheers us in the dullest weather, and adds to our happiness, by amusing the young and delighting the recluse, charming the tediousness of the cloister, and gladdening the soul of the innocent and captivated." It breeds generally twice a year when domesticated; and it sometimes happens that the female lays her eggs for the second time before the first brood is fledged. The male then good-naturedly takes her place on the eggs while she feeds the young ones, and feeds them in his turn, when she sits in the nest. They are very easily tamed, when brought up with attention and kindness, and take their food out of the hand, often perching on the shoulder of their mistress, and feeding out of her mouth. The Canary-bird is sometimes, and with success, matched with the linnæus or the goldfinch; and the produce is a beautiful mule bird, partaking of the talents and plumage of both.

Canary-birds live twelve or thirteen years in our climate, and sing well to the end of their life.

The following curious anecdote of one of these birds is related by Dr. Darwin: "On observing a Canary-bird at the house of a gentleman near Tutbury, in Derbyshire, I was told it always fainted away when its cage was cleaned; and I desired to see the experiment. The cage being taken from the ceiling, and the bottom drawn out, the bird began to tremble, and turned quite white about the root of the bill: he then opened his mouth, as if for breath, and respired quick; stood up straighter on his perch, hung his wings, spread his tail, closed his eyes, and appeared quite stiff for half-an-hour; till at length, with much trembling and deep respirations, he came gradually to himself."

A few years ago, a Frenchman exhibited in London twenty-four Canary-birds, many of which, he said, were from eighteen to twenty-five years of age. Some of these balanced themselves, head downward, on their shoulders, having their legs and tail in the air. One of them taking a slender stick in its claws, passed its head between its legs, and suffered itself to be turned round, as if in the act of being roasted. Another balanced itself, and was swung backward and forward on a kind of a slack rope. A third was dressed in military uniform, having a cap on its head, wearing a sword and pouch, and carrying a firelock in one claw: after some time sitting upright, this bird, at the word of command, freed itself from its dress, and flew away to the cage. A fourth suffered itself to be shot at, and falling down as if dead, was put into a little wheelbarrow, and wheeled away by one of its comrades!



THE CHAFFINCH. (*Fringilla caelebs.*)

THE Chaffinch is of the same dimensions as the sparrow, but more lightly and elegantly formed. Its nest, which is of the most beautiful and elaborate construction, is com-

posed of mosses and lichens, interwoven and lined with wool, hair, and feathers. "Four or five eggs," says Mr. Waterton, "are the usual number which the Chaffinch's nest contains, and sometimes only three. The thorn, and most of the evergreen shrubs, the sprouts on the boles of forest trees, the woodbine, the whin, the wild-rose, and occasionally the bramble, are this bird's favourite places for nidification. Like all its congeners, it never covers its eggs on retiring from the nest, for its young are hatched blind. There is something peculiarly pleasing to me in the song of this bird. Perhaps association of ideas may add a trifle to the value of its melody; for when I hear the first note of the Chaffinch, I know that winter is on the eve of its departure, and that sunshine and fine weather are not far off. The Chaffinch never sings when on the wing; but it warbles incessantly on the trees, and on the hedgerows, from the early part of February to the second week in July; and then (if the bird be in a state of freedom,) its song entirely ceases."





THE BULLFINCH. (*Loxia pyrrhula.*)

THIS is a very docile bird, and will nearly imitate the sound of a pipe, or the whistle of man, with its voice, the mellowness of which is really charming. It is, by bird-fanciers, considered to excel all other small birds, except the linnet, in the softness of its tones, and in the variety of its notes. In captivity, its melody seems to be as great a solace to itself, as it is a pleasure to its master. By day, and even when the evening has called for the artificial light of candles, the Bullfinch pursues his melodious exertions, and if there be any other birds in the apartment, awakes them gently to the pleasing task of singing in concert with him. His notes are upon one of the lowest keys of the gamut of birds.

The plumage of the Bullfinch is beautiful, though simple and uniform, consisting only of three or four shades of colours. In the male, a lovely scarlet or crimson colour adorns the breast, throat, and jaws, as far as the eyes; the crown of the head is black; the rump and tail are white; the neck and back grey, or lead-coloured. The name of this bird originates from its head and neck being, like those of the bull, very large in proportion to the body. The female does not share with the male the brightness of colours in the plumage. Bullfinches build their nests in gardens and orchards, and particularly in places that abound in fruit-trees, as they are passionately fond of fruit, which they often destroy before it is ripe.



THE GOLDFINCH. (*Fringilla carduelis.*)

THIS bird is also called the Thistlefinch, from his fondness for the seed of that plant. He is very beautiful, his plumage being elegantly diversified, his form small, but pleasing, and his voice not loud, but sweet. He is easily tamed, and often exhibited as a captive, with a chain round his body, drawing up with trouble, but yet with amazing dexterity, two small buckets, alternately, one containing his meat, the other his drink. When caught old, the Goldfinch, after a few weeks, if well attended to, and gently treated, becomes as familiar as if he had been brought up by the hand of his keeper. Some have been taught to fire a small piece of artillery, and go through the drilling exercise, to the great astonishment of the spectators; but the cruel and severe treatment that animals undergo, when taught performances altogether contrary to their nature, should prevent us from encouraging such exhibitions.

This bird, as if conscious of the beauty of his plumage, likes to view himself in a glass, which is sometimes fixed for this purpose in the back of the cage. The art with which it composes and builds its nest is really worthy of admiration; it is generally interwoven with moss, small twigs, horsehair, and other pliant materials; the inside stuffed most carefully with fine down, and tufts of cotton-grass, or cannach. There the female deposits five or six eggs, which are whitish, marked at their upper end with purple dots.


“ The Goldfinch weaves, with willow down inlaid,
 And cannach tufts, his wonderful abode ;
 And oft suspended at the limber end
 Of plane-tree spray, among the broad-leaved shoots,
 The tiny hammock swings to every gale.
 Sometimes in closest thickets 'tis concealed ;
 Sometimes in hedge luxuriant, where the brier,
 The bramble, and the plum-tree branch
 Warp through the thorn, surmounted by the flowers
 Of climbing vetch, and honeysuckle wild.” GRAHAME.

The following lines were written by Cowper on a Goldfinch starved to death in its cage. The Goldfinch speaks :—

“ Time was when I was free as air,
 The thistle's downy seed my fare,
 My drink the morning dew ;
 I perched at will on every spray,
 My form genteel, my plumage gay,
 My strains for ever new.

“ But gandy plumage, sprightly strain,
 And form genteel were all in vain,
 And of a transient date ;
 For caught and caged, and starved to death,
 In dying sighs my little breath
 Soon passed the wiry grate.

“ Thanks, gentle author of my woes,
 Thanks for this most effectual close
 And cure of every ill.
 Never your cruelty repress !
 For I, if you had shown me less,
 Had been your prisoner still.”



THE STARE, OR STARLING (*Sturnus vulgaris*)

Is about the size and shape of a blackbird; the tips of the feathers on the neck and back are yellow; the feathers under the tail of an ash-colour; the other parts of the plumage are black, with a purple or deep blue gloss, changing as it is variously exposed to the light. In the hen, the tips of the feathers on the breast and belly, to the very throat, are white; which constitutes a material point in the choice of the bird, as the female is no singer. She lays four or five eggs, lightly tintured with a greenish cast of blue. Starlings build in hollow trees and clefts of rocks, are very easily tamed, and can add to their natural notes any words or modulations which they are taught.

In the winter season Starlings collect in vast flocks, and may be known at a great distance by their whirling mode of flight. The evening is the time when they assemble in the greatest numbers, and betake themselves to fens and marshes. Sterne has immortalized the Starling in his *Sentimental Journey*:—"The bird flew to the place where I was attempting his deliverance, and thrusting his head through the trellis, pressed his head against it, as if impatient.—'I fear, poor creature,' said I, 'I can't set thee at liberty.'—'No,' said the Starling, 'I can't get out.' 'Disguise thyself as thou wilt, still, slavery,' said I, 'still thou art a bitter draught!'"



THE RAVEN. (*Corvus Corax.*)

“The Raven sits
 On the raven-stone,
 And his black wing flits
 O'er the milk-white bone ;
 To and fro, as the night-winds blow,
 The carcass of the assassin swings :
 And there alone, on the raven-stone,
 The Raven flaps his dusky wings.
 The fetters creak—and his ebon beak
 Creaks to the close of the hollow sound :
 And this is the tune, by the light of the moon,
 To which the witches dance their round.”

BYRON'S MANFRED.

THE Raven is about twenty-six inches in length, and his weight about three pounds. The bill is strong, black, and hooked at the tip. The plumage of the whole body

of a shining black, glossed with deep blue; the back of the lower part inclining to a dusky colour. He is of a strong and hardy disposition, and inhabits all climates of the globe. He builds his nest in trees; and the female lays five or six eggs of a palish green colour, spotted with brown. It is said that the life of this bird extends to a century; and even beyond that period, if we can believe the accounts of several naturalists on the subject. The Raven unites the voracious appetite of the crow to the dishonesty of the daw and the docility of almost every other bird. He feeds chiefly on small animals; and is said to destroy rabbits, young ducks, and chickens, and sometimes even lambs, when they happen to be dropped in a weak state. In the northern regions, he preys on carrion, in concert with the white bear, the arctic fox, and the eagle. The faculty of scent in these birds must be very acute; for in the coldest of the winter days, at Hudson's Bay, when every kind of effluvium is almost instantaneously destroyed by the frost, buffaloes, and other beasts have been killed, where not one of these birds was seen; but in a few hours scores of them have been found collected about the spot, to pick up the blood and offal. The Raven possesses many diverting and mischievous qualities; he is active, curious, sagacious, and impudent; by nature a glutton, by habit a thief, in disposition a miser, and in practice a rogue. He is fond of picking up any small piece of money, bits of glass, or anything that shines, which he carefully conceals under the eaves of roofs, or in any other inaccessible place. He is easily tamed; and, like the parrot and the starling, can imitate the human voice, in articulating words. At the seat of the Marquess of Aylesbury, in Wiltshire, a tame raven, that had been taught to speak, used to ramble about in the park, where he was commonly attended and beset with crows, rooks, and others of his inquisitive tribe. When a considerable number of these were collected round him, he would lift up his head, and with a hoarse and hollow voice shout out *Holloa!* This would instantly put to flight, and disperse his sable brethren; while the Raven seemed to enjoy the fright he had occasioned.

When domesticated, the Raven is of great service, both as a scavenger and in keeping watch, in the last of which he is more alert and vigilant than almost any other animal. The Raven was the ensign of the invading Danes, and the prejudice thereby engendered against the bird is not yet quite extinct. Of its perseverance in the act of incubation, Mr. White relates the following singular anecdote :

“ In the centre of a grove near Selborne, there stood an oak, which, though on the whole shapely and tall, bulged out into a large excrescence near the middle of the stem. On this tree a pair of Ravens had fixed their residence for such a series of years, that the oak was distinguished by the title of ‘The Raven-tree.’ Many were the attempts of the neighbouring youths to get at this nest: the difficulty whetted their inclinations, and each was ambitious of surmounting the arduous task; but when they arrived at the swelling, it jutted out so in their way, and was so far beyond their grasp, that the boldest lads were deterred, and acknowledged the undertaking to be too hazardous. Thus the Ravens continued to build, nest upon nest, in perfect security, till the fatal day on which the wood was to be levelled. This was in the month of February, when those birds usually sit. The saw was applied to the trunk, the wedges were inserted into the opening, the wood echoed to the heavy blows of the mallet, the tree nodded to its fall; but still the dam persisted in sitting. At last, when it gave way, the bird was flung from her nest; and though her parental affection deserved a better fate, was whipped down by the twigs, which brought her dead to the ground !”

The croaking of the Raven was formerly considered a note of ill omen,—

“ The Raven croaked as she sat at her meal,
And the old woman knew what he said,
And she grew pale at the Raven’s tale,
And sickened and went to her bed.”



THE CARRION CROW. (*Corvus corone.*)

THIS bird is less in size than the raven. The bill is strong, thick, and straight. The general colour is black, except the extremities of the feathers, which are of a greyish tint. His delight is to feed upon carcasses, and dead animals, or malefactors exposed on the gibbet. Like the raven, he has a very acute scent, and is said to smell gunpowder afar off. He roosts upon trees, and takes both animal and vegetable food. Crows, like rooks, are gregarious, and often fly in large companies in the fields or in the woods. On the upland moors, Crows occupy the place which rooks fill in the low country; and as the Crow has a very coarse and uncouth voice, the Lowlanders of Scotland are in the habit of saying that the Highland rooks "speak Gaelic." They are great destroyers of partridges, as they generally pierce their eggs with their bills, and carry them in that manner through the air, to a great distance, to feed their young. The female lays five or six eggs.

Mr. Montagu states that he once saw a Crow in pursuit of a pigeon, at which it made several pounces, like a hawk; but the pigeon escaped by flying in at the door of a house. He saw another strike a pigeon dead from the top of a barn. The Crow is so bold a bird that neither the kite, the buzzard, nor the raven can approach its nest without being driven away. When it has young ones, it will even attack the peregrine falcon, and at a single pounce sometimes bring that bird to the ground.



THE ROOK. (*Corvus frugilegus.*)

THE cawing of these birds, on the tops of high trees near gentlemen's houses, and in the middle of cities, is not very pleasing; yet old habits, to which we are reconciled, have as much influence upon us, as if they were productive of amusement. Hence it has been seldom attempted to destroy a rookery; although the noise and other inconveniences that accompany these birds render their vicinity often troublesome. They feed entirely on corn and insects, and are little bigger than the common crows. In Suffolk and in some parts of Norfolk, the farmers find it their interest to encourage the breed of Rooks, as the only means of freeing their grounds from the grub which produces the cockchafer, and which in this state destroys the roots of corn and grass to such a degree, that instances have been known where the turf of pasture land might be turned up with the foot. The farmers in a northern county, a good many years ago, waged a war of extermination against the Rooks, but the very next year the crops were so completely cut up by grubs, that the same proprietors were at considerable expense in getting Rooks back again. Young Rooks are good eating, but should be skinned before they are dressed. The colour is black, but brighter than that of the crow, which the Rook resembles in shape. The female lays the same number of eggs; and the male shares with her the trouble of fetching sticks, and interweaving them to make the nest, an operation which is attended with a great deal of fighting and disputing with the other Rooks.

New comers are often severely beaten by the old inhabitants, and are even frequently driven quite away; of this an instance occurred near Newcastle, in the year 1783. A pair of Rooks, after an unsuccessful attempt to establish themselves in a rookery at no great distance from the Exchange, were compelled to abandon the attempt, and take refuge on the spire of that building; and, though constantly interrupted by other Rooks, they built their nest on the *top of the vane*, and reared their young ones, undisturbed by the noise of the populace below. The nest and its inhabitants were of course turned about by every change of the wind! They returned and built their nest every year on the same place, till 1793, soon after which year the spire was taken down. A small copperplate was engraved, of the size of a watch-paper, with a representation of the spire and the nest; and so much pleased were the inhabitants and other persons with it, that as many copies were sold as produced to the engraver a profit of ten pounds. The wood-cut by Bewick, in the title-page to his *Select Fables*, gives a view of the old Exchange, with the Rook's nest on the vane.

It is amusing to see Rooks coming at sunset as thick as a cloud, hovering over a grove, and, after several eddies described in the air, and incessant cawings, each repairing to its own nest, and settling in a few minutes to rest, till the dawn calls them up again to their pasture in the neighbouring fields.


Dr. Darwin has remarked, that an instinctive feeling of danger from mankind is much more apparent in Rooks than in most other birds. Any one who has in the least observed them will see that they evidently distinguish that the danger is greater when a man is armed with a gun, than when he has no weapon with him. In the spring of the year, if a person happened to walk under a rookery with a gun in his hand, the inhabitants of the trees rise on their wings, and scream to the unfledged young to shrink into their nests from the sight of the enemy. The country people observing this circumstance so uniformly to occur, assert that Rooks can smell gunpowder.



THE JACKDAW. (*Corvus monedula.*)

THIS bird is much less than the crow. He has a large head and long bill, in proportion to the size of his body. The colour of the plumage is black, but on some parts inclining to a bluish hue; the fore part, of the head is of a deeper black. The Jackdaw feeds upon nuts, fruits, seeds, and insects; and builds in ancient castles, towers, cliffs, and all desolate and ruinous places. The female lays five or six eggs, smaller, paler, and marked with fewer spots than those of the crow.

Jackdaws are easily tamed, and may with little difficulty be taught to pronounce several words. They conceal such parts of their food as they cannot eat, and often, along with it, small pieces of money or toys, frequently occasioning, for the moment, suspicions of theft in persons who are innocent. In Switzerland there is found a variety of the Jackdaw, which has a white ring round its neck. In Norway, and other cold countries, they have been seen entirely white. In a state of nature, Jackdaws and rooks frequently feed together, and the Jackdaws come to meet the rooks in the morning, and also accompany them for some distance on their retreat at night.



THE MAGPIE. (*Caudata pica.*)

“From bough to bough the restless Magpie roves,
And chatters as he flies.”
GISBORNE.

THIS bird resembles the daw, except in the whiteness of the breast and wings, and the length of the tail. The black of the feathers is accompanied with a changing gloss of green and purple. It is a very loquacious creature, and can be taught to imitate the human voice as well as any of the feathered creation.

Plutarch relates a singular story of a Magpie belonging to a barber at Rome, which could imitate, to a wonderful extent, almost every noise that it heard. Some trumpets happened one day to be sounded before the shop; and for a day or two afterwards the Magpie was quite mute, and seemed pensive and melancholy. This surprised all who knew it; and they supposed the sound of the trumpets had so stunned the bird as to deprive it at the same time of voice and hearing. This, however, was not the case; for, says the writer, the bird had been all the time occupied in profound meditation, and was studying how to imitate the sound of the trumpets; accordingly, in the first attempt, it perfectly imitated all their repetitions, stops, and changes. This new lesson, however, made it entirely forget everything that it had learned before.

The Magpie feeds on everything; worms, insects, meat, cheese, bread, milk, and all kinds of seeds; and also on small birds, when they come in its way: the young of the blackbird and of the thrush, and even a strayed

chicken, often fall a prey to its rapacity. It is fond of hiding pieces of money or wearing apparel, which it carries away by stealth, and with much dexterity, to its hole. Its cunning is also remarked in the manner of making its nest, which it covers all over with hawthorn branches, the thorns sticking outward; within it is lined with fibrous roots, wool, and long grass, and then plastered all round with mud and clay. The canopy above is composed of the sharpest thorns, woven together in such a manner as to deny all entrance except at the door, which is just large enough to permit egress and regress to the owners. In this fortress the birds bring up their brood with security, safe from all attacks, but those of the climbing schoolboy, who often finds his torn and bloody hands too dear a price for the eggs or the young ones.


There are many superstitions respecting Magpies; and it is singular that in all the southern and middle districts of England, two Magpies together are thought to betoken luck; while in Lancashire, and other northern counties, they are thought to betoken misfortune. The chattering of Magpies was formerly supposed to foretell the arrival of strangers.





THE CROSSBILL. (*Loxia*.)

THE Crossbill is a native of the vast pine forests of the high latitudes. The bill of this singular bird is of considerable length, and the mandibles towards the point are very sharp and strong, curved in opposite directions, so that when closed the points cross each other, from which the bird derives his name. This curious organization enables them to obtain their food, which chiefly consists of the seeds of the cones of the fir, with the greatest facility. These seeds, for a considerable time after they have ripened, are so firmly enclosed within their ligneous scales that the bill of no ordinary bird could reach them. Fixing itself across the cone, the Crossbill brings the mandibles of its beak immediately over each other, and insinuates them between the scales, then forcing them laterally, the scales open. The mandibles are again brought in contact, keeping the scales open, whilst the bird scoops out the seed with its tongue.



THE JAY (*Garrulus glandarius*)

Is less than the magpie, and resembles him more in the habits of his life than in the shape and colour of his body. Like him, he is talkative, and ready to imitate all sounds, but boasts of ornamental colours, which the magpie is deprived of. The ablest painter can produce no colour to equal the brightness of the chequered tablets of white, black, and blue, which adorn the sides of his wings. His head is covered with feathers, which are moveable at will, and the motion of which is expressive of the internal affections of the bird, whether he is stimulated by fear, anger, or desire.

A Jay, kept by a person in the north of England, had learned at the approach of cattle to set a cur dog upon them, by whistling and calling him by his name. One winter, during a severe frost, the dog was by this means excited to attack a cow that was big with calf, when the poor animal fell on the ice, and was much hurt. The Jay was complained of as a nuisance, and its owner was obliged to destroy it.


The hen lays five or six eggs, of a dull white colour, mottled with brown.



THE CORNISH CHOUGH (*Pyrrhcorax graculus*)

Is like the jackdaw in shape and colour, but somewhat larger. The bill and legs are of a red colour, and hence the bird is frequently called the red-legged Crow. It is an inhabitant of Cornwall, Wales, and all the western coasts of England, and is generally to be found among rocks near the sea, where it builds, as well as in old ruinous castles and churches on the sea-side. The voice of the Chough resembles that of the jackdaw, except that it exceeds it in hoarseness and strength.

Mr. Montagu describing a Chough in the possession of a friend, says, "his curiosity is beyond bounds, never failing to examine everything new to him: if the gardener be pruning, he examines the nail-box, carries off the nails, and scatters the shreds about. Should a ladder be left against the wall, he instantly mounts, and goes all round the top of the wall; and if hungry descends at a convenient place, and immediately travels to the kitchen window, where he makes an incessant knocking with his bill, until he is fed or let in. If allowed to enter, his first endeavour is to get up-stairs; and if not interrupted, goes as high as he can, and gets into any room on the attic story: but his intention is to get upon the top of the house. He is excessively fond of being caressed, and would stand quietly by the hour to be smoothed; but resents an affront with violence and effect, by both bill and claws, and will hold so fast by the latter, that he is with difficulty disengaged."



THE SATIN BOWER-BIRD.

(*Ptilonorhynchus holosericeus*.)

THIS singular bird was first brought before the notice of the public by Mr. Gould, in his splendid work, the *Birds of Australia*, from which the following extracts are given by permission of its author. The most remarkable circumstance relating to this bird, is its construction of a bower-like tenement, the object of which, it should seem, is a sort of playing-ground, or hall of assembly.

“The Satin Bower-bird,” says Mr. Gould, “is not a stationary species, but appears to range from one part of a district to another, either for the purpose of varying the nature, or of obtaining a more abundant supply of food. Judging from the many specimens I dissected, it would seem that it is, altogether granivorous and frugivorous; or, if not exclusively so, that insects form but a small portion of its diet. The brushes it inhabits are studded with enormous fig-trees, some of them towering to the height of two hundred feet; among the lofty branches of which the Satin Bower-bird finds, in the small wild fig with

which the branches are loaded, an abundant supply of a favourite food: this species also commits considerable depredation on ripening corn. It appears to have particular times in the day for feeding, and when thus engaged among the low shrub-like trees, I have approached within a few feet without creating alarm; but at other times I have found this bird extremely shy, especially the old males, which not unfrequently perch on the topmost branch of the loftiest tree, whence they can survey all around, and watch the movements of the females and young in the brush below. Besides the loud liquid call peculiar to the male, both sexes frequently utter a harsh, unpleasant, guttural note, indicative of surprise or displeasure. The old black males are exceedingly few in number, as compared with the females and young male birds in the green dress, from which and other circumstances I am led to believe that at least two, if not three years, elapse before they attain the rich satin-like plumage, which, when once perfectly assumed, is, I believe, never again thrown off. The extraordinary bower-like structures alluded to above, are usually placed under the shelter of the branches of some overhanging tree in the most retired part of the forest, and differ considerably in size. The base consists of an extensive and rather convex platform of sticks, firmly interwoven, on the centre of which the bower itself is built: this, like the platform on which it is placed, and with which it is interwoven, is formed of sticks and twigs, but of a more slender and flexible description, the tips of the twigs being so arranged as to curve inwards and nearly meet at the top: in the interior of the bower the materials are so placed, that the forks of the twigs are always presented outwards, by which arrangement not the slightest obstruction is offered to the passage of the birds. The interest of this curious bower is much enhanced by the manner in which it is decorated at and near the entrance with the most gaily-coloured articles that can be collected, such as the blue tail-feathers of the Rose-bill and Pennantian parrots, bleached bones, the shells of snails, &c.; some of the feathers are stuck in among the twigs, while others with the bones and shells are strewed about near the en-

trances. The propensity of these birds to pick up and fly off with any attractive object, is so well known to the natives, that they always search the runs for any small missing article, as the bowl of a pipe, &c., that may have been accidentally dropped in the brush. I myself found at the entrance of one of them a small, neatly-worked stone tomahawk, of an inch and a half in length, together with some slips of blue cotton rags, which the birds had doubtless picked up at a deserted encampment of the natives. For what purpose these curious bowers are made is not yet, perhaps, fully understood; they are certainly not used as a nest, but as a place of resort for many individuals of both sexes, which, when there assembled, run through and around the bower in a sportive and playful manner, and that so frequently, that it is seldom entirely deserted."



THE BIRD OF PARADISE. (*Paradisea apoda.*)


THERE are about seven distinct species of Birds of Paradise. They are generally seen on the wing; and flying in

the torrid zone at a small distance from the land, their appearance is therefore most welcome to the tired sailor and anxious passenger, by its foretelling the vicinity of *terra firma*. The head is small, but adorned with colours which vie with the brightest hues of the feathered tribe; the neck is a beautiful fawn, and the body very small, but covered with long feathers of a browner hue, tinged with gold: the two middle feathers of the tail are little more than filaments, except at the point and near the base. Although the body is no larger than that of a thrush, the total length is two feet. This bird has long been esteemed by ladies as a head-dress, and as those sent to Europe for this purpose always had the legs cut off for the convenience of packing, it was reported, and at one time believed, that the Bird of Paradise had no legs, but that it lived always on the wing. Indeed, a very fierce controversy arose on this subject among the earlier naturalists.

The native place of these birds is New Guinea and the neighbouring islands, where they are generally found in flocks of thirty or forty, roosting on fig or teak-trees. They always fly against the wind, that it may not ruffle their light and spreading plumage, as, if the wind came from behind it would blow their long tails over their backs. They take shelter from storms in the most dense thickets, and feed principally on figs, the berries of the teak, and insects. The note of the Bird of Paradise is very unpleasant, and resembles the cawing of a raven; it is chiefly heard in windy weather, when they dread being thrown on the ground.


In their flights they are led, it is alleged, by a single bird, which the natives call their king, but which is said to be of a different species. It is further pretended that when this bird settles, the whole flight settle also, in consequence of which they sometimes perish, as they are unable to rise again, owing to the peculiar structure of their wings. M. Lesson saw two of these birds at Amboyna, in a cage, and Mr. Beale, of Macao, kept one in his aviary for more than nine years: they were fed with living insects and would not touch them when dead.

NUTHATCH.



THE NUTHATCH, OR NUTJOBBER,
(*Sitta Europaea*.)

Is less than the chaffinch. The head, neck, and beak are of an ash-colour; the sides under the wings red; the throat and breast of a pale yellow; the chin white, and the feathers under the tail red, with white tips. The Nuthatch feeds upon nuts, which he hoards in the hollow part of a tree: and it is pleasing to see him fetch a nut out of the hole, place it first in a chink, and standing above it with his head downwards, striking it with all his might, break the shell, and catch up the kernel. He feeds also upon flies and other small insects. The hen is so attached to her brood, that, when disturbed from her nest, she flutters about the head of the depredator, and hisses like a snake. The Nuthatches are shy and solitary birds, and, like the woodpecker, frequent woods, and run up and down the trees with surprising facility. They often move their tails in the manner of the wagtail. They do not migrate, but during the winter approach nearer to inhabited places, and are sometimes seen in orchards and gardens. The female lays her eggs in holes of trees, frequently in those which have been deserted by the woodpecker. The young ones are said to afford very delicate food.



THE WRYNECK. (*Yunx torquilla.*)

THIS bird, Mr. Gould tells us, has received its English name from its habit of moving its head and neck in various directions, and with an undulating motion, like that of a snake; indeed, in some parts of England it is called the snake-bird. When found in its usual retreat in the hole of a tree, it makes a loud hissing noise, raises the feathers of the crown, and writhing its head and neck towards each shoulder alternately, with grotesque contortions, becomes an object of terror to a timid intruder; and the bird, taking advantage of a moment of indecision, darts with the rapidity of lightning from a situation where escape appeared impossible.

The Wryneck deposits its eggs on fragments of decayed wood within a hollow tree, and makes scarcely any nest. The birds when caught young are easily tamed.

THE CREEPER. (*Certhia familiaris.*)

THE Creepers form a family in themselves. They are dispersed through most countries of the globe, and feed chiefly on insects, in search of which they run in a spiral direction round the stems and branches of trees, with great agility. The *Wall Creeper*, or *Spider Catcher*, is larger than a house-sparrow. It has a long, slender, black bill; the head, neck, and back are of an ash-co-

lour; the breast white; the wings a compound of lead-colour and red. It is a brisk and cheerful bird, and has a pleasant note. It builds its nest in the holes of trees.

The *Ox-eye Creeper*, is not much bigger than a wren, and has a long, slender, sharp bill. The throat and breast are white; the head, back, and wings of a fox colour; the middle parts of the wings whitish; above the eyes, on each side, there is a white spot. It is commonly seen in England, and builds in hollow trees. The smaller the bird, in general, the greater number of eggs the female lays: the number of the Creeper's eggs is sometimes above twenty. In America the Creeper hatches twice during the summer, and has generally from eighteen to twenty eggs at a time.



THE HUMMING-BIRDS. (*Trochilus colubris.*)

THERE are numerous species of Humming-Birds, but that represented in the plate is one of the most common. They are abundant in South America, particularly in Brazil; and are so small and so brilliant in their colours, that when seen fluttering about in the brilliant rays of a tropical sun, they look like flying gems. They are extremely active, darting about, and thrusting their long beaks and flexible tongues into every flower they see, in search of food. Sometimes they will remain suspended in the air for an hour or two together, vibrating their wings with

such velocity, that they cannot be seen distinctly, but appear like a mist round the body of the bird, while they make that curious humming noise from which the bird takes its name. Sometimes they quarrel, when their little throats become distended, their crests, tails, and wings expand, and they fight with inconceivable fury, till one of them falls exhausted on the ground. The most common species is *Trochilus colubris*, the Ruby-throated Humming Bird, and one of them has been kept alive in a cage for more than three months, by feeding it with sugar and water.




THE HOOPOE. (*Upupa epops.*)

THIS is a small bird, measuring no more than twelve inches from the point of the bill to the end of the tail. The bill is sharp, black, and somewhat bending. The head is adorned with a very beautiful, large, moveable crest, a kind of bright halo, the radiation of which places the head nearly in the centre of a golden circle. This pleasing ornament, which the bird sets up or lets fall at

pleasure, is composed of a double row of feathers, reaching from the bill to the nape of the neck, which is of a pale red. The breast is white, with black streaks tending downwards; the wings and back are varied with white and black cross-lines. The food of the Hoopoe consists chiefly of insects, with the remains of which its nest is sometimes so filled as to become extremely offensive. This beautifully crested bird is not at all common in this country, and is solitary, two of them being seldom seen together, while in Egypt, where Hoopoes are very common, they are often seen in small flocks. The female is said to have two or three broods in the year; and makes no nest, but lays five or six eggs in the hollow of a tree, without any sort of preparation. The name alludes to the note of the bird, which resembles the word "hoop" repeated several times in a low voice.

Though this bird is found occasionally both in England and Scotland, it never breeds with us. It is common in Italy, where its strange startling cry is often heard, without the bird being seen, as it keeps itself concealed among trees.

This bird is called *bubbola* by the Italians, most probably from its peculiar cry. It keeps itself concealed among the trees; but it is constantly heard repeating the syllable *bu, bu*, with such a strong sonorous voice that it may be heard at a great distance. Its song, properly so called, is only uttered during the honeymoon. Although the Hoopoe lives and builds in woods, it may be often seen, in search of insect food, in fields and pastures. The nest is generally placed either in the natural hollow of a tree, or in the deserted excavation of a woodpecker. It is composed outwardly of feathers, and is lined with the hair of cows and horses.



SECTION III.—Scansores, or Climbers.

THE CUCKOO. (*Cuculus canorus.*)

“ Hail, beauteous stranger of the wood,
 Attendant on the spring!
 Now Heaven repairs thy rural seat,
 And woods thy welcome sing.

“ Soon as the daisy decks the green,
 Thy certain voice we hear;
 Hast thou a star to guide thy path,
 Or mark the rolling year!

“ Delightful visitant! with thee
 I hail the time of flowers,
 When heav'n is fill'd with music sweet,
 Of birds among the bowers.”

LOGAN.

THE well-known notes of this bird, in spite of their monotony, are heard with pleasure in spring, as a sure prognostic of fine weather. The Cuckoo is generally first heard about the middle of April, and ceases towards the end of June. This bird is so shy, that he is seldom seen when uttering his singular note. The female does not build a nest, but lays her eggs in that of some other bird, generally in that of the sparrow.

The Cuckoo is somewhat less than the magpie, his length being about twelve inches from the tip of the bill to the end of the tail. He is remarkable for his round prominent nostrils; the lower part of the body is of a yellowish colour, with black transverse lines on the throat and across the breast; the head and upper part of the body and wings are beautifully marked with black and tawny stripes, and on the top of the head there are a few white spots. The tail is long, and on the exterior part, or edges of the feathers, there are several white marks; the ground colour of the body is a sort of grey. The legs are short, and covered with feathers, and the feet are composed of four toes, two before and two behind.

We are indebted to the observations of Dr. Jenner for the following account of the habits and economy of this singular bird in the disposal of its egg. He states that, during the time the hedge-sparrow is laying her eggs, which generally occupies four or five days, the Cuckoo contrives to deposit her egg among the rest, leaving the future care of it entirely to the hedge-sparrow. This intrusion often occasions some disorder; for the old hedge-sparrow, at intervals, while she is sitting, not only throws out some of her own eggs, but sometimes injures them in such a way that they become addle, so that it frequently happens that not more than two or three of the parent bird's eggs are hatched: but, what is very remarkable, it has never been observed that she has either thrown out or injured the egg of the Cuckoo. When the hedge-sparrow has set her usual time, and has disengaged the young Cuckoo, and some of her own offspring from the shell, her own young ones and any of her eggs that remained unhatched are soon turned out: the young Cuckoo

then remains in full possession of the nest, and is the sole object of the future care of the foster parent. The young birds are not previously killed, nor are the eggs demolished; but they are left to perish together, either entangled in the bush that contains the nest, or lying on the ground beneath it. On the 18th June, 1787, Dr. Jenner examined a nest of a hedge-sparrow, which then contained a Cuckoo's and three hedge-sparrow's eggs. On inspecting it the day following, the bird had hatched; but the nest then contained only a young Cuckoo and one young hedge-sparrow. The nest was placed so near the extremity of a hedge, that he could distinctly see what was going forward in it; and, to his great astonishment, he saw the young Cuckoo, though so lately hatched, in the act of turning out the young hedge-sparrow. The mode of accomplishing this was curious; the little animal, with the assistance of its rump and wings, contrived to get the bird upon its back, and making a lodgment for its burden by elevating its elbows, climbed backward with it up the side of the nest, till it reached the top; where, resting for a moment, it threw off its load with a jerk, and quite disengaged it from the nest. After remaining a short time in this situation, and feeling about with the extremities of its wings, as if to be convinced that the business was properly executed, it dropped into the nest again. Dr. Jenner made several experiments in different nests, by repeatedly putting in an egg to the young Cuckoo, which he always found to be disposed of in the same manner. It is very remarkable, that nature seems to have provided for the singular disposition of the Cuckoo, in its formation at this period; for, different from other newly-hatched birds, its back, from the scapulæ downward, is very broad, with a considerable depression in the middle, which seems intended for the express purpose of giving a more secure lodgment to the egg of the hedge-sparrow or its young one, while the young Cuckoo is employed in removing either of them from the nest. When it is about twelve days old, this cavity is quite filled up, the back assumes the shape of that of nestling birds in general, and at that time the disposition of turning out

its companion entirely ceases. The smallness of the Cuckoo's egg, which in general is less than that of the house-sparrow, is another circumstance to be attended to in this surprising transaction, and seems to account for the parent Cuckoo's depositing it in the nests of such small birds only as these. If she were to do this in the nest of a bird that produced a larger egg, and consequently a larger nestling, the design would probably be frustrated, the young Cuckoo would be unequal to the task of becoming sole possessor of the nest, and might fall a sacrifice to the superior strength of its partners. Dr. Jenner observes, that the eggs of two Cuckoos are sometimes deposited in the same nest; and gives the following instance which fell under his observation. Two Cuckoos and a hedge-sparrow were hatched in the same nest; one hedge-sparrow's egg remained unhatched. In a few hours a contest began between the Cuckoos for possession of the nest; and this continued undetermined till the afternoon of the following day, when the one which was somewhat superior in size, turned out the other, together with the young hedge-sparrow and the unhatched egg. The contest, he adds, was very remarkable: the combatants alternately appeared to have the advantage, as each carried the other several times nearly to the top of the nest, and again sank down oppressed by the weight of its burden; till at length, after various efforts, the strongest of the two prevailed, and was afterwards brought up by the hedge-sparrow.

The American Cuckoo, or Cow-bird, is quite different in its habits to the European Cuckoo, as it builds a nest for its eggs, and hatches its young itself, like other birds.



THE ROLLER (*Coracias garrula*)

Is about the size of the jay. Its bill is black, sharp, and somewhat hooked. The head is of a dirty green, mingled with blue; of which colour is also the throat, with white lines in the middle of each feather; the breast is of a pale blue, like that of a pigeon; the middle of the back, between the shoulders, is red; the rump, and less coverts of the wings are dark blue: the feet are short, and, like those of a dove, of a dirty yellow colour.

The Roller is wilder than the jay, and frequents the thickest woods; it builds its nest chiefly on birch-trees. Buffon says it is a bird of passage, and migrates in the months of May and September. In those countries where it is common, (namely, in Asia, Africa, and South America,) it is said to fly in large flocks in the autumn, and is frequently seen on cultivated grounds, with rooks and other birds, searching for worms, insects, seeds, berries, roots, and, in cases of necessity, small frogs.



THE KINGFISHER (*Alcedo ispida*)

Is the Halcyon of the ancients, and his name recalls to

our mind the most lively ideas. It was believed, that, as long as the female sat upon her eggs, the god of storms and tempests refrained from disturbing the calmness of the waves, and *Halcyon days* were, for navigators of old, the most secure times to perform their voyages.


“As firm as the rock, and as calm as the flood,
Where the peace-loving Halcyon deposits her brood.”

But although this bears analogy to a natural coincidence between the time of brooding assigned to the Kingfishers and a part of the year when the ocean is less tempestuous, yet Mythology would exercise her fancy, and turn into wonders that which was nothing else than the common course of nature.

This bird is nearly as small as a common sparrow, but the head and beak appear proportionally too big for the body. The bright blue of the back and wings claims our admiration, as it changes into deep purple or lively green, according to the angles of light under which the bird presents itself to the eye. It generally haunts the banks of rivers, for the purpose of seizing small fish, on which it subsists, and which it takes in amazing quantities, by balancing itself at a distance above the water for a certain time, and then darting on the fish with unerring aim. It dives perpendicularly into the water, where it continues several seconds, and then brings up the fish, which it carries to land, beats to death, and afterwards swallows. When it cannot find a projecting bough, it sits on some stone near the brink, or even on the gravel; but the moment it perceives the fish, it takes a spring upwards of twelve or fifteen feet, and drops from that height upon its prey.

The Kingfisher lays its eggs, to the number of seven or more, in a hole in the bank of the river or stream that it frequents. Dr. Heysham had a female brought alive to him at Carlisle by a boy, who said he had taken it the preceding night when sitting on its eggs. His information on the subject was, that “having often observed these birds frequent a bank upon the river Peteril, he had watched them carefully, and at last he saw them go into a small hole in the bank. The hole was too narrow to

admit his hand ; but, as it was made in soft mould, he easily enlarged it. It was upwards of half-a-yard long ; at the end of it the eggs, which were six in number, were placed upon the bare mould, without the smallest appearance of a nest." The eggs were considerably larger than those of the yellow-hammer, and of a transparent white colour. It appears, from a still later account that the direction of the holes is always upward ; that they are enlarged at the end, and have there a kind of bedding formed of the bones of small fish, and some other substances, evidently the castings of the parent animals. This bedding is generally half an inch thick, and mixed with earth ; and on it the female deposits and hatches her eggs. When the young ones are nearly full feathered, they are extremely voracious ; and as the old birds do not supply them with all the food they can devour, they are continually chirping, and may be discovered by their noise.



THE COMMON GREEN WOODPECKER

(*Picus viridis*)

RECEIVES his name from his habit of pecking the insects from the chinks of trees and holes in the bark. The bill is straight, strong, and angular at the end ; and in most of the species is formed like a wedge, for the purpose of piercing the trees. The nostrils are covered

with bristles. The tongue is very long, slender, cylindrical, bony, hard, and jagged at the end. The toes are placed two forward and two backward; and the tail consists of ten hard, stiff, and sharp-pointed feathers. A Woodpecker is often seen hanging by his claws, and resting upon his breast against the stem of a tree; when after darting his beak against the bark, with great strength and noise, he runs round the tree with great alacrity, which manœuvre has made the country people suppose that he goes round to see whether he has not pierced the tree through; though the fact is, the bird is in search of the insects, which he hopes to have driven out by his blow.


The following lines, from Moore's beautiful song, allude to the noise which the Woodpecker makes in searching for its food:

"I knew by the smoke that so gracefully curl'd
 Above the green elms, that a cottage was near,
 And I said, if there's peace to be found in the world,
 A heart that was humble might hope for it here.
 Every leaf was at rest, and I heard not a sound,
 But the Woodpecker tapping the hollow beech-tree."

The fact is, that this beating against the bark is for no other purpose than to rouse the insects which the chink contains, and to force them to come out, which they do from their alarm at the noise, when the Woodpecker turning round takes them unawares, and feeds upon them: if the insects do not answer the delusive call, he darts his long tongue into the hole, and brings out, by this means, his reluctant prey. The plumage of this bird is a compound of red and green, two colours, the approximation of which is always productive of harmony in the works of nature. They nestle in the hollows of trees, where the female lays five or six whitish eggs, without making any nest, trusting to the natural heat of her body to hatch them.

The Green Woodpecker is seen more frequently on the ground than the other kinds, particularly where there are ant-hills. It inserts its long tongue into the holes through which the ants issue, and draws them out in abundance.


Sometimes with its feet and bill it makes a breach in the nest, and devours the ants and their eggs at its ease. The young ones climb up and down the trees before they are able to fly; they roost very early, and repose in their holes till day. There are many different kinds of Woodpecker, five of which are common to this country.



THE TOUCAN (*Ramphastos tucanus*)

Is a native of South America, very conspicuous on the magnitude and shape of its bill; which, in some of the species, is nearly as long and as large as the body itself. The length of its body is about eighteen inches, (the size of the magpie;) the head is large and strong, and the neck short, in order the more easily to support the bulk of such a beak. The head, neck, and wings are black; the breast of a most lovely orange saffron colour; the lower part of the body and the thighs are vermillion; the tail black. Mr. Gould's specimen represents a narrow straw-coloured belt across the centre of the breast, dividing the orange tint from the vermillion. One of these

birds that was kept in a cage was very fond of fruit, which it held for some time in its beak, touching it with great delight with the tip of its feathery tongue, and then tossing it into its throat by a sudden upright jerk; it also fed on small birds, insects, caterpillars, &c.



THE GREY PARROT. (*Psittacus erythacus*.)

THE tongue of the Parrot is not unlike a black soft bean, and fills so completely the capacity of its beak, that the bird can easily modulate sounds and articulate words; the beak is composed of two pieces, both moveable, which is a peculiarity belonging almost exclusively to this tribe of birds. The bill of the Parrot is strongly hooked, and assists it in climbing, catching hold of the boughs of the trees with it, and then drawing its legs upwards; then again advancing the beak, and afterwards the feet, for its legs are not adapted for hopping from bough to bough, as other birds do. Several stories are told of the sagacity of these birds, and of the aptitude of their interrogatories and answers, but they have been no doubt the effect of chance.

Dr. Goldsmith says that a Parrot, belonging to King Henry the Seventh, having been kept in a room next the Thames, in his palace of Westminster, had learned to repeat many sentences from the boatmen and passengers. One day, sporting on its perch, it unluckily fell into the water. The bird had no sooner discovered its situation,


than it called out aloud, "A boat! twenty pounds for a boat!" A waterman, happening to be near the place where the Parrot was floating, immediately took it up, and restored it to the king; demanding, as the bird was a favourite, that he should be paid the reward the bird had called out. This was refused; but it was agreed that, as the Parrot had offered a reward, the man should again refer to its determination for the sum he was to receive. "Give the knave a groat," screamed the bird the instant the reference was made.

The memory of Parrots is very astonishing, and they can not only imitate discourse, but can sing verses of songs, and mimic gestures and actions. Scaliger saw one that performed the dance of the Savoyards, at the same time that it repeated their song. The song was well imitated, but when the bird tried to caper, it was with the worst grace imaginable, as he turned in his toes, and kept tumbling back in a most clumsy manner.

Willoughby tells us of a Parrot, which, when a person said to it, "Laugh, Poll, laugh," laughed accordingly, and the instant after screamed out, "What a fool to make me laugh!" Another, which had grown old with its master, shared with him the infirmities of age. Being accustomed to hear scarcely anything but the words, "I am sick;" when a person asked it, "How do you do, Poll?" "I am sick," it replied in a doleful tone, stretching itself out, "I am sick."

Parrots are very numerous in the East and West Indies, where they assemble in companies, like rooks, and build in the hollows of trees. The female lays two or three eggs, marked with little specks, like those of the partridge. They never breed in our climate, though they live here to a great age. They feed entirely upon vegetables, but when tame, will take from the mouth of their master or mistress any kind of chewed meat, and chiefly eggs, of which they seem particularly fond. They bite or pinch very hard, and some of them possess so much strength in their beak, that they could easily break a man's finger. The Parrot is sensible of attachment, as well as of revenge; and if in their mimic attitudes they show great

pleasure at the sight of their feeders, they also fly up with anger to the face of those who once have affronted or injured them.



THE MACAW (*Psittacus*, or *Macrocerus aracanga*)

Is the largest of the parrot tribe, and painted with the finest colours Nature can bestow. The beak is uncommonly strong; and the tail proportionally longer than that of any of the parrot tribe. Its voice is fierce and tremulous, sometimes sounding like the laugh of an old man; and it seems to utter the word "Arara," which occasions its bearing that name in its native country. The Macaw of St. Domingo is of a fine red colour, except some blue and yellow feathers about the wings. The flesh of this bird, like that of the parrots, is much esteemed as an article of food in that island.

When tame, it eats almost every article of human food, and is particularly fond of bread, beef, fried fish, pastry, and sugar. It cracks nuts with its bill, and dexterously picks out the kernels with its claws. It does not chew the soft fruits, but sucks them by pressing its tongue against the upper part of its beak: and the harder sort of food, such as bread and pastry, it bruises, or chews, by pressing the tip of the lower upon the most hollow part of the upper mandible.



THE PAROQUET (*Palæornis*)


Is less than the common parrot, and furnished with a longer tail. There are several varieties of this bird, the habits and qualities of which are similar to those of the parrots. They are even more easily tamed; and, although they do not articulate so readily, are the amusement of sedentary ladies in all countries where they are known. Parroquets are natives of the East Indies and Australia, particularly Van Diemen's Land, where they are found in great abundance.



THE RING PAROQUET. (*Palæornis Alexandri*.)

This beautiful species, no less remarkable for the elegance of its form than for its docility and imitative powers, is supposed to have been the first of the parrot species known to the ancients, from the time of Alexander the Great down to the age of Nero. It is about fifteen inches long; its bill is thick and red; the head and the body a bright green; the neck, breast, and the whole of the under side of a paler tint. It has a red circle, or ring, which encompasses the neck, and is about

the breadth of a little finger at the back ; but grows narrower by degrees towards the sides, and ends under the lower bill. The lower part of the body is of so faint a green, that it seems almost yellow. The tail also is of a yellowish green, and the legs and feet ash-coloured.



THE COCKATOO. (*Ptyctolophus galeritus.*)

THIS bird is distinguished from the parrots, by a beautiful crest, composed of a tuft of elegant feathers, which he can raise or depress at pleasure. We meet with some of a beautiful white plumage, and the inside feathers of the crest of a pleasing yellow, with a spot of the same colour under each eye, and one upon the breast. This bird often repeats the word Cockatoo, and has also been taught some others, but does not articulate so well as the parrot. The Cockatoos are natives of the Indian Islands and Australia, where they are found in great abundance.

Mr. Bennet, in speaking of the large black Cockatoo, of New Holland, says, that if this bird observes on the trunk of a tree indications of a larva being within, it diligently labours to get at it with its powerful beak and should the object of its pursuit be deep within the wood, as often happens, the trunk becomes so extensively hacked, that a slight gust of wind will lay the tree prostrate.



SECTION IV.—Gallinaceous Birds.

THE PEACOCK. (*Pavo cristatus*.)

ASTONISHED at the unparalleled beauty of this bird, the ancients could not help indulging their lively and creative fancy, in accounting for the magnificence of his plumage. They made him the favourite of imperial Juno, sister and wife to Jupiter; and not less than the hundred eyes of Argus were pulled out to ornament his tail; indeed, there is scarcely anything in nature that can vie with the transcendent lustre of the Peacock's feathers. The changing glory of his neck eclipses the deep azure of ultramarine; and at the least evolution, it assumes the green tint of the emerald, and the purple hue of the amethyst. His head, which is small and finely shaped, has several curious stripes of white and black round the eyes, and is sur-

mounted by an elegant plume, or tuft of feathers, each of which is composed of a slender stem and a small tuft at the top. Displayed with conscious pride, and exposed under a variety of angles to the reflections of versatile light, the broad and variegated disks of his tail, of which the neck, head, and breast of the bird become the centre, claim our admiration. By an extraordinary mixture of the brightest colours, it displays at once the richness of gold, and the paler tints of silver, fringed with bronze-coloured edges, and surrounding eye-like spots of dark brown and sapphire. The hen does not share in the beauty of the cock, and her feathers are generally of a light brown. She lays only a few eggs at a time, generally at an interval of three or four days; they are white and spotted, like the eggs of the turkey. She sits from twenty-seven to thirty days.


The loud screamings of the Peacock are worse than the harsh croakings of the raven, and a sure prognostic of bad weather; and his feet, more clumsy than those of the turkey, make a sad contrast with the elegance of his plumage.

“ Though richest hues the Peacock’s plumes adorn,
Yet horror screams from his discordant throat.”

The spreading of the tail, the swelling of the throat, neck, and breast, and the puffing noise which they emit at certain times, are proofs that the turkey and the Peacock stand nearly allied in the family chain of animated beings.

The flesh of the Peacock was anciently esteemed a princely dish; and the whole bird used to be served on the table, with the feathers of the neck and tail preserved; but few people could now relish such food, as it is much coarser than the flesh of the turkey. The Italians have given this laconic description of the Peacock: “ He has the plumage of an angel, the voice of a devil, and the stomach of a thief.”

THE TURKEY.



THE TURKEY (*Meleagris Gallo-Pavo*)

WAS originally an inhabitant of America, whence he was brought to Europe by some Jesuit missionaries, which accounts for his being called a Jesuit in some parts of the continent. Except as regards the plumage and the tuft on the head, he is like the peacock in many particulars. The general colour of the feathers is brown and black; and turkeys have about the head, especially the cock, naked and tuberos lumps of flesh of a bright red colour. A long fleshy appendage hangs from the base of the upper mandible, and seems to be lengthened and shortened at pleasure. The hen lays from fifteen to twenty eggs, which are whitish and freckled. The chickens are very tender, and require great care and attentive nursing, until they are able to seek their food. In the county of Nor-


folk the breeding of Turkeys, which is there a considerable branch of trade, is brought to great perfection; and some weighing upwards of twenty pounds each have been raised there. They appear to have a natural antipathy to everything of a red colour.

Though extremely prone to quarrel among themselves, they are, in general, weak and cowardly against other animals, and fly from almost every creature that ventures to oppose them. On the contrary, they pursue everything that appears to dread them, particularly small dogs and children; and after having made these objects of their aversion scamper, they evince their pride and satisfaction by displaying their plumage, strutting about among their female train, and uttering their peculiar note of self-approbation. Some instances, however, have occurred, in which the Turkey-cock has exhibited a considerable share of courage and prowess; as will appear from the following anecdote:—A gentleman of New York received from a distant part a Turkey-cock and hen, and with them a pair of bantams; which were put all together into the yard with his other poultry. Some time afterward, as he was feeding them from the barn-door, a large hawk suddenly turned the corner of the barn, and made a pounce at the bantam hen: she immediately gave the alarm, by a noise which is natural to her on such occasions; when the Turkey-cock, who was at the distance of about two yards, and without doubt understood the hawk's intention, flew at the tyrant with such violence, and gave him so severe a stroke with his spurs, as to knock him from the hen to a considerable distance; by which means the bantam was rescued from destruction.

The wild Turkey-cock is, in the American forests, an object of considerable interest. It perches on the tops of the deciduous cypress and magnolias.

“ On the top

Of yon magnolia, the loud Turkey's voice
Is heralding the dawn; from tree to tree
Extends the wakening watch-note far and wide,
Till the whole woodlands echo with the cry.” SOUTHEY.



THE GUINEA FOWL, OR PINTADO.

(*Numida Meleagris.*)

THIS bird, which is also called the *Pearled Hen*, was originally brought from Africa, where the breed is common, and seems to have been well known to the Romans, who used to esteem the flesh of this fowl as a delicacy, and admit it at their banquets. It went then by the name of Numidian Hen, or *Meleagris*, because it was fabled that the sisters of Meleager, who unceasingly deplored his death, were metamorphosed into Guinea Hens by Diana. In fact, although they are now domesticated with us, they still retain a great deal of their original freedom, and have a stupid look. Their noise is very disagreeable; it is a creaking note, which, incessantly repeated, grates upon the ear, and becomes very teasing and unpleasant. They belong to the class of birds called *pulveratores*; as they scrape the ground and roll themselves in the dust like common hens, in order to get rid of small insects which lodge in their feathers.

The Pintado is somewhat larger than the common hen; the head is bare of feathers, and covered with a naked skin of a bluish colour; on the top is a callous protuberance of a conical form. At the base of the bill on each side hangs a loose wattle, red in the female and bluish in the male. The general colour of the plumage is a dark bluish grey, sprinkled with round white spots of different sizes, resembling pearls, from which circumstance the epithet of *pearled* has been applied to this bird; which,


at first sight, appears as if it had been pelted by a strong shower of hail. These spots, which we find of a larger dimension upon some of the feathers of the pheasant, and still more so on the tail of the peacock, are convincing proofs of a near relationship between these fowls.

If trained when young, these birds may easily be rendered tame. M. Bruë informs us, that when he was on the coast of Senegal, he received, as a present from an African princess, two Guinea fowls. Both these birds were so familiar that they would approach the table and eat out of his plate; and, when they had liberty to fly about upon the beach, they always returned to the ship when the dinner or supper bell rang.

In a wild state, it is asserted that the Pintado associates in large flocks. Dampier speaks of having seen between two and three hundred of them together in the Cape de Verd Islands. They were originally introduced into our country from the coast of Africa, somewhat earlier than the year 1260.

They are sometimes caught, Mr. Gosse tells us, by the following stratagem: a small quantity of corn is steeped for a night in proof rum, and is then placed in a shallow vessel, with a little fresh rum, and the water expressed from a bitter cassava, grated. This is deposited within an enclosed ground to which the depredators resort. A small quantity of the grated cassava is then strewed over it, and it is left. The fowls eat the medicated food greedily, and are soon found reeling about intoxicated, unable to escape, and content with thrusting their heads into a corner. It is almost unnecessary to observe that in this state they become an easy prey. Pigeons are sometimes caught in this manner in Germany, by the poachers.

This bird has, of late years, greatly increased in this country, and is often seen hanging at the poultry shops and in the markets; the great abundance of them has considerably reduced their value, and they now sell, proportionally, like other fowls. The eggs are smaller and rounder than those of the common hen, and of a speckled reddish-brown colour. They are esteemed a very delicate food.



THE BRUSH TURKEY OF AUSTRALIA.

(*Megapodius tumulus.*)

It is remarkable, that this bird does not hatch its eggs by incubation. It collects together a great heap of decaying vegetables as the place of deposit of its eggs, thus making a hotbed, arising from the decomposition of the collected matter, by the heat of which the young are hatched. This mound varies in quantity from two to four cart-loads, and is not the work of a single pair of birds, but is the result of the united labour of many.

I have been credibly informed, both by natives and settlers living near their haunts, that it is not an unusual event to obtain nearly a bushel of eggs at one time from a single heap; and as they are delicious eating, they are eagerly sought after.

Mr. Gould, in his *Birds of Australia*, gives the following account of his discovery of one of these nests.

“ I landed beside a thicket, and had not proceeded far from the shore, ere I came to a mound of sand and shells, with a slight mixture of black soil, the base resting on a sandy beach, only a few feet above high-water mark; it

was enveloped in the large yellow-blossomed Hibiscus, and was of a conical form, twenty feet in circumference at the base, and about five feet in height. On pointing it out to the native, and asking him what it was, he replied, 'Oooregoorga Rambal,' Jungle-fowls' house or nest. I then scrambled up the sides of it, and, to my extreme delight found a young bird in a hole about two feet deep; it was lying on a few dry withered leaves, and appeared only a few days old. So far I was satisfied that these mounds had some connexion with the bird's mode of incubation; but I was still sceptical as to the probability of these young birds ascending from so great a depth as the natives represented, and my suspicions were confirmed by my being unable to induce the native, in this instance, to search for the eggs, his excuse being that he knew it would be no use, as he saw no traces of the old birds having recently been there. I took the utmost care of the young bird, intending to rear it if possible; I therefore obtained a moderate-sized box, and placed in it a large portion of sand. As it fed rather freely on bruised Indian corn, I was in full hopes of succeeding; but it proved of so wild and intractable a disposition, that it would not reconcile itself to such close confinement, and effected its escape on the third day. During the period it remained in captivity, it was incessantly occupied in scratching up the sand into heaps, and the rapidity with which it threw the sand from one end of the box to the other, was quite surprising for so young and small a bird, its size not being larger than that of a small quail.

"At night it was so restless, that I was constantly kept awake by the noise it made in its endeavours to escape. In scratching up the sand it only used one foot, and having grasped a handful, as it were, the sand was thrown behind it, with but little apparent exertion, and without shifting its standing position on the other leg; this habit seemed to be the result of an innate restless disposition, and a desire to use its powerful feet, and to have but little connexion with its feeding; for although Indian corn was mixed with the sand, I never detected the bird in picking any of it up while thus employed.


“ I continued to receive the eggs without having any opportunity of seeing them taken from the mound until the sixth of February ; when, on again visiting Knocker’s Bay, I had the gratification of seeing two taken from a depth of six feet, in one of the largest mounds I had then seen. In this instance the holes ran down in an oblique direction from the centre towards the outer slope of the hillock, so that although the eggs were six feet deep from the summit, they were only two or three feet from the side. The birds are said to lay but a single egg in each hole, and after the egg is deposited, the earth is immediately thrown down lightly, until the hole is filled up ; the upper part of the mound is then smoothed and rounded over. It is easily known when a Jungle-fowl has been recently excavating, from the distinct impression of its feet on the top and sides of the mound, and from the earth being so lightly thrown over, that with a slender stick the direction of the hole may readily be detected ; the ease or difficulty of thrusting the stick down indicating the length of time that has elapsed since the birds’ operations. Thus far it is easy enough ; but to reach the eggs requires no little exertion and perseverance. The natives dig them up with their hands alone, and only make sufficient room to admit their bodies, and to throw out the earth between their legs : by grubbing with their fingers alone, they are enabled to follow the direction of the hole with greater certainty, which will sometimes, at a depth of several feet, turn off abruptly at right-angles, its direct course being obstructed by a clump of wood, or some other impediment.”

In all probability, as Nature has adopted this mode of reproduction, she has also furnished the tender birds with the power of sustaining themselves from the earliest period ; and the great size of the egg would equally lead to this conclusion, since in so large a space it is reasonable to suppose that the bird would be much more developed than is usually found in eggs of smaller dimensions. The eggs are perfectly white, of a long, oval form, three inches and three quarters long by two inches and a half in diameter.

a gay and sprightly air upon some neighbouring branch to ascertain the cause of the disturbance: immediate advantage must be taken of this circumstance, or the next moment it may be half-way down the gully. So totally different is the shooting of this bird to anything practised in Europe, that the most expert shot would have but little chance, until well experienced in the peculiar nature of the country, and the habits of the bird. The *Menura* seldom, if ever, attempts to escape by flight; it easily eludes pursuit by its extraordinary power of flying. None are so efficient in obtaining specimens as the naked black, whose noiseless and gliding steps enable him to steal upon it unheard and unperceived, and with the gun in his hand, he rarely allows it to escape, and in many instances he will even kill it with his own weapons.

“The Lyre-bird is of a wandering disposition, and although it probably keeps to the same brush, it is constantly engaged in traversing it from one end to the other, from mountain-top to the bottom of the gullies, whose steep and rugged sides present no obstacle to its long legs and powerful muscular thighs: it is also capable of performing extraordinary leaps; and I have heard it stated, that it will spring ten feet perpendicularly from the ground. It appears to be of solitary habits, as I have never seen more than a pair together, and these only in a single instance; they were both males, and were chasing each other round and round with extreme rapidity, apparently in play, pausing every now and then to utter their loud shrill calls; while thus employed they carried the tail horizontally, as they always do when running quickly through the bush, that being the only position in which this great organ could be conveniently borne at such times. Among its many curious habits, the only one at all approaching to those of the *Gallinacæa*, is that of forming small round hillocks, which are constantly visited during the day, and upon which the male is constantly trampling, at the same time erecting and spreading out his tail in the most graceful manner, and uttering his various cries, sometimes pouring forth his natural notes, at others mocking those of other birds, and even

the howling of the native dog, or dingo. The early morning and the evening are the periods when it is most animated and active."



THE PHEASANT. (*Phasianus colchicus.*)

THE name of this bird implies that he was originally a native of the banks of the river Phasis in Armenia; how and when he emigrated, and began to frequent our groves, is unknown. He is of the size of the common cock; the bill is of a pale horn colour; the nostrils arched; the eyes yellow, and surrounded by a naked warty skin, of a beautiful scarlet, finely spotted with black; immediately under each eye there is a small patch of short feathers, of a dark glossy purple; the upper parts of the head and neck are of a deep purple, varying to glossy green and blue; the lower parts of the neck and breast are of a reddish chestnut, with black indented edges; the sides and lower part of the breast are of the same colour, with tips of black to each feather, which, in different lights, vary to glossy purple; indeed, the whole colour of this half-domesticated fowl is very beautiful, uniting the brightness of deep yellow gold to the finest tints of the ruby and turquoise, with reflections of green;


the whole being set off by several spots of shining black ; but in this as in every other kind of gorgeously-feathered birds, nature has for some wise purposes, yet unknown to us, denied the female that admirable beauty of plumage which belongs to the male. The Pheasant lives in the woods, which he leaves at dusk to perambulate corn-fields and other sequestered places, where he feeds with his females, upon acorns, berries, grain, and seeds of plants, but chiefly on ants' eggs, of which he is particularly fond. His flesh is justly accounted better meat than any of the domestic or wild fowls, as it unites the delicacy of the common chicken to a peculiar taste of its own. The female lays eighteen or twenty eggs once a year, in the wild state ; but it is in vain that we have attempted to domesticate this bird entirely, as she never will remain patiently confined, and if she ever breeds in confinement, is very careless of her brood.

There are great varieties of Pheasants, of extraordinary beauty and brilliancy of colours : many of these, brought from the rich provinces of China, are kept in aviaries in this kingdom : the common Pheasant is likewise a native of the East, and is the only one of its kind that has multiplied in our island ; of late years they have become so numerous in our preserves, that at feeding time in the morning, they may be seen in the neighbouring fields in large numbers, running about almost as tame as chickens.

This beautiful bird is elegantly described in the following passage :—

“ See! from the brake the whirring Pheasant springs,
 And mounts exulting on triumphant wings ;
 Short is his joy ; he feels the fiery wound,
 Flutters in blood, and panting beats the ground :
 Ah ! what avails his glossy, varying dyes,
 His purple crest, his scarlet-circled eyes,
 The vivid green his shining plumes unfold,
 His painted wings, and breast that flames with gold !”

POPE'S WINDSOR FOREST.



THE RED-LEGGED PARTRIDGE.

(Perdix rufus.)

THESE Partridges are natives of Guernsey and Jersey; but are also very frequently found on the adjoining coasts of France and England. In the Western districts of France they are very abundant, and their flesh is plump and juicy. In England it is as white as in France, but more dry. The side-feathers are very handsomely speckled, and there is a rich black mark beginning behind the eye and forming a kind of gorget on the breast. The eyelids are of a bright red, as are the legs and toes, and the claws are brown. They build their nests on the ground; but are sometimes found perched on trees, or on a fence or paling.


The common Partridge (*Perdix cinerea*), is in weight about fourteen ounces. The shape of this bird is peculiar to itself and the quail, both having a very short tail; the plumage, although it cannot boast of gaudiness, is very pleasing to the eye, being a mixture of brown


and fawn-colour, interspersed with gray and ash-colour tints. The head is small and pretty; the beak strong, but short, and resembling that of all other granivorous birds. The female lays fifteen or eighteen eggs, and leads her brood in the corn-fields with the utmost care. Young Partridges are among the birds which run fleetly the moment they come out of the shell, and may sometimes be found running with a piece of the shell still remaining on their heads. The affection of Partridges for their offspring is peculiarly interesting. Both the parents lead them out to feed: they point out to them the proper places for their food, and assist them in finding it by scratching the ground with their feet. They frequently sit close together, covering the young ones with their wings; and from this position they are not easily roused. If, however, they are disturbed, most people acquainted with rural affairs know the confusion that ensues. The male gives the first signal of alarm, by a peculiar cry of distress; throwing himself at the same moment more immediately into the way of danger, in order to mislead the enemy. He flutters along the ground, hanging his wings, and exhibiting every symptom of debility. By this stratagem he seldom fails of so far attracting the attention of the intruder as to allow the female to conduct the helpless unfledged brood into some place of security.

The nest is usually on the ground; but on the farm of Lion Hall, in Essex, belonging to Colonel Hawker, a Partridge, in the year 1788, formed her nest, and hatched sixteen eggs, on the top of a pollard oak-tree! What renders this circumstance the more remarkable is, that the tree had, fastened to it, the bars of a stile, where there was a footpath; and the passengers, in going over, discovered and disturbed her before she sat close. When the brood was hatched, the birds scrambled down the short and rough boughs, which grew out all around the trunk of the tree, and reached the ground in safety. It has long been a received opinion among sportsmen, as well as among naturalists, that the female Partridge has none of the bay feathers of the breast like the male. This,

however, is a mistake; for, Mr. Montagu happening to kill nine birds in one day, with very little variation as to the bay mark on the breast, he was led to open them all, and discovered that five of them were females. On carefully examining the plumage, he found that the males could only be known by the superior brightness of colour about the head; which alone, after the first or second year, seems to be the true mark of distinction. They fly in coveys till about the third week in February, when they separate and pair; but if the weather be very severe, it is not unusual to see them collect together again. We are told that a gamekeeper, in Dorsetshire, hearing a partridge utter a cry of distress, was attracted by the sound into a field of oats, when the bird ran round him very much agitated; upon his looking among the corn, he saw in the midst of her infant brood a large snake, which he killed; and perceiving its body much distended, he opened it, when to his astonishment two young Partridges ran from their prison, and joined their mother; two others were found dead in its stomach. Partridges have ever held a distinguished place at the tables of the luxurious: we have an old distich:—

“ If the Partridge had the woodcock’s thigh,
’T would be the best bird that e’er did fly.”






THE QUAIL (*Coturnix vulgaris*)

Is a small bird, being in length no more than seven inches. The colour of the breast is a dirty pale yellow, and the throat has a little mixture of red: the head is black, and the body and wings have black stripes upon a hazel-coloured ground. Its habits and manner of living resemble those of the partridge, and it is either caught in nets by decoy birds, or shot by the help of the setting-dog, its call being easily imitated by tapping two pieces of copper one against another. The flesh of the Quail is very luscious, and next in flavour to that of the partridge. Quails are birds of passage, the only peculiarity in which they differ from all other of the poultry kind; and such prodigious numbers have sometimes appeared on the western coast of the kingdom of Naples, that one hundred thousand have been caught in one day, within the space of three or four miles. In some parts of the south of Russia, they abound so greatly, that at the time of their migration they are caught by thousands, and sent in casks to Moscow and St. Petersburg. The female seldom lays more than six or seven eggs.

The ancient Athenians kept this bird, merely for the sport of fighting with each other, as game cocks do, and never ate the flesh. The Quail was that wild fowl which God thought proper to send to the chosen people of Israel, as a sustenance for them in the desert.

The Chinese Quail is a beautiful little bird, and often kept in cages in China, for the singular purpose, as it is said, of warming people's hands in winter; as taking the soft, warm body of the bird in the hand diffuses through it an agreeable warmth.

The American Quail (*Ortyx borealis*) is larger than the common Quail, and is something between a Quail and a partridge.



THE RED GROUSE. (*Lagopus Scoticus.*)


“High on exulting wing the Heath-Cock rose,
And blew his shrill blast o'er perennial snows.”

ROGERS.

THIS bird is called by some ornithologists the *Moor-Cock*, and by others *Red Game*. The beak is black and short; over the eyes there is a bare skin of a bright red. The general colour of the plumage is red and black, variegated, and intermixed with each other, except the wings, which are brownish, spotted with red, and the tail, which is black; the legs are covered with long, thick feathers down to the very toes. It is common in the north of England, in Scotland, and in Wales; and not

only affords great diversion to the noblemen and gentlemen of those countries who are fond of shooting, but also repays them well for their trouble, as the flesh is very delicate, and holds on our table an equal place with the partridge and the pheasant. The season of Grouse shooting commences August 20. In winter they are usually found in flocks of sometimes fifty to one hundred in number, which are termed by sportsmen *packs*, and become remarkably shy and wild, seldom allowing the sportsman to approach them within one hundred yards. They keep near the summits of the heathy hills, and seldom descend to the lower grounds. Here they feed on the mountain berries, and on the tender tops of the heath. The hen lays seven or eight eggs of a reddish black colour.

Besides the red Grouse, there are the following of the same species, but they are now found only in the wildest parts of the Highlands of Scotland.



THE BLACK COCK (*Tetrao tetrix*)

Is about four pounds in weight; but the female, which is usually called the Grey Hen, is often not more than two. The plumage of the whole body of the male is black, and glossed over the neck and rump with shining blue;

the coverts of the wings are of a dusky brown, with the quill feathers black and white. The tail is much forked. These birds never pair; but in the spring the males assemble at their accustomed haunts on the tops of heathy mountains, where they crow and clap their wings.

“ And from the pine’s hig top brought down
The giant Grouse, while boastful he display’d
His breast of varying green, and crow’d and clapp’d
His glossy wings.”

GISBORNE.

The females, at this signal, resort to them. The males are very quarrelsome, and fight together like game cocks. On these occasions, they are so inattentive to their own safety, that two or three have sometimes been killed at one shot; and instances have occurred of their having been knocked down with a stick.

Like the Capercailzie, or Cock of the Woods, a larger species of this genus, these birds are common in Russia, Siberia, and other northern countries, chiefly in wooded and mountainous situations; and in the northern parts of our own island on uncultivated moors.

The *Ptarmigan*, or *White Grouse*, is somewhat larger than a pigeon; its bill is black, and its plumage in summer is of a pale brown colour, elegantly mottled with small bars and dusky spots. The head and neck are marked with broad bars of black, rust-colour, and white, the wings and belly are white. The *White Grouse* is fond of lofty situations, where it braves the severest cold. It is found in most of the northern parts of Europe, even as far as Greenland. In this country it is only to be met with on the summits of some of our highest hills, chiefly in Scotland, and in the Hebrides and Orkneys, but sometimes in Cumberland and Wales.



THE COMMON COCK. (*Gallus domesticus.*)

“ While the Cock, with lively din,
Scatters the rear of darkness thin ;
And to the stack, or the barn door
Stoutly struts his dames before.”

MILTON.

THIS bird is so well known, that it would be needless to say much of him. His plumage is various and beautiful, his courage very great and proverbial, his sobriety astonishing, and his intuitive knowledge of the period of sun-rise has baffled the most scrutinizing researches of naturalists. When of a good breed, and well taught to fight, he will die rather than yield to his adversary. The hen lays a great number of eggs, and will hatch as many as thirteen at one sitting; but this is considered the extreme number, being as many as she can well cover. When in the secluded state of incubation she eats very little; and yet is so courageous and strong, that she will rise and fight any men or animals that dare to approach her nest. It is impossible to conceive how, with such


a scanty sustenance as she takes, she can, for twenty-one days, emit constantly from her body, as much heat as would raise Fahrenheit's thermometer to ninety-six degrees. The flesh of this bird is delicate and wholesome, and universally relished as nourishing and agreeable food.

There are several varieties or families of this fowl. The Hamburgh Cock has a beautiful tuft of feathers about his ears, and on the top of his head; and the Bantam has his legs and toes entirely feathered, which is more an impediment than an ornament to the bird.

The cruel sport of cockfighting may be traced back to the earliest antiquity. The Athenians seem to have received it from India, where it is even now followed with a kind of frenzy; and we are told that the Chinese will sometimes risk not only the whole of their property, but their wives and children on the issue of a battle. The religion of the Greeks could not see that game with pleasure, and therefore cockfighting was allowed only once a year; but the Romans, who had given to the bird the name of their earliest enemies, the Gauls, adopted the practice with rapture, and introduced it into this island. Henry VIII. delighted in this sport, and caused a commodious house to be built for the purpose, which, although now applied to a very different use, still retains the name of the Cockpit. The part of our ships so called, seems also to indicate that in former times the diversion of cockfighting was permitted in order to beguile the tedious hours of a long voyage. It is a great pity that a custom, which originated in barbarous times, should still continue, to the disgrace of a philosophic and enlightened age.—The Cock has been a subject of considerable interest with the poets; and has been very commonly called by them "Chanticleer."

"Within this homestead lived, without a peer
For crowing loud, the noble Chanticleer." DRYDEN.

"The feathered songster, Chanticleer,
Had wound his bugle-horn,
And told the early villager
The coming of the morn." CHATTERTON.



**BANKIVA COCK.—JAGO COCK AND HEN.
SPANISH COCK AND HEN.**

FROM the Bankiva fowl nearly all the various kinds of fowls found in British poultry-yards are said to be sprung. It is a native of the island of Java, and is characterised by a red indented comb, red wattles, and ash-grey legs and feet. The cock has a thin indented or scalloped comb, and wattles under the mouth. The feathers of the neck are long, falling down, and rounded at the tips, and are of the finest gold colour. The head and neck are fawn-coloured, the wing-coverts dusky brownish and black; the tail and belly black. The hen is of a dusky ash-grey and yellowish colour, and has a much smaller comb and beard than the cock.

THE PADUAN, OR JAGO FOWL.

(Gallus Giganteus.)

The wild species, termed by Marsden the Jago Fowl, is a native of Java and Sumatra, and is supposed by Temminck to be the original of this fine breed, though little is known of the wild sort, further than that it is double the size of the Bankiva, or dunghill fowl. Marsden says he has seen in the East a cock of this species tall enough to pick crumbs from a dining-table. They are said to weigh from eight to ten pounds. The combs of both the cock and hen are large, frequently double, of the form of a crown, with a tufted crest of feathers, which is largest in the hen; the voice is stronger and harsher than that of other fowls; but the most singular peculiarity is, that they do not come into full feather till about half-grown. The Cochinchina fowls, in the possession of her Majesty, are said to be a variety of the Jago Fowls. There are numerous hybrids and varieties of the Jago Fowl found under different names in poultry-yards, but all of them lay fine large eggs, and are highly esteemed for the excellent flavour of their flesh. One of the most interesting of these varieties is called

THE SPANISH FOWL,

the body and tail feathers of which are of a rich black, with occasionally a little white on the breast. The cock of this variety is a most majestic bird; its deportment is grave and stately, and its eyes are encircled with a ring of brown feathers, from which rises a black tuft that covers the ears. There are other similar feathers behind the comb and beneath the wattles. The legs and feet are of lead colour, except the sole of the foot, which is yellowish.

THE BANTAM FOWL

Is a small variety, with short legs, most frequently

feathered to the toes, so as sometimes to obstruct walking. Many Bantam fanciers prefer those which have clear bright legs, without any vestige of feathers. The full-bred Bantam cock should have a rose comb, a well-feathered tail, full hackles, a proud lively carriage, and ought not to weigh more than a pound. The nankeen coloured and the black are the greatest favourites. If of the latter colour, the bird should have no feathers of any other sort in his plumage. The nankeen bird should have his feathers edged with black, his wings barred with purple, his tail feathers black, his hackles slightly studded with purple, and his breast black, with white edges to the feathers. The hens should be small, clean-legged, and match in plumage with the cock.



THE DODO. (*Didus ineptus*.)

SWIFTNESS has generally been considered the attribute of birds; but the Dodo appears never to have had any title to this distinction. Instead of exciting the idea of swiftness by its appearance, in the drawings that have been preserved of it, it strikes the imagination as a thing the most unwieldy and inactive of all nature. Its body is massive, almost round, and covered with gray feathers. It is just barely supported upon two short thick legs, like pillars; while its head and neck rise from it in a manner truly grotesque. The neck, thick and puffy, is joined to the head, which consists of two immense jaws, which open far beyond the eye. The Dodo formerly inhabited the Isle of France; but it has been long extinct.

The Dodo was supposed by the earliest naturalists who described it, to be a kind of turkey, as in the flavour of its flesh it resembled that bird. Later naturalists supposed it to be a kind of swan, and this opinion was followed by the celebrated Buffon. Others thought it was a kind of vulture; and others, judging from the shortness of its wings, placed it in the ostrich tribe. Modern naturalists, however, having carefully examined the bones of the bird

which have been preserved, are of opinion that it is a gigantic pigeon. An entire specimen existed about a hundred years ago in the Ashmolean Museum at Oxford, but only part of the bird, and one of the feet remain; there is also a foot preserved in the British Museum. There is a reference to this extinct species in Humboldt's *Cosmos*, see Bohn's edition, vol. i. page 29, and a note on the Dodo by Dr. Mantell at the end of the volume.

The *Solitaire* is another remarkable bird which was formerly found in the Mauritius and the adjoining islands; but which has now become extinct.




THE RINGDOVE, CUSHAT, OR WOOD PIGEON,

(*Columba palumbus*,)

Is the largest Pigeon found in our island, by which it may be distinguished from all others; its weight is about twenty ounces; its length eighteen inches, and its circumference about thirty. It is usually known as the Wood Pigeon. This bird is of a pale chocolate or cream-colour, with a black ringlet, having white edges, round the neck; the breed is common in Britain. Its habits are like those of other birds of the tribe; but it is so strongly attached to its native freedom, that all attempts to domesticate it, with a few rare exceptions, have hitherto proved ineffectual.

These birds build their nest chiefly on the pine, or holly, with dried sticks thrown rudely together; and the eggs, which may frequently be seen through the bottom of the nest, are larger than those of the domestic Pigeon. Attempts have been made to domesticate them by hatching their eggs under the common Pigeon, but as soon as they were able to fly, they always escaped to their proper haunts.

Mr. Montague bred up a curious assemblage of birds, which lived together in perfect amity: it consisted of a common pigeon, a Ringdove, a white owl, and a sparrow-hawk; the Ringdove was master of the whole.



THE STOCKDOVE, OR WILD PIGEON.

(*Columba oenas.*)

“ The Stockdove, recluse, with her mate,
 Conceals her fond bliss in the grove,
 And murmuring seems to repeat,
 That May is the mother of love.” CUNNINGHAM.

THIS bird is called the Stockdove, because it builds in the stocks of trees, which have been headed down and are become thick and bristly; and not, as some have supposed, because it is the stock, or original, from which all

the tame pigeons have sprung. Sometimes these birds lay their eggs in deserted rabbit-warrens, on the sod, without making any nest.


The colour of the Stockdove is generally of a deep slate or lead tint, with rings of black about the feathers. Multitudes of them visit this country in the winter, from their more northerly summer retreats. They appear about November and retire again (except a few that breed with us) in the spring. While the beech woods were suffered to cover large tracts of ground, these birds used to haunt them in myriads, frequently extending above a mile in length, as they went out in the morning to feed. They are, however, still found in considerable quantities in many parts of England, but never in Scotland, forming their nests in the hollows of trees; not, like the ring-dove, on boughs. Their murmuring strains, or cooings, in the morning and at dusk, are highly pleasing, and throw an agreeable melancholy on the solitude of the grove. The poet of the Seasons expresses this in the following lines, with a beautiful instance of imitative harmony.

“————— the Stockdove breathes
A melancholy murmur through the whole.”

SPRING.

Wordsworth also gives a pleasing description of the mournful cooing of these birds:

“ I heard a Stockdove sing or say
His homely tale this very day ;
His voice was buried among trees,
Yet to be come at by the breeze ;
He did not cease ; but cooed, and cooed ;
And somewhat pensively he wooed ;
He sang of love with quiet blending,
Slow to begin, and never ending ;
Of serious faith and inward glee,
That was the song,—the song for me ”

THE TAME PIGEON. (*Columba livia*.)

The shape of this bird is well known, but the colour varies so much, that it defies the rules of classification. They prefer a gregarious life, and abide often, to the number of five or six thousand, in a cot built expressly for them in the neighbourhood of a farm-yard, with proper holes to nestle in. The female Pigeon, through the whole species, lays two eggs at a time, which produce generally a male and a female. It is pleasing to see how eager the male is to sit upon the eggs, in order that his mate may rest and feed herself. The young ones when hatched, require no food for the first three days, warmth being their only nourishment; they are then fed from the crop of the mother; who has the power of forcing up the half-digested peas which she has swallowed to give them to her young. The young ones, open-mouthed, receive this tribute of affection, and are thus fed three times a day.


There are upwards of twenty varieties of the domestic Pigeon, and of these the carriers are the most celebrated. They obtain their name from being sometimes employed to convey letters or small packets from one place to another. The rapidity of their flight is very wonderful. Lithgow assures us that one of them will carry a letter from Babylon to Aleppo (which, to a man, is usually thirty days' journey) in forty-eight hours. To measure their speed with some degree of exactness, a gentleman, many years ago, on a trifling wager, sent a Carrier Pigeon from London, by the coach, to a friend at Bury St. Edmond's; and along with it a note, desiring that

the Pigeon, two days after its arrival there, might be thrown up precisely when the town clock struck nine in the morning. This was accordingly done, and the Pigeon arrived in London at half-past eleven o'clock on the same morning, having flown seventy-two miles in two hours and a half. From the instant of its liberation, its flight is directed through the clouds, at a great height, to its home. By an instinct altogether inconceivable, it darts onward, in a straight line, to the very spot whence it was taken, but how it can direct its flight so exactly will probably for ever remain unknown to us.

“ Led by what chart, transports the timid Dove
 The wreaths of conquest, or the vows of love?
 Say through the clouds what compass points her flight?
 Monarchs have gazed, and nations blessed the sight.
 Pile rocks on rocks, bid woods and mountains rise,
 Eclipse her native shades, her native skies;—
 'Tis vain! through ether's pathless wilds she goes,
 And lights at last where all her cares repose.
 Sweet bird, thy truth shall Harlem's walls attest,
 And unborn ages consecrate thy nest.” ROGERS.

The Carrier Pigeon is easily distinguished from the other varieties, by a broad circle of naked white skin round the eyes, and by its dark blue or blackish colour.

It would be as fruitless as unnecessary to attempt to describe all the varieties of the Tame Pigeon; for human art has so much altered the colour and figure of this bird, that pigeon-fanciers, by pairing a male and female of different sorts, can, as they express it, “breed them to a feather.” Hence we have the various names of Carriers, Tumblers, Jacobins, Croppers, Pouters, Bunts, Turbits, Shakers, Fantails, Owls, Nuns, &c., all of which may at first have accidentally varied from the stock-dove, and these have been further improved by crossing, food, and climate. An actual post-system, in which pigeons were the messengers, was established by the Sultan Nouredin Mahmoud, which lasted about a century, and ceased in 1258, when Bagdad fell into the hands of the Moguls.



THE TURTLE DOVE. (*Columba turtur.*)

“ Go, beautiful and gentle Dove,
 And greet the morning ray;
 For lo! the sun shines bright above,
 And the rain is pass'd away.” BOWLES.

THIS Dove brings to the heart and mind the most pleasing recollections; its name is nearly synonymous with faithfulness and unvariable affection. The male or female is so much attached to its respective mate that it is said, perhaps with more poetry than truth, that if one die the other will never survive; however, the author of these observations was an eye-witness to the death of a female Turtle Dove, who was unfortunately killed by a spaniel, in the absence of the male; the disconsolate sur-

vivor, after having in vain searched everywhere for his mate, came and mournfully perched upon the wonted trough, waiting patiently for her to repair thither in order to get food; but, after two days of unavailing expectation, he, by spontaneous abstinence, pined and died on the place. Such examples are not common; and, we believe, that, when not domesticated, the appearance of another female, in the time of coupling, sets at defiance all natural propensity to constancy, and puts an end to the much-famed disconsolate widowhood. Their general colour is a bluish grey; the breast and neck of a whitish purple, with a ringlet of beautiful white feathers with black edges about the sides of the neck. Nothing can express the sensation which is excited in a feeling mind, when the tender and sweetly plaintive notes of the Turtle Dove breathe from the grove on a beautiful spring evening.

“ Deep in the wood, thy voice I list, and love
Thy soft complaining song, thy tender cooing;
Oh! what a wining way thou hast of wooing,
Gentlest of all thy race—sweet Turtle-dove!
Thine is a note which doth not pass away
Like the light music of a summer's day,
Hushing the voice of mirth and staying folly,
And waking in the breast a gentle melancholy.”—INGLIS.





SECTION V.—Grallatores, or Waders.

THE OSTRICH. (*Struthio camelus*.)

THIS bird is a native of Africa, and is so tall that when it holds up its head it is eleven feet in height. The head is very small in comparison with the body, being hardly bigger than one of the toes; it is covered, as well as the neck, with a kind of down, or thin-set hairs, instead of feathers. The sides and thighs are entirely bare and flesh-coloured. The lower part of the neck, where the feathers begin, is white. The wings are short, and of no use in flying, but when the bird runs, which it does with a strange jumping kind of motion, it raises its short wings, and holds them quivering over its back, where they seem to serve as a kind of sail to gather the wind, and

carry the bird onwards. The feathers of the back, in the cock, are coal black; in the hen only dusky, and so soft that they resemble a kind of wool. The tail is thick, bushy, and round; in the cock whitish, in the hen dusky with white tops. These are the feathers so generally in requisition, to decorate the head-dress of ladies, and the helmets of warriors.

The Ostrich swallows anything that presents itself, leather, glass, iron, bread, hair, &c.; and the power of digestion in the stomach is so strong that even iron is very much affected by it. An Ostrich in the Zoological Gardens in the Regent's Park, was, however, killed by swallowing a lady's parasol.

“ O'er the wild waste the stupid Ostrich strays,
In devious search to pick her scanty meal,
Whose fierce digestion gnaws the temper'd steel.”

MICKLE'S LUSIAD.

They are polygamous birds; one male being generally seen with two or three, and sometimes with five females. The female Ostrich, in the tropical regions, after depositing her eggs in the sand, trusts them to be hatched by the heat of the climate, and leaves the young ones to provide for themselves; and in the book of Job there is a beautiful passage relating to this habit of the Ostrich, “ which leaveth her eggs in the earth, and warmeth them in the dust; and forgetteth that the foot may crush them, or that the wild beast may break them. She is hardened against her young ones, as though they were not hers. Her labour is in vain; without fear, because God hath deprived her of wisdom; neither has he imparted to her understanding. What time she lifteth up her head on high, she scorneth the horse and his rider.” In colder regions, however, the female Ostrich sits upon her eggs like other birds, and brings up her young. The eggs are as large as a young child's head, with a hard stony shell; and one has been known to weigh upwards of three pounds. The time of incubation is six weeks. That Ostriches have great affection for their offspring, may be inferred from the assertion of Professor Thunberg, who says that he

once rode past the place where a hen Ostrich was sitting in her nest; when the bird sprang up and pursued him, evidently with a view to prevent his noticing her eggs or young. Every time he turned his horse towards her, she retreated ten or twelve paces; but as soon as he rode on again, she pursued him, till he had got to a considerable distance from the place where he had started her. In the tropical regions, some persons breed Ostriches in flocks; for they may be tamed with very little trouble; and in their domestic state, few animals may be rendered more useful. When M. Adanson was at Podar, a French factory on the southern bank of the river Niger, two young, but full-grown Ostriches, belonging to the factory, afforded him a very amusing sight. They were so tame, that two little blacks mounted both together on the back of the largest. No sooner did he feel their weight, than he began to run as fast as possible, and carried them several times round the village; and it was impossible to stop him otherwise than by obstructing the passage. This sight pleased M. Adanson so much, that he wished it to be repeated; and to try their strength, directed a full-grown negro to mount the smaller, and two others the larger of the birds. This burden did not seem at all disproportioned to their strength. At first, they went at a tolerably sharp trot; but when they became a little heated, they expanded their wings, as though to catch the wind, and moved with such fleetness that they scarcely seemed to touch the ground. The foot of the Ostrich has only two toes, one of which is extremely large and strong to make its way through the moving sands of the desert. There is a remarkably fine Ostrich at the Surrey Zoological Gardens.




THE CASSOWARY (*Casuarus galeatus*)

Is next in size to the ostrich, but of a different nature. His wings are hardly perceptible, being very short, and entirely concealed under the plumage. The general tint of his feathers is brown, with some spots of vermilion; his head is small and depressed, with a horny crown; the head and neck are deprived of feathers, and only set with a kind of hairy down. From the bill to the claws the body measures about five feet and a half; about the neck are two protuberances of a bluish colour, in shape like the wattles of a cock. Unlike other birds, the feathers of the wings, and other parts of the body, are exactly the same; so that at a distance he looks as if he were entirely covered with the hairs of a bear, rather than with the plumage of a bird. The Cassowary is as voracious as the Ostrich, and eats indiscriminately whatever comes in his way, and does not seem to have any

sort of predilection in the choice of his food. The Dutch travellers assert that he can devour not only glass, iron, and stones, but even burning coals, without testifying the smallest fear, or sustaining the least injury; and it is said that the passage of his food is performed so speedily, that even eggs will pass unbroken. He is a native of the southern parts of India; the eggs of the female are nearly fifteen inches in circumference, of a greyish ash-colour, marked with green. It has been said of the Cassowary, that he has the head of a warrior, the eye of a lion, the armament of a porcupine, and the swiftness of a courser.

A Cassowary once kept in the menagerie of the museum at Paris, devoured every day between three and four pounds weight of bread, six or seven apples, and a bunch of carrots. In summer it drank about four pints of water in the day; and in winter somewhat more. It swallowed all its food without bruising it. This bird was sometimes ill-tempered and mischievous; and much irritated when any person approached it of a dirty or ragged appearance, or dressed in red clothes; and frequently attempted to strike at them by kicking forward with its feet. It has been known to leap out of its enclosure, and to tear the legs of a man with its claws.

The Cassowary is very vigorous and powerful; its beak being, in proportion, much stronger than that of the ostrich, it has the means of defending itself with great advantage, and of easily pulling down and breaking in pieces almost any hard substance. It strikes in a very dangerous manner with its feet either behind or before, not unlike the kicking of a horse, at any object which offends it, and runs with surprising swiftness.



THE EMEU. (*Dromaius Novæ Hollandiæ.*)

THE head of this bird is without any horny crest, and feathered, but the cheeks and throat are nearly naked. The general colour is a dull brown, mottled with a dingy grey; and the young are striped with black. In appearance it closely resembles the ostrich, next to which it is the tallest bird known, but is of a more thick-set and clumsy make; though at the same time very swift and strong, and able to make a formidable defence against its hunters and their dogs, by kicking in a very vigorous and dangerous manner. It is, however, very docile, and if taken young may be easily tamed. The flesh is considered excellent eating, and is said to possess a flavour something between a sucking-pig and a turkey. The only sound that this bird emits is a low drumming noise, which it does by means of a valve attached to the lungs. The female Emeu lays her eggs in different places, but they are afterwards collected by her, or the male, by rolling them to one place, when the two birds sit on them alternately.



THE CRANE. (*Grus cinerea.*)

CRANES frequent marshy places, and live upon small fish and water-insects. Their long beaks enable them to search the water and mud for their prey, and their long necks prevent the necessity of their stooping to pick up from between their feet the objects of their search. The top of the head is black, destitute of feathers, and covered with much hairy down; the throat and sides of the neck are of a black hue; the back, the coverings of the feathers, and the body are ash-coloured. They used to be common in the fen countries, Lincolnshire and Cambridgeshire; but are not now so frequently seen in England as formerly. In their flight, Cranes mount high in

the air, and their voices can be heard even when the birds cease to be perceptible to the eye; and it is said that their sight is so keen, that they discover at a great distance any field of corn, or other food which they are fond of, and presently alight and enjoy it. These depredations they generally commit during the night, trampling down the ground as if it had been marched over by an army. They generally form themselves in the air into the shape of a wedge, in order to cut the adverse winds with greater facility.

“—— Part more wise,

In common, ranged in future, wedge their way,
 Intelligent of seasons, and set forth
 Their airy caravan high over seas
 Flying, and over lands, with mutual wing
 Easing their flight. So steers the prudent Crane
 Her annual voyage, borne on winds. The air
 Floats as they pass, fann'd by unnumber'd wings.”

MILTON.

This bird lives to a considerable age; and, as it is easily tamed, it has been ascertained that the Crane often reaches his fortieth year.


According to Kolben, they are often observed in large flocks on the marshes about the Cape of Good Hope: He says, he never saw a flock of them on the ground that had not some placed apparently as sentinels, to keep a look out, while the others are feeding; who, on the approach of danger, immediately give notice to the rest. These sentinels stand on one leg; and at intervals stretch out their necks, as if to observe that all is safe. On notice being given of danger, the whole flock are in an instant on the wing. Kolben also adds, that in the nighttime each of the watching Cranes, which rest on their left legs, hold in their right claw a stone of considerable weight, in order that, if overcome by sleep, the falling of the stone may awake them.



THE BALEARIC CRANE, OR CROWNED
DEMOISELLE, (*Anthropoides pavonina*.)

Is originally, as the name expresses, a native of Majorca and Minorca, in the Mediterranean sea, which were formerly called the Balearic Isles, but is chiefly found now in the Cape Verd Islands. The shape of its body is not unlike that of the common Crane, but it has a principal and distinctive mark on the head; which is, a tuft of hairs, or rather strong greyish bristles, standing out like rays in all directions, from which peculiarity this species take its other name of the Crowned Heron. They roost and feed in the manner of peacocks.

The Demoiselle, or Numidian Crane, (*Anthropoides Virgo*.) is remarkable for the grace and symmetry of its form, and the elegance of its deportment. It is rather larger than the species above described, and is a native of many parts of Asia and Africa. It frequents damp and marshy places, in search of small fishes, frogs, &c., which are its favourite food. It is easily domesticated.



THE STORK. (*Ciconia alba*.)

THE neck, head, breast, and body of this bird are white, the rump and exterior feathers of the wings black; the eyelids naked; the tail white, and the legs long, slender, and of a reddish colour. Storks are birds of passage. When leaving Europe they assemble together on some particular night, and all take their flight at once. As they feed on frogs, lizards, serpents, and other noxious creatures, it is not to be expected that man should be inimical to them, and therefore they have been generally a favourite with the nations they frequent. The Dutch have laws against destroying them: they are therefore very common in Holland; and build their nests and rear

their young on the tops of houses and chimneys in the middle of its most frequented and populous cities, and may be seen by dozens familiarly walking about the markets, where they feed on the offal.

The Stork much resembles the crane in its conformation, but appears somewhat more corpulent. The former lays four eggs, whereas the latter lays but two.

It is said that Storks visit Egypt in such abundance, that the fields and meadows are white with them. The Egyptians, however, are not displeased with the sight; as frogs are there generated in such numbers, that did not the Storks devour them, they would overrun everything. Between Belba and Gaza, the fields of Palestine are often rendered desert on account of the abundance of mice and rats; and were they not destroyed, the inhabitants could have no harvest. The disposition of the Stork is mild and placid; it is easily tamed, and may be trained to reside in gardens, which it will clear of insects and reptiles. It has a grave air, and a mournful aspect; yet, when roused by example, exhibits a certain degree of gaiety; for it joins in the frolics of children, hopping about and playing with them.

During their migrations, Storks are observed in vast quantities. Dr. Shaw saw three flights of them leaving Egypt, and passing over Mount Carmel, each of which appeared to be nearly half-a-mile in width; and he says they were three hours in passing over.

The Stork, like the ibis, was an object of worship among the ancients, and to kill them was a crime punishable with death. The Stork is remarkable for its great affection towards its young, but more especially for its attention to its parents in old age.



THE ADJUTANT (*Leptoptilos argala*)

Is a bird of the stork kind, and a native of India, and other warm countries. Its bill is very different from that of other birds, and is composed of two long sticks, like wood, in order to clean which it makes a loud clattering noise. The head and neck are bare of feathers, like the ostrich; the former looking as if made of wood; the latter of a flesh-colour. The eye seems to be without lids, or any cartilaginous appendages. Under the chin is a kind of bag or purse, that seems adapted to receive the water they are obliged to swallow with the substance they feed on. The coverts of the wings and the back are black, with a bluish cast; the under part of the body whitish; the legs are long, without feathers, and of a greyish hue, as are the thighs, which seem to be as slender as the leg.

THE HERON. (*Ardea Herodias.*)

THE habits of the Heron are peculiar. Perched on a stone, or the stump of a tree, by the solitary current of a brook, his neck and long beak half-buried between his shoulders, he will wait the whole day long, patient and unmoved, for the passing of a small fish, or the hopping of a frog; but his appetite is insatiable.

This bird is about four feet long from the tip of the bill to the end of the claws; to the end of the tail about thirty-eight inches; its breadth, when the wings are extended, is about five feet. The male is distinguished by a crest or tuft of black feathers hanging from the

hinder part of his head, which in chivalrous times was of great value, and held as a peculiar mark of distinction when worn above the plume of ostrich feathers. The back is clothed with down instead of feathers.

Virgil places the Heron among the birds that are affected by and foretell the approaching storm :

“ When watchful Herons leave their watery stand,
And mounting upwards with erected flight,
Gain on the skies, and soar above the sight.”


DRYDEN.

The Heron, though living chiefly in the vicinity of marshes and lakes, forms its nest on the tops of the loftiest trees, or on the pointed cliffs of the sea-shore. It resembles the rook in its habits : a great number of Herons living together in what is called a Heronry, as rooks do in a rookery. The female lays four large eggs, of a pale green colour; and the natural term of this bird's life is said to exceed sixty years.

In England, Herons were formerly ranked among the royal game, and protected as such by the laws ; and when falconry was in fashion, the pursuit of the Heron was a favourite amusement.

—————“ Now like a wearied stag,
That stands at bay, the Hern provokes their rage ;
Close by his languid wing, in downy plumes
Covers his fatal beak, and cautious hides
The well-dissembled fraud. The falcon darts
Like lightning from above, and in her breast
Receives the latent death ; down plumb she falls,
Bounding from earth, and with her trickling gore
Defiles her gaudy plumage. See, alas !
The falconer in despair, his favourite bird
Dead at his feet ; as of his dearest friend,
He weeps her fate ; he meditates revenge,
He storms, he foams, he gives a loose to rage :
Nor wants he long the means ; the Hern fatigued,
Borne down by numbers, yields, and prone on earth
He drops ; his cruel foes wheeling around
Insult at will.”

SOMERVILLE.



THE BITTERN (*Botaurus Stellaris*)

Is not quite so large as the common heron; its head is small, narrow, and compressed at the sides. The crown is black, the throat and sides of the neck red, with narrow black lines, and the back of a pale red, mixed with yellow. The claws are long and slender, the inside of the middle one being serrated, the better to enable it to hold its prey. The bill is about four inches in length. The most remarkable character in this bird is the hollow and yet loud rumbling of his voice; his bellowing is heard at the distance of a mile, at the time of sunset, and it is hardly possible to conceive at first how such a body of sound, resembling the lowing of an ox, can be produced by a bird comparatively so small. The booming


noise was formerly believed to be made while the bird plunged its bill into the mud; hence Thomson,—

—————“ So that scarce
The Bittern knows his time, with bill ingulf'd
To shake the sounding marsh.”

And Southey also describes the peculiar noise of this bird in his poem of Thalaba:

“ And when at evening, o'er the swampy plain,
The Bittern's boom came far,
Distinct in darkness seen—
Above the low horizon's lingering light,
Rose the near ruins of old Babylon.”


Sometimes in the evening the Bittern soars on a sudden in a straight, or, at other times, in a spiral line, so high in the air, that it ceases to be perceptible to the eye. When attacked by the buzzard, or other birds of prey, it defends itself with great courage, and generally beats off such assailants; neither does it betray any symptoms of fear when wounded by the sportsman, but eyes him with a keen undaunted look; and, when driven to extremity, will attack him with the utmost vigour, wounding his legs, or aiming at his eyes with its sharp and piercing bill. It was formerly held in much estimation at the tables of the great, and is again recovering its credit as a fashionable dish. The flesh is considered delicious. In autumn it changes its abode, always commencing its journey at sunset. Its precautions for concealment and security seem directed with great care and circumspection. It usually sits in the reeds with its head erect; and thus, from its great length of neck, sees over their tops, without being itself perceived by the sportsman. The principal food of these birds, during summer, consists of fish and frogs; but in autumn they resort to the woods in pursuit of mice, which they seize with great dexterity, and always swallow whole. About this season they usually become very fat.



• THE SPOONBILL (*Platalea leucorodia*)

Is a large bird; the colour of the whole body is white, and the resemblance of the bill to a spoon has caused the denomination of the bird. In some species the plumage inclines from white to pink colour. On the hind part of the head is a beautiful white crest, reclining backward. The legs and thighs are black. The wisdom of Providence is most conspicuous in the conformation of the bill, which is entirely adapted to the habits and manner of feeding of these birds; the frog and the lizard, which constitute the principal food of the Spoonbill, often escape the thin and narrow beak of the heron and other birds, but the mandibles of this bird are so large at the end, that the prey cannot slip aside. Like rooks and herons, Spoonbills build their nests on the tops of high trees, and lay three or four eggs, the size of those of a hen, which are white, sprinkled with pale red. These birds are very noisy during the breeding season. The Spoonbill migrates northward in the summer; and returns to southern climes on the approach of winter; and is found in all the intermediate low countries between the Faroe Isles and the Cape of Good Hope.

The *American* or *Roseate Spoonbill* is very beautiful. Its colour is white, tinged with rose, which deepens in the wings and tail into the richest carmine. The feet are half-webbed, and the bird is generally found on the sea-coast, where it wades into the sea in quest of the small shell-fish of different kinds, on which it feeds.



THE CURLEW. (*Numenius arquata*.)

“ Soothed by the murmurs of the sea-beat shore,
His dun-gray plumage floating to the gale,
The Curlew blends his melancholy wail
With those hoarse sounds the rushing waters pour.”

MISS WILLIAMS.

“ Wild as the scream of the Curlew,
From rock to rock the signal flew.” SIR WALTER SCOTT.

THE Curlew is a large bird, weighing about twenty-four ounces; and is found on the sea-shore on all sides of England. The middle parts of the feathers of the head, neck, and back are black, the borders or outsides ash-coloured, with a mixture of red; and the lower part of the body white. The beak has a regular curve downward, and is soft at the point. This bird's flesh may challenge for flavour and delicacy that of any other water-fowl, and the people of Suffolk say proverbially:


“ A Curlew, be she white, be she black,
She carries twelve pence on her back:”

but it must be confessed that the quality and goodness of the flesh of Curlews depend on their manner of feeding, and the season in which they are caught. When they dwell on the sea-shore, they acquire a kind of rankness, which is so strong, that, unless they are basted on the spit with vinegar, they are not agreeable eating.



THE REDSHANK. (*Scolopax*, or *Totanus calidris*.)

THIS bird has received its name from the colour of its legs, which are of a crimson red. In size it is between the lapwing and the snipe, and is sometimes called the *Pool Snipe*. The head and back are of a dusky ash-colour, spotted with black, the throat party-coloured black and white, the black being drawn down along the feathers. The breast is whiter, with fewer spots. The Redshank delights in the fen countries, and in wet and marshy grounds, where it breeds and rears its young. The female lays four whitish eggs, with olive-coloured dashes, and marked with irregular spots of black. Pennant and Latham say, that it flies round its nest when disturbed, making a noise like a lapwing. It is not so common on the sea-shore as several others of its kindred. We must here observe, that this bird has often been mistaken for others. The fact is, that several birds changing their plumage, and increasing or diminishing their size according to their age, the season of the year, and the climate they live in, set all nomenclators at defiance, and confound all classifications.



THE GODWIT (*Limosa algocephala*)

Is met with in various parts of Great Britain, and is rather larger than the woodcock, which it much resembles in appearance. In spring and summer it resides in the fens and marshes, where it rears its young, and feeds on small worms and insects; but in winter it seeks the salt marshes and the sea-shore, where it feeds upon the shell-fish and marine insects left by the retiring tide. A peculiarity belonging to this bird is the shape of its bill, which is a little turned upwards. The head, neck, and back are of a reddish brown; the under part of the body white; and the legs dusky, and sometimes black.

The Godwit is much esteemed by epicures as a great delicacy, and sells very high. It is caught in nets, to which it is allured by a *stale* or stuffed bird, in the same manner and in the same season as the ruffs and reeves.



THE SNIPE. (*Scolopax gallinago*.)

“ The Snipe flies screaming from the marshy verge,
And towers in airy circles o’er the wood.

Still heard at intervals ; and oft returns,
And stoops as bent to alight ; then wheels aloft
With sudden fear, and screams and stoops again,
Her favourite glade reluctant to forsake." GIBBORNE.

THE Snipe weighs about four ounces. A pale red line divides the head longways ; the chin under the bill is white ; the neck is a mixture of brown and red ; the lower part of the body is almost all white. The back and wings are of a dusky colour. The flesh is tender, sweet, and in flavour next to that of the woodcock. Snipes feed especially upon small red worms, and insects, which they find in muddy and swampy places, on the banks of rivulets and brooks, and on the clayish margin of ponds. It is said that some Snipes remain with us all the summer, and build in moors and marshes, laying four or five eggs. The others are migratory, and when forced by severe frosts to sheltered springs, are often seen in large flights. Mr. Daniel states, that, about thirty years ago, Snipes were so abundant in the fens of Cambridgeshire, that as many were taken in Milton fen, by means of a lark-net, in one night, and by a single man, as could be contained in a small hamper.






THE WOODCOCK (*Scolopax rusticola*)

Is somewhat less than the partridge. The upper side of the body is partly-coloured of red, black, and grey, and very beautiful. From the bill almost to the middle of the head it is of a reddish ash-colour. The lower part of the body is grey, with transverse brown lines; under the tail the colour is somewhat yellowish; the chin is white, with a tincture of yellow. Woodcocks are migratory birds, coming over into Britain in autumn, and departing again in the beginning of spring; they pair before they go, and are seen flying in braces.

The colours of this timid bird render it difficult to discern him among the withered stalks and leaves of fern, sticks, moss, and grass, which form the back-ground of the scenery, by which he is sheltered in his moist and solitary retreats. By habit only is the sportsman enabled to discover him, and his leading marks are the full eye and glossy silver white-tipped tail of the bird. The flesh is held in high estimation, and hence he is eagerly sought after. It is hardly necessary to observe that in dressing a Woodcock for the spit the entrails are not drawn, but are allowed to drop upon slices of toasted bread, and are relished as a delicious kind of sauce. By some late observations, it appears that several individuals of the species remain with us the whole year. They frequent especially wet and swampy woods, the thick hedges near rivulets, and places affording them their allotted food, which consists of very small insects found in the moist ground.

“ The Woodcock’s early visit and abode
Of long continuance, in our temperate clime,
Foretell a liberal harvest.”

PHILIPS.



THE LAPWING, OR PEEWIT.

(*Vanellus cristatus.*)

THIS well-known bird is found in nearly all countries, and is of the size of a common pigeon. The female lays four or five eggs, of a yellow colour, varied all over with large black spots and strokes. Lapwings build their nests on the ground in the middle of some field or heath, open and exposed to view, laying only some few straws under the eggs: as soon as the young are hatched, they instantly forsake the nest, running away with the shell on their back, and following the mother, only covered with a kind of down, like young ducks. The parents have been impressed by nature with the most attentive love and care for their offspring; for if the fowler, or any other enemy, should come near the nest, the female, panting with fear, lessens her call to make her enemies believe that she is much further off, and thereby deceives those that search for her brood; she also sometimes pretends to be wounded, and utters a faint cry as she limps away, to lead the fowler from her nest. This bird is really beautiful, although it does not exhibit that gaudiness of colours of which other species of the feathered tribe can boast: it weighs about half-a-pound. The head and the crest which elegantly adorns it are black; this crest, composed of unwebbed feathers, is about four inches in length. The back is of a dark green, glossed with blue shades; the throat is black; the hinder part of the neck and the breast are white. The Lapwing, when

in search of food, stamps with his feet upon the ground, and when the earth-worms, alarmed at the noise, appear he seizes and devours them. His voice, on the swampy places along the sea-shores, heard at night, resembles the sound of *peewit*, or *teewit*, and hence his name in several parts of Great Britain; he is also called the *Great Plover* by several ornithologists. This bird is one of those who attract the fowler's attention in winter.

“ With slaughtering gun th' unwearied fowler roves,
 When frosts have whiten'd all the naked groves ;
 Where doves in flocks the leafless trees o'ershade,
 And lonely woodcocks haunt the watery glade.
 He lifts his tube, and levels with his eye ;
 Straight a short thunder breaks the frozen sky ;
 Oft, as in airy rings they skim the heath,
 The clamorous Lapwings feel the leaden death :
 Oft, as the mounting larks their notes prepare,
 They fall, and leave their little lives in air.” POPE.

The following anecdote, from Bewick's History of Birds, exhibits the domestic nature of the Lapwing, as well as the art with which it conciliates the regard of animals materially differing from itself, and generally considered as hostile to every species of the feathered tribe. Two Lapwings were given to a clergyman, who put them into his garden ; one of them soon died, but the other continued to pick up such food as the place afforded, till winter deprived it of its usual supply. Necessity soon compelled it to draw nearer to the house, by which it gradually became familiarized to occasional interruptions from the family. At length one of the servants, when she had occasion to go into the back-kitchen with a light, observed that the Lapwing always uttered his cry of “*pee-wit*,” to obtain admittance. The bird soon grew more familiar : as the winter advanced, he approached as far as the kitchen, but with much caution, as that part of the house was generally occupied by a dog and cat, whose friendship, however, the Lapwing at length conciliated so entirely, that it was his regular custom to resort to the fire-side as soon as it grew dark, and spend the evening

and night with his two associates, sitting close by them, and partaking of the comforts of a warm hearth. As soon as spring appeared, he discontinued his visits to the house, and betook himself to the garden; but, on the approach of winter, he had recourse to his old shelter and friends, who received him very cordially. Security was productive of insolence; what was at first obtained with caution, was afterwards taken without reserve: he frequently amused himself with washing in the bowl which was set for the dog to drink out of; and while he was thus employed, he showed marks of the greatest indignation if either of his companions presumed to interrupt him. He died in the asylum he had thus chosen, being choked with something that he had picked up from the floor.



THE RUFF AND REEVE. (*Machetes pugnax*.)

It is curious to see, in our observation of natural objects, how the creative power of Providence seems to have tried all forms and shapes in the composition of species. In the cock bird of this species a circle or collar of long feathers, somewhat resembling a ruff, encompasses the neck under the head, whence the bird has received the name of Ruff. It is about a foot in length, with a bill about an inch long. There is a wonderful and almost infinite variety in the colours of the feathers of the males;

so that in spring there can scarcely be found two exactly alike, but after moulting they become all alike again. The hens are smaller than the cocks, and their feathers undergo no change. The female is called a Reeve, and its flesh is thought a great delicacy for the table. The males are very fond of fighting, and they attack each other with such fury, that the bird has received the name of *pugnax*, which signifies fighting. It is a bird of passage, and arrives in the fens of Lincolnshire, and other similar places, in the spring. Mr. Pennant tells us, that in the course of a single morning more than six dozen have been caught in one net; and that a fowler has been known to catch between forty and fifty dozen in a season.



THE KNOT (*Tringa Canutus*)

Is a small bird, whose head and back are of a dusky ash-colour, or dark grey; while the lower part of the body is pure white, or white varied by black lines. The sides under the wings are spotted with brown. The bird weighs about four ounces and a half, and generally makes its appearance in Lincolnshire in the beginning of winter, and abides there for two or three months, after which they fly off in flocks. They are caught in great numbers by nets, into which they are decoyed by carved wooden figures, painted to represent themselves, and placed within them, much in the same way as the ruff. When the Knot is fat, its flesh is considered excellent food. It is also fattened for sale, and then considered equal to the ruff in

flavour. The season for taking it is from August to November, after which the frost compels it to disappear. —This bird is said to have been a favourite dish with Canute the Great; and Camden observes that its name is derived from his—Knut, or Knout, as he was called— which, in process of time, has been changed to Knot.




THE GREY PLOVER (*Tringa squatarola*)

Is about the size of the lapwing: the head, back, and coverts of the wings are black, with tips of a greenish grey; the chin white; the throat spotted with brown or dusky spots; the breast and thighs white. The flavour of the flesh, when the bird is caught in the proper season, is delicate and savoury; at other times it is hard, and has a strong and rank flavour.

The *Green Plover* is about the size of the former. The colour of the whole upper side is black, thick set with yellowish green spots; the breast brown, with spots as on the back; the body is white. The flesh is sweet and tender, and therefore esteemed a choice dish, in this and other countries.

The "Plover eggs," frequently seen at the tables of the opulent and luxurious, are not those of the Plover, but of the Lapwing.



THE DOTTREL. (*Charadrius Morinellus*)

Is proverbially accounted a foolish bird, yet why so it is hardly possible to say. Its length is about ten inches; the bill is not quite an inch long, and is black. The forehead is mottled with brown and grey; the top of the head is black; and over each eye there is an arched line of white. The back and wings are a light brown; the breast is a pale dull orange; the middle of the body is black, and the rest and the thighs are of a reddish white. The tail is brown, black towards the end, and tipped with white. This bird seems to be migratory, and makes its appearance in Lincolnshire, Cambridgeshire, and Derbyshire in April, but soon leaves those counties and retires in June to places which are totally unknown. In April, and sometimes in September, Dottrels are seen in Wiltshire and Berkshire. They are generally caught, like other birds, by night; when, dazzled by the light of a torch, they are at a loss to know where to fly for safety, the whole place being in darkness, and generally select the very spot which they should avoid. Many ridiculous stories have been propagated about the gestures of this bird, and its endeavouring to imitate the actions of the fowler, and thereby falling into the snare laid for him; but they ought to be entirely exploded.




THE WATER-HEN (*Gallinula chloropus*)

Is larger than the Plover, and is also called the *Moor-Hen*, or *Moor-Coot*, and the *Gallinule*. The breast is of a lead-colour, the lower part of the body inclining to ash-colour, and the back blackish. As she swims or walks, she often flirts up her tail. Water-hens feed upon watery grass and roots, and upon the small insects which adhere to them: they grow fat about the latter end of September, and their flesh is then considered nearly equal to that of the teal; yet it can seldom be entirely deprived of its fishy taste. They build their nests upon low trees and shrubs by the water-side, breeding twice or thrice in the course of a summer; the eggs are white, with a tint of green, dashed with brown spots.

There are very few countries in the world where these birds are not to be found. They generally prefer the cold mountainous regions in summer, and lower and warmer situations during winter.


“The fish are leaping, and the water-hen
Dives up and down. A storm is coming on.”

SCHILLER. WILLIAM TELL.




THE CORN-CRAKE, OR LAND-RAIL,
(*Ortygometra crex*)

Is a migratory bird, appearing in England in April, and departing in October. At the time of its arrival it is very lean, but becomes excessively fat before it quits the island. Their favourite haunts are cold and humid upland-districts, corn-fields in the vicinity of water, and marshy grass-lands. Their cry is a peculiar roll of short notes, all in the same key and of the same length. The sound, crecq, crecq, crecq, has been compared to the noise made by drawing the finger along the teeth of a comb. The legs of the Corn-crake are unusually long for the size of the bird, and hang down while it is on the wing. Its flesh is greatly esteemed for its delicate flavour. This bird is never seen on the wing in this country, and is extremely difficult to capture; they cannot be made to rise like partridges and many other birds, nor is it of much use to invade their cover. They glide through the corn, without the least perceptible rustle, and if the sportsman follows in the direction of the sound, it ceases for a while, and then, perhaps, is heard far in the rear; if he follows it again, it is not long before the sound is heard setting in its former or some other direction.

THE COOT. (*Fulica atra.*)

THIS bird has so many traits in its character, and so many features in its general appearance, like the rails and water-hens, that to place it after them seems a natural and easy gradation; and accordingly this has been done by Cuvier, though it was considered by Linnæus to be a genus distinct from those birds, and from the waders in general, on account of its being fin-footed, and its constant attachment to the waters, which, indeed, it seldom quits. The manner in which Coots build their nest is very ingenious. They form it of interwoven aquatic weeds, and place it among the rushes, in such a way that it may occasionally rise with, but not be washed away by the stream; and if ever this accident happens, steady on her nest the hen does not desert her brood, but follows with them the destiny of their floating cradle. This bird, in the figure and shape of its body, resembles the water-hen, and weighs about twenty-four ounces. The feathers about the head and neck are low, soft, and thick. The colour about the whole of the body is black, but of a deeper hue about the head. The cere rises upon the forehead in a peculiar manner, and appears as if Providence had designed it for a means of defence. It changes its whitish colour to a pale red or pink in the breeding season. Coots are very shy, and seldom venture abroad before dusk. When attacked, they defend themselves with their feet, and they do this so energetically, that sportsmen say, "Beware of a winged Coot, or he will scratch you like a cat."

SECTION IV.—Palmipedes, or Web-footed Birds.

THE PELICAN (*Pelecanus onocrotalus*)

Is in size nearly equal to the swan; the colour of the body is white, inclining to pink; the beak is straight and long, with a sharp hook at the end; the gullet, or skin of the lower mandible, is so capable of distention, that it may be dilated to contain fish in large quantities, and some say fifteen quarts of water. This pouch Providence has allotted to the bird, that he may bring to his eyrie sufficient food for several days, and save himself the trouble of travelling through the air, and watching and diving so often. The legs are black, and the four toes palmated. It is a very indolent, inactive, and inelegant bird, often sitting whole days and nights on rocks or branches of trees, motionless and in a melancholy posture, till the resistless stimulus of hunger spurs it on, and forces it to the sea in search of nourishment: when thus incited to exertion, the Pelican flies from the spot, and,

raising itself thirty or forty feet above the surface of the water, turns its head with one eye downward, and continues to fly in that position till it sees a fish near the surface. It then darts down with astonishing swiftness, seizes its prey with unerring certainty, and stores it in its pouch. Having done this, it rises into the air, and repeats the same action till it has procured a sufficient stock. The Pelican is one of the most heavy, sluggish, and voracious of all the feathered tribes, and is but ill fitted to take those vast flights, or to make those cautious provisions which have been ascribed to it. It is, however, by no means destitute of natural affection, either towards its young ones, or towards others of its own species. Clavigero, in his History of Mexico, says, that sometimes the Americans, in order to procure, without trouble, a supply of fish, cruelly break the wing of a live Pelican, and, after tying the bird to a tree, conceal themselves near the place. The screams of the miserable bird attract other Pelicans to the place, which, he assures us, eject a portion of the provisions from their pouches for their imprisoned companion. As soon as the men observe this, they rush to the spot, and, after leaving a small quantity for the bird, carry off the remainder.

In America, Pelicans are often rendered domestic, and are so trained, that at command they go in the morning and return before night with their pouches distended with prey, part of which they are made to disgorge, while the rest is left them for their trouble. The bird is said to live sometimes a hundred years.

Our forefathers attributed extraordinary affection to this bird, more than is attested by any save heraldic evidence. Thus, in several crests, it is represented in the act of feeding its young with its own blood, which it procures by striking its breast with the sharp point of its beak.



THE CORMORANT (*Phalacrocorax carbo*)

Is a large water-bird, nearly allied to the pelican, possessed with a very voracious appetite, and consequently of a very rapacious disposition. It lives upon all sorts of fish; the fresh water, and the briny waves of the sea both paying a large contribution to its craving stomach. The bill is about five inches in length, and of a dusky colour; the predominant tints of the body are black and dark green. The smell of these birds when alive is excessively rank and disagreeable; and their flesh is so disgusting, that even the Greenlanders, among whom they are very common, will scarcely eat it. They were formerly tamed in England for the purpose of catching fish, as falcons and hawks were for chasing the fleet inhabitants of the air. We are told that the custom is still in practice in China. This bird, although of the aquatic kind, is often seen, like the pelican, perched upon trees. Milton tells us that Satan

“ _____ on the tree of life,
The middle tree, and highest there that grew,
Sat like a Cormorant.”

In the year 1793, one of them was observed sitting on the vane of St. Martin's steeple, Ludgate Hill, London, and was shot there in the presence of a great number of people.

The crested Cormorant (*Phalacrocorax cristatus*) is of a dark green, with a singular tuft on the front of the head. It breeds in rocky caves on the sea-coast.



THE GANNET, OR SOLAN GOOSE.

(*Sula Bassana.*)

THESE birds are insatiably voracious, but are somewhat particular in their choice of prey; disdaining, unless in great want, any food worse than herrings or mackerel. No fewer than one hundred thousand Gannets are supposed to frequent the rocks of St. Kilda; and of these, including the young ones, at least twenty thousand are annually killed for food by the inhabitants. The Gannet is somewhat more than three feet in length, and weighs about seven pounds. The bill is six inches long, straight almost to the point, where it is a little bent; its edges are

irregularly jagged, to enable it the better to secure its prey; and about an inch from the base of the upper mandible there is a sharp process pointing forward. The general colour of the plumage is dingy white, with a greyish tinge. Surrounding each eye there is a naked skin of a fine blue colour; from the corner of the mouth a narrow slip of naked black skin extends to the hind part of the head; and beneath the chin there is a pouch capable of containing five or six herrings. The neck is long; the body flat, and very full of feathers. On the crown of the head, and the back part of the neck, is a small buff-coloured space. The quill-feathers, and some other parts of the wings, are black; as are also the legs, except a fine pea-green stripe in front. The tail is wedge-shaped, and consists of twelve sharp-pointed feathers.

These birds chiefly resort to those uninhabited islands where their food is found in plenty, and man seldom comes to disturb them. The Islands to the north, Ailsa Craig, on the west coast of Scotland, the Skelig Islands off the coasts of Kerry in Ireland, and those that lie in the North Sea off Norway, abound with them. But it is on the Bass Rock, in the Frith of Forth, that they are seen in the greatest abundance. "There is a small island," says the celebrated Harvey, "called the Bass, not more than a mile in circumference: the surface is almost wholly covered during the months of May and June with the nests of the Solan Geese, their eggs, and their young. It is scarcely possible to walk without treading on them: the flocks of birds upon the wing are so numerous as to darken the air like a cloud; and their noise is such, that one cannot without difficulty be heard by the person next to him. When one looks down upon the sea from the precipice, its whole surface seems covered with infinite numbers of birds of different kinds, swimming and pursuing their prey. If, in sailing round the island, one surveys its hanging cliffs, in every craig or fissure of the broken rocks may be seen innumerable birds, of various sorts and sizes, more than the stars of heaven when viewed in a serene night. If they are viewed at a distance, either receding, or in their approach to the island, they seem like one vast swarm of bees."



THE SWAN. (*Cygnus ferus and olor.*)

" Fair is the Swan, whose majesty prevailing
 O'er breezeless water, on Locarno's lake,
 Bears him on, while, proudly sailing,
 He leaves behind a moon-illumined wake :
 Behold ! the mantling spirit of reserve
 Fashions his neck into a goodly curve—
 An arch thrown back between luxuriant wings
 Of whitest garniture, like fir-tree boughs,
 To which, on some unruffled morning, clings
 A flaky weight of winter's purest snows !
 Behold ! as with a gushing impulse heaves
 That downy prow, and softly cleaves
 The mirror of the crystal flood,
 Vanish inverted hill, and shadowy wood,
 And pendent rocks, where'er in gliding state
 Winds the mute creature, without visible mate
 Or rival, save the queen of night,
 Showering down a silver light
 From heaven upon her chosen favourite ! "

WORDSWORTH.

THERE are two distinct species of this elegantly formed and majestic bird, the wild and the mute, or tame; both bearing the general characters of the genus, yet not exactly tallying with each other. The beak of the wild Swan is surmounted with a yellow skin, which runs up to the eye. In the tame one this appendage (the use of which has not been yet sufficiently explored) is jet black, as are the feet in both species. The tame Swan is the largest of all web-footed water-fowl, sometimes weighing about twenty pounds: the whole body of the full-grown Swan is covered with a beautiful pure white plumage, but the young ones are grey; under the feathers is a thick but soft down, which is of very great use, and often employed as an ornament. The elegance of form which this bird displays, when, with his arched neck and half-displayed wings, he sails along the crystal surface of a tranquil stream, which reflects, as he passes, the snowy beauty of his dress, is worthy of admiration. Thomson describes the Swan in the following beautiful manner:

“ ——— the stately sailing Swan
Gives out his snowy plumage to the gale,
And arching proud his neck, with oary feet,
Bears forward fierce, and guards his osier isle,
Protective of his young.”


Swans have for ages been protected on the river Thames as royal property; and it continues at this day to be accounted felony to steal their eggs; by this means their increase is secured, and they prove a delightful ornament to that noble river. Latham says the estimation in which they were held, in the reign of Edward IV., was such, that no one who possessed a freehold of less than the clear yearly value of five marks, was permitted even to keep any. In those times, hardly a piece of water was left unoccupied by these birds, as well on account of the gratification they gave to the eye of their lordly owners, as that which they also afforded when they graced the sumptuous board at the splendid feasts of that period: but the fashion of those days has passed away, and Swans are by no means so common now as

they were formerly, being by most people accounted a coarse kind of food, and consequently held in little estimation: but the Cygnets (so the young Swans are called) are still fattened for the table, and are sold very high, commonly for a guinea each, and sometimes more; hence it may be presumed they are better food than is generally imagined.

At Abbotsbury there was formerly a noble Swannery, the property of the Earl of Ilchester, where six or seven hundred birds were kept; but the collection has of late been much diminished. The Swannery belonged anciently to the abbot, and, previously to the dissolution of monasteries, the Swans frequently amounted to double the above number.

From the whiteness of this bird, the expression of a "Black Swan" was used in ancient times as equivalent to a nonentity; but a species nearly entirely black has been lately discovered in Australia. This bird is larger than the white Swan, and its bill of a rich scarlet. The whole plumage (except the primaries and secondaries, which are white) is of the most intense black.

Swans are very long lived, sometimes attaining the great age of a century and a half.



THE WILD GOOSE. (*Anser ferus.*)

"The farmer's Goose, who in the stubble
Has fed without restraint or trouble,

Grown fat with corn, and sitting still,
 Can scarce get o'er the barn-door sill;
 And hardly waddles forth to cool
 Her body in the neighbouring pool;
 Nor loudly cackles at the door,
 For cackling shows the Goose is poor."

SWIFT.

THE Goose is very different in outward appearance from the last-named bird. Stupidity in her look, uncouthness in her walk, and heaviness in her flight, are her principal characteristics. But why should we dwell upon these defects? they are not such in the great scale of the creation. Her flesh feeds many, and is not disdained even by the great; her feathers keep us warm; and the very pen I hold in my hand was plucked from her wing.

These birds are kept in vast quantities in the fens of Lincolnshire; several persons there having as many as a thousand breeders. They breed in general only once a year, but if well kept they sometimes hatch twice in a season. During their sitting, the birds have spaces allotted to each, in rows of wicker pens placed one above another; and the Goose-herd, who has the care of them, drives the whole flock to water twice a day, and bringing them back to their habitations, places every bird (without missing one) in its own nest. It is scarcely credible what numbers of Geese are driven from the distant counties to London for sale, frequently two or three thousand in a drove; and, in the year 1783, one drove passed through Chelmsford, in its way from Suffolk to London, that contained more than nine thousand. However simple in appearance or awkward in gesture the Goose may be, it is not without many marks of sentiment and understanding. The courage with which it protects its offspring and defends itself against ravenous birds, and certain instances of attachment and even of gratitude, which have been observed in it, render our general contempt of the Goose ill-founded.

The Goose was held in great veneration among the Romans, as having by her watchfulness saved the Capitol


from the attack of the Gauls. Virgil says, in the seventh book of the *Æneid*,

“ The silver Goose before the shining gate
There flew, and by her cackle saved the state.”

DRYDEN.

The colour of this useful bird is generally white; though we often find them of a mixture of white, grey, black, and sometimes yellow. The feet, which are palmated, are orange-coloured, and the beak is serrated. The male of the Goose is called the Gander; and the young ones Goslings. Geese are very long-lived, one is known to have lived above seventy years.


The *Wild Goose* is, of course, the original of the tame one, and differs much in colour from her, the general tint of its feathers being a greyish black. Wild Geese fly by night in large flocks to more southern countries; and their clang is heard from the regions of the clouds, although the birds are out of sight.



THE DUCK. (*Anas boschas.*)


THE common Duck is of two kinds, the wild and the tame, the latter being but the same species altered by domestication; the difference between them is very trifling, save that the colour of the Mallard, or male wild Duck, is constantly the same in all the individuals, whereas the Drakes, or tame ones, are varied in their plumage. The females do not share with the males in beauty of plumage: the admirable scarf of glossy green and blue, which sur-

rounds the neck of Drakes and Mallards, being an exclusive prerogative of the male sex. There is also a curious and invariable peculiarity belonging to the males, which consists of a few curled feathers rising upon the rump.



Wild Ducks are caught by decoys in the fen countries, and in such prodigious numbers, that in only ten decoys in the neighbourhood of Wainfleet, as many as thirty-one thousand two hundred have been caught in one season. They do not always build their nests close to the water, but often at a considerable distance from it; in which case the female will take the young ones in her beak, or between her legs to the water. They have sometimes been known to lay their eggs in a high tree, in a deserted magpie's or crow's nest; and an instance has been recorded of one being found at Etchingam, in Sussex, sitting upon nine eggs in an oak, at the height of twenty-five feet from the ground: the eggs were supported by some small twigs laid cross ways.

The tame Ducks, reared about mills and rivers, or wherever there is a sufficient quantity of water for them to indulge their sports and to search for food, become a branch of trade, which proves very profitable to their owners.



THE WIGEON (*Anas Penelope*)

WEIGHS about twenty-two ounces, and feeds upon grass and roots growing at the bottom of lakes, rivers, and ponds. The plumage of this bird is much variegated, and its flesh esteemed a great delicacy, though not so highly praised as that of the teal. The bill of the Wigeon is black; the head and upper part of the neck of a bright bay; the back and sides under the wings waved with black and white; the breast purple, the lower part of the body white; and the legs are dusky. The young of both sexes are grey, and continue in this plain garb till the month of February; after which a change takes place, and the plumage of the male begins to assume its rich colourings, in which, it is said, he continues till the end of July; and then again the feathers become dark and grey, so that he is hardly to be distinguished from the female.

Wigeons commonly fly in small flocks during the night, and may be known from other birds by their whistling note, while they are on the wing. They quit the desert morasses of the north on the approach of winter, and as they advance towards the end of their destined southern journey, they spread themselves along the shores, and over the marshes and lakes, in various parts of the continent, as well as those of the British isles; and it is said that some of the flocks advance as far south as Egypt.


The Wigeon is easily domesticated in places where there is plenty of water, and is much admired for its

beauty, sprightly look, and busy frolicsome manners: yet it is generally asserted that they will not breed in confinement, or at least that the female will not make a nest and perform the act of incubation; but that she will lay eggs, which are generally dropped into the water.



THE TEAL (*Querquedula crecca*)

Is the least of the duck tribe, weighing only twelve ounces. The lower part of the body is of a dingy white, inclining to a grey tint. The back and sides under the wings are curiously varied with lines of white and black; the wings are all over brown, and the tail of the same colour. This bird is common in England during the winter months, and it is still uncertain whether it does not breed here as it does in France. Dr. Heysham says, it is known to breed in the neighbourhood of Carlisle. The female makes her nest of reeds interwoven with grass; and, as it is reported, places it among the rushes, in order that it may rise and fall with the water. Their eggs are of the size of those of a pigeon, six or seven in number, and of a dull white colour, marked with small brownish spots; but it appears that they sometimes lay ten or twelve eggs, for Buffon remarks, that that number of young are seen in clusters on the pools, feeding on cresses, chervil, and some other weeds, as well as upon seeds and small insects that swarm in the water. The flesh of the Teal is a great delicacy in the winter season, and has less of the fishy flavour than any of the wild duck kind. It is known to breed and remain throughout the year in various temperate climates of the world, and is in the summer met with as far northward as Iceland.



THE STORMY PETREL, OR MOTHER CARY'S
CHICKEN. (*Thalassidroma pelagica.*)

“ O'er the deep ! o'er the deep,
Where the whale, and the shark, and the sword-fish sleep,
Outflying the blast and the driving rain,
The Petrel telleth her tale in vain ;
For the mariner curseth the warning bird,
Who bringeth him news of the storm unheard !
Oh ! thus does the prophet, of good or ill,
Meet hate from creatures he serveth still ;
Yet he ne'er falters :—So, Petrel ! spring
Once more o'er the waves on thy stormy wing !” PROCTOR.

THE Stormy Petrel is not larger than a swallow ; and its colour is entirely black, except the coverts of the tail, the tail itself, and the vent-feathers, which are white : its legs are long and slender. Ranging over the expanse of the ocean, and frequently at a vast distance from the land, this bird is able to brave the utmost fury of the storms. Even in the most tempestuous weather it is frequently observed by the mariners skimming with almost incredible velocity along the billows, and sometimes over their summits. They often follow vessels in great flocks, to pick up anything that is thrown overboard ; but its appearance is looked upon by the sailors as the sure presage of stormy weather in the course of a few hours. It seems to seek protection from the fury of the wind in the wake of the vessels ; and it is probably that for the same reason it often flies between two surges. The nest of this bird is found in the Orkney Islands, under loose stones, in the months of June and July. It lives chiefly

on small fish; and although mute by day, it is very clamorous by night.

The length of the Norfolk Island Petrel is about sixteen inches. The bill is about an inch and a half long, black, and much hooked at the end. The head as far as the eyes, and the chin, are mottled, in waves of brown and white: the rest of the body is of a sooty brown above, and a deep ash-colour beneath. The wings, when closed, exceed the tail by an inch. The legs are of a pale yellow; and part of the toes and webs are black. This variety of the Petrel burrows in the sand like rabbits.

Mudie, in his very entertaining work on British Birds, says that they are called Petrels, or "little Petrels," because they move along the surface as if they were literally walking on the water. He also informs us that they are at times very full of oil, and that the Faroese, taking advantage of this circumstance, convert them into lamps, by fixing them in an upright position and drawing a wick through their bodies, which they light at the mouth.



THE AUK, OR NORTHERN PENGUIN

(*Alca impennis*)

Is considered, by some observers of the wonderful operations of the Creator, a link between the bird and the fish; and we place its description here, in order to exhibit more explicitly and conspicuously the uninterrupted chain of beings which constitute the whole sum of created nature. These birds have very short wings, not exceeding four

inches and a quarter from the tip of the longest quill-feather to the first joint; with short black legs, placed near the vent; and from the awkwardness with which they fly or walk, they are seldom seen out of the water, and rarely venture beyond soundings. They breed in caverns and rocky cliffs, and the female lays one egg only, which she deposits and hatches upon a ledge close to the sea-mark: it is about six inches in length, white streaked with purple lines. When on shore, this bird and all of the same species, stand upright on their rump and short legs, which gives them a very awkward appearance.

Mr. Pennant has asserted that the affection of these birds for their young is so great, that when laid hold of by the wings, they will give themselves the most cruel bites on any part of their body that they can reach, as if actuated by despair; and that, when released, instead of flying away, they will often hurry again into their burrows. Bingley says, "When I was in Wales, in the summer of 1801, I took several Puffin Auks out of the holes that had young ones in them, for the purpose of ascertaining this fact. They bit me with great violence, but none of them seized on any parts of their own body: a few, on being released, ran into the burrows, but not always into those from which I had taken them. If it was more easy for them to escape into a hole than raise themselves into the air, they did so; but if not, they ran down the slope of the hill in which their burrows were formed, and flew away. The noise they make when with their young is a singular kind of humming, much resembling that produced by the large wheels used for the spinning of worsted. On being seized, they emitted this noise with greater violence: and from its being interrupted by their struggling to escape, it sounded not much unlike the efforts of a dumb man to speak. The young ones are entirely covered with a long blackish down, and in shape, are altogether so different from the parent birds, that no one would at first sight suppose them of the same species. Their bill also is long, pointed, and black, with scarcely any marks of furrows."

BOOK III.
INHABITANTS OF THE WATER.
§ 1. CETACEA, or *Sea Mammalia*.



THE WHITE WHALE. (*Beluga leucas*.)

“ Nature’s strange work, vast Whales of different form,
Toss up the troubled flood, and are themselves a storm ;
Uncouth the sight, when they in dreadful play,
Discharge their nostrils, and refund a sea ;
Or angry lash the foam with hideous sound,
And scatter all the watery dust around :
Fearless, the fierce destructive monsters roll,
Ingulf the fish, and drive the flying shoal ;
In deepest seas these living isles appear,
And deepest seas can scarce their pressure bear ;
Their bulk would more than fill the shelvy strait,
And fathom’d depths would yield beneath their weight.”


THE Whale is not properly a fish ; since, though it lives in the sea, and has fins and a tail instead of legs and feet,


it resembles in most other respects a seal, and differs from fishes, properly so called, in many important points. Indeed, it is always included in the class *Mammalia*, by zoologists, as it brings forth its young alive, and nourishes them with its milk; and hence a conceited person, who said he knew every fish from the shrimp to the Whale, was justly laughed at, as neither the Whale nor the shrimp are included in the fishes by zoologists.

The general form of the Whale's body is that of a fish; but the tail is placed horizontally instead of vertically, and the skeleton of the fins exactly resembles that of a hand affixed to a contracted arm, though it is covered with so thick a skin that no trace of the formation of the bones can be discovered externally. There are only two fins, which are very small, and close to the head. The Whale, however, differs from fishes most materially in its having warm blood; and in its lungs, which are exactly the same as those of quadrupeds. Hence, though the Whale can remain a long time under water without breathing, it is compelled to come to the surface whenever it does breathe, and for this purpose it is furnished with two large nostrils, or blow-holes as they are called. The blow-holes are most beautifully and curiously contrived to close when the animal sinks under water; so that not a drop of water can enter the lungs, however great the pressure may be. The Whale is also provided with a very thick skin, containing an immense quantity of liquid oil, called the blubber, which is so easily detached from the flesh, that when a Whale is killed, the blubber, which is sometimes two feet thick, is taken off by passing a common spade between it and the body. This thick oily skin is a non-conductor of heat, and is thus admirably adapted for preventing the warm blood of the Whale from being chilled by the cold of the water. The true fishes, which are unprovided with such a covering, have cold blood, and are therefore not susceptible of chills.

The White Whale, or Beluga, was formerly included in the dolphins, and is still considered by some to belong to that family. The body is white, tinged with yellow, or

rose-colour, and its proportions are more agreeable than those of most of the cetacea. White Whales are gregarious, assembling in flocks or herds, and playing about with rapid and graceful movements. The female has two young ones at a time, over which she watches with the greatest apparent affection. They follow all her movements, and do not quit her till they are nearly full grown. This Whale is generally confined to the northern latitudes, though one was taken in the Firth of Forth in 1815. The oil is of excellent quality, and the flesh eats like beef. According to some writers the flesh, when pickled with vinegar and salt, is as well tasted as pork; and thus the body, which is generally thrown away when the sailors have cut off the blubber, might be used by them as food. The internal membranes are used by the Greenlanders for windows, and the sinews for thread, and the fins and tail, when properly prepared, are said by some of the old writers to be good eating.





THE COMMON OR GREENLAND WHALE.

(*Balæna mysticetus.*)

THESE creatures have no teeth in either jaw, but their mouths are furnished with a kind of fringe of numerous long horny laminae, which are what we call whalebone, and which form a kind of strainer, admitting only the small fish on which the Whale feeds. This Whalebone is one of the valuable products of the white Whale, though the oil is most important.

“ As when enclosing harpooners assail,
 In hyperborean seas, the slumbering Whale;
 Soon as the javelins pierce the scaly side,
 He groans, he darts impetuous down the tide;
 And, rack'd all o'er with lacerating pain,
 He flies remote beneath the flood in vain.”

FALCONER.

Whales are taken in great numbers about Spitsbergen, or Greenland, and other northern countries by the English, the Dutch, &c. The South Sea Company, for se-

veral years, used to send annually on this expedition about twenty sail of ships, every ship being above three hundred tons of burden, and carrying forty-five men: this fleet usually sailed about the end of March, but seldom began to fish till May. When they begin their fishery, each ship is fastened or moored with nose-hooks to the ice. Two boats, each manned with six men, are ordered by the commodore to look out for the coming of the fish for two hours, when they are relieved by two more, and so by turns: the two boats lie at some small distance from the ship, each separated from the other, fastened to the ice with their boat-hooks, ready to let go in an instant at the first sight of the Whale. Here the dexterity of the Whale hunters is to be admired; for as soon as the animal shows itself, every man is at his oar, and they all rush on the Whale with prodigious swiftness; at the same time taking care to come behind its head, that it may not see the boat, which sometimes so alarms it, that it plunges down again before they have time to strike it. But the greatest care is to be taken of the tail, with which it many times does very great damage, both to the boats and seamen. The harpooner, who is placed at the head or bow of the boat, seeing the back of the Whale, and making the onset, thrusts the harpoon with all his might into its body by the help of a staff fixed to the iron for this purpose, and leaves it in, a line being fastened to it of about two inches in circumference, and one hundred and thirty-six fathoms long. Every boat is furnished with seven of these lines, from the motion of which, when let run, they observe the course of the Whale.

As soon as the Whale is struck, the third man in the boat holds up his oar, with something on the top, as a signal to the ship; at the sight of which the man who is appointed to watch gives the alarm to those that are asleep, who instantly let fall their other four boats, which hang on the tackles, two at each side, ready to let go at a minute's warning, all furnished with six men each, harpoons, lances, lines, &c. Two or three of these boats row to the place where the Whale may be expected to come up again; the other to assist the boat that first struck

it, with line; as the Whale will sometimes run out three more boats' lines, all fastened to each other, for when the lines of the first boat are almost run out, they throw the end to the second to be fastened to theirs, and the second boat does the same to the third, and so on. In this manner line is supplied to such an extent that a large Whale has been known to carry off three miles of it.

A Whale, when he is first struck, will run out above a hundred fathoms of line, before the harpooner is able to take a turn round the boat's stern; and with such swiftness that a man stands ready to throw water on the line to quench it, in case it should take fire, which it frequently does. There was, many years ago, a boat to be seen in the South Sea dock at Deptford, the head of which was sawed off by the swiftness of the line running out. The harpoon would be of but little avail in the destruction of this animal; but part of the rowers, either at the first onset, or when, in order to fetch his breath, he rises to the surface and discovers himself to view, throwing aside their oars, and taking up their very sharp lances, thrust them into his body, till they see him spurt the blood through the blow-holes, the sight of which is a sign of the creature's being mortally wounded. The fishermen, upon the killing of a Whale, are each entitled to some small reward. After the Whale is killed, they cut all the lines that were fastened to it, and then cut off the tail; upon this it instantly turns on its back; and in this manner they tow it to the ship, where they fasten ropes to keep it from sinking; and, when it is cold, begin to cut off the blubber.

The blubber of a Whale is frequently found to be eighteen or twenty inches thick; which yields fifty or sixty puncheons of oil, each puncheon containing seventy-four gallons; and the upper jaw yields about six hundred pieces of whalebone, most of which are about twelve feet long, and six or eight inches broad; the whole produce of a Whale being worth one thousand pounds, sometimes more or less according to the size of the animal. Whilst the men are at work on the back of the Whale they have spurs on their boots, with two prongs, which come down

on each side of their feet, lest they should slip, the back of the Whale being very slippery. All the ships have orders to quit the North Seas by the 24th of June, for then the Whales begin to be very mischievous.

When the Whale feeds, it swims with considerable velocity below the surface of the sea, with its jaws widely extended. A stream of water consequently enters its mouth, carrying along with it immense quantities of cuttle-fish, sea-blubber, shrimps, and other small marine animals. The water escapes at the sides; but the food is entangled, and, as it were, sifted by the fringe of Whalebone within the mouth; this kind of strainer is rendered necessary by the very small gullet, which in a Whale of sixty feet long, does not exceed four inches in width. The sailors say that a penny-loaf would choke a Whale.

The Whale bellows fearfully when wounded or in distress; and its young is called a cub.



THE SPERMACETI WHALE, OR CACHALOT.


(*Physeter macrocephalus.*)

THIS animal has teeth in the lower jaw only; and no whalebone. The substance called spermaceti is extracted from its immense head, which is nearly half the size of the entire animal; and the throat is so large that it could swallow a shark. This must therefore have been the

Whale that swallowed the prophet Jonah, as described in Holy Writ.

The quantity of oil produced from the Spermaceti Whale is not so considerable as that obtained from the common or Greenland Whale, but in quality it is far preferable, as it yields a bright flame, without exhaling any nauseous smell. The substance known by the name of ambergris is also obtained from the body of this animal. It is generally found in the stomach, but sometimes in the intestines; and, in a commercial point of view, is a highly valuable production. The spermaceti is in a fluid state while the animal is living, and as soon as it is dead a hole is made in the head, and the liquid taken out with buckets. It becomes solid as it cools, and it is afterwards made into candles, &c.

When we reflect that the same Power whose will has formed the immense bulk of this marine monster has also given animation, senses, and passions to the smallest of the microscopic animalcules, how lowered must be the pride of man, who, standing in the middle, and nearly at equal distance from both, is yet unable to comprehend the mechanism which puts them in motion, and much less that intelligence and power which has given them life, and has assigned to them their respective stations in the universe! Let us then exclaim, with astonishment and gratitude, with the Psalmist: "O Lord, how inscrutable are thy ways, how magnificent thy works!"



THE DOLPHIN. (*Delphinus delphis.*)

THIS animal, like the whale, is not considered a fish, though it lives in the water, as it has warm blood and suckles its young, which are born alive. It has also lungs instead of gills, and is therefore obliged to raise its head above the surface of the water to breathe.


The Dolphin is from six to ten feet in length. The body is roundish, gradually diminishing towards the tail; the nose is long and pointed, the skin smooth, the back black or dusky blue, becoming white below. It has eight small teeth in each jaw; a dorsal and two pectoral fins, and a tail in the shape of a half-moon. The snout of the Dolphin is most useful to it when in search of eels and other fishes which harbour in the mud: This beak-like snout has probably made the French call the Dolphin the sea-goose.

Several curious stories have been related of this animal, most of which are fabulous. The anecdote of Arion, the musician, who, being thrown overboard by pirates, was indebted for his life to one of these animals, is well known, and acquired great credit among ancient poets, as it was said to be by his music that Arion charmed the Dolphin. There are several other fables mentioned by ancient authors to prove the philanthropy of the Dolphin. Since the province of *Dauphiné* in France has been united to the crown, the heir-apparent has been called "Dauphin," and quarters a Dolphin on his shield. Falconer, in his

beautiful poem, "The Shipwreck," describes the death of this fish in the following elegant manner :

"—— Beneath the lofty vessel's stern
 A shoal of sporting Dolphins they discern,
 Beaming from burnish'd scales refulgent rays,
 Till all the glowing ocean seems to blaze.
 In curling wreaths they wanton on the tide ;
 Now bound aloft, now downward swiftly glide.
 Awhile beneath the waves their tracks remain,
 And burn in silver streams along the liquid plain ;
 Soon to the sport of death the crew repair,
 Dart the long lance, or spread the bated snare.
 One in redoubling mazes wheels along,
 And glides, unhappy, near the tripple prong.
 Rodmond, unerring, o'er his head suspends
 The barbed steel, and every turn attends :
 Unerring aim'd, the missile weapon flew,
 And plunging, struck the fated victim through.
 The upturning points his pond'rous bulk sustain ;
 On deck he struggles with convulsive pain ;
 But while his heart the fatal javelin thrills,
 And fleeting life escapes in sanguine rills,
 What radiant changes strike the astonish'd sight,
 What glowing hues of mingled shade and light !
 No equal beauties gild the lucid west
 With parting beams all o'er profusely dress'd ;
 No lovelier colours paint the vernal dawn,
 When orient dews impearl the enamell'd lawn ;
 Than from his sides in bright suffusion flow,
 That now with gold empyreal seem to glow ;
 Now in pellucid sapphires meet the view,
 And emulate the soft celestial hue ;
 Now beam a flaming crimson to the eye,
 And now assume the purple's deeper dye :
 But here description clouds each shining ray ;
 What terms of art can Nature's power display !"

Unfortunately for poetry, the beautiful colours of the dying Dolphin exist entirely in the fancy of the poet ; as the Dolphin in a dying state displays no tints but black and white.




THE PORPOISE. (*Phocæna vulgaris.*)

THE Porpoise is one of the cetaceæ, and nearly allied to the dolphin, but it has not the beaked snout of that animal. The length of the Porpoise, from the tip of the snout to the end of the tail, is from four to eight feet, and its girth about two feet and a half. The figure of the whole body is conical; the colour of the back is deep blue, inclining to shining black; the sides are grey, becoming white below. The tail is forked, and composed of several rays united by a membrane. The body is covered with a rough skin, but has no scales: there are only three fins, one on the back, and one on each shoulder. The eyes are very small. When the flesh is cut up, it looks very much like pork; but although it was once considered a sumptuous article of food, and is said to have been occasionally introduced at the tables of the old English nobility, it certainly has a disagreeable flavour. Porpoises live on small fish, and appear generally in large shoals, particularly in the mackerel and herring seasons, at which time they do very great damage to fishermen, by breaking and destroying the nets to get at their prey. Their motion in the water is a kind of circular leap; they dive deep, but soon again rise up in order to breathe. They are so eager in the pursuit of their prey, that they sometimes ascend large rivers, and have even been seen above Westminster Bridge. They have no gills, and blow out

the water with a loud noise, which in calm weather may be heard at a great distance. They are seen nearly in all seas, and are very common upon the British coasts, where they sport with great activity, chiefly at the approach of a squall.

The Grampus, or Thresher, is a species of Porpoise, and a decided and inveterate enemy to whales; which they attack in great flocks, fastening round them like so many bull-dogs, making them roar with pain, and frequently killing and devouring them. They are usually from twenty to twenty-five feet in length, and in general form and colour resemble the common Porpoise; but the lower jaw is considerably wider than the upper, and the body in proportion, is somewhat broader and more deep. The back-fin sometimes measures six feet in length. In one of the poems of Waller, a story (founded on fact) is recorded of the parental affection of these animals. A Grampus and her cub had got into an arm of the sea, where, by the desertion of the tide, they were enclosed on every side. The men on shore saw their situation, and ran down upon them with such weapons as they could at the moment collect. The poor animals were soon wounded several places, so that all the immediately surrounding water was stained with their blood. They made many efforts to escape; and the old one, by superior strength, forced itself over the shallow into the ocean. But though in safety herself she would not leave her young one in the hands of assassins. She therefore again rushed in; and seemed resolved, since she could not prevent, at least to share the fate of her offspring. The story concludes with poetical justice; for the tide coming in, conveyed them both off in safety; and it is probable from the great thickness of their skins, that their wounds had not been very deep.




THE SEA-UNICORN, OR NARVHAL,

(*Monodon monoceros*,)

A MARINE animal, differing from all the cetacea to which it belongs, in not having any teeth, properly so called, and in being armed with a horn of seven or eight feet in length, which projects from the head. This horn is white, spirally striated throughout its whole length, and tapering to a point: it is harder, whiter, and more valuable than the ivory of the elephant, and was formerly in high repute for its supposed medical properties: it may be sometimes seen set with an elegant head as a walking-stick. The animal itself is from twenty to forty feet in length, and is occasionally found with two of these horns; indeed, there is always the germ of a second horn both in the male and female, though it is rarely developed in the former, and never in the latter, from which we may conjecture that the females trust entirely to the males for their defence, as we know is the case with several of the mammalia. When there is only one horn, it is always on the left side of the head; and when there are two, the horn on the left side is always larger than the other. This animal chiefly inhabits the arctic seas, and its food is said to consist of the smaller kinds of flat fish and other marine animals; its horn is useful in breaking away the ice when it wants to come up to breathe. The blubber supplies a small quantity of very fine oil, and the Greenlanders are very partial to the flesh.

SECTION II.—Cartilaginous Fishes.

THE SHARK. (*Squalus carcharias.*)

“Increasing still the terrors of the storms,
His jaws horrific arm'd with threefold fate,
Here dwells the direful Shark.”

THE Shark differs from the whale in not being one of the mammalia. It is cold-blooded, and does not suckle its young. It has no lungs, and its mode of breathing is like that of other fishes, except that its gills are fixed, and the water escapes by five elongated apertures on each side. Some of the Sharks produce their young alive, and others lay eggs or, rather, horny cases of an oblong shape, with long tendrils at each of the four corners. After the young Sharks are hatched, these curious cases are often washed on shore, and are called mermaids' purses.

The bones of the Shark are like gristle, and very different from those of most other fishes. Hence all the fishes with bones similar to those of the Shark are placed in a separate order, and called cartilaginous fishes.

The White Shark is sometimes found weighing nearly two thousand pounds. The throat is often large enough to swallow a man; and a human body has sometimes been found entire in the stomach of this tremendous animal.

He is furnished with six rows of sharp triangular teeth, which amount in all to a hundred and forty-four, serrated on their edges, and capable of being erected or depressed at pleasure, owing to a curious muscular mechanism in the palate and jaws of the Shark. The whole body and fins are of a light ash-colour; the skin rough, and employed to smooth cabinet work, or to cover small boxes or cases. His eyes are large and staring, and he possesses great muscular strength in his tail and fins. Whenever he spies, from the deepest recesses of the sea, a man swimming or diving, he darts from the place, up to his prey, and if unable to take in the whole, or snatch away a limb, he follows for a long time the boat or vessel in which the more nimble swimmer has found a safe and opportune retreat: but seldom does he let any one escape his jaws, and get off entire. Sir Brook Watson was swimming at a little distance from a ship, when he saw a Shark making towards him. Struck with terror at its approach, he cried out for assistance. A rope was instantly thrown; but even while the men were in the act of drawing him up the ship's side, the monster darted after him, and, at a single snap, tore off his leg.


We are told that, in the reign of Queen Anne, some of the men of an English merchant-ship, which had arrived at Barbadoes, were one day bathing in the sea, when a large Shark appeared, and was rushing upon them. A person from the ship called out to warn them of their danger; on which they all immediately swam to the vessel, and arrived in perfect safety, except one poor man, who was cut in two by the Shark, almost within reach of the oars. A comrade and intimate friend of the unfortunate victim, when he observed the severed trunk of his companion, was seized with a degree of horror that words cannot describe. The insatiate Shark was seen traversing the bloody surface in search of the remainder of his prey, when the brave youth plunged into the water, determining either to make the Shark disgorge, or to be buried himself in the same grave. He held in his hand a long and sharp-pointed knife, and the rapacious animal pushed furiously towards him; he had turned on his side,

and had opened his enormous jaws, in order to seize him, when the youth, diving dexterously under, seized him with his left hand, somewhere about the upper fins, and stabbed him several times in the belly. The Shark, enraged with pain, and streaming with blood, plunged in all directions in order to disengage himself from his enemy. The crews of the surrounding vessels saw that the combat was decided: but they were ignorant which was slain, until the Shark, weakened by loss of blood, made towards the shore, and along with him his conqueror; who, flushed with victory, pushed his foe with redoubled ardour, and, by the aid of an ebbing tide, dragged him on shore. Here he ripped up the bowels of the animal, obtained the severed remainder of his friend's body, and buried it with the trunk in the same grave. This story, however incredible it may appear, is related in the History of Barbadoes, on the most satisfactory authority.

Had nature allowed this fish to seize his prey with as much facility as many others, the Shark tribe would have soon depopulated the ocean, and reigned alone in the vast regions of the sea, till hunger would have forced them to attack and ultimately destroy each other; but the upper jaw of this devouring animal, is so constructed as to offer, by its prominency, an impediment to the Shark's easily seizing his prey; and consequently when on the point of catching hold of anything, he is obliged to turn on one side, which troublesome evolution often gives the object of his pursuit time to escape. The flesh of this fish is of a disagreeable taste, and cannot be eaten with any kind of relish, except the part near the tail. The Greenland Shark is another very voracious species; and one extremely difficult to kill. It is the great enemy of the whale, and devours the bodies of those left by the fishers after cutting out all the blubber.

The *Dog-fishes* are so excessively voracious, that they are altogether fearless of mankind. They follow vessels with great eagerness, seizing with avidity everything eatable that is thrown overboard; and have sometimes been known to throw themselves on fishermen, and on persons bathing in the sea. As, however, they are much smaller

and weaker than most of the other Sharks, they do not always attack their enemies by open force, but generally have recourse to stratagem. They, consequently, conceal themselves in the mud, and lie in ambush, like the ray or skate-fish, (also one of the cartilaginous fishes,) until they have an opportunity of successfully attacking their prey. On the coasts of Scarborough, where haddocks, cod, and Dog-fish are in great abundance, the fishermen universally believe that the Dog-fish make a line or semicircle to encompass a shoal of haddocks and cod, confining them within certain limits near the shore, and eating them as occasion requires: they are therefore considered very destructive to this fishery. The flesh of the Dog-fish is hard and disagreeable; its skin, when dried, is made into the well-known *shagreen*, and from the liver a considerable quantity of oil may be extracted. Shagreen is also made from the skin of other cartilaginous fishes.



THE SKATE, OR MAID (*Raia batis*)

Is a species of the Ray, which was long disregarded in this country as a coarse, bad-tasted food, but which now appears upon our best tables. It is still, however, disregarded in Scotland and the north of England, where its flesh is principally used as a bait for other fish. On some parts of the continent, where these fish are caught in great abundance, they are dried for sale. The best season for Skate is the spring of the year. The body is broad and flat, of a brown colour on the back, and white on the lower side: the head is not distinct from the

body, so that this fish and all belonging to this genus are apparently acephalous, or without a head. Dr. Monro has remarked, that in the gills of a large Skate there are upwards of one hundred and forty-four thousand subdivisions, or folds; and that the whole extent of this membrane, whose surface is nearly equal to that of the whole human body, may be seen by a microscope to be covered with a network of vessels, that are not only extremely minute, but exquisitely beautiful. The tail of the Skate is long and generally prickly: the eggs are often found on the beach after a storm, in the shape of a square bag, with two horns at each end, as here represented.




In this horny case the embryo is contained, and grows till it has acquired strength enough to burst through its prison. The colour of the bag is maroon, and the substance like thin brown parchment or leather. The female begins to drop these singly in the month, of May, and continues to do so for several months, to the number of two or three hundred. These cases are often washed on shore, and when empty are called Skate-barrows.

The Skate sometimes attains a very large size. Willoughby speaks of one so huge that it would have served one hundred and twenty men for dinner. Some naturalists are of opinion that these fishes are the largest inhabitants of the deep, and that only the smallest of them come near the surface of the water, the biggest remaining flat at the bottom of the sea, where an unfathomable depth secures them against the wiles of man.

The *Thornback* resembles the Skate in its general appearance; the principal difference consists in the latter

having sharp teeth, and a single row of spines upon the tail, while the former has blunt teeth, and several rows of spines both upon the back and tail. A Thornback was caught near the island of St. Kitt's, in the year 1634, which measured twelve feet in length, and nearly ten in width. It is sometimes eaten in England, but as its flesh is inferior to that of the Skate, it is generally sold at a low price. The young ones, however, which have the denomination of *Maids*, are delicate eating.




THE TORPEDO, OR ELECTRIC RAY.

(*Torpedo vulgaris.*)

THIS curious fish is capable of giving a violent shock, like that produced by the electrical machine, to the person who handles it. The body is nearly circular, and thicker than any other of the Ray kind, and is sometimes so large as to weigh between seventy and eighty pounds. The skin is smooth, of a dusky brown colour, and white underneath. The ventral fins form on each side, at the end of the body, nearly a quarter of a circle. The tail is short, and the two dorsal fins are near its origin. The mouth is small, and as in the other species, there are on each side below it five breathing apertures.

The shock imparted by the touch of the Cramp-fish, as the Torpedo is vulgarly called, is often attended with a sudden sickness at the stomach, a general tremor, a kind of convulsion, and sometimes a total suspension of the

faculties of the mind. Such power of self-defence has Providence allowed this lumpish and inactive fish. Whenever an enemy approaches, the Torpedo emits from its body that benumbing shock, which incapacitates the other instantly, and it thereby gets time to escape. Nor is it merely a means of defence, but an advantage in other respects, for the Torpedo thus benumbs its prey, and easily seizes upon it. The animals thus killed also become more easy of digestion.




THE HAG-FISH, (*Gastrobanchus glutinosa*.)

A CARTILAGINOUS fish, which in its general appearance bears a near resemblance to the Lamprey. Its colour is dusky bluish above, and reddish towards the head and tail; its length from four to six inches. The Hag-fish is remarkable for its total want of eyes; its mouth is of an oblong form, with two beards or cirri on each side, and on the upper part four. On the top of the head is a small spout-hole, furnished with a valve, by which it can be closed at pleasure. A double row of pores extends beneath the body, from one extremity to the other, which on pressure exude a quantity of viscid fluid, which, when attacked by large fish, the Hag throws out, so as to cloud the surrounding element in such a manner as to render itself invisible to its assailants. "The habits of this fish are highly singular: it will enter the bodies of such fishes as it happens to find on the fishermen's hooks, and which consequently have lost the power of escaping its attack; and gnawing its way through the skin, will devour all the internal parts, leaving only the bones and the skin.


If put into a large vessel of sea-water, it is said in a very short space to render the whole water so glutinous that it may easily be drawn out in the form of threads."

SECTION III.—Bony Fishes.



THE PILOT-FISH. (*Naucrates ductor*.)


THE body of this fish is long, the head compressed, rounding off in front, without scales as far as the operculum. The mouth is small, the jaws of equal length, and furnished with small teeth; the palate has a curved row of similar teeth in front, and the tongue has teeth all along. The colour varies in several species. The Pilot-fish will frequently attend a ship during its course at sea for weeks, or even months together; and there are many curious stories told respecting its habits, in occasionally directing a shark where to find a good meal, and also in warning him how to avoid a dangerous bait. Whether this be true or not will be difficult to determine; but it is certain that this voracious fish is generally found in company with the shark, and picks up the smaller pieces of food which his predatory master drops, either by accident or design.



THE REMORA, OR SUCKING-FISH,
(*Echeneis remora*.)

RESEMBLES the herring; its head is thick, naked, depressed, and marked on the upper side with transverse rough lines. The fins are seven in number; the under jaw is longer than the upper, and both furnished with teeth. This fish is provided by nature with a strong adhesive power, and, by means of the grooved lines on its head, can attach itself to any animal or body whatever. We might suppose that a small fish with seven acting fins, armed like a galley with oars, would have a great power of motion in the water, but, for some reason unknown to us, Providence has contrived for him an easier way of travelling, by enabling him to fix himself to the hull of a ship, and even to the body of a larger animal than himself, as the whale, the shark, and others. Our forefathers believed that, small as he is, this fish had the power of arresting the progress of a ship in its fastest sailing by adhering to the bottom.

“ The Sucking-fish beneath, with secret chains,
Clung to the keel, the swiftest ship detains.
The seamen run confused, no labour spared,
Let fly the sheets, and hoist the topmost yard.
The master bids them give her all the sails,
To court the winds and catch the coming gales.
But, though the canvass bellies with the blast,
And boisterous winds bend down the cracking mast,
The bark stands firmly rooted in the sea,
And will, unmoved, nor winds nor waves obey :
Still, as when calms have flatted all the plain,
And infant waves scarce wrinkle on the main.
No ship in harbour moor'd so careless rides,
When ruffling waters tell the flowing tides ;
Appall'd, the sailors stare, through strange surprise,
Believe they dream, and rub their waking eyes.”



THE SEA-WOLF, OR SEA-BAT,
(*Anarrhichas lupus*,)


Is often caught at Heligoland, an island not far from the mouth of the Elbe; and is about three feet in length, and having a larger and flatter head than the shark. The back, sides, and fins are of a bluish colour; the body is nearly white; the whole skin is smooth and slippery, without any appearance of scales. It is of a very voracious nature, and has a double row of sharp and round teeth, both in the upper and lower jaw. Its appetite, however, does not lead it to destroy fishes similar in shape to itself, as it is supposed to feed entirely on crustaceous animals, whose shells it breaks easily with its teeth. It is sometimes found in the northern seas exceeding twelve feet in length, and owes its name to its natural fierceness and voracity. The fishermen dread its bite, and endeavour as speedily as possible to strike out its fore-teeth, which are so strong, that they are capable of leaving an impression on an anchor. The fins nearest the head spread themselves, when the animal is swimming, in the shape of two large fans, and their motion contributes considerably to accelerate its natural swiftness. The flesh is good, but not often eaten.



THE HORNED SILURE


Grows to a large size, weighing sometimes three hundred pounds, and measuring eight to ten feet in length, and two in breadth. It has a broad, flat, thin head; and the horns, which are on each side of the upper lip, are armed with short crooked spines, like teeth. A remarkable peculiarity in this fish is the dorsal fin, which is close to the head, and is long, stiff, dentated like the horns, and is, no doubt, an instrument of defence. In colour it resembles the eel, and has no scales; only one small fin on the back, and a forked tail; its flesh is esteemed next to that of the eel, and has a similar flavour. This fish is a great depredator, and makes considerable havoc among the smaller inhabitants of the rivers and lakes which it inhabits. The Danube, and several other rivers of Germany, and the lakes of Switzerland and Bavaria, contain numerous species of the genus *Silurus*.

One of these fish, caught at Surinam, on examination, was found to have its mouth filled with yellow eggs, none of which contained a fish completely formed; hence it is concluded that the Silure, to defend her eggs from her voracious enemies, hatches them instinctively in her mouth. But she is supposed at times to emit them, when in search of food; and when satisfied, to take them into her mouth again.



THE FATHER LASHER. (*Cottus bubalis.*)

THE whimsical denomination of Father Lasher, given to this fish, cannot be easily accounted for; perhaps it may be ascribable to the quick and repeated lashings of its tail, when the fish is caught and thrown upon the sand. The length is about eight or nine inches, and it is usually found under stones, on the rocky coasts of our island. In Greenland these fish are so numerous, that the inhabitants depend largely upon them for their food. When made into soup, they are nutritive and wholesome. The head is large, and armed with vast spines, by which this fish combats every enemy that attacks it, swelling out its cheeks and gill-covers to an unusual size. Its colour is a dull brown, mottled with white, and sometimes mixed with red; the fins and tail are transparent, and the lower part of the body a shining white.



THE MONK-FISH, OR ANGEL-FISH,
(*Squatina Angelus.*)

Is very voracious, and feeds upon all kinds of flat fish,


as soles, flounders, &c. It is often caught on the coasts of Great Britain, and of such a size as to weigh sometimes a hundred pounds. This fish seems to be of a middle nature between the rays and sharks, and is called by Pliny the Squatina; a name which seems to bring this species near that of the skate. Its head is large; the mouth has five rows of teeth, which are capable of being raised or depressed at pleasure. The back is of a pale ash-colour; the belly white and smooth. The shores of Cornwall are often frequented by this fish, but its flesh does not deserve to be praised, being hard, and of a very indifferent flavour.

It is supposed to have acquired the name of Angel-fish, from its extended pectoral fins bearing some similarity to wings; and of monk-fish, from its rounded head appearing as if enveloped in a monk's hood. The skin is rather rough, and is used for polishing, and other works in the arts. Mr. Donovan says that the Turks of the present day make shagreen of it.



THE SAW-FISH. (*Pristis antiquorum.*)

This fish is found in the European and Atlantic seas. Its body is flattened anteriorly with four or five branchial openings below on each side; two spiracles behind the eyes; no anal fin; the head prolonged into a depressed bony beak, with strong pointed spines on each side; the lips are rough and sharp like a file, supplying the place of teeth. With its formidable weapon which resembles a toothed saw, this fish attacks the largest whales, and inflicts very severe wounds. The colour of its body is of a grayish brown above, and paler below;—its length about fifteen feet, the saw being about a third of the whole.




THE SWORD-FISH (*Xiphias gladius*)

HAS received its name from its long snout resembling the blade of a sword. It sometimes weighs above one hundred pounds, and is fifteen feet in length. The body is of a conical form, black on the back, white under the body, a large mouth, and no teeth; the tail is remarkably forked. The Sword-fish is often taken off the coast of Italy, in the Bay of Naples, and about Sicily. They are struck at by the fishermen, and their flesh is considered as good as that of the sturgeon by the Sicilians, who seem to be particularly fond of it. Other European seas are not destitute of this curious animal.

The Sword-fish and the whale are said never to meet without coming to battle; and the former has the reputation of being always the aggressor. Sometimes two Sword-fishes join against one whale; in which case the combat is by no means equal. The whale uses his tail in his defence; he dives deeply into the water, head foremost, and makes such a blow with his tail, that, should it take effect, it kills the Sword-fish at a single stroke; but the latter is in general sufficiently adroit to avoid it, and immediately rushes at the whale, and buries its weapon in his sides. When the whale discovers the Sword-fish darting upon him, he dives to the bottom; but is closely pursued by his antagonist, who compels him again to rise to the surface. The battle then begins afresh, and lasts until the Sword-fish loses sight of the whale, who is at length compelled to swim off, which his superior agility enables him to do. In piercing the whale's body with

the tremendous weapon at his snout, the Sword-fish seldom inflicts a dangerous wound, not being able to penetrate beyond the blubber. This animal can drive its sword with such force into the keel of a ship, as to bury it wholly in the timber. A part of the bottom of a vessel, with the sword imbedded in it, is to be seen in the British Museum.



THE FLYING SCORPION.


How admirable is Nature! how extensive her power, and how various the forms with which she has surrounded the united elements of animated matter! From the uncouth shape of the wallowing whale, of the unwieldy hippopotamus, or ponderous elephant, to the light and elegant form of the painted moth or fluttering humming-bird, she seems to have exhausted all ideas, all conceptions, and not to have left a single figure untried. The fish represented above, is one of those, in the outlines and decorations of which appear the discordant qualities of frightfulness and beauty. Armed *cap-a-pié*, surrounded with spines and thorns bristling on his back, and fins like an armed phalanx of lancebearers; and decorated on the body with yellow ribands, interwoven with white fillets; and on the purple fins of his breast, with the milky dots of the pintado, the Scorpion presents a very extraordinary contrast. His eyes, like those of which poets sang when celebrating the

Nereids and Naiads, consist of black pupils, surrounded with a silver iris, radiated with alternate divisions of blue and black. The rays of the dorsal fin are spiny, spotted brown and yellow, conjoined below by a dark brown membrane, and separate above; the ventral fins are violet with white drops, and the tail and anal fins are a sort of tessellated work of blue, black, and white, united with the greatest symmetry, and not unlike those ancient fragments of Roman pavements often found in this island.

This variegated fish is found in the rivers of Amboyna and Japan; its flesh is white, firm, and well tasted, like our perch, but it does not grow so large; it is of a very voracious disposition, feeding on the young of other fish, some of which, two inches in length, have been found in its craw. The skin has both the appearance and smoothness of parchment. To the tremendous armour of its back, fins, and tail, this fish owes the name of Scorpion.

The *Butterfly Fish* is about six or seven inches long, and inhabits the Adriatic Sea. In October it is not uncommon at Venice, where it is offered for sale among the great quantity of various fish which the coasts of Italy afford. It has no apparent scales, and is of a faint blue or ash-colour; the dorsal fin is elegantly spotted with black, and the flesh is well tasted and tender. This fish bears some resemblance and apparent affinity to the *Scorpion*, the *Gurnard*, and *Father Lasher*.

The *Bull-head*, or *Cottus*, has a body of a wedge-like form; the head is flat and broader than the body; the fin covering the gills has six spines; the head is furnished with prickles, knobs, and beards. It is of the prickly-finned thoracic tribe.



THE TOAD-FISH.

THE figure of the animal above has been correctly copied at the foot of Blackfriars Bridge in the month of May, 1812, from a specimen exhibited there for several days by the fishermen who had caught it below bridge: it is about three feet and a half long, the head forming more than the third part of the whole; the mouth is uncommonly large, and armed with several rows of white and sharp teeth, not very regularly set, and seemingly moving in the cartilaginous jaws. The tongue appears fleshy and broad, the pectoral fins widely extended, and placed a little below the eyes; and between them and the mouth a sort of suture in black stitches and in vandyke or zigzag shape, appears very conspicuous upon the silvery white of the skin in this part of the animal. The back is armed with several rows of tubercles; but the most curious peculiarity belonging to this ill-shaped citizen of the deep is, that he seems to have received the 'gift of feet'; for the ventral fins are exactly in that shape, with divided toes, the use of which appears to be that of opening the ooze at the bottom of the sea, or the sand on the shores, where this gluttonous fish conceals himself with his jaws wide open, to catch the imprudent flounder or sole, or any others which their giddiness leads to the dangerous abyss. The colour is brownish, and the scales hardly perceptible.


The *Sun Fish*, (*orthagoriscus*) appears like the forepart of the body of a large fish, which has been amputated in the middle. The mouth is small, with two broad teeth only in each jaw. Its nearly circular form, and the

silvery whiteness of the sides, together with their brilliant phosphorescence during the night, have obtained for it very generally the appellations of sun or moon fish. While swimming it turns round like a wheel, and sometimes floats with its head above water, when it appears like a dying fish. It grows to a large size; sometimes being seven or eight feet in circumference, and weighing from three to five hundred pounds. The back of this curious marine animal is of a rich blue colour. It frequents the coasts of both the ancient and new continent, and has been found on the shores of England. There are several species of this genus of fish, but the difference between them is so inconsiderable, that it would not be interesting to our readers to find here a minute description of each.



THE TRUMPET-FISH (*Centriscus scolopax*)

Is found in the Mediterranean, and is from four to eight inches in length. He has a large snout, long and narrow at the end; the eyes are large, the irides red, and the body is covered with rough cinereous scales. The anterior part of the body has two bony substances like fins, and another situate on the belly. This fish is often found in the ocean, where he seems to be driven by storms, as he is seldom seen there in any other than tempestuous weather. His breathing the water out of his snout with a sounding noise, has occasioned our sailors to call him the *Trumpet*. This species belongs to the *Centriscus* genus.



THE LUMP-SUCKER, OR SEA-OWL.

(*Cyclopterus lumpus.*)

THIS odd-shaped fish derives its name chiefly from the clumsiness of its form; it is also called the Cock Paddle. Its colour when in the highest perfection combines various shades of blue, purple, and rich orange; the abdomen is red; it has no scales, but on all sides sharp black tubercles, in shape like warts; on each side are three rows of sharp prickles, and on the back two distinct fins. The great resort of this species is on the Northern seas, about the coast of Greenland; it is also caught in many parts of the British seas, during the spring season, when it approaches the shore for the purpose of depositing its spawn; and in the month of March it may be seen at the stalls of the London markets. This unseemly fish is usually about a foot in length, and ten or more inches in breadth, and sometimes weighs seven pounds. The flesh is but indifferent.

The Lump-sucker is very remarkable for the manner in which its ventral fins are arranged. They are united by a membrane so as to form a kind of oval and concave disc, by means of which it is enabled to adhere with great force to any substance to which it fastens itself. Pennant says, that, on throwing an individual of this species into a pail of water, it adhered so firmly to the bottom, that on taking the fish by the tail, the whole pail was lifted up, though it held some gallons.



THE OCCELLATED SUCKER.

(*Cyclopterus Ocellatus*.)

ANOTHER Malacopterygious fish, a relative of the lump-sucker, and chiefly remarkable for the singular appendage observable on its head. It possesses similar tenacity of suction. The utility of this faculty to animals inhabiting the rocky shores and turbulent seas of Greenland, is sufficiently obvious.

In the Northern seas great numbers of the different species of Lump-suckers are devoured by the seals, who swallow all but the skins, quantities of which thus emptied are seen floating about in the spring months; and, it is said, that the spots where the seals carry on their depredations can be readily distinguished by the smoothness of the water.



THE ANGLER. (*Lophius piscatorius*.)

THIS extraordinary fish is occasionally met with on our coasts, and is commonly known by the names of the Fishing Frog, Toad Fish, and Sea Devil. In shape it is the most uncouth and unsightly of the piscatory tribe, resembling the frog in its tadpole state. It grows to a large size. A specimen taken in the sea, near Scarborough, was between four and five feet in length, the head considerably larger than the body, round at the circumference, flat above; the mouth is of a prodigious size, being a yard in width, and armed with sharp teeth. It lives as it were in ambush at the bottom of the sea, and by means of its fins stirs up the mud and sand, so as to conceal itself from other fishes on whom it preys. The manner in which it procures its prey is very extraordinary, the peculiarity of its construction forbidding the possibility of rapid movement. Two long tough filaments are placed above the nose, each of them furnished with a thin appendage, closely resembling a fishing-line when baited and flung out. The back is provided with three others, united by a web and forming the first dorsal fin. Pliny notices these remarkable appendages, and explains their use. "The Fishing Frog," says he, "puts forth the slender horns situated beneath his eyes, enticing by that means the little fish to play around till they come within

his reach, when he springs upon them." But it is not only the lesser inhabitants of the water that the Angler ensnares! even the fierce and voracious sword-fish often falls a prey to his artifices. For this reason, the fishermen regard the Angler with peculiar respect, and carefully release, and restore him to his usual haunts, when accidentally entangled in their nets.


Cicero also notices this extraordinary creature, in his treatise on the nature of the Gods. He observed its wonderful construction when musing on the shores of Sicily.



THE FOUR-HORNED TRUNK-FISH.

(*Ostracion quadricornis*.)

THESE singular fishes are distinguished from all others by the bony covering which envelopes them. The head and body are covered with plates of bone, forming an inflexible cuirass, and leaving exposed only the tail, fins, mouth, and a portion of the gill opening. They have no ventral fins, and the dorsal and anal are placed far back. Their liver is large, and abounds with oil. The Trunk-fish is a native of the Indian and American seas. Some of the species are considered excellent eating.




THE GLOBE FISH (*Tetraodon hispidus*)

Is an oblong fish, inhabiting the seas of Carolina, and

endowed with the extraordinary property of swelling its under surface into a large globe. This sudden enlargement not only alarms the enemies of the Tetrodon, but prevents them from making good their hold, by presenting to their grasp little more than an inflated bag. It is also covered with spines, which merely adhere to the skin, and are capable of being erected on any sudden emergency; thus giving to an innocent and defenceless creature a most formidable appearance.

When inflated they roll over on their backs, floating in this position, without any power of directing their course. Some species are reckoned poisonous. One is electrical, (*Tetraodon lineatus*,) and is found on the Nile, cast on shore by the inundations, and picked up by the children as a plaything.



THE CAVALLO-MARINO, OR SEA-HORSE.

(*Hippocampus brevisrostris*.)

THIS is a small fish of a curious shape. The length is from six to ten, and sometimes reaches twelve inches: the head bears some resemblance to that of a horse, whence originates its name. A series of longitudinal and transverse ridges run from the head to the tail, which is spirally curved and prehensile.

The following account of two specimens taken alive at Guernsey, in June, 1835, by F. C. Lukis, Esq., is extracted from Yarrell's *British Fishes*. These creatures

were kept about twelve days in a glass vessel, and their actions were equally novel and amusing. "An appearance of search for a resting-place induced me," says Mr. Lukis, "to consult their wishes, by placing seaweed and straws in the vessel: the desired effect was obtained, and has afforded me much to reflect upon in their habits. They now exhibit many of their peculiarities, and few subjects of the deep have displayed, *in prison*, more sport or more intelligence.

"When swimming about, they maintain a vertical position; but the tail is ready to grasp whatever meets it in the water, quickly entwines in any direction round the weeds, and when fixed, the animal intently watches the surrounding objects, and darts at its prey with the greatest dexterity.

"When the animals approach each other, they often twist their tails together, and struggle to separate or attach themselves to the weeds; this is done by the under part of their cheeks or chin, which is also used for raising the body when a new spot is wanted for the tail to entwine afresh. The eyes move independently of each other, as in the chameleon, and this, with the brilliant changeable iridescence about the head, and its blue bands, forcibly reminds the observer of that animal."




THE STURGEON (*Accipenser sturio*)

SOMETIMES grows to the length of eighteen feet, and has been found to weigh five hundred pounds. It has a long, slender, pointed nose, small eyes, and a small mouth, destitute of teeth, placed beneath and unsupported by the

maxillæ; so that when the animal is dead, the mouth remains always open. The body is covered with five rows of large bony tubercles, and the under side is flat; it has one dorsal fin, two pectoral, two ventral, and one anal. The upper part of the body is of a muddy olive colour, and the under part silvery. The tail is bifurcated, the upper part being much longer than the under. Sturgeons subsist principally on insects and marine plants, which they find at the bottom of the water, where they mostly resort.

The Sturgeon annually ascends our rivers in the summer, particularly those of the Eden and Esk; and when caught, as it sometimes is, in the salmon-nets, it scarcely makes any resistance, but is drawn out of the water apparently lifeless. One of the largest Sturgeons ever caught in our rivers was taken in the Esk a good many years ago: it weighed four hundred and sixty pounds. This fish is found in most of the rivers in Europe; it is also common in those of North America, and especially in the lakes and rivers of Northern Asia.

The flesh of the Sturgeon is delicious; and it was so much valued in the time of the Emperor Severus, that it was brought to table by servants with coronets on their heads, and preceded by music. In London, every Sturgeon that is caught in the Thames is presented by the Lord Mayor to the Sovereign. The roe, when preserved with salt and oil, is called *caviar*, and is a favourite dish with many persons; the best is made in Russia. The flesh is also pickled or salted, and sent all over Europe. So prolific is this fish, that Catesby says, the females frequently contain a bushel of spawn each; and Leeuwenhoek found in the roe of one of them no fewer than one hundred and fifty thousand million eggs!




THE FLYING FISH OF THE OCEAN.

(*Exocoetus volitans.*)

THIS fish has a slender body, a projecting under-lip, and very large and prominent eyes. The ventral fins are small, but the fins on the back are so long and wide as to answer the purpose of wings, and aided by them the fish is enabled to rise out of the water, and support itself in the air. It must not be supposed, however, that the Flying-fish can soar like a bird; on the contrary, it can only spring from the water to a considerable height, (sometimes as much as twenty feet,) and fly about a hundred and fifty, or two hundred yards; most commonly, however, it does not rise above two or three feet from the water, and remains fluttering over the surface for about a hundred yards, when it again drops into its native element.

There is another Flying-fish (*Exocoetus exiliens*) in the Mediterranean, nearly allied to the gurnard. This fish is about a foot long; it is brown above, reddish below, and has blackish fins spotted with blue; and it

is armed with a long and pointed spine on the opercle, which it can raise, and with which it can inflict very serious wounds.



THE GURNARD. (*Trigla cuculus.*)


THIS genus is divided into several species. The Red Gurnard has fins and body of a bright red colour; and the head is large, and covered with strong bony plates. The eyes are large, round, and vertical. The mouth is large; and the palate and jaws are armed with sharp teeth. The gill-membrane has seven rays. The back has a longitudinal spinous groove on each side. There are slender articulate appendages at the base of each pectoral fin. This fish is not unfrequently met with on the southern shores of England; and is often seen exposed in the fish-markets of the maritime towns of Dorset and Devonshire, as well as in Cornwall. It is a pleasant-tasted fish, when properly stuffed and baked, the flavour being similar to that of the haddock.

Whilst in the water, the colours of the Red Gurnard are almost inconceivably brilliant and beautiful, particularly in the broad glare of sunshine, as they then vary, in the most pleasing manner, with every motion of the fish.

The *Grey Gurnard* usually measures from one to two feet in length. The extremity of the head, in front, is armed on each side with three short spines. The forehead and the covers of the gills are silvery; the latter being finely radiated. The body is covered with small scales; the upper parts are of a deep grey, spotted with white and yellow, and sometimes with black; and the lower parts

silvery. About the months of May and June the Grey Gurnards approach the shores in considerable shoals, for the purpose of depositing their spawn in the shallows; at other times they reside in the depths of the ocean, where they have a plentiful supply of food in crabs, lobsters, and other shell-fish, on which it is supposed they for the most part feed. They are occasionally found on the shores of Great Britain and Ireland, in the spawning season.

The *Lucerna* is caught in the Mediterranean sea, and is of a very curious shape; its fins about the gills being so large, and spreading so much like a fan on each side, that they appear somewhat like wings. The tail is bifid, and the scales very small. The flesh is esteemed among the Italians, and the *Lucerna* is often seen in the fish-markets of Naples, Venice, and other towns on the seashore. This fish much resembles the Father Lasher and the Gurnard; and it is called *Lucerna*, because it shines in the dark.




THE JOHN DORY. (*Zeus faber*)

It would be an inexcusable neglect to pass this fish unnoticed, not on account of its disputing with the haddock the honour of having been pressed by the fingers of the apostle, nor of its having been trodden upon by the gigantic foot of St. Christopher, when he carried on his shoulders a divine burden across an arm of the sea, but for the excellence of its flesh. It has been for some years

in such favour with our epicures, that one of them, a comedian of high repute, (Quin,) took a journey to Plymouth merely to eat this fish in perfection. Its body presents the shape of a rhomboid, but the sides are much compressed; the mouth is large, and the snout long, composed of several cartilaginous plates, which wrap and fold one over another, in order to enable the fish to catch its prey. The colour is a dark green, marked with black spots, with a golden gloss, whence the name originated. They inhabit the coasts of England, and particularly Torbay, whence they are sent to the fish-markets of London.

When the Dory is taken alive out of the water, it is able to compress its internal organs so rapidly that the air, in rushing through the openings of the gills, produces a kind of noise somewhat like that which, on similar occasions, is emitted by the gurnards.



ZEUS OPAH.

This species of the Dory is of a bright silver colour, with a cast of bluish-green on the back. Several of the last rays, both of the dorsal and anal fin, extend beyond the membrane, reaching even further than the tail itself. It has been supposed that the smaller kind of fishes may be attracted with these long flexible filaments, and mistake them for worms, while the Zeus, concealed among the sea-weeds, lies in wait for its prey. It is a native of the Indian seas.



THE COD-FISH (*Gadus morrhua*)

Is a noble inhabitant of the seas; not only on account of its size, but also for the goodness of its flesh, either fresh or salted. The body measures sometimes above three, and even four feet in length, with a proportionable thickness. The back is of a brown olive colour, with white spots on the sides, and the lower part of the body is entirely white. The eyes are large and staring. The head is broad and fleshy, and esteemed a delicious dish.

The fecundity of all fishes must be an object of the greatest astonishment to every observer of nature. In the year 1790, a Cod-fish was sold in Workington market, Cumberland, for one shilling: it weighed fifteen pounds, and measured two feet nine inches in length, and seven inches in breadth; the roe weighed two pounds ten ounces, one grain of which contained three hundred and twenty eggs. The whole, therefore, might contain, by fair estimation, three million nine hundred and four thousand four hundred and forty eggs. From such a trifle as this we may observe the prodigious value of the fishing trade to a commercial nation, and hence draw a useful hint for increasing it; for, supposing that each of the above eggs should arrive at the same perfection and size, its produce would weigh twenty-six thousand one hundred and twenty-three tons; and consequently would load two hundred and sixty-one sail of ships, each of one hundred tons burden. If each fish were brought to market, and sold as the original one, for one shilling, the produce then would be one hundred and ninety-five thousand pounds; that is to say, the first shilling would produce twenty times one hundred and ninety-five thousand, or three million nine hundred thousand shillings.

In the European seas, the Cod begin to spawn in

January, and they deposit their eggs in rough ground among rocks. Some continue in roe until the beginning of April. Cod-fish are reckoned best for the table from October to Christmas. The air-bladders, under the name of sounds, are pickled, and sold separately.

The chief fisheries for Cod are in the Bay of Canada, on the great bank of Newfoundland, and off the isle of St. Peter, and the isle of Sable. The vessels frequenting these fisheries are from a hundred to two hundred tons burden, and will each catch thirty thousand Cod, or more. The best season is from the beginning of February to the end of April. Each fisherman takes only one Cod at a time, and yet the more experienced will catch from three to four hundred in a day. It is fatiguing work, owing particularly to the intense cold they are obliged to suffer during the operation.

Cod frequently grow to a very great size. The largest that is known to have been caught in this kingdom was taken at Scarborough, in the year 1775; it measured five feet eight inches in length, and five feet in circumference, and weighed seventy-eight pounds. The usual weight of this fish is from fourteen to forty pounds.



THE HADDOCK (*Gadus aeglefinus*)

Is much less in size than the cod-fish, and differs somewhat from it in shape; it is of a bluish colour on the back, with small scales; a black line is carried on from the upper corner of the gills on both sides down to the tail; in the middle of the sides, under the line a little beneath

the gills, is a black spot on each shoulder, which resembles the mark of a man's finger and thumb; from which circumstance it is called *St. Peter's fish*, alluding to the fact recorded in the seventeenth chapter of St. Matthew: "Go thou to the sea, and cast a hook, and take up the fish that first cometh up, and when thou hast opened his mouth, thou shalt find a piece of money; that take and give unto them for me and thee." And while St. Peter held the fish with his fore-finger and thumb, it is fabled, that the skin received, and preserved to this day, the hereditary impression.

Haddocks migrate in immense shoals, which usually arrive on the Yorkshire coasts about the middle of winter. These shoals are sometimes known to extend from the shore nearly three miles in breadth, and in length from Flamborough Head to Tynemouth Castle, a distance of fifty miles; and, perhaps, even further. An idea of the number of Haddocks may be formed from the following circumstance: three fishermen, within a mile of the harbour of Scarborough, frequently loaded their boat with these fish twice a day, taking each time a ton weight of them.


The flesh of the Haddock is harder and thicker than that of the whiting, and not so good; but it is often brought upon the table, either broiled, boiled, or baked, and is by many much esteemed. The Haddocks caught on the Irish coast, near Dublin, are unusually large, and of a fine flavour, and unite to the firmness of the turbot much of its sweetness. They are in season from October to January.





THE WHITING (*Gadus Merlangus*)

Is seldom more than twelve inches in length, and of a slender and tapering form. The scales are small and fine. The back is silvery, and when just taken out of the sea, reflects the rays of light with great lustre and gloss. The flesh is light, wholesome, and nourishing; and is often recommended to sick or convalescent patients, when other food is not approved of. The Whiting is found on the coasts of England, and is in its proper season from August to February.



THE LING (*Gadus molva*)


Is from three to four feet in length, though some have been caught much larger. The body is long, the head flat, the teeth in the upper jaw small and numerous, with a small beard on the chin; it has five fins, namely, two dorsal, two pectoral and one ventral.

These fish abound on the coasts of Great Britain and Ireland, and great quantities are salted for home consumption and exportation. On the eastern coasts of England they are in their greatest perfection from the

beginning of February to the end of May. They spawn in June: at this season the males separate from the females, who deposit their eggs in the soft oozy ground at the mouth of large rivers.

In a commercial point of view, the Ling may be considered a very important fish. Nine hundred thousand pounds weight are annually exported from Norway. In England these fish are caught and cured in somewhat the same manner as the cod. Those which are caught off the shores of America are by no means so much esteemed as those which frequent the coasts of Great Britain and Norway; and the Ling in the neighbourhood of Iceland are so bad, that the inhabitants are unable to find a sale for them in any country except their own. The roe and air-bladders, or sounds of the Ling, are pickled, and sold separately.

The Hake (*Gadus merluccius*) is a coarse fish, nearly allied to the Ling, and is caught in great abundance on the Devonshire and Cornwall coast. It is also found on the coasts of Ireland and Scotland, where it is called stock-fish, and often confounded with cod.




THE SMELT, OR SPARLING. (*Osmerus eperlanus*.)

THIS fish is in length about eight or nine inches, and nearly one in breadth; the body is of a light olive green, inclining to silver white. The smell, when the fish is fresh and raw, is not unlike that of ripe cucumbers, but it goes off in the frying-pan, and the Smelt then yields a tender and most delicious food. Smelts are sea-fish, and inhabit

the sea-coast and harbours; but they are often taken in the Thames, the Medway, and other large rivers, which they ascend in the spawning season. The skin of this fish is so transparent, that with the help of a microscope, its blood may be seen to circulate.

Smelts are found on the coasts of all the northern countries of Europe, and even in the Mediterranean. They vary considerably in size. Mr. Pennant states that the largest he had ever heard of measured thirteen inches in length, and weighed half a pound.



THE MACKEREL (*Scomber Scomber*)

Is taken and well known in all parts of the world. It is usually about a foot or more in length; the body is thick, firm, and fleshy, slender toward the tail; the snout sharp, the tail forked, the back of a lovely green, beautifully variegated, or, as it were, painted with black strokes; the under part of the body is of a silvery colour, reflecting, as well as the sides, the most elegant tints of the opal and the mother-of-pearl. Nothing can be more in-

teresting and pleasing to the eye than to see Mackerel, just caught, brought on shore by the fishermen, and spread, with all their radiancy, upon the pebbles of the beach, at the first rays of the rising sun; but when taken out of their element, they quickly die.


Mackerel visit our shores in vast shoals; but, from being very tender and unfit for long carriage, they are found less useful than other gregarious fish. The usual bait is a bit of red cloth, or a piece of the tail of the Mackerel. The great fishery for them is in some parts of the south and west coasts of England; this is of such an extent as to employ, in the whole, a capital of nearly two hundred thousand pounds. The fishermen go out to the distance of several leagues from the shore, and stretch their nets, which are sometimes miles in extent, across the tide during the night. A single boat has been known to bring in, after one night's fishing, a cargo that has been sold for nearly seventy pounds. The roes of the Mackerel are used in the Mediterranean for *caviar*. In Cornwall, and also in several parts of the continent, Mackerel are preserved by pickling and salting; and in this state possess a flavour somewhat like that of the salmon. Their voracity has scarcely any bounds; and when they get among a shoal of herrings, they make such havock, as frequently to drive it away. Mackerel are in season from March to June.



THE GAR-FISH, (*Esox belone*.)

OF which the figure above is an exact representation, is of a very extraordinary form. The body, in shape and colour, is not unlike that of a mackerel; but the nose, or upper jaw, is protracted into a kind of lance, nearly as long as the rest of the body. It is vulgarly supposed that

this fish leads the phalanxes of mackerel through the regions of the deep; and, like a faithful and experienced pilot, traces their journey, points out their dangers, and conducts them to their destination. A curious singularity of this creature is, that its bones are of a bright green colour; the flesh is not so firm nor of so good a flavour as that of the mackerel, but sells well whenever it comes to market.



THE HERRING. (*Clupea harengus*.)

This fish is somewhat like the mackerel in shape, as well as in delicacy of taste, although it differs much in flavour. It is about nine or ten inches long, and about two and a half broad, and has bloodshot eyes; the scales large and roundish; the tail forked; the body of a fat, soft, delicate flesh, but more rank than that of the mackerel, and therefore less wholesome. Yet some people are so very fond of it, that they call the Herring *the King of Fishes*. They swim in shoals, and spawn once a year, about the autumnal equinox, at which time they are the best. They come into shallow water to spawn, like the mackerel; and hence they periodically visit our coasts, retiring again to the deep waters when the spawning season is over.

The fecundity of the Herring is astonishing. It has been calculated, that if the offspring of a single pair of Herrings could be suffered to multiply unmolested and undiminished for twenty years, they would exhibit a bulk

ten times the size of the earth. But, happily, Providence has so contrived the balance of nature by giving them innumerable enemies. All the monsters of the deep find them an easy prey; and, in addition to these, immense flocks of sea-fowl watch their outset, and spread devastation on all sides.

In the year 1773, the Herrings for two months were in such immense shoals on the Scotch coasts, that it appears, from tolerably accurate computations, no fewer than one thousand six hundred and fifty boat-loads were taken in Loch Torridon in one night. These would, in the whole, amount to nearly twenty thousand barrels.

This fish is prepared in different ways, in order to be kept for use through the year. The white, or pickled Herrings, are washed in fresh water, and left the space of twelve or fifteen hours in a tub full of strong brine, made of fresh water and sea-salt. When taken out, they are drained, and put in rows or layers in barrels, with salt.

Red Herrings are prepared in the same manner, with this difference, that they are left in the brine double the time above mentioned; and when taken out, placed in a large chimney constructed for the purpose, and containing about twelve thousand, where they are smoked by means of a fire underneath, made of brushwood, for the space of twenty-four hours.



THE SPRAT, (*Clupea sprattus*.)

A WELL-known fish, between four or five inches in length, the back fin very remote from the nose; the lower jaw longer than the upper, and the eyes bloodshot, like those

of the herring, to which it is nearly allied. Sprats arrive yearly in the beginning of November in the river Thames; and generally a large dish of them is presented on the table at Guildhall, on Lord Mayor's Day, November 9th. They continue through the winter, and depart in March. They are sold by measure, and yield a great deal of sustenance to poor people in the winter season. It is reported that they have been taken yearly about Easter time in a lake in Cheshire, called Kostern Meer, and in the river Mersey, in which the sea ebbs and flows seven or eight miles below the lake.

This fish is caught on the southern shores of France, where it is held in great repute; and from its abounding in the neighbourhood of the island of Sardinia, it is in that country called the Sardine. It is sent here pickled in the same way as herrings, and packed in barrels. Most naturalists, however, now consider the Sardine as a different species from the Sprat, and it is certainly much more delicate in flavour.


THE PILCHARD. (*Clupea pilchardus*.)

THE chief difference between this fish and the herring is, that the body of the Pilchard is more round and thick; the nose shorter in proportion, turning up; and the under jaw shorter. The back is more elevated, and the belly not so sharp. The scales adhere very closely, whilst those of the herring easily drop off. It is also, in general, of considerably smaller size.

About the middle of July Pilchards appear in vast shoals off the coasts of Cornwall. These shoals remain till the latter end of October, when it is probable they retire to some undisturbed deep, at a little distance, for the winter.

The Pilchard fishery is an important branch of commerce. From a statement of the number of hogsheads exported each year, for ten years, from 1747 to 1756 inclusive, from the four ports of Fowy, Falmouth, Penzance, and St. Ives, it appears that Fowy exported yearly one thousand seven hundred and thirty-two hogsheads;

Falmouth, fourteen thousand six hundred and thirty-one; Penzance and Mount's Bay, twelve thousand one hundred and forty-nine; St. Ives, one thousand two hundred and eighty-two; in all, twenty-nine thousand seven hundred and ninety-four hogsheads. Every hogshead, for ten years last past, together with the bounty allowed for exportation, and the oil made out of it, has amounted, one year with another, at an average, to the price of one pound thirteen shillings and three-pence; so that the cash paid for Pilchards exported has, at a medium, annually amounted to the sum of forty-nine thousand five hundred and thirty-two pounds. The above was the state of the fishing several years ago; at present it is still more extensive.




THE ANCHOVY. (*Engraulis encrasicolus.*)

LIKE the herring and sprat, these fish leave the depths of the open sea, in order to frequent the smooth and shallow places of the coasts for the purpose of spawning. The fishermen generally light a fire on the shore, for the purpose of attracting the Anchovies, when they fish for them in the night. After they are cleaned, and their heads cut off, they are cur'd in a particular way, and packed in small barrels for sale and exportation. Anchovies are occasionally found both in the North Sea and in the Baltic; but it is supposed that they are in much greater number in the Mediterranean than in any other part of the world. They have sometimes, though rarely, been caught in the river Dee, on the coasts of Flintshire and

Cheshire. The upper jaw of this fish is longer than the under; the back is brown; the sides silvery; fins short; the dorsal fin, opposite the ventrals, transparent; the tail fin-forked. Its length is about three inches.

THE WHITE BAIT. (*Clupea alba*.)

THIS beautiful little fish is a pure white, without spots on either side. Immense quantities are caught from the beginning of April to the end of September in the Thames; but they are so delicate, as scarcely to bear carriage, and are therefore thought best when eaten as near as possible to the place where they were taken; and hence the custom of having White-Bait dinners at the taverns at Greenwich and Blackwall. It was long supposed that the White-Bait was the fry of the shad, but it is now proved to be a distinct species.



THE TURBOT. (*Pleuronectes*, or *Rhombus maximus*.)

THE Turbot is a well-known fish, and much esteemed for the delicate taste, firmness, and sweetness of its flesh. Juvenal, in his fourth satire, gives us a very ludicrous description of the Roman emperor Domitian assembling the senate, to decide how and with what sauce this fish should be eaten. The Turbot is sometimes two feet and a half long, and about two broad. The scales on the skin are so very small that they are hardly perceptible. The colour of the upper side of the body is a dark brown, spotted with dirty yellow; the under side a pure white, tinged on the edges with a somewhat flesh-colour, or pale

pink. There is a great difficulty in baiting the Turbot, as it is very fastidious in its food; nothing can allure it but herrings, or small slices of haddocks, and lampreys; and as it lies in deep water, flirting and paddling on the ooze at the bottom of the sea, no net can reach it, so that it is generally caught by hook and line. It is found chiefly on the northern coasts of England, Scotland, and Holland.



THE PLAICE, (*Pleuronectes platessa*,)

A WELL-known English fish, nearly allied to the turbot. It has smooth sides, an anal spine, and the eyes and six tubercles are placed on the same side of the head. The body is very flat, and the upper part of the fish of a clear brown colour, marked with orange-coloured spots, and the belly white. Plaice spawn in the beginning of February; and when full-grown assume something like the shape of a turbot, but the flesh is very different, being soft and nearly tasteless.

When near the ground, they swim slowly and horizontally, but if suddenly disturbed, they change the horizontal to the vertical position, darting along with meteor-like rapidity, and then again, quickly resuming their inactive habits at the bottom of the water. Plaice feed on small fish and young crustacea, and have sometimes been taken on our coasts weighing fifteen pounds, but a fish half that weight is considered very large. The finest kind, called Diamond Plaice, are caught on the Sussex coast. These fish are in considerable esteem as food, though by no means equal to the Turbot and Sole. Those of a moderate size are reckoned the best eating.



THE FLOUNDER. (*Pleuronectes flesus.*)

THE principal distinction between the plaice and the Flounder consists in the former having a row of six tubercles behind the left eye, of which this fish is entirely destitute; it is also a little longer in the body, and, when full grown, somewhat thicker. The back is of a dark olive colour, spotted. In taste they are reckoned more delicate than the plaice. They live long after being taken out of their element, and are often cried in the streets of London, but they seldom appear on the tables of the rich and dainty. They are common in the British rivers, and in all large rivers which obey the impression of the tide, and they feed upon worms bred in the mud at the bottom of the water.



THE SOLE (*Pleuronectes solea*)

Is well known as a very excellent fish, whose flesh is firm, delicate, and of a pleasing flavour. Soles grow to the length of eighteen inches, and even more in some of our seas. They are often found of this size and superiority in Torbay, whence they are sent to market at Exeter, and several other towns in Devonshire and the

adjacent counties. They are found also in the Mediterranean and several other seas; and, when in season, are in great requisition for the most luxurious tables. The upper part of the body is brown; the under part white; one of the pectoral fins is tipped with black; the sides are yellow, and the tail rounded at the extremity. It is said that the small Soles, caught in the northern seas, are of a much superior taste to the large ones, which the southern and western coasts afford.


This fish has also the quality of keeping sweet and good for several days, even in hot weather, and is thought to acquire a more delicate flavour by being thus kept. On this account it is that Soles in the London markets are frequently more esteemed than those which are cooked immediately after they are taken out of the sea.

In the economy of flat fish we have an account of one circumstance which is very remarkable: among various other marine productions, they have been known to feed on shell-fish, although they are furnished with no apparatus whatever in their mouth for reducing these to a state calculated for digestion.



THE SALMON-PINK, BRANDLING, PAR, OR SKEGGER.

THIS brilliant little fish is the smallest of the *salmonidæ*, and is only found in rivers frequented by salmon; for whenever a river becomes deserted by them, the samlet also disappears. This fish is considered to be the fry of the true salmon, and Mr. Young, in a recent essay, has, we think, fairly established the fact; but Mr. Yarrell, and other naturalists, assert it to be a distinct species.




THE SALMON (*Salmo salar*)

Is the boast of large rivers, and one of the noblest inhabitants of the sea, if we esteem it by its bulk, colour, and the sweetness of its flesh. Salmon are found of a great weight, and sometimes measure five feet in length. The colour is beautiful, a dark blue dotted with black spots on the back, merging to silvery white on the sides, and white with a little shade of pink below. The fins are comparatively small. These fish, though they live principally in the sea, come up the rivers at the spawning season, to a considerable distance inland, where the female deposits her eggs. Soon after, both she and the male take an excursion to the vast regions of the sea, and do not visit any of the land streams again till the next year, when they return for the same purpose. They are so powerfully impelled by this natural impulse, that, if they are stopped when swimming up a river by a fall of water, they spring up with such a force through the descending torrent, that they stem it till they reach the higher bed of the stream; and on this account small cascades on the Tweed and other rivers are often called Salmon-leaps. The Salmon is in a great measure confined


to the northern seas, being unknown in the Mediterranean, and in the waters of other warm climates. The flesh is red when raw, rather paler when salted or boiled: it is an agreeable food, fat, tender, and sweet, and excels in richness all other fresh-water fish: however, it does not agree with every stomach, and is often injurious when eaten by sick persons.

In the river Tweed, about the month of July, the capture of Salmon is astonishing: often a boat-load, and sometimes nearly two, may be taken at a tide; and in one instance more than seven hundred fish were caught at a single haul of the net. From fifty to a hundred at a haul are very common. Some of these are sent to London by the steam-boats; but part are slightly salted and pickled, in which state they are called kipper. The season for fishing commences in the Tweed in February, and ends about old Michaelmas day. On this river there are about forty considerable fisheries, which extend upwards, about fourteen miles from the mouth; besides many others of less consequence. These, several years ago, were let at an annual rent of more than ten thousand pounds; and to defray this expense, it has been calculated that upwards of two hundred thousand Salmon must be caught there, one year with another. The principal Salmon fisheries in Europe are in the rivers, or on the sea-coasts adjoining the large rivers of England, Scotland, and Ireland. The chief English rivers, in which they are now caught, are the Tyne, the Trent, the Severn, and the Tweed. They were formerly found in the Thames, but none have been taken there for the last twenty years. The Salmon fry go down the river to the sea in April. A young Salmon under two pounds in weight is called a Salmon Peel, and a larger one a Grilse. Salmon cannot be eaten too fresh; and is very unwholesome when stale.



THE SALMON TROUT, (*Salmo Trutta*.)

ALSO called the *Bull Trout*, or *Sea Trout*, is thicker in the body than the common trout, and weighs about three pounds; it has a large smooth head, which, as well as the back, is of a bluish tint, with a green gloss; the sides are marked with numerous black spots; and the tail is broadest at the end. It is said that in the beginning of summer the flesh of this fish reddens, and remains this colour till the month of August; which is very probably owing to their being on the point of spawning. Like the salmon, this fish inhabits the sea; but in the months of November and December it enters the rivers, in order to deposit its roe; and consequently, in the spawning season, it is occasionally found in lakes and streams, at a great distance from the sea. It is very delicate, and much esteemed on our tables; but, as it contains a great deal of fat, it ought to be dressed as soon as possible, for it would soon become putrid. Some people prefer this fish to salmon; but they are both apt to cause illness when eaten in too great a quantity.



THE TROUT. (*Salmo fario*.)


THIS fish, in figure, resembles the salmon; it has a short roundish head, and a blunt snout. Trouts are fresh-water fish, and they breed and live constantly in rivers and small pellucid streams which sparkle over clean pebbles and beds of sand.

They feed on river flies and other water insects, and are so fond of them, and so blindly voracious, that anglers deceive them with artificial flies made of feathers, wool, and other materials, which resemble very closely the natural ones. In Lough Neagh, in Ireland, Trouts have been caught weighing thirty pounds; and we are told, that in the lake of Geneva, and in the northern lakes of England, they are found of a still larger size. It holds the first place among the river fish, and its flesh is very delicious, but difficult of digestion when old, or kept too long. They spawn in the month of December, and deposit their eggs in the gravel at the bottom of rivers, dykes, and ponds. Unlike most other fish, the Trouts are least esteemed when near spawning. They are properly in season in the months of July and August, being then fat and well-tasted.

The beautiful silvery Trout is the most voracious of fresh-water fish, and will devour every living thing which the water produces; even its own spawn in all its stages, and will lie upon the bed or hill, watching to seize its young fry, as they become vivified and rise from under their gravelly birth-place. Neither does he confine himself to any given sort of fish, but luxuriates his rapacious stomach upon all the varieties, from instinct, occasionally changing his food to larvæ, caddis, ephemera, worms, and even the young of the water-snail, all of which act as alteratives. Owing to his large fins and broad tail, his movements are extremely rapid, and from his muscular power and pliability he seldom misses his prey. His habits are solitary, being only accompanied by one, and that at some distance from him, in the summer season; and as the autumn approaches, when larvæ &c., are diminishing, he keeps entirely alone, until the pairing season returns. The period of spawning differs in various rivers from natural causes, such as snow, cold rains, or inclement weather; for, as Trout, like salmon, spawn on gravel beds in shallow water, the cold readily affects them. When they cannot reach the spot prepared for the deposit of their eggs, they frequently abstain from spawning for yeeks. The younger Trout generally hill, as it is termed, earlier than those of larger growth. They begin to throw up their bed early in December, when the female and male may be seen working together, the former mostly in advance. By constant labour they dig a hollow in the gravel, throwing it up on each side, and at last forming a heap, which is called a hill, or bed. At this period they are very shy and stupid, and even the shadow of a cloud will frighten them from their hill, when they retreat into deeper water; but upon finding all quiet they return. This preparation generally occupies two or three weeks; and frequently the hill is shared both in labour and occupation by several pairs of Trout; it often measures many feet in diameter, and is two or three feet higher than the bed of the stream. From the middle of December to the end of January the Trout is in full spawning operation; when the fish deposit their eggs in the hollow, and afterwards work the gravel over them to the depth of about three inches. If the temperature of the water is not altered

during the period of incubation, the young make their appearance on the fiftieth day; never earlier, frequently later. Nature has endowed the young fry with so much instinct of self-preservation, that for many days they keep under the gravel, and it is curious to see the shoal hiding together under large stones to protect themselves from danger: this they continue to do until the egg-shell, in which they remain partially enveloped, falls off from their delicate frames. This shell, which adheres to them for fourteen days, contains a proportion of fluid necessary for their support during this period of helplessness. After this they resort to the shallows and scours to avoid the larger fish, where they remain solitary for a year, during which time, in good keep, they attain the weight of three to four ounces; the second year, eight to ten ounces, after which they begin to breed. Fish, like every animal, becomes fat when it has abundance of food with little or no exertion; so that the growth is entirely regulated by the relative proportion of food and labour. I have observed this difference in the same brood of Trout, artificially bred upon my system: the one brood being placed in water well supplied with food, the other in a spring-stream where little food existed; the former, at ten months old, were four inches long, and three and a half ounces in weight, while the latter were only an inch and a half long, and less than an ounce in weight. Although Trout are not migratory, yet, when they become large, they run up stream to purer water. The small Trout are carried down the stream, against their habit, by the flushes of water or floods during the autumn months, being unable to stem the thickened torrent which fills their gills with alluvial deposit, and hinders their respiration, whence they become weak and sickly. In this state of water all fish sicken more or less, and it destroys vast numbers in the very young state. I have known thousands destroyed by the overflowing of a river, as well old as young. The cause of all our rivers falling off in the quantity of fish, is from the increasing impurity of the water, as fish especially require pure water.


The above interesting notice of the Trout has been communicated to the publisher by MR. BOCCIS, who devotes himself professionally to the increase of fish in rivers and ponds, and has performed marvels.



THE CHARR, OR ALPINE TROUT,

(*Salmo salvelinus*,)

Is not unlike the trout; the scales are very small; the colour of the body marked with numerous spots and points of black, red, and silver, mixed with yellow, and without a circle; the back tinged with olive-green; the belly white, the snout bluish. All the fins, except those of the back, are reddish, and the adipose one is red on its edge. This fish is about twelve inches in length, and is esteemed very delicate as an article of food, especially by the Italians. It is abundant in the Lago di Garda, near Venice; and is also found, not only in our northern lakes in Westmoreland and Scotland, but also in the large sheets of water at the foot of the mountains in Lapland. The potted Char enjoys a high and deserved reputation in several parts of the continent, as well as in England. The Char is a fresh-water fish, and is generally found in the deepest parts of lakes; it is never taken by the angler, only by the net.




THE GRAYLING. (*Salmo thymallus.*)

THIS fish never exceeds fifteen inches in length, and seldom arrives at three pounds weight. The back and sides are of a silvery grey, and when the fish is first taken out of the water, slightly varied with blue and gold. The coverts of the gills are of a glossy green, and the scales are large.

The Grayling is a fresh-water fish, and delights chiefly in clear and not too rapid streams, where it affords great amusement to the angler, as it is very voracious, and rises eagerly to the fly. They are bolder than trout, and even, if missed by the hook several times successively, they will still pursue the bait. They feed principally on worms, insects, and water-snails; and the shells of the latter are often found in great quantity in their stomachs. They spawn in the months of April and May. The largest fish of this species ever heard of was one caught in the Severn, and weighed five pounds.

Ancient writers strongly recommended this fish as food for sick persons, as they considered it peculiarly wholesome and easy of digestion.



THE PIKE. (*Esox lucius.*)

THE body of this fish is a pale olive-grey, deepest on the back, and marked on the sides by several yellowish spots or patches; the abdomen white, slightly spotted with black; its length is from one to eight feet, and its weight from one or two to forty or fifty pounds. The flesh is white and firm, and considered very wholesome; the larger and older it is, the more it is esteemed. There is scarcely any fish of its size in the world that in voracity can equal the Pike.* It lives in rivers, lakes, and ponds; and in a confined piece of water will soon destroy all other fish, as it generally does not feed upon anything else, and often swallows one nearly as big as itself; for through its greediness in eating, it takes the head foremost, and so draws it in by little and little at a time, till it has swallowed the whole. A gudgeon of good size has been found in the stomach of a large Pike, the head of which had already received clear marks of the power of digestion, whilst the rest of the fish was still fresh and unimpaired.

"I have been assured (says Walton) by my friend Mr. Seagrave, who keeps tame otters, that he has known a Pike, in extreme hunger, fight with one of his otters for a


* Mr. Boccius has, however, shown that the Trout is even more voracious.

carp that the otter had caught, and was then bringing out of the water."

Boulker, in his *Art of Angling*, says, that his father caught a Pike, which he presented to Lord Cholmondeley, that was an ell long, and weighed thirty-six pounds. His lordship directed it to be put into a canal in his garden, which at that time contained a great quantity of fish. Twelve months afterwards the water was drawn off, and it was discovered that the Pike had devoured all the fish, except a large carp that weighed between nine and ten pounds, and even this had been bitten in several places. The Pike was again put in, and an entire fresh stock of fish for him to feed on: all these he devoured in less than a year. Several times he was observed by workmen who were standing near, to draw ducks and other water-fowl under water. Crows were shot and thrown in, which he took in the presence of the men. From this time the slaughtermen had orders to feed him with the garbage of the slaughter-house; but being afterwards neglected, he died, as is supposed, from want of food.

In December, 1765, a Pike was caught in the river Ouse, that weighed upwards of twenty-eight pounds, and was sold for a guinea. When it was opened, a watch with a black riband and two seals were found in its body. These, it was afterwards found, had belonged to a gentleman's servant, who had been drowned in the river about a month before.


The Pike is a very long-lived fish. In the year 1497, one was caught at Heilbrun in Swabia, to which was affixed a brazen ring, with the following words engraved on it in Greek characters: "I am the fish, which was first of all put into this lake, by the hands of the governor of the universe, Frederick the Second, the fifth of October, 1230."



THE PERCH (*Perca fluviatilis*)

SELDOM grows to any great size; yet we have an account of one which is said to have weighed nine pounds. The body is deep, the scales rough, the back arched, and the side-lines placed near the back. For beauty of colours, the Perch vies with the gaudiest inhabitants of the waters; the back glows with the deep reflections of the brightest emeralds, divided by five broad stripes; the abdomen imitates the tints of the opal and mother-of-pearl; and the ruby hue of the fins completes an assemblage of colours most harmonious and elegant. It is a gregarious fish, and is caught in several rivers of these islands; the flesh is firm, delicate, and much esteemed.

It is generally believed that a pike will not attack a full-grown Perch: he is deterred from so doing by the spiny or dorsal fin on the back, which this fish always erects at the approach of an enemy. Perch are so voracious, that, if an expert angler happens to find a shoal of them, he may catch every one. If, however, a single fish escape that has felt the hook, all is over; as this fish becomes so restless, as soon to occasion the whole shoal to leave the place. Perch are so bold, that they are generally the first fish caught by a young angler; they will also soon learn to take bread thrown into the water to feed them. A large-sized Perch weighs about three pounds; but generally the Perches caught in ponds do not exceed eight or ten ounces in weight.




THE POPE, OR RUFFE.

THE Pope is very like a small perch, but with a curiously formed single dorsal fin: the colour of the back is a dusky olive green; the sides light brownish green and copper colour; and small brown spots are spread over the dorsal fin, the back, and tail. The pectoral, ventral, and anal fins are pale brown. This fish rarely exceeds six inches in length; but it is nearly as good as a perch of the same size; which it resembles, both in its haunts and habits: it spawns in April, and feeds on small fry, worms, or aquatic insects.

Cuvier assigns the credit of the first discovery of this fish to an Englishman of the name of Caius, who found it in the river Yare, near Norwich, and called it *Aspredo*, a translation of our name Ruffe, (rough,) which is well applied to it, on account of the harsh feel of its denticulated scales.




THE CARP (*Cyprinus carpio*)

Is famous for the sweetness of its flesh, when of moderate size, that is, when measuring about twelve to fifteen inches in length, and weighing about three pounds. The scales are large, with a golden gloss upon a dark green ground. These fish sometimes grow to the length of three or four feet, and contain a great quantity of fat. The soft roe of the Carp is esteemed a great delicacy among epicures. In the canals of Chantilly, formerly the seat of the Prince of Condé, Carps have been kept for above one hundred years, most of them appearing hoary through old age, and so tame that they answered to their names when the keeper called them to be fed. This fish has large molar teeth only, situate at the back part of the head or throat, and a broad tongue; the tail is widely spread as well as the fins, which are inclined to a reddish tint. Carp that live in rivers and running streams are preferred for the table, as those which inhabit pools and ponds have generally a muddy and disagreeable taste. Though so cunning in general as to be called the River Fox, yet at spawning time they suffer themselves to be tickled and caught without at-

tempting to escape. It is said that Carp were first brought to England about three hundred years ago. They are very tenacious of life, and at the inns in Holland are often kept alive a month or six weeks, by being fed with bread and milk, and laid on wet moss in a net, which is hung from the ceiling in an airy place. The moss is kept moist, and water is thrown over the fish twice a day.

Carp is always considered a delicacy for the table, especially when stewed in port wine; and it appears to have been long held in high estimation on that account, as we find, from the privy purse expenses of Henry VIII., that the bluff king was exceedingly fond of Carp.

The Tench, (Cyprinus tinca,) like the carp, is remarkably tenacious of life. Its body is thick and short, and seldom exceeds twelve inches in length, or four pounds in weight. The eyes are red; the back, dorsal, and ventral fins, dusky; the head, sides, and abdomen of a greenish hue, mixed with gold; and the tail very broad. The Tench delights in still water, in the muddy parts of ponds, where it is the most secure from the voracious ramblings and fierce attacks of the tyrant pike, and from the hook of the angler; here it lives nearly motionless, lurking beneath flags, reeds, and weeds. This inactive life has enabled some individuals of this species to attain an extraordinary bulk. We have read, as a well-authenticated fact, that in the northern part of England, in a piece of water, which having been long neglected, was filled with timber, stones, and rubbish, two hundred Tench, and as many perch of good size were found; and that one fish in particular, which seemed to have been shut up in a nook, had not only surpassed all the others in size, but had also taken the form of the hole in which it had been accidentally confined. The body was in the shape of a half-moon, conforming in the convexity of its outlines to the concavity of the dungeon where this innocent sufferer had been immured for a number of years, it weighed eleven pounds.



THE GOLD-FISH, OR GOLDEN CARP,
(*Cyprinus auratus*,)

WAS originally brought from China, and first introduced into England in 1661, but is now become quite common and will breed as freely in ponds as the carp. The average size is about five inches, and it scarcely ever exceeds seven and a half. Gold-fish are highly prized in China, and are extensively introduced in the ornamental waters of our own country. Nothing is more pleasing than to see them glide along and play in the transparent crystal, whilst their broad and glittering scales reflect the rays of the sun. They are often kept within the small compass of a glass bowl, where they become tame and docile, and after a short time seem to recognise their feeders.

The smallest fish are preferred, not only from their being the most beautiful, but because a greater number of them can be kept in a small circumference. These are of a fine orange red colour, appearing as if sprinkled over with gold-dust. Some, however, are white, like silver; and others, white spotted with red.


When Gold-fish are kept in ponds, they are often taught to rise to the surface of the water at the sound of a bell, to be fed.



THE GUDGEON, (*Cyprinus gobio*,)

A WELL-known fresh-water fish, generally found in gentle streams, on gravelly scours: it is between five and six inches in length. The back is brown, the abdomen white,

and the sides tinged with red; the tail is forked. Gudgeons spawn in summer, and feed upon worms and aquatic insects. Their flesh is white, of excellent flavour, and easy of digestion. In the months of September and October, these fish are taken in the rivers of some parts of the continent in great abundance; and the markets are well supplied with them. They are caught both in nets, and with hooks and lines.



THE CHUB (*Cyprinus cephalus*)

Is of a coarse nature, and full of bones; it seldom exceeds the weight of five pounds. The body is of an oblong shape, nearly round; the head, which is large, and the back are of a deep dusky green; the sides silvery, and the abdomen white; the pectoral fins are of a pale yellow, the ventral and anal ones red; and the tail brown, tinged with blue at its extremity, and slightly forked. This fish frequents the deep holes of rivers, but in the summer, when the sun shines, it rises to the surface, and lies quiet under the shade of the trees, that spread their foliage on the verdant banks; but yet, though it seems to indulge itself in slumber, it is easily awakened, and at the least alarm dives rapidly to the bottom. Although a leather-mouthed fish, it takes every species of food, including small fish, the same as a trout, though not so voracious. In March and April this fish may be caught with large red worms; in June and July with flies, snails, and cherries; in August and September with cheese pounded in a mortar, mixed with saffron and butter. When the Chub seizes a bait, it bites so eagerly, that its jaws are often heard to chop like those of a dog. It, however, seldom breaks its hold, and, when once struck, is soon tired.



THE BARBEL. (*Cyprinus barbus*.)

THE Barbel is readily distinguished from all other fresh-water fish by the barbs or wattels attached to its mouth. Its upper jaw is very considerably extended beyond the lower jaw. The Lea, the Thames, and various other rivers in the neighbourhood of London abound in this fish, which affords excellent sport to the angler. "During summer," says Mr. Gorrell, "this fish, in shoals, frequents the weedy parts of the river; but as soon as the weeds begin to decay in autumn, it seeks the deeper water, and shelters itself near piles, locks, and bridges, which it frequents till the following spring." It is sometimes found to weigh from fifteen to eighteen pounds, and to measure three feet in length, but its usual length is from twelve to eighteen inches. The flesh is coarse and unsavoury, and held in no estimation.



THE DACE (*Cyprinus leuciscus*)

RESEMBLES the chub in its form, but is smaller, and of a lighter colour; it is gregarious and remarkably prolific. It is seldom more than ten inches in length; the back is of a dusky colour, tinged with yellow and green, and the sides have a silvery cast.

Dace spawn in March, and are in season about three weeks afterwards. They improve, and are good about

Michaelmas; but in February they are best. The flesh is, however, at all times woolly and insipid. They are very lively creatures, and, if kept in ponds, may live a considerable time.




THE ROACH (*Cyprinus rutilus*)

BELONGS also to the carp family, and is remarkable for its numerous progeny. It is a deep yet thin made fish, in shape somewhat resembling the bream, but approaching the carp in the breadth and shape of its scales, which are large and deciduous. The soundness of the flesh is become proverbial, and pleases the taste by a peculiar delicacy of flavour. The ventral fins are, like those of the perch, of a bright crimson, and the irides of the eye sparkle like rubies and garnets. The length of the Roach is commonly between nine and ten inches, but sometimes much greater.



THE BLEAK (*Cyprinus alburnus*)

Is nearly allied to the roach. It is a small glittering fish, familiar to most persons from its playing about on warm summer evenings on the surface of rivers in chace of flies, bread-crumbs, &c. The scales are employed in making artificial pearls.



THE BREAM (*Cyprinus brama*)

Is a flattish fish, not unlike the carp in several points, but much broader in proportion to its length and thickness. Its head is truncated, the upper jaw a little projecting; the forehead a bluish black; cheeks yellowish; body olive, paler below; fins obscure, with an oblong conical process at the base of the ventral fins; twenty-nine rays in the anal fin; its length is about two feet. The scales are large and of a bright colour; the tail has the form of a crescent. It frequents the deepest parts of rivers, lakes, and ponds. These fish spawn in May, secluding themselves at that time so carefully in the ooze at the bottom of the water, that they are seldom found with either soft or hard roe in them, so that in some countries the name is often used to denote sterility. The flesh is not comparable to that of the carp.

The white Bream never exceeds a pound in weight, and is consequently much smaller than the common or Carp Bream, which frequently weighs seven or eight pounds.

In some of the lakes of Ireland great quantities of Bream are taken, many of them of very large size, sometimes weighing as much as twelve or even fourteen pounds each. A place conveniently situated for the fishing is baited with grain, or other coarse food, for ten days or a fortnight regularly, after which great sport is usually obtained. The party frequently catch several hundred weight, which are distributed among the poor of the vicinity, who split and dry them with great care, to eat with their potatoes.



THE MINNOW. (*Cyprinus phoxinus.*)

THE body of the Minnow is of a blackish green, with blue and yellow variegations; the abdomen silvery; scales small; ten rays in the ventral, anal, and dorsal fins; tail forked, and marked near the base with a dusky spot. Its length is about three inches.

This beautiful and well-known fish is gregarious, and is frequent in clear gravelly streams and rivulets in many parts of Europe. In Britain it appears in March, and is seldom seen after October. It spawns in June, and is, indeed, found in roe during the greater part of the summer. It is easily tamed; and, in captivity, may be taught to pick flies or filaments of beef from the hand.

The flesh of the Minnow is extremely delicate, but the fish is so small that it would take a great number to make a dish, and consequently it is seldom used for human food. Its chief value is as a bait for catching other fish. In some parts of England it is so abundant, as sometimes to be used as manure.




THE BULL-HEAD, OR MILLER'S THUMB,

(*Cottus gobio,*)

Is found in clear brooks and rivers in most parts of

Europe. It is from four to five inches long, the head is large in proportion to the body, broad and depressed; the gill fins round, and beautifully notched. The mouth is large and full of small teeth, the general colour of the body is a dark brownish black. This fish is remarkably stupid, and may be caught with ease by the most inexperienced angler, even with a bent pin and coarse thread. Its hiding places are among loose stones, under which the peculiar flattened form of its head enables it to thrust itself. Its popular name appears to have suggested itself from the resemblance the head of the fish is supposed to bear to the form of a miller's thumb, the peculiar conformation of which is produced by his mode of testing samples of meal.



THE ELECTRICAL EEL. (*Gymnotus electricus*.)

This very remarkable fish is about three feet in length, and twelve inches in circumference, in the thickest part of the body. The head is broad, flat, and large; the mouth wide and destitute of teeth; the rostrum obtuse and rounded; the eyes small and of a bluish colour; the back of a darkish brown, the sides grey, and the abdomen of a dingy white. Across the body there are several annular divisions, or rather ridges of the skin, which give the fish the power of contracting or dilating itself at pleasure. It is able to swim backwards as well as forwards.

Mr. Bryant mentions an instance of the shock from one


of these fish being felt through a considerable thickness of wood. One morning, while he was standing by, as a servant was emptying a tub, in which an Electrical Eel was contained, he had lifted it entirely from the ground, and was pouring off the water to renew it, when he received a shock so violent as occasioned him to let the tub fall. He then called another person to his assistance, and they lifted up the tub together, each laying hold only on the outside. When they were pouring off the remainder of the water, they each received a shock so smart that they were compelled to desist.

Persons have been knocked down with a stroke. One of these fish having been taken from a net and laid upon the grass, an English sailor, notwithstanding all the persuasions that were used to prevent him, would insist on taking it up; but the moment he grasped it, he dropped down in a fit; his eyes were fixed, his face became livid, and it was not without difficulty that his senses were restored. He said that the instant he touched it "the cold ran swiftly up his arm into his body, and pierced him to the heart."

Humboldt tells us that when the Indians wish to catch these Eels they drive some wild horses through the pools which the fish inhabit; and that when the Eels have exhausted their electrical power upon the horses, the Indians take them without difficulty. He relates an instance in which he says that the horses, stunned with the shocks they received, sank under water, but most of them rose again, and gained the shore, where they lay stretched out on the ground, apparently quite exhausted and without the power of moving, so much were they stupified and benumbed. In about a quarter of an hour, however, the Eels appeared to have exhausted themselves, and instead of attacking fresh horses that were driven into the pond, fled before them. The Indians then entered the water and caught as many fish as they liked.*

This most singular fish is peculiar to South America, where it is found only in stagnant pools, at a great distance from the sea.

* See a very animated account of the capture of this fish, in Humboldt's *Views of Nature*.—*Bohn's Edition*, page 16.



THE EEL. (*Muræna Anguilla.*)

THE Eel resembles a serpent in its form, though no two animals can be more different in every other respect. Eels are fresh-water fish; but as they are very susceptible of cold, those which inhabit rivers go down every autumn towards the sea, which is always warmer than a river, and return in spring. Those that live in ponds seek the deep water for their winter-quarters, and sometimes bury themselves in the mud at the bottom. They are very tenacious of life, and will live for a long time out of water; they are even sometimes found on the grass, passing from one pond to another, in search, as it is said, of food.

They are voracious feeders, eating frogs, snails, and other molluscous animals, worms, the fry of fishes, and the larva of various insects, as well as grass and aquatic weeds. Eels are ovo-viviparous; that is, their young are produced alive, but hatched before birth from eggs.

The common Eel often weighs upwards of twenty pounds. The flesh is tender, soft, and nourishing, but does not agree with all stomachs.

The *Conger*, or *Sea Eel*, is very large and thick. Its body is dusky above, and silvery below; the dorsal and

anal fins are edged with black; and the lateral line is dotted with white. Its flesh is firm, and was much esteemed by the ancients.

The voracity of the Conger Eel is very great, and it is one of the most powerful enemies with which the fishermen of the British islands have to contend. Being usually caught by a hook and line, it requires some care to land and kill the large ones without danger. We are informed that, on such occasions, they have been known to entwine themselves round the legs of a fisherman, and fight with the utmost fury. A Conger, six feet in length, was caught in the Wash at Yarmouth, in April, 1808; but not without a severe contest with the man who had seized it. The animal is stated to have risen half erect, and to have actually knocked the fisherman down before he could secure it. This Conger weighed only about sixty pounds: but some of the largest exceed even a hundred-weight.


THE LAMPREY. (*Petromyzon marinus*.)

THE Lamprey belongs to the last family of cartilaginous fishes, and is the lowest in the scale of vertebrated animals. It grows to the length of about three feet, although the British species, with which we are best acquainted, seldom exceeds twelve inches. To avoid the constant muscular exertion necessary to prevent their being carried away by the current, they attach themselves by the mouth to stones or rocks, and hence are called *Petromyzon*, Stone-suckers. The Lamprey, although no longer maintaining its ancient repute, is still considered a delicacy; those taken in the Severn being preferred to all others. Henry the First, as is well known, died of a surfeit of them; and in the reign of Henry the Fourth their importation was encouraged by immunities. The Roman epicures prized this fish so highly, that they bestowed the utmost care, and expended enormous sums in rearing them. Pliny tells us that Lucullus formed a fish-pond of such extent, that the fish it contained were, at his death, sold for four million sesterces.

BOOK IV.

REPTILES.

§ 1. *Serpents or Ophidian Reptiles.*



SERPENTS.

SERPENTS are characterized by an elongated body, clothed in scales and destitute of limbs, but furnished with a tail. They move by lateral undulations of the body; and in this manner they glide with equal ease along the bare ground, through entangled thickets or water, and up the trunks of trees. They possess the power of fasting a great length of time, and when they feed, always swallow their prey whole, which they are enabled to accomplish by their faculty of dilating their bodies to an enormous size. This power is carried to such an extent that a Boa Constrictor can swallow a bullock whole, suffering no other inconvenience than that of lying in a state of torpor, while digestion is proceeding. Serpents generally roll themselves up when in a state of repose, with the head in the centre; and when disturbed raise the head, before they uncoil the body. They also frequently raise themselves upright, supporting themselves on the tail. The Serpent is often made a subject of poetry; and as it was the form adopted by the arch fiend to seduce Eve, it is generally considered the emblem of insinuation and flattery.

“ ————— on his rear,

Circular base of rising folds that tower'd
Fold above fold, surprising maze, his head
Crested aloft, and carbuncle his eyes.

With burnish'd neck of verdant gold, erect
 Amidst his circling spires that on the grass
 Floated redundant; pleasing was his shape
 And lovely Oft he bow'd
 His turret crest and sleek enamell'd neck,
 Fawning, and lick'd the ground whereon she trod."

PARADISE LOST.

The ancients paid great honours to Serpents, and sometimes called them good genii: they frequented sepulchres and burying-places, and were addressed like the tutelary divinities of these places. We read, in the fifth book of the *Æneis*, that, when the Trojan hero sacrificed to his father's ghost, a serpent of this kind made his appearance,

" ——— and from the tomb began to glide
 His huge bulk on seven high volumes roll'd;
 Blue was his breadth of back, and streak'd with scaly gold,
 Thus riding on his curls he seem'd to pass
 A rolling fire along, and sifge the grass;
 More various colours through his body run,
 Than Iris when her bow imbibes the sun.
 Between the rising altars and around,
 The sacred monster shot along the ground;
 With harmless play among the bowls he pass'd,
 And with his lolling tongue assay'd the taste:
 Thus fed with holy food, the wondrous guest
 Within the hollow tomb retired to rest." DRYDEN.

This animal was exalted to the honour of being an emblem of prudence, and even of eternity; and is often represented as the latter in Egyptian hieroglyphics, biting his tail, so as to form a circle. Serpents are very numerous in Africa; and Lucan, in his *Pharsalia*, gives us a very extraordinary account of the different species, which he seems to have drawn partly from ancient Greek authors, partly from actual traditions. He says,

" Why plagues like these infect the Libyan air;
 Why deaths unknown in various shapes appear;
 Why, fruitful to destroy, the cursed land
 Is temper'd thus by Nature's secret hand;
 Dark and obscure the hidden cause remains,
 And still deludes the vain inquirer's pains."

ROWE'S LUCAN.

Serpents differ very much in size. We are told of Serpents in the isle of Java measuring fifty feet in length; and in the British Museum there is a skin of one thirty-two feet long. Though all Serpents are amphibious, some are much fonder of the water than others; and, though destitute of fins or gills, remain at the bottom or swim along the surface with great ease. From their internal structure they are well adapted for either element, and their blood is as capable of circulating at the bottom of ponds as that of the frog or the tortoise. Some Serpents are not voracious, but others are so to an extraordinary degree.



THE VIPER, OR ADDER, (*Vipera berus*,)

Is a venomous species of serpent that seldom exceeds the length of two or three feet, and is of a dull yellow colour with black spots, the abdomen being entirely black; the head is nearly in the shape of a lozenge, and much thicker than the body. The Viper is viviparous, yet it is


ascertained that the eggs are formed, though they are hatched in the body of the mother.

The Reverend Mr. White of Selborne, in company with a friend, surprised a large female Viper, as she lay on the grass, basking in the sun, which seemed very heavy and bloated. As Vipers are so venomous that they should be destroyed, they killed her; and afterwards, being curious to know what made her so large, they opened her, and found in her abdomen fifteen young ones, about the size of full-grown earth-worms. This little fry issued into the world with the true Viper spirit about them, showing great alertness as soon as they were disengaged from the body of their parent. They twisted and wriggled about, set themselves up, and gaped very wide when touched with a stick; exhibiting manifest tokens of menace and defiance, though as yet no fangs could be discovered, even by the help of glasses.

Vipers attain their full growth in seven years; they feed on frogs, toads, lizards, and other animals of that kind, and it is even asserted that they catch mice and small birds, of which they seem very fond. They cast their skin every year. The teeth of the Viper are surrounded with a small bladder containing poison. There is no doubt but this poison, which appears to have been infused into the jaws of the Viper and other serpents by Providence, as a means of revenge upon their enemies, is so harmless to the animal itself, that when swallowed by it it only serves to accelerate its digestion; and, indeed, if the venom were hurtful to it, how could it masticate frogs and mice without breaking the baneful bladder, and being injured by the poisonous liquor? The Viper is very patient of hunger, and may be kept more than six months without food. When in confinement, it refuses all sustenance, and the sharpness of its poison decreases in proportion: when at liberty, it remains torpid throughout the winter; yet, when confined, it has never been observed to take its annual repose.

The Viper is a native of many parts of this island, chiefly the dry and chalky counties. Its flesh was formerly used for broth, and much esteemed in medicine, particularly to

restore debilitated constitutions. It was also used as a cosmetic, being supposed to render the complexion fair. It was probably, that from the use made by the ancients of this animal in medicine, Esculapius is represented with a serpent. The best remedy against the bite of the Viper is to suck the wound, which may be done without danger, and after this to rub it with sweet oil, and poultice it with bread and milk.



THE HORNED VIPER. (*Vipera cerastes.*)

THIS species of Viper is nearly allied to the asp, and has a pointed and solid horny substance on each eyelid, formed of two projecting scales: its body is of a pale yellowish or greyish colour, with distant sub-ovate transverse brown spots; and in length it is from one to two feet. It appears to be in some measure domesticated in Egypt, where it enters houses when the family are at table, picks up the crumbs from the floor, and retires without doing any injury to the company.

This species is found in the deserts of Africa, and is often mentioned by the ancients. Pliny tells us that "the serpent Cerastes hath many times four small horns, standing out double; with moving whereof she amusethe the birds, and traineth them unto her for to catch them, hiding all the rest of her body."



THE ASP. (*Coluber aspis*.)

THE Asp is well known as having poisoned Cleopatra, but naturalists do not appear quite certain what animal was meant by the ancients when they spoke of the Asp; some suppose it to have been the Aspic of the Mediterranean, (*Vipera aspis*), a kind of viper; but this creature is not found in Egypt; and others refer it to the haje, (*Coluber haje*), which is closely allied to the *cobra di capello*, or hooded snake of India, the most poisonous of all the serpent family.

The Haje, or Egyptian Asp, is from three to six feet in length; it has two teeth longer than the rest, through which the venom flows. The body is covered with small round scales, and is of a greenish colour, bordered with brown; its neck is capable of inflation. The jugglers of Egypt, by pressing this Asp on the nape of the neck with the finger, throw the animal into a kind of catalepsy, which renders it stiff and immoveable; when they say

that they have changed it into a rod. The habit which this species has of raising itself up when approached, induced the ancient Egyptians to believe that it guarded the fields where it was found; and it is sculptured on the gates of their temples as an emblem of the protecting divinity of the world.

The *Hooded Serpent*, or *Cobra di Capello*, called by the Indians the *Nagao*, is from three to eight feet long, with two long fangs hanging out of the upper jaw. It has a broad neck, and a mark of dark brown on the forehead; which, when viewed frontwise, looks like a pair of spectacles; but behind, like the head of a cat. The eyes are fierce and full of fire; the head is small, and the nose flat, though covered with very large scales, of a yellowish ash-colour; the skin is white, and the large tumour on the neck is flat and covered with oblong smooth scales. This serpent is extremely dreaded by the British residents in India, as its bite has hitherto been found to be incurable, and the sufferer generally dies in half an hour.


Of this genus are the dancing-snakes, which are carried in baskets throughout Hindoostan, and procure a maintenance for a set of people, who play a few simple notes on the flute, with which the snakes seem much delighted, and keep time by a graceful motion of the head; erecting about half their length from the ground, and following the music with gentle curves, like the undulating lines of a swan's neck. It is a well-attested fact, that, when a house is infested with these snakes, and some others of the coluber genus, which destroy poultry and small domestic animals, as also by the larger serpents of the boa tribe, the musicians are sent for; who, by playing on a flageolet, find out their hiding places, and charm them to destruction: for no sooner do the snakes hear the music, than they come softly from their retreat, and are easily taken. I imagine these musical snakes were known in Palestine, from the Psalmist comparing the ungodly to the deaf adder, which stoppeth her ears, and refuseth to hear the voice of the charmer, charm he never so wisely.



THE SNAKE (*Natrix torquata*, or *Coluber natrix*)

Is the largest of all English serpents, sometimes exceeding four feet in length. The colour of the body is variegated with yellow, green, white, and regular spots of brown and black. They seem to enjoy themselves when basking in the sun, at the foot of an old wall. This animal is perfectly innocuous, although many reports have been circulated and believed to the contrary; it feeds on frogs, worms, mice, and various kinds of insects, and passes the greater part of the winter in a state of torpidity. In the spring they re-appear, and at this season uniformly cast their skins. This is a process that they also seem to undergo in autumn. Mr. White says: "About the middle of September we found in a field, near a hedge, the slough of a large snake, which seemed to have been newly cast. It appeared as if turned wrong side outward, and as if it had been drawn off backward, like a stocking or a woman's glove. Not only the whole skin, but even the scales from the eyes were peeled off, and appeared in the slough like a pair of spectacles. The reptile, at the time of changing his coat, had entangled himself intricately in the grass and weeds, in order that the friction of the stalks and blades might promote this curious shifting of his exuvia."

The snake of warmer countries is not so harmless as that of this island, as will be easily perceived by what will follow.



THE RATTLE-SNAKE (*Crotalus horridus*)

Is a native of the new world, and grows to five and sometimes to six feet in length, and is nearly as thick as a man's thigh; it is not unlike the viper, having a large head and small neck, and inflicting a very dangerous wound. Over each eye is a large pendulous scale, the use of which has not yet been ascertained; the body is scaly and hard, variegated with several different colours. The principal characteristic of this justly dreaded serpent is the rattle, a kind of instrument resembling the curb-chain of a bridle, at the extremity of the tail; it is formed of thin, hard, hollow bones, linked together, and rattling on the least motion. When disturbed, the creature shakes this rattle with considerable noise and rapidity, striking terror into all the smaller animals, which are afraid of the destructive venom that this serpent communicates to the wounded limb with his bite. The wound the Rattle-snake inflicts, through the uncommon sharpness and rapid fluency of the poison, generally terminates the torment and life of the unhappy victim in the course of six or seven hours.

A Snake of this kind exhibited in London at a menagerie of foreign animals, in the year 1810, wounded a carpenter's hand, who was repairing its cage, and seeking for his rule: the man suffered the most excruciating pain, and his life could not be saved, although medical assistance was immediately applied, and every effort made to prevent the dire effect of the poison. The proprietor was condemned to pay a deodand for the injury done by the serpent.



THE AMPHISBÆNA. (*Amphisbæna fuliginosa.*)

THIS name is now applied only to a genus of South American reptiles, which are of a harmless nature, being destitute of those fangs which prepare the venom in poisonous serpents. The head is so small, and the tail so thick and short, that at first sight it is difficult to distinguish one from the other; and this circumstance, united to the animal's habit of proceeding either backwards or forwards as occasion may require, gave rise to the supposition throughout the native regions of the Amphisbæna, that it had two heads, one at each extremity, and that it was impossible to destroy one by simple cutting, as the two heads would mutually seek one another, and reunite! The ancients gave the name of Amphisbæna to what they considered a two-headed serpent; but it is not known with certainty which of the serpent tribe they meant, as their amphisbæna is described by Lucan as venomous, though in his lines elegance of language, beauty of versification, and liveliness of fancy, have perhaps a greater claim than truth to the admiration of the reader.

“ With hissings fierce, dire Amphisbænas rear
 Their double heads, and rouse the soldier's fear.
 Eager he flies: more eager they pursue;
 On every side the onset quick renew!
 With equal swiftness face or shun the prey,
 And follow fast when thought to run away.
 Thus on the looms the busy shuttles glide,
 Alternate fly, and shoot at either side.”



THE BOA CONSTRICTOR.

THIS immense animal is often twenty feet in length, and sometimes even thirty-five; the ground colour of its skin is yellowish grey, on which is distributed, along the back, a series of large chain-like, reddish brown, and sometimes perfectly red, variegations, with other smaller and more irregular marks and spots. It is a native of South America, where it chiefly resides in the most retired situations in woods and marshes.

The bite of this snake is not venomous; nor is the animal believed to bite at all, from any other impulse than that of anger. It kills its prey, by twining round it, and crushing its bones.

These monsters will attack and devour the largest animals, of which the following is an instance. A Boa had, for some time, been waiting near the brink of a pool, in expectation of its prey, when a buffalo appeared. Having darted upon the affrighted beast, it instantly began to encircle him with its voluminous twistings; and at every twist the bones of the buffalo were heard to crack as loud as the report of a gun. It was in vain that the animal struggled and bellowed; its enormous enemy entwined it so closely, that at length all its bones were crushed to pieces, like those of a malefactor on the wheel,

and the whole body was reduced to one uniform mass: the serpent then untwined its folds in order to swallow its prey at leisure. To prepare for this, and also to make it slip down the throat the more smoothly, it licked the whole body over, covering it with a mucilaginous substance. It then began to swallow it, at the end that afforded the least resistance; and in the act of swallowing, the throat suffered so great a dilation, as to take in a substance that was thrice its own ordinary thickness.

The Anaconda (*Boa murina*) is larger than the Boa Constrictor, being generally nearly thirty feet long. It is also a South American species; but the Python, which is of equal or greater size, is a native of India.

SECTION II.—Batrachian Reptiles.




THE FROG. (*Rana temporaria*.)

WHEN this reptile issues from the egg it is merely a black, oval mass, with a slender tail. This tadpole, as it is then

called, is the embryo of the Frog, and when it has attained a certain size, bursts its case, and appears with head and legs, leaving its tail with the rest of the exuviae of its former shape. This metamorphosis is one of the most curious in nature, and deserves our observation. Like other reptiles, it is not necessary for it to breathe in order to put its blood into circulation, as it has a communication between the two ventricles of the heart. It lives during spring in ponds, brooks, muddy ditches, marshy grounds, and other watery places; in summer in corn-fields and pasture land. Its voice proceeds from two bladders, one on each side of the mouth, which it can fill with wind: when it croaks it puts its head out of the water, or else it could not be heard. The hinder legs of the Frog are much longer than the fore ones, to help it in its repeated and extensive leaps. The whole of the body bears a little resemblance to some of the warm-blooded animals, principally about the thighs and the toes. The frog is extremely tenacious of life, and often survives the abscission of its head for several hours. It is supposed that Frogs spend the whole winter at the bottom of some stagnant water, in a state of torpidity.

There are several species of the Frog; they are all oviparous, and the eggs are gelatinous. The *Edible Frog* is the species used in France and Germany for food; it is considerably larger than the common kind, and though rare in England, is very plentiful in France, Germany, and Italy. Its colour is olive green, marked with black patches on the back and on its limbs with transverse bars of the same. From the tip of the nose, three distinct stripes of pale yellow extend to the extremity of the body; the middle one slightly depressed, and the lateral ones considerably elevated. The upper parts are of a pale whitish colour, tinged with green, and marked with irregular brown spots. These creatures are brought from the country, thirty or forty thousand at a time, to Vienna, and sold to the great dealers, who have froggeries for them, which are pits four or five feet deep, dug in the ground, the mouth covered with a board, and, in severe weather, with straw. In the year 1793, there were but

three great dealers in Vienna, by whom those persons who brought them to the markets ready for the cook were supplied: only the legs and thighs are eaten, and these are always skinned. They are rather dear, being considered a great delicacy. The Edible Frogs are caught in various ways; sometimes in the night, by means of nets, into which they are attracted by the light of torches that are carried out for the purpose; and sometimes by hooks, baited with worms, insects, flesh, or even a bit of red cloth. They are exceedingly voracious, and seize everything that moves before them.



THE TOAD, (*Bufo vulgaris*.)

WHOSE very name seems to carry with it something of an opprobrious meaning, is not unworthy the attention of the keen observer of nature: for though prejudice and false associations have affixed a stigma on certain species of animals, none of the works of our Creator are despicable; but all, the more minutely they are examined, are found to have a great claim to our admiration. Somewhat like the frog in the body, it also resembles that animal in its habits; but the frog leaps, while the Toad crawls. It is an error to suppose the Toad to be a noxious and venomous animal; it is as harmless as the frog, and, like some of the human kind, only labours under the stigma of undeserved calumny. Several stories have been related of its spitting poison, or knowing how to expel the venom it may have received from the spider or any other animals; but these fables have been long exploded. A curious, and yet inexplicable phenomenon is, that Toads have been


found alive in the centre of large blocks of stone, where they must have subsisted without food and respiration for a number of years. In the year 1719, M. Hubert, professor of philosophy at Caen, was witness to a living Toad being taken from the solid trunk of an elm-tree. It was lodged exactly in the centre, and filled the whole of the space that contained it. The tree was in every other respect firm and sound. Dr. Bradley saw a Toad taken from the trunk of a large oak. In the year 1733, a live Toad was discovered by M. Grayburg, in a hard and solid block of stone, which had been dug up in a quarry in Gothland. On being touched with a stick upon the head, he informs us, it contracted its eyes, as if asleep; and, when the stick was moved, gradually opened them. Its mouth had no aperture, but was closed round by a yellowish skin. On being pressed with the stick on the back, a small quantity of clear water issued from it behind, and it immediately died. A living Toad was found in a block of marble at Chillingham castle, belonging to Lord Tankerville, near Alnwick, in Northumberland.

With regard to the length of life of these animals, it is impossible to state anything decisive, as several facts prove that some of them have been gifted with astonishing longevity.

A correspondent of Mr. Pennant's supplied him with some curious particulars respecting a domestic Toad, which continued in the same place for *thirty-six* years. It frequented the steps before the hall-door of a gentleman's house in Devonshire. By being constantly fed, it was rendered so tame as always to come out of its hole in the evening when a candle was brought, and to look up as if expecting to be carried into the house, where it was frequently fed with insects. An animal of this description being so much noticed and befriended, excited the curiosity of all who came to the house, and even females so far conquered the horrors instilled into them by their nurses, as generally to request to see it fed. It appeared most partial to flesh-maggots, which were kept for it in bran. It would follow them on the table, and, when within a proper distance, would fix its eyes and remain

motionless for a little while, apparently to prepare for the stroke which was to follow, and which was instantaneous. It threw out its tongue to a great distance, and the insect, stuck by the glutinous matter to its tip, was swallowed by a motion quicker than the eye could follow. After having been kept more than thirty-six years, it was at length destroyed by a tame raven, which one day seeing it at the mouth of its hole, pulled it out, and so wounded it that it died.

SECTION III.—Saurian Reptiles.



THE LIZARD. (*Lacerta vivipara*.)

THIS is a British species, and is one of the very few reptiles found in Ireland. Its movements are very graceful. It comes out of its hiding-place during the day to bask in the sun; and when it sees an insect, it darts like lightning upon it, seizing it with its sharp little teeth, and soon swallowing it. The young are produced in eggs, which are generally hatched the moment they are laid, the skin of the egg being so thin that the young Lizard can be seen through it. The green Lizard is a beautiful creature. Its colours are more brilliant and beautiful than those of any other European species, and exhibit a rich and varied mixture of darker and lighter green, interspersed with specks and marks of yellow, brown, black, and sometimes even red. The head is covered with large angular scales; and the rest of the upper parts with very small ones. The tail is generally much longer than the body. Beneath the throat there is a kind of

collar, formed by scales of much darker colour than the rest of the animal.

The Lizard seems occasionally to lay aside its natural gentleness of disposition, but no further than for the purpose of obtaining food. Mr. Edwards once surprised a Lizard in the act of fighting with a small bird, as she sat on her nest in a vine against a wall, with newly-hatched young. He supposed that the Lizard would have made a prey of the latter, could it have driven the old bird from her nest. He watched the contest for some time; but, on his near approach, the Lizard dropped to the ground, and the bird flew off.




THE CHAMELEON. (*Chamaeleo vulgaris.*)

“ A lizard’s body, lean and long,
A fish’s head, a serpent’s tongue;
Its foot with triple claw disjoin’d;
And what a length of tail behind!
How slow its pace! and then its hue!”—MERRICK.

THE Chameleon is a small animal, about ten inches long, and its tail nearly the same length. Its body is shagreened

with small compressed scaly granules; its back is edged, and its tail round, long, and tapering. Its feet have each five toes, which are situated three one way and two another, in order to enable it to lay firm hold of the branches; but wherever it happens that these are too large for the animal to grasp with its feet, it coils round them its long, prehensile tail, and fixes its claws strongly into the bark. When walking on the ground, it steps forward in an extremely cautious manner, seeming never to lift one foot until it is well assured of the firmness of the rest. From these precautions, its motions have a ridiculous appearance of gravity, when contrasted with the smallness of its size, and the activity that might be expected from an animal so nearly allied to some of the most lively in the creation. Though the Chameleon is repulsive in its appearance, it is perfectly harmless. It feeds only on insects, for which the structure of its tongue is well adapted, being long and protrusive, and furnished with a dilated, glutinous, and somewhat tubular tip. With this it seizes on insects with the greatest ease, darting it out and immediately retracting it, with the prey thus secured, which it swallows whole. The strange notion that Chameleons were able to feed on air, seems to have arisen merely from the circumstance of these animals, like all others of the Lizard family, being able to subsist for a great length of time without food. The eyes of the Chameleon have the singular property of looking at the same instant in different directions; one of them may be seen to move when the other is at rest, or one will be directed forward, whilst the other is attending to some object behind; or in a similar manner upward and downward. It has the power of inflating its body to double its ordinary size, and at these times it is transparent. It can undoubtedly change its colour; but it is not true that it takes that of any object it may be near. On the contrary, its change of colour depends on its being exposed to a very strong light; and it only changes from its natural dull grey to a beautiful green, spotted unequally with red. Africa is the native country of the Chameleons, of which there are fourteen species; but two of them are found also in different parts of Asia and New Holland, and one (*C. vulgaris*) in the

south of Europe; but this animal has never been found in any part of America. Chameleons live in trees, hanging to the branches by their long prehensile tails.



THE CROCODILE OF THE NILE.

(*Crocodilus vulgaris*.)

THIS animal is frequently thirty feet long. The female lays about twenty eggs twice or thrice a year in the sand, where they are hatched by the heat of the sun; and the mother is said to take no care of the young ones. The head of this species, as of all the true Crocodiles, is twice as long as it is broad; the snout is pointed and unequal; and the eyes, which are small, are placed very far asunder. The colour is a greenish bronze, speckled with brown, and of a yellowish green underneath: six rows of nearly equal-sized plates run along the back. This Crocodile is less ferocious than some of the other kinds, and when taken young, may be tamed. It is common in Senegal and other parts of Africa, as well as in the Nile.


The double-crested Crocodile of the Ganges is very ferocious; and when gorged with prey, will bury what it can-

not eat in the mud. This is also done occasionally by the other species.

The *Crocodile* of South America is more flexible in its body than the other kinds; the female lays her eggs in the sand, but when the young are hatched, and she hears them, she scrapes away the sand to aid their escape, and watches over and feeds them till they are able to take care of themselves.

The method which the African adopts to kill this formidable creature displays considerable ingenuity and courage. Having wrapped a thick cloth round his arm, and provided himself with a long knife, he proceeds to the known haunt, usually a reedy swamp or river. The moment the crocodile perceives him it rushes at him with open mouth, but is coolly received by its antagonist, who thrusts his covered arm between its jaws. The teeth cannot pierce through the thick folds of the cloth, so that his arm only gets a smart squeeze, and before the creature can disengage itself, he adroitly cuts its throat.

The *Garvials* have very long, sharply-pointed snouts, and are feet webbed to the toes. They are much smaller than the crocodile, and not destructive to human life, as they live principally on fish. They have only been found on the Ganges.



THE ALLIGATOR, OR CAYMAN.

(*Alligator lucius.*)

THE habits of the Alligator are much the same as those of

the crocodile. The principal mark of distinction is, that the former has its head and part of the neck more smooth than the latter, and the snout is considerably more wide and flat, as well as more rounded at the extremity. The largest of these animals do not exceed eighteen feet. Alligators are natives of the warmer parts of America, and are the dread of all living animals. Their voracity is so great, that they do not spare even mankind. A short time before M. Navarette was at the Manillas, he was told that, as a young woman was washing her feet at one of the rivers, an Alligator seized and carried her off. Her husband, to whom she had been but just married, hearing her screams, threw himself headlong into the water, and, with a dagger in his hand, pursued the robber. He overtook and fought the animal with such success, as to recover his wife; but, unfortunately for her brave rescuer, she died before she could be brought to the shore.

The voice of the Alligator is loud and harsh. They have an unpleasant and powerful musky scent. M. Pagés says, that near one of the rivers in America, where they were numerous, their effluvia was so strong as to impregnate his provisions, and even to give them the nauseous taste of rotten musk. This effluvia proceeds chiefly from four glands, two of which are situated in the groin, near each thigh, and the other two at the breast, under each fore leg. Dampier informs us that, when his men killed an Alligator, they generally took out these glands, and, after having dried them, wore them in their hats by way of perfume.

The following anecdote of the voracity of this animal is related by Waterton in his "Wanderings in South America:"—"One Sunday evening, some years ago, as I was walking with Don Felipe de Ynciarte, governor of Angustura, on the bank of the Oroonoque, 'Stop here a minute or two, Don Carlos,' said he to me, 'while I recount a sad accident. One fine evening last year, as the people of Angustura were sauntering up and down here, in the Alameda, I was within twenty yards of this place, when I saw a large Cayman rush out of the river, seize a man, and carry him down, before anybody had it in his

power to assist him. The screams of the poor fellow were terrible as the Cayman was running off with him. He plunged into the river with his prey; we instantly lost sight of him, and never saw or heard him more.’”

SECTION IV.—Chelonian Reptiles.




THE COMMON, OR GREEK TORTOISE.

(*Testudo Græca.*)

THIS animal has a small head, four feet, and a tail, which it can gather within the shell in such a way that the top and under part meet together, and so closely, that the greatest strength cannot separate them. The eye is destitute of an upper lid, the under one serving to defend that organ. The upper shell, composed of thirty-seven compartments, is convex, and so strong, that a loaded cart can pass over it without injuring the creature inside. In winter Tortoises are said to bury themselves in the ground, or retire to some cavern or hole, which they line with moss, grass, and leaves, and where they pass in safe and solitary retirement the whole of this season. The Tortoise is very tenacious of life, and is no less remarkable for its longevity, as it is ascertained that one lived upwards of one hundred and twenty years in the garden of Lambeth Palace.

This animal is found in most of the countries near the Mediterranean Sea, in Corsica, Sardinia, and some of the islands of the Archipelago, as well as in many parts of the north of Africa.

The Tortoise-shell which is produced by this animal, as well as by the hawk's-bill turtle, is employed in several articles of useful and ornamental utensils, and when of a good colour, sells at a considerable price.



THE GREEN TURTLE. (*Chelonia mydas.*)

ALL the Turtles are considered very delicate food, especially the green, the hawk's-bill, and the loggerhead. Some of them are so large as to weigh from four to eight hundred pounds. Dampier mentions an immensely large one that was caught at Port Royal, in the Bay of Campeachy. It was nearly six feet in width, and four in thickness. A son of Captain Roch, a boy about ten years old, went in the shell, from the shore to his father's ship, which was about a quarter of a mile distant.

Turtle generally ascend from the sea, and crawl on the beach, either for food, or for the purpose of laying their eggs, (which are as large sometimes as those of a common hen,) sometimes to the number of fifty or sixty at a time. The young ones, as soon as they are hatched, crawl down to the water. Turtles are caught, when sleeping on land, by turning them on their backs; for as they cannot turn themselves over again, all means of escape is denied them. The lean of the Green Turtle tastes and looks like veal,

without any fishy flavour. The fat is as green as grass, and very sweet. The introduction of Turtle as an article of food into England, appears to have taken place within the last eighty or ninety years. They are common in Jamaica, and in most of the islands of the East and West Indies. Green Turtles are sometimes caught on the shores of Europe, driven thither by stress of weather. In the year 1752, one, six feet long and four feet broad, weighing between eight and nine hundred pounds, was caught in the harbour of Dieppe, after a storm. In 1754, a still larger one, upwards of eight feet long, was caught near Antioche, and was carried to the Abbey of Longveau, near Vannes, in Brittany; and, in the year 1810, a small one was caught amongst the sub-marine rocks, near Christchurch, in Hampshire.

The reader will remember how delighted Robinson Crusoe was, to find a large Turtle, which, he says, contained three score eggs. Behold him dragging it home,



BOOK V.—MOLLUSCOUS ANIMALS.

§ I. BIVALVES, or those having two shells.




THE PEARL OYSTER. (*Avicula Margaritifera*.)

Who that sees the beauty and delicacy of pearls would imagine that they were the production of disease? Such, however, is the case, as they are either formed in the body of the oyster which inhabits the shell; or they rise from cracks in the shell itself, the delicate, silvery, half-transparent lining of which forms the substance generally called Mother-of-pearl, or Nacre. These shells are found in the Persian Gulf and at Ceylon, where they form an important article of commerce.

The Chinese form pearls by casting into the shell of a certain kind of muscle artificial beads, which at the end of a year become covered with a pearly crust, in such a manner that they cannot be distinguished from the natural pearl.*

* For a very interesting article on this subject, see Beckmann's History of Inventions, vol. I. p. 259. *Bohn's Standard Library.*



THE COMMON OYSTER (*Ostrea edulis*)

HAS long been in favour with man for its delicacy as an article of food; the Lucrine lake used to be as much in renown among the Romans for the choicest kind of Oysters, as Canelle Bay with the French, and the Colchester beds with us. The two shells of the Oyster are generally unequal in size; the hinge is without teeth, but furnished with a somewhat oval cavity, and generally with lateral transverse grooves. Oysters sometimes grow to a very large size; in the East Indies they are said sometimes to measure nearly two feet in diameter.

The principal breeding season of Oysters is in the months of April and May, when they cast their young, which are enveloped in slime, and in this state called *spats* by the fishermen, upon rocks, stones, shells, or any other hard substance that happens to be near the place where they lie; and to these the spats immediately adhere. Till they obtain their film or crust, they are somewhat like the end of a candle, but of a greenish hue. The substances to which they adhere, of whatever nature, are called *cultch*. From the spawning time till about the end of July, Oysters are said to be sick; but by the end of August they become perfectly recovered; from May till August they are out of season, and unwholesome. The Oyster-fishery of our principal coasts is regulated by a court of admiralty. In the month of May the fishermen are allowed to take the Oysters, in order to separate the spawn from the cultch, the latter of which is thrown in again, for the purpose of preserving the bed for the

future. After this month it is felony to carry away the culch, and otherwise punishable to take any Oyster, between whose shells, when closed, a shilling will rattle. The reason of the heavy penalty on destroying the culch is, that when this is taken away, muscles and cockles will breed on the bed; and, by gradually occupying all the places on which the spawn should be cast, will destroy the Oysters.

The Oyster has been represented, by many authors, as an animal destitute not only of motion, but of every species of sensation. It is able, however, to perform movements which are perfectly consonant to its wants, to the dangers it apprehends, and to the enemies by which it is attacked. The gills, through which the Oyster breathes, are what is commonly called the beard, and are very indigestible. The scallop is nearly allied to the Oyster.

THE COMMON COCKLE. (*Cardium Fimbria.*)

FEW of our shell-fish are more common, in inlets and bays near the mouths of rivers, than these. In such situations they are usually found immersed at the depth of two or three inches in the sand, the place of each being marked by a small, circular, depressed spot. When they open their shells, the entrance into them is protected by a soft membrane, which entirely closes up the front, except in two places, at each of which there is a small, yellow, and fringed tube; by means of which they receive and eject the water which conveys to their body the nutriment necessary for their support.

Cockles are in great request as food among the labouring classes, and are caught chiefly in the winter months. Their size varies from five or six inches to half an inch in diameter. The shell is generally white; it has twenty-six longitudinal ridges, is transversely wrinkled, and has somewhat imbricated striæ. The foot of these animals is largely developed, and is to them a most important organ, as they use it not merely for progression, but in the excavation of hollows in the sand or mud in which they dwell.



1. THE MUSCLE. (*Mytilus edulis.*)

LIKE the oyster, the Muscle inhabits a bivalve shell, to which it adheres by a strong cartilaginous tie. The shells of several of the species are beautiful. The Muscle possesses the property of locomotion, which it performs with the member called its tongue, by which it gets hold of the rock, and is enabled to draw itself along; it has also the property of emitting a kind of thread, called the byssus, which, fixing the sides of the shell upon the ground, answers the purpose of a cable, to keep the body of the fish steady.

The *Chama*, which is akin to the Muscle, was used by the ancients to engrave various figures upon, from which circumstance those small bas-reliefs, so valued now, have obtained among the Italians and collectors the name of *Cameos*. The shells of some of these are decorated with red or yellow stripes, diverging from the hinge, and spreading to the edges. The *Giant Chama* has been found to weigh more than five hundred pounds, and the oyster-like animal within was large enough to furnish a meal for one hundred and twenty men. The animals which inhabit these shells are sometimes called Clams.

SECTION II.—Univalves.

2. THE ADMIRAL.

ONE of the cone-shells, the inhabitant of which is a kind of snail, with a very distinct head. If nature has taken a delight in painting the wings of birds, the skins of quadrupeds, and the scales of fishes, she seems not to have

been less pleased in pencilling the shells of these inhabitants of the deep. The variety, brightness, and versatility of the colouring have long been deservedly the object of man's admiration; and we cannot help being astonished at the richness which a cabinet of well-selected shells presents to the eye.

The manner of preparing the shells, and of bringing out their beautiful colours is simple, and yet requires great care. The crust must be gently rubbed with spirit of salt or *muriatic acid*, and then washed with clear water; this process will cleanse the shell, and display the wonderful brilliancy which was hidden under the first coat.



THE SNIPE, (*Murex haustellus*, or *cornutus*.)

So called on account of the length of a prominency coming out of the shell. It is surrounded with blunt prickles, and the colour of the whole is elegantly variegated.




THE LIMPET. (*Patella*.)

THE shape of this shell is pyramidal; it adheres to the rock with such strength, that it can only be removed by means of a knife or a strong blow. The apex of the shell is sometimes sharp, sometimes obtuse, and often

surrounded with points and sharp prickles. When thoroughly cleansed the shell is generally of a beautiful purple tint of great brilliancy, though the animal that lives under this magnificent roof is a kind of snail, disagreeable to the eye and insipid to the palate. They are found on the rocks, which are incessantly beaten by the surges and breakers, on the sea-shores of almost every country in the world. It is not by any glutinous liquid, as it has been asserted, that this fish adheres so strongly to the rock; but by the simple process of expelling the air from between its cloak and the rock to which it affixes itself.

The variety which is thrown into the sum of animated beings is so wonderfully great, that naturalists have reckoned more than a hundred and twenty-nine species of limpets, and nearly allied genera; the difference arising principally out of the diversity of the shells in form and colour.



THE GARDEN SNAIL (*Helix aspersa*)

Is furnished with four tentacula, two of which are smaller than the others; at the end of these tentacula, which

the animal pushes out or draws back, like telescopes, are blackish knobs, which are the eyes. The snail lays eggs, which are about the size of small peas, semi-transparent, and of a soft substance. By closely examining with a magnifying lens the eggs which a Water Snail, kept in a bottle of water, had deposited against the glass, the young Snail was seen in the egg, with its embryo shell on its back; two have also been observed in one egg, each of them with the rudiments of the shell.

The Garden Snail is extremely tenacious of life, and remains in a state of torpor during the winter. It is said, indeed, that it can remain in this state for many years, and the following instance is probably without parallel in any other animal. Mr. S. Simon, a merchant of Dublin, whose father, a Fellow of the Royal Society, and a lover of natural history, left him a small collection of fossils and other curiosities, had, among them, the shells of some Snails. About *fifteen years* after his father's death, he gave to his son, a child of ten years old, some of these Snail-shells to play with. The boy placed them in a flower-pot, which he filled with water, and the next day put them into a basin. Having occasion to use this, Mr. Simon observed that the animals had come out of their shells. He examined the child respecting them, and was assured that they were the same which had been in the cabinet. The boy said he had a few more, and brought them. Mr. S. put one of these into water, and, in an hour and a half afterwards, observed that it had put out its horns and body, which it moved but slowly, probably from weakness. Major Vallancy, Dr. Span, and other gentlemen, were afterwards present, and saw one of these snails crawl out; the rest being dead, probably from their remaining some days in the water.

The reproduction of the Snail is most curious; at a certain time of the year they meet in pairs, and stationing themselves an inch or two apart, they launch at each other several little darts, not quite half an inch long. These are of a horny substance, and sharply pointed at one end. The animals, during the breeding season, are furnished with a little reservoir for them, situated in the

neck, and opening on the right side. After the discharge of the first dart, the wounded Snail immediately retaliates on its aggressor by ejecting at it a similar one; the other renews the battle, and in turn is again wounded. Thus are the darts of Cupid, metaphorical with all the rest of the creation, completely realized in Snails. After the combat they both lay eggs.

The *Small Grey Slug* (*Limax cinereus*) resembles a Snail in all points except that it has no shell, consequently the brown skin of the back is rougher and stronger than that of the Snail. Its progress on the ground may easily be traced by the slime which it leaves in its track. Few animals are more destructive to vegetation than these.

The *Black Slug* is a well-known inhabitant of our fields and meadows, during the summer season. The country people consider its appearance as an indication of approaching rain; but this is rather to be accounted for by the moisture of the ground and plants. Indeed it very seldom appears abroad during dry weather. The Black Slug feeds on the leaves of different kinds of plants.


THE SEPIA, OR CUTTLE-FISH. (*Sepia octopodia.*)

THE structure of these animals is very remarkable. Their body is cylindrical, and, in some of the species, entirely covered with a fleshy sheath; in others the sheath reaches only to the middle of the body. They have eight arms, or rather legs, and in general two feelers, much longer than the arms. Both the feelers and arms are furnished with strong circular cups or suckers. The mouth is hard, strong, and horny, resembling in texture the beak of a parrot. The body is of a jelly-like substance, and usually covered with a coarse skin, having the appearance of leather. By means of the numerous circular cups or suckers with which the arms are furnished; they seize their prey, and firmly attach themselves to the rocks. Their adhesive power is so great, that it is generally more easy to tear off the arms, than to separate them from the substance to which they are affixed: if the arms happen to be broken off, they are

soon reproduced. The size to which this creature grows has been variously stated; and, although evidently exaggerated by some authors, it undoubtedly attains to a very considerable magnitude. When attacked in its own element, it has been known to overcome a large dog. Its jaws are extremely strong and powerful, and with its beak it can crush in pieces the shells of the fish on which it feeds. In the body is a bladder filled with a dark inky fluid, which it emits when alarmed, and which not only tinges the water so as to conceal its retreat, but is so bitter as immediately to drive off its enemies. This inky fluid, when dried, forms a very valuable colour, used by artists, and known as Sepia.

The bone, or calcareous plate of the *Sepia Officinalis*, a species common on our coasts, is a well-known substance, and is much employed in the manufacture of tooth-powder; and by silversmiths for moulds, to cast their small work, such as rings, &c. It is also converted into that useful article of stationery, called pounce.


The animal of the ARGONAUT, or PAPER NAUTILUS, is a kind of Sepia.



BOOK VI.

ARTICULATED ANIMALS.

§ I. ANNELIDA, or ringed animals.



WORMS. (*Vermes.*)

THESE creatures constitute a class by themselves, under the name of annelida, in the works of modern naturalists. They are distinguished from the caterpillar and maggot, by undergoing no change, and crawling by means of the annular structure of their bodies.

The *Earth Worm* has neither bones, eyes, ears, or brain; it has a round, annulated body, with generally an elevated fleshy belt near the head. Though considered a

great nuisance by gardeners, Earth Worms perforate, and loosen the soil, and render it pervious to rains and the fibres of plants, by drawing into it straws and the stalks of leaves; and chiefly by throwing infinite numbers of lumps called worm-casts, which form a fine manure for grass and corn. They are, however, very injurious to plants in pots.



THE LEECH (*Hirudo medicinalis*)


Is about three inches in length, and in its exterior form somewhat resembles the worm, when extended, but often contracts itself greatly in length, at the same time expanding in thickness. It has a small head, a black skin, with six yellow lines above, and spotted with yellow below. The mouth of the Leech is of curious construction; it has three jaws, each of which is armed with two ranges of very fine teeth, with which it pierces the skin; and then draws up, as through a siphon, the blood, upon which it feeds. The progressive movement of the Leech is effected by sticking, by suction, its mouth to a certain spot, then bringing its tail, which also has the property of sticking, in the same manner as the head, and then advancing its head further on, quickly followed by the tail, and so on. The common Leech is very often met with in brooks and rivulets. Its uses in medicine are well known, as by its means the blood can be extracted from diseased parts, to which the lancet cannot be applied.

The blood which the Leech sucks out of the wound it makes, supplies it with nutriment for so great a period of time, that a Leech, after having been satisfied with blood, has been known to live three years without any food. It


is usual, however, to make them disgorge the greater part of the blood they have swallowed by sprinkling them with salt; as otherwise they would not bite again till the blood they had taken was fully digested.

Leeches lay eggs, which are covered with a kind of membrane, which serves to protect them when they are deposited in the clay and holes in the sides of ponds. They appear to live on the eggs of fish or frogs, but eagerly attach themselves to the legs of human beings, horses, or cows, whenever they have an opportunity. As there is a prejudice among the country people that Leeches, never breed well till they have tasted blood, it is said that they drive their horses and cows into the water inhabited by the Leeches, and consequently that the Leech districts are remarkable for their wretched-looking horses and cattle. Leeches must be five years old before they are fit for medical purposes; and they are caught in shallow water in spring by people going in with naked feet and ankles, to which the Leeches adhere, when they are picked off and put in baskets provided for the purpose. In summer a raft is made of twigs, and the waters being disturbed with a stick, the Leeches rise to the surface, and get entangled in the raft. When caught, they are washed in water with a very little salt in it, and packed in wet linen cloths, which are put into a barrel with a canvas cover, and sent away for sale. London is chiefly supplied from the fenny districts of Lincolnshire. They are also, when kept in a glass bottle with water, a good barometer, as they always come up to the neck of the bottle when rainy weather is approaching, remain at the bottom in dry weather, and move anxiously up and down when the weather is stormy.—Horse-Leeches are larger than the common species, more voracious, and narrower at each extremity.

SECTION II.—Crustacea.


THE LOBSTER (*Astacus marinus*)

HAS a cylindrical body, long antennæ, and a broad tail. Its large claws enable it to seize on its prey, to fix itself on the small prominences of rocks in the sea, to resist the motion of the waves, and to defend itself against its enemies. When the Lobster wants to spring off the rocks, it makes a fulcrum of its tail, which has the action of a powerful spring. Its gait is awkward, as in all the crustacea. Besides its claws, it has four small legs on each side, to assist it in its movements. Under the tail the hen Lobster preserves her eggs till they are hatched. They are extremely prolific. Dr. Baxter says he counted twelve thousand four hundred and forty-four eggs under the tail of a female Lobster, besides those that remained in the body undeveloped. Lobsters deposit their eggs in the sand, where they are soon hatched. Like the rest of their tribe, they cast their shells annually, previous to which they appear languid and restless: they acquire an entirely new covering in a few days.



THE CRAYFISH (*Astacus fluviatilis*)

MAY be called the lobster of fresh water, and its presence is generally esteemed an evidence of the goodness of the water. Crayfish are considered a very strengthening food. They are caught in shallow brooks, hid under large stones, out of which they crawl backwards to seek for their prey, which consists of small insects; the hooks employed to catch them are baited with liver or flesh, which they nibble most greedily.



THE CRAB. (*Cancer pagurus*.)

CRABS are of various sizes, some weighing several pounds, and others only a few grains, all of different species. They do not move forward, but sideways. They have a small tail closed on the body; which forms a considerable

and essential difference between them and the lobsters, prawns, shrimps, and cray-fish.

The most remarkable circumstance in the history of these animals, is the changing of their shells and the renewal of their broken claws. The former, as it is stated, take place once a year, and usually between Christmas and Easter. During the operation they retire among the cavities of rocks, and under great stones. Crabs are naturally quarrelsome amongst themselves, and frequently have serious contests, by means of those formidable weapons, their great claws. With these they lay hold of their adversary's legs; and wherever they seize, it is not easy to make them forego their hold. The animal seized has, therefore, no other alternative but to leave part of the leg behind in token of victory.

An experiment was tried to prove the extremely tenacious disposition of the Crab. By irritating it, a fisherman made a Crab seize one of its own small claws with a large one. The animal did not distinguish that it was itself the aggressor, but exerted its strength, and soon cracked the shell of the small claw. Feeling itself wounded, it cast off the piece in the usual place, but continued to hold it with the great claw for a long time afterwards.



The *Violet Land Crabs* of the Caribbee Islands are most singular in their habits; they descend in annual and

regular caravans from the mountains, their natural abode, to the sea-shores, in order to deposit their spawn, after which they again return to the mountains. These Crabs form, in their procession, a body of fifty paces broad, wheeling along three miles in depth. This battalion moves slowly, but with regularity and uniformity, either when they descend or ascend the hills. They abound in Jamaica, where they are accounted a great delicacy by the natives, and are common in the adjacent islands.




THE SOLDIER CRAB, OR HERMIT CRAB,

(*Pagurus miles*.)

Is a curious animal, and ought to be noticed here for its singular habits. It is somewhat like a lobster divested of its shell; it is about four inches in length, and has no shell on the hinder part, but is covered down to the tail with a rough skin; it is also armed with strong hard nippers. This Crab has not been provided by nature with a shell, and is obliged to seek for one which has been deserted by its legitimate tenant; but as this covering cannot grow of course proportionally with him, he is forced out of it by his increasing size, and finds himself under the necessity of looking out for a new one: it is curious to see him when in want of a new house, crawling from one empty shell to another, examining and trying his new habitation. Sometimes, when two competitors happen to eye the same premises, a great contest

arises, and of course the strongest gets the manor. These Crabs live on sea-weed, and the smaller insects which they find in the puddles left by the receding tide.



1. THE SHRIMP. (*Crangon vulgaris*.)

THE Shrimp is a well-known small crustaceous animal, nearly allied to the lobster, which it resembles in shape. It has long slender feelers, between which are two projecting laminae; ten feet and five fins, but no claws. This animal breeds on all the sandy shores of Great Britain: it is frequently found in harbours, and even in the ditches and ponds of salt marshes. During life the body is semi-transparent, and so much resembles sea-water, that the animal is distinguished with difficulty. Its ordinary motion consists of leaps. Its flavour is very delicate.

2. THE PRAWN. (*Palemon serratus*.)

THE Prawn is not unlike the shrimp, but exceeds it considerably in size; and has a projecting ridge down the back, furnished with sharp teeth. Its colour, when boiled, is a most beautiful pink. The flesh is very delicate although perhaps inferior in flavour to that of the shrimp.

Prawns are chiefly found among sea-weed, and in the vicinity of rocks, at a little distance from the shore. They seldom enter the mouths of rivers. Their usual

mode of swimming is on their backs; but when threatened with danger, they throw themselves on one side, and spring backward to very considerable distances. They feed on all the smaller kinds of marine animals, which they seize and devour with great voracity. In their turn, they are the prey of numerous species of fish, although the sharp and serrated horn in front of their head constitutes a powerful weapon of defence against the attacks of all the smaller kinds. At the side of the head there is frequently to be observed a large and apparently unnatural lump. This, if examined, will be found to contain, under the thoracic plate, a species of parasitic animal, which occupies the whole cavity, and there feeds and perfects its growth. The same tumour or lump may be also observed on the shrimp.

Being in great request for the table, both shrimps and Prawns are eagerly sought for by fishermen, who catch them either in osier baskets, similar to those employed in catching lobsters, or in a kind of net called a *Putting-net*. These, which are well known to all frequenters of the sea-coast, are five or six feet in width, and flat at the bottom; and are pushed along in the shallow water, upon the sandy shores, by a man who walks behind.

SECTION III.—Arachnida.

THIS order, according to Lamarck, and other modern zoologists, contains the Spiders, Scorpions, Centipedes, Mites, and other wingless insects which do not undergo any metamorphoses. These creatures differ from the true insects in the number of their feet, which are generally eight, and frequently very numerous, while those of the true insects never exceed six. They also differ in their manner of breathing, the spiders and scorpions being furnished with air-sacks, or bags, which act like lungs, while the other animals belonging to the class have tracheæ, or air-passages, like the true insects.



SPIDERS. (*Arachnida.*)

ALL the Spiders are distinguished by having no antennæ, eight legs, and generally eight eyes; mandibles terminated by a moveable claw, which sometimes emits poison; and an abdomen without rings, furnished at its point with four or six spinnarets, from which the Spider emits the threads used in spinning its web. This web is wonderful in its formation.

“The Spider’s touch, how exquisitely fine!
Feels at each thread, and lives along the line.” POPE.

The Spider sits in the middle, and at the least motion caused by a fly or other insect pressing against it, rushes on his prey, and sucks its juices; if, however, it should appear at all formidable, the Spider carefully encloses it in a shroud of web, which, of course, quite disables it; and then feasts on it at his convenience. The most difficult part of the business is to eject the remains, which is often attended with great detriment to the net. The female generally lays from nine hundred to a thousand

eggs, which are contained in a kind of bag, and thus an immense number of Spiders are hatched every year, which would soon become troublesome from their numbers, if they were not kept in check by the numerous birds which prey upon them. The silk which the Spider produces is not strong enough to be employed for any useful purpose, though, out of curiosity, gloves and stockings have been woven out of it. A great difficulty, however, arises in the pugnacious habits of Spiders, as when a number of them are kept together, they fight so dreadfully, that in a short time only a very few are left alive; and a great number would be required, as twelve Spiders do not produce so much silk as a single silkworm. Spiders resemble the crustacea in having the power of reproducing the legs which they lose.



THE TARANTULA. (*Lycosa tarantula*.)

THIS Spider is a native of the South of Europe. It lives in fields, and its dwelling is about four inches deep in the ground, half an inch wide, and closed at the mouth with a net. They lay about seven hundred and thirty eggs, which are hatched in the spring. These Spiders do not live quite a year; the parents never survive the winter.

Inflammation, difficulty of breathing, and sickness, are said to be the invariable consequences of the bite of this animal. Dr. Mead, and other medical men, have countenanced the popular story of these effects being counter-

acted by the power of music. It is, however, now well known, that this singular mode of cure was nothing more than a trick frequently practised on credulous travellers, who were desirous of witnessing it. Mr. Swinburne, when he was in Italy, minutely investigated every particular relative to the Tarantula. The season was not far enough advanced, and it was pretended that no persons had as yet been bitten that year: he, however, prevailed upon a woman, who had formerly been bitten, to dance the part before him. Several musicians were summoned, and she performed the dance, as every one present assured him, to perfection. At first she lolled stupidly on a chair, while the instruments played a dull strain. They touched at length the chord supposed to vibrate to her heart; and up she sprung with a hideous yell, staggered about the room like a drunken person, holding a handkerchief in both hands, raising them alternately, and moving in very true time. As the music grew brisker, her motions quickened, and she skipped about with great vigour, and in a variety of steps, every now and then shrieking very loud. The scene was unpleasant, and, at his request, an end was put to it before the woman was tired.

He informs us, that, whenever they are to dance, a place is prepared for them, hung round with bunches of grapes and ribbons. The patients are dressed in white, with red, green, or yellow ribbons; on their shoulders they have a white scarf; they let their hair fall loose about their ears, and throw the head quite back. He says that they are exact copies of the ancient priestesses of Bacchus. The introduction of Christianity abolished all public exhibitions of heathenish rites; but the women, unwilling to give up their darling amusement, in performing the frantic character of Bacchantes, devised other pretences; and he supposes that accident led them to the discovery of the Tarantula, of which they took advantage for that purpose.

THE CHEESE MITE. (*Acarus domesticus*, or *siro*.)

THESE destructive little creatures differ from spiders in having the thorax and abdomen united and covered with

the same skin, though it is contracted in one part. They have also, when young, only six legs, though the two others appear afterwards; and their feet are armed with strong hooks, which enable them to retain hold of the cheese or other food, in which they take up their abode. Their bodies are covered with hair, and their mouths are furnished with strong mandibles, with which they soon hew down huge rocks and mountains of cheese. The eggs of these mites are so small, that it has been computed that a pigeon's egg would contain thirty millions of them. It must be observed that this mite is only found in dry cheese, in which it looks like reddish dust. The cheese-hopper, found in moist rotten cheese, is the maggot of a kind of fly. (*Prophila Casei*.)

SECTION IV.—Insects.

LINNÆUS included under the name of Insects all those animals which have their legs and bodies divided into segments or joints; the term insect being derived from the Latin word *insectare* to cut or notch. Modern Zoologists, however, call these animals the articulata, and confine the term Insect to those creatures which undergo metamorphoses.

These Insects have all six legs and two antennæ or feelers; and though the transformations they undergo differ slightly in the different kinds, the following is the order in which they occur:—the perfect insect lays eggs, which when hatched produce larvæ; which are called grubs when they belong to beetles, maggots to flies, and caterpillars to butterflies. These larvæ eat voraciously; and, as they rapidly increase in size, they generally moult, that is, change their skins, two or three times. When the larvæ are full grown, they go into the pupa state, in which they remain torpid and without food for a considerable length of time, sometimes first spinning a loose covering for the pupa called a cocoon. The pupa is generally called a chrysalis; but it is also sometimes called a nymph, and sometimes an aurelia. The last

transformation is when the insect breaks from its covering in a perfect form, when it is called the imago. The perfect insect is divided into three segments, or parts, called the head, the thorax, and the abdomen.

ORDER I.—Coleoptera, or Beetles.

THE larva of the beetle is a grub, which generally continues in that state three or four years, eating voraciously during the whole period. When full grown it in most cases either descends into the ground, where it undergoes its transformations, first into a nymph, or pupa, and then into a beetle; or it makes itself a rough cocoon of bits of stick and dead leaves, in which it changes into a pupa, and afterwards into a beetle. The wood-eating beetles undergo their transformations in the tree on which they feed. The pupa of the beetle is termed incomplete, because all the parts of the insect are visible in it, instead of being enclosed in one thick covering, as in the moths and butterflies. The head of the beetle is furnished with two compound eyes; two antennæ; (differing in shape in the various species, but always having at least eleven joints;) and a mouth, consisting of a labrum, or upper lip, a labium, or under lip, two mandibles, or upper jaws, and two maxillæ, or under jaws. There is also the mentum, or chin, and a part called the clypeus, to which the upper jaw is attached.

The thorax is the part which supports the legs and wings. The legs are divided into five portions, of which the part terminated by the claw is called the tarsus. There are two membraneous wings, which are covered by two hardened wings or wing-cases called the elytra, which generally open by a straight line down the back; and hence the name of Coleoptera, which signifies wing in a case: the abdomen is simply the body.

The number of beetles is very great, and indeed Mr. Westwood informs us that more than thirty thousand species have been described, of which about three thousand five hundred are natives of Britain.



THE COCKCHAFER. (*Melolontha vulgaris*.)

THE Cockchafer is one of the lamellicorn beetles. The female lays her eggs in the ground, and the grubs, when hatched, are soft, thick, and whitish. It is from its white appearance that the grub of the Cockchafer is called *le ver blanc* by the French. These grubs, sometimes in immense numbers, work between the turf and the soil in the richest meadows, devouring the roots of the grass to such a degree that the turf rises, and will roll up with almost as much ease as if it had been cut with a turfing knife; the soil underneath appearing, for more than an inch in depth, like the bed of a garden. In this, the grubs lie, on their backs, in a curved position, the head and tail uppermost, and the rest of the body buried in the mould. It is also said that a whole field of fine flourishing grass has become, in a few weeks, withered, dry, and as brittle as hay, in consequence of these grubs devouring the roots.

In the year 1688, great numbers of Cockchafers appeared on the hedges and trees of the south-west coast of the county of Galway, in clusters of thousands, clinging to each others' backs, in the manner of bees when they swarm. During the day they continued quiet, but towards sunset, the whole were in motion; and the humming noise of their wings sounded like distant drums. Their numbers were so great, that, for the space of two or three square miles, they entirely darkened the air. Persons travelling on the roads, or who were abroad in the fields, found it difficult to make their way home, as the insects were continually beating against their faces, and occasioned great pain. In a very short time the

leaves of all the trees, for several miles round, were destroyed, leaving the whole country, though it was near midsummer, as naked and desolate as it would have been in the middle of winter. The noise which these enormous swarms made, in seizing and devouring the leaves, was so loud as to have been compared to the distant sawing of timber. Swine and poultry destroyed them in vast numbers; waiting under the trees for the clusters of insects to drop, and then devouring such swarms as to become fat upon them alone. Even the native Irish, from the insects having eaten up the whole produce of the ground, adopted a mode of cooking them, and thus used them as food. Towards the end of the summer, they disappeared so suddenly, that in a few days there was not one left.

Rooks are very fond of eating these grubs, and often, when they are seen in a newly-sown field, apparently devouring the grain, they are, in fact, rendering the greatest service to the farmer, by destroying his great enemy, the white worm.


THE DOR, OR BLIND BEETLE.

(*Geotrupes stercorarius.*)

THIS well-known insect, which is sometimes also called "the shard borne beetle," has been often noticed by the poets. Amongst others, Shakespeare makes Macbeth say,

" Ere to black Hecate's summons
The shard borne beetle, with its drowsy hum,
Hath rung night's yawning peel, there shall be done
A deed of dreadful note."

This beetle, which is a British insect, generally lays its eggs in a mass of cow-dung, which it afterwards buries in the earth. It makes a dull drowsy noise when it flies, and often strikes itself against any person or object it may meet, as though it were blind. It has also the habit of stretching out its limbs and pretending to be dead when caught.



THE STAG BEETLE. (*Lucanus cervus*.)

“ See the proud giant of the Beetle race ;
 What shining arms his polish'd limbs encase !
 Like some stern warrior, formidably bright,
 His steely sides reflect a gleamy light ;
 On his large forehead spreading horns he wears,
 And high in air the branching antlers bears :
 O'er many an inch extends his wide domain,
 And his rich treasury swells with hoarded grain.”

BARBAULD.

THIS insect is the largest, and most singular in shape, of any in this country. It is known by two horn-like mandibles, projecting from its head, and resembling those of a stag, with which it is able to pinch very severely. These mandibles are strongly dentated from the root to the point. The wing-cases have neither streaks nor spots. The whole insect is of a deep brown. It is sometimes found in hollow oaks and beeches, near London ; but is much smaller than the same species found in the country.


The larvæ, or grubs, lodge under the bark, or in the hollow of old trees ; which they bite and reduce to fine powder. The larvæ are supposed to exist three or four years before they form their cocoons. These insects are mostly found in Kent and Sussex. In Germany there is a popular but idle notion, that they sometimes, by means of their jaws, carry burning coals into houses ; and that, in consequence of this mischievous propensity, dreadful fires have been occasioned. The Stag Beetle is one of the lamellicorn Coleoptera.



THE ELEPHANT BEETLE

(*Scarabeus, or Dynastes elephas*)

Is found in South America, particularly in Guiana, and Surinam, as well as near the river Orinoko. It is the largest beetle of its kind hitherto known; it is black, and the whole body is covered with a very hard shell, quite as thick and as strong as that of a small crab. Its length, from the hinder part to the eyes, is almost four inches; and from the same part to the end of the proboscis, or trunk, four inches and three quarters. The transverse diameter of the body is two inches and a quarter; and the breadth of each case for the wings, upwards of an inch. The antennæ, or feelers, are quite horny; for which reason the proboscis, or trunk, is moveable at its insertion into the head, and seems to supply the place of feelers. The horns are about an inch long, and terminate in points. The proboscis is an inch and a quarter long, and turns upwards, making a crooked line terminating in two horns, each of which is nearly a quarter of an inch long; but they are not perforated at the end, like the proboscis of other insects; the feet are all forked. Above the head is a prominence, or small horn, which, if the rest of the trunk were away, would cause this part to resemble the horn of a rhinoceros. There is, indeed, a beetle named after that animal, whose lower horn resembles this: its scientific name is *Oryctes Rhinoceros*.



THE MUSK BEETLE, OR GOAT CHAFFER.

(*Cerambyx moschatus.*)

THIS is one of the longicorn beetles. It is a very beautiful insect, of a glossy bluish green colour, with a cast of shining gold. The wings under the case are black. The legs are of the same bluish green colour, only somewhat paler; and the breast is pointed at each extremity. Between these points are three little tubercles near the wings, and three smaller towards the head. The cases of the wings are oblong, and somewhat in the shape of a lance; with three ribs a little raised and running lengthways. The feelers are nearly as long as the body, composed of many small joints, which grow smaller near the ends. It is found among old willow-trees, and has a kind of musky smell.



THE GROUND BEETLE. (*Carabeus clathratus.*)

THE Ground Beetle is not only one of the largest but the most beautiful and brilliant that this country produces. The head, breast, and wing-cases are of a coppery green;

the latter having three longitudinal rows of oblong raised spots. All the under part of the insect is black. Having only very short wings beneath the cases, Nature has providentially supplied it with such legs as enable it to run with amazing swiftness. This insect is frequently found in damp places, under stones and heaps of decayed plants in gardens. There are several species, one of which (*Carabeus violaceus*) is of a beautiful purple.

The larvæ live under ground, or in decayed wood where they remain until metamorphosed to their perfect state, when they proceed to devour the larvæ of other insects, and all weaker animals that they can conquer.

The Ground Beetles are found as early as the beginning of March, in paths and near old walls, where the sun warms the earth with its vivifying beams. Many of the large species have been found between the decayed bark and wood of willow-trees.



THE GLOWWORM. (*Lampyris noctiluca*.)

It is only the female Glowworm which produces the beautiful light, for which the insect is so well known, and she frequently communicates this light to her eggs. She is without wings or wing-cases, and possesses no beauty when seen by daylight. The male has wings, and leathery elytra. The larva is a very ugly, and very voracious grub, which will eat any vegetable matter it can find, and will even feed on snails and slugs.



THE DEATH-WATCH. (*Anobium tessellatum*.)

THIS creature is called the Death-Watch, from a superstitious notion, that when its beating is heard, it is a sign that some one in the house is going to die. The insect lives in wood, and the noise is produced by its striking its head against whatever is near it. These insects, in the larva state, do a great deal of mischief to old furniture, in which they perforate numerous round holes. To enable them to do this they are furnished with two maxillæ formed like two cutting pincers, with the help of which they bore the holes so neatly that the French call them *vrilletes*, from *vrille*, a gimlet. Sometimes two of these insects may be heard ticking, answering each other; and sometimes the Death-watch may be made to tick, by tapping with the finger-nail upon a table. These creatures imitate death with great exactness when they are caught, or when they think themselves in danger.



THE GRAINED BULL-HEAD, AND SPANISH FLY, OR CANTHARIDIS. (*Cantharis vesicatoria*.)

THESE insects are natives of the South of Europe, and are very important in medicine for their blistering powers. They have a very disagreeable smell, and emit a fluid of so corrosive a nature, that many persons have suffered greatly from gathering them; and it is said to be extremely dangerous to sleep under a tree infested by them, as their smell produces a lethargic sleep, which frequently terminates in death. They are generally caught by laying linen cloths under the trees they infest, and beating the boughs; they are then put into hair sieves, and held over vessels of boiling vinegar, till the vapour

kills them. After this they are dried in ovens, or on hurdles, exposed to the sun, and then packed up for sale. When dried, fifty of them hardly weigh a drachm.

They have sometimes been administered internally, by reckless people, as aphrodisiacs. They are of a highly poisonous nature, and there are many instances, some even recent, of their producing violent hæmorrhage and death.



THE LADY BIRD, OR LADY COW.

(*Coccinella septem punctata.*)

THE larva of this well-known and beautiful little beetle is disagreeable and almost disgusting in its appearance; but to compensate for this it is extremely useful in destroying the aphid, or green fly. In the perfect insect the elytra are scarlet, beautifully spotted with black; some species having seven, and others five spots, and one of the most beautiful, eighteen. The head is very small, the antennæ and legs very short, and the body nearly round. The English name is said to be derived from the quiet habits of the insect, as it is so gentle that it may crawl on a lady's hand without her knowing it.

ORDER II.—Orthoptera.

In this order the elytra, or wing cases, are much softer and more flexible than in the beetles; they are frequently membranous or webbed, and when closed they do not form a straight line down the back. The mouth is also different; the maxillæ being terminated by a horny, toothed piece called the galea. There is also a kind of tongue, and the metamorphosis is only half complete.




THE EARWIG. (*Forficula auricularia.*)

UNLIKE those of most other insects the eggs are hatched

and the young Earwigs are fostered by the parent. At the beginning of the month of June, M. de Geer found, under a stone, a female Earwig, accompanied by many little ones, evidently her young. They continued close to her, and often placed themselves under her body, as chickens do under a hen.

This little animal is very nimble, and perfectly harmless, except to flowers, notwithstanding the fabulous charge which was so long believed against it, of entering the human ear, and depositing its eggs there, which were said to cause intolerable pain when hatched, and the young began to gnaw the inside of the ear. The Earwig possesses wings, which, when extended, cover nearly the whole insect. The elytra, or wing-cases, are short, and do not extend along the whole body, but only over the breast. The wings are concealed beneath these, and are somewhat of an oval shape. There is great elegance in the manner in which the insect folds its wings beneath its elytra.

The *Black Beetle*, or Cockroach, (*Blatta Orientalis*), so common in London kitchens, is nearly allied to the Earwig



THE WALKING LEAF, OR MANTIS.

(*Phyllium Siccifolium*.)

THIS insect is remarkably shaped. The head is joined to

the body by a neck longer than the body itself. It has two polished eyes, and two short feelers. The breast is long, narrow, and margined. The wing-cases, which cover two-thirds of the body, are veined and reticulated, or netted. The wings are veined and transparent. The hinder legs are very long, the next shorter; and the foremost pair of thighs are terminated with spines: the others have membraneous lobes, which serve them as wings in their flight. The top of the head is membraneous, shaped like an awl, and divided at its extremity. This animal is one of the innumerable instances which Nature affords of the infinite wisdom of the Creator; for, whenever an animal is found to deviate in shape from the general system, it is still formed to answer the design of its existence. Thus, this insect having such long legs, could never have sustained itself in the air, had not Providence bestowed on the legs themselves a species of wings to balance their weight. These are instances with which Nature teems; and which would make the atheist tremble, would he but contemplate the admirable design and system with which they are characterised as

“Parts of one stupendous whole;
Whose body Nature is, and God the soul.”

These insects are partly of a pale yellowish green, and partly brown; so that they look like dead leaves, whence their English name. They are found in the East Indies and China.




THE GRASSHOPPER (*Gryllus*, or *Acrydium*)

Is of a bright green, the head somewhat resembling that of a horse; the corselet is armed with a strong buckler.

Of its six legs the hinder two are much longer than the others, to assist the insect in leaping. The males make a chirping noise, which is caused by the thighs being rubbed alternately against the tip of the wing-cases: if handled roughly, the Grasshopper bites very sharply.

Toward the end of autumn the female deposits her eggs in a hole, which she makes in the earth for this purpose, by means of an instrument at her tail, called an ovipositor. These eggs sometimes amount to a hundred and fifty; they are about the size of carawayseeds, white, oval, and of a horny substance. The female, having thus performed her duty, soon languishes and dies. In the beginning of May following a small white grub issues out of each egg. The creature passes about twenty days under this humble and creeping form; after which, having assumed the pupa shape, whilst all the rudiments of the future Grasshopper are concealed under a thin outward skin, it retires under a thistle or a thorn-bush, most likely in order to be more secure; and there, after a variety of laborious exertions, writhings, and palpitations, the temporary covering divides, and the insect jumps out of its *exuvia*.



THE LOCUST, MOLE CRICKET, AND CRICKET.

1. THE LOCUST. (*Gryllus, Migratorius.*)

THE Bible, which was written in a country where the Locust made a distinguished figure among natural pro-

ductions, has given us several very striking images of these animals' numbers and rapacity. It compares an army to a swarm of locusts: it describes them as rising out of the earth, where they are produced; as pursuing a settled march to destroy the fruits of the earth, and as the frequent instruments of Divine indignation.

The native countries of the Locust are Central Asia and the North of Africa, but they migrate every year to Europe, where they destroy every green thing they meet with.

When the Locusts take the field they have a leader at their head, whose flight they observe, and to whose motions they pay a strict attention. They appear at a distance like a black cloud, which, as it approaches, gathers upon the horizon, and almost hides the light of the day. It often happens that the husbandman sees this imminent calamity pass away without doing him any mischief; and the whole swarm proceed onward, to settle upon the labours of some less fortunate country. But wretched is the district upon which they fix; they ravage the meadow and the corn land; strip the trees of their leaves, and the gardens of their beauty; the visitation of a few minutes destroys the expectations of a year; and a famine but too frequently ensues. In their native climates they are not so injurious as in the south of Europe, for in Syria and Palestine, though the plain and the forest be stripped of their verdure, the power of vegetation is so great, that an interval of three or four days repairs the calamity: but our verdure is the produce of a season; and we must wait till the ensuing spring repairs the damage. Besides, in their long flights to this part of the world, the Locusts are famished by the tediousness of their journey, and are therefore more voracious wherever they happen to settle. But it is not by what they devour that they do so much damage, as by what they destroy. Their very bite contaminates the plant, and injures its future vegetation. To use the expression of the husbandman, they burn whatever they touch, and leave the marks of their devastation for two or three years ensuing. And if so noxious while living, they are still more so when dead; for wherever they fall they infect the air in such a manner that the smell is insupportable.

In the year 1690 a cloud of Locusts was seen to enter Russia in three different places; and thence to spread themselves over Poland and Lithuania, in such astonishing multitudes, that the air was darkened, and the earth covered with their numbers. In some places they were seen lying dead, heaped upon each other to the depth of four feet; in others they covered the surface like a black cloth: the trees bent beneath their weight, and the damage which the country sustained exceeded computation. In Barbary their numbers are formidable, and their visits frequent. In the year 1724, Dr. Shaw was a witness of their devastations in that country. Their first appearance was about the latter end of March, when the wind had been southerly for some time. In the beginning of April their numbers were so much increased, that in the heat of the day they formed themselves into large swarms, which appeared like clouds, and darkened the sun. In the middle of May they began to disappear, retiring into the plains to deposit their eggs. In the next month, being June, the young brood began to make their appearance, forming many compact bodies of several hundred yards square; which marching forward, climbed the trees, walls, and houses, eating everything that was green in their way.

“ ——— To their general's voice they soon obey'd
 Innumerable. As when the potent rod
 Of Amram's son, in Egypt's evil day,
 Waved round the coast, upcall'd a pitchy cloud
 Of Locusts, warping on the eastern wind,
 That o'er the realms of impious Pharaoh hung
 Like night, and darkened all the land of Nile;
 So numberless were those bad angels seen,
 Hovering on wings, under the cope of Hell,
 'Twixt upper, nether, and surrounding fires.”—MILTON.

2. THE MOLE CRICKET. (*Gryllotalpa vulgaris*.)

THE two fore feet of this insect, placed very near the head, are short and broad, and, like those of the mole are contrived to help the insect in burrowing under ground. The Mole Cricket is very destructive in gardens,

as it attacks the roots of young plants, and causes them soon to rot and die. The female forms a nest of clammy earth, in which she lays from two to four hundred eggs. The nest is carefully closed up on every side, to secure the brood from the incursions of grubs and other subterraneous depredators. The song of the Mole Cricket is harmonious.

3. THE CRICKET. (*Gryllus domesticus*.)

THE domestic Crickets generally inhabit houses, selecting for their place of retirement the chimneys or backs of ovens; and feeding upon anything that comes in their way, flour, bread, meat, and especially sugar, of which they seem to be particularly fond. The chirping noise, which they make nearly without intermission, proceeds only from the males, who produce it by rubbing their legs against the base of their wing-cases.

Crickets are generally of a brown rusty colour, and the organ of vision appears in them to be very weak and imperfect, as they find their way much better in the dark than when dazzled by the sudden light of a candle. The Field Cricket (*G. campestris*) has the same form, but is of a different species to the House Cricket, and is black, with a fine gloss. Its noise is heard at a great distance, and is so similar to that of the grasshopper, that it is difficult to distinguish one from the other.

ORDER III.—Hemiptera.

THESE insects have neither mandibles nor maxillæ, but in lieu of them they have a tubular articulated rostrum, adapted for suction. Insects thus formed are called haustellated. The four wings are all membraneous, but the outer ones are leathery at the base. Some of the species are without wings. The antennæ are very small, and sometimes scarcely perceptible. The metamorphoses of these insects are incomplete.



THE LANTERN FLY OF THE WEST INDIES.

(*Fulgora lanternaria.*)

THIS Lantern Fly is a nocturnal insect, with a hood, or bladder on its head, which appears like a lantern in the night; but by day it is clear and transparent, and very curiously ornamented with red and green stripes. Such a shining light issues from this part of the insect, that it is possible to read by it. The wings and whole body are elegantly adorned with a mixture of red, green, yellow, and other splendid colours. The creature can contract or dilate the hood, or bladder, as it pleases.

These flies are as luminous as a lighted torch, and reflect a lustre on all neighbouring objects. They are in continual motion during the night; but this motion is various and uncertain, alternately rising and sinking. They commonly hover about six feet from the ground. It is said that not a night in the year occurs in which they are not seen; in winter more frequently so than in summer. Neither rain nor snow prevents their appearance. From all these circumstances, many have erroneously supposed them to be merely the ignis fatuus, or jack-of-the-lantern, which exhales from the marshy lands, over which it is seen to hover in the night.



THE COCHINEAL INSECT. (*Coccus cacti.*)

THE Cochineal insect is of the same genus as the scale

insect on the vine, which looks like a little bit of wool attached to the branch, but which, when pressed, stains the fingers with a red liquid. The Cochineal insect in the like manner affixes itself to the leafy stems of the nopal-tree, a kind of opuntia, or prickly-pear, common in Mexico and South America, whence the Cochineal used in Europe is principally imported.

When the Mexicans have gathered the Cochineal insects, they put them into holes in the ground, where they kill them with boiling water, and afterwards dry them in the sun; or they kill them by putting them into an oven, or laying them upon hot plates. From the various methods of killing them arise the different colours in which they appear when brought to us. While they are living, they seem to be sprinkled over with a white powder, which they lose when the boiling water is poured upon them, but preserve when killed in an oven. Those dried upon hot plates are the best.

The quantity of Cochineal annually exported from Mexico and South America is said to be worth more than five hundred thousand pounds sterling, a vast sum to arise from so minute an insect; and the present annual consumption of Cochineal in England has been estimated at about one hundred and fifty thousand pounds weight. The Mexicans think so highly of their trade in this insect, that the republic has adopted the nopal-tree as part of its arms.

It is for dyeing scarlet that Cochineal is chiefly in demand; but, although a peculiarly brilliant dye is now obtained from it, this substance gave only a dull crimson colour, until a chemist of the name of Kuster, who lived at Bow, near London, about the middle of the seventeenth century, discovered the art of preparing it with a solution of tin. Cochineal, if kept in a dry place, may be preserved without injury for a great length of time. An instance has been mentioned of some of this dye, one hundred and thirty years old, having been found to produce the same effect as though it had been perfectly fresh.



THE PLANT LOUSE, OR GREEN FLY.

(*Aphis.*)

THE Aphides are sometimes viviparous, and at other times oviparous, according to the season of the year. Those of the rose-tree have been particularly noticed, and of ten generations produced in one spring, summer, and autumn, the first generation was oviparous, the eight following viviparous, and the last oviparous. The first nine generations consisted of females only; but in the tenth there were some males. In this singular aberration from the common laws of nature, this insect is a remarkable anomaly. They multiply at such an extraordinary rate—the whole ten generations within three months, each comprising nearly one hundred individuals,—so that from a single *Aphis* ten thousand million millions may be produced in that short period.

The moss-rose, the hop, the vine, the apple-tree, the bean, the willow, and privet are all particularly liable to be infested with this insect; the various species of which take their names according to the plants on which they are usually found. The red tumours, commonly called galls, which are seen on the surfaces of leaves, especially on those of the willow, varying from the size of a lady-bird to that of a pigeon's egg, are produced by the *Aphis*, and contain thousands of small lice. The oval tubercles of this insect secrete a saccharine fluid, of which ants are very fond; and it is this fluid dropped upon the adjacent leaves, or the extravasated sap flowing from the wounds

caused by the punctures of the insects, which is known under the name of honey dew.

After a mild spring, most of the species of *Aphis* become so numerous as to destroy all the young shoots of the plants on which they are found. No successful mode of destroying them has yet been discovered, but the best remedy against them is to wash the infested shoots with tobacco water or soap lees; and to repeat the operation when any *Aphides* are seen.

ORDER IV.—Neuroptera.

THESE insects have four transparent wings, strongly and beautifully varied, so as to resemble net-work. The mouth has mandibles and maxillæ. The abdomen of the female has neither ovipositor nor sting.



THE ANT-LION. (*Myrmeleon formicaleo*.)

THIS insect is hatched from an egg laid in soft moving ground, or sand; the larva soon increases in size, and assumes the shape of a small spider, with this difference, that the legs are constructed in such a way, that it can only proceed backwards or sideways. The abdomen is very large and fleshy; and the head, which is small, is

armed with two long jaws like horns, somewhat resembling those of the stag-beetle. What must create our utmost admiration is, that this insect, which can only move in a retrograde direction, is doomed by Nature to feed upon flies and ants, the quickness and agility of which would at all times deprive him of his prey, were he not endowed with an uncommon instinct, which prompts him to the following stratagem. He makes a kind of funnel-shaped hole in loose earth or sand, and placing himself at the bottom of it, waits there with the utmost patience for several days, till an incautious ant or giddy fly falls into the deathful pit. Then all his skill is put in requisition; he throws out, by the shaking of his large jaws, a great quantity of sand upon the insect, to prevent its climbing up the steep sides of the hole; and when the prey appears strong and nimble, he gives such a general commotion, that the whole construction crumbles down, and the unfortunate insect, overwhelmed with the ruins, falls into the jaws of the Ant-lion, which open like a pair of forceps. When the Ant-lion has sucked out the blood and inside of his prey, he takes it upon his head, and, by a sudden jerk, throws the carcase to a distance from his abode. When the larva has attained its full size, it spins for itself a cocoon of white shining silk, with an external covering of sand. In about three weeks there bursts from this pupa case a slender-waisted winged insect, which, after fluttering about for a few weeks, and depositing eggs in the sand, resigns its life. The winged insect resembles a beautiful dragon-fly; it has a head of a chestnut colour; the body is of a pearly grey, the legs short, and the wings, which resemble the finest lace, are beautifully marked with dark lines and spots. This fly is often seen fluttering about the sides of roads and dry banks exposed to the east, in the months of June and July; it continues for a little time, and then entirely disappears. The Ant-lion is very rare in this country; but in France and Italy there is not a bank on the sides of a public road, or a sandy ridge at the foot of an old wall, which does not harbour a great number of these insects.



THE GREAT DRAGON FLY. (*Libellula grandis*.)

THIS genus of insects is well known to every one. The larva lives in the water, and wears a kind of mask, which it moves at will, and which serves to hold its prey while it devours it. The period of transformation being come, the larva goes to the water-side, and fixes on a plant, or sticks fast to a piece of dry wood, in which position it remains during its pupa or nymph state. When the period for the final transformation arrives, the skin of the nymph, grown parched, splits at the upper part of the thorax, the winged insect issues forth gradually, throws off its slough, expands its wings, flutters, and then flies off with gracefulness and ease. The elegance of its slender shape, the richness of its colours, the delicacy and resplendent texture of its wings, render it a beautiful object. It is in length about four inches.

The female deposits her eggs in the water, from which spring the larvæ, which afterwards undergo the same transformations.

The *Day Fly*, (*Ephemera*,) so called on account of the shortness of its life, is a small insect originating from a larva residing in rivers. After remaining several months in the creeping state, a nymph is formed, from which the perfect insect bursts, three or four hours after midday, into the fly form, and dies soon after. This fly has the singular

characteristic of casting off its entire skin, very soon after it has attained its perfect state; and hence persons often find their clothes covered with its white exuviae.

ORDER V.—Hymenoptera.

IN this order the wings are neither so large nor so strongly veined as in the previous. The mouth is furnished with mandibles, maxillæ, and an upper and lower lip; and the abdomen of the female is always terminated either with an ovipositor or a sting. The metamorphosis of these insects is complete.

This order contains the Bees, of which there are three kinds, the working Bees, the Drones, and the Queen. The Queen, or Parent Bee, is the soul of the hive; to her all the rest are so attached, that they will follow her wherever she goes. If she dies all their labours are at an end, an universal mourning ensues, and all her subjects reject their food and follow her. Should, however, a new queen arise, before this catastrophe attends the hive, joy renovates their spirits, and their toils are renewed. This has been observed by removing the chrysalis of a Queen Bee from one hive to another which had lost its own empress. But this attachment is only in proportion to the utility she affords to the commonwealth. She is so prolific as to lay fifteen or eighteen thousand eggs, which produce about eight hundred males or drones, four or five Queen Bees, and the rest Working Bees or Neuters. Their cells differ in size; the largest being for the males, the royal cells for the Queens, and the smallest for the neuters. The Parent Bee deposits in these cells such eggs as will produce the species for which they are respectively destined. In two or three days the eggs are hatched, when the Neuters turn nurses to the rest, whom they feed, most tenderly, with bee-bread and honey. After twenty-one days, the young Bees are able to form cells, with such indefatigable activity, that they will then do more in one week than during all the rest of the year. Sometimes there are


Bees who, less laborious than the others, support themselves by pillaging the hives of the rest; upon which a battle ensues between the industrious and the despoiling insects. Frequently contentions will arise among them, when a new colony seek their habitation in a hive already occupied. Their foes are the wasp, the hornet, and various kinds of birds.

The Bee collects the honey by means of its proboscis, or trunk, which is a most astonishing piece of mechanism, consisting of more than twenty parts. Entering the hive, the insect disgorges the honey into cells, for winter subsistence; or else presents it to the labouring Bees. A Bee can collect, in one day, more honey than a hundred chymists could extract in a hundred years.

When they begin to fill their hive, they divide into four parties: one is deputed to search the fields for materials, another to work on these materials, a third to polish the rough work of the cells, and a fourth to provide food for the labourers. There are waiters always attending to serve the artisan with immediate refreshment, lest his work should be retarded by his going to gather it for himself.

So expert are the Bees, that a honeycomb, composed of a double range of cells backed one against another, and which is a foot long, and six inches broad, is completed in one day, so as to contain three thousand Bees. The cells are most curiously composed of little hexagonal sides, which exactly conform to the similar portions of the adjoining cells. At the entrance of every cell the Bee architect forms a ledge, which fortifies the aperture, and prevents the injuries it might receive from the frequent ingress and egress of the Bees.

Bees produce honey, which they lay up for winter consumption; wax, of which they form their cells, and a substance called bee-bread, which they extract chiefly from the pollen of the flowers, and which they use for feeding their young.



In the above cut are given a representation of, first, the *Queen Bee*, placed on the left hand side; and, on the right side of her, the *Drone*; below them both is represented the *Working Bee*.



THE WASP (*Vespa vulgaris*)

Is a very fierce, dangerous, and rapacious insect; it is much larger than the bee, and furnished with a powerful sting. The abdomen is striped with yellow and black. All kinds of Wasps make curious nests; some attach them to the beams of a barn or other building, or place them in the hollow of a large tree, but the common Wasp digs in a hole in the ground. Wasps do not construct cells with the same care and accuracy as the bee; and as they do not lay up a store of honey for their support during winter, they mostly die at that season; and the few that live remain in a torpid state till spring. Their sting is very large; and the poisonous liquor of it, when introduced into the human body, excites inflammation, and creates very considerable pain. Wasps have no queen and do not swarm.



THE ICHNEUMON FLY. (*Pimpla persuasoria*.)

THE mouth of this insect has jaws, without any tongue. The antennæ contain more than thirty joints; and the abdomen is generally joined to the body by a pedicle. The ovipositor is enclosed in a cylindrical sheath, composed of two valves.

One distinguishing and striking characteristic of all the species of this kind of fly is the almost continual agitation of their antennæ. The name of Ichneumon has been applied to them from the service they do us by destroying caterpillars, plant-lice, and other insects; as the Ichneumon or mangouste destroys the crocodile in the East. The tip of the abdomen of the females is armed with an ovipositor, visible in some species, though not in others; and this instrument, though so fine, is able to penetrate through mortar and plaster. The female fly uses it to deposit her eggs in the body of other insects when in the caterpillar, or pupa state; so that the young as soon as they are hatched, may feed upon the caterpillar, penetrating to its very entrails. These larvæ, however, contrive to suck out the nutritious juices of their prey, without attacking its vitals; for the caterpillar continues to live for a long time, so as to afford them food till they have attained their full size. It is not uncommon to see caterpillars fixed upon trees, as if they were sitting upon their eggs; when it is afterwards discovered that the lar-

væ, which were within their bodies, have spun their threads, with which, as with cords, the caterpillars are fastened down, and so perish miserably.

“A friend of mine,” says Dr. Derham, “put about forty large caterpillars collected from cabbages on some bran and a few leaves in a box, and covered it with gauze to prevent their escape. After a few days we saw, from more than three-fourths of them, about eight or ten little caterpillars of one of the Ichneumon flies come out of their backs, and spin each a small cocoon of silk, and in a few days the large caterpillars died.”

The Ichneumons performed great service in the years 1731 and 1732, by multiplying in the same proportion as the caterpillars, and their larvæ destroyed more of these destructive creatures than could any efforts of human industry.



THE LABOURING AND SOLDIER ANT.

(*Formica rufa.*)

THE colour of the Ant is in general a dark red or brown, with a fine gloss on the abdomen. They are like the bees, divided into three tribes, male, female, and neutral. The females and neuters are furnished with stings for their defence; the males are wholly destitute of them. The males and females are in proper season furnished with wings, but the neuters have none, and they are doomed always to labour and drudgery on the hill. This hill is constructed with considerable art and labour; it is composed of leaves, bits of wood, sand, earth, and gum from the trees, which are all united into a mass, perforated with galleries to give access to the numerous cells which it contains. From this hill there are several paths, worn by

the constant passing and repassing of these creatures; and it is worthy the admiration of the naturalist to consider how busy the whole legion appears, in bringing bits of straw, dead bodies of other insects, or in carrying away their eggs, if any danger threatens their republic. Their sense of smell is very keen, and they discover at a great distance any food they may be in search of.

ORDER VI.—*Lepidoptera*. The Moths and Butterflies.

THE insects included in this order are all remarkable for their beauty. Their wings are membranous and veined, like those of the dragon flies and their allies, but instead of being naked they are covered by close-set scales of the most delicate texture and most brilliant colours. The mouth is furnished with a spiral trunk or tongue, by which nectar is sucked from the flowers; but in other respects it only differs from the mouths of the masticating mandibulated orders in the smallness of its parts. The antennæ vary in the different kinds: but those of all the diurnal lepidoptera, or butterflies, are terminated by a small inflation or knob; while those of the nocturnal species, or moths, taper to a point, and are often feathery, or comb-shaped. The transformations of the species belonging to this order are all complete.

Over the larvæ of this order the ichneumons reign with undisputed sway; attacking all indiscriminately, from the minute insect that forms its labyrinth within the thickness of a leaf, to the giant caterpillar of the hawk-moth. The most useful of all, however, the silkworm, appears, at least with us, to be exempted from this scourge. De.Geer, out of fifteen larvæ that were mining between the two cuticles of a rose-leaf, found that fourteen were destroyed by one of these parasites; the most useful species of which is that which destroys the clothes-moth.



THE MOTH WITH ITS CHRYSALIS AND CATERPILLAR.

THE larva of all the lepidoptera is a Caterpillar composed of twelve ring-like segments, exclusive of the head, which is harder than the other parts, and always of a deeper colour than the body. Each Caterpillar has nine breathing-holes on each side; and each of the three segments nearest the head are furnished with a pair of short legs, ending in a kind of claw, which are the true legs of the insect. The Caterpillar has, however, eight or ten other legs on the hinder segments of its body. The head has twelve eyes, and two very short conical antennæ; and the mouth is furnished with two strong mandibles, two maxillæ, a labrum, and four palpi.


The habits of Caterpillars differ; some, which are called surveyors, fix themselves to a leaf with their fore and hind feet close together, and their bodies bent in the form of a loop; others assemble in great numbers, and spin themselves a silken tent, under which they live; and others stretch themselves out like a piece of dry stick, which they very much resemble. Many make themselves cocoons; but others have no other covering in the pupa state than a smooth shining skin, or a dark mummy-like cerement. The chrysalis of a butterfly is generally angular, and that of a Moth cylindrical.



TORTOISE-SHELL BUTTERFLY.

(*Vanessa urticae.*)

THE Caterpillar which feeds on the nettle is about an inch in length, covered with bristles, and of a reddish brown colour. After having changed its skin three times when in the shape of a Caterpillar, it crawls up to a branching part of the stalk; and, hanging itself by the hinder part or tail, swells and bursts in such a curious way, that the Caterpillar's skin drops to the ground, and the chrysalis, or aurelia, remains suspended; till after a fortnight of torpor it bursts its skin again, and escapes in the vast plains of the air, under the beautiful form of a variegated Butterfly. The golden line which shines through the pupa case of this Butterfly are supposed to have suggested the words chrysalis and aurelia, both of which signify golden. The wings of the perfect insect are about two inches in extent, of a deep orange colour above, and their base and hinder margin black, with a series of blue crescents. These Butterflies, which are very common in England, appear in spring, and at the end of June and beginning of September.




THE CABBAGE BUTTERFLY.

(*Papilio*, or *Pieris Brassicæ*.)

WHEN the colewort and cauliflower begin to heart, the perfect insect of this Caterpillar is found depositing her eggs upon the leaves. The heat of the sun soon vivifies them and brings forth the Caterpillars, which immediately proceed to consume the vegetables on which they received being. They bear the heat of the sun without inconvenience; but cannot endure long rains, and in wet weather they soon disappear. There are several species of this Butterfly, but the common white, with a black spot on each of the under wings, is the earliest seen in our gardens. It lays its eggs in May; and its Caterpillars, which are soon hatched, feed together till the end of June, when they go into the pupa state, from which the perfect Butterfly appears in July. The eggs laid by the second brood of Butterflies produce Caterpillars which feed during the remainder of the summer, and remain in the pupa state all winter, to be hatched the following spring.

From the astonishing fecundity of these insects, it may be wondered that they do not, in the course of time, completely overspread the face of the earth, and totally consume every green plant. This would certainly be the case, if Providence had not put a check to their progress. One of the kinds of the ichneumon-fly deposits her eggs

within the caterpillar of this Butterfly, and they are there hatched. In their larva state they continue preying on the vitals of the animal; they then pass to the pupa condition, and eventually emerge the perfect insects. So greatly are we indebted to this apparently contemptible little parasite, for keeping down the increase of an insect which would otherwise become a serious and alarming evil.




THE MAGPIE, OR CURRANT MOTH.

(*Phalæna, or Abraxas Grossulariata.*)

THIS Moth is said by most entomologists to lay its eggs in the ground; but others assert that it lays them on the leaves, and covers them with a kind of white silk, which it spins for the purpose. The Caterpillar is one of the kind called loopers, and is very destructive. The chrysalis is naked and shining; and its colour is a bright yellow with black bands. The Moth is white spotted with black, and hence its name of Magpie.


The black and white caterpillar of this Moth is very destructive to currant and gooseberry bushes, and in some seasons particularly so. Mr. Kirby, especially, cites the devastations at Hull in the spring of 1814. He also confirms Boerhaave's assertion, that the severity of winter has no effect in destroying the larvæ of those insects, as these abounded even more after a winter, when Fahrenheit stood at zero, than after a winter which was remarkably mild.



THE WINTER MOTH. (*Geometra brumata*.)

THIS Caterpillar delights in newly opened leaves; it is not so ravenous as many others, making long intervals between its meals, but it seldom quits a leaf until it has entirely consumed it. The colour is very elegant. The upper part of the body is of a fine yellowish green; but it is by no means so beautiful after as before feeding, its skin being so thin as to imbibe the hue of whatever food it eats. They are also called looper Caterpillars, because when they crawl they draw their hind and fore-feet together, so as to form their bodies into a loop. They go into the pupa state towards the end of June, burying themselves for that purpose in the earth; and in November or December the perfect insect is brought forth.

It is evident that they possess great muscular power, and hence their positions during repose are very striking. Fixing themselves by their anal feet alone, they extend their bodies in a straight line, holding it in that position for a long time. This, together with their obscure colours, and the warts on their bodies, renders it often difficult to distinguish them from the twigs of the trees on which they feed. When alarmed these Caterpillars have the instinct to drop from the leaves, and suspend themselves by a thread, which enables them to remount when the danger is over.



THE BLACK ARCHES MOTH.

(*Bombyx*, or *Psilura Monocha*.)

THESE Caterpillars feed on the leaves of red roses and gooseberry bushes, and, in short, on almost every kind of plant. The chrysalis is of a shining brown, and is enveloped in a slight cocoon. When they wish to change from one situation to another, they ascend by attaching themselves to the bough with their feet, by which they raise the body like a serpent, and thus gain their desired situation. They hold themselves so fast by their feet that they can scarcely be taken from the part to which they adhere.



THE SILKWORM. (*Bombyx mori*.)

WITHOUT entering into a very minute description of this

Caterpillar, we shall confine ourselves to what we think will be at once more interesting and more useful; as the Silkworm is an insect of universal service, and not of singular beauty, we are induced to prefer giving an account of its utility, rather than any elaborate description of its figure or colour.

This larva feeds on the leaves of the mulberry-tree, and when first produced, is extremely small, and entirely black. In a few days it appears in a new habit, which is white, tinged with the colour of its food; and before it goes into its chrysalis state, it changes its skin several times. When full grown it spins its cone of silk, which is its cocoon, in the same manner as other insects. The Moth possesses no beauty.

The art of manufacturing silk was known to the ancients. We are informed that, in the third century, the wife of the Roman emperor Aurelian entreated him to give her a robe of purple silk, which he refused on account of its enormous price.

It is not certain at what precise period the manufacture of silk was first introduced into England; but in the year 1242, we are told that part of the streets of London were covered or shaded with silk, for the reception of Richard, the brother of Henry III., on his return from the Holy Land. In 1454 the silk manufactures of England are said to have been confined merely to ribbons, laces, and other trifling articles. Queen Elizabeth, in the third year of her reign, was furnished by her silk-woman with a pair of black knit silk stockings, which she is stated to have admired as "marvellous delicate wear;" and after the using of which she no longer had cloth ones as before. James the First, whilst King of Scotland, requested of the Earl of Mar the loan of a pair of silk stockings to appear in before the English ambassador, enforcing his request with the cogent appeal, "For ye would not, sure, that your king should appear as a scrub before strangers."

THE CLOTHES MOTH. (*Tinea vestianella*.)

THE larva of this little Moth is well known from the damage it commits in woollen cloth and furs. These sub-

stances constitute the principal support of the Caterpillar, and therefore the parent is, by its natural instinct, directed to deposit its eggs in them. As soon as it quits the egg, the Caterpillar begins to form for itself a nest: for this purpose, after having spun a fine coating of silk immediately around its body, it eats the filaments of the cloth or fur, close to the thread of the cloth, or to the skin. This operation is performed by its jaws, which act in the manner of scissors. The pieces are cut into convenient lengths, and applied, with great dexterity, one by one, to the outside of its case; and to this it fastens them by means of its silk. Its covering being thus formed, the little Caterpillar never quits it but on the most urgent necessity. When it wants to feed, it puts out its head at either end of its case, as best suits its conveniency. When it wishes to change its place, it puts out its head and its six fore legs, by means of which it moves forward, taking care first to fix its hind legs into the inside of the case, so as to drag it along. After having changed within its case into a chrysalis, it issues, in about three weeks, a small, winged, mealy-looking Moth, of silvery drab colour, well known to almost every mistress of a family. The best mode of destroying this insect, when in the cloth, is to place a saucer of oil of turpentine with the articles affected in a close place, when the vapour raised by the warm air will immediately destroy it. Should the Caterpillar be old and strong, it may be necessary to brush the clothes with a brush, the points of which have been dipped in turpentine. Camphor wrapped up with furs will protect them from the Moth.



THE SMALL SILVER-COLOURED BOOK-
WORM MAGNIFIED. (*Crambus pinquinella*.)

“ Insatiate brute, whose teeth abuse
The sweetest servants of the muse !

His roses nipt in every page,
 My poor Anacreon mourns thy rage ;
 By thee my Ovid wounded lies ;
 By thee my Lesbia's sparrow dies ;
 Thy rabid teeth have half destroy'd
 The work of love in Biddy Floyd ;
 They rent Belinda's locks away,
 And spoil'd the Blouzelind of Gay ;
 For all, for every single deed,
 Relentless justice bids thee bleed.
 Then fall a victim to the Nine,
 Myself the priest, my desk the shrine."

PARNELL.

THIS small, white, silvery-shining Worm, or Caterpillar is found among books and papers, corroding and eating holes through the leaves and covers : it appears, to the naked eye, a small, glittering, pearl-coloured Moth, which, upon the removing of books and papers in the summer, is often observed to go into some corner, where it may the better protect itself from danger. Its head is blunt, and its body tapers towards the tail, being divided into fourteen segments, having the appearance of shells or shields. These shells are again covered or tiled over with a multitude of thin transparent scales, which, from the multiplicity of their reflecting surfaces, make the whole animal appear of a perfect pearl colour. This insect is furnished on each side of its head with a cluster of eyes ; and each of these clusters is beset with a row of small bristles, very like the cilia or hairs on the eyelids ; and, perhaps, serving for the same purpose. It has two long palpi, curiously ringed or knobbed, having at each knob small hairs or bristles, here and there, like knots of fringe : besides these, it has two shorter palpi, knotted and fringed like the former. It has three tails, in every particular resembling the two longer palpi that grow out of the head. The body is covered with sharp-pointed bristles, like spears.

ORDER VII.—Diptera, or Flies.

THIS order is characterised by having only two wings, which are transparent, and which have two little move-

able bodies, called halteres or balancers, placed close behind them. The head is covered with numerous eyes; and the mouth is furnished with a bristly proboscis or sucker. The legs are very long in proportion to the body, and are in many species terminated by two or three small cushion-like expansions, which are supposed to enable them to walk on glass. Each foot has also two hooks or claws.



THE HOUSEFLY. (*Musca domestica*.)

THIS insect lays its eggs in sinks, or any other place where there is decaying vegetable matter tolerably moist. The larvæ, or maggots, are thick and fleshy, without legs, but having the mouth furnished with hooks, by means of which they drag themselves along when they wish to move. They go into the pupa state without throwing off the skin of the maggot; and when the perfect insect appears, it forces off a kind of cap from one end of the pupa case, in order to make its escape. There are two other varieties of this insect, one called *Musca Grossa*, which is as large as a humble bee, and is chiefly found among flowers and bushes; the other, known as the Meat-fly, or Blue-bottle (*Musca vomitoria*.)

THE GNAT. (*Culex pipiens*.)

THIS is an insect which deserves the observation of the naturalist, not only for the very curious conformation of its proboscis, (which so quickly and powerfully penetrates into our flesh, and through which it sucks our blood into its body,) but also for the several metamorphoses it undergoes before it arrives at its winged state. The Gnat deposits its eggs upon the slimy surface of stagnant water, and sets them upright one against another, in the form of a small boat: after floating upon the water for several days in the first warm weather in the spring, as soon as

they feel the time of hatching, the maggots, which the eggs contain, precipitate themselves to the bottom, and there feed for a fortnight; after which time they undergo another transformation. The result is a curious animal, with a kind of wheel in incessant motion, by means of which it imbibes the air at the surface of the water, to which it is obliged to repair every moment: having passed about ten days in this state, its increase being at an end, it assumes another form, keeps longer near the surface, and at last the outer skin bursts, and the winged insect, standing upon the *exuvix* it is going to leave behind, smooths its new-born wings, springs into the air, and begins its depredations. The fecundity of the Gnat is so remarkable, that in the course of one summer they might increase to the amazing number of five or six hundred thousands, if Providence had not ordered that they should become the prey of birds, who by this means prevent their multiplying more than they generally do. These insects are very annoying from their blood-sucking propensities; and as the sucker is horny at the tip, it inflicts a severe wound, into which the insect emits a small quantity of poison, which occasions the pain and inflammation always felt from a Gnat bite.

ORDER VIII.—Suctoria.

THESE insects are without wings. The mouth is furnished with a trunk or beak, formed to wound as well as to suck.



THE FLEA (*Pulex irritans*)

Is one of those little creatures with which want of cleanliness in mankind is punished. It is one of the most

annoying insects that infest the human race, as, by its leapings, it often escapes being caught. It is oviparous, and the egg, which is hardly discernible by the naked eye, contains at maturity a small white worm, beset with hairs. This worm soon spins for itself a little silk cocoon, from which the perfect insect issues. The Flea is an active, troublesome, blood-thirsty insect; it has a small head, large eyes, and roundish, but compressed body, which is covered with a kind of armour resembling the tortoise-shell in colour and transparency. The plates of which this skin is composed are also armed with spines or bristles. It has six legs, two of which are much longer than the others, in order to enable the insect to make such wondrous leaps, as to raise the body above two hundred times its diameter. It is observed, that in one day the Flea will devour above ten times its own bulk. The great strength and agility of the Flea are well known, from the exhibition of the industrious Fleas.

BOOK VII.—RADIATA.



THE STAR-FISH. (*Asterias.*)

THIS animal is often found adhering to rocks on the sea-shores. The common species is furnished with five rays,

and is of a yellow or red colour. It has a slow and progressive motion, and is often found on the beach among sea-weeds after a storm; but when caught it very soon decays, and can scarcely be carried a hundred yards before it becomes soft and putrid.

Mr. Bingley describes an animal of this kind, which he kept by him for some time alive; it had more than four thousand tentacula on the under sides of the rays. These it frequently retracted, and again pushed out, as a snail does its horns; and by means of them it was enabled firmly to adhere to the dish containing the salt-water in which it was kept. Whenever he touched the tentacula with his finger, all those of that ray or limb were gradually withdrawn, but those of the other rays were not in the least affected by it.

Like the other radiata, these animals breathe by means of a kind of branchiæ or gills, which consist of a fringed substance extended along each ray, and communicating with the stomach. These all unite under a small circular and striated operculum, or cover, which may be observed on the back.

THE SEA-URCHIN. (*Echinus*.)

THIS animal, which lodges in the cavities of rocks just beneath low-water mark, on most of the British coasts, is nearly of a globular shape, not much unlike that of an orange, having its shell marked into ten partitions, with rows of projections like beads, which divide it. On the outside of the shell there are a great number of sharp, moveable spines, of a dull violet and greenish colour, curiously articulated, like balls and sockets, with tubercles on the surface, and connected by strong ligaments to the skin or epidermis, with which the shell is covered. The mouth is situated in the under part, and is armed with five strong and sharpened teeth. The animal can move from place to place by means of its contractile tubular feet and its spines; but its movements are slow and laborious. So tenacious of life are the Sea-urchins,

that the ancients, according to Appian, believed that the body retained life even when cut to pieces.

“ Sea-urchins, who their native armour boast,
 All stuck with spikes, prefer the sandy coast.
 Should you with knives their prickly bodies wound,
 Still the crude morsels pant upon the ground ;
 You may e'en then, when motion seems no more,
 Departing sense and fleeting life restore.
 If in the sea the mangled parts you cast,
 The conscious pieces to their fellows haste ;
 Again they aptly join, their whole compose,
 Move as before, nor life nor vigour lose.”

In Marseilles, and some other towns on the continent, the Sea-urchin is exposed for sale in the markets, as oysters are with us, and is eaten boiled as an egg. The Romans adopted it as food, and dressed it with vinegar, mead, parsley, and mint.

ZOOPHYTES.

ZOOPHYTES hold a middle station between animals and vegetables. Most of them, deprived altogether of the power of locomotion, are fixed by stems that take root in crevices of rocks among sand, or in such other situations as Nature has destined for their abode: these, by degrees, send off branches, till at length some of them attain the size and extent of large shrubs. The Zoophytes were placed by Linnæus in two divisions. The stony branches of the first division, which have the general appellation of coral, are hollow and full of cells, which are the habitations of animals. The next division consists of such zoophytes as have softer stems, and are, in general, not merely inhabitants of a stem or branches, but are themselves in the form of a plant. Those of this division which are best known are the corallines, the sponges, and the polypes. Modern naturalists divide the Zoophytes into numerous families.



Magnified branch, exhibiting the Animals.

Gorgonia Nobilis.

THE CORAL, AND CORALLINES.

THE Coral, or Gorgonia, is a hard, stony, branched, and cylindrical substance, which is formed at the bottom of the sea by animals called polyps, or, to use the Latin and now established term, *polypi*. The whole form a living mass, or polypidom, all the polypi in which are united under one skin, and have one common stomach. Each of these polypi resides in a distinct cell; they are generally dormant during winter, and like the blossoms of plants, push forth buds, and expand in the summer season. The stems and branches of the Gorgonia, which are of a somewhat horny and flexible nature, may be considered as the true skeletons of the

nests of the sea-polypi, being covered with a fleshy or pulpy substance, the interior surface of which is porous. These pores are the mouths or openings of the cells, in which the polypi are lodged; and it is the number, disposition, and varied structure of these, in addition to the general aspect of the plant-like nest of habitations, that constitute the distinguishing difference of the species. The figure of the animal, when it can be ascertained, forms the secondary character. Corals differ exceedingly in size, some being three feet in height, while others, in deep bogs, or in marine situations favourable to their growth, attain the gigantic height of ten or twelve feet; and from their number, as well as magnitude, their elegantly branched appearance, interwoven structure, and brilliant tints, they form a conspicuous portion of those vast sub-marine groves of Coral, which are seen by navigators in the tropical regions of the globe.

The bone of the Red Coral constitutes that beautiful and much esteemed production, the true or red coral of the jewellers. It is found in the Mediterranean, Adriatic, and Red Sea, and appears to be nowhere more abundant than in the seas about Marseilles, Corsica, Sicily, the coasts of Africa, and in the vicinity of Barbary; where the Coral fisheries are carried on with great spirit, and prove very lucrative. It is equal in hardness and durability to the most compact marble; and these qualities, in addition to its beautiful texture and colour, have rendered it valuable in all ages. Thus in the book of Job, "No mention shall be made of corals, or of pearls; for the price of wisdom is above rubies."

The mode of obtaining Coral is by a very simple machine, consisting of two strong bars of wood or iron, tied across each other, with a weight suspended from their centre of union. Each of the bars is loosely surrounded, throughout its whole length, with twisted hemp; and, at the extremity, there is a small open net. The machine is suspended by a rope, and dragged along those rocks where the Coral is most abundant; and such as is broken off either becomes entangled in the hemp, or falls into the nets.

Coral is bought by weight, and its value increases according to its size. Beads of large size are worth about forty shillings an ounce, whilst small ones do not sell for more than four shillings. Large pieces of Coral are sometimes cut into balls, and exported to China, to be worn as insignia in the caps of officers of state. These, if perfectly sound and of good colour, and upwards of an inch in diameter, have been known to produce in that market, as much as three to four hundred pounds sterling each. There are extant many beautiful pieces of sculpture in coral, as this substance has in all ages been considered an admirable material on which to exhibit the artist's taste and skill. Probably the finest specimen of sculptured Coral yet known, is a chess-board and men, in the palace of the Tuilleries.


The Chinese have, within the last few years, succeeded in cutting Coral beads of much smaller dimension than has hitherto been effected by any European artist. These, which are not larger than small pins' heads, are called Seed Coral, and are now imported from China into this country in very considerable quantity for necklaces. Nearly the whole of the Coral of commerce is of a red colour, white Coral being considered of little value. There are modes by which Coral may be so exactly imitated, that without a close inspection, it is sometimes impossible to detect the counterfeit.

In the Pacific Ocean several of the Coral reefs are extremely beautiful, and the voyager is astonished with the curious and fantastic forms of the various marine productions of which they are composed. Wheat-sheaves, mushrooms, cabbage-leaves, with innumerable plants and flowers, are vividly represented by different kinds of Coral, and glow beneath the water in brilliant tints of brown and purple, white or green; each with a peculiar form and shade of colouring, equal in richness and variety to the most beautiful productions of the vegetable world. Corals and funguses start from between the fissures of the rocks; while large portions of the former, in a dead state, connected into a solid mass, of a dull white colour, compose the stone-work of the reef. Solid

masses, termed negro heads, of different dusky hues, and generally dry and blackened by exposure to the weather, are also occasionally conspicuous. Even these are not without ornament, for nature delights in the variety of her decorations. They are studded with small shells, and beautifully marked with outlines expressive of their origin. The edges of the reefs, particularly those exposed to the waves, partake of a considerable degree of lightness, and form small coves and caverns, the resort of live corals, sponges, sea-eggs, and trefangs, or sea traces, (valued in China, for their invigorating quality,) and enormous cockles, which are scarcely to be distinguished from the rock, excepting when they suddenly close their shells, and discharge living fountains, which rise to the height of four or five feet.

With regard to the formation of coral reefs, it has been conjectured, from the appearance of the low islands in some parts of the South-Sea and Indian Ocean, (where they occur in rows or groups, while they are totally absent in other parts of the same seas,) that Coral insects rear their habitations on marine shoals, or, to speak more properly, on the top of sub-marine mountains. They evidently adhere together by virtue, either of the glutinous nature of the insect, or some property in the water, and increase by the labours of successive generations, both in extent and height. Floating substances of every kind are caught by their stony tree-like fabrics, till at length a solid mass of rock is formed, which gradually advances to the surface of the water. The deposits of the ocean no longer tenaciously adhere, but remain in a loose state, and form what is termed by mariners a key upon the summit of the reef; while the sea, by throwing up sand and mud on the top of these animal rocks, progressively raises them above its level. The new island, for such it may now be called, is soon visited by sea-birds; plants successively appear, and carpet the sterile soil with a luxuriant covering. As these decay, vegetable mould is gradually deposited; cocoa-nuts, or some floating seeds, flung on shore by the impetuosity of the waves, take root, and soon begin to grow; land-birds, attracted by

the verdant appearance of the bank, fly thither in quest of provisions, and deposit the seeds of shrubs and trees ; every high tide and every gale adds some new treasure : the appearance of an island is gradually assumed, and at length man comes to take possession.



CORAL INSECTS, MAGNIFIED.


1. Coralline of the *Astrea annanas*.
2. Animal of the *Caryophylla solitaria*.
3. Animal of the *Tubipora musica*.

4. Animal and dwelling of the *Cellepora hyalina*.
5. Animal and central axis of the *Gorgonia patula*.

THIS elegant species is highly ornamental in collections of natural history ; and one of them, the *Corallina Officinalis*, is valuable in medicine. Fifty-seven species, varying in size and structure, are now known ; some of which are extremely rare and beautiful.

The south of Devon presents an exquisite variety of these interesting marine productions. They are often left by the receding tide upon the shore, and float on the surface of the waters, either attached to a group of

matted sea-weed, fragments of wood, or the empty muscle-shell.



CORALLINES are some of the most beautiful productions of the deep; they are closely allied to corals in their construction, and are one step higher in the scale of Zoophytic life, than their immoveable neighbour, the rock-adhering Sponge.

They attach themselves to stones, and other solid bodies, and are concretions, formed by the Polype insects which inhabit them. They are of a plant-like form, with elevated branches, and exhibit an elegant appearance from their symmetry and proportions, being subdivided, calcareous, and composed of little joints, like beads strung on a necklace; which peculiarity easily distinguishes them from the coral family. The joints are perforated, or full of minute cells; and in these the Polypi reside, the openings serving them as door-ways, through which to seize their prey. When a branch is

immersed in vinegar, the calcareous crust dissolves, and leaves the interior uninjured, by which means its beautiful formation may be easily examined.



SPONGE is a substance of a soft, light, porous, and elastic nature, which is found adhering to rocks at the bottom of the sea, in several parts of the Mediterranean, and particularly near the islands of the Grecian Archipelago; and which, in its native state, is filled with animal jelly. The general uses of Sponge, arising from its ready absorption of fluids, and distension by moisture, are well known and of great importance. It is collected from rocks, in water five or six fathoms deep, chiefly by divers. When first taken from the sea, it has a strong and fishy smell, from the animal matter it contains, of which it is divested by being washed in clear water. No other preparation than this is requisite previously to its being packed up for exportation and sale. The growth of Sponge is so rapid, that it is frequently found in perfection on rocks, from which, only two years before, it had been entirely cleared.

The creatures which inhabit the Sponge belong to the

class Vermes, order Zoophyte; and are fixed, and flexile, composed of net-like fibres, or small spines, and covered with minute mouths, by which they absorb and reject water.

As they are never designed to remove from their places of abode, they are provided with a dwelling capable of close adhesion to the surface of the rocks: and what is still more wonderful, as this peculiarity of construction forbids them to go in quest of prey, innumerable mouths are arranged in all directions, like fishing-nets.

The inherent chemical properties of this curious Zoophyte are very remarkable. When a Sponge has been immersed for fourteen or sixteen days in nitric acid (diluted with three parts of distilled water) it becomes nearly transparent, and when touched with ammonia, assumes a deep orange colour, inclining to a brownish red. But if much softened by the acid, the whole fabric immediately disappears, on being immersed in ammonia, and forms a deep orange-coloured solution. A Sponge, when boiled, gives out a considerable portion of animal jelly. The infusion of a small quantity of oak bark, causes this to fall to the bottom of the vessel, as a sediment, and so entirely changes the nature of the Sponge, that, when dry, it crumbles between the fingers; and, when moist, it may be torn like wetted paper. In this state we should naturally conclude that it is entirely useless: but no; the operations of chemistry resemble a magic wand. Boil the same in water, with caustic potash, its latent qualities will be called forth; and, behold, a deposition of animal soap!

THE LONG-ARMED, AND THE FRESH-WATER GREEN POLYPI. (*Hydroida.*)


THESE are two species, which will fully illustrate the nature of the whole tribe. They are found in clear waters, and may generally be seen in small ditches and trenches of fields, especially in the months of April and May. They affix themselves to the under-parts of leaves, and to the stalks of such vegetables as happen to grow in the same water; and feed on the various species of small

worms, and other aquatic animals within their reach. When any of these pass near a Polype, the latter suddenly catches it with its arms, and dragging it to its mouth, swallows it by degrees, much in the same manner as a snake gorges its prey. Two Polypi may occasionally be seen in the act of seizing the same worm at different ends, and dragging it in opposite directions with great force. It sometimes happens, that while one is swallowing the end it has seized, the other is employed in the same manner; and thus they continue swallowing, each his part, until their mouths meet. They then rest for some time in this situation, till the worm breaks between them, and each goes off with his share. But sometimes when the mouths of both are thus joined together a combat ensues, and the largest Polype usually swallows his antagonist; the animal thus swallowed, however, seems to be a gainer by its misfortune, as after it has lain in the conqueror's body for about an hour, it issues unhurt, and often in possession of the prey that had been the original cause of contention. The remains of the animal, on which the Polype feeds, are evacuated at the mouth, the only opening in the body. The species are multiplied by a kind of vegetation, one or two, or even more young ones, emerging gradually from the sides of the parent animal; and these young ones are frequently again prolific before they drop off; so that it is no uncommon thing to see two or three generations at once on the same Polype. But the most astonishing fact respecting this animal is, that if a Polype be cut in pieces, it is not destroyed, but is multiplied by dissection. It may be cut in every direction that fancy can suggest, and even into very minute divisions, and not only the parent stock will remain uninjured, but every section will become an animal. Even when turned inside out, it suffers no material injury; for, in that state, it will soon begin to take food, and to perform all its other natural functions.

M. Trembley, of Geneva, ascertained that different portions of one Polype could be engrafted on another. Two transverse sections brought into contact will quickly

unite and form one animal, though each section should belong to a different species. The head of one species may be engrafted on the body of another. When one Polype is introduced by the tail into another's body, the two heads unite and form one individual. Pursuing these strange operations, M. Trembley gave scope to his fancy by repeatedly splitting the head and part of the body; he thus formed hydras more complicated than ever struck the imagination of the most romantic fabulist.

Though so difficult to destroy by division, all the Polyypes, even those which form the corals, may be easily killed by depriving them of moisture, when they soon shrivel up, and the tissue of their skins is completely destroyed.



THE SEA POLYPL.

This interesting genus is very generally diffused, and while some are scarcely distinguishable from the seaweeds to which they adhere, others stand forth in all their


native beauty, and gaily decorate the rocks to which they cling.

Such, especially, are the Sea Anemones; an elegant genus, closely resembling the flower from which it derives a name. These when gathered will continue in great beauty for a considerable time, if kept in salt water, and fed with small pieces of fish. Their colours in different lights are as changeable as those of the chameleon.

There are several species of this family, all equally distinguished for their varied tints and elegant construction. The Sea Carnation presents the appearance of a long white fig, delicately wrinkled and curvated at the top, like the petals of a garden carnation; the Sunflower Anemone grows on rocks, and expands its saffron-coloured petals, like the stately flower to which it owes a name; and the common Sea Anemone (such as are gathered from off the rocks at Sidmouth) is distinguished by a red colour, and rough internal surface, while the centre is often of the purest white, elegantly marked with numerous carmine streaks. But the most extraordinary of the species is the Cluster Animal Flower, consisting of many tube-like bodies, surmounted by a mouth, which gently swell towards the upper part, and is surrounded by one or two rows of tentaculæ. These, on contracting, look like a circle of beads; while the lower extremity resembles that of a bulbous root, which closely adheres to the rock, and sends forth other tubes, that creep like roots in various directions.

AN
APPENDIX
OF
FABULOUS ANIMALS,

OUR object in the previous pages has been to combine interest with amusement, and to present truth unmixed with fable. Yet considering that some fictitious animals are conventionally recognized in poetry and painting, we have thought it desirable to subjoin an account of them. The Sphinx, the Dragon, the Unicorn, Pegasus, and the Centaur are so familiar to us, both in sculpture and fable, that some notice of these mythological creations seems indispensable.



THE SPHINX.

PROVIDENCE has ordered, that as the plains of Egypt are not visited by showers, they should be fertilized by the

overflowing of the Nile, which takes place annually, a little after the summer solstice. This phenomenon, the source of unfailling fertility in the vales of the Delta up to Memphis, and around the bases of the majestic and venerable pyramids, was of the greatest importance to the people of Misraim, from the far-famed Pharos to the frontiers of Ethiopia. It was therefore their interest to calculate correctly the season, the month, and nearly the hour, when the flood should begin; the more so, as the sudden invasion of the waters was dangerous to the inhabitants of the low lands, the meadows and the fens, and often destroyed the cottages, and drowned the flocks and the improvident villagers. The star Sirius was remarked to emerge from the blazing halo of the sun about the time of the rising of the Nile; it was a warning, and was accordingly called the Dog-star, as if barking from the heavens to apprise the inhabitants of the valleys of the impending rise of the waters. The Egyptian astronomers, to mark the period, combined the signs of the zodiac answering to the two months during which the overflowing took place. These signs happening to be Leo and Virgo, the mystical fancy of the ancient Egyptians united them in one, and thus formed the figure of the Sphinx, which is the head and breast of a virgin, and the body of a lion. This was a great enigma to the Greeks and Phœnicians, who travelled to Egypt; they saw the monster, but could not comprehend its meaning. On returning to their respective countries, they invented the fable of the Sphinx offering riddles at the gates of Thebes, and destroying those who could not unravel them; having probably been told by the supercilious sages of that nation, that they who could not guess the meaning of the Sphinx were to forfeit their life in atonement for their ignorance. Long afterwards, the real sense of the symbol was forgotten, and Egypt, in her superstition, began to worship the emblem, of which innumerable figures still exist in that once flourishing country.

The Sphinx has been introduced in heraldry to adorn the gorgets of those general officers who distinguished themselves against the French on the banks of the Nile;

it has also been adopted as an ornament in various decorations ; and two specimens, exquisitely wrought, are seen on the front wall of Syon House, at Brentford, the seat of his Grace the Duke of Northumberland.

This chimerical figure is generally represented as sitting and at rest ; a graceful attitude adopted by Egyptian sculptors, and imitated by the Greeks and Romans.



THE DRAGON.

THIS fabulous animal, which figures largely in ancient romances, was supposed to be the tutelary genius of fresh-water springs in the bosom of dark forests and enchanted rocks. Dragons were harnessed to the car of Ceres ; they were the guardians of the golden apples of the Hesperides, and of the golden fleece of Colchis ; and in several parts of the world set as protectors to the carbuncles and other precious stones, hidden at the bottom of wells and fountains. They are represented as scaly serpents, with webbed feet, and with wings similar to those of a bat ; having been, it seems, originally a hieroglyphic emblem of the dangerous influence of an undue combination of air and water. Thus the serpent Python was the allegory of a pestilence, originating from a union of mephitic air and moisture. They have been long supporters to the arms of the city of London, as if the guardians of the wealth which commerce brings hither from all the parts of the world. Four of them are placed in fanciful attitudes, and beautifully carved, on the pedestal of the monument of London.




THE WIVERN, WOLVERINE.

THIS fabulous animal somewhat resembles the dragon, only that, instead of four, it has two legs, which are webbed, and armed with claws. There is no doubt that this imaginary being was originally conceived in the brains of the poets and romancers, in times of chivalry, when the Crusaders overran the plains of Palestine and Assyria. The heat of the climate in some vales at the foot of the mountains, which intersect the deserts of those countries, was favourable to the breeding of all sorts of serpents, some of an immense size. The European soldiers of Godfrey and Richard, unaccustomed to such sights, were easily frightened, whenever they met those monsters on the sedgy banks of small lakes, under the shade of cedars and palm-trees, where they appeared as if posted to guard the sacred waters, so precious in so hot a country; and magnified in their idle tales, when inactive in camps, the bulk of the serpent they had seen. The castle of Lusignan, in the province of Poitou, was supposed to contain one of those winged serpents. It is a very ancient armorial bearing, and now stands as supporter to the arms of several illustrious houses.



THE COCKATRICE, OR BASILISK.

THE fruitful imagination of man knows hardly any bounds. The animal which bears the name of Basilisk was originally supposed to be a serpent, with a sort of comb or crown on its head: but that was not sufficiently marvellous. It was supposed also to be hatched from a cock's egg, upon which a snake had performed the office of incubation; and the animal had the head of a cock, and the wings and tail of a dragon. Hatched near a spring of water, the common resort of serpents, it was asserted, that, frightened at his own extraordinary shape, he soon precipitated himself to the bottom, whence, by the mortal look from his fiery eyes, he had the power of killing whoever dared to gaze at him. The name of *Cockatrice* alludes to its heterogeneous origin. This animal also is found among ancient armorial devices.



THE GRYPHON, OR GRIFFIN,

WAS originally an emblem of life. It was used to adorn funeral monuments and sepulchres. The upper part of this allegorical animal resembles the eagle, the king of the birds, and the rest the lion, the king of beasts; which is said to imply that man, who lives upon the earth, cannot subsist without air. In latter times it was supposed that the Gryphon was posted as a jailor at the entrance of enchanted castles and caverns where subterraneous treasures were concealed. Milton compares Satan in his flight to the Gryphon, in the following beautiful passage :

“ As when a Gryphon through the wilderness,
 With winged course o'er hill or moory dale,
 Pursues the Arimasian, who, by stealth,
 Had from his wakeful custody purloin'd
 The guarded gold ; so eagerly the fiend,
 O'er bog or steep, through strait, rough, dense, or rare,
 With head, hands, wings, or feet, pursues his way,
 And swims, or sinks, or wades, or creeps, or flies.”

The *Arimasians* were Asiatic wizards, who, by magic, used to obtain a knowledge of the places where treasures lay hidden. Their incessant wranglings with the Gryphons about gold-mines are mentioned by Herodotus and Pliny. Lucan says that they inhabited Scythia, and adorned their hair with gold ; that they had but one eye in the middle of the forehead, and lived on the banks of the gold-sanded river Arimaspes.

Virgil, in his eighth Pastoral, mentions this animal as if really existing, but does not give us any description of it; and Claudian, in his Epistle to Serena, alludes to the supposed fact of their keeping watch over masses of gold in the bosom of northern mountains.

THE PHŒNIX.

HERODOTUS, Pliny, and nearly sixty other classical authors, have related marvellous stories of this bird, all of which are of course fabulous. The Phœnix, they say, inhabits the plains of Arabia, and is about the size of an eagle, with gorgeous plumage of purple and gold. He is the only one of his kind in the world. At the approach of death, he builds himself a nest of aromatic herbs, and on it yields up his life. From his marrow proceeds a worm, which shortly becomes a young Phœnix, whose first duty is to discharge the obsequies of his sire. For this purpose he collects a quantity of myrrh, which he moulds into the shape of an egg, as large as he can conveniently carry, and then scooping it out, he deposits the body of his sire in the inside. Having stopped it up again with myrrh, he carries it to the Temple of the Sun in Egypt, where he devoutly places it on the altar. This is the only time that he is seen during his life, which lasts five hundred years. According to others, after preparing a funeral pile of rich herbs and spices, he burns himself, but from his ashes revives in all the freshness of youth.

From late mythological researches, it is conjectured that the Phœnix is a symbol of five hundred years, of which the conclusion was celebrated by a solemn sacrifice, in which the figure of a bird was burnt. His being restored to youth signifies that the new springs from the old.

THE MERMAID, OR SIREN.

THE existence of this animal, half a woman and half a

fish, has long been talked of, believed, disbelieved, and doubted. Homer is the first who speaks of such beings, which he styles *Sirens*; but we do not find that he gives any description of their shape; however, it was soon asserted that the Sirens were, as Horace, in his "Art of Poetry," describes them,

"Above, a lovely maid, a fish below."


The Sirens were three sisters, whose voice was so delightfully harmonious, and enticing, that no resistance could be made against its powerful charms; but "'twas death to hear," for they led the navigators and their ships to certain destruction among the rocks that bordered the dangerous coasts which they inhabited, near the shores of Italy.

The belief in the existence of Mermaids has been current at different periods; indeed, some years ago, several persons made depositions before a magistrate, that they had seen Mermaids come out of the sea and play on the rocks, but that they sprang into their element before they were able to secure them.

A creature, said to be a dried Mermaid, was exhibited in London about the year 1828; but it was afterwards discovered to be the body of a monkey artfully attached to the dried tail of a salmon.

THE KRAKEN.

THIS creature is another fabulous inhabitant of the sea. It is said to be three or four miles in breadth, and to live generally at the bottom of the sea, on the Norway coast. When it moves the commotion of the sea is so violent that it upsets boats and even small ships; and when it comes to the surface, it is generally mistaken for an island.



THE DOLPHIN.

THIS is the Dolphin of heraldry, and as fabulous an animal as any here mentioned, as may be seen by comparing it with the figure of the real Dolphin, given with the description in a former part of this work. This fish was said to curl up his back to carry his favourites over the seas without wetting them; and to assume the most brilliant colours in dying, changing from a bright blue to as bright a yellow, and then to red and green, &c. &c.



THE UNICORN.

THIS is another offspring of the lively and fruitful fancy of man. It is represented as a compound of the horse and stag, the head and body belong to the former, and the hoofs to the latter, while the horn, the tufts, and the

tail are anomalies. This animal holds a high rank in heraldry, and is one of the supporters of the royal arms of England.

The Unicorn is often mentioned in the Scriptures, and by many commentators is supposed to be the rhinoceros. From the book of Job we learn that it was not only an animal of considerable strength, but also of a very fierce and intractable disposition—"Will the Unicorn be willing to serve thee, or abide by thy crib? Canst thou bind the Unicorn with his band in the furrow? or will he harrow the valleys for thee? Will thou trust him, because his strength is great? or wilt thou leave thy labour to him? Wilt thou believe him, that he will bring home thy seed, and gather it into thy barn?" Ch. xxxix. ver. 9—11. In the book of Psalms, xcii. ver. 10. "My horn shalt thou exalt like the horn of a Unicorn."

THE PEGASUS.

ANOTHER liberty has been taken with the horse. Mythology has added wings to its elegant figure, and called it *Pegasus*. This animal, it is said, sprang from the blood of Medusa, when Perseus had cut off her head; and immediately afterwards flew upwards towards heaven, but stopped short, and alighted on Mount Helicon, where he struck the ground with his foot, and instantly the fountain Hippocrene burst from the ground. During his residence on Mount Helicon, Pegasus became a great favourite with the Muses, who resided occasionally on that lofty mountain; and still, when any one attempts extravagant flights of poetry, he is said to have mounted on his Pegasus, as it was difficult to approach the Muses when raised so high. On the contrary, the Castalian fountain on Mount Parnassus was more accessible, and inspired poetry of a gentler nature. But to return to Pegasus; he was at length tamed by Neptune, or Minerva, and lent by the latter to Bellerophon, to enable him to conquer the horrid monster called the Chimera, which was always shifting its place, and vomiting forth flames and smoke. After the victory was achieved, Bellerophon attempted

to fly up to heaven; but Pegasus threw his rider, and flying up to heaven without him, was changed into the constellation of stars which still bears his name. Pegasus is sometimes confounded with the Hippogriff, or *Ippogrifo* of Ariosto, which is often seen in coats of arms.

THE CENTAUR.

LIKE the Sphinx, this creature is a compound of the brute and human form, exhibiting the body of a man united to that of a horse, the former rising from the chest of the latter. Absurd as such a combination must appear to the anatomist, and ill adapted as it seems for agility, it is not wholly devoid of grace, and is very frequently met with in antique sculpture. According to Grecian mythology, these beings inhabited Thessaly; and poetry has celebrated their combats with Hercules, Theseus, and Pirithous, the latter of whom was the leader of the Lapithæ, a people who vanquished the Centaurs. Their fabulous existence had its origin in that love of the marvellous, which is always found to exist in the earlier stages of society. Hence the natives of Thessaly being distinguished for their skill in horsemanship, at a time when their neighbours were unacquainted with the art of riding, they would be described as combining the powers both of the human and the equine race; in the same manner as some of the American tribes, when they first beheld the Spaniards mounted on horses, mistook them for a different race of beings from themselves, supposing them to be half men and half quadrupeds. It is by such errors that fiction, whether poetry or painting be its vehicle, creates those fanciful beings and shapes which delight the imagination.

THE SATYR.

ALTHOUGH the Satyr of the ancient poets can hardly be termed an animal, as the human form predominates, he may be introduced here as our final example of fabulous creatures. Satyrs and Fauns are represented as men

with goats' legs and horns, and were supposed to be the attendants of Bacchus, with whose worship they are generally connected. The idea of such beings was probably derived from some of the larger species of apes. They are described as inhabiting woods and forests, of which they were regarded as the protecting deities. Probably they were partly personifications, intended to express the debasing influence of animal propensities and sensual indulgence; and as nothing tends more than intoxication to reduce man to a level with the brutes, since it deprives reason of all control over the passions, the form of the Satyr may have been ingeniously intended as a visible representation of the degraded state of those who surrender up the noblest prerogative of man. Whether such was really or not the idea of those who first feigned the existence of such creatures, we may very rationally adopt this explanation, and thereby deduce an important moral lesson from what is in itself an extravagant fiction.

INDEX.

	Page		Page
A.		Asp	430
Abraxas grossulariata	501	Ass	109
Acarus domesticus, or Siro	469	— Wild	112
Acipenser sturio	382	Astacus marinus	461
Adder	427	— fluviatilis	462
Adjutant	310	Asterias	509
Admiral	452	Auchenia glama	152
Alauda arborea	215	Auk, or Northern Penguin	344
— arvensis	213	Australian Hedgehog	95
Alca impennis	344	Avicula Margaritifera	449
Alcedo ispida	255	Axis	143
Alligator lucius	444		
— or Cayman	444	B.	
Alpine Trout	408	Baboon	155
— Rat, or Marmot	79	Badger	44
Amphisbena	434	Bald Eagle	165
Anarrhichas lupus	368	Balkena mysticetus	349
Anas boschas	313	Balearic Crane	307
— Penelope	341	Bankiva Cock	288
Anchovy	398	Bantam Cock	290
Angler	379	Barbary Ape	154
Annelida	458	Barbel	417
Anobium tessellatum	478	Barn Owl	187
Anser ferus	337	Basiliak	527
Ant, Labouring	496	Bat	67
— Soldier	496	Batrachian Reptiles	436
— Eater, great	93	Bear, Black	38
— Lion	489	— Brown	38
Antelope	131	— Polar	41
Anthropoides pavonia	307	Beaver	71
— Virgo	307	Bee	492
Antilope cervicapra	131	Beluga leucas	346
— Dorcas	132	Bird of Paradise	244
— Gnu	135	Bison	123
— picta	134	Bittern	313
— rupicapra	133	Bivalves	449
Ape, Barbary	154	Black Arches Moth	505
Aphis	488	Black American Bear	38
Apis mellifica	492	Black Beetle	480
Aquila chrysaetos	161	Blackbird	191
Arachnida	467	Black Cap	201
Arctomys Marmotta	79	Black Cock	284
Ardea Herodias	311	Black Eagle	167
Argali, or Wild Sheep of	129	Black Martin	211
— Asia	129	Black Slug	456
Argonaut	457	Blatta orientalis	480
Armadillo	92	Bleak	418
Articulated animals	458	Blind Beetle	473
Arvicola amphibia	84		

	Page		Page
Bloodhound	19	Canis lupus	34
Blue Ox	134	— molossus	23
Boa Constrictor	435	— sanguinarius	19
Boar, Wild	103	— sensilis	29
Bombyx, or Psilura Monocha	503	— vulpes	32
— mori	503	Cantharis vesicatoria	478
Bookworm, small silver co- loured	505	Capra hircus	129
Boquetin	130	Caprimulgus Europæus	212
Bos Bonasus	123	Capuchin Monkeys	158
— Indicus	125	Carabus clathratus	477
— Scoticus	119	Caracal Lynx	16
— Taurus	116	Cardium fimbria	451
Botaurus Stellaris	313	Carduelis canaria	222
Bradypus tridactylus	90	Carp	413
Brahmin Bull	125	Carrion Crow	233
Brandling	402	Cassowary	302
Bream	419	Castor Fiber	71
Brown European Bear	38	Casuaris galeatus	302
Brush Turkey	271	Cat	17
Bubalus Caffer	121	Caudata pica	237
Bubo Virginianus	185	Cavallo - Marino, or Sea- Horse	381
Budytes Flava	205	Cavia Cobaya	80
Bufo vulgaris	438	Cayman	444
Buffalo	121	Centaur	533
Bull	116	Centruscus scolopax	376
— Wild	119	Cerambyx moschatus	476
Bulldog	24	Cercopithecus Diana	156
Bullfinch	229	Certhia familiaris	247
Bullhead	420	Cervus Alces	140
Butcher-bird, or Shrike	188	— axis	143
— Little	189	— canadensis	144
Butterfly fish	374	— capreolus	138
Buzzard	172	— dama	139
— Honey	174	— Elaphas	136
		— Tarantus, or rangi- fer	141
C.		Chaffinch	224
Cabbage Butterfly	500	Chamaeleo vulgaris	441
Cachalot	352	Chameleon	441
Camel	148	Chamois	133
Camel of America	152	Charadrius Morinellus	326
Camelopard	145	Charr, or Alpine Trout	407
Camelopardalis Giraffa	145	Cheese Mite	469
Camelus Bactrianus	148	Cheetah, or Hunting Leo- pard	13
— Dromedarius	150	Chelonian reptiles	446
Canary Bird	222	Chelonia mydas	447
Cancer pagurus	462	Chimpansee	153
Canis avicularis	27	Chincella lanigera	88
— aureus	36	Chinchilla	88
— domesticus molossus	24	Chouga	241
— familiaris	25	Chub	416
— familiaris avicularis	22	Ciconia alba	308
— familiaris graius	31		
— hyæna	37		

	Page		Page
Clinus aquaticus	190	Crangon vulgaris	465
Circus cyaneus	183	Crayfish	462
Civet Cat	46	Creepers	247
Clothes Moth	504	Crickets	485
Clupea alba	399	Crocodile of the Nile	443
— harengus	396	Crocodylus vulgaris	443
— pilchardus	397	Crossbill	239
— sprattus	396	Crotalus horridus	433
Coati-Mondi	45	Crowned Demoiselle	307
Coccinella septempunctata	479	Crustacea	461
Coccus cacti	486	Cuckoo	251
Cochineal Insect	486	Cuculus canorus	251
Cockatoo	264	Culex pipiens	507
Cockatrice, or Basilisk	527	Curlew	316
Cock, common	286	Currant Moth	501
— Bankiva	288	Cushat	292
— Bantam	289	Cuttle-fish	456
— Black	284	Cyclopterus lumpus	377
— Jago	289	— ocellatus	378
— Spanish	289	Cynocephalus	155
Cockchafer	472	Cygnus ferus and olor	335
Cockle	451	Cyprinus alburnus	418
Cockroach	485	— auratus	415
Cod-Fish	388	— barbus	417
Coleoptera	471	— brama	419
Coluber aspis	430	— carpio	413
— natrix	432	— cephalus	416
Columbaenas	293	— gobio	415
— livia	295	— leuciscus	416
— palumbus	292	— phoxinus	420
— Turtur	297	— rutilus	418
Condor	169	— tinca	414
Coot	329	Cypselus apus	211
Coracias garrula	255		
Coral and Corallines	512	D.	
Cormorant	332	Dace	417
— crested	333	Dasyptus sexcinctus	92
Corn Crake, or Land Rail	328	Day Fly	491
Cornish Chough	241	Death-Watch	478
Corvus corone	233	Deer, Fallow	139
— Corax	230	Delphinus Delphis	354
— frugilegus	234	Demoiselle	307
— monedula	236	Diana Monkey	156
Cottus bubalis	370	Didelphis Virginiana	70
— gobio	420	Didus ineptus	291
Coturnix vulgaris	282	Dipas Jerboa	86
Cow	118	Pipper	190
Crab	462	Diptera	506
— violet land	463	Dodo	291
— soldier, or hermit	464	Dog-fishes	361
Crambus pinguinella	505	Dogs :	
Crane	395	— Bloodhound*	19
— Balearic	307	— Bulldog	24
— Numidian	307	— Foxhound	21

	Page		Page
Dogs:		Exocetus volitans	384
— Greyhound	31		
— Mastiff	23	F.	
— Newfoundland	29	Falcon	179
— Pointer	22	— Ger	179
— Shepherd's	25	— Peregrine	179
— Spaniel	27	Falco palumbarius	175
— Water Spaniel	28	— æsalon	182
Dolphin	354	— Buteo	172
— of mythology	531	— communis	179
Domestic Rabbit	67	— Milvus	178
Dormouse, or Sleeper	85	— nisus	176
— American	86	— pavorus	174
Dor, or Blind Beetle	473	— Haliaetus	184
Dottrel	326	Fallow Deer	139
Dragon	525	Father-Lasher	370
Dragon fly	497	Felis Carracal	16
Dromaius Novæ Hollandiæ	304	— Catus, var. domestica	17
Dromedary	150	— concolor	15
Duck	339	— jubata	19
Duck-billed Platypus, or		— Leo	1
Water-mole	94	— Leopardus	10
		— Lynx	16
E.		— Onca	14
Eagle, Black	167	— Pardus	11
— Golden	161	— Tigris	8
— Sea	164	Ferret	51
— Whiteheaded or Bald	165	Fiber Zibethicus	85
Earth Worm	458	Fieldfare	194
Earwig	479	Fishing Hawk	184
Echeneis remora	367	Flea	508
Echidua setosa	95	Flounder	401
Echinus	510	Flusher	189
Eel	423	Flying-fish of the Ocean	384
Electrical Eel	421	Flying Scorpion	373
Electric Ray	364	Forficula auricularia	479
Elephant	96	Formica rufa	496
Elephant Beetle	475	Fox	32
Elephas Indicus	96	Foxhound	21
Elk	140	Fresh-water Green Polypes	509
Emberiza citrinella	217	Fringilla cœlebs	224
— hortulana	217	— carduelis	227
Emeu, or Australian Cas-		— Linota	221
sowary	304	Frog	436
Engraulis encrasicolus	398	Fulgora laternaria	486
Ephemera	491	Fulica atra	329
Equus Asinus	109		
— Caballus	106	G.	
— Hermionus	112	Gadus æglefinus	389
— Zebra	114	— Merlangus	391
Erinaceus Europæus	61	— merlucius	392
Esox lucius	409	— molva	391
— belone	394	— morghua	388
		Gallinula chloropus	327

	Page		Page
Gallus domesticus	286		
—— giganteus	289		
Gannet, or Solan Goose	333		
Garden Snail	454		
Gar-Fish	394		
Garrulus glandarius	240		
Gastrobanchus glutinosa	365		
Gazelle	132		
Gennet	52		
Geometra brumata	502		
Geotrupes stercorarius	473		
Giraffe, or Camelopard	145		
Globe Fish	380		
Glowworm	477		
Gnat	507		
Gnu	135		
Goat	129		
Goat Chaffer	476		
Goatsucker	212		
Godwit	318		
Golden Carp	415		
Golden Eagle	161		
Goldfinch	227		
Gold-Fish, or Golden Carp	415		
Goose	337		
Goshawk	175		
Grampus	357		
Grasshopper	481		
Grayling	408		
Great Ant-eater	93		
Great Dragon Fly	491		
Greek Tortoise	446		
Green Fly	488		
Greenland Whale	349		
Green Plover	325		
Green Turtle	447		
Green Woodpecker	257		
Greyhound	31		
Grey Parrot	260		
Grey Plover	325		
Griffin	528		
Ground Beetle	477		
Grouse, red	283		
Grus cinerea	305		
Gryllotalpa vulgaris	484		
Gryllus domesticus	485		
—— Migratorius	482		
——, or Acrydium	481		
Gryphon, or Griffin	528		
Gudgeon	415		
Guinea Fowl	269		
Guineapig	80		
Gurnard	385		
Gymnotus electricus	421		
		H.	
		Haddock	389
		Hag Fish	365
		Hake	392
		Halietus albicilla	164
		—— leucocephalus	165
		Hare	73
		Harfang, or Snowy Owl	186
		Hedgehog	61
		—— Australian	95
		Helix aspersa	454
		Hemiptera	485
		Hen Harrier	183
		Hermit Crab	464
		Heron	311
		Herpestes Pharaonis	47
		Herring	395
		Hippocampus brevisrostris	381
		Hippopotamus amphibius	99
		Hippopotamus, or River Horse	99
		Hirudo Medicinalis	459
		Hirundo rustica	206
		—— urbica	209
		Hog, domestic	101
		Honey-Buzzard	174
		Hoopoe	249
		Horned Owl	185
		—— Silure	369
		—— Viper	429
		Horse	106
		House-fly	507
		House Martin, or Window Swallow	209
		Humming-Birds	248
		Hyæna	37
		Hydra	509
		Hydroida	519
		Hymenoptera	492
		Hystrix cristata	89
		I.	
		Ibex, or Boquetin	130
		Ichneumon Fly	495
		——, or Egyptian Man- goustè	47
		Insects	470
		J.	
		Jackal	36
		Jackdaw	236
		Jago Cock and Her	288
		Jaguar	14
		Jay	240

	Page		Page
Jerboa	86		
John Dory	386		
Jungle Fowl, or Brush Turkey	271		
		K.	
Kangaroo	68		
Kingfisher	255		
Kite	178		
Knot	324		
Kraken	530		
		L.	
Lacerta vivipara	440		
Lady Bird, or Lady Cow	479		
Lagopus Scoticus	283		
Lamprey	424		
Lamproloma noctiluca	477		
Land Rail	328		
Lanius excubitor	288		
Lantern-Fly of the West Indies	486		
Lapwing, or Peewit	321		
Leech	459		
Lemur albifrons	160		
nigrifrons	160		
Leopard	10		
Lepidoptera	497		
Leptoptilos argala	310		
Lepus cuniculus	76		
timidus	73		
Libellula grandis	491		
Limax cinereus	456		
Limosina ægocephala	318		
Limpet	453		
Ling	391		
Linnet	221		
Lion	1		
Lioness and Cubs	6		
Lizard	440		
Llama, or Camel of America	152		
Lobster	461		
Locust	482		
Long-armed, and the Fresh-water Green Polytes.	509		
Lophius piscatorius	379		
Loxia	239		
pyrrhula	226		
Lucanus cervus	474		
Lump-sucker, or Sea-owl	377		
Lycosa tarantula	461		
Lynx, caracal	16		
Lyre-Bird of Australia, or Pheasant of the woods	274		
		M.	
Macaw	262		
Machetes Pugnax	323		
Mackerel	393		
Macropus major	68		
Magpie	237		
Magpie, or Currant Moth	501		
Maid	362		
Mantis	480		
Marikina Monkey	159		
Marmot, or Alpine Rat	79		
Martin, or Window Swallow	209		
Mastiff	23		
Magapodius tumulus	271		
Meleagris Gallo Pavo	267		
Melix aspersa	455		
Melolontha vulgaris	472		
Menura superba	274		
Merlin	182		
Mermaid, or Siren	529		
Miller's Thumb	420		
Minnow	420		
Missel Thrush	192		
Mole	63		
Mole Cricket	482		
Molluscous animals	449		
Mongoos	160		
Monkey, Capuchin	158		
Diana	156		
Oustiti	159		
Marikina	159		
Proboscis	156		
Spider	158		
Monk-fish, or Angel-fish	370		
Monodon monoceros	358		
Morse	59		
Motacilla boarula	204		
Mother Cary's Chickens	343		
Moth with its Chrysalis and Caterpillar	498		
Mouse	81		
Mule	113		
Muraena Anguilla	423		
Murex haustellus	453		
Musca domestica	507		
Muscle	452		
Mus decumanus	82		
Musculus	81		
Musk Beetle, or Goat Chafer	476		
Musk Rat	85		
Mustela furo	51		
Lutra	55		
Zibellina	54		
putorius	53		

	Page		Page
Mustela vulgaris	49	Palemon serratus	465
Myoxis glis	85	Panther	11
Myrmecophaga jubata	93	Papilio, or Pieris Brassicæ	500
Myrmeleon formicaleo	489	Par	402
Mytilus edulis	452	Paradise, Bird of	244
N.		Paradisea apoda	244
Narwhal	358	Paroquet	263
Nasalis larratus	156	—— Ring	263
Natrix torquata	432	Partridge	279
Naucrates ductor	316	Parrot, Grey	260
Nautilus, Paper	457	—— red-legged	279
Neuroptera	489	Parus major	216
Newfoundland Dog	29	Patella	453
Nightingale	198	Pavo cristatus	265
Northern Penguin	344	Peacock	265
Numenius arquata	316	Pearl Oyster	449
Numida Meleagris	269	Peewit	321
Numidian Crane	307	Pegasus	532
Nathatch, or Nutjobber	246	Pelecanus onocratulus	330
Nutjobber	246	Pelican	330
Nyl Ghau, or Blue Ox	134	Penguin	344
O.		Perca fluviatilis	411
Ocellated Sucker	378	Ferch	411
Ortodymetia grex	328	Perdix cinerea	279
Opossum Virginian	70	—— rufus	279
Ornithorhynchus paradoxus	94	Petromyzon marinus	424
Orthagoriscus	375	Phalacrocorax carbo	332
Orthoptera	479	—— cristatus	333
Ortolan	21	Phalæna, or Abraxas Gros- sulariata	501
Ortygometra Crex	328	Phasianus colchicus	277
Ortyx borealis	203	Phœnix	519
Osmerus eperlanus	392	Pheasant	277
Osprey, or Fishing Hawk	184	Philomela lusciniæ	198
Ostracion quadricornis	380	Phoca vitulina	57
Ostrea edulis	450	Phœcæna vulgaris	356
Ostrich	299	Phœnix	529
Otter	55	Phyllium siccifolium	480
Ounce	12	Physeter macrocephalus	352
Ouistiti Monkey	159	Picus viridis	257
Ourang Outang, or Chim- pansee	153	Pigeon, tame	295
Ouzel, Ring	195	—— wild	293
Ouzel, Water	190	—— wood	292
Ovis Aries	126	Pike	409
Owl, Horned	185	Pilchard	397
—— White, Barn, or Screech	187	Pilot Fish	366
Oyster, Pearl	449	Pimpla persuasoria	495
—— common	450	Pintado	269
P.		Plaice	400
Pagurus miles	464	Plantlouse, or Green Fly	488
Palæornis torquatus	263	Platalea leucorodia	315
Palæornis Alexandri	263	Platypus, duck-billed	94
		Pleuronectes flesus	401
		—— fluviatilis	379

	Page		Page
Pleuronectes platessa	400	Remora, or Sucking-fish	367
solea	401	Rhamphast tucanus	259
Plover, green	325	Rhinoceros	100
grey	325	Indicus	100
Plyctolophus galeritus	264	Rhombus maximus	399
Polypi	519	Ring Dove, or Cushat	292
sea	521	Ring Ouzel	195
Proboscis Monkey	166	Ring Paroquet	263
Pointer	22	River Horse	99
Polar, or White Bear	41	Roach	418
Polecat	53	Robin, or Redbreast	195
Pope, or Ruffe	412	Roebuck	138
Porcupine	89	Roller	255
Porpoise	356	Rook	234
Prawn	465	Rubecula familiaris	195
Pristis antiquorum	371	Ruff and Reeve	323
Procyon, or Ursus lotor	42	Ruffe	412
Psittacus erythacus	260	S.	
, or Macrocerus ara-		Sable	54
canga	262	Salmon	403
Ptarmigan	285	Pink	402
Ptilonorhynchus	242	Salmon Trout	405
Holoseri-		Salmo alpinus	408
ceus	242	fario	406
Pulex irritans	508	salar	403
Puma	15	salvelinus	407
Pyrgita domestica	220	thymallus	407
Pyrhocorax graculus	241	trutta	405
Q.		Satin Bower Bird	242
Quagga	115	Satyr	533
Quail	282	Saurian Reptiles	440
American	283	Saw Fish	371
Querquedula crecca	342	Saxicola Ananthe	218
R.		Scarabeus, or Dynastes cle-	
Rabbit	76	phantis	475
domestic	77	Scirus vulgaris	78
Raccoon	42	Scelopax gallinago	318
Radiata	509	or Totanus calidris	317
Raia batis	362	rusticola	320
Ram	128	Scomber Scomber	393
Wallachian	128	Screech Owl	187
Ramphastos tucanus	259	Seal	57
Rana temporaria	436	Sea-Bat	368
Rat	82	Sea-Horse	381
Water	84	Sea-Owl	377
Musk	85	Sea-Unicorn, or Narvhal	358
Rattle Snake	433	Sea-Urchin	510
Raven	230	Sea-Wolf, or Sea-Bat	368
Redbreast	195	Sepia octopodia	456
Red Gröuse	283	Sepia, or Cuttle-fish	456
Redshank	317	Serpents	425
Redwing	193	Shark	359
Reeve	323	Sheep	126
Rein Deer	141	Shepherd's Dog	25

	Page		Page
Shrew	65	Sucking-fish	367
Shrike	188	Suctorina	508
Shrimp	465	Sula bassana	333
Silkworm	503	Sun-fish	375
Simia troglodytes	153	Surnia nyctea	186
Siren	529	Sus scrofa	101
Sitta Europæa	246	Swallow	206
Skate, or Maid	362	Swan	335
Skegger	402	Swift	211
Skylark	213	Sword Fish	372
Sloth	90	Sylvia atricapilla	201
Slug, small grey	456	trochilus	203
black	456		
Smelt, or Sparling	392	T.	
Snail, Garden	454	Talpa Europea	63
Snake	432	Tame Pigeon	295
Snipe	318	Tapir	105
Snipe-shell	453	Tapir Americanus	105
Snowy Owl	186	Tarantula	468
Solan Goose	333	Teal	342
Soldier Crab, or Hermit Crab	464	Tench	414
Sole	401	Testudo Græca	446
Song Thrush	192	Tetrao Tetrix	284
Sorex araneus	65	Tetraodon Hispidus	380
Spaniel	27	Thalassidroma pelagica	343
Spanish Cock and Hen	288	Throstle	192
Spanish Fly	478	Thrush, or Throstle, Song.	192
Sparling	372	missel	192
Sparrow	220	Tiger	8
Sparrowhawk	176	Tinea vestianella	504
Spermaceti Whale, or Ca- chalot	352	Titmouse, or Tom Tit	216
Sphinx	523	Toad	438
Spider Monkey	158	Toad Fish	375
Spiders	467	Tom Tit	216
Sponge	518	Torpedo, or Electric Ray	364
Spoonbill	315	Tortoise, common, or Greek	446
Sprat	396	Tortoiseshell Butterfly	499
Squalus carcharias	359	Toucan	259
Squatina angelus	370	Trichechus Rosmarus	59
Squirrel	478	Trigla cuculus	385
Stag	136	Tringa Canutus	324
Stag Beetle	474	squatarola	325
Stare, or Starling	229	Trochilus colubris	248
Starfish	509	Troglodytes vulgaris	202
Starling	229	Trout	406
Stockdove, or Wild Pigeon	293	Trumpet Fish	376
Stork	308	Trunk-fish	380
Stormy Petrel, or Mother Cary's Chickens	343	Turbot	399
Strix flammea	187	Turdus iliacus	193
Struthio camelus	299	Merula	191
Sturgeon	382	parus	194
Sturnus vulgaris	229	torquatus	195
		viscivorus	192
		Turkey	207

	Page		Page
Dogs:		Exocetus volitans	384
— Greyhound	31		
— Mastiff	23	F.	
— Newfoundland	29	Falcon	179
— Pointer	22	— Ger	179
— Shepherd's	25	— Peregrine	179
— Spaniel	27	Falco palumbarius	175
— Water Spaniel	28	— asalon	182
Dolphin	354	— Buteo	172
— of mythology	531	— communis	179
Domestic Rabbit	67	— Milvus	178
Dormouse, or Sleeper	85	— nisus	176
— American	86	— pavorus	174
Dor, or Blind Beetle	473	— Haliaetus	184
Dottrel	326	Fallow Deer	139
Dragon	525	Father-Lasher	370
Dragon fly	491	Felis Carracal	16
Dromaius Novæ Hollandiæ	304	— Catus, var. domestica	17
Dromedary	150	— concolor	15
Duck	339	— jubata	19
Duck-billed Platypus, or		— Leo	1
Water-mole	94	— Leopardus	10
		— Lynx	16
E.		— Onca	14
Eagle, Black	167	— Pardus	11
— Golden	161	— Tigris	8
— Sea	164	Ferret	51
— Whiteheaded or Bald	165	Fiber Zibethecus	85
Earth Worm	458	Fieldfare	194
Earwig	479	Fishing Hawk	184
Echeneis remora	367	Flea	508
Echidua setosa	95	Flounder	401
Echinus	510	Flasher	189
Eel	423	Flying-fish of the Ocean	384
Electrical Eel	421	Flying Scorpion	373
Electric Ray	364	Focficula auricularia	479
Elephant	96	Formica rufa	496
Elephant Beetle	475	Fox	32
Elephas Indicus	96	Foxhound	21
Elk	140	Fresh-water Green Polyptes	509
Emberiza citrinella	217	Frangilla œlebs	224
— hortulana	217	— carduelis	227
Emeu, or Australian Cas-		— Linota	221
sowary	304	Frog	436
Engraulis encrasicolus	398	Fulgora laternaria	486
Ephemera	491	Fulica atra	329
Equus Asinus	109		
— caballus	106	G.	
— Hermionus	112	Gadus aeglefinus	389
— Zebra	114	— Merlangus	391
Erinaceus Europæus	61	— merlucius	392
Esox lucius	409	— molva	391
— belone	394	— morghua	388
		Gallinula chloropus	327



DOUGLAS BARNES

THE
NATURE

PHILIP

NATURE

PHILIP

NATURE

PHILIP

NATURE

PHILIP

NATURE