



JOURNAL
OF THE
ASIATIC SOCIETY OF BENGAL.

VOL. LX.

PART II. (NATURAL HISTORY, &c.)

(Nos. I to IV.—1891.)



EDITED BY

W. J. SCLATER, ESQ., M. A., F. Z. S.

AND

SURGEON-CAPTAIN J. H. TULL WALSH, J. M. S.

NATURAL HISTORY SECRETARIES.

It will flourish, if naturalists, chemists, antiquaries, philologists, and
in different parts of Asia, will commit their observations to writing, and send
the Asiatic Society at Calcutta. It will languish, if such communications shall be
intermitted; and it will die away, if they shall entirely cease." SIR WM. JONES.

CALCUTTA:

PRINTED AT THE BAPTIST MISSION PRESS,
AND PUBLISHED BY THE
ASIATIC SOCIETY, 57 PARK STREET.

1892,

868 92



Acc no - 985
9/01/07

523
H. I.

LIST OF CONTRIBUTORS.

	Page
BARCLAY, A., M. B.;— <i>Additional Uredinæ from the neighbourhood of Simla</i> ,.....	211
BIGOT, J. M. F.;— <i>Catalogue of the Diptera of the Oriental region, Part I</i> ,	250
DOHERTY, WILLIAM;— <i>A List of Butterflies of Engano with some remarks on the Danaidæ</i> ,.....	4
—————;— <i>New and rare Indian Lycœnidæ</i> ,	32
—————;— <i>Butterflies of Sumba and Sambawa with some account of the Island of Sumba</i>	141
EATON, A. E., M. A., F. E. S.;— <i>Notes on some native Ephemeroïdæ in the Indian Museum, Calcutta</i> ,	406
KING, GEORGE, M. B., LL. D., F. R. S., C. I. E.;— <i>Materials for a Flora of the Malayan Peninsula, No. III</i> ,	38
PRAIN, D.;— <i>On an undescribed oriental species of Nepeta</i> ,	204
—————;— <i>Noviciæ Indiæ IV. Two additional species of Glyptopetalum</i>	206
—————;— <i>Natural History Notes from H. M.'s I. M. Survey Steamer "Investigator," Commander R. F. HOSKYN, R. N., Commanding—No. 25. The Vegetation of the Coco Group</i> ,....	283
SCLATER, W. L., M. A.;— <i>Notes on the collection of Snakes in the Indian Museum with descriptions of several new species</i> ,	230
WALSH, J. H. TULL;— <i>On certain Spiders which mimic Ants</i> ,.....	1
—————;— <i>Natural History Notes from H. M.'s Indian Marine Survey Steamer "Investigator" Cammander R. F. HOSKYN, R. N., Commanding. No. 24. List of Deep-sea Holothurians collected during seasons 1887 to 1891, with descriptions of new species</i> ,	197



Dates of Issue. Part II, 1891.

- No. I.—Containing pp. 1—140, with Plate I, was issued on May 19th, 1891.
- No. II.—Containing pp. 141—210, with Plates II and III, was issued on July 16th, 1891.
- No. III.—Containing pp. 211—282, with Pls. IV, V and VI, was issued on September 28th, 1891.
- No. IV.—Containing pp. 283—414, was issued on March 12th, 1892.

LIST OF PLATES.

- I New oriental Butterflies (Doherty).
II Butterflies of Sumba and Sumbawa (Doherty).
III *Nepeta Bellevii* (Praia).
IV } Uredineæ from Simla (Barclay).
V }
VI New Indian Snakes (Selater).
-

JOURNAL
OF THE
ASIATIC SOCIETY OF BENGAL.



Part II.—NATURAL SCIENCE.

No. I.—1891.

I.—On certain Spiders which mimic Ants.—By SURGEON J. H. TULL
WALSH, I. M. S.

[Received 25th February; read 4th March.]

Whoever has studied the structure and habits of the various genera of the *Formicidae* must have been struck by the "fitness" which these little creatures possess for "the struggle for existence." Even in those individuals with a rudimentary and useless sting there are still the powerful biting mandibles, the acid poison which can be ejected, often to a considerable distance, on to the enemy and various protective odours, such as those secreted by the anal glands of most of the *Dolichoderidae*. The pangolin and other ant-eaters certainly cause havoc among the ants who make their dwellings in the ground, but smaller animals and birds cannot attack ants with impunity, and it is noticeable that the ants most frequently mimicked live and feed on trees. Such being the case the ant is well protected, and any other creature that, by an accident of natural selection, resembled an ant in form and colour would have obtained an advantage through this resemblance, all unconscious as regards the individual but conscious, if one may use such an expression, in relation to the orderly complexity of nature. The advantageous resemblance would, according to generally accepted laws, be transmitted and strengthened until the mimic reaped the full benefit accruing from its likeness to the ants among which it lived. Such instances of mimicry are seen among a certain sub-family of spiders,

the *Attideæ*. These spiders, ant-like in form, and partially ant-like in habit, do not spin webs for the purpose of catching prey, but, wandering about in company with the ants they resemble, spring upon their victims from behind, (hence called by some Entomologists *Saltigradæ*). Their home is generally fixed to the under surface of a leaf and consists of a small oval, whitish, silky nest just big enough to accommodate the spider. Attention has been drawn to the presence of these spiders in America* and Africa†; Mr. Wood-Mason collected two or three specimens in Assam some years ago and Mr. Rothney‡ notes the occurrence of a *Salticus* in company with *Sima rufo-nigra* in the neighbourhood of Barrackpur. I have found these spider mimics in Orissa, and also in and near Calcutta, and have, during the last eighteen months, collected or acquired some eight or ten species or varieties belonging to genera of the sub-family *Attideæ*. With one or two exceptions all these spiders were found hunting with the ants they so closely resemble. The two most common are a variety of *Salticus formicarius* Linn. which mimics *Sima rufo-nigra* Jerd. and a pretty *Salticus* (sp. ?) which may be found in company with *Ocophylla smaragdina* Fabr. whose nests are extremely common on the trees in the Royal Botanic Gardens at Sibpur.

The resemblance in form and colour is so great that collectors have been deceived, and indeed except with a lens it is difficult often to say which is the ant and which is the spider; but at the same time it must be remembered that the likeness is greater when both are alive and moving than when the dead spider is compared with the dead ant. While the body in most sub-families of spiders is short and rounded with a constriction only between the cephalothorax and the abdomen, the mimic has a long thin body like that of an ant. There is a partial constriction marking off the cephalic from the thoracic portion of the cephalothorax, and that part of the spider's body which joins the cephalothorax to the abdomen is drawn out into a pedicle having on its upper surface nodes mimicking closely those on a ant's pedicle. The colouring of the spider is also a more or less correct imitation of that of the ant. A superficial resemblance could hardly go farther, but there is a still more wonderful point to notice. The spider has four pair of legs and no antennæ; the ant has three pair of legs and a pair of long an-

* Bates, Trans. Linn. Soc. Vol. XXIII.

Belt "Naturalist in Nicaragua," p. 314.

Peckham "Protective Resemblances in Spiders." I have not been able to read this in the original and know of it only from references found in Poulton's "The Colours of Animals."

† J. P. Mansel Weale Nature Vol. iii. p. 508.

‡ Jour. Bomb. Nat. Hist. Soc. Vol. V, p. 44.

tennæ which are generally kept in motion as the little animal runs along. In adapting themselves to circumstances, the spiders have learned to use their first pair of legs to represent antennæ. In all the cases that I have noticed, the spider when moving holds its first pair of legs aloft to simulate antennæ, and certainly in the case of a *Salticus* (sp. ?) which mimics *Camponotus micans* Mayr these legs are kept in continual motion. On one of the bottles presented to me by Mr. J. Wood-Mason I find the following note made at the time the spiders were captured:—

“(Cachar, J. W.-M.) smaller one mimics and runs about with a little brown ant carrying its palpi like the open mandibles of the ants, and its first pair of legs off the ground and elbowed, as the ants do their antennæ.”

This note draws attention to another curious resemblance which is produced by the flattening of the terminal joint of the palpi. In the spider found by Mr. Wood-Mason and in several other species, this formation occurs and the falces, which are small, are partly hidden by the palpi. In other species and notably in *Salticus formicarius* the palpi are small and the resemblance to mandibles is produced by the large flattened first joint of the falces; thus the same end, as far as the resulting mimicry is concerned, is attained by two very different morphological variations. Belt (*l. c.*) notices the fact that in the Nicaraguan species the fore-legs are raised from the ground and J. P. Mansel Weale (*l. c.*) makes some interesting remarks which I will quote. He says:—

“The most perfect cases of mimicry I know of are two spiders (specific nature unknown to me) which have the closest resemblance to ants. They belong to the *Salticidæ* and are apparently related to *Salticus formicarius*. The one is smooth black and shining and runs rapidly on the ground and bark of trees, and resembles the ant which builds its nest in *Acacia horrida* and is used by the Kafirs for the purposes of torture. The other is larger and has its cephalothorax dull black and its abdomen covered with short yellowish hairs. It is generally found running on the stems of herbaceous plants and small bushes and closely resembles an ant found in similar situations. The fore-legs in both species are larger than the second pair are frequently held up when they closely resemble the antennæ of ants.”

As a general rule therefore most observers agree that the first pair of legs is used to simulate antennæ, but an exception must be noted. E. G. Peckham (*l. c.*) records that an American species *Synageles picata* “holds up its second pair of legs to represent antennæ.” This peculiarity of habit has apparently produced or been produced by a change in the relative length of the legs in this species. The general

formula for the *Attidæ* seems to be 4, 1, 3, 2; the fourth pair of legs being the longest and the second pair the shortest. In *Synageles picata* the formula, to judge from a figure (*l. c.*), is 4, 2, 3, 1. *Synemosyna formica*, another American spider observed by Peckham, has the usual formula, but is said to use its second pair of legs as antennæ!

I have mentioned that the spiders are probably protected from birds and other enemies by their resemblance to ants, but there can be no doubt that frequently they also thereby gain another very considerable advantage. The ants with which these spiders "most do congregate" are fairly omnivorous feeders, but shew a decided preference for sweet juices often to be found exuding from trees, fruit or flowers. To these juices come also flies, small beetles and other insects which form the natural prey of the spider, and which do not, under the circumstances, particularly fear the ants. Thus while the flies are sucking up sweetness in company with the ants, the spider is no doubt able, under cover of his disguise, to approach near enough to make a spring upon his unsuspecting victim and fix his sharp falces into its body. As regards the ants themselves, they do not seem to take any particular notice of the spiders, and do not apparently attack them. One spider, a mimic of *Ocophylla smaragdina* was found by me in a nest of these ants with its little silky shelter attached to one of the leaves which formed part of the abode of probably the most fierce of all the ants found in Bengal. It may also be supposed that the spider does not attack or annoy the ants.

II.—*A List of the Butterflies of Engano, with some Remarks on the Danaidæ.*—By WILLIAM DOHERTY, Cincinnati, U. S. A. Communicated by the NATURAL HISTORY SECRETARY.

[Received 21st February, 1891, read 4th March, 1891.]

(With Plate I, figs. 1—4.)

The long parallel lines of upheaval which characterize Burma are continued far into the Malayan region in the form of three great chains of islands and mountains. The most eastern of these, and the oldest, being chiefly composed of primitive rocks, consists of the Malay Peninsula, itself built up of several parallel ranges, the Riouw and Lingga groups, Banka and Billiton. The most western includes the Audamans and Nicobars, and the line of islands which may be called the Nias group, lying west of Sumatra, extending perhaps to western Java. Between these two the large island of Sumatra has been formed, probably in times geologically recent. No doubt some parts of Sumatra are composed of older rocks, but till the great volcanic up-

heaval occurred, its place was probably occupied by a few isolated islets, and to this day its fauna is chiefly made up of immigrants from the Malay Peninsula, and in the extreme south-west from Java, the number of endemic species being small. To understand the faunal relations of the Nias Islands, one must eliminate all idea of Sumatra from the mind. Until quite recent times, their relations were, I think, wholly with Java on the one side and the Nicobars on the other.

The Nias Islands lie like a broken breakwater along the western coast of Sumatra, in a line between five and six hundred miles in length, from Pulo Babi to South Pageh. They are separated from Sumatra by a deep channel nearly a hundred miles in breadth, but in two places more or less bridged by islands. Their united area is now only about 6000 square miles, equal to that of Yorkshire. But they seem to be the remains of a much larger mass of land. The deep sea that surrounds them swallows up all the alluvium from their streams; the tremendous surf on their western shore steadily undermines their hills, and under this process the islands have long been wearing away.

Engano lies much further south, and is wholly surrounded by deep sea, in which it might long ago have disappeared but for the coral reefs that protect most of its coast. It is only eighty miles from Sumatra, to which it has no faunal resemblance whatever. On the other hand, it is 180 miles from South Pageh in the Nias group, and 210 from Java. But on the side of Java there is only open sea, while on that of Nias there are three or four reefs and islets, and as might be expected from this, Engano may be zoologically considered as an outlying member of the Nias group, with certain Javan affinities.

An excellent description of the people and products of Engano has lately appeared in the *Tijdschrift van Nederlandsch Indië*, but so little is said of the island itself, that I may be permitted a few remarks on the subject. The area seems to be incorrectly stated; it is about a hundred and twenty square miles. The eastern coast is low and flat, bordered in places by mangrove swamps. The western, where the hills attain a height of nearly a thousand feet, descends precipitously into a narrow lagoon filled with branching corals and coral-haunting fishes, and on the reef beyond, the surf of one of the bluest and deepest of seas beats continually with such violence that the whole island seems to shake under it. Engano seems now to be sinking; it has formerly been more or less submerged. A shell of coral rock covers almost the whole of it, thick over the eastern lowlands, thinning gradually to the westward, so that the streams generally break through it there, and flow through deep gorges. In some places the crust has given way so as to form deep little round dells, with stalactite caves piercing their sides,

and a subterranean stream roaring at the bottom, hidden by dense vegetation. The people believe these places to be haunted by evil spirits, and when I wanted to be lowered into one to look for shells in the caves, they all objected most vehemently.

Beneath the coral, the rock seems everywhere a coarse, friable sandstone. If the Nias islands consist generally of this soft rock, it will go far to explain their present fragmentary state.

Until the present generation, the population consisted of a number of hostile tribes, and the older men are still proficient in the use of the spear, and delight in mimic battles and dramatic representations of the surprise of a sleeping enemy. In recent times native vessels have taken to coming to Pulo Dua, two small islands a mile or two off the western coast, for cargoes of cocoa-nuts, and it is probably the growth of this trade that broke down the tribal system. At present Javanese and Malays have now settled in several villages, though Pulo Dua still remains their headquarters, the plague of sandflies making life unendurable to them on the main island. It is to the trading vessels that the Enganese also owe the introduction of diseases that are rapidly killing them off. An outbreak of small-pox carried off all the inhabitants of the two villages near the southern end of the island, and to this day this district is called the Land of Ghosts, and no Enganese will set foot on it. No alcoholic liquors have been brought to Engano; whether this is owing to the precautions of the Dutch, or the religious scruples of the Sumatran traders, I do not know. But syphilis was introduced many years ago, and through it the race has lost all reproductive power. In some of the villages there are hardly any children, and the area of cultivated land decreases every year. The Dutch government, I believe, now contemplates leasing the island to capitalists, and in another century it seems likely that the natives will be entirely replaced by Javanese coolies working under European supervision.

The islanders are of about the same degree of civilization as the Nicobarese. Till lately they are said to have gone about almost naked, and their island is always called Pulo Telanjang or the Naked Island by the Malays. They grow no rice, nor do they make use of the pandanus and cycas as food like the Nicobarese. They make little temporary clearings, surrounded by a stout fence to keep out wild pigs. Their staple food is the taro or *kaladi*,* which they call *ayudapa*. They also have plantains, papayas and pumpkins, and eat quantities of cocoa-nuts, and drink the water in the nuts, the springs near the coast being usually brackish. They also draw toddy from the tree, though its use is not

* Our word *Calladium*, though it looks Greek enough, is derived from this, the Malay name of the plant.

habitual as in the Nicobars. They keep fowls and a few pigs, feeding both on cocoa-nuts, and are expert in spearing fish and turtle. Thanks to this abundant diet and the rarity of famines, they have lost the woodman's instincts, and are little better acquainted with the forest and its vegetable and animal products, than are the rice-growing Malays of Borneo and Sumatra, who scarcely know the names of the commonest trees.

Their origin will perhaps puzzle future investigators. Though their physiognomy is odd and characteristic, they seem to be more or less allied to the Nicobarese, but without the negrito strain which seemed to me obvious in some members of that race. The theory has been advanced that the Nicobarese are of Shan or Siamese blood, no doubt on philological grounds, as there is no personal resemblance. On the other hand, the Malays are physically almost indistinguishable from the Siamese, and may roughly be defined as a Shan people, just touched with Polynesian blood in a few localities, and speaking a Polynesian language slightly mongolized.

A list of Enganese words is given in the article I have mentioned. The enunciation is curiously different from the Malay, and is difficult to follow, the vowels appearing to be uttered in several different tones, as in Shan or Chinese. As in Nicobarese, euphony is spoilt by the excessive number of imperfect k's and ng's (the French *n nasal*). These consonants, which rarely occur in Polynesian languages, except the Malay, are generally absent in the personal and place-names, which usually have a strikingly Polynesian air. I spent much of my time at Kayapn, where Pahakela (the *p* is always pronounced halfway between a *p* and an *f*) was chief: the names would be natural in New Zealand or Hawaii.

There seem to be no indigenous mammals on the island except bats, wild pigs and a *Paradoxurus*. This absence of four-footed enemies may have been one cause of the excessive multiplication of birds. One sees more birds in a day in Engano, than in a month in Borneo or Sumatra, and coming from the latter island I was struck with their exceeding tameness. I saw four sorts of parrots and three of pigeons; the latter are never out of sight, the former keep up a deafening noise all day. The species seem generally different from the Sumatran, and no doubt some are undescribed. I think an ornithologist could do good work in Engano.

I think I got nine or ten sorts of land shells of which the greater number must be new. My small collections of moths and beetles were sent to England. The striking features of the latter were the prevalence of *Elateride* and the extreme rarity of *Phyllophaga*. No *Cassida* was taken at all.

Frogs, toads, snakes and lizards all abounded, and it is evident that the island has been at one time connected with the mainland, especially as a true cyprinoid fish appears to occur in the streams of the highlands. The coral crust does not imply the entire submergence of the island, while the number of peculiar butterflies shows its long isolation.

The forest is finest on the western coast, where the coral crust is thin or broken. The vegetation is there nearly as grand as on the mountain slopes of Sumatra. This side of the island is evidently very rainy. Even the comparatively barren eastern coast seems well suited to some plants, and in some places the pandani, which do not form thickets by themselves, as in the Nicobars, but grow among other trees, reach a surprising height, one I saw being fully a hundred feet high.

I caught fifty-eight kinds of butterflies, and saw two or three more on the day I landed. Three days of heavy rain followed, after which insects were scarce, and I obtained good sets of only a few species. The heavy forests of the interior scarcely produced anything but *Cyrestis perianther*, *Mycalis mineus*, *Amathusia amythaon*, *Epoxyglides tharis*, and *Paragerydus unicolor*. Most of the peculiar *Danaidæ* occurred only close to the shore. Examination of the inland forest at another season may produce true endemic species, such as have been found in Nias. Should any future collector visit the island, I recommend Bua-bua, near the western coast, as the best collecting ground, and April or May as the best season. My own visit was in September, 1890, and lasted three weeks.

The species are mostly local forms of widely spread species. I have felt obliged to give names to thirteen of them, including nearly all of the *Danaidæ*, and these should in most cases rather be called subspecies than species, but as I always give the name of the parent form, this ought to cause no difficulty. My types will be placed in the collection of the Hon. L. W. Rothschild. I have endeavoured to make in the text such comparisons as I could with the species taken by Herr Kheil in Nias, but I find it difficult to sum up the results.

While I have a few criticisms to make on Mr. Moore's well-known monograph of the *Danaidæ*,* I do not wish to appear ungrateful for the help he has there given to all students of this group. Though some have objected, he seems to have done right in giving names to the numerous subdivisions he has made. At the same time, most of them seem scarcely worthy of a higher rank than that of subgenera, and some are founded on minute and unreliable characters.† His classification

* In the Proceedings of the Zoological Society for 1883.

† Such as the rudimentary recurrent vein in the cell of the forewing, a feature

is dubious, and some of his species worse than dubious. But in any case he has let a flood of light into one of the darkest corners of entomology, and all future work in this family must start from his as a base.

I do not see any reason for accepting his term *Euploina*, instead of the commonly received *Danaidæ* or *Danainæ*. The name *Danaus* necessarily falls, having been used for *Pieride*. But *Danaida* and *Danais* are both older than *Limnas*, and there can be no similar objection to them. Mr. Scudder, with his usual conscientiousness, has adopted *Danaida*, the earlier of the two names. But I think his verdict may be reasonably traversed on the ground of the irregular formation of *Danaida* and the universal use of *Danais*. It can hardly be seriously maintained that the latter name cannot be used on account of its resemblance to *Danaus*. So the group obviously remains the *Danaidæ* (or *Danainæ*), the genus *Danais*, and its type *plexippus*,* *Anosia* falling before it.

Mr. Moore's primary division of the family is into two groups, the *Limnaina*, including *Danais* and *Hestia*, and the *Euploina* including

which seems to vary in different specimens. *Mahintha* was founded on a local race of *Euplaa core*. *E. simulatrix* was placed first in *Vadebra* (*Crastia*) and then in *Menama*, though it obviously cannot come into either, or into the "section" in which both are placed.

* The name now applied to a butterfly known to every American farmer as *archippus*. These useless changes of name now so much insisted on, especially by American naturalists, are bringing scientific nomenclature into well-deserved disrepute among all outsiders. Surely there ought to be a statute of limitations; security that some one, turning over musty volumes of pre-scientific times, shall not make all existing works obsolete. At present we stop short at Linnæus. This is purely an arbitrary line. The next generation will perhaps go back to Ray and Swammerdam; with the aid of a little zeal and imagination quite a number of generic names can be found in their books. The first false step taken was the acceptance of Hübner's childish work as an authority. There was *Adolias*, a genus described by such a profound and discriminating writer as Boisduval, and accepted by all naturalists. Finally, some one discovered that a few years before the date of Boisduval's great work, Hübner, a contemporary, not of Linnæus, but of Latreille, had invented a genus *Euthalia*, described merely as "dark with white and red spots," containing *lubentina* and *adonia*, and placed in an imaginary family, prettily named *die Fröhliche* or The Joyful Ones. *E. acantha*, and *E. evelina* (the latter along with an African *Aterica* and a European *Apatura*) were at the same time placed in different genera of another imaginary family called *die Munters*, or The Lively Ones. And so, to the confusion of naturalists all over the world, *Euthalia* took the place of *Adolias*. Lepidopterists have yielded to an infinity of similar changes. It remains to be seen whether coleopterists will be equally submissive. Mr. Crotch now proposes to alter the names of a number of the best-known genera of beetles, names consecrated by a century of use. I cannot help wishing his opponents all success in the struggle against *ce malencontreux droit de priorité*, as M. Deyrolle calls it.

Euploea and *Hamadryas*. The only difference mentioned, is that in the former there is usually a precostal cell in the hindwing, wanting in the latter. Now, it is true, an obscure rudiment of one is present in some species, but better distinctions can easily be found. *Hamadryas* probably does not belong to this family, but to the *Neotropidæ*.* *Hestia* has undoubted affinities with *Danais* in its neuration and markings, but in its anal tufts, its egg, and the structure of its feet, it is more like *Euploea*, while its antennæ show how remote it is from either. So I would suggest the following classification instead of Mr. Moore's.

Tarsi with large paronychia and pulvillus. Anal tufts two or four.

Antennæ filiform, HESTIA Group.

„ clavate, EUPLOEA Group.

Tarsi with rudimentary paronychia and pulvillus†. Anal tufts two.

DANAIS Group.

Mr. Butler, to whom we owe the classification of the *Euploea*s by the male marks, accepted, as all previous writers had done, Cramer's core as the type of the genus, and formed the genus *Macroploea*, with *elisa* as its type, applying Hübner's name *Crastia* to *climena* and its allies. This arrangement has been generally adopted, and it is most unfortunate that Mr. Moore should alter all three names on what may be called archæological grounds. *Core* has been accepted for a century as the

* I do not know whether Schatz's name can stand, as it is not derived from that of any genus, and has no type. The name *Erycinidæ* has been dropped by some writers for this reason, the generic name *Erycina* having been superseded.

† The appendages of the last joint of the tarsi of butterflies are, 1st, the claws, 2nd, the paronychia, 3rd, the pulvillus with its shield. The *paronychia* are tactile organs, one on each side of the foot, slender and flexible (not jointed as some writers have said), covered with short sensitive hairs or *papille*. In most cases they are split into two long finger-like process, the longer following the course of the claws, the shorter curling round the edge of the pulvillus-sheath, obviously with the object of examining the surfaces touched by these two organs. The *pulvillus* is a soft, muscular projection, resembling an additional tarsal joint, but not armour-clad like the others. It is evidently intended to act as a buffer, to break the shock of alighting, and no doubt owes its name *pulvillus*, or cushion, to this function. Its lateral surface is sensitive, with papillæ like those on the paronychia. But its lower surface is covered with a round corneous plate, or shield, which no doubt acts as a *sucker*, flexible in life, slightly concave in the middle. This is easily detached in dried specimens.

Now in certain butterflies of aerial habits like *Danais*, the claws have been greatly lengthened, so that the pulvillus did not touch the surface on alighting. Thus losing all function, it has shrunk into a small, hard, inflexible mass. The paronychia have similarly lost their use, and remain in various stages of obsolescence. They are still distinct in *Ideopsis*, and retain their bifid shape, though too small to be of use.

type of *Euplœa*, and although it would be more pedantically accurate to follow Fabricius rather than Cramer, it seems a pity to confine the use of *Euplœa* to two or three species (with their local forms) forming a small aberrant genus, now well-known as *Macroploea*, while hundreds of species cluster around *core* as their natural type and centre. *Macroploea* (Mr. Moore's *Euplœa*) is a true genus, and must be ultimately recognized as such. But unless we accept M. Moore's little groups as genera, we must find some general name for the large mass of species remaining. It is unlikely that naturalists will consent to call them all *Crastia*. Surely they had better keep the name of *Euplœa* as before, with *Crastia* (= *Vadebra*,* Moore) as a subgenus under it.

As to the position of the family as a whole, I understand that Mr. Scudder now gives it a low one, below the *Apaturidæ*, *Nymphalidæ* and *Satyridæ*. Much, however, may be said in favour of retaining it at the head of butterflies. The shoulder plates are in most butterflies shaped more or less like a human foot as seen from the side, having a long posterior process. It is only in the *Danaidæ* and the *Acræidæ* that they are simply semicircular in shape (in the *Danaidæ* more than half a circle) without the posterior process. The process is also greatly reduced in *Cynthia* and *Cethosia*, the nearest genera of the *Apaturidæ*, while in the lower forms of the *Tetrapoda*, the plate greatly resembles that of other butterflies. The long claws of the *Danaidæ*, culminating in *Danais* with its aborted paronychial, may also show a higher development than other *Tetrapoda*. But they are also found in the *Papilio*s, and seem to go along with a prolonged and aerial flight. The fore-feet of the male still retain the tarsus, which is lost in most *Neotropidæ* (*Ithomiadæ*), but it is less developed than in the *Morphidæ* and most other higher butterflies.

The striking feature of the *Danaidæ*, however, is the more perfect development of the fore-feet of the female into sensory organs much resembling the antennæ, but set with highly specialized tactile hairs or papillæ.† I have taken a good many notes on this subject, on which so far as I know nothing has yet been written, and when they are more complete, I hope to publish them. So far as examination has yet shown me, there seems to be a gradual development upwards from the *Morphidæ*, in which the joints are tapering, the last the slenderest, with a conspicuous pulvillus and rudimentary claws with their appertaining hairs. In the higher groups, these tarsi become less and less like a foot, and

* Mr. Moore applied this name to two different genera of butterflies in the same issue of the Proceedings of the Zoological Society.

† These also occur in all *Tetrapoda*, and in some other groups, as the *Nemobiadæ*, but are there few and small.

more and more like a palpus. The *Satyridæ* and *Nymphalidæ* are more developed in this respect than the *Morphidæ*, and the *Apaturidæ* much more. The *Neotropidæ* according to Mr. Godman have the fore-tarsi of the female not clavate, and hence probably less developed. In the *Danaidæ* and the nearest *Apaturidæ* (*Cethosia* and *Cynthia*), they attain their largest development. This can hardly be an organ of touch, because these feet are poorly provided with muscles, and are capable of but little movement. It may be an organ of hearing, but it is more likely one of smell, correlated with the scent-producing glands of the male. It is significant that in the *Neotropidæ*, where this foot is more normal, the male has no anal tufts.

I think the special development of this organ, which is found in all Tetrapod butterflies, though but little specialized in the lowest, is an argument for the high rank of the *Danaidæ*. But I am aware that the feet of butterflies often vary most irregularly, independently of the general organism. The aborted forefoot of the male shows some such inconsistencies, rudiments being, as Darwin has shown, eminently variable. *Melanitis* and *Bletogona* in the *Satyridæ*, many *Pieridæ*, and at least one genus of the *Lycanidæ* have the claws bifid. The number of joints in the forefeet of female Tetrapods also varies. And in *Pseudergolis*, a genus of the *Apaturidæ*, differing but little from its neighbours in other points, the fore-tarsi of the female, though small, are quite perfect, with claws, pulvillus and bifid paronychia. This fact would place it quite outside of the Tetrapod butterflies, near the *Nemeobiadæ*, if the feet offered really reliable characters.

The prehensors of the *Danaidæ* are chiefly remarkable for the aborted uncus and the broad flat clasps. They do not vary much in the different species; most in *Hestia* and some kinds of *Danais*. My notes are insufficient to permit me to generalize on the subject. It is remarkable that the only species in which they are really very aberrant is *Danais aglaioides*, where the clasps are prolonged downwards, and exposed to view even in dried specimens. This butterfly is closely allied to *D. aglaia*, differing chiefly in the shape of the wings, and they are found together over much of their range. It seems likely that in this case the prehensors have been strongly differentiated to prevent hybridism, and keep the species separate.*

* This may also be the case with the genus *Lampides*, where a number of protected species, remarkably alike in colour and markings, have the prehensors extremely unlike, and so differently armed with hooks and pincers, that the union of different species would presumably bring about serious consequences. Great confusion has been caused here by the German lepidopterists, who have recently figured a great many species with very brief descriptions, and no account of the prehensors. The

The egg also varies little in the *Danaidæ*, and affords few characters for classification. It is slightly like that of *Libythea* (which more resembles that of the *Pieridæ*), and is nearly identical with that of the *Acroïdæ*, and with those of *Oynthia* and *Cethosia*, the first genera of the *Apaturidæ*, which are again connected by several links with *Argynnis*, where the series seems to end. In the *Danaidæ* it is large, soft, cartridge-shaped, more than half again as high as wide, yellowish, or sometimes pearl-white. It has a variable number of upright ribs anastomosing near the apex (where the reticulations are more or less hexagonal over a varying extent of surface), and with horizontal cross-lines varying greatly in number and in distinctness, and generally hard to count. I append a list of species with the number of vertical ribs on the egg of each, and with that of the cross-lines in parenthesis, whenever I know it. The numbers always vary slightly in the same species, and that of the cross-lines can only be given approximately. The egg of *Nectaria* and *Macroploea* I forgot to examine. That of *Hestia* is unlike the others in having the reticulations more or less hexagonal, the ribs being slightly zigzag. The figures support the idea that *Hestia* is nearer *Euploea* than *Danais*. It will be observed that in *Danais* (including *Ideopsis* and *Radena*, which are practically identical) the cross-lines are usually more numerous than the ribs, while in *Hestia* and *Euploea* they are fewer, *Stictoploea* coming near *Hestia* in this respect, as well as in the anal tufts. The figures for *Danais chrysippus* and perhaps *limniaca* and *tytia* are doubtful.

HESTIA Group.	<i>Hestia cadellii</i> ,	21 (14).
	<i>hadenii</i> ,	23 (14-15).
	<i>lynceus</i> ,	21 (14).
EUPLOEA Group.	<i>Stictoploea lacordairei</i> ,	27 (13).
	<i>Salpinx novaræ</i> ,	26.
	<i>kollarii</i> ,	22-23 (20).
	<i>splendens</i> ,	22 (18-19).
	<i>Euploea esperii</i> ,	26.
	<i>camorta</i> ,	31.
	<i>simulatrix</i> ,	30.
	<i>core</i> ,	34 (23-25).
	<i>midamus</i> ,	24-25 (20).
	<i>alcathoë</i> ,	39 (26-27).
	<i>rhadamanthus</i> ,	21-23 (15).
	<i>pinwillii</i> ,	26 (18).
DANAIS Group.	<i>Danais genutia</i> ,	32 (30).

revision of this genus, which is further complicated by the prevalence of local and seasonal forms, will prove a heavy task to future naturalists.

<i>Danais hegesippus</i> ,	31-34 (29-30).
<i>nesippus</i> ,	34.
<i>chrysippus</i> ,	34 ("nearly 50" ?).
<i>gautamoides</i> ,	23-24.
<i>limniace</i> ,	25 (25).
<i>septentrionis</i> ,	19-21 (30).
<i>melaneus</i> ,	16-18 (27).
<i>larissa</i> ,	20-21 (30).
<i>tytia</i> ,	20-21 (38).
<i>melanoleuca</i> ,	19 (29).
<i>aglaia</i> ,	17 (26-27).
<i>aglaioides</i> ,	15-16 (25).
<i>Ideopsis daos</i> ,	15 (25).
<i>Radena nicobarica</i> ,	16.
<i>vulgaris</i> ,	14-16 (25).

Next comes the vexed question of scent-glands. I was so discouraged by the loss of all my notes on this and other structural matters in 1887, that I have taken but few since, so I cannot now speak with much authority on the subject.

It seems probable that there is a disagreeable odour in all *Danaida*, apart from that of the anal tufts and alar glands. Though infinitely less in strength than it is in the *Agaristidae* and other moths, it is still quite perceptible on pinching some species, such as *Danais genutia*, *crocea* and *vulgaris* (which smells of sorrel). In others I cannot make it out at all. This smell is probably associated with a taste highly disagreeable to spiders, chickens, etc. The results of my experiments made some years ago in the Celebes, Java, etc., are now lost, but in the Malay Peninsula I lately made a few on spiders, with the following results. Except in the case of the first two species, they cannot be entirely trusted.

Species always rejected, *Danais genutia*,
Radena vulgaris,
Ornithoptera ruficollis,
 (only two offered).

Species sometimes rejected, *Euploea nidamus* ;
 „ *rhadamanthus*,
Neptis varmona,
Cethosia hypsina,
Loxura atymnus,
Lampides ælianus.

Species never rejected, *Neptis* sp.
Athyma sp. etc.

It is to be observed that the *Radenas* are perhaps the most perfectly

protected of the *Danaidæ*, and yet they are not brightly coloured, they have no alar scent-glands, and the anal tufts are the smallest in the whole family, and give the least fragrance.

Apart from those producing the odour pervading the whole body, four different classes of scent-glands have been pointed out.

The first are the impressed silky streaks of altered scales on the forewings of some *Euploas*, often called *brands*. Mr. Distant oddly enough calls these pseudo-scent-glands, taking the falsity of their claims for granted, while assuming to leave the question open. The marks in question seem to consist of scales only, and I have not yet observed any glands connected with them. Though not very conspicuous in some cases, they may be addressed to the eye of the female, and help her to recognize the male of her species. For odours vary but little in the *Euploas*, and the amount of mimicry is so great that in most localities there are several species similarly coloured but bearing different sex marks. In any case these brands can hardly be protective, and the assumption by some writers that a species furnished with them is better protected, and more likely to be a mimicked than a mimicking form, is without foundation.

The variously-coloured velvety patches on the hindwing of *Trepsichrois*, *Salpinx* etc., are no doubt true scent-organs. In the case of *Trepsichrois midamus* the odour is sometimes quite perceptible in the detached hindwing, while in *Salpinx* and *Calliphea* it is apparently excited by friction against the forewing. In some cases, such as *Euploea oceanis*, described below, the velvety area is on the underside of the forewing.

The "pouches" on the hind-wings of certain species of *Danais*, such as *D. limniace*, are probably true scent-organs, though neither Professor Wood-Mason, who has given them special attention, nor myself, have been able to detect any odour. He has kindly shown me a microscopic section of the pouch. The cavity is lined with a semi-circle of long cylindrical cells radiating from it, their nuclei near their inner ends, the outer wall of the mass thick and chitinous, the inner membranous, with the attached scales aborted. According to Professor Wood-Mason, the odour-giving fluid is presumably manufactured in these cells, and not merely drawn from the body and stored in them. An account of these pouches, apparently made from the study of dried specimens, has appeared in the work on the "Duftapparate indo-australischer Schmetterlinge," by Dr. Erich Haase of Dresden.

Finally there are the abdominal tufts which we find in this family alone of all butterflies. They resemble those of the *Callidulidæ*, but are more specialized, though proportionately smaller than in these and

many other moths.* They exist in a very rudimentary state in the female, and can as in the male be forced to the surface by the action of the fluids in the body. From some such small beginning, they have grown to their greatest development, several stages of the process remaining as fixed characters in the different genera. The smallest tufts occur in *Kadena vulgaris*; they are much longer in *R. juvena*. The greatest development of simple tufts occurs perhaps in *Euploea* (*Trepischrois*) *midamus*. In all the species of the *Danais* group the tufts are simple and single, only in *Tirumala* the hairs are curled at the tips. They are also simple in the majority of the old genus *Euploea*, and I limit the name to these. The gland in all these consists, when protruded, of a finger-like projection extending laterally on each side of the abdomen, and bent round forwards (*i. e.*, towards the head) in a semicircle, bearing long odoriferous hairs to the very tip.

But in the genus *Salpinx*, as limited by Mr. Butler (including Mr. Moore's *Salpinx*, *Isamia*, *Pademna*, *Satanga* and *Selinda*, but not apparently *Danisepa*), and also in his genus *Calliptea*,† while the gland is similar (somewhat longer and more bent), the long hairs are gathered chiefly around its base, the outer part being naked, except at the extreme tip, where there is a brush of short, stiff bristles, sometimes coloured differently from the long basal hairs. This terminal brush is only seen when the gland is fully protruded. The development of this form from the first is obvious. This group is also large, but the number of species seems to me to have been greatly exaggerated.

In more advanced forms, a new gland, generally somewhat shorter than the other, has been pushed out from its hairy base on the anterior side. In this case both glands are covered with hair to the tip, the space between their bases being naked. This occurs in the genus *Hestia*, and apparently also in *Macroploea* and *Stictoploea*‡ as defined by Mr. Butler.

Finally in the single genus *Nectaria*, while the posterior tuft remains unchanged, the anterior or last developed one is again absorbed, and is represented by a microscopic tuft at the base of the other. This can hardly be the incipient stage, as it is too small to be of any use, and the nature of its origin is obviously as given above. It hence appears

* In most moths these tufts are attached to the prehensors or clasping-organs of the male.

† Herr Georg Semper and Dr. Haase consider *Calliptea* identical with *Euploea* (*i. e.*, *Macroploea*), but I cannot agree with them. *Calliptea* scarcely differs from *Salpinx*.

‡ My drawings of the glands of *Macroploea* and *Stictoploea* were lost in 1887, and I have not examined those organs since. But my impression is that they are very similar to those of *Hestia*, though somewhat less developed.

that *Nectaria* has attained the furthest development in this regard, and may still with good reason be retained at the head of the *Danaïdo*, and of all butterflies.

The following scheme will serve to show the relations of the genera of this family. Mr. Moore's genera may be reduced to subgenera, though *Parantica* and perhaps one or two others may be retained.

- A. Feet with well-developed paronychia and pulvillus.
 - B. Forewing with a pre-discoidal cell.
 - C. Anal glands four, two aborted,..... NECTARIA.
 - C. Anal glands four, none aborted, HESTIA.
 - B. Forewing with no pre-discoidal cell.
 - C. Glands four (?), none aborted.
 - D. Forewing of male with no brand, ... MACROPLÆA.
 - D. Forewing of male with two brands, ... STICTOPLÆA.
 - C. Glands two, tufts four, SALPINX.*
 - C. Glands two, tufts two, EUPLÆA.
- A. Feet with rudimentary paronychia and pulvillus.
 - B. Hindwing triangular, with scent-pouches, DANAIIS.
 - B. Hindwing triangular, without scent-pouches, ... RADENA.
 - B. Hindwing rounded, generally without scent-pouches, IDEOPSIS.

There seem to be grounds for believing that the scent-tufts in the male serve three distinct purposes, 1st, to warn off enemies, 2nd, to attract the female by colour, 3rd, to attract her by odour.†

1st. The odours given forth are in some cases unpleasant. In *Danaïis limniace*, it is that of turmeric, and is downright disagreeable. Other species of *Danaïis* have the peculiar aromatic fragrance characteristic of so many protected Mediterranean plants. In the genus *Salpinx*, in which the tufts reach their greatest development in size, the odour, though exceedingly sweet, is almost overpowering. Now it may at first sight seem absurd to say that the scent of honey may protect an insect from its enemies. But we know this to be sometimes the case. The male of the singular Hesperian, *Calliana pieridoides* has assumed conspicuous white colours along with a delicious odour, the female remaining dark and odourless.‡ The fine Malayan Morphid,

* Including *Calliptera*.

† The odour is not always present, but comes and goes, whether by the exhaustion of the supply, or by the will of the insect, it would be interesting to learn. I have sometimes examined the tufts of several males one after the other, without detecting the perfume.

‡ Mr. de Nicéville recently discovered the female, and it turned out to be dark, as I had predicted.

that *Nectarina* has attained the furthest development in this regard, and may still with good reason be retained at the head of the *Danaida*, and of all butterflies.

The following scheme will serve to show the relations of the genera of this family. Mr. Moore's genera may be reduced to subgenera, though *Parantica* and perhaps one or two others may be retained.

- A. Feet with well-developed paronychia and pulvillus.
 - B. Forewing with a pre-discoidal cell.
 - C. Anal glands four, two aborted, NECTARIA.
 - C. Anal glands four, none aborted, HESTIA.
 - B. Forewing with no pre-discoidal cell.
 - C. Glands four (?), none aborted.
 - D. Forewing of male with no brand, ... MACROPLEA.
 - D. Forewing of male with two brands, ... STICTOPLEA.
 - C. Glands two, tufts four, SALPINX.*
 - C. Glands two, tufts two, EUPLAEA.
- A. Feet with rudimentary paronychia and pulvillus.
 - B. Hindwing triangular, with scent-pouches, DANAIS.
 - B. Hindwing triangular, without scent-pouches, ... RADENA.
 - B. Hindwing rounded, generally without scent-pouches, IDEOPSIS.

There seem to be grounds for believing that the scent-tufts in the male serve three distinct purposes, 1st, to warn off enemies, 2nd, to attract the female by colour, 3rd, to attract her by odour.†

1st. The odours given forth are in some cases unpleasant. In *Danais limnæa*, it is that of turmeric, and is downright disagreeable. Other species of *Danais* have the peculiar aromatic fragrance characteristic of so many protected Mediterranean plants. In the genus *Salpinx*, in which the tufts reach their greatest development in size, the odour, though exceedingly sweet, is almost overpowering. Now it may at first sight seem absurd to say that the scent of honey may protect an insect from its enemies. But we know this to be sometimes the case. The male of the singular Hesperian, *Calliana pieridoides* has assumed conspicuous white colours along with a delicious odour, the female remaining dark and odourless.‡ The fine Malayan Morphid,

* Including *Calliploea*.

† The odour is not always present, but comes and goes, whether by the exhaustion of the supply, or by the will of the insect, it would be interesting to learn. I have sometimes examined the tufts of several males one after the other, without detecting the perfume.

‡ Mr. de Nicéville recently discovered the female, and it turned out to be dark, as I had predicted.

Melanocyna faunula, is saturated with sweet perfumes, and drifts feebly about the country, fearless of enemies. The fragrant butterflies of the genera *Stichopthalma* and *Tanaris*, though no doubt not so completely protected, are so conspicuously coloured and fly so feebly, that one cannot suppose them to be eaten by birds. The *Ornithopteras*, though I have shown by experiment that they are protected from some enemies, smell of nothing worse than petunias. I have heard that cattle will not touch peppermint or pennyroyal, though the scent of those plants is so pleasant to us. On the whole I do not doubt that slow-flying, brilliantly-coloured insects like *Trepsichrois* and *Salpinx* are more or less protected by their scent-tufts.

2nd. Colour in the tufts seems a later development than odour. In *Radena*, where they are least developed, the tufts are of very inconspicuous hues. In *Danais* they are usually not at all brilliant, though it must be admitted that the odours are also less developed in these. But in some species the hairs are curled,* and this may be an ornamental development. In the *Euplaeus* the growth of colour and odour go on *pari passu*. The brilliant ochreous tufts of *Trepsichrois* contrast finely with the black and shining blue of the wings, and are visible from a long distance. This colour appears in a great many species. In *Salpinx* the tufts are still larger, and the small terminal pair are frequently coloured differently from the others. In *Hestia* the anterior tuft seems usually coloured otherwise than the posterior, though in *H. cadellii* they are both dull grey.

3rd. The great majority of the family smell of honey or of flowers—vanilla, tuberoses, jasmine, etc., and outside of the *Danais* group, the only aberrant perfume I can remember is that of winter-green,† which is also found in butterflies of other families, and in Pyralid and Geometrid moths. But honey- and flower-like smells are the rule. This suggests the possibility that the odour-producing particles may not be manufactured by the insects, but be derived directly from the plants they frequent. At any rate the attractive scent must often be identical with that of the flowers on which they feed. So that it does not seem incredible that the female should sometimes be wooed under false pretences, and led to expect a dinner instead of a lover.

Male *Euplaeus* often meet in great swarms, haunting some particular spot in the forest for many successive days, some perching on leaves and flowers, but most circling slowly around, many of them displaying their tufts, so that the air is noticeably permeated with their fragrance. Many different species meet on these occasions, as if recognizing the

* This is also the case with a few *Euplaeus*, such as *alcatooi*.

† The same smell is given out by several tropical plants.

family relationship of all. The females may at the same time be lurking hard by in the jungle, though in smaller numbers, a few of them sometimes joining in the flight of the males. I have often observed males flying alone with expanded tufts, and I suppose they are trying to attract the female from a distance, appealing rather to her sense of smell than that of sight. In the presence of the female, the male keeps his tufts in continual action, and whether of dark or bright colours, they may well be as attractive to her eyes, as the plumes of the cock grouse or peacock are to his heus. The eyes of butterflies are so prominent, that though the female never faces the male, but keeps gliding on just before him in coy retreat, I cannot think that any grace of his escapes her notice. But her sight is probably not so strong as to recognize these ornaments at a great distance. So that the attractive colours are presumably intended to take effect at close quarters only, while with a favourable wind the odour is no doubt perceptible far away. I myself cannot generally make it out more than two or three feet away. But the forefeet of female *Danaidæ* seem to be developed into a powerful organ of smell, and even apart from this, that sense must be kept in all butterflies, since they are attracted from long distances by the scent of flowers. And few flowers are so fragrant as these insects.

List of Enganese Butterflies.

Family DANAIDÆ.

1. NECTARIA LEUCONOË, Erichson, var. ENGANIA. This slight variety seems darker than the typical *N. leuconoë*, as figured by Doubleday, just as *N. clara* (as figured by Herr Semper) is much lighter. The base is but slightly touched with creamy, and more so in the female than in the male. The dark lines in the cell of the forewing are distinct, the black transverse area there narrower and more quadrate than in *leuconoë*, the discal dark markings are more connected, and those on the hindwing are more triangular, the wedge-shaped white spot near the lower angle of the forewing is distinct, and in general the markings are very clearly cut and distinctly outlined. The male has two large whitish abdominal tufts, each with a minute red tincture of another near its outward base.

2. MACROPLEA CORUS, Fabricius, var. MICRONESIA. Somewhat resembling *M. corus* (*elisa*), with the size and shape of *M. phœbus*. Male, forewing with the spots minute, one in the cell, seven in a bent discal series, and ten submarginal dots. Hindwing with two or three discal, and one or two subapical dots. *Below*, forewing with a dot in the cell, two large discal spots with a dot below them, two subapical dots, and

a submarginal row of thirteen dots. Hindwing with three discal dots subapically, and four or five submarginal ones. The single female has the spots still fewer and less conspicuous, the submarginal ones wanting except at the apex.

As I took only two or three specimens, I can hardly describe this form as a distinct species. It seems greatly to resemble *M. semicirculus* from the Moluccas. The extreme smallness of the spots easily distinguishes it from *M. pharetena*, described by Herr Kheil from Nias. *M. pharetena* (the name was perhaps intended as an anagram of *phaenarete*) seems to be simply the female of *M. phobus*, which he also includes in his list of Nias butterflies.

3. *SALPISX PHANE*, n. sp. Male, above rich velvety blue, resplendent in some lights, the hindwing nearly equally brilliant, whereas in *S. novara* and *S. vestigiata* it is all brown. Forewing, with one costal and the usual interno-median pale blue spot, and a row of seven outer-discal ones, of which the first three are united, the third the largest. Hindwing generally with two subapical spots. *Below*, rich violet brown; the spots are sometimes wholly wanting in both sexes, but generally one or two remain subapically on the hindwing. The male before me has five subapical spots in two series on the forewing and four larger ones on the hindwing. The larger tufts are dark brown, the small terminal ones whitish. The female is less richly coloured, the hindwing being brown: there are two elongate blue spots in the interno-median space of the forewing above.

This is a local form of *S. leucostictos*, Gmelin, but it seems more distinct from the Javanese type than are either the Malaccan (*vestigiata*) or the Nicobarese (*novarae*) forms.

4. *EUPLÆA (TREPSICHOIS) MALAKONI*, n. sp. Male like *E. mulciber* but with the hindwing blue like the forewing, though less brilliant, whereas in the other species it is brown. The odoriferous patch on the hindwing occupies all the end of the cell, extending to the upper median vein. The pale blue spots of the forewing, though variable, are few and inconspicuous; in the specimen before me there are only nine in all, some very minute. The female varies greatly in the distinctness of the spots, but in all cases the forewing has no trace of blue, which distinguishes it from the other local forms of *E. midamus*. Tufts bright yellow.

5. *EUPLÆA (CRASTIA*) ENGANENSIS*, n. sp. It greatly resembles

* Mr. Moore's *Vadebra*.

E. clinena from the Moluccas, and is near *E. sepulchralis*, Butler, from Java, but with the forewing imperceptibly paler outwardly, while the hindwing is very distinctly so, both above and below. Both sexes are rich brown above, without any trace of blue; the female has a single whitish spot in the middle of the disc. *Below*, there are usually on the forewing, one large violet-white spot in the cell, two or three subapical dots, two submarginally in the median spaces; five discal ones, the lower two larger and quadrate, the middle one slender, the upper two usually minute. *Hindwing* with one spot in the cell, a semicircle of six beyond it, then a curved series of about eight, (those below the middle median vein minute or wanting), and four submarginal spots near the apex, occasionally five or six. Tufts bright ochre.

This common species outwardly resembles *Euploea simulatrix*, Wood-Mason, from the Nicobars, which, however, has a large sex-mark, somewhat as in *Salpina*, on the hindwing above. Mr. Moore placed this species first in *Vadebra* (*Crastia*) and then in *Menama*, where it certainly does not belong.

6. *EUPLEA* (*CRASTIA*?) *OCEANIS*, n. sp. Male. Above dark brown with dull bluish reflections in some lights, the margin not visibly paler. *Below* uniform dark violet-brown, the markings varying greatly, most of them elongate but small. Forewing with one spot in the cell and seven beyond it, the upper ones slender, sometimes minute; a line of five outer-discal spots, the upper ones frequently absent; the female has sometimes a sixth spot in the interno-median space, beyond the line of the others. Hindwing with a small spot in the cell, and a row of 6-7 larger ones beyond it, and then a very variable series of 5-10, all but the first (which is sometimes absent) forming nearly a straight line: beyond these there are generally a few submarginal dots, sometimes forming an irregular line of ten.

The female has an obscure whitish spot in the cell of the forewing above, and seven similar ones beyond it, the upper ones slender. Beyond this the outer discal spots are slightly indicated. Hindwing with the outer discal line of spots indicated in various degrees.

The form of the wing in this species is as in Mr. Moore's genus *Gamatoba*, but the sex-mark is peculiar. The female has an elongate white mark on the underside of the forewing, below the lower median vein. Both sexes have the hind margin broadly whitish and shining below the internal vein. In the male there is a slightly velvety grey-brown patch of altered scales forming an elongate, quadrate mass, from the internal vein nearly to the lower median, equally divided by the wrinkle representing the submedian vein.

This species can scarcely be included in any of Mr. Moore's subgenera, and I know no described form resembling it. It was unluckily out of season when I was in Engano, and I only got two or three in fair condition, the rest being worn and faded.

7. *EUPLOEA PAHAKELA*, n. sp. Male, above, forewing rich brown, not at all lighter outwardly; hindwing paler brown, slightly darkening near the margin; a single "brand" like that of *E. core*; no other markings above in either sex. *Below* there is a rounded violet white spot in the cell, and a larger pinkish one in the lower median space, one in the upper median space, and sometimes one in the space above. Hindwing with a small spot in the cell, and 3-5 minute ones beyond it. There are sometimes traces of obscure subapical and subanal dots. Tufts bright ochre. The female is very pale brown and remarkably translucent; it has a long white streak below the lower median vein on the forewing below.

This species seems to have no near allies except *E. camorta*, Moore, from the Nicobars, which obviously differs in the pale borders of the wings.

Besides the *Euploea*s mentioned, I believe I saw a species of *Calliploea*, near *mazares*, flying at Kayapu on the day of my arrival.

The *Euploea*s of Nias, as described by Herr Kheil, differ greatly from those of Engano; there can hardly be more than one species (a *Salpinx*) common to both. I append the list.

Nias.	Engano.
<i>Macroploea phaeus</i> .	<i>Macroploea micronesia</i> .
" <i>phaeretena</i> (♀ of preceding?)	
<i>Euploea</i> (<i>Tronga</i>) <i>niasica</i>	-----
" (<i>Trepsichrois</i>) <i>verhuelli</i> .	<i>Euploea</i> (<i>Trepsichrois</i>) <i>malakoni</i> .
" (<i>Penoa</i>) <i>menetriesii</i> .	-----
-----	<i>Euploea</i> (<i>Crastia</i>) <i>enganensis</i> .
-----	" " <i>oceanis</i> .
-----	" <i>pahakela</i> .
<i>Salpinx leucostictos</i> (perhaps the same as <i>phane</i>).	<i>Salpinx phane</i> .
" (<i>Isamia</i>) <i>lowii</i> .	-----
" " <i>staudingerii</i> .	-----
" (<i>Tiruna</i>) <i>ochsenheimeri</i> .	-----
-----	? <i>Salpinx</i> (<i>Calliploea</i>) sp.

The absence of the true *Euploea* (*Core* group) and of *Crastia* (*Climena* group) from Nias, and of *Tronga* and *Isamia* from Engano, are interesting.

8. *DANAIS PIETERSII*, n. sp. Pl. I, Fig. 1. Male and female, above black, forewing with the lower part of the cell, the basal part of the lower median, and the greater part of the interno-median space dull ferruginous, the veins black. Margin touched with white at intervals; a line of four submarginal white dots in the median spaces; another larger and diffused near the lower angle; an inner line of seven white marks, of which the pair in the upper median space is of some size, the other dots: a subapical band of six widely-separated diffused whitish spots from the costa, the lower two largest; below these two obscure spots in the upper median space. *Hindwing* black, a large, pale ferruginous area in the cell, and others in the submedian and internal spaces extending three-fourths towards the margin; shorter, slender ferruginous streaks in the other discal spaces: two lines of submarginal whitish dots mostly obsolete, more distinct in the median spaces; cilia alternately black and white. *Below*, similarly marked but with the spots more distinct and numerous and violet-white instead of whitish; the lower part of the forewing ferruginous, the apex, as well as the outer part of the hindwing, suffused, with dull, dark reddish, which takes the place of the black. *Hindwing*, with pale ferruginous areas in all the spaces, most of them bordered with silvery white, the veins dark; the two submarginal series of the white spots distinct and complete. In some specimens the cell is nearly white, the inner ferruginous area evanescent.

I am not able to compare this species with *D. eurydice* from Nias, but judging from the forms with which Mr. Butler compares it, that species is without the ferruginous areas on the hindwing above.

I name this for my kind friend Mijubeer Pieters, Controleur of Kroë in Sumatra. It is a common and conspicuous species in Engano, whereas Herr Kheil describes *D. eurydice* as rare in Nias.

9. *DANAIS (BAHORA) CHRYSÆA*, n. sp. Pl. I, Fig. 3. A local form of *D. philomela* from Java, differing in the subapical spots being more elongate, and in all the markings of the forewing being yellow except the two outer series, the yellow area of the cell heavily clouded with black scales except at its lower angle, in this resembling *D. crocea*. The interno-median yellow area shows is divided by a line of black scales in the middle, the quadrate discal spots are partly joined, as in *crocea*. *Hindwing* marked as in *crocea* and coloured as in *philomela*. The wings are very long and narrow. The female has the yellow area more restricted, all the spots larger and more conspicuous.

I think I saw *Parantica aglaia*, or its representative, on the day of my arrival at Engano, but no specimens were taken.

10. *RADENA LONGA*, n. sp. Pl. I, Fig. 2. A local form of the Javanese *R. juvena*. Forewing very long, falcate, the white markings mostly elongate and reduced, cell of forewing with the basal mark slender, clavate at tip, the outer one small with the upper streak absent, the discal spots beyond the cell much reduced. Hindwing, with the white area in the cell broken by two dark lines, the upper wide, the lower slender, not quite continuous; all the discal spots beyond the cell slender and elongate.

11. *RADENA MACRA*, n. sp. Pl. I, Fig. 4. A local form of *R. vulgaris*, the forewing elongate (but shorter than in *R. longa*), falcate, cell with the basal mark bifid, the upper streak slender, not so long as the lower, the outer spot narrow at its upper end, with a minute spot adjoining it above, and without the slender prolongation present in *R. vulgaris*; the three spots beyond the cell greatly, and all the others more or less reduced in size. Hindwing with all the spots reduced, that in the cell simply bifid, without any spot in the bifurcation (such as occurs in *R. vulgaris*). The submarginal dots are wanting on the apex of the forewing above and below.

These two *Radenas* are about equally common, and are found together. In the island of Sembawa, east of Java, two *Radenas* also occur, but in this case the *juvena* form is restricted to higher altitudes, though there is a zone in which both are found. These facts surprised me as I had supposed *juvena*, like *vulgaris*, to be simply a local form of *R. similis*.

The anal tufts of *R. vulgaris* and *macra* are shorter than in any other Danaid, the hairs white at the base and grey outwardly. In *R. juvena* and *longa*, they are nearly twice as long and grey throughout.

All the Enganese species of *Danais* are distinguished by their elongate form. This is also the case with several groups of butterflies in the Celebes, as shown by Mr. Wallace.

Family SATYRIDÆ.

12. *LETHE EUROPA*, Fab. scarce.

13. *MELANITIS LEDA*, Linn. (*ismene*). My specimens were dark and nearly unmarked above, but of the usual shape. Both the ocellate and the non-ocellate forms were taken.

14. *MYCALESIS MINEUS*, Linn. Common.

Family ELYMNIADÆ.

15. *ELYMNIAS DOLOROSA*, Butler, var. *ENGANICA*. A local form of

Elymnias panthera, Java, resembling *dolorosa*, as described by Mr. Butler from Nias. The forewing, however, is scarcely paler outwardly, its lower angle generally suffused with reddish; the ocelli of the hindwing are placed in diffused pale spots. Underside with two or three minute ocelli on the forewing, and six on the hindwing the first nearly all white, the others blue with the outer end dark and the pupil white. The outer part of the wing is but slightly paler than the inner, and is not "stone yellow striated with blackish" as in *dolorosa*, but pale grey-brown with red striae similar to those of the base. *E. dolorosa* has the hindwing more strongly dentate than in *panthera*, with a decided tail, but in the Enganese form the tail is much less distinct than in *panthera*. I took only females. The sex of the type of *dolorosa* is not stated.

FAMILY MORPHIDÆ.

16. *AMATHUSIA AMYTHAON*, Doubleday, var. *INSULARIS*. Above black, the disc glossed with blue in some lights up to the border of the hindwing, but not apically on the forewing, the marginal line white. The lilac band is somewhat intermediate between that of *A. portheus*, or *dilucida*, and that of *A. westwoodii*, tapering less than in the latter, extending down to the internal, and up to the upper median vein near the outer margin, very broad costally. Below, the extensive space beyond the middle striga of the hindwing almost to the submarginal line is dusky in both sexes, without the lilac gloss of the rest. On the forewing there is a smaller dusky patch chiefly costal, beyond the fifth striga. These dark spaces are scarcely visible in the allied forms. The underside seems generally darker than in *westwoodii*, the ocelli have broader dark borders, the tails are much broader and more marked with white than in any other variety; the anal black spots are touched with blue in both sexes.

The ochreous band of the female is narrow, bifid near its lower end, deeply incised outwardly along the middle median vein and partly enclosing a dark spot in its inner margin above the same vein.

One of the numerous local forms of *Amathusia amythaon*, which are by some held as species. I am unluckily unable to compare it with Felder's Javanese varieties. It differs from *portheus* and *dilucida* in the broad tails and narrower blue band. It is like Westwood's *amythaon* (now called *westwoodii*), of uncertain habitat, but the band is less tapering, broader outwardly, the tails much broader, and the outer margin of the forewing convex instead of concave.

FAMILY APATURIDÆ.

17. *CUPHA ERYMANTHIS*, Drury.

18. *ATELLA ALCIPE*, Cramer. The specimens are quite normal. Herr Kheil mentions *Atella phalanta*, and not *alcippe*, as occurring in Nias, possibly a mistake.

19. *CYRESTIS PEGIANDER*, Fabr. Resembles the Malaccan form (*themire*, Honrath), but still darker, much darker than the Tenasserim one, which is perhaps a distinct variety. I have not seen the Javanese form, but if Horsfield's drawing is correct, and the insect has only the single space between the two outermost ochreous bands white, *themire* may be considered a distinct species.

A common species in the high forest.

20. *PRECIS IDA*, Cramer. Only one taken.

21. *HYPOLIMNAS BOLINA*, Linn. The male is normal, the female infinitely variable. Occasionally it is not much unlike the male. Usually the bluish band across the forewing is obscure and there are traces of a reddish band from the disc of the forewing across the hindwing, with white discal spots beyond it. The white band across the forewing below is present or absent. Sometimes this form resembles *Danaüs pietersii* when flying. Mr. Woodford has given an interesting account of the variability of the female of this butterfly in the Solomon Islands.

22. *HYPOLIMNAS ANOMALA*, Wallace. My single male, taken at Malakoni, has only the costa and outer margin of the forewing purplish, with an outer-discal line of white spots on both wings, a submarginal line, and a few discal streaks. The hindwing has no white on the disc. No doubt the species is as variable in Engano as elsewhere.

23. *DOLESCHALLIA NIASICA*, Butler. I took two or three males which may be conspecific with the female described as *niastica*. They are much darker than males of *D. bisaltide*, the apex of the hindwing strongly suffused with black. Herr Kheil gives both *niastica* and *bisaltide* from Nias, apparently considering them distinct, but it is unlikely that two varieties so closely allied can be found together. I have recently taken typical *Doleschallia pratipa* and typical *polibete* in the same piece of jungle in Western Siam, but this is a case of two local forms overlapping at the limit of their ranges. The same is true of *Parthenos lilacinus* and *gambrisius* from Karenni to Lower Siam.

Family NYMPHALIDÆ.

24. *NEPTIS SOMA*, Moore, var. *MERIDIÆ*. Smaller than the Indian variety, and darker below, the markings of the upperside somewhat smaller and less fuliginous.

25. *NEPTIS OMBALATA*, Kheil, var. *ENGANO*. Like the Nias form described by Herr Kheil, but on the upperside the subapical white spots are smaller and better separated, the two lines of submarginal lunules distinct, and the upper white band of the hindwing much narrower and more broken. The colour of the underside is a rich red-brown, and the black borders of the white bands are very conspicuous.

FAMILY LYCENIDÆ.

•*Subfamily Aphneucinae.*

26. *HYPOLYCENA THECLOIDES*, Felder. This butterfly occurs locally in the Nicobars and the Malay Peninsula, and seems to feed on some shore-plant. It seemed scarce in Engano.

27. *EOÖXYLIDES THARIS*, Hübner. Common in the forests of the interior. The specimens are smaller than Sumatran ones, with the inner black fascia of the underside obsolescent.

28. *BINDAHARA SUGRIVA*, Horsf. Only two males taken, both quite normal. Herr Kheil mentions *B. phocides* in his Nias list, probably by mistake. I have taken *sugriva* in Great and Kar Nicobar, and again in Java.

Subfamily Deudoriginae.

29. *DEUDORIX EPIAREBAS*, Moore. The cell of the forewing is touched with red in the female.

Subfamily Lycœninae.

30. *NACADUBA ARDATES*, Moore. Common.

31. *NACADUBA VIOLA*, Moore. Only one or two seen.

32. *NACADUBA PROMINENS*, Moore. Scarce.

33. *NACADUBA MACROPHALMA*, Felder. Scarce. None of these species are mentioned as occurring in Nias, but Herr Kheil's *Plebeius kupu* is apparently the female of *N. viola*.

34. *CATOCHREYSOPS STRABO*, Fab., var. *LITHARGYRIA*. I think the true *strabo* was also seen, and no doubt *C. pandava* and *cneius* also occur. These species all seem to feed on a leguminous plant growing on the sea-beach, and are found on many small islands. *Catochrysops pandava* is per-

haps the commonest butterfly of the Nicobars. Herr Kheil mentions only *C. strabo* and *eneius* from Nias, and also *C. kandarpa* (= *strabo*).

35. EVERES PARRHASIUS, Fab., (stated by Mr. de Nicéville to be the same as the European *argiades*). Only one or two seen. This is Herr Kheil's *Plebeius polysperchinus*.

36. LAMPIDES BOCHUS, Cramer, (*Jamides bochus*). My specimens were identical with Indian ones. The species seems to vary but little throughout its range. The Nicobar form seems perfectly distinct, and should stand as *L. nicobaricus*, Wood-Mason and de Nicéville. Herr Kheil's *Plebeius siraha*, from Nias, is apparently the same as *L. bochus*, but why he should compare it with *Plebeius balliston* (*Lycenæsthes bengalensis*) I do not know.

37. LAMPIDES SUBDITUS, var. TELANJANG. Female. Above, with the inner border of the black outer area excised by an entering angle. Hindwing with the outer bluish rings bounded inwardly by a broad dark band which is suffused anally with reddish. Below, the submarginal pair of fasciæ diffused and lunular. Hindwing with the orange forming a large area discally, extending above the radial vein and to the submedian, only three of the black spots complete, the inner zigzag line obsolescent.

Only a single female taken, but *subditus* is so different from other species of *Lampides*, and this form is so distinct from *subditus*, that I have thought best to name it.

38. LAMPIDES ELPIDION, n. sp. A local form of *Lampides elpis*, the dark white-bordered submarginal spots of the hindwing separated from the basal blue by a broad unbroken dark wavy band: forewing with the blue pale and milky, the outer dark margin rather broad, the veins edged slenderly with black at the apex. Below the ground-colour is uniform pale reddish-brown, as in the dry season form of *elpis*. Prehensors as in *elpis*. I have compared this butterfly with a long series of Indian and with four Javanese males, and it seems a good local race, easily distinguished. It is very much larger than Herr Kheil's *Plebeius talinga*, which has the black border of the forewing much broader, and the inner band of the hindwing wanting.

39. LAMPIDES CELENO, Cramer. I identify this species with some doubt; a small pale-blue form, quite common along the shore. Another *Lampides*, which I have not been able to place, is pale grey above, with the border rather narrow on the forewing and reduced to a thread on the hindwing.

Herr Kheil describes a *Cyaniris* from Nias, *puspinus*, dubiously distinct from *C. puspa*. *Zizera karsandra* certainly occurs in Engano, but I do not seem to have taken it. Herr Kheil gives it in his Nias list under the name of *Plebeius lysimon*.

40. PITHECOPS HYLAX, Fab. Scarce.

*Subfamily Gerydinae.**

41. PARAGERYDUS UNICOLOR, Felder, (*horsfieldii*, Moore). The identity of *horsfieldii* and *unicolor* seems generally accepted by the German naturalists, and Mr. de Nicéville, who has examined Felder's types (three females) of *unicolor* at Vienna, has come to the same conclusion.

Paragerydus certainly does differ considerably in aspect from *Allotinus*. I think it may be kept distinct from it for the present, on account of the approximate second and third subcostal branches in the forewing of the male.

Family PIERIDÆ.

42. TERIAS HARINA, Horsf. Only one taken. Herr Kheil does not record it from Nias.

43. TERIAS HECABE, Linn.

44. TERIAS SARI, Horsf. Taken only on the hills.

45. APPIAS HIPPO, Cramer. Only females taken. It may be the Javanese *Appias lycida*, and not *hippo*.

46. HUPHINA ETHEL, n. sp. Male, above white, all the veins, including the internal and medians, black, and bordered with diffused black scales, the cell and the upper median vein heavily bordered with black, the costa tinged with lemon. Outer border rather widely and equally black, a dark, diffused outer-discal band parallel with the margin as far as the internal vein, cutting off seven submarginal spots, all white except the first, which is bright yellow, slender, the last large and out-

* In the 1889 volume of this Journal, by an unfortunate blunder I described the fore tarsi of the *Gerydinae* as like the middle and hind ones, in spite of my numerous drawings showing the contrary. I also gave Herr Kheil's *Allotinus aphocha* as equivalent to *horsfieldii*. *A. aphocha* may be distinct, though badly described, and figured only on the underside, where it is identical with *horsfieldii*. I now doubt if my proposed genus *Malais* is distinct from *Logania*, though a Bornean form resembling *L. sriwa* does have the tibiae short and thick.

wardly incised. *Hindwing*, rich ochreous yellow, (without any orange tinge) from the first subcostal vein to the hind margin; the outer border rather widely black (but not subanally), the outer part of the veins, except the submedian and especially the subcostals and radial, also black, bordered with diffused scales. *Below*, the white area is reduced on the forewing, the upper submarginal spots united in a large triangular bright yellow mass, only the lower two white, that above the upper median vein obsolescent. *Hindwing* lemon yellow, ochreous yellow on the extreme anal margin, with greenish areas above and below the cell, especially around the subcostal veins; the dark brown border very wide, attaining the cell, enclosing six yellow spots, all large except the fourth which is obsolescent, the first three more or less united.

The absence of all orange on the wings, the submarginal band of the forewing, the veins outlined with black on both wings, the very broad marginal dark band of the hindwing below, and the large extent of the yellow area, tinged with greenish below, easily distinguish this peculiar species from *Huphina lea* and *judith*.

Only one male, taken at Bua Bua.

FAMILY PAPILIONIDÆ.

47. ORNITHOPTERA NEREIS, n. sp. A local form of *O. pompeius*, Cramer. Male. Above, black, the cell immaculate, the veins of the disc black, bordered with rather conspicuous whitish rays. *Hindwing* golden yellow, bordered with a deeply scalloped black band, which is only about $\frac{1}{5}$ of an inch wide at the ends of the veins, the base black above the middle of the costal space, including the root of the cell; two (in one specimen five) black discal spots subanally in the gold. *Below*, the white streaks near the veins are more continuous, and the end of the cell is slightly touched with whitish; a little red at the base of the wings.

Female, with the outer third of the cell entirely dull whitish, the whitish streaks between the veins coalescing, and extending nearly to the outer margins, the black rays in the middle of the spaces not nearly reaching the cell; hindwing very dull golden, the border wide, the discal spots coalescing widely with each other and with the outer black band, so as to enclose small yellowish lanceolate spots in pairs divided by the veins. *Below*, the hindwing is dull pale whitish-yellow, without any golden tint; this area extends only to the upper subcostal vein and occupies two-thirds of the cell. Several males and two females were taken, but one of the latter was unluckily destroyed, and the other is worn.

This species seems nearest the South Indian *Ornithoptera minos*.

Herr Kheil calls the Nias form *O. amphrysus*, Cramer, but it seems unlikely that a Javan species should occur in Nias and not in Engano. *O. nereis* obviously differs from *amphrysus* in the absence of the yellow band across the forewing of the male. The opaque whitish-yellow of the underside of the female is a striking character, and distinguishes it from a Philippine form, which otherwise resembles it considerably, especially in the male sex.

48. PAPILIO (MENELAIDES) ARISTOLOCHIE, Fab. Only one specimen taken; it was quite small. The species seems wanting on the opposite coast of Sumatra, being apparently replaced by *Papilio antiphus*.

49. PAPILIO (ILIADES) OCEANI, n. sp. A local form of *Papilio memnon*, from Java. Male, above like *memnon*, but the lines of luteous scales on the forewing are nearly obsolete (slightly visible near the apex), and those of blue scales on the hindwing less conspicuous, and not nearly reaching the cell. Below, the red basal areas are wholly wanting, and the outer grey area much narrower, not at all enclosing the series of large black spots.

Female, forewing above heavily marked with pale luteous bands bordering the dark veins over the whole disc, entering the upper end of the cell, black rays in the middle of the spaces, the apex darker. Hindwing similar but with the luteous bands less conspicuous, enclosing a series of large black spots. Below, the red basal spots are present, but very small, the grey border slightly broader than in the male.

The absence of the red basal spots above and below in the male, above in the female, the obsolete markings of the upperside of the male, and the narrow band of the hindwing below in both sexes, easily distinguish this species from its allies.

Only one male and one female taken.

50. PAPILIO (CHARUS) HELENUS, Linn. var. ENGANIUS. The lines of luteous scales are conspicuous above and below, more or less whitish near the lower angle of the forewing; the red lunules of the upperside are absent, except the anal one, which is obscure. Below, the white area is large, the marginal white lunules are distinct, the submarginal orange ones small and obscure, wanting in the lower radial and upper median spaces, giving the insect somewhat the appearance of *Papilio prexaspes*; the female has a diffused white spot on each side of the radial vein.

51. PAPILIO (ZETIDES) AGAMEMNON, Linn. One tattered male, Bua Bua.

52. PAPILIO (ZETIDES) SARPEDON, Linn.

Family HESPERIADÆ.

53. HASORA BADRA, Moore. Two males and a female taken, normal.

54. PADRAONA PALMARUM, Moore, var. KAYAPU. The black area beyond and below the cell of the forewing is nearly obsolete, the yellow band of the hindwing very wide. This is perhaps a distinct species, but as only one male was taken, I cannot be sure.

55. CHAPRA MATHIAS, Fab.

56. UDASPES FOLUS, Cram.

57. HIDARI IRAVA, Moore. One male.

58. TAGIADES ATTICUS, Fab. The two hyaline spots below the three subapical ones are absent in the male, the two at the end of the cell are joined in the female. The white area of the hindwing is very large in both sexes, extending to the outer margin, where there are three black spots.

EXPLANATION OF PLATE I.

Fig. 1. *Danais pietersii*, n. sp.

„ 2. *Radena longa*, n. sp.

„ 3. *Danais chrysea*, n. sp.

„ 4. *Radena macra*, n. sp.

Figs. 5-8 refer to Mr. Doherty's other paper.

III.—*New and Rare Indian Lycænidae*.—By WILLIAM DOHERTY,
Cincinnati, U. S. A. Communicated by the NATURAL HISTORY SECRETARY.

[Received 9th March 1891:—Read 6th May, 1891.]

(With Plate I, Figs. 5-8.)

Family LYCÆNIDÆ.

Subfamily THECLINÆ.

1. ARHOPALA KHAMTI, n. sp. Pl. I, Fig. 5.

Near *A. aenea*, Hew., differing in the dark, dull indigo-blue of the upperside, and the darker shade of the underside, with the terminal cell-spot remote from that in the lower median space of the forewing; hindwing with a large subanal ocellus bordered with metallic green, which

extends to the lower median vein; a dark spot edged with whitish in the lower median space. Lobe and tail large.

Margherita, Upper Assam.

2. *FLOS ANAMUS*, n. sp. Pl. I, Fig. 6.

Female, like *Flos asoka*, but with the blue of the upperside pale, slightly violet outwardly, not reaching up to the upper radial vein. Below, forewing with the transverse fasciæ much duller, and more regular, those in the interno-median space of the forewing obscure. Hindwing with the pale basal stripe absent, the base all dark, touched with scarlet costally; the lower half of the wing very dark, especially subabdominally, the markings there obscured; beyond the dark base there is a broad lilac area transversely from the costa to the hind-margin, containing a sinuous irregular band from the costa to the median vein, continued by a small separate spot in the interno-median space; an obscure ocellus in the lower median space submarginally, no other metallic markings. The species also resembles the Himalayan form of *Flos fulgidus*, Hew., but obviously differs in the colour of the upperside; and on the underside in the whitish spot at the end of the cell of the forewing, which is narrow and conspicuous, (broad and dull in *fulgidus*); the lower part of the hindwing is much darker, and the pale costal band absent.

Margherita, Upper Assam.

3. *ACESINA ZEPHYRETTA*, n. sp.

Male, above dark brown, a small diffused light blue area occupying less than a sixth of the forewing, including part of the cell, the interno-median and the lower median space, extending just above the middle median vein, the veins dark; hindwing unmarked, the cilia whitish, especially apically. Below brown, slightly glossed with violet, except the spots, which are darker and encircled by broad whitish rings. There is no costal white spot on the hindwing, but the apex is chiefly whitish, the disc irrorated with whitish scales, the ocelli obsolescent.

This species, in which the male resembles a female, is obviously distinct, and seems to connect *A. paraganesa* with the other species of *Acesina*.

Margherita, Upper Assam.

4. *ACESINA ARIEL*, n. sp.

Male, above violet-blue (dull in some lights) over half of the hindwing, and rather more than half of the forewing, the blue areas rounded. Underside uniform brown, strongly glossed with violet, not irrorated with whitish scales, the markings annular, scarcely darker than the

ground-colour, with slender, pale violet-whitish rings; no costal white patch on the hindwing, the apex not whitish; three distinct subanal ocelli with metallic bluish-green irides. The violet hue of the upperside distinguishes it from *A. ammon* and *ammonides*, as well as the entire absence of the costal white patch on the hindwing below.

Margherita, Upper Assam.

5. *ACESINA AMMONIDES*, n. sp. (= *ammon*, mihi, nec Hewitson).

The Tenasserim form of *A. ammon*, which I have now been able to compare with a specimen from Pahang (Malay Peninsula) kindly lent me by Mr. de Nicéville, and one from Perak, taken by myself, seems a good local race, which I distinguish by the above name. In *A. ammon* the blue is slightly tinged with violet (but much less than in *A. ariel*), and occupies less than half of the forewing and hardly more than a quarter of the hindwing. In *ammonides*, the blue is more azure in tint, extends well beyond the cell all around it, and occupies half of the forewing and half of the hindwing. Below, in *ammon* there is only the conspicuous white costal mark on the hindwing. In *ammonides*, the apices of both wings, especially the hindwing, are strongly suffused with whitish, and there are whitish scales abdominally on the hindwing, and a quadrate discal white spot between the lower two median veins. In one specimen of *ammonides*, the three minute ocelli of the hindwing are touched with metallic gold; *ammon* is without metallic scales.

Tenasserim valley.

Besides those above mentioned, I took the following species of this subfamily at Margherita, Upper Assam. *Zephyrus distortus* (*Zinaspia distorta*, de Nicéville), *Flos moellerii*, *Darasana perimuta* and *paramuta*, *Arhopala teesta*, *singla*, *centaurus*, *amantes*, *rama*, *anarte*, *belphoebe*, *bazalus* and *camdeo*.

Subfamily *APHNÆINÆ*.

6. *DRINA MANEIA*, Hew.

The veins of the forewing are marked with raised lines of light-brown scales in the blue area, somewhat as in *Papilio ganesa* or *Argynnis childrenii*. The veins so marked are the three medians, the lower radial and the submedian, besides a line in the interno-median space, and two terminally in the cell. These are presumably scent-glands.

The venation agrees well with that of *Drina donina*, the type of the genus; so that Mr. de Nicéville, who had never seen the species, showed some acuteness in placing it here, in spite of its wholly different appearance.

Rare at Padang Rongas, Perak.

Genus *THEIX*, novum.

Male, forewing with the subcostal vein five-parted (including the vein itself as a branch), the first branch originating one-third before the end of the cell, the third a little before the end, the second nearer the first than the third, the fourth from the third halfway to the apex, the fifth close to the apex, very short. Discocellular veins nearly straight, the lower half again as long as the upper, cell slightly longest at its lower angle, second bifurcation of the median vein a little before its end. Submedian vein exceedingly remote from the median, straight for nearly half of its length, then bent downwards like a bow. In the interno-median space discally, there is a deep oblique depression on the upperside, covered with short grey down, and bearing a conspicuous extensile tuft of long orange hairs produced downwards and outwards over the depression. Hindwing with the two lower median branches forking simultaneously from the end of the cell; a long tail from the submedian, a short one from the lower median vein.

The curious scent-organ in the middle of the forewing of the male, resembling that of *Dacalana* and *Arrhenothrix*, has so distorted the venation that I have thought it advisable to separate this genus from *Neocheritra*. Whether the male has five and the female four subcostal branches, as in that genus, I do not know.

7. *THEIX GAMA*, Distant, (*Neocheritra gama*).

Above black, a tuft of orange hairs over a small grey cavity in the middle of the forewing. Hindwing with the lower part white, containing two black spots; above this grey, with three black spots on the boundary between the grey and the white; the upper part of the wing black. Below as in the female. The species apparently mimics *Boöxyliides tharis*.

Rare at Padang Rengas, Malay Peninsula. I have also taken it in the mountains of south-western Sumatra.

Subfamily *PORITINÆ*.

Genus *MASSAGA*, mihi. I find that in the male of *M. pediada*, the type of this genus, there is a narrow tuft of prostrate black hairs arising at the end of the cell, extending beyond it along the upper border of the upper median vein. This was pointed out to me in *M. pharygo* by Mr. de Nicéville and I afterwards found it in *M. pediada*. In *M. potina* it is apparently present, but very small and inconspicuous. In all these species the upper tuft is of considerable size, yellowish or whitish, turned upwards along the upper subcostal vein, in a large whitish patch. In

Poritia, the lower tuft is absent, the upper one black or dark brown, without the whitish patch.

Subfamily LYCÆNINÆ.

Genus PHENGARIS, novum. The splendid Chinese butterfly *Lycœna atroguttata*, Oberthür, deserves to be placed in a separate genus or subgenus, distinguished from *Lycœna* by the upper discocellular vein of the hindwing being short and angled outwardly, the lower discocellular meeting the median vein opposite its second forking.

This butterfly is certainly the finest of the subfamily, unless the *danis* group of *Cyaniris* be excepted. I was not able to detect any odour about it, but it has all the air of a protected species. I often saw it in the meadows of the Kutcha Naga country, Naga Hills, from 6000 to 8000 feet elevation, flying very slowly and visible from a great distance, so that I caught a good number, in spite of its rarity. The character of its markings, round black spots on a pure white ground, is very remarkable. It is hard to avoid thinking *Tajuria maculata*, Hew. a mimic of this species, though it seems to live at a lower elevation, and further to the westward. *Taraka hamada* is somewhat similarly marked, and is obviously protected.

I have taken the name *Phengaris*, which means a daughter of the moon, from the modern Greek.

Subfamily GERYDINÆ.

8. GERYDUS HERACLEION, n. sp.

Male, forewing less acute than in *G. symethus*, hindwing rounded. Above, brown, forewing with a slaty gloss, the apex darker, a broad oblique white band from the upper end of the cell and beyond it (above the cell it is obscure), almost to the middle of the interno-median space, the outer part dehiscient along the lower median vein, projecting furthest in the lower median space (unlike *symethus*). The band is much broader than in *Gerydus biggsii*, Distant, (*gopara*, de Nicéville). Hindwing all dark. Below, the white band of the forewing is obscure and broken, the spot in the lower median space quite separate from and more distinct than the rest, the transverse lunular band obsolete in the forewing, three costal ring-spots, three small subapical lunules. Hindwing, much less clouded with blackish than in *G. croton*, the lunular transverse band nearly regular, the basal spots quadrate, the submarginal black dots very distinct.

Perak, Malay Peninsula.

The species is larger than *G. symethus*, not quite so large as *G. ancon*. The upper median vein of the male is naked above and swollen from the end of the cell one-fourth towards the outer margin. This may be regarded as a generic character of *Gerydus*, since it occurs at any rate in *G. symethus*, *biggsii*, *boisduvalii*, *heracleion*, *irroratus* var. *assamensis*, and *ancon*. In *G. croton* the swelling is indistinct, and the vein is covered with black scales.

9. *GERYDUS IRRORATUS*, Druce, var. *ASSAMENSIS*, nov. Pl. I. Fig. 7.

Above, unmarked except by a small, pale, longitudinal area around the base of the upper median vein on the forewing. Below, pale grey-brown, without the dark markings of *G. boisduvalii*, a small pale area on the forewing below the middle median vein, the markings lunular, those in the cell of the forewing reduced, the transverse discal band of the forewing subapical, extending only to the upper median vein, a single conspicuous dark lunule near the lower angle; the transverse band of the hindwing regular, an undulated, continuous submarginal dark line.

Dhansiri Valley, Naga Hills.

It resembles *G. melanion* from the Philippines, but is without the white area near the lower angle of the forewing above. It may be conspecific with Mr. Druce's *G. irroratus* (from Siam) which has never been figured or properly described.

I have taken what may be the female of *Gerydus irroratus* in Perak. Some pale markings represent the broken white band of *G. boisduvalii*, the hindwing is angled at the upper median vein.

The figure represents the transverse band of the hindwing incorrectly; it is really composed of separate annular lunules.

10. *LOGANIA MASSALIA*, n. sp., Pl. I, Fig. 8.

Female. Above black, a round, dull white discal area on the forewing from just above the upper median vein almost to the submedian vein. Below irregularly speckled and variegated; forewing with the costal and apical parts ochreous-brown, the rest blackish. Hindwing also tinged with ochreous, a submarginal dark area, and obscure dark transverse bands. Hindwing not angled, the margin entire.

Nearest an undescribed *Logania* from Perak, Malay Peninsula, which, however, has the upperside marked as in *L. marmorata*, and the margin undulated,

Margherita, Upper Assam.

EXPLANATION OF PLATE I.

- Fig. 5. *Arhopala khamti*, n. sp. (Assam.)
 Fig. 6. *Flos ahamus*, n. sp. (Assam.)
 Fig. 7. *Gerydus irroratus*, Drace, var. *assamensis*, var. nov. (Assam.)
 Fig. 8. *Logania massalia*, n. sp. (Assam.)
 Figs. 1—5 refer to the previous paper on the Butterflies of Eugano.

IV.—*Materials for a Flora of the Malayan Peninsula.*—By GEORGE KING, M. B., LL. D., F. R. S., C. I. E., Superintendent of the Royal Botanic Garden, Calcutta. No. 3.

(Continued from page 206 of Vol. LIX of 1890.)

[Received 2nd March 1891. Read April 1st 1891.]

In the arrangement of the Natural families which is being followed in these papers (that of DeCandolle as modified by the late Mr. Bentham and Sir Joseph Hooker), the family *Dipterocarpeae* should have preceded *Malvaceae*. Delays have, however, occurred in the elaboration of that family; and, rather than postpone the publication of the remaining three *Thalamifloral* orders, I have decided to submit my account of these to the Society now, deferring my paper on the *Dipterocarpeae* and on the previously omitted *Anonaceae* to a future occasion.

ORDER XVII. MALVACEÆ.

Herbs, shrubs or trees; herbaceous portions often stellate-hairy or scaly. *Leaves* alternate, palminerved, simple, lobed, or rarely compound. *Stipules* free, sometimes caducous. *Bracteoles* 3 or more, free or combined, often forming an epicalyx. *Flowers* axillary or terminal, solitary, fascicled or cymose-paniculate, regular, hermaphrodite or 1-sexual. *Sepals* 5, valvate, free or connate. *Petals* 5, twisted-imbricate. *Stamens* ∞ , rarely definite, adnate to the base of the petals; filaments monadelphous, forming a tube; anthers oblong or reniform, cells sinuous or twisted, linear or annular, ultimately 1-celled bursting longitudinally. *Ovary* 2-many-celled, entire, or lobed, of 2-5 or usually more carpels whorled round a central axis; styles connate below or throughout their length; ovules 1 or more, curved, attached to the inner angle of each carpel. *Fruit* of dry cocci, or capsular and loculicidal, often large and woody. *Seeds* reniform or obovoid, sometimes arillate; albumen scanty, often mucilaginous or 0; embryo curved; cotyledons leafy, usually

folded or crumpled.—Distrib. Abundant in warm regions, common in temperate, absent from arctic. Genera 57; known species about 700.

A. Staminal tube entire, or but slightly divided at the apex.

Tribe I. *Malvaceae*. Herbs or shrubs. Ripe carpels separating from the axis. Styles as many as the carpels.

Ovules solitary; carpels with convergent, often beaked, apices 1. *Sida*.

Ovules 2 or more; carpels with divergent, not beaked, apices 2. *Abutilon*.

Tribe II. *Urenaeeae*. Styles or stigmatic branches twice as many as the carpels.

Fruit of indehiscent cocci 3. *Urena*.

Tribe III. *Hibisceae*. Herbs or shrubs. Fruit capsular. Sepals leafy. Staminal-tube truncate or 5-toothed at the apex.

Calyx toothed: stigmas distinct, spreading ... 4. *Hibiscus*.

„ truncate: stigmas united 5. *Thespesia*.

B. Staminal tube short or divided into single filaments to its base.

Tribe IV. *Bombacidae*. Trees. Sepals leathery: styles connate or free. Fruit capsular.

Leaves digitately compound, calyx truncate or irregularly 3 to 5-lobed; seed silky outside.

Anthers solitary 6. *Bombax*.

„ in groups of 2 or 3 7. *Eriodendron*.

Leaves simple, usually scaly; fruit woody, muricate; seeds arillate.

Calyx tubular or bell-shaped.

Anthers linear, cells sinuous ... 8. *Durio*.

Anthers globose, opening by a pore ... 9. *Boschia*.

Calyx dilated at the base.

Calyx finally forming a cushion-shaped annulus 10. *Neesia*.

Calyx 5-pouched at the base, petals inserted on the calyx 11. *Calostegia*.

1. *SIDA*, Linn.

Herbs or undershrubs. Leaves entire or lobed. Bracteoles 0. Calyx of 5 valvate sepals, tubular below. Corolla of 5 petals, free above, connate below and adnate to the tube of the stamens. Staminal-tube dividing at the summit into numerous anther-bearing filaments. Carpels 5 or more, whorled; styles as many as the carpels, stigmas terminal.

Ripe carpels separating from the axis, generally 2-awned at the summit, and dehiscing irregularly or by a small slit. *Seed* solitary, pendulous or horizontal; radicle superior.—*Distrib.* A genus of about 80 species, most of them being tropical weeds.

1. *S. MYSORENSIS*, W. & A. Prod. I, 59. A sub-erect, sometimes decumbent, herb 1 to 2 feet high, covered with more or less glutinous hairs. *Leaves* cordate-ovate, acuminate, coarsely serrate-crenate, 1.5 to 2.5 in. long and 1 to 1.5 in. broad; petiole about half as long as the blade. *Stipules* linear, less than half as long as the petiole. *Flowers* less than .5 in. in diam., in few-flowered axillary racemose cymes, corolla yellow; pedicels shorter than the petioles, jointed near the middle. *Carpels* shorter than the calyx, sub-glabrous, each with a short awn, or awnless. Mast. in Hook. fil. Fl. Br. Ind. I, 322; Thwaites Enum. 28. *S. hirta*, Wall. Cat. 1855, not of Lam. *S. urticifolia*, W. & A., l. c. *S. nervosa*, Wall. Cat. 1853 E. *S. olens*, Ham. in Wall. Cat. 1874. *S. glutinosa*, Roxb. Hort. Beng. 97; Fl. Ind. iii, 172; Wall. Cat. 1855, not of Cav. *S. tenax*, Ham. in W. & A. Prodr. i, l. c.; Wall. Cat. 1855, E. F. *S. fasciculiflora*, Miq. Fl. Ind. Bat. i, Pt. 2, 140. *S. radicans* Cav. Diss. i, 8; W. & A. Prod. i, 59.

A weed by roadsides; in Perak and probably in the other provinces. *Distrib.* India, Java.

2. *S. CARPINIFOLIA*, L. An undershrub 2 to 3 feet high; glabrous or sub-glabrous; a few minute stellate hairs on the stems and petioles. *Leaves* linear-lanceolate, acute, serrate, 2 to 3 in. long and .25 to .35 in. broad; petioles .1 to .2 in. *Stipules* subulate, nerved, much longer than the petiole. *Flowers* .5 in. in diam, solitary, axillary; corolla yellow, peduncles as long as the petiole, jointed, minutely bracteolate. *Carpels* shorter than the sub-globose ribbed calyx, glabrous, rugulose, each with 2 short awns. DC. Prod. i. 460. Mast. in Hook. fil. Fl. Br. Ind. i. 323; Wall. Cat. 1871. *S. acuta*, Burm.; Cav. Diss. i p. 15, t. 2, f. 3; DC. Prodr. i. 461; Wall. Cat. 1868, 1, 2, 3, 4, 5; Roxb. Fl. Ind. iii. 171; W. & A. Prodr. i. 57; Dalz. & Gibs. Bomb. Fl. 17; Thwaites Enum. 27; Miq. Fl. Ind. Bat. i. Pt. 2. p. 143; Wight Ic. t. 95; Bl. Bijdr. 55; Wall. Cat. 1868 G. *S. lanceolata*, Roxb. l. c. 175; Wall. Cat. 1868 F. *S. stipulata*, Cav. Diss. i. t. 3, f. 10; DC. Prodr. i. 460; W. & A. Prodr. l. c. *S. Stauntoniana*, DC. l. c.; *S. scoparia*, Lour. ex W. & A. l. c.

In all the provinces as a weed. *Distrib.* India and Tropics generally.

3. *S. RHOMBIFOLIA*, Linn. sp. 961. An erect under shrub 2 to 3 feet high, from glabrous to hoary, stellate-pubescent. *Leaves* varying

from ob-lanceolate or obovate to rhomboid, but always with tapering bases, serrate to crenate; under surface hoary, rarely green; length .5 to 2.5 in., petiole .1 to .2 in. *Stipules* setaceous, longer than the petioles. *Flowers* .5 in. in diam., axillary, solitary; corolla yellow, rarely white; peduncles much longer (sometimes six times) than the petioles, variously and sometimes indistinctly jointed, ebracteolate. *Carpels* smooth or pubescent, or reticulate, each usually with 1 or 2 rather long awns, sometimes awnless, generally longer than the calyx. Mast. in Hook. fl. Fl. Br. Ind. i. 323; Miq. Fl. Ind. Bat. i. pt. 2. p. 142; DC. Prodr. i. 462; Roxb. Fl. Ind. iii. 176; Wall. Cat. 1862, 2; Thwaites Enum. 28. *S. canariensis*, Willd.; DC. Prodr. i. 462. *S. compressa*, Wall. Cat. 1866; DC. Prodr. i. 462.

This very polymorphic species has been divided into varieties by Dr. Masters in Hooker's Fl. Br. Ind. l.c. as follows:—

"Var. 1. *scabrida*, W. & A. Prodr. i. 57 (sp.); sprinkled with rigid hairs, leaves concolorous, peduncles joined at the base, carpels awned.

"Var. 2. *retusa*, Linn. (sp.); leaves obovate retuse hoary underneath, peduncles equalling the leaves jointed above the middle, carpellary awns short.—Cav. Diss. i. t. 3, f. 4, and Diss. v. t. 131, f. 2; Bl. Bijdr. 75; W. & A. Prodr. i. 38; Wall. Cat. 1870; DC. Prodr. i. 462; Roxb. Fl. Ind. iii. 175; Dalz. & Gibs. Bomb. Fl. 17; Miq. Fl. Ind. Bat. i. pt. 2, 142. *S. chinensis*, Retz ex Roxb. Hort. Beng. 97; Fl. Ind. iii. 174. *S. philippica*, DC. Prodr. i. 462; W. & A. Prodr. l.c.; Wall. Cat. 1869; Rheede Hort. Mal. x. 18; Rumph. Amb. v. t. 19.—The *S. corynocarpa*, Wall. Cat. 1870, seems to be a form of this variety, with densely intricate woody branches, and long carpellary awns.

"Var. 3. *rhomboidea*, Roxb. Hort. Beng. 50; Fl. Ind. iii. 176 (sp.); leaves rhomboid hoary beneath, peduncles jointed at the base, carpellary awns very short inflected. DC. Prodr. i. 462; W. & A. Prodr. i. 57, Wall. Cat. 1862 E., 1863; Thwaites Enum. 28. *S. rhombifolia*, Wall. Cat. 1862 F.? *S. orientalis*, Cav. Diss. i. t. 12.—The flowers expand at noon (Roxb.).

"Var. 4. *obovata*, Wall. Cat. 1864 (sp.); leaves 1½ by 2 in., broadly obovate, hoary beneath, apex coarsely toothed, base cuneate, petiole ¼ in., peduncle longer than the petiole shorter than the blade.

"Var. 5. *microphylla*, Cav. Diss. i. t. 12, f. 2 (sp.); leaves small, elliptic dentate hoary beneath, peduncle slightly exceeding the petiole, carpels 5-7 awned.—Roxb. Fl. Ind. iii. 170; DC. Prodr. i. 461."

In all the provinces—a common weed. Distrib. The Tropics generally.

4. *S. CORDIFOLIA*, Linn. spec. 961. An erect softly hairy undershrub

2 to 3 feet high, the hairs on the branches and petioles long and spreading. *Leaves* oblong-cordate, obtuse, rarely acute, crenate; both surfaces, but especially the pale lower surface, soft hairy; length 1.25 to 2 in., breadth .8 to 1.25; petiole slightly longer than the blade. *Stipules* linear, less than half the length of the petiole. *Flowers* .6 in. in diam., axillary, solitary; corolla yellow; peduncles jointed near the apex, varying in length, the lower longer, the upper shorter, than the petioles. *Carpels* boldly 3-angled, reticulate, sub-glabrous, crowned by 2 strong, divergent, retro-hispid awns. DC. Prod. i. 464, Roxb. Fl. Ind., iii. 177; Wall. Cat. 1849; W. & A. Prodr. i. 58; Thwaites Enum. 28. Dalz. & Gibs. Fl. Bombay, 17; Mast. in Hook. fil. Fl. Br. Ind. i. 324, and in Oliver's Fl. Trop. Afr. i. 181; Miq. Fl. Ind. Bat. i. pt. 2, 140. *S. herbacea*, Cav. Diss. i. 19, t. 13, f. 1; DC. Prodr. i. 463. *S. micans*, Cav. Diss. i. 19, t. 3, f. 1. *S. rotundifolia*, Cav. Diss. i. 20, t. 3, f. 6, and Diss. vi. t. 194, f. 2; Wall. Cat. 1849, D; DC. Prodr. i. 464. *S. althai-folia*, Swartz, Guill. & Per. Fl. Seneg. i. 73.—Rheede Hort. Mal. x. t. 54.

In Malacca: and probably in all the Provinces as a weed. Distrib. The Tropics generally.

2. ABUTILON, Gærtn.

Herbs or undershrubs more or less covered with down. *Leaves* angled or palmately-lobed. *Inflorescence* axillary or terminal. *Bracteoles* 0. *Calyx* of 5 valvate sepals, tubular below. *Corolla* of 5 petals, free above, connate below and adnate to the tube of the stamens. *Staminal-tube* divided at the apex into numerous filaments. *Carpels* 5-8. *Styles* as many as the carpels. *Ripe carpels* separating from the axis, awned or not, 1- or more-seeded. *Seeds* reniform. Distrib. About 70 species, all tropical or subtropical.

A. INDICUM, G. Don. Gen. Syst. i. 504. An annual or perennial undershrub. *Leaves* broadly cordate, irregularly and coarsely toothed or sub-entire, pale and minutely pubescent on both surfaces, often with a few longer hairs intermixed, length 1 to 2 in., breadth 1 to 2 in.; petiole usually longer than the blade. *Flowers* 1 in. in diam, axillary, solitary, the peduncles longer than the petioles, jointed near the top; corolla yellow. *Sepals* ovate, acute, shorter than the spreading petals. *Carpels* 15 to 20, longer than the calyx, truncate or with short spreading awns, tomentose at first, ultimately sub-glabrous. *Seeds* dark brown, minutely stellate-hairy. Mast. in Hook. fil. Fl. Br. Ind. i. 326; *A. asiaticum*, W. & A. Prodr. i. 56, not *Sida asiatica*, Linn.; W. & A. Prodr. i. 56; Wight Ic. t. 12; Dalz. & Gibs. Bomb. Fl. 18; Thwaites Enum. 27; Mast. in Oliv. Fl. Trop. Afr. i. 186; Miq. Fl. Ind.

Bat. i. pt. 2, 146. *Sida indica*, L.; DC. Prodr. i. 471; Cav. Diss. i. p. 33, t. 7, f. 10; Roxb. Fl. Ind. iii. 179; Wall. Cat. 1859, 1, 2, D. F. *Sida populifolia*, W. & A. l.c. *A. populifolia*, G. Don. l.c. *Sida populifolia*, DC. Prodr. i. 470; Cav. Diss. i. t. 7, fig. 9; Roxb. Fl. Ind. iii. 179; Bl. Bijdr. 79. *S. Beloere*, L'Her. Stirp. i. 130. *S. Eteroomischos*, Cav. Diss. ii. 55 and v. p. 275, t. 128.

Singapore, Selangore and probably in all the other provinces. A weed.

3. URENA, Linn.

Herbs or undershrubs, more or less covered with rigid stellate hairs. *Leaves* angled or lobed. *Flowers* clustered. *Bracteoles* 5, adnate to the 5-cleft calyx, sometimes coherent at the base into a cup. *Petals* 5, often tomentose at the back, free above, connate below and united to the base of the tube of the stamens. *Staminal-tube* truncate or minutely toothed. *Anthers* nearly sessile. *Ovary* 5-celled, cells 1-ovuled, opposite the petals; stigmatic branches 10; stigmas capitate. *Ripe carpels* covered with hooked bristles or smooth, indehiscent, separating from the axis when ripe. *Seed* ascending; cotyledons bent and folded; radicle inferior. Distrib. Species 4-5, natives of tropical and sub-tropical countries, 2 only being confined to Asia.

U. LOBATA, Linn. Spec. 974. A herbaceous undershrub 1 to 3 feet high, more or less hairy. *Leaves* very variable; the lower rotund to reniform, more or less cordate at the base, the apex usually acute, edges with 5 to 7 shallow lobes or sub-entire, 5 to 7-nerved; length 1 to 2 in., breadth 1 to 2.5 in.; upper leaves smaller and sometimes ovate to linear-lanceolate, 3-nerved. *Petiole* shorter than the blade; bracteoles oblong-lanceolate, as long as the sepals. *Corolla* pink, .5 to 1 in. in diam. *Carpels* tomentose, and with many smooth hooked spines. Mast. in Hook. fil. Fl. Br. Ind. i. 329; Miq. Fl. Ind. Bat. i. pt. 2, p. 149; Cav. Diss. iv. p. 336, t. 185, fig. 1; Miq. Pl. Jungh. 283; DC. Prodr. i. 441; Roxb. Fl. Ind. iii. 182; W. & A. Prodr. i. 56; Wall. Cat. 1928; Dalz. & Gibs. Bomb. Fl. 18; Thwaites Enum. 25; Miq. Fl. Ind. Bat. i. pt. 2, 148. *U. cana*, Wall. Cat. 1930 B. *U. palmata*, Roxb. Fl. Ind. iii. 182. *U. tomentosa*, Bl. Bijdr. 65.

All the Provinces: a weed. Distrib. The tropics generally.

Var. 1. *sinuata*, Miq. Fl. Ind. Bat. l.c.; leaves deeply 5-lobed, the lobes narrowed at the base, serrate, often pinnatifid, bracteoles linear; flowers often smaller than in the typical plant. *U. sinuata*, Linn.; DC. Prodr. i. 441; Roxb. Hort. Beng. 50; Fl. Ind. iii. 182; Wall. Cat. 1933 E.; W. & A. Prodr. i. 46; Hook. Fl. Br. Ind. i. 329; Thwaites Enum.

Pl. Cey. 25; Dalz. & Gibs. Bomb. Fl. 18. *U. muricata*, DC. Prodr. i. 442. *U. Lappago*, DC. Prodr. i. 441. *U. morifolia*, DC. Prodr. i. 442? *U. heterophylla*, Smith in Rees' Cycl. 37; Wall. Cat. 1933 E, F, G, H, K. *U. tomentosa*, Wall. Cat. 1933 H.;—Burm. Zeyl. t. 69, f. 2.

Distributed like the last.

Var. 2. *scabriuscula*, DC. Prodr. i. 441 (sp.); herbaceous; leaves roundish, scarcely lobed, with 1-3 glands beneath; bracteoles linear, longer than the sepals. *U. scabriuscula*, Wall. Cat. 1928 F; W. & A. Prodr. i. 46; Dalz. & Gibs. Bomb. Fl. 18.

4. HIBISCUS, Linn.

Herbs, shrubs, or trees. *Leaves* stipulate, usually more or less palmately-lobed. *Inflorescence* axillary, rarely terminal. *Bracteoles* 5 or more, free, or connate at the base. *Calyx* 5-toothed or 5-fid, valvate, sometimes spathaceous. *Petals* 5, connate at the base with the staminal-tube. *Staminal-tube* truncate or 5-toothed at the summit; filaments many; anthers reniform, 1-celled. *Ovary* 5-celled, cells opposite the sepals, each with 3 or more ovules; styles 5, connate below; stigmas capitate or sub-spathulate. *Capsule* loculicidally 5-valved, sometimes with a separate endocarp, or with false dissepiments forming a spuriously 10-celled fruit. *Seeds* glabrous, hairy or woolly. About 150 species; distributed chiefly in the tropical regions of both hemispheres.

Calyx spathaceous, deciduous ... 1. *H. Abesmoschus*.

Calyx persistent, 5-cleft.

Bracteoles of involucre distinct, their

apices spatulate ... 2. *H. Surattensis*.

Bracteoles united at the base, nearly as

long as the calyx ... 3. *H. macrophyllus*.

Bracteoles united into a cup much shorter than the calyx.

Involucre and calyx softly pubescent 4. *H. tiliaceus*.

" " rugulose 5. *H. floccosus*.

1. *H. ABESMOSCHUS*, Linn. Spec. 980. A stout annual under-shrub 2 to 3 feet high: young branches and peduncles retro-hispid, all other parts hispid or stellate-hispid. *Leaves* variable, usually with 3 to 5, deep, oblong-lanceolate or linear, serrate-crenate, acute lobes, sometimes hastate or sagittate, the base always rounded; length and breadth 3 to 5 in.; petiole longer than the blade: stipules minute, subulate, fugaceous. *Flowers* 3 in. in diam., axillary, solitary; peduncles shorter than the petioles, ebracteate. *Involucres* 8 to 12, linear, .5 to .75 in. long. *Calyx* 1.25 in. long, toothed at the apex. *Corolla* yellow with a crimson

centre, glabrous. *Capsule* oblong, pointed, hispid, becoming subglabrous, 1 to 3 in. long. *Seeds* reniform, striate, glabrous, musky. Mast. in Hook. fl. Fl. Br. Ind. i. 342 (excl. syn. *H. sagittifolius*, Kurz.); DC. Prod. i. 452; Roxb. Fl. Ind. iii. 202; Griff. Not. iv. 521. *Abelmoschus moschatus*, Mönch; W. & A. Prod. i. 53; Wight Ic. t. 399; Wall. Cat. 1915, F, G, H, I, K, L; Thwaites Enum. 27; Miq. Fl. Ind. Bat. i. pt. 2, 151. *H. flavescens*, Cav. Diss. iii. t. 70, f. 2; DC. l.c. 454. *H. spathaceus*, Wall. Cat. K. *H. ricinifolius*, Wall. Cat. 1915. *Bamia chinensis*, Wall. Cat. 1616? *Hibiscus pseudo-abelmoschus*, Bl. Bijdr. 70. *H. longifolius*, Willd. Spec. iii. 827; DC. Prod. i. 450. *Bamia multififormis* and *betulifolia*, Wall. Cat. 1917 and 1918.

In all the Provinces; cultivated or naturalised. Distrib. the tropics generally.

2. *H. SURATTENSIS*, Linn. Spec. 979. A weak straggling undershrub; the branches, petioles and peduncles with small recurved prickles and a few soft spreading pale hairs. *Leaves* palmately 3 to 5-partite, rarely ovate, sub-entire, serrate, sparsely pilose; length and breadth 1.5 to 3 in.; petiole slightly longer than the blade. *Stipules* broadly ear-shaped. *Flowers* 2 to 2.5 in. long, solitary, axillary, corolla yellow with dark centre; bracts of involucre 10 to 12, linear with spatulate apices. *Capsules* membranous, the individual carpels with 3 bold aculeate nerves and a long terminal point. *Seeds* with long straight brittle yellowish hairs. Mast. in Hook. fl. Fl. Br. Ind. i. 334; Miq. Fl. Ind. Bat. i. pt. 2, 161; Bl. Bijdr. 68; DC. Prodr. i. 449; W. & A. Prodr. i. 48; Roxb. Fl. Ind. iii. 205; Wight Ic. t. 197; Cav. Diss. iii. t. 53, f. 1; Thwaites Enum. 26; Wall. Cat. 1893, 1, 2, 3, D, E, F, G; Dalz. & Gibs. Bomb. Fl. 20; Mast. in Oliv. Fl. Trop. Afr. i. 201; Miq. Fl. Ind. Bat. i. pt. 2, 161. *H. furcatus*, Wall. Cat. 1896 C, not of Roxb.

Malacca, Perak, and probably in the other Provinces. Distrib. The tropics generally.

This has a decumbent or even climbing habit.

3. *H. MACROPHYLLUS*, Roxb. Hort. Bēng. 51. A large shrub or small tree, all parts more or less covered with pale soft minute velvety tomentum; the young branches, petioles, pedicels, bracteoles and calyx bearing, in addition, numerous more or less deciduous tufts of long spreading stiff tawny hairs. *Leaves* large, on long petioles, cordate-orbicular to reniform, the apex shortly sharply and abruptly acuminate, the edges entire; palmately 7 to 9-nerved; length and breadth 7 to 12 in.; petiole usually longer than the blade. *Stipules* oblong, convolute, hispid-tomentose, 3 to 4 in. long. *Flowers* in terminal cymes, pedicels

1.5 to 2 in. long, articulate near the apex and bearing two large broadly ovate deciduous bracts. *Involucres* of the individual flower 10 to 12, linear-lanceolate, connate at the base, as long as the calyx, hispid-tomentose like the calyx. *Calyx* with 5 deep linear teeth; the tube 10-ribbed, 1 to 1.25 in. long. *Corolla* 4 in. in diam., purple. *Fruit* pointed, hispid, as long as the persistent calyx. *Seeds* reniform, their edges densely fulvous-sericeous. Mast. in Hook. fil. Fl. Br. Ind. i. 337; Kurz For. Fl. Br. Barm. i. 126; DC. Prod. i. 455; Wall. Pl. As. Rar. i. 44, t. 51; Wall. Cat. 1903. *H. setosus*, Roxb. Fl. Ind. iii. 194. *H. vestitus*, Griff. Notul. iv. 519.

Penang, Perak. Distrib. Java, India.

4. *H. TILIACEUS*, Linn. Spec. 976. A small much branched tree; young branches minutely pubescent. *Leaves* sub-coriaceous, broadly cordate to reniform, minutely crenulate or entire, rarely lobed, acute; upper surface scaly, minutely pubescent, glabrescent or glabrous; lower densely and minutely hoary-pubescent; nerves 7 to 9 pairs, palmate; length and breadth 3.5 to 6.5 in., petioles .5 to 2 in., stipules oblong, oblique, shorter than the petiole. *Flowers* solitary; or in pedunculate, solitary, 2 to 3-flowered, axillary cymes; the peduncles 2 or 3 times as long as the petioles, with 2 obliquely oblong, opposite, pubescent, caducous bracts. *Involucres* 7 to 10, acute, united above the middle. *Sepals* 5, like the involucres but twice as long, with an elongated gland externally. *Corolla* campanulate, 4 in. in diam., yellow with crimson centre. *Fruit* as long as the calyx or shorter, ovate-acute, stellate-pubescent, spuriously 10-celled. *Seeds* few, obovate-reniform, faintly striate, sparsely scaly, pubescent, or glabrous. Mast. in Hook. fil. Fl. Br. Ind. i. 343; Kurz For. Fl. Burm. i. 126; DC. Prod. i. 454; Cav. Diss. iii. p. 151, t. 55, f. 1; Bl. Bijdr. 72; Roxb. Fl. Ind. iii. 182; Miq. Fl. Ind. Bat. i. pt. 2, 153; Beddome Fl. Sylvat. Anal. Gen. t. 4. *Paritium tiliaceum*, A. Juss. in St. Hil. Fl. Bras. Med. i. p. 156; (excl. syn. *H. elatum*) W. & A. Prodr. i. 52; Wight Ic. t. 7; Wall. Cat. 1912; Thwaites Enum. 26; Dalz. & Gibs. Bomb. Fl. 17; Griff. Notul. iv. 523. *H. tortuosus*, Roxb. Fl. Br. Ind. iii. 192; Wall. Cat. 1912 G, 1913 B.

All the provinces; near water. Distrib. The tropics generally near the coasts.

5. *H. FLOCCOSUS*, Mast. in Hook. fil. Fl. Br. Ind. i. 343. A tree 30 to 40 feet high; young branches, petioles, peduncles and outer surfaces of involucres and calyx rugulose and minutely rusty-puberulous. *Leaves* sub-coriaceous, cordate-reniform, 5-angled, acute, irregularly and

distantly sub-crenate; both surfaces minutely and sparsely stellate-pubescent, glabrescent when old, harsh; length and breadth 2 to 6 in., petiole less than half as long as the blade. *Flowers* in stout few-flowered terminal racemes longer than the leaves; peduncles stout, very rugulose, ebracteate, .75 to 1.5 in. long. *Involucres* combined into a bluntly-lobed cup much shorter than the calyx. *Sepals* oblong-lanceolate, 1.5 in. long, coriaceous, united for half their length or more. *Petals* membranous, spathulate, 4 in. long, glabrous inside, boldly striate and hispid-pubescent externally. *Staminal-tube* stellate-pubescent. *Capsule* obovoid, truncate, shorter than the persistent closely adherent calyx, densely stellate-pubescent and very rugulose, 5-valved, dehiscing only at the apex. *Seeds* numerous, obovate, sub-compressed, with shortly pilose angles, the rest of the surface scaly.

Mount Ophir, Malacca; Maingay (Kew Distrib.) 216. Perak; King's Collector 7024.

I have not been able to detect stipules on any of the specimens I have seen. They are probably fugacious.

5. THESPESIA, CORR.

Trees or shrubs. *Leaves* entire. *Inflorescence* axillary. *Bracteoles* 5-8, arising from the thickened end of the peduncle, deciduous. *Calyx* cup-shaped, truncate, minutely 5-toothed. *Corolla* convolute. *Staminal-tube* 5-toothed at the apex. *Ovary* 4-5-celled; style club-shaped, 5-furrowed, entire or 5-toothed; ovules few in each cell. *Capsule* loculicidal or scarcely dehiscing. *Seeds* tomentose; cotyledons conduplicate, black-dotted.—Natives of tropical Asia, Madagascar, and Australasia; species about 6.

T. POPULNEA, CORR. in Ann. Mus. ix. p. 290. A tree 20 to 30 feet high, young shoots scaly. *Leaves* on long petioles, sub-coriaceous, broadly cordate, acuminate, entire, glabrous above, sparsely scaly on lower surface; the base 5 to 7-nerved with a glandular pore between the nerves; length 4.5 to 6 in., breadth 3 to 4 in. petiole 2.5 in. *Flowers* 2 to 3 in. in diam., solitary, axillary, on peduncles shorter than the petioles; *petals* bright yellow with a brown spot at the base; bracteoles close to the calyx, lanceolate, often abortive. *Capsule* 1 to 1.5 in. in diam., depressed-spheroidal, scaly, becoming glabrescent; pericarp of 2 layers. *Seeds* 1 to 3 in each cell, reniform, minutely tomentose or mealy. Mast. in Hook. fil. Fl. Br. Ind. i. 345; Kurz For. Fl. Burm. i. 128; Miq. Fl. Ind. Bat. i. pt. 2, 150; Pierre Fl. For. Coch-Chine x. 173; Bl. Bijdr. 73; Cav. Diss. iii. 152, t. 56, f. 1; DC. Prodr. i. 456; W. & A. Prodr. i. 54; Wight Ic. t. 8; Thwaites Enum. 27; Beddome Fl. Sylvat. t. 63;

Dalz. & Gibs. Bomb. Fl. 18; Wall. Cat. 1888, 1, 2, & C to H. Miq. Fl. Ind. Bat. i. pt. 2, 150. *Hibiscus populneus*, L.; Roxb. Hort. Beng. 51; Flor. Ind. iii. 190. *H. populneooides*, Roxb. l.c. *Malvaviscus populneus*, Gaertn. Fruct. ii. 253, t. 135. *Azanza acuminata*, Alefeld Bot. Zeit. 1861, 299.

In all the provinces, on the sea-shore. Distrib. Tropics generally.

6. BOMBAX, Linn.

Trees. *Leaves* digitate, deciduous. *Peduncles* axillary or subterminal, solitary or clustered, 1-flowered. *Flowers* appearing before the leaves. *Bracteoles* 0. *Calyx* coriaceous, cup-shaped, truncate or lobed. *Petals* obovate or oblong. *Stamens* in 5 bundles opposite the petals: filaments numerous; anthers reniform, 1-celled. *Ovary* 5-celled, multi-ovulate; style clavate, stigmas 5. *Capsule* loculicidally 5-valved, valves coriaceous, woolly within. *Seeds* silky, the testa thin, albumen small; cotyledons contortuplicate. About 10 species, all tropical and mostly American; 1 in Africa.

1. *B. INSIGNE*, Wall. Pl. As. Rar. i. 71, t. 79, 80; Cat. 1841. A tall tree; trunk without prickles; branchlets armed or not; all parts glabrous. *Leaves* 7-9-foliolate; leaflets sub-coriaceous, obovate or oblanceolate, shortly acuminate, attenuate at the base, glaucous beneath; length 5 to 8 in., breadth 2.5 to 3 in.; petiolules .5 to .75 in.; petioles longer than the leaflets. *Flowers* 5 or 6 in. long, solitary towards the end of the leafless branches; peduncles .75 in. long, stout, clavate. *Calyx* 1.5 in. long, thickly coriaceous, urceolate-globose, obscurely and irregularly lobed, ultimately 2-cleft, sub-glabrous outside, silky inside. *Petals* fleshy, oblong, obtuse, recurved, internally glabrous, externally shortly sericeous, red to orange or yellowish. *Stamens* many; filaments fleshy, united for .5 in. above the base into 4 or 5 bundles. *Capsule* oblong, 10 in. long by 1.5 in. thick, curved, glabrous. Mast. in Hook. fil. Fl. Br. Ind. i. 349; Kurz For. Fl. Burm. i. 130; Journ. As. Soc. Beng. 1873, ii. p. 61. *B. festivum*, Wall. Cat. 1841.

Andamans. Distrib. Burmah.

The earliest name of this is *B. festivum* (1828). But at p. 89 of his Catalogue, Wallich changed this to *B. insigne*, under which name he figured and described it. It comes very near to *B. malabaricum*, DC.; but Wallich says it is a much smaller tree, and Kurz says it has many more stamens, than the former. I include it as an Andaman plant solely on the authority of the late Mr. Kurz, but I have seen no specimen collected by him or by any other person in the Andamans. And I have a strong suspicion that what Kurz regarded

as *B. insigne* is really an undescribed species which Wallich issued as 1840-2 B of his Catalogue under the name *B. malabaricum*, var. *albiflora*. His No. 3 of the same name I have not seen. A tree with leaves exactly like Wallich's 1840-4 and with unarmed trunk and branches has recently been collected in the little Coco Island by Dr. D. Prain for the Calcutta Herbarium.

2. *B. MALABARICUM*, DC. Prod. i. 479. A tree with the general characters of the last, but much larger; and with the trunk and branches prickly, the leaflets much narrower (lanceolate not obovate) and the flowers and fruit smaller. Mast. in Hook. fil. Fl. Br. Ind. i. 349; Kurz For. Fl. Burm. i. 136; Bl. Bijdr. 81; Wight III. t. 29; W. & A. Prodr. i. 61; Wall. Cat. 1840 (exclude No. 4 and possibly No. 2 B); Beddome Fl. Sylvat. t. 82. *Salmalia malabarica*, Schott Meletem, 35; Thwaites Enum. 28; Dalz. & Gibs. Bomb. Fl. 22; Miq. Fl. Ind. Bat. i. pt. 2, 166. *Bombax heptaphylla*, Cav. Diss. v. p. 296; Roxb. Hort. Beng. 50; Cor. Pl. iii. t. 247; Fl. Ind. iii. 167. *B. Ceiba*, Burm. Fl. Ind. 145, excl. syn. *Gossampinus rubra*, Ham. in Trans. Linn. Soc. xv.
— Andaman Islands; common.

7. ERIODENDRON, DC.

Trees. *Leaves* digitate, deciduous. *Flowers* appearing before the leaves, tufted at the ends of the branches, or axillary, large white or rose-coloured. *Bracteoles* 0. *Calyx* cup-shaped, truncate, or 3-5-fid. *Petals* oblong. *Staminal* bundles 5, opposite the petals, connate at the base, each bearing 2-3 sinuous or linear anthers. *Ovary* ovoid, 5-celled; style cylindrical, dilated, stigma obscurely 5-lobed. *Capsule* oblong, coriaceous or woody, 5-celled, 5-valved, valves densely silky within. *Seeds* globose or obovoid; testa crustaceous, smooth with silky hairs, albumen scanty; cotyledons contortuplicate.—About eight species—1 Asiatic and African, the others American.

1. *E. ANFRACTUOSUM*, DC. Prod. i. 479. A tall tree, the trunk prickly when young; branchlets stout, smooth, glaucous. *Leaflets* 8 or 9, lanceolate, acuminate, entire or serrulate towards the apex, the base acute; glaucous beneath; length 3 to 4 in., breadth .75 to 1 in., petiolule .25 in. broad; petioles usually longer than the leaflets. *Flowers* pedunculate, in fascicles of 3 to 8 below the apices of the branches; peduncles 1 to 2 in. long, minutely bracteate: involucre none. *Calyx* cup-shaped, with 5 rounded lobes, glabrous externally, sericeous internally. *Petals* oblanceolate, tomentose externally, glabrous within, 1 to 1.5 in. long, whitish. *Filaments* shorter than the petals. *Capsule* oblong, 3 to 5 in. long, smooth. *Seeds* numerous, sub-ovoid, black. Mast. in Hook. fil.

Fl. Br. Ind. i. 350; Bl. Bijdr. 81; W. & A. Prodr. i. 61; Wight Ic. t. 400; Griff. Not. iv. 533; Dalz. & Gibs. Bomb. Fl. 22; Miq. Fl. Ind. Bat. i. pt. 2, 166; Beddome Fl. Sylvat. Anal. Gen. t. 4. Wall. Cat. 1839. *Bombax pentandrum*, Linn. Sp. Pl. 989; Cav. Diss. v. 293, t. 151; Roxb. Fl. Ind. iii. 165. *B. orientale*, Spreng. Syst. iii. 124. *Ceiba pentandra*, Gært. Fruct. ii. 244, t. 133; Ham. in Trans. Linn. Soc. xv. 126. *Eriodendron orientale*, Steud. Nomencl. 587; Thwaites Enum. 28; Kurz For. Fl. Br. Burm. i. 131.

In all the provinces. Distrib. Malayan Archipelago, British India, West Indies. Often planted.

8. DURIO, Linn.

Trees, with entire coriaceous penni-nerved leaves, scaly beneath (except in *D. Oxleyanus*). Flowers in lateral cymes: peduncles angular. Bracts 2 or 1, connate into a cup, or distinct below, tips free, deciduous. Calyx bell-shaped, leathery, like the bracteoles densely scaly, the sepals distinct, or 5-fid, lobes valvate oblong or rounded. Petals 5, contorted-imbricate, spatulate, longer than the sepals. Staminal-tube divided into 4-5 phalanges opposite the petals; filaments many, bearing a globose head of sinuous 1-celled anthers, or (in *D. Oxleyanus*) a single annular 1-celled anther. Ovary usually scaly externally, 4-5-celled; styles connate, stigmas capitate; ovules many and 2-seriate in each cell. Fruit very large, subglobose or oblong, spiny, indehiscent or localicidally 5-valved. Seeds arillate; cotyledons fleshy, often connate. Distrib. Malay Peninsula and islands; species 3.

1. *D. ZIBETHINUS*, Linn. Syst. Nat. edit. xiii. p. 581. A tall tree; young branches thin and, like all the soft parts except the upper surfaces of the leaves, minutely scaly. Leaves elliptic-oblong, rarely obovate-oblong, shortly and abruptly acuminate, the base rounded; both surfaces shining, the upper glabrous, the lower adpressed-lepidote; main nerves 10 to 12 pairs, thin, slightly ascending; length 4.5 to 6 in., breadth 1.5 to 1.8 in., petiole .4 to .5 in. Flowers 2 in. long, 2 to 3 in. in diam., on long slender pendulous dichotomous peduncles in fascicles from the stem and larger branches, globose in bud: peduncles lepidote, 3 in. long, the bracts embracing the calyx and shorter than it. Calyx tubular, ventricose at the base, the limb with 5 or 6 short broad teeth. Petals twice as long as the calyx, spatulate. Stamens in 5 bundles united only at the very base; the filaments in each bundle united for one-fourth of their length: anthers glomerulate, reniform, compressed. Ovary elongate-ovoid, scaly; style pubescent, as long as the stamens. Fruit ovoid-globose, 8 to 12 in. long, woody, densely covered with strong

smooth pyramidal spines, 5-valved. *Seeds* few, large, with copious succulent arillus. *Mast.* in Hook. fil. Fl. Br. Ind. i. 351, and Journ. Linn. Soc. xiv. 501; Beccari Malesia, iii. 230, t. xii. f. 1 to 5, xxxvi. f. 1 to 12; Kurz For. Fl. Burm. i. 131; DC. Prod. i. 480; Bl. Bijdr. 81; Koen. in Trans. Linn. Soc. vii. 266, t. 14—16; Roxb. Fl. Ind. iii. 399. Miq. Fl. Ind. Bat. i. pt. 2, 167. Griff. Not. iv. 528; Ic. t. 596. Wall. Cat. 1842.—Rumph. Amb. i. 99, t. 29.

In all the provinces except probably the Nicobars. cultivated. *Distrib.* Malayan Archipelago.

2. *D. LOWIANUS*, Scortechini MSS. A tree 50 to 60 feet high; young branchlets and petioles and lower surface of midrib with rather large loose scales. *Leaves* narrowly elliptic-oblong, shortly acuminate; the base rounded, not attenuate; upper surface glabrous, the midrib puberulous, lower quite covered with adpressed scales, mostly minute, but a few larger and loose; main nerves 14 to 18 pairs, faint, sub-horizontal; length 4.5 to 5.5 in., breadth 1.5 to 2 in.; petiole .5 in. stout. *Cymes* crowded on small tubercles on branches several years old, trichotomous, 3 in. in diam. and about as long. *Flower-pedicels* .5 to .75 in. long, angled, covered with loose coppery scales. *Flowers* 2 in. in diam.; bracts 2 or 3, .5 in. long, broadly ovate, connate, deciduous. *Calyx* campanulate, its base sub-inflated, .75 in. long, its mouth with 3 broad blunt, shallow teeth, glabrous inside, covered with large silvery scales outside. *Petals* 5, oblanceolate, glabrous inside, pubescent outside, 1.25 in. long. *Stamens* in 5 phalanges, dividing shortly above the base into about 8 processes each dividing at its apex into several short filaments, each bearing a single reniform anther with marginal dehiscence. *Ovary* broadly ovoid, densely covered with large loose scales, 5-celled with 4 ovules in each, biseriate. *Style* cylindric, tapering, pubescent; stigma capitate. *Fruit* unknown.

Perak. Scortechini No. 1969.

A species collected only once and named by the late lamented Father Scortechini in honour of Sir Hugh Low, representative of the British Government at Perak, and to whose enlightened help Malayan Botany owes very much. The species approaches *D. Zibethinus* in many respects.

3. *D. MALACCENSIS*, Planch. MSS. *Mast.* in Hook. fil. Fl. Br. Ind. i. 351. A tree; the young branches thin, very minutely adpressed-scaly. *Leaves* elliptic-lanceolate with acute apices; the base acute, sometimes slightly rounded; main nerves about 20 pairs, thin, almost horizontal; both surfaces shining, the upper glabrous, the lower very

minutely adpressed-scaly; length 5 to 6.5 in., breadth 1.5 to 1.8 in.; petiole .5 in., scaly like the branches. *Peduncles* .5 to 1 in. long, in fascicles from tubercles on the stem, angled, bifurcating at the apex and bearing two pedicellate flowers, sometimes bearing one or two pedicels below the apex: pedicels two or three times as long as the common peduncles, angled, loosely scaly. *Flowers* 2.5 to 3 in. long. *Bracts* 2, broadly ovate, acute, embracing the buds. *Sepals* 5, ovate-oblong, blunt, valvate, 1.25 in. long, glabrous internally but with numerous very loose scales externally. *Petals* nearly twice as long as the sepals, narrowly oblong, pubescent on both surfaces, the outer with a few loose scales. *Anthers* narrowly oblong, 1-celled, sessile in groups on the apices of groups of combined filaments which are again united into 5 phalanges which, for more than half their length, form a tube round the ovary and style. *Ovary* oblong, angled, densely covered with scales with long cylindrical stalks and flat heads. *Style* shorter than the staminal tube, pubescent, slightly scaly. *Stigma* capitate. *Young fruit* globular, densely covered with subulate pubescent spines. *Ripe fruit* unknown. *Mast. in Journ. Linn. Soc.* xiv. p. 501, t. xiv. fig. 17 to 20: *Beccari Malesia*, iii. 237, t. xii. fig. 6 to 8.

Malacca; Griffith, Maingay (No. 212, Kew Distrib.) Distrib. Burmah.

This is known only from Malacca and Burmah. It is distinguished from *D. Perakensis*, which in other respects it much resembles, by the stalked scales on the ovary, and by the larger and looser scales on the leaves. Doubtless when ripe fruit of both is found, better characters will be yielded by it. Beccari's specimen No. 852, and the same distinguished botanist's Nos. 2190 and 2590 from Borneo, have been referred by Masters (*Journ. Linn. Soc. l. c.*) to this species. But Beccari (in *Malesia* iii. 238, 244) founded his species *D. affinis* on the former, and his *D. testitudinarum* on the two latter.

4. *D. TESTITUDINARUM*, Becc. *Malesia*, iii. p. 244, t. xiii and xiv. A tall tree bearing flowers only near the base of the trunk; young branches rather slender, minutely sub-adpressed scaly. *Leaves* narrowly elliptic-oblong or oblanceolate-oblong, acute or shortly acuminate, the margins (in var. 2) sometimes with a single wide shallow indentation, the base rounded; upper surface glabrous, the lower densely covered with sub-adpressed scales: main nerves 18 to 22 pairs, rather bold, subhorizontal: length 4.6 to 8.5 in. (only 2.5 to 3.5 in. in var. 1 and much longer and broader in var. 2); breadth 1.4 to 2.2 in.; petiole 6 to .25 in., thickened at apex. *Flowers* 3 to 3.5 in. long, in short condensed bracteolate racemes from tubercles near the base of the trunk;

the axes, pedicels, bracteoles and bracts densely covered with large loose scales: bracts enveloping the buds 2, broadly ovate, blunt. *Sepals* 5, valvate, wide and saccate at the base, the apices narrowed, glabrous inside, densely covered outside with loose large scales. *Petals* narrowly oblong, obtuse, more than twice as long as the sepals. *Stamens* as in *D. Malaccensis*. *Ovary* oblong, densely covered with loose, flat, sessile scales. *Style* shorter than the stamens, pubescent, sparsely scaly. *Stigma* capitate. *Fruit* (according to Beccari) on long peduncles, globose, 4 in. in diam., with 4 or 5 slight superficial grooves, densely covered with short broad, pyramidal spines. *Seeds* sub-ovate, obtuse, angled; the arillus short, thin, cup-shaped.

Perak; at low elevations, Kunstler, Wray. Distrib. Borneo.

Var. 1. *Pinangiana*, Becc. l. c. 246. Leaves narrowly lanceolate, acuminate, 2.5 to 3.5 in. long by .6 to .9 in. broad. *Flowers* smaller than in the typical form: fruit unknown.

Penang, at 2,500 feet; Curtis No. 293. This variety, of which only imperfect specimens have as yet been obtained, will probably, when full material shall be forthcoming, prove to be a distinct species.

Var. 2. *macrophylla*, King. *Leaves* 10 to 17 in. long, 2.5 to 5.5 in. broad, the edge sometimes with a single shallow indentation. *Racemes* 3 in. long, many-flowered, with numerous bracteoles.

Perak; Kunstler 7497, Wray 3397. No fruit of this variety has as yet been collected. Like the last, it may prove to be a distinct species.

5. *D. WRAYII*, King, n. sp. A large tree; young branches very slender and, like the petioles and under surface of midrib, covered with rather large adpressed pale brown scales. *Leaves* narrowly elliptic-oblong with caudate acuminate apex and rounded base; upper surface quite glabrous, lower closely covered with thin adpressed silvery scales smaller than these on the midrib; main nerves 10 to 12 pairs, sub-horizontal, faint: length 5.5 to 8.5 in., breadth 2 to 2.5 in., petiole .75 in. *Flowers* nearly 2 in. long, from the branches; pedicels of individual flowers rather more than 1 in. long, with many large loose scales. *Bracts* 3, broadly ovate, connate. *Calyx* cup-shaped, the mouth with 5 broad, rather deep, sub-acute teeth; inside glabrous, outside covered with large adpressed silvery scales as are also the bracts. *Petals* 1.5 in. long, oblanceolate, or spatulate-clawed, the claw very narrow, pubescent on both surfaces but especially on the outer. *Stamens* in 5 phalanges united at the bases only, each phalange dividing into 5 or 6 processes at the apices of which are born about 8 narrow reniform anthers dehiscing by their edges. *Ovary* broadly ovoid, loosely scaly.

Style longer than the stamens, cylindric, pubescent, not scaly: stigma capitate. *Fruit* unknown.

Upper Perak at 300 feet; Wray.

The fruit of this is unknown. Mr. Wray describes the petals as pink. The caudate-lanceolate leaves of this are different from those of any other *Durio* of the Malayan Peninsula.

6. D. OXLEYANUS, Griff. Notul. iv. 531. A tree, the young branches, petioles and under surfaces of the midrib adpressed-lepidote. *Leaves* elliptic-oblong, rounded at base and apex; upper surface glabrous; the lower softly pubescent, not scaly except on the midrib, the 15 to 18 pairs of main nerves stout, sub-horizontal, prominent beneath; length 3.5 to 5 in., breadth 1.5 to 2 in., petiole .5 in. *Flowers* about 1 in. in diam., in few-flowered scaly cymes from the smaller branches. *Involucral bracts* 2, broadly ovate, pubescent, sparsely and minutely scaly. *Calyx* cup-shaped, the mouth with 4 broad shallow rather blunt teeth, inside glabrous, outside with many large loose scales. *Petals* 4, oblanceolate or spatulate, little longer than the calyx, pubescent on both surfaces, not scaly. *Stamens* 20, shorter than the petals; 5 free and alternating with 5 phalanges of 3 each which are slightly united by the bases of their filaments: *anthers* solitary, drum-shaped, the dehiscence circular. *Ovary* depressed-globular, 4-celled, densely stellate-hairy. *Style* cylindric, pilose; stigma capitate. *Fruit* unknown. Mast. in Hook. fil. Fl. Br. Ind. i. 351 and Journ. Linn. Soc. xiv. 501, t. xvi. fig. 13 to 16. Beccari Malesia, III, 252. *Neesia Griffithii*, Planch. MSS.

Malacca, Griffith No. 545. Maingay, No. 220, (Kew Distrib.)

This differs, as Beccari has well pointed out (Malesia l. c.), from the other species of *Durio* by the absence of scales from every part of the leaf except the petiole and midrib; by the single, not glomerulate, anthers; by the hairy, not squamose, ovary. Should the fruit when found also present differences, it may be desirable to create a new genus for this species.

9. BOSCHIA, Korth.

Trees. *Leaves* oblong, entire, scaly beneath. *Flowers* small, axillary. *Bracteoles* 2-3, connate at the base, deciduous. *Calyx* deeply 4-5 parted. *Petals* linear-ligulate, entire or laciniate. *Stamens* many, some free, others irregularly coherent, outermost without anthers; anthers globose, 1-celled, opening by a terminal pore, solitary, or in groups of 2-6. *Ovary* 3-5-celled, style elongate; ovules one or more in each cell. *Fruit* oblong, 3 to 5-celled, 3 to 5-valved, muricate. *Seeds* few, ovoid, half-covered by a fleshy, coloured, cup-shaped arillus; cotyledons foliaceous. *Species* 4: all Malayan.

1. *B. GRIFFITHII*, Masters in Hook. fil. Fl. Br. Ind. i. 352. A tree 40 to 60 feet high; young branches rather slender, pale, minutely farfuraceous. *Leaves* oblong, or elliptic-oblong, or obovate-oblong, shortly and abruptly acuminate, slightly narrowed towards the rounded base; upper surface quite glabrous; the lower pale, very minutely pubescent, the midrib and nerves slightly scaly; main nerves 8 to 11 pairs, spreading, prominent beneath and dark coloured; length 5 to 6.5 in., breadth 1.5 to 2.25 in., petiole .4 to .6 in.: stipules linear, deciduous. *Flowers* .75 in. in diam., solitary, or in 2 to 3-flowered cymes from the axils of leaves or of fallen leaves; pedicels shorter than the petioles, bracteolate. *Involucral bracts* 2, broadly-ovate, blunt, connate at the base, closely enveloping the buds; scaly externally, glabrous within. *Sepals* 4, ovate, spreading, pubescent on both surfaces, scaly also on the outer. *Petals* 4 to 8, nearly twice as long as sepals, linear or linear-spathulate, 1 in. broad. *Stamens* very numerous, unequal, slightly united by the bases of the filaments: the outer without anthers, some flat resembling the petals, a few of the inner longer and bearing at their apices 1 to 4 oblong obovoid anthers which dehisce by an apical pore. *Ovary* ovoid, 3-celled, densely covered with peltate, fimbriate, long-stalked scales. *Style* as long as the longest stamens. *Stigma* subcapitate. *Fruit* oblong, pointed at each end, 1.5 to 2 in. long, densely covered with sharp stout conical spines, 3-celled, dehiscent. *Seeds* 3 to 6, or fewer. *Mast.* in Journ. Linn. Soc. xiv. t. xv, fig. 29 to 39, t. xvi., fig. 40 to 42. *Beccari Malesia* III, p. 256. *Heteropyxis*, Griff. Not. iv. 524; *Ic. Pl. As.* t. 594.

Malacca; Griffith, Maingay. Perak, very common. *Distrib.* Sumatra, Forbes, No. 3068.

10. *NEESIA*, Blume.

Trees. *Branches* marked with large leaf-scars. *Leaves* entire, pinnate-veined. *Stipules* leafy. *Cymes* from the stem in the axils of the fallen leaves. *Bracteoles* 3, connate into a cup, deciduous, covered, like the sepals, with peltate scales. *Calyx* ventricose, conical above, opening by a circular irregularly crenulate orifice at the top, ultimately dilated and cushion-shaped at the base. *Petals* 5, free, imbricate. *Stamens* numerous, the filaments more or less united; anthers 2-celled, opening lengthwise, connective thick; staminodes 0. *Ovary* oblong, 5-celled; style short; stigma capitate; ovules numerous, 2-seriate, horizontal, anatropous. *Fruit* ovoid, woody, muricate, loculicidally 5-valved. *Seeds* albuminous; aril 0; cotyledons flat, leafy. *Distrib.* Seven species, all Malayan.

N. SYNANDRA, Mast. in Hook. fil. Fl. Br. Ind. i. 352. A tree 70 to 100 feet high; young branches stout, their bark dark lenticellate and with large cicatrices. *Leaves* large, crowded near the apices of the branches, coriaceous, oblong-elliptic to obovate-elliptic: the apex rounded, emarginate; the edges sub-undulate, slightly narrowed in the lower third to the sub-cordate base; upper surface glabrous, lower puberulous; nerves 13 to 22 pairs, spreading, stout and distinct on both surfaces, the reticulations also distinct; length 7 to 16 in., breadth 3.5 to 8 in.; petiole 1.5 to 3 in., thickened at base and apex; stipules foliaceous, with very stout midribs, 1.5 to 2.5 in. long. *Cymes* short (1.5 in. long), crowded, dichotomous, 8 to 12-flowered, from the axils of sub-apical fallen leaves; the pedicels short, scaly. *Flowers* about 6 in. long. *Bracts* connate into a 3-lobed cup surrounding the base of the flowers. *Calyx* ventricose with a contracted irregularly and minutely toothed mouth, densely pubescent inside, scaly outside as are the bracts, ultimately involute so as to form an annular cushion .5 in. or more in diam. *Petals* 5, free, much imbricate, ovate-lanceolate, glabrous. *Stamens* numerous, the filaments more or less connate at the base, unequal; anthers sub-globular, 2-celled. *Ovary* conical, sessile, densely pilose, not scaly: style slightly longer than the ovary; stigma capitate, 5-angled. *Fruit* 6 to 8 in. long and 4 to 5 in. in diam., ovoid-conic, pedunculate, with 5 bold rounded vertical angles: the pericarp very thick, woody, externally covered with stout pyramidal sharp spines, internally lined with a dense layer of stiff yellow hair; 5-celled, deliscent. Mast. in Journ. Linn. Soc. xiv. p. 504. *Beccari Malesia*, iii. 263.

Malacca, Maingay. Perak; Scortechini, Wray, King's Collector.

I have seen no specimens of the plant (*N. altissima*) on which Blume founded this genus. But, judging from his admirable description and fine coloured figure (Nov. Act. Acad. Caes. xvii. 83, t. vi), this species must be very closely allied to that. I find the stamens of this agree both with Blume's description above referred to, and with Sig. Beccari's, in his admirable and splendidly illustrated monograph in *Malesia* iii. pp. 258 to 268. Ripe fruit and seeds of this are as yet unknown.

11. CÆLOSTEGIA, Benth.

Tall trees. *Leaves* simple, entire, scaly beneath. *Flowers* small (scarcely .25 in. in diam.), cymose; the inflorescence, bracts and calyx scaly. *Bracts* connate into a toothed cup. *Calyx* with constricted tube, pouched above and constricted at the apex into 5 connivent lobes. *Petals* 5, free, inserted near the apex of the calyx tube, connivent. *Stamens* numerous; the filaments short, thick, slightly connate at the base, the apex constricted; the anthers globose, 3 to 4-celled. *Ovary*

partly immersed in the calyx-tube, globular or sub-globular, 5-celled; the ovules few, erect. *Style* short; stigma peltate, discoid, large. *Fruit* large, woody, muricate externally, hairy within, 5-celled, few-seeded, deliscent. Three species; all Malayan.

C. GRIFFITHII, Benth. in Benth. & Hook. fil. Gen. Plant. i. 213. A tree; the young branches rather slender, dark-coloured, striate, minutely and deciduously scaly. *Leaves* coriaceous, oval, shortly and bluntly acuminate, the base rounded; upper surface glabrous, lower sparsely adpressed-scaly; main nerves about 8 pairs, spreading, faint; length 2.4 to 3.75 in., breadth 1.25 to 1.6 in.; petiole .5 to .75 in., minutely adpressed-scaly. *Inflorescence* of fasciculate cymose racemes about 2 in. long, from the axils of fallen leaves, many-flowered; pedicels longer than the flowers. *Flowers* .25 in. in diam., scaly. *Bracts* connate into a 3-lobed cup less than half as long as the calyx. *Calyx* constricted at the base, then dilated into a 5-pouched sac which is contracted and 5-toothed at its apex. *Petals* 5, distinct, inserted on the calyx at the apex of its tube, triangular, acute, connivent, fleshy, glabrous. *Stamens* numerous, attached to the petals; the anthers small, globose, 3 or 4-celled. *Ovary* globular-obovate, densely covered with large loose scales. *Style* shorter than the ovary; stigma peltate, thick, its edges wavy. *Fruit* unknown. Mast. in Hook. fil. Fl. Br. Ind. i. 353 and Journ. Linn. Soc. xiv. 505, t. xvi, figs. 43 to 50. *Beccari Malesia*, iii. 270.

Malacca, Griffith; Perak, Scortechini, King's Collector.

Fruit was not known when this genus was first established by the late Mr. Bentham; and, of this species, fruit is still unknown. Sig. Beccari has, however, discovered two species in Sumatra and Borneo (*C. Sumatrana* and *Bornensis*) the fruit of which he describes and figures (*Malesia*, iii. 271, t. xxvii. to xxix); and from his description the generic description has been completed.

Order XVIII. STERCULIACEÆ.

Herbs, shrubs or trees; herbaceous portions usually more or less stellate-pubescent. *Bark* usually abounding in mucilage, inner fibrous. *Leaves* alternate, simple, often lobed, stipulate. *Inflorescence* axillary, rarely terminal, usually cymose. *Flowers* regular, uni- or bi-sexual. *Sepals* 5, often connate. *Petals* 5 or 0. *Andræcium* columnar or tubular, of many stamens; or stamens rarely few, free; anthers in heads, or in a single ring at the apex of the column, or dispersed on the outside of the tube, or arranged along the edge of a cup or tube, with intervening staminodes or sterile stamens; anther-cells always 2,

parallel or divergent. Ovaries 2 to 5, free, rarely 1, sessile or stalked; styles slightly united and becoming free or slightly coherent, as many as the ovaries. Ovules many or few, attached to the inner angles of the ovaries, anatropous, ascending or horizontal, raphe ventral or lateral. Fruit dry or fleshy, dehiscent or indehiscent. Seeds sometimes arillate, albuminous or exalbuminous: cotyledons leafy, flat, folded or convolute; radicle short, inferior, pointing towards, or remote from the hilum. Distrib. Abundant in the tropics of either hemisphere and in subtropical Africa and Australia. Genera 40—45; species from 500 to 600.

Tribe I. *Sterculiææ*. Flowers unisexual or polygamous. Petals 0. Andræcium columnar; the anthers clustered at its apex; or in a 1-seriate ring.

Anthers numerous.

Ovary with 2 or more ovules in each cell; fruit dehiscent

1. *Sterculia*.

Ovarian cells 1-ovuled; fruit indehiscent

2. *Tarrietia*.

Anthers 5, whorled; fruit indehiscent.

3. *Heritiera*.

Tribe II. *Helicterææ*. Flowers hermaphrodite. Petals deciduous. Andræcium columnar below, dilated above into a cup, margin bearing on it the anthers usually alternating with staminodes.

Capsule membranous, inflated... ..

4. *Kleinhovia*.

Capsule more or less woody, not inflated.

Anther-cells divaricate; seeds not winged

5. *Helicteres*.

Anther-cells parallel; seeds winged

6. *Pterospermum*.

Tribe III. *Hermannieææ*. Flowers hermaphrodite. Petals marcescent, flat. Andræcium tubular at the base only; stamens 5, staminodes 0.

Ovary 5-celled

7. *Melochia*.

Ovary 1-celled, 1-seeded

8. *Waltheria*.

Tribe IV. *Buettnerieææ*. Petals concave or unguiculate at the base; filaments in a tube with the anthers at its apex, solitary or in groups between staminodes.

Stamens in a single series.

Stamens in groups between the staminodes;

Petals unguiculate

9. *Abroma*.

- Stamens solitary between the staminodes.
- Petals unguiculate, with 2 lateral lobes and a long subterminal appendage 10. *Buettneria*.
- Petals linear not lobed, concave not unguiculate at the base 11. *Commersonia*.
- Stamens in several series 12. *Leptonychia*.

1. STERCULIA, Linn.

Trees or shrubs. *Leaves* simple, entire or palmately lobed, sometimes digitately compound. *Inflorescence* paniced or racemose, usually axillary and crowded towards the apices of the branches. *Flowers* male and hermaphrodite. *Calyx* campanulate or rotate, 4-5 lobed, often coloured. *Petals* 0. *Staminal column* bearing a head or ring of usually sessile, 2-celled, anthers at its apex, the cells often divergent. *Carpels* 5, distinct or slightly cohering, 2 to many-ovuled, borne on the apex of a more or less elongated gynophore; styles more or less connate: stigmas free or united so as to form a peltate lobed disc., *Ripe carpels* distinct, spreading, sessile or stalked, follicular, from membranous to woody, with several (rarely many) seeds; or navicular with a single seed. *Seeds* 1 to many, sometimes winged, rarely arillate; albumen bipartite, flat or lobed: cotyledons thin flat and adherent to the albumen, or fleshy; radicle near to or remote from the hilum. *Distrib.* About 70 species tropical and chiefly Asiatic.

Sect. I. *Eusterculia*, Endl. *Follicle* coriaceous or woody. *Seeds* two or more.

Leaves simple, orbicular or reniform.

Leaves lobed.

Follicles glabrous within, the edges only ciliate; gynophore and staminal tube glabrous 1. *S. villosa*.

Follicles hispid-pilose within; gynophore and staminal tube hairy 2. *S. ornata*.

Leaves not lobed 3. *S. macrophylla*.

Leaves simple, longer than broad; not orbicular or reniform.

Leaves quite glabrous.

Calyx-lobes not cohering by their apices 4. *S. laevis*.

Calyx-lobes cohering by their apices.

Flowers in racemes: nerves of leaves 6 pairs or fewer 5. *S. hyposticta*.

- Flowers in panicles: nerves of leaves more than 6 pairs
 Leaves narrowly oblong-lanceolate; follicles 1 to 1.25 in. long 6. *S. parvifolia*.
 Leaves ovate or obovate-oblong to narrowly elliptic.
 Ovaries 3, villous: stamens 7 7. *S. Kunstleri*.
 Ovaries 5, scaly; stamens 10 8. *S. parvifolia*.
 Leaves more or less hairy.
 Calyx-lobes not cohering by their apices: leaves glandular-dotted beneath 9. *S. Scortechinii*.
 Calyx-lobes slightly cohering by their apices: leaves white beneath ... 10. *S. bicolor*.
 Calyx-lobes spreading, connivent and cohering by their apices.
 Stigmas free, long, recurved ... 11. *S. angustifolia*.
 Stigmas united into a lobed disc.
 Leaves more or less obovate 12. *S. rubiginosa*.
 „ lanceolate 13. *S. ensifolia*.
 Species of uncertain position... .. 14. *S. pubescens*.
- Sect. II. *Firmiana*, Marsili; Br. in Benn. Pl. Jav. Rar. 235 (gen.). *Follicles* membranous, opening long before maturity. *Seeds* two or more.
 Calyx .75 in. long: staminal tube about the same length; adult leaves glabrous... 15. *S. colorata*.
 Calyx 1.25 in. long, staminal tube .5 in. longer: adult leaves minutely stellate-pubescent 16. *S. fulgens*.
- Sect. III. *Pterygota*, Endl. (gen.). *Follicles* woody. *Seeds* many, winged at the apex ... 17. *S. alata*.
- Sect. IV. *Scaphium*, Endl. *Anthems* 15, (sometimes 10). *Stigmas* lobed. *Follicles* large, membranous, boat-shaped, often gibbous, opening long before maturity, containing only 1 seed near the base.
 Leaves ovate-rotund, deeply cordate ... 18. *S. linearicarpa*.
 Leaves ovate to ovate-oblong: main nerves 2 to 4 pairs 19. *S. scaphigera*.

Leaves elliptic-oblong: main nerves 6 to
7 pairs 20. *S. afinis*.

Sect. V. *Pterocymbium*, Br. in Benn. Pl. Jav. Rar.
219 (gen.). *Flowers* sub-hermaphrodite.
Anthers 10. *Styles* coherent, stigmas re-
curved. *Follicles* 4—6, membranous, open-
ing long before maturity. *Seed* solitary.

Leaves broadly ovate, acuminate, the
base deeply cordate 21. *S. campanulata*.

Leaves elliptic-oblong; the base broadly
rounded or sub-truncate, not cordate... 22. *S. tubulata*.

1. *S. VILLOSA*, Roxb. Hort. Beng. 50. A tree 30 to 60 feet high: young branches thick, their apices tawny-tomentose and enveloped by the large sub-caducous stipules, the bark pale with large leaf-cicatrices. *Leaves* thickly membranous, rotund or reniform, with 5 to 7 broad abruptly acuminate often toothed lobes, the sinuses between the lobes acute; the base deeply cordate, the basal lobes rounded: upper surface at first minutely stellate-pubescent, ultimately glabrous, except the 5 to 7 radiating tomentose nerves: under surface uniformly and minutely tomentose; length and breadth from 12 to 18 inches: petiole ovate-lanceolate, acuminate, with cordate bases, pubescent, sub-caducous. *Panicles* from the axils of the previous year's leaves, solitary, from 6 to 12 in. long: branches short, many-flowered, tomentose. *Calyx* campanulate, 4 in. in diam., with 5 ovate acute spreading lobes as long as the tube, yellowish with purple fundus, veined, puberulous outside especially towards the base, almost glabrous inside. *Male flower*; staminal column longer than the calyx-tube, slightly curved, quite glabrous, bearing at its apex 10 sub-sessile anthers with thick connective and 2 divergent cells. *Female flower*; gynophore glabrous, thickened above; ovaries 5, conjoined, tomentose; styles conjoined, puberulous, curved; stigma small, lobed. *Follicles* 3 to 5, coriaceous, sessile, bright red when ripe, oblong, tapering to both ends; 2 to 2.5 in. long by 1 in. broad: shortly hispid-pubescent externally, smooth and shining internally and glabrous except along the placental edges which are strongly ciliate. *Seeds* 6 or more, oval, smooth. Roxb. Fl. Ind. i. 153; Kurz For. Fl. Burm. i. 136; Mast. in Hook. fil. Fl. Br. Ind. i. 355; Pierre Fl. Forest. Coch-Chine, t. 185, fig. D.; Wall. Cat. 1136, 2, 3, D.; W. & A. Prodr. i. 63; Dalz. & Gibs. Bomb. Fl. 22; Br. in Benn. Pl. Jav. Rar. 227.

Andamans, Prain. Distrib. British India.

2. *S. ORNATA*, Wall. in Herb. Calcutta. A tree 20 to 30 feet high : young branches thick, glabrous, pale, the leaf-cicatrices very large, the apices deciduously pilose, coccineous drying into brown. *Leaves* thickly membranous, reniform, more or less deeply divided into 5 or 7 acuminate lobes, the sinuses between the lobes wide, the base deeply cordate; upper surface minutely strigose, often stellate, minutely pitted; lower surface yellowish-brown, minutely and uniformly tawny-tomentose, minutely glandular-dotted under the hair; the 5 to 7 radiating main nerves and the ascending secondary nerves bold and distinct; length about 12 in., breadth about 15 in.; petiole 15 to 18 in. long, thickened at the base, minutely tomentose. *Panicles* from the axils of the previous year's leaves, solitary, 8 to 15 in. long, shortly branched, many-flowered, pulverulent reddish-tomentose. *Calyx* ochre-coloured with red fundus, veined, widely campanulate, sub-rotate, with 5 ovate acute spreading lobes longer than the tube, stellate-pubescent externally, puberulous internally; .75 in. in diam. *Male flower*; gynophore about as long as the tube, curved, sparsely glandular-hairy, bearing at its apex 10 small anthers with thick connective. *Female flower*; gynophore thickened above, densely tawny-tomentose as are the conjoined ovaries and curved style; the ovaries with a ring of about 10 sessile anthers at their base; stigma discoid, rugulose, 5-lobed. *Follicles* about 5, sessile, coriaceous, narrowly oblong, very shortly beaked, brilliant orange scarlet when ripe, outside glabrescent, inside densely coccineous-pilose; length 4 in., breadth 1.25 in. *Seeds* about 6, oval, smooth. Wall. in Voigt Hort. Calc. Suburb. 105 (*name only*); Kurz Journ. As. Soc. Beng. Vol. xlii. pt. 2, p. 258; Vol. xliii. pt. 2, p. 116; For. Fl. Burm. i, 136. *Sterculia armata*, Mast. in Hook. fil. Fl. Br. Ind. i. 357, *in part*. Pierre Fl. Forest. Coch-Chine, t. 185, fig. C.

Burmah; Wallich, Brandis, Kurz. Andamans, Kurz.

I include this species because, although the evidence of its having been collected in the Andamans is not very good, I think it extremely likely that it does occur there, and that good unmistakable specimens will soon be forthcoming. The species in many respects resembles *S. villosa*, with which it appears to have often been confused. The distinctive marks to separate it from *S. villosa* are that the leaves are minutely dotted and pitted; that the apices of the young branches have red hairs (becoming brown on drying); that after the hairs have fallen the young branches have pale polished bark with very large leaf-cicatrices and some warts, but no sub-persistent stipules; that the flowers are larger (.75 in. in diam. as against .4 in.); that the staminal column and gynophore are hairy; that the follicles are larger and paler; and that the whole of their inner surface is densely hispid-pilose.

3. *S. MACROPHYLLA*, Vent. Hort. Malm. ii. No. 91 (in note). A tree 80 to 120 feet high; young branches very thick, rough from the leaf cicatrices, the apices deciduously rufous or tawny-pilose. *Leaves* sub-coriaceous, broadly ovate to ovate-rotund or obovate-rotund, entire, narrowing to the slightly cordate 7-nerved base; upper surface sparsely and rather minutely pubescent, some of the hairs 2-branched, becoming glabrescent with age, the midrib and nerves always pubescent; under surface sub-tomentose, tawny, the midrib and 6 to 8 pairs of lateral nerves prominent, rufous-villose; transverse venation distinct, rather straight; length 8 to 16 in., breadth 6 to 12 in., petiole 3.6 to 6 in., softly hairy, tawny. *Panicles* solitary, axillary, nearly as long as the leaves, much-branched, many-flowered, hispidulous-pubescent, capillary, shorter than the flowers. *Flower-buds* minute, sub-globose. *Calyx* .15 in. long, campanulate, stellate-hairy, 5-lobed; the lobes triangular, erect, shorter than the tube. *Follicles* 3 to 5, shortly stalked, woody, sub-rotund, about 2.25 in. each way, crimson when ripe, outside pubescent and longitudinally rugose; inside smooth. *Seeds* oblong, black, smooth, .75 in. long. Mast. in Hook. fil. Fl. Brit. Ind. i. 356; R. Brown in Benn. Pl. Jav. Rar. 230.

Malacca; Maingay No. 233 (Kew Dist.). Perak; at elevations of 200 to 500 feet; King's Collector Nos. 6052 and 7923; Scortechini, No. 230. Distrib. Java, Brit. North Borneo.

4. *S. LAEVIS*, Wall. Cat. 1138. A shrub or small tree; young branches rather thin, with pale striate bark, the apices deciduously rusty-puberulous. *Leaves* membranous, narrowly ovate-oblong, sometimes slightly obovate, the apex shortly and bluntly acuminate; the base tapering, acute, rarely rounded, faintly 3-nerved: both surfaces glabrous, shining, the midrib and 6 to 9 pairs of spreading nerves prominent on the lower: length 4.5 to 9 in., breadth 2 to 3 in.; petiole 1.1 to 2.5 in., smooth, thickened at the apex. *Panicles* meagre, solitary, axillary, slender, puberulous, shorter than the leaves, few-flowered; pedicels about as long as the flowers. *Flower-buds* oblong. *Calyx* .5 in. long or more, pubescent on both surfaces but especially on the inner; the tube urceolate, divided at its apex into 5 linear-oblong sub-acute ascending lobes, longer than the tube, slightly connivent but not cohering by their apices, hispidulous on their inner surface. *Male flower*; staminal column shorter than the tube, glabrous; anthers 10, sessile at its apex, elongate-ovate. *Hermaph. flower*: gynophore very short; ovaries 5, boat-shaped, rusty-pubescent, sub-sessile, with a ring of 10 sessile anthers at their base outside: styles almost obsolete; stigmas 5, cylindric, free, radiating, recurved, pubescent beneath. *Follicles* 3 to 5, coriaceous, narrowly

oblong, with short straight beaks, bright red when ripe, puberulous externally, slightly curved, glabrous, shining and ridged internally, 2 in. long and about .5 in. broad. *Seeds* 3 or 4 oblong, black, shining. Mast. in Hook. fil. Fl. Br. Ind. i. 357. Pierre Fl. Forest. Coch-Chine t. 192, figs. 1 to 7; Br. in Benn. Pl. Jav. Rar. 230; Miq. Fl. Ind. Bat. i. pt. 2, 174. *S. coccinea*, Jack Mal. Misc. i. 286, not of Roxb.

Penang, Perak, Malacca, Singapore: at low elevations: but not common.

5. *S. HYPOSTICTA*, Miq. Fl. Ind. Bat. Suppl., 399. A shrub or small tree, all parts glabrous except the inflorescence: young branches slender, dark and smooth becoming (by the falling off of the bark) pale and striate. *Leaves* membranous, oblong, to oblong-lanceolate, sometimes slightly obovate, abruptly acuminate or even caudate-acuminate, entire, the base slightly narrowed and rounded, or not narrowed and truncate, emarginate, rarely acute, 3-nerved; both surfaces glabrous, shining: lateral main nerves 3 to 5 pairs, spreading, curved, inarching far from the margin, prominent beneath: length 3.5 to 5.5 in., breadth 1.5 to 2.25 in., petiole 1 to 1.5 in., thickened at base and apex. *Racemes* axillary, solitary, drooping, longer than the leaves, minutely whitish pubescent, with superficial brown stellate hairs: bracteoles linear, longer than the pedicels. *Calyx* with narrowly campanulate tube .25 in. long, densely rufous-pubescent externally and glabrous inside: lobes 5, not quite so long as the tube, linear, spreading, connivent, cohering from some time by their tips, the edges recurved, glandular-pilose inside, sub-pubescent outside. *Male flower*: staminal column short, glabrous, with 8 sessile oblong 2-celled anthers at its apex. *Female flower*: gynophore short; ovaries 4, ovoid, conjoined, shortly tomentose, with ring of 8 sessile anthers at their base. *Style* simple, curved, sparsely villous; stigma large, glabrous, with 4 fleshy oblong-obovoid curved lobes. *Follicles* 2 or 3, coriaceous, bright red when ripe, narrowly oblong, tapering to each end, 2 to 2.25 in. long and .65 in. broad; externally minutely rusty-pubescent; internally glabrous, wrinkled. *Seeds* 4, oblong, pointed, black. Kurz in Journ. As. Soc. Beng. Vol. xlv. pt. 2, p. 120.

Perak; King's Collector, Wray. Nicobars, Kurz.

6. *S. PARVIFOLIA*, Wall. Cat. 1123. A tree 20 to 30 feet high: young branches slender, striate, the older pale, the younger dark-coloured, glabrous. *Leaves* membranous, drying of a pale green, oblong-lanceolate, rarely ovate-oblong, bluntly acuminate, entire; the base acute or rounded, faintly 3-nerved; both surfaces glabrous: main nerves 6 to 8 pairs,

spreading, rather prominent on both surfaces as is the midrib: length 4 to 6.5 in., breadth 1 to 1.75 in.; petiole 1 to 1.75 in., smooth, slender, thickened at the apex. *Racemes* solitary, axillary, much shorter than the leaves, few-flowered, glabrous; flower-pedicels shorter than the flowers, capillary. *Flower-buds* oblong. *Calyx* less than .5 in. long, glabrescent externally, puberulous internally especially on the lobes; tube wide, cylindric, with 5 linear-lanceolate lobes about as long as itself, spreading, incurving and joined for some time by their tips. *Male flower*: staminal column shorter than the tube and bearing at its apex about 12 small oblong anthers with thick connective and diverging cells. *Herm. flower*: gynophore very short, glabrous; ovaries 5, broadly ovate, rusty-pubescent; styles united, recurved, with many white spreading hairs; stigmas clavate, flattened, recurved, spreading. *Follicles* 3 to 5, broadly oblong, with a straight beak, 1 to 1.25 in long, .6 in. broad. *Seeds* 2, broadly ovoid, black, shining. Mast. in Hook. fil. Fl. Br. Ind. i. 356; R. Brown in Benn. Pl. Javan. Rar. 229; Miq. Fl. Ind. Bat. Vol. i. pt. 2, p. 173.

Penang, Perak, Malacca.

Closely allied to *S. laevis*, Wall.: but with smaller flowers and follicles, and with calyx lobes coherent at their tips.

7. *S. KUNSTLERI*, King, n. sp. A tree 30 to 60 feet high; all parts (except the inflorescence and the tips of the young branches) glabrous; branches with pale smooth striate bark. *Leaves* thinly coriaceous, broadly ovate (or slightly obovate) to oblong or narrowly elliptic, the apex rounded, blunt, sub-acute or very shortly and sub-abruptly acuminate; slightly narrowed to the rounded or sub-truncate, rarely acute, 3 to 5-nerved, base; both surfaces shining; lateral nerves about 7 to 9 pairs, spreading, slightly prominent beneath: length 4 to 9 in., breadth 2 to 4.5 in.; petiole .75 to 2.75 in., slender, glabrous. *Panicles* solitary, narrow, in the axils of (and shorter than) the mature leaves, or supra-axillary, slender; the lateral branches short, 1-to 3-flowered, flocculent-tomentose, rusty; bracteoles lanceolate to ovate, caducous. *Calyx* .3 to .35 in. long, the tube urceolate, densely stellate-tomentose outside, sub-glabrescent inside; lobes 5, shorter than the tube, linear-lanceolate, villous on the inner surface, tomentose on the outer, spreading, connivent and slightly coherent by their tips. *Male flower*: staminal column slender, shorter than the calyx-tube, curved, bearing at its apex 5 to 7 sessile broad anthers. *Hermaph. flower*: gynophore short; ovaries 3, ovoid, villous, with a ring of adpressed sessile oblong anthers at their base: styles distinct, short, thin, sparsely villous; stigmas thick, fleshy, clavate, bent (outwards) on themselves. *Follicles* 2 or 3,

woody, from peach-coloured to carmine when ripe, oblong, rounded at at the base, the apex acute and slightly curved; externally rugose (the rugae mostly longitudinal), minutely tomentose, inside smooth; length 3 to 3.5 in., breadth 1.25 to 1.5 in. *Seeds* narrowly ovoid, nearly 1 in long, black.

Perak; King's Collector Nos. 3259, 7211, 7245, Scortechini No. 1805; at 100 to 300 feet elevation. Distrib. Sumatra; Forbes, No. 2679.

In externals this species closely resembles *S. parviflora*, Roxb. But, after numerous dissections, I conclude that the two species are quite distinct. The ovaries of this are never more than 3, and they are always densely villous; those of *parviflora* are invariably 5, and they are scaly, not villous. The stigmas of this are long and are bent outwards on themselves; these of *parviflora* are short and recurved outwards from their junction with the styles: they are not bent on themselves. The follicles of this are thicker and more woody and the seeds are larger than those of *S. parviflora*. Moreover this has never more than 7 stamens, while *S. parviflora* has 10. The leaves of this are rather thicker in texture and the young branches are thinner and paler than those of *S. parviflora*.

8. *S. PARVIFLORA*, Roxb. Hort. Beng. 50. A tree 20 to 50 feet high; young branches rather thick; the tips ferruginous-tomentose; the bark pale, rough, glabrous. *Leaves* membranous, oval, ovate or obovate-oblong, the apex rather abruptly shortly and bluntly acuminate, entire; the base rounded and slightly cordate, or sub-truncate and emarginate, 5-nerved; both surfaces glabrous, but not shining; the midrib and 7 or 8 pairs of spreading rather prominent lateral nerves sparsely stellate-pubescent on the lower when young; length 4 to 10 in., breadth 2 to 5.5 in.; petiole 1 to 4 in., deciduously rufous-tomentose. *Panicles* about as long as the leaves, slender, the lateral branches short and the flower-pedicels capillary, everywhere covered with rusty stellate tomentum, ebracteolate. *Calyx* 2 in. long with an urceolate tube, the mouth with 5 linear-lanceolate lobes almost as long as the tube, incurved and united by their apices, stellate-tomentose externally, glabrous within. *Male flower*: staminal column shorter than the calyx-tube, bearing at its apex 10 sessile short narrowly ovate anthers with thick connective. *Herm. flower*: ovaries 5, ovoid, scaly, with a ring of anthers at their base: ovules 4 or 5. *Styles* slightly united, slender, sparsely villous, short; stigmas united into a fleshy boldly 5-lobed disc, but easily separable into 5 fleshy flattish recurved stigmas. *Follicles* 1 to 5, thickly coriaceous, brilliant red to orange, pubescent to glabrescent, oblong, shortly beaked, 2.5 to 3.5 in. long and 1.25 to 1.5 in. broad; inside glabrous, shining,

boldly ridged. *Seeds* broadly ovoid, black, .6 in. long, smooth. Roxb. Fl. Ind. iii. 147; Brown in Bennett Pl. Jav. Rar. 232; Wall. Cat. 1121. Kurz For. Fl. Burm. i. 138. Pierre Fl. Forest. Coch-Chine, t. 195 F. *S. Maingayi*, Mast. in Hook. fil. Fl. Br. Ind. i. 359; Pierre Fl. Forest. Coch-Chine, t. 188 A.

Penang, Malacca, Perak; at low elevations, common. Distrib. Burmah and Sylhet in British India; Cochin China.

After careful dissection of the flowers of the types of the two species *S. parviflora*, Roxb. and *S. Maingayi*, Masters, and of flowers of many other specimens, I can come to no other conclusion than that they are one and the same. There is a curious tendency to inequality in size in the leaves, some being twice as large as others rising from the same twig within the distance of an inch. And the panicles usually follow the leaves in the matter of length.

9. *S. SCORTECHINII*, King, n. sp. A tall tree; young branches rather thick, their bark pale, rough, the youngest parts deciduously rusty-pubescent. *Leaves* thinly coriaceous, oblong, slightly obovate, the apex rounded, with an abrupt short blunt point, entire; the base slightly narrowed, rounded or minutely cordate, 3-nerved; upper surface glabrous, shining; the lower slightly paler, dull, thickly dotted with minute reddish flat shining glands, the midrib and 4 to 5 pairs of prominent ascending lateral nerves stellate-pubescent: length 2.5 to 3.5 in., breadth 1.25 to 1.65 in.; petiole .65 to 1 in., deciduously pulverulent-tomentose. *Panicles* racemes-like, axillary, solitary, shorter than the leaves, densely pulverulent-tomentose, rusty; pedicels as long as the buds; bracteoles ovate, .25 in. long, imbricate, caducous. *Calyx* campanulate, divided almost to its base into 5 broadly ovate spreading not connivent lobes, pubescent-tomentose both internally and externally. *Male flower*: staminal column shorter than the calyx, crowned by about 10 short anthers with thick sub-cuneate connective and short divergent cells. *Herm. flower*: *Ovary* 3-celled, obliquely ovoid, pubescent-scaly; ovules 3 or 4 in. each cell. *Styles* connate, pubescent. *Stigmas* 3, large, ovoid, spreading, glabrous, dark-coloured. *Follicles* not seen.

Perak; Scortechinii, No. 2068.

Collected only once, and without fruit.

10. *S. BICOLOR*, Mast. in Hook. fil. Fl. Br. Ind. i. 359. A tree 40 to 60 feet high; young branches rather thin, cinereous, striate, glabrous, rufous-pubescent at the very tips. *Leaves* small, membranous, obovate-oblong, acute or shortly mucronate, entire, slightly narrowed to the minutely 2 to 3-nerved rounded base; upper surface glabrous

when adult, with a few small scattered white stellate hairs when young; under surface pale from a layer of minute whitish hairs, the midrib and 16 to 18 pairs of sub-horizontal lateral nerves rufous-tomentose; length 2.5 to 3 in., breadth 1.2 to 1.4; petiole about 1 in., slender, scaly-tomentose. *Panicles* about as long as the leaves, slender, in the axils of young leaves, pulverulent-tomentose, sub-ferruginous; branches short, spreading. *Calyx* pedicellate, ovoid-oblong, pointed in bud, when adult .3 in. long, widely campanulate, with 5 linear incurved pubescent lobes as long as the tube. *Staminal column* shorter than the tube, glabrous; anthers about 12, sessile at the apex of the column, their connective thick, cuneate, the cells divergent. *Follicles* unknown.

Malacca; Maingay, No. 230 (Kew Distrib.) Perak. Wray, No. 2378.

Recognisable at once by its small leaves, white beneath. The figure named *S. bicolor*, Mast. by Pierre (Fl. Forest. Coch-Chine t. 187) agrees neither with M. Pierre's own description of it; nor with Masters' type-specimen. There may probably have been some printer's blunder in the matter.

11. *S. ANGUSTIFOLIA*, Roxb. Hort. Beng. 50. A small tree: young branches densely velvety rusty-tomentose; ultimately rather pale, glabrous, warty and striate. *Leaves* membranous, oblong-lanceolate rarely ovate-lanceolate, acuminate or acute, entire, slightly narrowed to the rounded 3-nerved base: upper surface glabrous, the lower more or less densely and softly rusty-tomentose: length 4 to 7 in., breadth 1.25 to 2.25; petiole .6 to 1.1 in., rusty-tomentose. *Panicles* solitary, axillary, crowded at the apices of the branches, lax, drooping, longer than the leaves, everywhere densely rusty-tomentose; pedicels much longer than the ovate pointed buds. *Calyx* .2 in. in diam., hispidulous-pubescent everywhere except the tube which inside is glabrous, deeply divided into 5 linear-lanceolate lobes; the lobes longer than the tube, spreading, connivent, cohering by their tips, their edges recurved. *Male flower*: staminal column as long as the tube, glabrous, recurved, bearing at its apex 10 oblong sessile anthers with large connective, the cells slightly divergent. *Herm. flower*: gynophore short, glabrous: ovaries 5, ovoid, rusty-tomentose with a ring of 10 sessile anthers at their base: style short, sparsely pilose: stigmas much longer than style, fleshy, spreading, recurved. *Follicles* 4 or 5, ovate-oblong with a short curved beak, 2.75 in. long and 1.35 in. broad, densely but minutely velvety rusty tomentose outside, smooth shining and rugose inside and with a few small scattered whitish hairs. Roxb. Fl. Ind. iii. 148. Pierre Fl. For. Coch-Chine, t. 190; Wall. Cat. 1133; R. Brown in Benn. Pl. Jav.

Rar. 231. Kurz For. Fl. Burm. i. 138, in part. *S. mollis*, Wall. Cat. 1131; R. Brown in Benn. Pl. Jav. Rar. 231. *S. Balanhas*, L. var. *mollis*, Mast. in Hook. fil. Fl. Br. Ind. i. 358.

Burmah; Griffith No. 578 (Kew Dist.); Helfer Nos. 579, 580; Falconer. Perak, King's Collector, No. 8360.

Roxburgh left in the Calcutta Herbarium an excellent coloured drawing of his *S. angustifolia*. In his Flora Indica he gives a very brief account of the species, drawn up from specimens flowering in the Botanic Garden and which he states came from Nepal. His description is too brief to be of any use: but his figure is so good that I have no hesitation in saying that no species of *Sterculia* collected since Roxburgh's time in any part of the outer Himalaya, or from the plain at its base, is in the least like this plant. I have little doubt that Roxburgh was deceived as to its origin by some changing of labels of the native gardeners at Calcutta (a sublimely inaccurate race!); and that the plant was really received, like so many others during the early years of the garden, from the Straits. Wallich, no doubt deceived by the alleged Himalayan origin of the plant, distributed (as No. 1133 of his list) specimens from the trees of it which were still in his time cultivated in the Calcutta Garden under Roxburgh's name, while specimens collected in Burmah he issued as No. 1131, under the name *S. mollis*, Wall. Pierre's figure above quoted does not agree very well with Roxburgh's, the panicles being by far too short and not nearly hairy enough.

12. *S. RUBIGINOSA*, Vent. Hort. Malmaison, ii. 91. A tree 20 to 50 feet high: young branches rather thick, their apices deciduously rufous-tomentose; the bark pale or brown, striate, glabrous. *Leaves* membranous, obovate-oblong, sometimes ovate-oblong, shortly and abruptly acuminate, entire; narrowed to the acute, rounded or minutely cordate, 3-nerved base: upper surface glabrous, or sparsely stellate-pubescent; the lower stellate-pubescent, most of the hairs pale and minute but these on the midrib and 7 to 10 pairs of spreading stout nerves larger and darker coloured: length 4.5 to 7.5 or rarely 12 in., breadth 2 to 3 in., rarely 4 in.; petiole varying with age from .3 to 1.5 in., rufous tomentose as are the linear caducous .5 in. long stipules. *Panicles* solitary in the axils of the crowded young leaves, many-flowered, shorter than, or as long as the leaves, rufous-tomentose like the outer surfaces of the flowers; flower-pedicels spreading, capillary. *Flower buds* broadly ovate. *Calyx* less than .5 in. long, widely campanulate, divided for half its length or more into 5 lanceolate spreading incurved lobes cohering by their tips, the lobes densely covered inside with white hispidulous hairs. *Male flower*; staminal column longer than the tube or about as

long, glabrous; *anthers* about 10, sessile at the apex of the column, 2-celled, the cells distinct. *Female flower*; gynophore very short; ovaries ovoid, villous (as are the united styles); with 10 sessile anthers at their base; stigma discoid, deeply 5-lobed. *Follicles* 5, coriaceous, crimson when ripe, oblong, shortly beaked, about 2 in. long and 1 in. broad; pubescent externally, glabrous shining and boldly ridged inside. *Seeds* oblong, ovoid, black. Mast. in Hook. fil. Fl. Br. Ind. i. 358; Kurz For. Fl. Burm. i. 138; Pierre Fl. Forest. Coch.-Chine, t. 194 B; Blume Bijdr. i. 82; Br. in Benn. Pl. Jav. Rar. 231; Miq. Fl. Ind. Bat. i. pt. 2, 175. *S. angustifolia*, Jack Mal. Misc. ex Hook. Bot. Misc. i. 287. *S. Jackiana*, Wall. Cat. 1134.

In all the Provinces except the Andaman and Nicobar Islands: at low elevations. Common. Distrib. Java and Sumatra, Cochinchina, Burmah.

Var. *glabrescens*, King: leaves 8 to 12 in. long, by 3 to 4.5 in. broad, softly pubescent beneath when young, much less narrowed to the (always sub-cordate or cordate) base than in the type; panicles much branched and sometimes longer than the leaves. *S. angustifolia*, Kurz (not Roxb.), in part, For. Fl. Burm. i. 138; *S. parviflora*, Kurz (not Roxb.) Journ. As. Soc. Beng. xliii. pt. 2, p. 116. *S. mollis*, Kurz (? of Wall.) l. c. xlv. pt. 2, p. 120. *S. Balanghas*, Linn. var. *glabrescens*, Mast. in Hook. fil. Fl. Br. Ind. i. 358, in part.

Andaman Islands; Helfer (Kew Distrib. No. 595); Kurz, Prain, Bot. Gard. Collectors. Nicobars, Kurz. Great Cocos, Prain. There are no Mergui or Eastern Peninsula specimens of this at Calcutta, and I believe the variety to be confined to the Islands above named.

In this species the petioles lengthen with the age of the leaf, many young leaves having petioles less than .25 in. long, while in old leaves the length varies from 1 to 1.5 in. And there is considerable variability in the size of the blade. Moreover, while in some the upper surface of the leaves is perfectly glabrous (except the midrib which is almost invariably rusty-tomentose), in others it is rough and scaberulous from the presence of scattered stellate hairs. The next species (*S. ensifolia*, Mast.) has, in my opinion, a very poor claim to specific rank; and I think it would be better to treat it a shrubby variety of this with narrower leaves and longer flowers. *S. parviflora*, Roxb. also differs very little from this, and might be reasonably enough regarded as a form of it with broader more glabrous cordate leaves with fewer nerves.

13. *S. ENSIFOLIA*, Mast. in Hook. fil. Fl. Br. Ind. i. 359. A shrub or small tree: young branches and petioles densely ferruginous-tomen-

tose. *Leaves* membranous, oblong-lanceolate or oblanceolate, shortly caudate-acuminate, entire, the base rounded, sometimes minutely cordate, rarely acute; upper surface glabrous, the midrib alone tomentose; under surface sparsely rusty-tomentose, the midrib and 8 to 10 pairs of spreading lateral nerves prominent: length 6 to 12 in., breadth 1·25 to 3·5 in., petiole ·3 to 1·5 in.; stipules erect, linear, half as long as the petiole, deciduous. *Panicles* or *racemes* axillary, solitary, lax, few-flowered, rusty-tomentose, hardly so long as the leaves; bracteoles linear, shorter than the pedicels. *Calyx* ·5 or ·6 in. long, broadly campanulate, pubescent on both surfaces, the tube much shorter than the linear-lanceolate spreading lobes the tips of which curve inwards and cohere. *Male flower*; staminal column longer than the calyx-tube but much shorter than its lobes, glabrous, curved, bearing at its apex 10 2-celled, oblong, nearly sessile, anthers. *Female flower*; gynophore very short: ovaries 5, ovoid, rusty-villous, surrounded at the base by 10 sub-sessile stamens. *Styles* short, united, densely covered with white hairs; stigmas united into a boldly 5-lobed disc. *Follicles* 1 to 5, shortly stalked, narrowly oblong, tapering to each end, the apex with a hooked beak, coriaceous, brownish-velvety, red when ripe, 2 to 2·5 in. long and ·75 in. broad. *Seeds* oval, black, smooth. Pierre Fl. Forest. Coch-Chine t. 194 C. *S. angustifolia*, Jack (not of Roxb.) Mal. Misc. ex Hook. Bot. Mis. i. 287.

Penang, Perak, at low elevations, common. Distrib. Burmah.

I have no doubt whatever that this is the plant described by Jack as the *S. angustifolia* of Roxb.

14. *S. PUBESCENS*, Mast. in Hook. fl. Fl. Br. Ind. i. 357. A tree, the younger parts rusty-pubescent. *Leaves* oblong, obtuse, or abruptly acuminate, entire, the base cordate; upper surface glabrous; lower densely and minutely pubescent, the nerves stellate-pilose: length 4 to 6 in., breadth 2 to 2·5 in.; petiole 1·25 in., sulcate: stipules subulate, ·25 in. long. *Panicle* erect, as long as or longer than the leaves, much branched: ultimate pedicels jointed, pubescent, spreading. *Calyx* ·25 in. long, campanulate; the lobes triangular acute, as long as the tube, hairy within. *Ovary* globose, downy.

Malacca, Maingay.

Except by Maingay's two specimens in the Kew Herbarium, this species is unknown. Specimens of it in good flower and in fruit are much wanted, so that a completer description than the foregoing may be prepared.

15. *S. COLORATA*, Roxb. Hort. Beng. 50. A tree 30 to 60 feet

high; young branches thick, rough, rather pale, glabrous. *Leaves* thinly coriaceous, roundish or reniform, usually palmately 3 to 5-lobed, the lobes triangular, acuminate; base deeply cordate, 5 to 7-nerved; both surfaces pulverulent-pubescent when young, glabrous when adult; length 4.5 to 9 in., breadth 5 to 12 in.; petiole 3.5 to 8 in., puberulous; stipules lanceolate, caducous. *Flowers* in axillary panicles or racemes from the axils of last year's fallen leaves, 2.5 to 4 in. long, densely covered, as is the exterior of the flowers, with coral-red, scaly tomentum. *Calyx* .75 in. long, funnel-shaped, curved, the mouth with 5 acute short triangular teeth, puberulous internally, villous at the base. *Staminal column* as long as, or longer than the calyx, slightly flattened, minutely furfuraceous-pubescent: anthers 20 to 25, sessile at the apex of the column, oblong, closely surrounding the 5 flask-shaped ovaries; styles 5, short, recurved: stigmas acute. *Follicles* 2 to 3 in. long, membranous, glabrous, veined, stipitate, open from an early age and bearing on their edges usually 2 smooth oval seeds. Roxb. Cor. Pl. i. 26, t. 25; Fl. Ind. iii. 146; Mast. in Hook. fil. Fl. Br. Ind. i. 359; Pierre Fl. Forest. Coch-Chine, t. 199; Kurz For. Fl. Burm. i. 138; Brand. For. Flora N. W. Ind. 34; Wall. Cat. 1119; Hook. Ic. Pl. 143; Dalz. & Gibs Bomb. Fl. 23; W. & A. Prodr. i. 63. *Firmiana colorata*. Br. in Benn. Pl. Jav. Rar. 235; Thwaites Enum. 29. *Erythropsis Roxburghiana*, Scott. & Endl. Melet. Bot. 33.

Andamans; Kurz, Prain. Distrib. India, Ceylon.

16. *S. FULGENS*, Wall. Cat. 1135. A tree 30 to 70 feet high; young branches rather thick, with smooth dark bark, at first pubescent, ultimately quite glabrous. *Leaves* large and with long petioles, thinly coriaceous, rotund with 5 shallow acuminate lobes, the base cordate: upper surface glabrous, harsh to the touch: lower densely and minutely stellate-pubescent, palmately 7-nerved, the nerves prominent beneath: length and breadth 15 to 18 in.; petiole 15 to 20 in., sulcate, minutely puberulous. *Racemes* or *panicles* 3 to 4 in. long, from the axils of last year's leaves, densely covered with orange or golden-yellowish scurfy tomentum as are the outer surfaces of the flowers. *Calyx* 1.25 in. long, funnel-shaped, slightly curved, the mouth with 5 short triangular teeth; internally minutely velvety-puberulous with a ring of long matted hair near the base. *Staminal tube* .5 in., longer than the calyx, 5-angled, sulcate, minutely tomentose; anthers 20, sessile, oblong, 1-celled, embracing the 5 flask-shaped ovaries; styles short, reflexed: stigmas acute. *Follicles* unknown. Mast. in Hook. fil. Fl. Br. Ind. i. 360; Kurz For. Fl. Burm. i. 139; Journ. As. Soc. Beng. pt. 2, 1874, p. 117; Wall. Cat. 1135; *Firmiana colorata* var. β , Br. in Benn. Pl. Jav. Rar. 235; Miq. Fl. Ind. Bat., i. pt. 2, 178.

Perak; King's Collector, No. 8673, Scortechini. Distrib. W. Sumatra, Forbes, No. 2105; Java, Burmah; Wallich.

There is no doubt this comes very close to *S. colorata*, Roxb. of which it might possibly be better to treat it as a variety characterised by larger flowers, with much more exerted staminal column, larger leaves, thinner and dark-coloured branchlets. Wallich, however, who saw the tree growing, regarded it as a species; and Robert Brown (Pl. Jav. Rar. p. 235), while treating it as a variety of *colorata*, remarks that it is probably worthy of specific rank. This plant (whether species or variety) is never found in British India proper. Its most northerly limit is Tenasserim, and from thence it extends southward into the Malayan Archipelago. In the Flora of British India, Dr. Masters gives the distribution of this as "Tropical Western Himalayan." The plant, however, which occurs in tropical valleys in that region is just as different from *S. fulgens*, Wall., as that is from *S. colorata*, Roxb. It is the tree to which Wallich gave the name *S. pallens*; and which he published (without describing) in Voigt's Hort. Suburb, Calcutta, p. 105. The leaves of *S. pallens* resemble those of *colorata* in shape; but their under surface is covered with dense pale yellow stellate tomentum. The calyx has a much wider mouth than that of *colorata*, and (like the axis and pedicels of the panicle) is densely covered with a very pale yellow tomentum, while the tomentum of *colorata* is of a vivid coral red. *S. pallens* is confined to the Western Himalaya, just as *S. fulgens* is limited to Burmah and Malaya.

17. *S. ALATA*, Roxb. Hort. Beng. 50. A tree 80 to 150 feet high; young branches rather stout, striate, glabrous. *Leaves* membranous, broadly ovate or ovate-oblong, acute or shortly acuminate, entire; the base deeply cordate, 5 to 7-nerved, some of the basal nerves pinnate on one side; both surfaces glabrous; lateral nerves 4 pairs, prominent on both surfaces as are the midrib and basal nerves; length 4 to 12 in., breadth 3 to 8 in., petiole 1.5 to 7 in.: stipules minute, subulate, caducous. *Racemes* from the axils of previous year's fallen leaves, usually in pairs, sometimes solitary, rarely terminal, about as long as the petioles, flocculent, rusty-tomentose, as are the flowers externally; bracts 3 to each flower, ensiform, caducous. *Calyx* .75 in. long, campanulate, deeply divided into 5 or 6 thick, fleshy, lanceolate segments. *Male flower*; staminal column thin, cylindric, much shorter than the calyx, glabrous, bearing at its apex 25 elongate anthers in five groups of 5 each; ovaries imperfect. *Female flower*; staminodes in 5 phalanges, sessile, embracing the bases of the 5 sub-ovate, multi-ovulate-ovaries; stigmas broad, emarginate. *Follicles* pedunculate, woody, pulverulent.

pubescent, 5 in. in diam., sub-globular, slightly compressed. *Seeds* oblong, compressed, the testa spongy, 1 in. long, with a large obovate thick spongy terminal wing 2.5 in. long and 1.25 broad. Roxb. Corom. Pl. iii. 84, t. 287; Fl. Ind. iii. 152; Kurz Fl. Br. Burm. i. 134; Pierre Fl. Forest. Coch-Chine, t. 196; Wall. Cat. 1125. *Pterygota Roxburghii*, Schott & Endl. Melet. *P. alata*, Br. in Benn. Pl. Jav. Rar. 234. *S. coccinea*, Wall. Cat. 1122, partly. *S. Heynii*, Beddome Flor. Sylvat. t. 230.

Perak, Scortechini; Andamans, Kurz. Distrib. Brit. India, Cochinchina.

18. *S. LINEARICARPA*, Mast. in Hook. fil. Br. Ind. i. 360. A tree 60 to 80 feet high: young branches thick, striate, deciduously pulverulent-tomentose, leaf-cicatrices large. *Leaves* coriaceous, ovate-orbicular, blunt or very slightly narrowed at the apex, edges entire, base deeply cordate, 7-nerved; upper surface glabrous, shining; lower deciduously pulverulent, hairy, almost glabrous when old, minutely reticulate, the midrib and 4 or 5 pairs of lateral nerves prominent: length and breadth 6 to 12 in.; petiole 2.5 to 6 in., sulcate, pulverulent-tomentose. *Panicles* axillary, solitary, stout, erect, as long as or longer than the leaves, rusty pulverulent-tomentose as are the outer surfaces of the calyces, the lateral branchlets short; bracteoles numerous, rotund, concave, caducous; flower-buds globose, sessile. *Calyx* rotate, the tube 1 in. long, with 5 slightly longer ovate acute lobes, tomentose externally, glabrous within. *Staminal column* not so long as the calyx-tube, glabrous; anthers 10, each with a short filament, cuneate, 2-celled; ovaries (rudimentary in some flowers) about 3, free, each 1 or 2-ovulate; style short; stigma entire, small. *Follicles* (? ripe) linear-lanceolate, 3 to 4 in. long and .6 in. broad, stipitate, longitudinally ridged and covered outside and inside with yellowish tomentum as is also the single oblong seed.

Malacca, Maingay. Perak; Scortechini, King's Collector.

The flowers and follicles of this are, in my opinion, those of *Scaphium* rather than of *Firmania*, to which section Dr. Masters has referred it.

19. *S. SCAPHIGERA*, Wall. Cat. 1130. A tree 90 to 120 feet high: young branches rather thick; the bark pale, minutely warted and striate, glabrous. *Leaves* coriaceous, glabrous, ovate to oblong-ovate, sub-acute or bluntnish-acuminate, entire; the base rounded or sub-truncate, often faintly cordate or emarginate, 3 to 5-nerved; main nerves 2 to 4 pairs, sub-erect, prominent on both surfaces; length 5 to 10 in., breadth 2.75 to 4.5 in.; petiole 2 to 5 in., thickened at both ends. *Panicles* only at

the ends of the branchlets, puberulous, shorter than the petioles, robust, with many short spreading branches, many-flowered; pedicels short, pubescent; bracteoles subulate, deciduous. *Calyx* from $\frac{3}{8}$ to $\frac{1}{4}$ in. long, deeply 5-lobed and almost rotate when expanded, stellate-puberulous externally, glabrous internally, the lobes lanceolate. *Male flower* with 15 to 30 anthers almost sessile round the apex of the column and surrounding the rudimentary villous ovary. *Female flower*; ovaries 5, bi-ovulate; styles united; stigma 5-lobed. *Follicles* 1 to 5, on rather stout pubescent stalks, when ripe 6 to 8 in. long and 1.25 to 2.5 in. broad, membranous, boat-shaped, gibbous about the middle, conspicuously veined and more or less puberulous externally especially on the nerves. *Seeds* 1 (rarely 2), ovoid, glabrous, shining, $\frac{5}{8}$ to 1 in. long, attached to the very base of the follicle. *Mast.* in Hook. fil. Fl. Br. Ind. i. 361; Kurz For. Fl. Burm. i. 140; Pierre Fl. Forest. Coch-Chine, t. 201. *Scaphium Wallichii*, R. Br. in Benu. Pl. Jav. Rar. 226.

Malacca, Griffith. Distrib. Sumatra, Burmah.

M. Pierre is in doubt whether his fine figure (l. c. t. 201), represents really the true plant of Wallich. In my opinion it does so most decidedly: R. Brown was right in describing the ovaries as five, and there is a specimen in the Calcutta Herbarium with 5 follicles.

20. *S. AFFINIS*, Mast. in Hook. fil. Fl. Br. Ind. i. 361. A tree: young branches rather stout, rough, dark in colour, the leaf cicatrices large, the very youngest minutely rusty-tomentose. *Leaves* thinly coriaceous, elliptic-oblong, with rather straight edges; the apex broad, suddenly acute; the base truncate (sometimes obliquely so), 3-nerved; both surfaces glabrous, the upper shining, the lower pale and rather dull: main nerves 6 or 7 pairs, conspicuous beneath as is the midrib; length 5 to 9 in., breadth 4.75 to 5.5 in.; petiole 4.5 in., thickened at each end. "*Panicle* erect, as long as the leaves, its branches downy, flattened or angular; peduncles thickly striated, angular, sub-pilose, spreading; ultimate pedicels downy, densely crowded. *Flowers* very small, the buds ovoid. *Flowers* $\frac{1}{25}$ in. *Calyx-lobes* ovate, longer than the funnel-shaped tube. *Follicle* a span long, falcate, leafy, glabrescent, shining within. *Seeds* $\frac{6}{5}$ in. long, solitary, oblong, black." *Scaphium affine*, Pierre Fl. Forest, Coch-Chine, t. 195 E.

Malacca; Maingay, No. 225 (Kew. Distrib.)

The only Maingayan specimen of this in the Calcutta Herbarium consists of leaves only, with a single detached fruit; and I have seen no specimen from any other collector. The foregoing description (as regards inflorescence, flower and fruit) is therefore copied verbatim from Masters (in F. B. I. l. c.).

21. *S. CAMPANULATA*, Wall. A tree 50 to 60 feet high: young branches rather slender, rusty-tomentose, soon becoming glabrous. *Leaves* membranous, broadly ovate, shortly acuminate, entire; the base usually deeply cordate, 3 to 7-nerved; sometimes 3 to 5-lobed; lateral nerves 3 or 4 pairs; upper surface glabrous, the midrib and nerves pubescent or puberulous; lower surface pubescent; length 4 to 6 in., breadth 3.75 to 5.5 in.; petiole 2.25 to 5 in. puberulous: stipules lateral, subulate, caducous. *Panicles* 3 or 4 in. long, in clusters of 2 or 3 at the apices of the branches, few-flowered, glabrous, erect, sub-corymbose; pedicels jointed, about .3 in. long, bracteoles caducous. *Calyx* widely campanulate, more than .75 in. across, green, pruinose, glabrous, veined, its mouth cut half-way down into 5 triangular velvety-edged lobes: *Staminal column* pubescent below. *Ovaries* gibbous at the apex: styles short, cohering; stigmas filiform, recurved: ovules 2, erect. *Follicles* 3 to 6, on slender puberulous stalks, membranous, veined, 2 to 3 in. long, boat-shaped, saccate with a sub-terminal lanceolate wing. *Seeds* sub-globose, with a shining crustaceous testa, .5 in. long or less. Mast. in Hook. fil. Fl. Br. Ind. i. 362; Kurz For. Fl. Br. Burm. i. 139. *Pterocymbium Javanicum*, Br. in Benn. Pl. Jav. Rar. 219, t. 45; Miq. Fl. Ind. Bat. i. pt. 2, 179. *Pt. campanulatum* and *Javanicum*, Pierre Fl. Forest. Coch-Chine, t. 195.

Perak; Fr. Scortechini, King's Collector. Nicobars, Kurz. Distrib. Malayan Archipelago, Burmah.

M. Pierre (l. c.) remarks that, in his opinion, the two species *campanulatum* and *Javanicum*, although closely related, are distinct species; but he does not mention the characters on which he relies for separating them. After dissecting many flowers of the tree (until recently growing in the Botanic Garden, Calcutta), on which Wallich founded his species *campanulatum*, I cannot see any respect in which they differ from Robert Brown's minute and excellent description and figures of *Pt. Javanicum*. I therefore agree with Dr. Masters in considering the two as one and the same species.

22. *S. TUBULATA*, Mast. in Hook. fil. Fl. Br. Ind. i. 362. A tree; young branches about as thick as a goose-quill, tomentose at the very points, the bark dark and rather rough. *Leaves* thinly coriaceous, elliptic-oblong, with a short abrupt rather blunt apiculus; edges entire; the base broadly rounded or sub-truncate, very slightly cordate; when adult both surfaces glabrous except the midrib and main nerves which are minutely rusty-tomentose; main nerves 5 to 7 pairs, spreading, slightly prominent below: length 4 in., breadth 1.75 in.; petiole .75 in. slender, deciduously rusty-tomentose. *Cymes* terminal, as long as the

leaves, many-flowered. *Calyx* 5 in. long, glabrous, narrowly tubular below, the mouth slightly expanded and with 5 ovate-lanceolate lobes shorter than the tube. *Staminal column* pilose; anthers in a ring. *Ovaries* 5; styles inflexed, cohering by their tips. *Follicles* 5, from 2 to 3 in. long and 1 in. broad, on tomentose stalks, oblong, acute, dilated at the base. *Seed* ovoid.

Malacca, Maingay.

At once distinguished by the singular calyx, tubular in its lower, lobed and spreading in its upper, half.

2. TARRIETIA, Blume.

Tall trees. *Leaves* digitate or simple, glabrous or scaly. *Flowers* unisexual, paniced. *Calyx* tubular, small, 5-toothed. *Petals* 0. *Staminal-column* short, bearing a ring of 10-15 very densely clustered anthers, cells parallel. *Ovary* of 3-5 nearly free carpels opposite the sepals; styles as many, short, filiform, stigmatose within; ovules 1 in each cell. *Ripe carpels* of stellately spreading samaras with long falcate wings. *Seeds* oblong; albumen bipartible; cotyledons flat; radicle next the hilum.—*Distrib.* Known species 5 or 6, Australian and Malayan.

Leaves digitately compound.

Under-surface of leaflets persistently
stellate-tomentose 1. *T. Perakensis*.

Under-surface deciduously tomentose,
the hairs simple 2. *T. Penangiana*.

Leaves simple.

Fruit glabrous 3. *T. simplicifolia*.

„ tomentose. 4. *T. Kunstleri*.

1. *T. PERAKENSIS*, King, n. sp. A tree 40 to 60 feet high: young branches, petioles, petiolules, under surface of leaves (when young) and inflorescence with minute deciduous rusty tomentum. *Leaves* digitately compound; leaflets 5 or 6, the lower smaller, obovate-elliptic to obovate-rotund, shortly and rather abruptly acuminate, the edges entire, slightly wavy; the base narrowed; upper surface minutely areolate, glabrous except the very minutely tomentose midrib and nerves; lower glabrous except the midrib: main nerves 10 to 14 pairs, stout and prominent beneath: length of the middle leaflet 4.5 to 5.5 in., of the lower 2 to 3.5 in.: breadth of the middle 2 to 3 in., of the lower 1.25 to 1.5 in.; petiolules .5 to 1 in.; petioles .3 to .45 in. *Inflorescence* in solitary, axillary, cymose racemes or panicles more than half as long as the

leaves, much crowded at the points of the branches. *Flowers* .15 in. long: pedicels slender, three times as long. *Calyx-tube* tomentose externally, sparsely pubescent within; *staminal tube* less than half its length. *Female calyx* rather longer than the male, otherwise the same. *stamens* 0: *ovaries* 5, obliquely ovoid, glabrous, each with a pubescent conic style crowned by a small hooked stigma. *Ripe fruit* compressed-ovoid, 1.25 to 1.5 in. long, and 1 to 1.15 in. in diam., glabrous, the wing falcate, 2 in. long and .5 in. broad, striate.

Perak, at low elevations; King's Collector, Penang, Curtis, No. 2229.

In its leaves this much resembles *T. Javanica*, Bl. (*Rumphia* iii. t. 127, fig. 1); but the leaves of Blume's plant are smaller and have more wavy edges. The flowers, however, of the two differ much in size, those of this being twice as large as the flowers of *T. Javanica*.

2. *T. CURTISII*, King, n. sp. A tree 20 to 40 feet high; young branches, petioles, petiolules and under surfaces of leaves densely covered with rusty stellate, non-deciduous tomentum. *Leaves* digitately 5 or 6-foliolate, the lower smaller, obovate, entire, wavy, apex retuse, base acute; upper surface minutely areolate, glabrous except the stellate tomentose midrib and main nerves; under surface, and especially the midrib, stellate-tomentose: main nerves 9 or 10 pairs, spreading, prominent beneath: length of the middle leaflet 3.5 to 4.5 in., of the lower 1.5 to 2.5 in.; breadth of the middle 2.25 to 2.5 in., of the lower .8 to 1.5 in., petiolules .5 to .75 in., petioles 2 to 2.5 in. *Inflorescence* in solitary, axillary, cymose racemes or few-flowered panicles, more than half as long as the leaves. *Ripe fruit* glabrous, compressed-ovoid, 1 in. long and .8 in. broad; wing narrowly falcate, 1.25 in. long and .25 in. broad, striate.

Penang at 2000 feet: Curtis No. 1427.

This is known only by Curtis's scanty specimens which are in fruit only. Its flowers are unknown. In leaves it closely approaches *T. Perakensis*, but the tomentum is stellate and persistent; whereas in *T. Perakensis*, the hairs are simple and deciduous. The leaflets of this are also smaller, fewer-nerved, more decidedly obovate, less elliptic than in *T. Perakensis*, and they are mucronate rather than acuminate.

3. *T. SIMPLICIFOLIA*, Mast. in Hook. fil. Fl. Br. Ind. i. 362. A tree, young branches pale, sub-glabrous, striate. *Leaves* simple, coriaceous, elliptic or obovate-elliptic, apex truncate or emarginate, shortly mucronate, entire, rather suddenly narrowed at the base or rounded; upper surface glabrous, shining; lower dull, rusty, minutely puberulous,

and slightly scaly; main nerves 16 to 20 pairs, prominent below, spreading; length 4.5 to 7 in., breadth 3 to 4.5 in.; petiole 2 to 3 in., thickened towards the apex. *Cymes* axillary, solitary, many-flowered, 1.5 to 2.5 in. long, minutely rusty-tomentose. *Flowers* 1 in. long; the pedicels shorter, stout. *Calyx-tube* campanulate, minutely tomentose externally, puberulous within; staminal tube short. *Fruit* (including wing) 3 in. long, obliquely spathulate, glabrous.

Malacca; Griffith, Maingay (Kew Distrib.) No. 231.

4. *T. KUNSTLERI*, King, n. sp. A tree 50 to 70 feet high: young branches petioles and peduncles minutely stellate-pubescent and lenticellate. *Leaves* elliptic to obovate-oblong, blunt, mucronate, entire, the base rounded or slightly narrowed: upper surface smooth, shining; the lower pale, sparsely stellate-puberulous on the midrib and nerves, otherwise (under a lens) minutely puberulous: main nerves 7 to 10 pairs, ascending, prominent beneath. *Fruit* at the apex of a solitary stellate-hairy peduncle, ovoid with an oblique sub-spathulate wing, minutely but densely velvety fulvous-tomentose; length of body 1 in. or more; wing about the same length and .6 in. broad.

Perak, near Laroot; King's Collector No. 7581.

Flowers of this are at present unknown. The leaves are at once distinguished from those of *T. simplicifolia* by their pale under surface, and the fruits by their tomentum.

3. *HERITIERA*, Aiton.

Trees. *Leaves* coriaceous, simple, scaly beneath. *Flowers* small, unisexual, in axillary panicles. *Calyx* 5, rarely 4-6 toothed or cleft. *Petals* 0. *Anthers* in a ring at the top of the column, cells 2, parallel. *Ovaries* 5-6, almost free: style short, stigmas 5, thick; ovules solitary in each cell. *Ripe carpels* woody, indehiscent, keeled or winged. *Albumen* 0; cotyledons thick; radicle next the hilum.—A genus of 6 or 7 species, natives of the Tropics of the old world, and of Australia.

H. LITTORALIS, Dryand. in DC. Prod. i. 484. A tree: young branches stout, rough. *Leaves* oblong or elliptic, the apex rounded or acute; the edges entire; base rounded or slightly cordate; lower surface pale; main nerves 7 to 9 pairs, slightly prominent beneath: length 5 to 10 in., breadth 2.25 to 4 in., petiole .5 to .75 in.: stipules lanceolate, caducous. *Flowers* .2 in. long, in many-flowered axillary cymose panicles shorter than the leaves. *Calyx* 5-toothed, puberulous, half as long as the pedicel. *Ripe fruit* 1.5 to 3.5 in. long, woody, compressed ovoid, boldly keeled at apex and on dorsum, glabrous, shining. Mast. in Hook. fil. Fl.

Br. Ind. i. 363; Kurz For. Fl. Burm. i. 140; Pierre Fl. Forest. Coch-Chine, t. 203; Miq. Fl. Ind. Bat. i. pt. 2, p. 179; Blume Bijdr. 84; Roxb. Fl. Ind. iii. 142; W. & A. Prodr. i. 63; Thwaites Enum. 28; Br. in Benn. Pl. Jav. Rar. 237; Miq. Fl. Ind. Bat. i. pt. 2, p. 179. *As Fomes*, Wall. Cat. 1139, partly. *Balanopteris Tothila*, Gaertn. Fruct. ii. 94, t. 99.

All the Provinces, on the coasts. Distrib. Malayan Archipelago and coasts of the tropics of the old world generally, and of Australia.

The plant originally issued by Wallich as *Trochetia contracta* (Cat. No. 1162) and afterwards named by him *Heritiera macrophylla*, (Pierre l. c. t. 204) has by some writers been reduced to *H. littoralis*. But Wallich's species was originally found in the interior of Burmah, and it has since been found in Cachar, far from the sea coast to which *H. littoralis* is strictly confined. *H. macrophylla* has moreover leaf-petioles more than twice as long as those of *H. littoralis*, and its fruit is warted and not smooth. I believe *H. macrophylla* to be a perfectly distinct species; as is also, in my opinion, the other Sylhet and Khasia small-leaved plant which Wallich issued as *H. acuminata*. (Cat. No. 7836.)

4. KLEINHOVIA, Linn.

A tree. *Leaves* 5 to 7-nerved and often cordate at the base. *In-florescence* a terminal, lax, cymose panicle. *Bracteoles* small. *Sepals* 5, much longer than the petals, linear-lanceolate, deciduous. *Petals* 5, unequal, the upper short, ovate-round, saccate, the middle pair concave and obliquely oblanceolate, the lower pair flat with convolute edges. *Stamens* 20, in 5 phalanges of 3 each with five solitary, free, often non-antheriferous, filaments between the phalanges; the filaments of all conjoined below into a long, externally hairy, narrowly cylindrical tube which surrounds the gynophore: anthers 4-celled, divergent. *Ovary* at the apex of the long gynophore and surrounded by the staminal tube, 5-lobed, 5-celled. *Capsule* turbinate-pyriform, membranous, inflated, 5-celled, loculicidal. *Seeds* 1 or 2 in each cell, tubercled: cotyledons convolute, radicle inferior. Distrib. One species. Tropics of the old world.

K. HOSPITA, L. Spec. 1365. *Leaves* ovate-rotund, acuminate, entire, palmately 3-5-nerved at the base, glabrous: length 3 to 6 in., breadth 2.5 to 5 in., petiole 1.5 to 2.5 in. DC. Prodr. i. 488; W. & A. Prodr. i. 64; Roxb. Fl. Ind. iii. 141; Miq. Fl. Ind. Bat. i. pt. 2, 186; Blume Bijdr. 86; Hassk. Pl. Jav. Rar. 313; Mast. in Hook. fil. Fl. Br. Ind. i. 364. Pierre Fl. Forest. Coch-Chine, t. 177.

In all the Provinces, but usually planted. Distrib. Malaya, Australasia, Br. India.

Apparently a variable plant. Dr. Masters (in Oliver's Flora of Trop. Africa, i. 226), describes the African specimens as having no stamens or staminodes alternating with the 5 phalanges of stamens. A specimen in the Calcutta Herbarium from Java has the under surface of the leaves softly hairy.

5. HELICTERES, Linn.

Trees or shrubs, more or less stellate-pubescent. Leaves simple. Flowers axillary, solitary or fascicled. Calyx tubular, 5-fid, often irregular. Petals 5, clawed, equal or unequal, the claws often with ear-shaped appendages. Staminal column surrounding the gynophore, 5-toothed or lobed at the apex; anthers at the top of the column, 2-celled. Five staminodes below the apex of the column. Ovary at the top of the column, 5-lobed, 5-celled; styles awl-shaped, more or less united, slightly thickened and stigmatose at the tips; ovules many in each cell. Follicles spirally twisted, or straight. Seeds tubercled; albumen scanty; cotyledons leafy, folded round the radicle which is next the hilum.—Distrib. About 30 species, natives of the tropics of both hemispheres.

Fruit spirally twisted 1. *H. Isora*.

Fruit not twisted.

Leaves ovate to oblong-lanceolate,
oblique; fruit more than 1 in. long ... 2. *H. hirsuta*.

Leaves lanceolate or oblanceolate, not
oblique: fruit less than 1 in. long ... 3. *H. angustifolia*.

1. *H. ISORA*, Linn. Spec. 1366. A shrub or small tree; young branches minutely tomentose. Leaves ovate-rotund, oblique; the apex rounded, abruptly acuminate; the edges irregularly serrate-dentate, sometimes lobed; the base cordate or rounded, rarely acute, palmately 5- to 7-nerved; upper surface scabrous, minutely hispid; lower pubescent or tomentose; length 2 to 4 in., breadth 1.25 to 3 in.; petiole .3 in. long, tomentose; stipules linear, about as long as the petioles. Flowers axillary, solitary, or in few-flowered minutely bracteolate cymes, 1.5 in. long. Calyx narrowly campanulate, laterally compressed, 2-lipped, 5-toothed, tomentose outside. Petals reflexed, the lower two much shorter and broader than the three upper. Staminal column longer than the petals, curved, very narrowly cylindrical, bearing at its apex 10 to 12 elongate-ovate stamens, and more internally 5 flat bifid staminodes. Ovary ovoid, sulcate, tomentose: styles slender, glabrous, united. Fruit cylindrical, twisted, crowned by the persistent styles, pubescent; 1.5 in. long, .4 in. in diam. Mast. in Hook. fil. Fl. Ind. i. 365; Bl.

Bijdr. 79; Pierre Fl. Forest. Coch-Chine, t. 208, figs. 12 to 25; DC. Prodr. i. 475; Roxb. Fl. Ind. iii. 143; W. & A. Prodr. i. 60; Wight Ic. t. 180; Miq. Fl. Ind. Bat. i. pt. 2, 169; Kurz For. Fl. Burm. i. 142; Brand. For. Flor. 34. *H. chrysocalyx*, Miq. in Pl. Hohen. *Isora corylifolia*, Wight, Hassk. in Tijds.-Nat. Gesch. xii. 107.

Perak; and probably in all the provinces. Distrib. Brit. India.

2. *H. HIRSUta*, Lour. Fl. Coch-Chine, 648. A shrub 6 or 8 feet high; the young branches velvety-tomentose. *Leaves* ovate, or ovate-rhomboid, sub-oblique (oblong to oblong-lanceolate in vars.) acuminate, irregularly crose-separate; the base sub- truncate or rounded, rarely sub-emarginate; upper surface scabrid-pubescent, the midrib and nerves tomentose; lower velvety-tomentose; nerves 4 or 5 pairs, prominent beneath; length 3.5 to 6 in., breadth 1.75 to 2.5 in.; petiole .4 in., tomentose. *Cymes* scorpioid, few-flowered, axillary, solitary, twice as long as the petiole. *Flowers* .75 in. long. *Calyx* narrowly cylindrical-campanulate, coarsely stellate-tomentose externally, the mouth with 5 acute unequal teeth. *Petals* linear, sub-spathulate, two rather broader than the others with slight horn-like appendages about the middle and all longer than the calyx and about as long as the stamens. *Staminal column* and pistil—as in *H. Isora*. *Fruit* cylindrical, acuminate, not twisted, the carpels firmly coherent; externally densely covered by long villous and stellately pilose soft prickles, 1.2 in. long and .35 in. in diam. Pierre Fl. Forest. Coch-Chine, t. 208, figs. 1 to 11; Kurz For. Fl. Burm. i. 143. *H. hirsuta*, Bl. Bijdr. 80. *H. spicata*, Colebr. in Wall. Cat. 1182; Mast. in Hook. fil. Fl. Br. Ind. i. 366; *Oudemansia hirsuta*, Miq. Fl. Ind. Bat. i. pt. 2, p. 171; Hassk. Retzia, i. p. 184; *Orthothecium hirsutum*, Hassk. Pl. Jav. Rar. 308.

Selangore, King's Collector. Penang, Curtis; and probably in the other provinces at low elevations. Distrib. Malayan Archipelago, China, Brit. India.

Var. *oblonga*, (species Wall. Cat. 1183). *Leaves* oblong, 5 or 6 in. long and 1.35 to 1.75 in. broad, sparsely stellate-tomentose beneath.

Penang, Andamans.

Var. *vestita*, (species Wall. Cat. 1844). *Leaves* oblong-lanceolate, oblique at the base; 3.5 to 5.5 in. long and 1 to 1.5 in. broad.

Burmah: ? Andamans.

There seems to be little doubt that Loureiro and Blume independently of each other gave this species the same specific name. Wallich's distribution of it under Colebrooke's MSS. name *spicata* took place many years subsequently, and that name must (although adopted by Dr. Masters) I think fall to the ground.

3. *H. ANGUSTIFOLIA*, L. sp. 1366. A shrub 4 to 6 feet high: young branches, petioles, under surfaces of leaves and peduncles minutely and more or less densely pubescent. *Leaves* lanceolate or oblanceolate, acute (or obtuse and mucronate in var. *obtusa*); entire; the base narrowed 3-nerved; upper surface glabrescent or glabrous; lateral nerves 5 or 6 pairs, not prominent; length 1.5 to 2 in., breadth .4 to .8 in., petiole 2 to .3 in. *Cymes* axillary, solitary, not much longer than the petioles, few-flowered. *Flowers* .4 or .5 in. long. *Calyx* densely stellate-tomentose externally, cylindric, the mouth slightly expanded, with 5 acute triangular teeth, 2-lipped. *Petals* longer than the calyx, linear-subspathulate, with 2 or 3 horned appendages below the middle. *Staminal column* shorter than the petals, narrowly cylindric and otherwise as in *H. Isora*, the stamens smaller. *Ovary* inserted near the apex of the staminal tube, sub-globular, ridged, tomentose. *Fruit* ovoid-cylindric, apiculate, not twisted, the carpels closely coherent, .75 in. long and .4 in. in diam., densely covered with stellate, villous soft prickles as in *H. hirsuta*. DC. Prodr. i. 476; Mast. in Hook. fil. Fl. Br. Ind. i. 365; Bl. Bijdr. 80; Pierre Fl. Forest. Coch-Chine, t. 210 and 211; Wall. Cat. 1180. *H. lanceolata*, DC. Prodr. i. 476; Pierre, l. c. 210 B. *H. virgata*, Wall. Cat. 1181. *Oudemansia integerrima*, Miq. Pl. Jungh. i. 296; Fl. Ind. Bat. i. pt. 2, 170. *Oud. Javensis*, Hassk. Retzia, i. 134. *Orthothecium Javense*, Hassk. Pl. Jav. Rar. 307.

Malayan Archipelago, China.

Var. *obtusa*, (species Wall. Cat. 1184); Pierre, l. c. 211 B, 14 to 25. Kurz in Journ. As. Soc. Beng. 1873, pt. ii. 62. *Leaves* obtuse, mucronate.

Perak; Nicobar Islands.

6. PTEROSPERMUM, Schreb.

Trees or shrubs, scaly or stellate-tomentose. *Leaves* usually bifarious, leathery, oblique, simple or lobed, penninerved. *Peduncles* 1-3, axillary and terminal. *Bracteoles* entire, laciniate, persistent or caducous. *Calyx* of 5 valvate, coriaceous, more or less connate, sepals. *Petals* 5, imbricate, membranous, deciduous with the calyx. *Staminal column* short, bearing opposite to the sepals 3 linear 2-celled anthers, and opposite to the petals 5 ligulate staminodes; cells parallel; connective apiculate. *Ovary* inserted within the top of the staminal column, 3-5 celled; style entire, stigma 5-furrowed; ovules many in each cell. *Capsule* woody or coriaceous, terete or angled, loculicidally 5-valved. *Seeds* winged above, attached in two rows to the inner angle of the cells of the capsule; albumen thin or 0; cotyledons plaited or corrugated,

radicle inferior. Distrib. A genus of about 18 species, confined to tropical Asia.

- | | |
|---|------------------------------|
| Flowers 6 in. long | 1. <i>P. diversifolium</i> . |
| „ 2 in. long. | |
| Sepals shortly pubescent inside,
capsule 3 to 4 in. long. | 2. <i>P. Blumeanum</i> . |
| Sepals with silky hairs inside;
capsule 1.5 in. long, with scaly
hairs | 3. <i>P. Jackianum</i> . |
| „ less than 2 in. long; capsule 2 to
2.5 in., glabrous | 4. <i>P. aceroides</i> . |

1. *P. DIVERSIFOLIUM*, Blume, Bijdr. 88. A tree 60 to 100 feet high: young branches, petioles, under surfaces of leaves and outer surface of sepals and fruit covered with a layer of minute, tawny tomentum with many, more or less deciduous, rufous, stellate hairs on its surface. *Leaves* coriaceous, varying from obovate-oblong to elliptic-rotund; the apex broad, blunt, or sub-truncate, suddenly contracted into a triangular point; the edges entire or sinuous, rarely lobed; the base always cordate or emarginate, 3 to 7-nerved and often oblique: upper surface shining, glabrous, except the tomentose midrib: main nerves 8 to 10 pairs, straight, sub-erect, prominent on both surfaces; length 6 to 9 in., breadth 3.5 to 6 in., petiole 1 to 1.25 or even 2 in., stipules small linear, caducous. *Flowers* 6 to 7 in. long, buds narrowly cylindrical, solitary, or in 3 to 4-flowered sub-sessile axillary cymes; pedicels .2 in. long, each with a minute recurved lanceolate bracteole. *Sepals* coriaceous, slightly shorter than the petals, linear, blunt, adpressed-sericeous internally. *Petals* membranous, linear, glabrescent. *Staminal tube* and gynophore 2 in. long; the free part of the filaments slightly longer; fertile anthers about 10, linear; staminodes 5, pubescent. *Ovary* fusiform, tomentose, 5-celled. *Style* less than 2 in. long, angled, pubescent; stigma fusiform. *Capsule* woody, oblong, pointed, acutely 5-angled, suddenly constricted at the base, about 4 to 5 in. long and 1.5 to 2 in. in diam. *Seeds* flattened, 1.5 to 2 in. long. Mast. in Hook. fil. Fl. Br. Ind. i. 367; Pierre Fl. For. Coch-Chine, t. 179; Miq. Fl. Ind. Bat. i. pt. 2, p. 192; Hassk. Pl. Jav. Rar. 316; Korth. Ned. Kruik. Arch. i. 312. *P. acerifolium*, Zoll. et Mor. Syst. Verz. p. 27 (excl. syn. Willd.)

Perak, Malacca; common; at low elevations. Distrib. Java, Philippines, Cochiu-China.

The leaves on young shoots of this are often peltate and deeply lobed.

2. *P. BLUMEANUM*, Korth. Ned. Kruik. Arch. ii. p. 311. A tree

40 to 50 feet high: young branches slender, almost black when dry; when very young covered by deciduous fuscescent rufous stellate hairs. *Leaves* thinly coriaceous, very inequilateral, oblong to ovate or lanceolate-oblong, entire, acuminate; the base broad, unequally cordate, one side auriculate or sub-auriculate; upper surface very dark when dry, glabrous, shining; the lower densely but minutely tawny or rufous-tomentose with many deciduous cinnamomeous stellate hairs on the surface; main nerves 5 to 7 pairs, prominent beneath; length 3 to 5.5 in., breadth 1.35 to 2 in.; petiole .15 in.; stipules subulate-lanceolate. *Flowers* 2 in. long, solitary, or in 2-3-flowered cymes, axillary, or (by the suppression of the leaves) in terminal racemes: pedicels .5 in. long, bracteate, cylindric in bud. *Sepals* coriaceous, narrowly linear, acute, scurfy, stellate-pubescent externally as are the pedicels and bracteoles, pubescent internally. *Petals* membranous, obliquely oblong-ob lanceolate or sub-spathulate, shorter than the sepals, glabrescent. *Staminal tube* and gynophore about .5 in. long, the free part of the filaments rather longer; fertile anthers about 10; staminodes 5, scaly-pubescent above. *Ovary* ovoid, villous, 5-celled. *Style* shorter than the staminal tube, glabrous: stigma narrowly ovoid. *Capsule* woody, oblong, 5-angled, sub-acute, gradually and slightly narrowed at the base, glabrous when ripe; 3 to 4 in. long and 1.5 in. in diam. *Seeds* flat, 1.5 in. long. Miq. Fl. Ind. Bat. i. pt. 2, p. 191. *Pterospermum lanceaefolium*, Bl. (not of Roxb.) Bijdr. 87. *P. cinnamomeum*, Kurz, For. Fl. Burm. i. 147. *P. Javanicum*, Jungh. Kurz, l. c. i. 147.

Perak, Penang; common at low elevations. Distrib. Sumatra, Java, Borneo, Burmah, Assam.

A very common tree in Perak. Korthal's Bornean species *P. fuscum* appears to me to be nothing more than a very cinnamomeous-tomentose form of this. And the Peninsular-Indian *P. rubiginosum*, Heyne, (Mast. in Hook. fil. Fl. Br. Ind. i. 368) cannot be very different. I should be induced to reduce both to the oldest described species which is this. Of the absolute identity of Kurz's *P. cinnamomeum* with this I have no doubt whatever.

3. *P. JACKIANUM*, Wall. Cat. 1164. A tree: the small branches slender, rather dark, when young covered by a layer of white minute tomentum with many rufous stellate hairs on its surface. *Leaves* sub-coriaceous oblong or elliptic-oblong, slightly inequilateral, entire, or sinuate towards the rather abruptly acuminate apex; the base sub-acute, or truncate and minutely cordate or emarginate, never auricled; upper surface pale brown when dry, glabrous except the puberulous midrib and nerves; under surface pale brown or buff, with

a layer of minute tomentum and on the surface (and especially on the midrib and nerves) many minute deciduous rusty stellate hairs; nerves 10 to 12 pairs, prominent beneath, spreading; length 4 to 5.5 in., breadth 1.5 to 2 in., petiole .25 in.; stipules caducous. *Flowers* 2 in. long: the buds cylindric, acute, solitary, axillary; pedicels .1 in. long, tomentose like the exterior of the sepals, minute, linear-subulate. *Sepals* linear-lanceolate, adpressed-sericeous within. *Petals* shorter than the sepals, oblanccolate, scaly, puberulous externally. *Staminal tube* and gynophore .25 in. long, the free part of the filaments more than twice as long; fertile anthers about 12; staminodes 5. *Ovary* fusiform. *Style* longer than the stamens, pubescent below; stigma cylindric. *Capsule* (*vide* Masters) shortly stalked, ovoid, terete, acute, 1.5 in. long and 1 in. in diam., covered with flat scaly hairs. Mast. in Hook. fil. Fl. Br. Ind. i. 367; *P. oblongum*, Wall. Cat. 1165.

Penang; Jack, Wallich, Curtis. Malacca; Stolickza, at low elevations.

This species does not appear to be a common one. *P. Blumeanum* has probably been mistaken for it.

4. *P. ACEROIDES*, Wall. Cat. 1171. A tree 35 to 50 feet high: young branches rather slender, covered (as are the petioles and under surfaces of the leaves) by a thin felted layer of minute white tomentum, above which is a superficial deciduous layer of loose stellate rufous hairs. *Leaves* coriaceous, more or less elliptic, sometimes obovate-elliptic, the apex abruptly and shortly acuminate, the edge often straight at the sides, sometimes waved, never lobed: the base sub-truncate, often cordate, 5 to 7-nerved; upper surface (when adult) glabrous: main lateral nerves 12 to 15 pairs, straight, oblique; length 5 to 10 in., breadth 3.25 to 5.5 in., petiole .4 to .5 in. *Flowers* 1.5 to 1.75 in. long; solitary, or in 3 to 4-flowered sub-sessile axillary cymes; pedicels .2 in. long, each with a deeply lobed tomentose bract; the buds narrowly cylindric, ribbed. *Sepals* very coriaceous, recurved, longer than the petals, linear, acute, scurfy-tomentose outside, adpressed-pubescent within. *Petals* membranous, obovate, glabrous in the inner, scurfy on the outer, surface. *Stamens* as long as the petals or shorter, the tube only .25 in. long; fertile anthers about 15, linear. *Style* shorter than the stamens, glabrous; stigma clavate; ovary densely sericeous, 5-angled. *Capsule* woody, oblong, pointed at both ends, angled, glabrous, 2 to 2.5 in. long. Kurz in Journ. As. Soc. Beng. 1873, pt. 62; For. Flora Burm. i. 145. *P. acerifolium*, Mast. (not of Willd.) in Hook. fl. Fl. Br. Ind. i. 368, in part. Miq. Ill. Arch. Ind. 84, in part.

Andaman Islands; Helfer, No. 568 (Kew Distrib.), Kurz, King's Collectors. Distrib. Burmah; Wallich.

The nearest ally of this is no doubt *P. acerifolium*, Willd., to which it has been reduced by Dr. Masters. But (having had living trees of both under observation in the Botanic Garden, Calcutta, for many years) I have no hesitation in saying that the two species are quite distinct. *P. aceroides* has entire, not lobed, leaves; much smaller flowers (less than 2 in. long) which expand during December and January: while those of *P. acerifolium* measure 6 in. length and open in March or April. The capsule of *P. aceroides* is moreover only 2 to 2.5 in. long and quite glabrous; while that of *P. acerifolium* is 4 to 6 in. long, with a rough densely stellate tomentose exterior.

7. MELOCHIA, Linn.

Herbs or undershrubs, more or less downy. *Leaves* simple. *Flowers* small, clustered or loosely paniced. *Sepals* 5, connate below. *Petals* 5, spathulate, marcescent. *Stamens* 5, opposite to the petals, connate below into a tube; anthers extrorse, 2-lobed, lobes parallel. *Ovary* sessile, 5-celled; cells opposite the petals, 2-ovuled; styles 5, free or connate at the base. *Capsule* loculicidally 5-valved. *Seeds* ascending, albuminous; embryo straight, cotyledons flat, radicle next the hilum.—*Distrib.* Species about 50, natives of the warmer regions of both hemispheres.

1. *M. CORCHORIFOLIA*, Linn. sp. 944. A pubescent, branching herb or undershrub. *Leaves* membranous, variable, broadly ovate, to ovate-oblong or lanceolate, acute, serrate or obscurely lobed; the base rounded, truncate or sub-hastate, 5-nerved, often plaited; petiole from 4 to 1 in.; stipules linear, minute. *Flowers* 2 in. in diam., in crowded terminal or axillary heads with many villous bracteoles intermixed. *Sepals* lanceolate, acuminate, ascending. *Petals* obovate. *Ovary* villous; styles glabrous. *Capsule* pisiform, pubescent, exceeding the calyx. Willd. Sp. Pl. iii. 604; Roxb. Fl. Ind. iii. 139; Wall. Cat. 1196, in part; Mast. in Hook. Fl. Br. Ind. i. 374. *M. truncata*, Willd. Sp. Pl. iii. 601. *M. supina*, L. Sp. Pl. 944. *M. affinis*, Wall. Cat. 1198. *M. pauciflora*, Wall. Cat. 1199. *Riedleia corchorifolia*, DC. Prodr. i. 491; W. & A. Prodr. i. 66; Miq. Fl. Ind. Bat. i. pt. 2, 188. *R. truncata*, W. & A. l. c. 66. *R. supina*, DC. Prodr. i. 491. *R. concatenata*, DC. Prodr. i. 492. *Visenia corchorifolia*, Spreng. Syst. iii. 30. *V. concatenana*, Spreng. Syst. iii. 30. *V. supina*, Spreng. Syst. iii. 31. *Melochia concatenata*, Wall. Cat. 1197. *Sida cuneifolia*, Roxb. Hort. Beng. 50,

In all the provinces, a common weed. *Distrib.* The Tropics generally.

2. *M. VELUTINA*, Bedd. Fl. Sylvat. t. 5. A large shrub or small tree, all parts pubescent, and with many of the hairs stellate. *Leaves* membranous, long-petioled, broadly ovate, acuminate, coarsely and irregularly serrate; the base 5 to 7-nerved, rounded or cordate; 4 to 9 in. long, by 3.5 to 8 in. broad: petioles 2.5 to 4.5 in.; stipules rounded, .25 in. long. *Cymes* on peduncles longer than the petioles, much branched, spreading, many-flowered, terminal and axillary. *Flowers* .25 in. in diam., pink. *Calyx* campanulate, with 5 deep broad abruptly acuminate teeth. *Petals* narrowly oblong, longer than the calyx, membranous. *Stamens* inserted on a hypogynous disk as are the petals; filaments flat. *Ovary* villous, as are the lower parts of the styles. *Capsules* .3 to .5 in. long, ovoid-cylindric, apiculate, deeply 5-grooved, bristly-tomentose. *Seed* solitary in each cell, its wing ascending. Mast. in Hook. fil. Fl. Br. Ind. i. 374; Kurz For. Fl. Burm. i. 148. *Visnia indica*, Houtt. Linn. Syst. vi. p. 287, t. 46; Miq. Fl. Ind. Bat. i. pt. 2, p. 189. *V. umbellata*, (Houtt.) Bl. Bijdr. 88; Wight Ic. 509. *V. Javanica*, Jungh. in Tijdsch. Nat. Gesch. viii. 302. *Glossospermum velutinum*, Wall. Cat. 1153. *G. ? cordatum*, Wall. Cat. 1155.

In all the Provinces at low elevations—a tree-weed appearing in abandoned fields. Distrib. Malayan Archipelago, British India, Mauritius.

8. WALTHERIA, Linn.

Herbs or undershrubs. *Leaves* simple. *Stipules* linear. *Flowers* small, in dense axillary or terminal clusters. *Sepals* 5, connate below into a bell-shaped tube. *Petals* 5, oblong-spathulate. *Stamens* 5, tubular below; anthers 2-lobed, lobes parallel. *Staminodes* 0. *Ovary* sessile, 1-celled; 2-ovulate. *Styles* 2, distinct, clavate. *Capsule* 2-valved, 1-seeded. *Seeds* ascending, albuminous; embryo straight, cotyledons flat. Distrib. About 15 species, one or two of which are weeds in the Tropics generally; the others are Tropical S. American.

W. INDICA, Linn. sp. 941. A pubescent undershrub. *Leaves* ovate-oblong, obtuse, serrate or crenate, the base rounded or cordate; nerves 5 to 7 pairs, prominent beneath. *Flowers* .25 in. in diam., sessile; bracts linear. *Calyx* campanulate, villous, 10-nerved, the mouth with 5 acuminate teeth. *Petals* oblanceolate, clawed, longer than the calyx. *Capsule* membranous, pubescent. DC. Prod. i. 493; W. & A. Prod. i. 67; Mast. in Hook. fil. Fl. Br. Ind. i. 374; Miq. Fl. Ind. Bat. i. pt. 2, p. 187; Wall. Cat. 1194. *W. Americana*, L. DC. Prod. i. 492. *W. elliptica*, Cav. Diss. vi. 171; Wall. Cat. 1195.

In all the Provinces: a weed. Distrib. The Tropics generally.

9. *ABROMA*, Jacq.

Trees or shrubs. *Leaves* cordate, ovate-oblong, serrulate, sometimes angled. *Peduncles* opposite the leaves, few-flowered. *Sepals* 5, connate near the base. *Petals* 5, purplish, concave below, prolonged above into a large spoon-shaped lamina. *Staminal-cup* of 5 fertile and as many sterile divisions; fertile filaments opposite the petals, 3-antheriferous; anthers 2-lobed, lobes divergent. *Staminodes* longer than the fertile filaments, obtuse. *Ovary* sessile, pyramidal, 5-lobed; cells many-ovuled, styles 5. *Capsule* membranous, 5-angled, 5-winged, truncate at the apex, septicidally 5-valved, valves villous at the edges. *Seeds* numerous, albuminous; embryo straight, cotyledons flat, cordate, radicle next the hilum.—Distrib. 2 or 3 species, natives of Tropical Asia.

1. *A. AUGUSTA*, Linn. fl. Suppl. 341. A pubescent large shrub or small tree: young branches pale. *Leaves* 5 to 7-nerved at the base, 3.5 to 6 in. long and 3 to 5 in. broad; petiole 1.75 to 2.5 in., the upper much smaller and narrower. *Stipules* linear, deciduous. *Flowers* 2 in. in diam., peduncles 1.5 in., extra-axillary. *Sepals* 1 in. long, lanceolate, free to nearly the base. *Petals* longer than the sepals, imbricate, deciduous. *Capsule* 1.5 to 2 in. in diam., glabrous or nearly so when ripe. DC. Prod. i. 485; Mast. in Hook. Fl. Br. Ind. i. 375; Bl. Bijdr. 85; Roxb. Hort. Beng. 50; Fl. Ind. iii. 156; Miq. Fl. Ind. Bat. i. pt. 2, 183; Beddome Flor. Sylvat. Anal. Gen. t. 5; W. & A. Prodr. i. 65; Wall. Cat. 1142. *A. angulata*, Lam. Ill. 636. *A. Wheeleri*, Retz. Obs. v. 27; Willd. Sp. Pl. iii. 1425. *A. fastuosum*, Gærtn. Fruct. i. 307, t. 64.

In all the Provinces at low elevations: usually near cultivation. Distrib. Malayan Archipelago, Philippines, China, Brit. India.

The bark yields a stout fibre.

10. *BUETTNERIA*, Linn.

Erect climbing or tomentose shrubs, herbs, or trees; sometimes prickly. *Leaves* various. *Flowers* minute, in axillary or terminal much-branched, umbellate cymes. *Sepals* 5, slightly connate near the base. *Petals* 5, unguiculate, concave, inflexed, with 2 small lateral lobes, and a long sub-terminal simple linear or narrowly lanceolate appendage. *Staminal tube* with 5 broad truncate or emarginate teeth and, between them, five 2-celled extrorse anthers (mouth entire in *B. Curtisii*). *Ovary* sessile, 5-celled, the cells 2-ovulate. *Style* entire, 5-fid. *Capsule* globose, echinate, septicidally 5-valved, the cells 1-seeded. *Seed* ascending, exalbuminous: cotyledons folded round the radicle. Distrib. About 48 species, mostly tropical American: a few tropical Asiatic and one African.

Leaves longer than broad, their bases not cordate or only minutely so.

Staminal tube with entire mouth ... 1. *B. Curtisii*.
Staminal tube with its mouth 5-lobed.

Leaves quite glabrous.

Capsule less than 1 in. in diam., covered with glandular barbed spines ... 2. *B. uncinata*.

Capsule more than 1 in. in diam., covered with short subulate spines ... 3. *B. Maingayi*.

Leaves more or less minutely hispid on both surfaces ... 4. *B. elliptica*.

„ hispid on the upper, hispid-tomentose on the lower, surface ... 5. *B. Jackiana*.

Leaves about as broad as long, deeply cordate at the base.

Leaves glabrous, or glabrescent, not lobed ... 6. *B. aspera*.

„ sparsely pubescent, often lobed ... 7. *B. Andamanensis*.

1. *B. CURTISII*, Oliver in Hook. Ic. Pl. t. 1761. A slender woody creeper, 10 to 15 feet long; young branches minutely puberulous. Leaves linear-lanceolate or oblanceolate-oblong, rarely ovate-oblong, acuminate, entire, narrowed to the sub-obtuse, minutely cordate, 5-nerved base: lateral nerves numerous, unequal and spreading at various angles, reticulations distinct: both surfaces glabrous, the lower with tufts of stellate hairs in the axils of the leaves; length 2.5 to 7 in., breadth .5 to 2 in.; petiole .25 in., pubescent. Cymes in axillary fascicles of 2 to 4, slender, 3 to 7-flowered, puberulous; peduncles about 1 in. long: flower pedicels .25 in. Flowers .45 in. in diam., buds conical. Calyx deeply 5-partite, the segments ovate-lanceolate, acuminate. Petals strap-shaped with 2 rather broad inflexed lateral lobes, and a long cylindrical curved sub-terminal appendage, about as long as the sepals. Capsule globular, about 1 in. in diam., veined, pubescent, and armed with numerous straight smooth bristles.

Penang, Curtis, Nos. 817 and 1166; Perak, common at low elevations.

This is closely allied to the Bornean *B. lancifolia*, Hook. fil. The leaves vary a good deal in shape, the most prevalent form in the Perak

specimens being linear-lanceolate. The Penang specimens are, on the other hand, as figured by Professor Oliver, oval-oblong.

2. *B. UNCINATA*, Mast. in Hook. fil. Fl. Br. Ind. i. 377. A woody climber: young branches at first scurfy and hispid, but very soon glabrous. *Leaves* sub-coriaceous, elliptic-oblong; gradually tapering in the upper third to the acuminate apex, entire, the base slightly cuneate, 3-nerved; both surfaces glabrous and shining, nerves 9 or 10 pairs, spreading, thin but prominent beneath: length 9 to 11 in., breadth 3 in., petiole nearly 3 in., thickened at the apex, glabrous. *Sepals* lanceolate, spreading, hispid. *Fruiting peduncles* (fide Masters) "half the length of the leaves. *Capsule* depressed-spheroidal, the size of a hazelnut, covered with hooked gland-tipped barbed hispid spines, 3-celled."

Malacca, Maingay, No. 242 (Kew Distrib.).

I have seen only Maingay's Malacca specimens.

3. *B. MAINGAYI*, Mast. in Hook. fil. Fl. Br. Ind. i. 377. A woody climber: young branches glabrous. *Leaves* sub-coriaceous, elliptic to elliptic-oblong, shortly bluntly and rather abruptly acuminate, entire; the base rounded with 3 bold and 2 minute nerves: both surfaces quite glabrous; lateral nerves about 2 or 3 pairs, prominent beneath as are the reticulations; length 7 or 8 in., breadth 3 to 3.5 in.; petioles 1.2 in., thickened towards the apex, glabrous. *Umbels* in axillary fascicles of 6 or 8, their peduncles about 1 in. long, slender, glabrescent; pedicels .25 in. *Sepals* .25 in. long, ovate-lanceolate. "*Petals* shorter than the sepals, with a long linear appendage. *Staminodes* erect, oblong, obtuse, bifid. *Style* as long as the ovary. *Fruiting peduncle* as long as the petiole. *Capsule* globose, 1.25 in. in diam., obscurely 5-lobed, studded with short subulate prickles."

Malacca; Griffith, Maingay.

Of this species I have seen no good specimens in flower or fruit, and the above account of these parts is taken from Masters' description.

4. *B. ELLIPTICA*, Mast. in Hook. fil. Fl. Br. Ind. i. 377. A woody climber; young branches minutely rusty-tomentose. *Leaves* broadly elliptic, abruptly and shortly acuminate, entire; the base 5-nerved, rounded or minutely cordate; upper surface minutely scabrid-hispid, the midrib and nerves hispid-tomentose; lower minutely pubescent on the veins, the midrib and longer nerves tomentose: lateral nerves 3 pairs, oblique, curving, prominent beneath as are the secondary nerves and reticulations: length 5.5 to 7.7 in., breadth 3.5 to 4.75 in., petiole

2 to 3 in., tomentose. *Umbels* pedunculate, solitary or in fascicles of 5 or 6, axillary, few-flowered, stellate-tomentose: peduncles .5 to 1 in.; pedicels .1 to .25 in., both slender. Buds .1 in. in diam. *Sepals* ovate, acute. *Petals* rounded, with long cylindric inflexed apices longer than the sepals. *Fruit* unknown.

Malacca, Maingay: No. 241 (Kew Distrib.). Perak; Scortechini.

Evidently a rare species; for I have seen, besides Maingay's, only Scortechini's solitary specimen.

5. *B. JACKIANA*, Wall. in Roxb. Fl. Ind. (ed Carey) ii. 386. A stout woody creeper, the young branches with densely minute ferruginous tomentum some of which is stellate. *Leaves* narrowly or broadly elliptic, acuminate, entire; the base boldly 3-nerved, rounded, sometimes slightly cordate, rarely acute; upper surface sparsely and shortly hispid; under surface rufous, hispid-tomentose especially on midrib and nerves, many of the hairs on both surfaces stellate; lateral nerves 3 or 4 pairs, curved, spreading; length 3.5 to 6 in., breadth 2 to 3.75 in.; petiole to 1 in. tomentose. *Umbels* pedunculate, solitary or in groups of 3 from the leaf-axils, few-flowered, tomentose: peduncles .25 to 1 in. long; pedicels about .35, slender. *Sepals* linear-subulate, spreading, hispid, about .4 in. long. *Petals* sub-rotund, lobed, each with a single long cylindric terminal appendage as long as the sepals. *Capsule* globose, slightly 5-furrowed, 1 in. in diam., black, glabrescent, armed with many straight spines. *Seeds* oblong, black. Mast. in Hook. fil. Fl. Br. Ind. i. 376; Wall. Cat. 1147.

Penang, Perak and Singapore; at low elevations.

6. *B. ASPERA*, Colebr. in Roxb. Fl. Ind. (ed. Carey), ii. 383. A powerful woody climber often with a tree-like stem; young branches glaucous, minutely and deciduously pubescent. *Leaves* sub-orbicular or ovate-orbicular, shortly acuminate, entire, the base cordate, 5 to 7-nerved; upper surface glabrous, shining; the lower glabrescent, the midrib and nerves puberulous; lateral nerves 4 to 6 pairs, prominent beneath as are the stout transverse veins; length 4.5 to 7.5 in., breadth about the same; petiole 2 to 5 in., glaucous-pubescent at first, afterwards glabrous. *Oxympes* axillary, solitary or fasciculate, pedunculate, much branched, many-flowered, pubescent to tomentose, 3 or 4 in. long; ultimate pedicels .5 in. long, slender. *Sepals* lanceolate, acute, spreading, .15 in. long, puberulous externally. *Petals* cuneate, shorter than the sepals, 3-lobed, the middle lobe linear-lanceolate, reflexed. *Staminodes* truncate. *Ovary* globular, scabrid. *Capsules* globular, 1.5 to 2 in. in diam., slightly depressed, pubescent when young, glabrous when ripe,

armed with many long, nearly straight, sharp spines. *Seeds* oblong, .5 in. or more long. Wall Cat. 1144; Mast. in Hook. fil. Fl. Br. Ind. i. 377; Kurz For. Fl. Burm. i. 151; Pierre Fl. Forest. Coch-Chine, t. 206, figs. 1 to 8. *B. grandifolia*, DC. Prodr. i. 486. *B. nepalensis*, Turcz. in Bull. Mosc, 1858, 207.

Andaman Islands. Distrib. Brit. India, China, Cochin-China.

7. *B. ANDAMANENSIS*, Kurz in Journ. As. Soc. Bengal, 1871, ii. 47. A woody climber: young branches scaberulous. *Leaves* sub-orbicular, crenate and palmately 3 to 5-lobed, the lobes acuminate; or ovate-rotund, acuminate and irregularly serrate-crenate and not lobed; the base always deeply cordate, 5 to 7-nerved; lateral nerves about 5 pairs, opposite, prominent beneath as are the midrib and straight transverse veins; both surfaces sparsely pubescent at first, but afterwards glabrous. *Cymes* umbellate, 2 or 3 times branched, spreading, many-flowered, solitary, or 2 or 3 in a fascicle, axillary: the common peduncle stout, .6 to .75 in. long; secondary peduncles about the same length, tertiary half as long; flower-pedicels .15 in., all slender and slightly pubescent. *Sepals* ovate acuminate or deltoid, puberulous externally. *Petals* with 2 obscure lateral lobes, and a long lanceolate inflexed middle lobe. *Staminodes* truncate. *Capsule* globose, less than 1 in. in diam, glaucous when young, armed with a few unequal, rather short, smooth, stiff spines. Kurz in Flora, 1871, p. 277; For. Fl. Br. Burm. i. 152; Mast. in Hook. fil. Fl. Br. Ind. i. 377; Pierre Fl. Forest. Coch-Chine, t. 207, figs. 1 to 9.

Andaman Islands. Distrib. Burmah, Siam, Cochin-China.

11. *COMMERSONIA*, Forsk.

Trees or shrubs. *Leaves* simple, oblique. *Inflorescence* cymose, terminal or axillary or leaf opposed. *Calyx* 5-cleft. *Petals* 5, concave at the base, prolonged into a long strap-shaped appendage at the apex. *Fertile stamens* 5, opposite the petals; anthers subglobose, 2-celled, cells diverging; *staminodes* 5, opposite to the sepals, lanceolate. *Carpels* 5, opposite to the sepals, connate; *styles* connate; *ovules* 2-6. *Capsule* loculicidally 5-valved, covered with bristly hairs. *Seeds* ascending, albuminous, strophiolate; cotyledons flat, radicle next the hilum. Distrib. A genus of about 8 species, some of which are natives of the Malay peninsula and Archipelago, others of Australia.

C. PLATYHYLLA, Andr. Bot. Rep. t. 603 (note). A low tree; young branches softly rusty-tomentose. *Leaves* membranous, inequilateral, ovate-acuminate, irregularly dentate-serrate; the base more or less cordate, one side sub-auriculate, upper surface sparsely and minutely

stellate-hairy, lower softly hoary tomentose: length 5 to 8 in., breadth 3 to 4.5 in., petioles .2 to .3 in.; stipules shorter than the petioles, scarious, lobed. *Cymes* corymbose, much shorter than the leaves, spreading, much branched, tomentose. *Flowers* .2 or .25 in. in diam. *Calyx* pubescent, cut nearly to the base into 5 ovate-lanceolate lobes. *Petals* as long as the sepals but much narrower, concave at the base; the terminal appendage elongate, narrowly oblong, its edges inflexed. *Stamens* 5, the anthers broad, extrorse. *Staminodes* 5, lanceolate, spreading, reflexed; shorter than the petals. *Ovary* 5-celled. *Capsule* globose, .4 or .5 in. in diam., densely covered with long soft, flexuose, pubescent bristles. *Mast.* in Hook. fil. Fl. Br. Ind. i. 378. *C. Javensis*, G. Don. Gen. Syst. i. 523; Hassk. Pl. Jav. Rar. 312. *C. echinata*, Blume Bijdr. 86; Wall. Cat. 1143; Andr. Bot. Rep. t. 519, not of Forst. *C. echinata*, var. β . Miq. Fl. Ind. Bat. i. pt. 2, 182. *Buettneria hypoleuca*, Turcz. in Mosc. Bull. 1858, 207.

In all the provinces except the Andamans and Nicobars. Distrib. Malayan Archipelago, Philippines.

12. LEPTONYCHIA, Turcz.

Shrubs or trees. *Leaves* simple, entire. *Flowers* in small axillary cymes. *Sepals* 5, valvate, united near the base. *Petals* 5, valvate, short, orbicular, concave. *Andræcium* tubular below, filamentiferous above, filaments 3-seriate, outer series of 5 to 10 ligulate staminodes opposite the petals, middle of 10 fertile stamens also opposite the petals, innermost of 5 very short fleshy subulate staminodes opposite the sepals; anthers linear-oblong, introrse, dehiscing at the sides longitudinally. *Ovary* sessile, 3-4-celled; placentas axile; styles connate, stigmas capitate; ovules many in each cell, anatropous. *Capsule* 2-3-celled, or by abortion 1-celled, dehiscing septicidally or loculicidally, or both simultaneously or irregularly. *Seeds* black, with a fleshy yellowish arillus; albumen fleshy, cotyledons foliaceous, radicle superior. Distrib. three or four species—Indo Malayan and Tropical African.

L. GLABRA, Turcz. in Mosc. Bull. for 1858, p. 222. A tree: the young branches glabrous. *Leaves* ovate-oblong or oblong-lanceolate, rather abruptly acuminate, the base slightly narrowed or rounded, 3-nerved; both surfaces glabrous, shining; main nerves 4 to 7 pairs, thin: length 4 to 8 in., breadth 1.6 to 3 in.; petiole .4 to .5 in., glabrous. *Flowers* .25 in. in diam.; the buds oblong, obtuse, 5-ridged. *Sepals* ovate-lanceolate or oblong, rather obtuse, spreading, pubescent on both surfaces, not veined. *Petals* about one-fourth the size of the sepals, broad, truncate, villous. *Stamens* 10, in five phalanges of two each,

nearly as long as the sepals. *Staminodes* 10 to 20, glabrous, the outer 5 to 15 shorter than, or as long as, the stamens, filiform; the inner invariably 5, short, subulate, internal to, and alternating with, the phalanges of stamens. *Ovary* broadly obovate, obtuse, obscurely 4-grooved, with a few scattered hairs near the apex, 4-celled. *Style* cylindrical, tapering, with sparse spreading hairs. *Capsule* coriaceous, depressed-obovoid, pale greyish, .5 in. long, rugose; within shining pale and wrinkled. *Seed* solitary, oblong, black, less than half covered by a thin arillus proceeding from its side. Mast. in Hook. fil. Fl. Br. Ind. i. 379; Kurz För. Fl. Burm. i. 150; Oudem. in Compt. Rend. Ac. Roy. Sc. Amsterd. 2 Ser., 11, 8, cum ic; Walp. Ann. vii. 449. *Grewia? caudata*, Wall. Cat. 1099. *L. heteroclita*, Kurz For. Fl. Burm. i. 150. *G. heteroclita*, Roxb. Fl. Ind. ii. 590. *Bimindykia trichostylis*, Kurz in Nat. Tijdsch. Ned. Ind., Ser. 3, iii. 164. *Turraea trichostylis*, Miq. Fl. Ind. Bat. Suppl. 502.

Malacca, Penang, Perak, Andamans; at low elevations. Distrib. Malayan Archipelago, Burma.

Var. *Mastersiana*, young branches, midribs and petioles of leaves puberulous; flowers .5 in. in diam., the buds pointed; sepals 3-veined: outer staminodes varying from 5 to 15, often pubescent in the upper half: ovary oblong-ovoid, villous, 3-celled: style glabrous: capsule black. *L. acuminata*, Mast. in Hook. fil. Fl. Br. Ind. i. 379.

Malacca and Perak. Distrib. Sumatra, Borneo, Burmah.

This shrub or small tree is common, and I have thus had the advantage of being able to examine a large number of flowers. The result of my examination of these is that, whereas the inner staminodes are invariably 5 in number, the outer series varies in number in the most perplexing way from 5 to 15. Where there are 10, they are always arranged in pairs united at the base: and where there are 15, they are arranged in threes united at the base. The proper view to take of these staminodes is I believe therefore that they are single organs, but sometimes deeply cleft into 2 or 3 linear and equal segments. On this account, and also on account of the similarity of the other organs, I am induced to think that there is but one species of *Leptonychia* and that Masters' species *acuminata* and Beddome's *L. moacurroides* are merely forms of the species on which Turczaninow originally founded the genus.

ORDER XIX. TILIACEÆ.

Trees, shrubs or herbs. *Leaves* alternate, rarely opposite, simple or lobed. *Stipules* free, usually caducous. *Flowers* usually cymose, or in cymose panicles, or racemose. *Flowers* regular, hermaphrodite, rarely unisexual. *Sepals* 3-5, free or connate, valvate. *Petals* as many as the sepals, rarely absent, imbricate or valvate. *Stamens* numerous, rarely

definite, usually springing from a prolonged or dilated torus, free or sometimes 5-adelphous, filaments filiform; anthers 2-celled. *Ovary* free, 2-10-celled; styles columnar, or divided into as many divisions as there are cells to the ovary, stigmas usually distinct, rarely confluent or sessile. *Ovules* attached to the inner angle of the cells of the ovary; if few in number, often pendulous from the apex or ascending from the base; if more numerous, disposed in 2 or more ranks, anatropous; raphe ventral or lateral. *Fruit* fleshy or dry, dehiscent or indehiscent, 2-10 or by abortion 1-celled (cells sometimes divided by false partitions); carpels separable or always united. *Seeds* 1 or many, ascending, pendulous or transverse, with no arillus; testa leathery or crustaceous or pilose; albumen fleshy, abundant or scanty, rarely wanting; embryo straight or slightly curved, cotyledons leafy, rarely fleshy, radicle next the hilum.—*Distrib.* about 370 species; most abundant in the tropics of either hemisphere.

Series A. *Holopetalæ*. *Petals* glabrous or rarely downy, coloured, thin, unguiculate, entire or nearly so, imbricate or twisted in the bud. *Anthers* globose or oblong, opening by slits.

Tribe I. *Brownlowiæ*. *Sepals* combined below into a cup. *Anthers* globose, cells ultimately confluent at the top.

* *Staminodes* 5.

Carpels distinct, 2-valved 1. *Brownlowia*.
Carpels combined, indehiscent, winged ... 2. *Pentace*.

** *Staminodes* 0.

Stamens on a raised torus 3. *Schoutenia*.
Stamens on a contracted torus ... 4. *Berrya*.

Tribe II. *Grewiæ*. *Sepals* distinct. *Petals* glandular at the base. *Stamens* springing from the apex of a raised torus.

Fruit drupaceous, not prickly 5. *Grewia*.

Fruit dry indehiscent or 3-5 coccous, prickly 6. *Triumfetta*.

Tribe III. *Tiliæ*. *Sepals* distinct: petals not glandular. *Stamens* springing from a contracted torus.

Herbs or undershrubs with 3 or 5-celled capsules: seeds without hairs ... 7. *Corchorus*.

Trees with 2-celled capsules; seeds with marginal hairs 8. *Trichospermum*.

Series B. *Heteropetalæ*. *Petals* usually incised, rarely entire or absent, induplicate or imbricate not twisted: anthers linear, opening by a terminal pore often with an apical awn or tuft of hairs.

Stamens on a raised torus; fruit drupaceous 9. *Elæocarpus*.

1. BROWNLOWIA, Roxb.

Trees. Pubescence stellate or scaly. Leaves entire, 3-5-nerved, feather-veined. Flowers numerous, small, in large terminal or axillary panicles. Calyx bell-shaped, irregularly 3-5-fid. Petals 5, without glands. Stamens many, free, springing from a raised torus. Staminodes 5, within the stamens, opposite the petals and petaloid. Anthers subglobose. Ovaries 5, each 2-ovulate; styles awl-shaped, slightly coherent; ovules ascending. Carpels ultimately free, 2-valved, 1-seeded. Albumen 0; cotyledons thick, fleshy.—Distrib. Nine species confined to Tropical Asia.

Leaves not peltate.

- | | | | |
|---------------------------------------|-----|-----|-------------------------------|
| Leaves lanceolate | ... | ... | 1. <i>B. lanceolata</i> . |
| „ broadly elliptic to elliptic-rotund | | | 2. <i>B. Kleinhowioidea</i> . |

Leaves peltate.

- | | | |
|-------------------------------|-----|-----------------------------|
| Leaves minutely hairy beneath | ... | 3. <i>B. Scortechinii</i> . |
| „ glabrous on both surfaces | ... | 4. <i>B. macrophylla</i> . |

1. BROWNLOWIA LANCEOLATA, Benth. in Journ. Linn. Soc. V. Suppl. ii. 57. A tree 25 to 30 feet high; young branches pale when dry, sublepidote. Leaves thinly coriaceous, lanceolate or oblong-lanceolate, acuminate, the base obtuse; upper surface when adult glabrous, shining, the lower covered by a dense layer of minute whitish yellow shining scales: main nerves 6 to 8 pairs (1 pair of them basal), not prominent: length 4.5 to 6 in., breadth 1.5 to 1.75 in., petiole .25 to .4 in. Panicles axillary or terminal, 1 to 3 in. long, and less than 1 in. across, few-flowered. Flowers .25 in. long, their pedicels about as long. Calyx .2 in. long, scaly like the pedicel, its lobes lanceolate. Petals longer than the calyx, oblong, blunt, slightly narrowed to the shortly unguiculate base, glabrous. Anther-cells sub-divaricate, sub-confluent when adult. Ovary deeply 3 to 5-lobed, pubescent, the cells 2-ovuled. Ripe carpels distinct, sub-globose, truncate, compressed on their inner surfaces, minutely lepidote and pubescent, .5 in. in diam. Seed solitary, with thin testa and large sub-hemispheric cotyledons. Hook. fil. Fl. Br. Ind. i. 381: Kurz For. Flora Burm. 154.

Malacca, Griffith. Distrib. Burmah and Bengal; in tidal forests and mangrove swamps.

The young parts are covered with rusty or pale brown scales, but the adult branchlets leaves and flowers are as above described.

2. BROWNLOWIA KLEINHOWIOIDEA, King, n. sp. A tree 40 to 50 feet high: young branches rather slender, covered with a dense thin layer of

minute pale brown hair. *Leaves* thinly coriaceous, broadly elliptic to elliptic-rotund, slightly narrowed to the obtuse apex, very little narrowed to the more or less cordate base: upper surface glabrous, very sparsely lepidote, the lower covered with a thin layer of very minute pale hair; basal nerves 4 or 6 (two of them small): main lateral nerves 3 pairs; transverse secondary nerves distinct: length 5 to 7 in., breadth 3.5 to 4.5 in.; petiole 2.5 to 3 in., thickened towards the apex, pubescent like the under surfaces of the leaves. *Panicles* mostly terminal, rarely axillary, 9 to 15 in. long (the axillary ones much smaller) the branches rather few, spreading little, the flowers rather closely clustered on the branchlets. *Flowers* .25 in. long, on pedicels about half as long. *Calyx* widely campanulate, cut for a third of its length into 5 acute triangular teeth, minutely tomentose externally, glabrous inside. *Petals* longer than the calyx, oblong, very obtuse, slightly narrowed but thickened towards the rather long basal claw. *Staminodes* linear, flat, about as long as the filaments. *Ovaries* 3 to 5, sub-globose, laterally compressed, pubescent. *Styles* subulate, a little longer than the stamens, slightly coherent. *Fruit* unknown.

On Gunong Bubu in Perak, at elevations of 600 to 1000 feet; King's Collector.

A species with leaves not unlike these of *Kleinhovia hospita*: in many respects closely allied to *B. elata*, but with much smaller flowers.

3. *BROWNLOWIA SCORTECHINII*, n. sp., King. A small slender tree: young branches stout, pale, sparsely lenticellate, pubescent at first but soon glabrous. *Leaves* coriaceous, ovate-elliptic, peltate, slightly narrowed to the acute or sub-acute apex; the edges sub-undulate; very little narrowed to the rounded, or sometimes sub-emarginate, base; upper surface glabrous; the lower pale from a thin continuous layer of very minute hairs; petiole attached 2.5 to 3 in. above the base, nerves radiating from it about 9, lateral nerves from the midrib about 4 pairs; all rather prominent beneath, as are the transverse secondary nerves: length 10 to 15 in., breadth 5.5 to 7 in., petiole 7 to 9 in. long, thickened at both ends. *Panicle* terminal, 6 to 12 in. long and about 6 in. broad, or sometimes small narrow panicles in terminal clusters of 6 to 10: branches spreading, compressed, puberulous; bracteoles ovate, fugaceous; pedicels, stout, .15 in. long in the bud but lengthening as the flower expands, puberulous. *Flowers* .6 in. long, crowded. *Calyx* narrowly campanulate; its teeth half as long as the tube, lanceolate, sub-acute, tomentose-lepidote externally. *Petals* longer than the calyx, narrowly obovate, much narrowed to the clawed base. *Staminodes* linear, about as long as the filaments. *Ovaries* 5, ovoid, compressed, stellate-pubescent.

Styles slightly longer than the stamens, subulate, bent at the apex. *Fruit* unknown.

Perak; Scortechini, No. 1918.

Collected only once by the late Fr. Scortechini and referred by him to *B. elata*, Roxb. The species is, however, quite distinct from *B. elata*; and also from *B. peltata*, which it more resembles in its leaves.

4. *BROWNLOWIA MACROPHYLLA*, King n. sp. A tree 30 to 40 feet high: young branches very stout, deciduously rufous-puberulous. *Leaves* very coriaceous, rotund, those on the older branches elliptic, the apex rounded or very slightly and shortly apiculate, the edges subundulate, the base broad, emarginate or slightly cordate, both surfaces glabrous; main nerves 7 to 9 basal and about 2 pairs lateral, prominent on both surfaces, secondary nerves transverse and very distinct: length of the rotund leaves 11 to 17 in., breadth 10 to 14: of the elliptic, length 6 to 10 in., breadth 3.5 to 5.5 in.: petiole 2.25 to 4.5 in., thickened at both ends. *Panicle* terminal, almost as long as the leaves, its branches numerous, compressed, grooved, spreading, scurfy and rusty-pubescent: bracts few, linear-lanceolate, nearly 1 in. long, persistent. *Flowers* .65 in. long: their peduncles shorter than the calyx, stout, deeply grooved. *Calyx* rather widely cylindric-campanulate, its teeth about half as long as the tube, acute, triangular, rusty-tomentose and scurfy externally. *Petals* oblong, obtuse, very little narrowed to the base and without any very distinct claw. *Ovaries* 3 to 5, narrowly ovoid, compressed, vertically ridged, lepidote as are the conjoined styles. *Fruit* sub-globose, much compressed, covered with a layer of very minute pale hairs, .75 in. in diam.

Perak, at low elevations and in moist ground; Scortechini, Wray, King's Collector.

2 PENTACE, Hassk.

Trees. Herbaceous portions sometimes pubescent or scaly, ultimately glabrous. *Leaves* entire, leathery, the lower surface (except in one species) pale from a thin layer of minute adpressed scaly hair. *Flowers* numerous, small, in terminal panicles. *Calyx* bell-shaped, usually 5-fid. *Petals* 5, membranous, glabrous, longer than the calyx, glandless. *Stamens* numerous, on a slightly raised torus, usually pentadelphous. *Staminodes* 5, opposite the sepals. *Anthers* subglobose; pollen globose, 3-pored. *Ovary* 5-celled, cells 2-ovuled; ovule pendulous, raphe next the placenta. *Styles* united, rarely free. *Fruit* dry, indehiscent, 3-10-winged, 1-celled, 1-seeded by abortion. *Seed* solitary, albuminous.—Distrib. About 15 species, all Malayan.

- Leaves with pinnate nervation.
- Ovary 3-ridged... 1. *P. triptera*.
- Ovary 5-ridged.
- Leaves with 6 or 7 pairs of nerves... 2. *P. Hookeriana*.
- " " 3 or 4 " ... 3. *P. Kunstleri*.
- Ovary 10-ridged 4. *P. perakensis*.
- Leaves boldly 5-nerved at the base, lateral nerves from the central nerve (midrib) 2 pairs; 7 to 14 in. long; ovary 5-ridged... 5. *P. macrophylla*.
- Leaves boldly 8-nerved at the base; the central nerve (midrib) with 1 or 2 pairs of lateral nerves: rarely more than 7 in. long.
- Ovary not visibly ridged 6. *P. floribunda*.
- Ovary 8 or 9-ridged.
- Styles quite confluent 7. *P. Curtisii*.
- " free.. 8. *P. eximia*.
- Ovary 10-ridged.
- Leaves glabrous on both surfaces... 9. *P. Scortechinii*.
- " with a dense layer of minute adpressed hair on the under surface 10. *P. Griffithii*.
- Leaves boldly 3-nerved at the base, the central nerve (midrib) without lateral nerves, only 3 or 4 in. long; ovary 5-ridged 11. *P. strychnoidea*.

1. PENTACE TRIPTERA, Mast. in Hook. fil. Fl. Br. Ind. i. 382. A large tree: young branches pubescent, speedily becoming glabrous, their bark dark-coloured. *Leaves* ovate to ovate-rotund, sometimes ovate-oblong, sub-acute or shortly and bluntly acuminate, the margins undulate, the base rounded; upper surface glabrous, the lower pale, minutely scaly; basal nerves one or two pairs; lateral 5 to 7 pairs, ascending, straight; length 4 to 5 in., breadth 2 to 2.75 in.; petiole .6 to 1.2 in. thickened towards the apex, pubescent. *Panicles* terminal and axillary, 6 to 8 in. long, with short many-flowered branches minutely and softly stellate-tomentose. *Flowers* nearly .2 in. long, on pedicels shorter than the calyx. *Calyx* with 5 lanceolate teeth, tomentose outside. *Petals* spatulate-oblong, obtuse. *Stamens* 5-delphous, longer than the style. *Staminodes* subulate, shorter than the stamens. *Ovary* densely tomentose, shortly 3-winged. *Style* filiform, glabrous, bent at the apex. *Fruit* oblong, narrow, .6 in. long, with 3 spreading membranous rounded wings .5 in. broad.

Malacca: Griffith, Maingay. Perak: Scortechini.

This approaches the Javan *P. polyantha*, Hassk., which has, however, larger flowers with a shallower calyx with longer teeth, a shorter style, and a 5-lobed ovary.

2. *PENTACE HOOKERIANA*, n. sp., King. A tree 30 to 40 feet high: young branches cinereous, glabrous. *Leaves* elliptic-oblong, slightly obovate, acute, the base narrowed and slightly unequal; upper surface glabrous, the lower dull; lateral main nerves about 6 pairs (one of the pairs basal), prominent on both surfaces; the intermediate nerves, transverse veins and reticulations prominent only on the lower: length 5 to 7.5 in.; breadth 2 to 2.75 in.; petiole .25 in., stout. *Panicles* terminal and in the axils of the upper leaves, 2.5 to 5 in. long, the branches spreading, everywhere scurfy-tomentose. *Flowers* rather crowded, .2 in. long, on pedicels shorter than the calyx. *Calyx* campanulate, cut half-way down into 3 or 4 broadly triangular sub-acute spreading teeth, scaly and minutely tomentose outside. *Petals* narrowly obovate. *Stamens* 15 in 5 bundles of 3 each, very much shorter than the petals. *Staminodes* thick, orbicular, embracing the ovary. *Ovary* depressed-globose, densely pubescent, obscurely 5-lobed, 5-celled. *Styles* 5, free, shorter than the ovary. *Fruit* unknown.

Perak, on the banks of the Kinta river: King's Collector, No. 815.

3. *PENTACE KUNTLERI*, n. sp., King. A tree 30 to 40 feet high: young branches slender, dark-coloured, glabrous. *Leaves* broadly ovate, with an abrupt short broad blunt acumen, the base rounded: upper surface shining, glabrous, the lower dull; lateral nerves 3 or 4 pairs, curved, prominent beneath; sometimes a pair of short slender sub-marginal nerves at the base: length 4.5 to 6 in., breadth 2.5 to 3.5 in.; petiole .75 to 1 in., stout, thickened at the apex. *Panicles* terminal, 3.5 to 6 in. long, puberulous, much-branched. *Flowers* numerous, .15 in. long, the pedicels slightly shorter. *Calyx* tubular-campanulate, minutely stellate-hairy and lepidote outside; the teeth triangular, acute, erect. *Petals* spatulate with a very long claw. *Stamens* in 5 bundles. *Staminodes* linear-lanceolate, as long as the filaments. *Ovary* depressed-globose, with 5 blunt angles, lepidote and pubescent, 5-celled. *Style* straight, glabrous. *Fruit* unknown.

Perak, at a very low elevation; King's Collector, No. 6871.

4. *PENTACE PERAKENSIS*, n. sp., King. A tree 30 to 40 feet high: young branches cinereous, glabrous. *Leaves* ovate-elliptic, slightly oblique, bluntly acuminate, the base rounded or sub-cuneate: upper surface shining, glabrous: the lower dull; lateral nerves about 5 pairs,

ascending, curved; length 5 to 6 in., breadth 2.5 to 3 in.; petiole .75 to 1 in., stout, and thickened at the apex. *Panicles* terminal, 4 to 5 in. long and less than 2 in. wide, little branched and few-flowered. *Flowers* about .1 in. long, their pedicels about as long. *Calyx* rotate, minutely lepidote outside; the teeth triangular, spreading. *Petals* ovate, narrowed to a short claw. *Stamens* about 30, 5-delphous. *Staminodes* lanceolate, as long as the filaments. *Ovary* globose, slightly pointed, 10-ridged, slightly hairy, 5-celled. *Style* about as long as the ovary, cylindrical. *Stigma* terminal, small. *Fruit* unknown.

Perak, King's Collector, No. 3428.

5. *PENTACE MACROPHYLLA*, n. sp., King. A tree usually from 20 to 30 feet high, but occasionally as much as 50 feet. Young branches rather slender, pale brown, glabrous. *Leaves* large, ovate-elliptic to almost rotund, the apex very shortly and abruptly blunt-acuminate, the base rounded: upper surface glabrous; the lower dull; basal nerves 2 pairs, the upper branched on one side; lateral nerves from the midrib 2 to 3 pairs, all ascending and little curved, prominent beneath; length 7 to 14 in., breadth 5 to 12 in.; petiole 2.5 to 3 in., stout. *Panicles* terminal and axillary, 6 to 15 in. long, lax, spreading, minutely yellowish-pubescent and scurfy. *Flowers* .15 in. long and .2 in. in diam., on pedicels about as long as the calyx. *Calyx* almost rotate, cut for two-thirds of its length into 5 lanceolate acute teeth, minutely yellowish-tomentose outside. *Petals* oblanceolate, obtuse. *Staminodes* linear, as long as the stamens. *Stamens* in 5 bundles of about 15 each. *Ovary* ovoid, scaly and pubescent, obtusely 5-angled. *Style* rather shorter than the stamens, cylindrical, pointed. *Fruit* .75 in. long with 10 radiating semi-elliptic striate sparsely scaly wings each .4 in. wide.

Perak at elevations up to 500 feet; King's Collector, Scortechini, Wray: common.

Distinguished from all the other known species by the large size of its leaves.

6. *PENTACE FLORIBUNDA*, n. sp., King. A tree 40 to 70 feet high; young branches slender, sparsely stellate-puberulous, the bark dark-coloured. *Leaves* elliptic-oblong to elliptic-rotund, the apex shortly and rather abruptly apiculate; the base rounded or slightly narrowed; upper surface glabrous, lower cinereous and with some scattered pubescence; basal nerves 2 pairs, one of them branching on one side; lateral nerves 2 or 3 pairs, all ascending and all rather prominent: transverse veins not prominent; length 5 to 6.5 in., breadth 2.5 to 3.75 in., petiole 1 to 2 in., thickened towards the apex. *Panicles* towards the apices of the

branches, axillary and terminal, stellate-pubescent, slender, spreading, many-branched. *Flowers* very numerous, $\frac{1}{2}$ in. long, the pedicels slender and rather longer. *Calyx* when expanded rotate, cut half way down into triangular very acute or acuminate spreading teeth, densely stellate-tomentose outside. *Petals* broadly oblanceolate, obtuse, narrowed to the base. *Stamens* 5-delphous. *Staminodes* apparently absent. *Ovary* globose, densely tomentose, not visibly ridged, 5-celled. *Style* filiform, tapering, straight, glabrous. *Fruit* unknown.

Perak, at elevations from 600 to 1000 feet: King's Collector, Nos. 7616 and 7730.

A species distinguished by its slender hoary panicles, with flowers by far more numerous than in any of the other species described here.

7. *PENTACE CURTISII*, n. sp. King. A large tree: the young branches slender, with dark-coloured bark, very minutely adpressed-lepidote, not hairy. *Leaves* ovate elliptic, with a short abrupt blunt acumen, the base rounded: upper surface glabrous, the lower cinereous; basal nerves 2 pairs, one pair slender and close to the margin, the other branching on one side: lateral nerves 2 or 3 pairs; all ascending and rather prominent beneath: length 3.25 to 5.5 in., breadth 2.25 to 2.75 in.; petiole .75 to 1 in. slender, slightly thickened at apex. *Panicles* mostly terminal (a few smaller axillary) 4.5 to 6 in. long with sparse cinereous stellate tomentum and scales, few-branched, and few-flowered. *Flowers* .15 in. long, on pedicels shorter than themselves. *Calyx* widely campanulate, stellate-tomentose outside; its teeth as long as the tube, broadly triangular, rather blunt. *Petals* oblanceolate or obovate-obtuse, much narrowed to the base. *Staminodes* lanceolate, acuminate, half as long as the filaments. *Stamens* in 5 bundles. *Ovary* turbinate, with 8 or 9 blunt ridges, lepidote-pubescent: style rather stout, cylindrical, shorter than the stamens. *Fruit* .5 in. long, with 8 semi-elliptic membranous wings .2 in. broad.

Penang; Curtis, No. 1573.

8. *PENTACE EXIMIA*, n. sp., King. A tree 50 to 70 feet high: young branches slender, dark-coloured, glabrous. *Leaves* ovate-elliptic to ovate-rotund, shortly and abruptly acuminate, the base slightly narrowed or rounded: upper surface glabrous, shining: lower paler and dull; basal nerves 1 pair, bold and reaching to the apex, often with a slender small sub-marginal pair: lateral nerves usually only one pair, short and curving; all rather bold beneath: length 4 to 5.5 in., breadth 2 to 4 in.; petiole .75 to 1.1 in., thickened at the apex. *Panicles* terminal, 3.5 to 5 in. long, (longer in fruit), minutely scurfy-tomentose, with

rather numerous spreading branches. *Flowers* numerous, 1 in. long, the pedicels about the same length. *Calyx* densely scaly outside, the teeth triangular. *Petals* cuneate, obtuse, narrowed to a broad claw. *Stamens* about 25, in groups of 5. *Staminodes* lanceolate. *Ovary* sub-globular, 10-ridged (the ridges in pairs), scaly and pubescent, 5-celled, the cells with imperfect septa and thus falsely 10-celled. *Styles* 10, much shorter than the stamens, free, or united when young at the base only. *Fruit* about 5 in. long, with 8 radiating semi-elliptic wings 1 to 1.5 in. broad, minutely adpressed-scaly.

Perak; at elevations under 1000 feet, King's Collector, Nos. 3482 and 3649.

This agrees with *P. Curtisii* in having 8-winged fruit, but the flowers are much smaller. The styles moreover are shorter than the ovary and quite distinct, which is the case in no other species of this genus which I have yet met with.

9. *PENTACE SCORTECHINII*, n. sp. King. A tree? young branches slender, glabrous, dark-coloured. *Leaves* elliptic-oblong, shortly caudate-acuminate, the base more or less cuneate: both surfaces quite glabrous, concolorous; basal nerves 1 pair very bold, as is the midrib; lateral nerves (from the midrib) 1 or 2 pairs, not conspicuous: length 7 to 9 in., breadth 2.75 to 3.5 in.; petiole less than .5 in., stout. *Panicles* terminal and axillary, slender, only about half the length of the leaves, few-branched, minutely tomentose. *Flowers* rather crowded, .25 in. long, on pedicels shorter than the calyx. *Calyx* widely tubular-campanulate, minutely scurfy-tomentose outside, cut a third of its depth into 5 small triangular reflexed teeth. *Petals* obovate, obtuse, much narrowed to the base. *Stamens* in 5 bundles of 15 each. *Staminodes* lanceolate, half as long as the stamens. *Ovary* ovoid, obscurely 5-ridged, scaly, 5-celled. *Style* cylindric, tapering, longer than the stamens. *Fruit* unknown.

Perak, Father Scortechini, No. 119b.

Only once collected and without fruit. A very distinct species.

10. *PENTACE GRIFFITHII*, n. sp., King. A tree: young branches slender, dark-coloured, glabrous. *Leaves* ovate-elliptic, tapering about equally to the acute apex and base; upper surface shining, lower dull; basal nerves 2 pairs, the lower pair slender and sub-marginal, the upper branched on one side and bold (as is the midrib), ascending, curved; lateral nerves (from the midrib) 2 pairs; length 4 to 7 in., breadth 2.25 to 3 in.; petiole nearly 1.5 in. long, thickened at both ends, but especially at the apex. *Panicles* terminal, slender, few-branched, lax, minutely

cinereous-tomentose. *Flowers* not very numerous, large for the genus (.25 in. long and .25 in. in diam.), on pedicels about as long as the calyx. *Calyx* widely campanulate, almost rotate, minutely stellate-tomentose outside; the teeth as long as the tube, spreading. *Petals* ovate, obtuse, rather suddenly contracted into a linear claw. *Stamens* in 5 groups of 12 or 13 each. *Staminodes* lanceolate, as long as the filaments. *Ovary* sub-globose, slightly compressed, minutely stellate-tomentose and scaly, obtusely 5-angled, 5-celled. *Style* cylindric, rather shorter than the filaments. *Fruit* nearly 1 in. long, with 10 radiating membranous, horizontal striate, minutely scaly, semi-elliptic, membranous wings, each .35 in. broad.

Tavoy in Tenasserim; Griffith, Aplin.

A very distinct species only once collected within recent years, by Mr. Aplin. There is, however, in the Kew Herbarium a twig of it collected by Griffith many years ago bearing this note in Griffith's handwriting "*Tiliacearum gen. nov. capsulis pluri-alatis.*" Although this plant has hitherto been found only in territory which is politically Burmese, yet Tavoy (being at the southern extremity of Tenasserim) is practically Malayan in its Flora and Fauna. I therefore include it here.

11. PENTACE STRYCHNOIDEA, n. sp., King. A tree 60 to 80 feet high: young branches slender, cinereous, glabrous. *Leaves* ovate-elliptic rarely ovate-oblong, shortly and abruptly acuminate, the base rounded or slightly narrowed; upper surface shining, glabrous; lower pale and dull; boldly 3-nerved and often with a slender sub-marginal pair of nerves; length 3 to 4 in., breadth 1.75 to 2.25 in.; petiole .75 in. slightly thickened at the apex. *Panicles* terminal, 3 to 6 in. long, few-branched, lax, minutely lepidote-puberulous. *Flowers* rather large for the genus (.2 in. long). *Calyx* cup-shaped, tomentose outside, cut more than half way down into 5 triangular acute teeth. *Petals* oblanceolate, slightly oblique, much narrowed to the base. *Stamens* in 5 bundles of about 20 each. *Staminodes* linear-lanceolate. *Ovary* ovoid-globose, obtusely 5-ridged, minutely tomentose and lepidote, 5-celled. *Style* filiform, as long as the stamens. *Fruit* unknown.

Perak; at elevations of from 500 to 1000 feet, King's Collector, No. 3478.

3. SCHOUTENIA, Korth.

Trees with alternate simple pinnately-nerved leaves. *Flowers* axillary, solitary or in clusters; or in terminal few-flowered panicles. *Calyx* campanulate, 5-lobed; lobes valvate, accrescent, coloured. *Petals*

cinereous-tomentose. *Flowers* not very numerous, large for the genus (.25 in. long and .25 in. in diam.), on pedicels about as long as the calyx. *Calyx* widely campanulate, almost rotate, minutely stellate-tomentose outside; the teeth as long as the tube, spreading. *Petals* ovate, obtuse, rather suddenly contracted into a linear claw. *Stamens* in 5 groups of 12 or 13 each. *Staminodes* lanceolate, as long as the filaments. *Ovary* sub-globose, slightly compressed, minutely stellate-tomentose and scaly, obtusely 5-angled, 5-celled. *Style* cylindric, rather shorter than the filaments. *Fruit* nearly 1 in. long, with 10 radiating membranous, horizontal striate, minutely scaly, semi-elliptic, membranous wings, each .35 in. broad.

Tavoy in Tenasserim; Griffith, Aplin.

A very distinct species only once collected within recent years, by Mr. Aplin. There is, however, in the Kew Herbarium a twig of it collected by Griffith many years ago bearing this note in Griffith's handwriting "*Tiliacearum gen. nov. capsulis pluri-alatis.*" Although this plant has hitherto been found only in territory which is politically Burmese, yet Tavoy (being at the southern extremity of Tenasserim) is practically Malayan in its Flora and Fauna. I therefore include it here.

11. PENTACE STRYCHNOIDEA, n. sp., King. A tree 60 to 80 feet high: young branches slender, cinereous, glabrous. *Leaves* ovate-elliptic rarely ovate-oblong, shortly and abruptly acuminate, the base rounded or slightly narrowed; upper surface shining, glabrous; lower pale and dull; boldly 3-nerved and often with a slender sub-marginal pair of nerves; length 3 to 4 in., breadth 1.75 to 2.25 in.; petiole .75 in. slightly thickened at the apex. *Panicles* terminal, 3 to 6 in. long, few-branched, lax, minutely lepidote-puberulous. *Flowers* rather large for the genus (.2 in. long). *Calyx* cup-shaped, tomentose outside, cut more than half way down into 5 triangular acute teeth. *Petals* oblanceolate, slightly oblique, much narrowed to the base. *Stamens* in 5 bundles of about 20 each. *Staminodes* linear-lanceolate. *Ovary* ovoid-globose, obtusely 5-ridged, minutely tomentose and lepidote, 5-celled. *Style* filiform, as long as the stamens. *Fruit* unknown.

Perak; at elevations of from 500 to 1000 feet, King's Collector, No. 3478.

3. SCHOUTENIA, Korth.

Trees with alternate simple pinnately-nerved leaves. *Flowers* axillary, solitary or in clusters; or in terminal few-flowered panicles. *Calyx* campanulate, 5-lobed; lobes valvate, accrescent, coloured. *Petals*

small, linear without claw, or absent. *Stamens* numerous, free, sometimes inserted on the apex of a short gynophore; anthers oblong, 2-celled: cells parallel, with longitudinal sutural dehiscence. *Staminodes* 0. *Ovary* sessile or shortly stalked, imperfectly 3 to 5-celled; cells with 2 ovules from the base of the axile placentas, style filiform; stigmas 3 to 5, linear fleshy, reflexed. *Capsule* with crustaceous fragile pericarp, dehiscing irregularly, 1-celled (by abortion), 1- to 3-seeded. *Seeds* subglobose, with leathery smooth testa, exalbuminous: the cotyledons large, leafy, thin, crumpled: embryo straight. *Distrib.* 5 species, of which 4 are Malayan and 1 Cambodian.

Flowers in panicles or solitary, axillary.

- | | |
|---|------------------------------|
| Calyx very accrescent very deeply lobed | 1. <i>S. Mastersii</i> . |
| „ slightly accrescent not deeply lobed | 2. <i>S. Kunstleri</i> . |
| Flowers in dense axillary glomeruli | ... 3. <i>S. glomerata</i> . |

1. *SCHOUTENIA MASTERSII*, King. A tree 60 to 80 feet high: young branches slender, dark-coloured, at first scaly but soon glabrous. *Leaves* thinly coriaceous, ovate-lanceolate, slightly obovate, shortly and bluntly acuminate, the base rounded; upper surface glabrous, the lower minutely and softly tawny-tomentose; nerves slightly prominent beneath, about 3 pairs lateral and 1 pair basal: length .75 to 3.25 in.; breadth .4 to 1.1 in.; petiole less than .1 in. *Flowers* solitary and axillary, or in terminal leafy panicles; the pedicels from .35 to .75 in. according to age, tawny-tomentose, jointed below the middle. *Calyx* membranous, pink, conspicuously veined, at first widely campanulate, .35 in. long, with 5 shallow teeth becoming with the ripening of the fruit, rotate, flat 1.5 to 2 in. in diam., and 5-angled; pubescent outside, glabrous within. *Filaments* very slender, longer than the style. *Ovary* obovoid-globose, tawny-tomentose. *Style* stout, three times as long as the ovary, tomentose: stigmas scaly. *Fruit* depressed-globose, .3 in. in diam., minutely tomentose. *Chartacalyx accrescens*, Mast. in Hook. fil. Fl. Br. Ind. i. 382.

Malacca, Penang, Perak. *Distrib.* Borneo.

On this plant the late Dr. Maingay founded his genus *Chartacalyx*. The only points, however, in which it differs from *Schoutenia* (as defined by Bentham and Hooker) are the absence of petals and the presence of a stalk to the ovary on the upper part of which the stamens are inserted; and these appear to me to be, in this order, differences of quite minor importance. Maingay never saw the fruit of this; but copious fruiting specimens have recently been collected and the fruit is found to be exactly that of *Schoutenia*. As regards the structure of the seeds of

Schoutenia, Korthals (the author of the genus) says nothing: nor does Bennet who (Pl. Jav. Rar. p. 239, t. 46) describes at greater length than Korthals the species *S. ovata*, the only one then known. Bennett neither describes nor figures albumen in the seed. Hasskarl (Retzia 1, 136) describes the seeds as exalbuminous, and I find none in the seeds of these species of which I have been able to examine ripe fruit. The only other known species are *S. ovata*, Korth. from Java; and *S. hypoleuca*, Pierre (Fl. Cochinchine t. 134) from Cambodia.

2. *SCHOUTENIA KUNSTLERI*, n. sp., King. A tree 60 to 70 feet high: young branches cinereous, rather rough-glabrous. *Leaves* thinly coriaceous, narrowly obovate-oblong or oblanceolate, acute, the margin slightly wavy, slightly narrowed to the rounded 3- to 5-nerved base; upper surface glabrous, shining: lower sub-silvery; the lateral nerves 4 or 5 pairs, spreading, curving, inter-arching near the margin, prominent on the lower surface as are the basal nerves and the numerous slightly curved transverse veins. *Flowers* crowded towards the ends of the branches, in numerous short few-flowered scurfy-tomentose racemes or cymes: pedicels from .5 to .75 in. long, jointed and bracteolate above the base, the bracteole oblanceolate. *Calyx* campanulate, membranous, coloured and veined, stellate-hairy on both surfaces, .5 to .75 in. long, according to age, cut to the base into 5 ovate spreading lobes. *Petals* 0. *Stamens* on a slightly elevated torus. *Ovary* sessile, sub-globose, densely tomentose, 5-celled. *Style* longer than the stamens. *Stigmas* 5, short, fleshy. *Fruit* 1-celled, 1-seeded, surrounded by the slightly accrescent persistent calyx.

Perak at elevations of from 300 to 800 feet: King's Collector, No. 3409: on Ulu Tupa, Wray, No. 2692.

According to the field notes of Messrs. Kunstler and Wray, the calyx is yellow when young, but becomes brown when the fruit ripens.

3. *SCHOUTENIA GLOMERATA*, n. sp., King. A tree from 40 to 60 feet high: young branches slender, cinereous, minutely pubescent. *Leaves* membranous, glabrous, elliptic-oblong, acute or shortly and bluntly acuminate, the margins slightly wavy; the base broad, rounded or emarginate, 3-nerved, the upper pair of nerves very strong, running to the apex of the leaf and joined to the midrib by numerous prominent curving transverse secondary nerves, all very prominent on the pale silvery shining under surface: length 10 to 15 in., breadth 3.5 to 5.5 in.; petiole only .25 in. long, stout, wrinkled. *Cymes* condensed, very crowded, axillary, 1 to 1.5 in. in diam. *Flowers* .25 in. long and .3 in. wide, on tomentose rufous pedicels about .2 in. long. *Calyx* widely

campanulate, densely rufous-tomentose; teeth 5, broadly triangular, sub-erect. *Petals* 0. *Stamens* numerous; the filaments slender, longer than the calyx. *Ovary* ovoid-globose, densely tawny-tomentose, 5-celled: style longer than the stamens; stigmas short, sub-globose. *Fruit* depressed globose, .75 in. in diam., sparsely stellate-tomentose, becoming glabrous, covered only at the base by the slightly accrescent calyx.

Johore; on Gunong Panti, King's Collector, No. 159.

4. BERRYA, Roxb.

A tree. *Leaves* alternate, ovate, acuminate, glabrous; base cordate, 5-7-nerved. *Panicles* large, many-flowered, terminal and axillary. *Calyx* campanulate, irregularly 3-5-lobed. *Petals* 5, spatulate. *Stamens* many, inserted on a short torus; anthers didymous, lobes divergent, opening lengthwise. *Staminodes* 0. *Ovary* 3-4-lobed, cells 4-ovuled; style consolidated, stigma lobed; ovules horizontal. *Fruit* loculicidally 3-4-valved, each valve 2-winged. *Seeds* pilose; albumen fleshy; cotyledons flat leafy, radicle superior next the hilum.—Distrib. The following is the only species.

BERRYA AMMONILLA, Roxb. Hort. Beng., 42. A large tree, glabrous except the inflorescence. *Leaves* membranous, broadly ovate, acuminate, the base slightly narrowed and cordate: both surfaces shining, minutely reticulate: basal nerves 2 or 3 pairs, lateral 5 or 6 pairs: length 4 to 8 in., breadth 3 to 5 in.; petiole .75 to 2.75 in. *Panicles* terminal, or in the upper axils, branching, 6 to 10 in. long, scurfy-pubescent: flowers .35 in. in diam.; their pedicels slender, .3 to .5 in. long. *Petals* longer than the calyx, narrowly oblong, obtuse, glabrous. *Anthers* half as long as the petals. *Ovary* ovoid, truncate, depressed at the origin of the styles, 6 to 8-ridged, pubescent. *Fruit* with 6 radiating, falcate, membranous, striate, deciduously stellate-tomentose wings .8 in. long. *Seeds* small: 1 to 4 in. each cell, covered with prurient pale brown hairs. Roxb. Fl. Ind. ii. 639; Corom. Plants, ii. t. 264; Wall. Cat. 1068; W. & A. Prodr. i. 81; Wight Ill. t. 34; Thwaites Enum. 32; Beddome Flor. Sylvat. t. 58; Kurz Fl. Burm. i. 155; Hook. fil. Fl. Br. Ind. i. 383.

South Andamans. Distrib. Burmah, Southern Peninsula, India, Ceylon.

5. GREWIA, Lindl.

Trees or shrubs more or less stellate-pubescent. *Leaves* entire, 1-9-nerved. *Flowers* axillary and few, or more numerous and panicked. *Sepals* distinct. *Petals* 5, glandular at the base, sometimes 0. *Stamens*

many on a raised torus. *Staminodes* 0. *Ovary* 2-4-celled, cells opposite the petals, 2-many-ovuled; style subulate, stigma shortly lobed. *Drupe* fleshy or fibrous, entire, or 2-4-lobed; stones 1-4, 1-2-seeded, with false partitions between the seeds. *Seeds* ascending; albumen fleshy or rarely 0; cotyledons flat. *Distrib.* About 60 species, chiefly tropical.

Sect. I. *Grewia* proper. *Flowers* axillary or terminal. *Fruit* fleshy or crustaceous, usually lobed

1. *G. umbellata.*

Sect. II. *Microcos.* *Inflorescence* terminal, in paniced cymes. *Flowers* involucrate. *Drupe* fleshy, entire

2. *G. paniculata.*

Sect. III. *Omphacarpus.* *Inflorescence* terminal, or terminal and axillary. *Flowers* involucrate. *Drupe* with a corky or fibrous rind.

Fruit minutely tomentose: mesocarp thick, soft, pulpy, and with many fibres; pyrene single, small.

Pyrene membranous: leaves softly tomentose beneath

3. *G. fibrocarpa.*

Pyrene cartilaginous: leaves sparsely stellate-hispid beneath

4. *G. globulifera.*

Fruit glabrous: mesocarp with thin pulp and a few fibres: pyrenes 2 or 3, bony.

Leaves sparsely-stellate pubescent beneath: drupe not narrowed into a pseudo-stalk...

5. *G. latifolia.*

Leaves glabrescent or pubescent beneath: drupe narrowed into a long pseudo-stalk...

6. *G. antidesmaefolia.*

Leaves quite glabrous.

Basal nerves bold and reaching beyond the middle.

Fruit .5 in. long, furrowed, not compressed

7. *G. laurifolia.*

Fruit 1.4 in. long, not furrowed, compressed

8. *G. calophylla.*

Basal nerves slender, not reaching to the middle: drupe .75 in. long

9. *G. Miqueliana.*

1. *GREWIA UMBELLATA*, Roxb. Hort. Beng. 42: Fl. Ind. ii. 591. A shrubby climber 10 to 20 feet long; whole plant except the upper surfaces of the leaves sparsely stellate-puberulous, the bark of the young

branches dark-coloured. *Leaves* oblong-ovate or elliptic, shortly and bluntly acuminate, minutely serrate; base rounded, 3-nerved; upper surface glabrous; the lower pale with the transverse veins prominent and straight: lateral nerves about 3 pairs: length 3 to 4.5 in., breadth 1.5 to 2 in., petiole .25 in. *Umbels* pedunculate, axillary or terminal, 6 to 8-flowered; the peduncle from .6 to 1 in. long, with a whorl of small lanceolate glabrous bracteoles at its apex. *Flowers* .75 in. long when expanded; their pedicels hirsute, unequal, from .2 to .5 in. long. *Sepals* ribbed and tomentose outside, glabrous inside, linear-oblong, reflexed. *Petals* much shorter than the sepals, oblong, each springing from the back of a large orbicular claw with hirsute edges. *Torus* long, ridged, tomentose. *Fruit* depressed-globular, obtusely 2- to 4-angled and with 2 to 4 shallow lobes, pericarp sparsely stellate-puberulous; endocarp pulpy; pyrene 2 to 4-celled; its loculi 1-seeded, the endocarp bony. Wight Ic. 83; Wall. Cat. 1084; Mast. in Hook. fil. Fl. Br. Ind. i. 385.

Malacca, Penang, Griffith, Maingay. Perak, King's Collector, Wray.

Roxburgh has left an excellent coloured drawing of this in the library of the Calcutta Herbarium, and there is no doubt about his plant. I cannot agree in identifying with this *G. pedicellata*, Roxb., which that author received from Amboyna: nor do I think that any *Grewia* from the Peninsula of Hindustan is referable to this species

2. *GREWIA PANICULATA*, Roxb. Fl. Ind. ii. 591. A bushy tree 15 to 30 feet high: young branches scurfy stellate-tomentose, ultimately glabrous, their bark brown. *Leaves* coriaceous, cuneate-obovate to elliptic; the apex blunt, shortly and abruptly acuminate, sometimes 3-lobed and unequal, obscurely serrate-dentate; the base rounded, 3-nerved; upper surface powdered with minute sparse stellate pubescence, the midrib and nerves tomentose: lower surface uniformly stellate-tomentose; the veins transverse, little curved, bold; lateral nerves 4 or 5 pairs, ascending, rather straight, prominent beneath: length 3 to 6 in., breadth 1.5 to 2.75 in.; petiole .25 in., tomentose: stipules glabrescent, lanceolate, often united in pairs, rather shorter than the petioles. *Panicles* 2.25 to 3.5 in. long, terminal or axillary, rusty-tomentose; bracteoles numerous, linear, sometimes bifid: branches spreading. *Flowers* .25 in. long, the pedicels rather shorter. *Sepals* spreading, concave, obovate narrowed to the base, the edges thin; tomentose on the outer, pilose on the inner, surface. *Petals* shorter than the sepals, oblong, blunt, expanded at the base into a concave claw, hirsute especially outside. *Torus* cup-shaped, short, the lip tomentose. *Ovary* ovoid, stellate-tomentose, 4-celled, each cell with several ovules. *Fruit* ob-

ovoid, recurved, with many curved striae, pericarp membranous, minutely and sparsely stellate-pubescent, the mesocarp fibrous with an outer layer of pulp: pyrene 1-celled, 1-seeded; endocarp stony. Wall. Cat. 1097, partly; Miq. Fl. Ind. Bat. i. pt. 2, 203; Mast. in Hook. fil. Fl. Br. Ind. i. 393. *G. Blumei*, Hassk. Tijdschr. Nat. Gesch. xii. 130; Miq. Fl. Ind. Bat. i. pt. 2, 203. *Microcos tomentosa*, Smith in Rees, Cycl. *G. affinis*, Hassk. Cat. Hort. Bog. 207, not of Lindl.

Singapore; Malacca, Maingay, No. 250. Griffith, No. 634 (Kew Distrib.). Perak. Penang; common.

I retain for this plant the name adopted for it by Masters in Hooker's Flora of British India. But Blume's *G. paniculata* (Bijdr. 115) was published seven years before Roxburgh's. I have not seen any specimen of Blume's plant: but if it be the same as this, then Blume's name must be substituted for that of Roxburgh as the author of the specific name. If Blume's plant, however, be different from Roxburgh's, then some other name must be found for the latter. That the plant above described is what Roxburgh meant to call *G. paniculata*, his coloured drawing in the Calcutta Herbarium leaves no room for doubt.

3. *GREWIA FIBROCARPA*, Mast. in Hook. fil. Fl. Br. Ind. i. 391. A tree 15 to 40 feet high; young branches, under surfaces of leaves, petioles, inflorescence and fruit densely clothed with yellowish-brown stellate tomentum. *Leaves* membranous, ovate-oblong or elliptic, shortly and abruptly acuminate, minutely and obscurely serrulate, the base rounded and boldly 3-nerved; upper surface scaberulous, the midrib and nerves tomentose, under surface softly tomentose; the 5 to 7 pairs of lateral nerves and the transverse veins rather prominent beneath: length 4.5 to 9 in., breadth 1.75 to 4 in., petiole .25 to .5 in., stout: stipules deeply and narrowly lobed. *Panicles* terminal and in the upper axils, crowded, .5 to 2 in. long: involucre lanceolate, curved, tomentose. *Flowers* .25 in. long, their pedicels much shorter. *Sepals* obovate-elliptic, very tomentose externally, the edges inflexed, sparsely pilose internally. *Petals* minute, sub-orbicular, sometimes absent. *Torus* short, hirsute. *Ovary* ovoid-globose, tomentose; the style short, conical, glabrous. *Fruit* soft, ovoid or obovoid, compressed, 1.25 in. long and .75 in. in diam., the pericarp membranous and densely tomentose outside, mesocarp fibrous and pulpy; pyrene small, solitary, leathery, 1-celled, 1-seeded. *G. paniculata*, Wall. (Cat. No. 1097 partly) not of Roxb.

Penang; Wallich, Curtis. Malacca; Griffith; Maingay, No. 248, (Kew Distrib.). Perak; Scortechini, King's Collector, Wray. Common.

In the fruit both of this and of *G. globulifera*, the mesocarp forms a thick pulp with many fibres intermixed, and the solitary pyrene is small with a soft coat.

4. *GREWIA GLOBULIFERA*, Mast. in Hook. fil. Fl. Br. Ind. i. 391. A small shrubby tree; young branches densely covered with short yellowish-brown tomentum. *Leaves* thinly coriaceous, broadly elliptic, sometimes slightly obovate and unequal-sided, shortly and abruptly acuminate, entire, the base rounded, boldly 3-nerved: upper surface scaberulous, glabrous except the minutely tomentose midrib and nerves: under surface shortly and sparsely stellate-hispid: main nerves 7 to 8 pairs, spreading, prominent beneath, the transverse nerves rather thin, the reticulations minute but distinct: length 4.5 to 10 in., breadth 3 to 6 in.; petiole .4 to .75 in., tomentose. *Panicles* often on long peduncles, axillary and terminal, narrow, few-flowered, covered with soft yellowish stellate tomentum: length 2.5 to 4.5 in. (of which the peduncle may be more than half). *Flowers* .35 in. long, their pedicels much shorter. *Sepals* oblong, spreading, curved inwards, tomentose on both surfaces, the edges much incurved. *Petals* much shorter than the sepals, glabrous, linear-lanceolate, without any distinct claw but sometimes more or less thickened and hairy at the base. *Torus* a very shallow cup with hirsute edge. *Ovary* ovoid, pointed, tomentose; style as long as the ovary, cylindric, glabrous. *Fruit* usually solitary at the apex of a branch of the panicle, sub-obovoid, compressed, 1.25 in. long and .65 in. in diam.; pericarp membranous minutely tomentose, the mesocarp pulpy and very fibrous; the single pyrene much smaller, endocarp cartilaginous, 1-celled, 1-seeded.

Malacca; Griffith, No. 635; Maingay, No. 245, (Kew Distrib.); Harvey. Perak; Scortechini, King's Collector, Wray: at low elevations.

In its fruit this much resembles *G. fibrocarpa*. The drupe, however, of this is obovoid not ovoid, and the stone is larger with cartilaginous not membranous endocarp. The leaves also differ in being sparsely shortly hispid-pubescent instead of softly tomentose. A near ally of this species is also *G. latifolia*, Mast. from which this differs in its petals having no distinct claw, whereas in those of *G. latifolia* the claw is larger than the limb. This also differs in the shape of its ovary and style, and in the very different appearance of its drupe.

5. *GREWIA LATIFOLIA*, Mast. in Hook. fil. Fl. Br. Ind. i. 392. A shrubby tree 20 to 40 feet high: young branches rather stout, minutely but harshly tawny-or cinereous-tomentose. *Leaves* coriaceous, drying a dark brown, broadly elliptic, shortly and abruptly sub-acuminate,

entire, slightly narrowed to the rounded 3-nerved base: upper surface glabrescent, the midrib sub-tomentose, lower surface rather sparsely rusty stellate-pubescent: main lateral nerves 5 to 8 pairs, prominent beneath as are the rather straight transverse veins: length 6 to 9 in., breadth 3.5 to 4.5 in.; petiole .5 to .75 in. stout, tomentose. *Panicles* short, axillary or terminal, rusty-tomentose 1.5 to 2.5 in. long and 1 in. or more broad, few-flowered: involucre ovate-lanceolate. *Flowers* 2.5 in. long, their pedicels shorter. *Sepals* oblong, tomentose on both surfaces. *Petals* shorter than the sepals, oblong, acute, the hirsute claw larger than the glabrescent limb. *Torus* cup-shaped, with hirsute margin. *Ovary* depressed-globose: style cylindric, puberulous. *Drupe* obovoid, .75 in. long and .5 in. in diam., pericarp at first sparsely pubescent, afterwards glabrous, mesocarp fibrous and pulpy: pyrene single, 1-celled, 1-seeded: endocarp bony.

Malacca; Griffith, (Kew Distrib.) 638/1; Maingay. Perak; King's Collector, Scortechini, Wray.

6. *GREWIA ANTIDESMEFOLIA*, n. sp., King. A tree usually 30 to 40, but sometimes 50 to 60 feet, high: young branches glabrous, their bark cinereous. *Leaves* membranous, glabrescent when young, when old quite glabrous, elliptic-oblong, acute or shortly acuminate, entire, the base usually cuneate but sometimes rounded, boldly 3-nerved; lateral main nerves 5 or 6 pairs, little curved, ascending, prominent beneath; length 4.5 to 8 in., breadth 1.5 to 2.75 in.; petiole .4 to .6 in. slender. *Panicles* pedunculate, axillary and terminal, slender, the branches short, spreading, few-flowered, densely but minutely cinereous, velvety, 2 to 3 in. long. *Flowers* .25 in. long, their pedicels shorter. *Sepals* elliptic, slightly obovate, their edges in the upper half much incurved, tomentose outside, pubescent inside. *Petals* much shorter than the sepals, oblong, blunt, the glabrescent limb about as long as the broad thickened claw; claw pilose behind, with hirsute edges in front. *Torus* cylindric, glabrous, with wide wavy hirsute mouth. *Ovary* ovoid-globose, pilose when young, glabrescent when adult, shorter than the cylindric glabrous style. *Fruit* pyriform, obtusely 3-angled, narrowed to a long pseudo-stalk, .75 in. long (including the narrowed portion) about .35 in. in diam.; pericarp glabrous, mesocarp slightly fleshy with a thin fibrous inner layer. *Pyrenes* 3, with bony endocarp, two of them abortive and the third 1-celled, 1-seeded.

Perak: at low elevations; common, Scortechini, King's Collector, Wray.

Var. *hirsuta*; young branches, lower surfaces of leaves, and ovary pubescent to tomentose.

Perak; King's Collector.

7. *GREWIA LAURIFOLIA*, Hook. in Hook. fil. Fl. Br. Ind. i. 392. A tree 20 to 30 feet high; all parts except the inflorescence glabrous: young branches with dark-coloured bark. *Leaves* thinly coriaceous, oblong-lanceolate or lanceolate, acuminate or acute, entire; the base rounded boldly 3-nerved: both surfaces shining; lateral nerves 1 or 2 pairs, alternate; length 4 to 6 in., breadth 1·5 to 2·5 in., petiole ·5 to ·7 in. *Panicles* terminal and axillary, 1·5 to 4 in. long, lax, few-flowered, puberulous: bracteoles few, linear, fugaceous. *Flowers* ·2 in. long, their pedicels about ·15 in. *Sepals* ovate, concave, the edges much inflexed, minutely tomentose on both surfaces. *Petals* much shorter than the sepals, oblong, often absent. *Torus* cup-shaped, its rim hirsute. *Ovary* globose, sub-glabrous, 4-celled. *Style* thick, cylindric, tapering, glabrous. *Drupe* ovoid, ·5 in. long, the pedicel about as long, with 1 or 2 vertical furrows, pericarp glabrous, endocarp fleshy and fibrous: pyrene 1 to 3-celled, but usually only one cell containing a single seed: endocarp bony.

Malacca; Griffith, Maingay. Penang; Curtis. Perak; Scortechini, King's Collector. Distrib. Sumatra.

8. *GREWIA CALOPHYLLA*, Kurz Andam. Rep. App. B. iii; Flor. Burm. i. 157. A tree 20 to 30 feet high: all parts glabrous except the minutely velvety tawny inflorescence: young branches slender, dark-coloured. *Leaves* thinly coriaceous, shining, ovate-lanceolate to ovate-elliptic, acuminate, entire; the base rounded or slightly cuneate, 3-nerved; lateral nerves 3 or 4 pairs, ascending; transverse nerves slender: length 4 to 7 in., breadth 1·75 to 3 in., petiole ·3 to ·75 in. *Panicles* pedunculate, axillary or terminal, few-flowered, 1·5 to 3 in. long. *Flowers* ·5 in. long, their pedicels very short. *Sepals* narrowly oblong, the edges much incurved, minutely velvety, much reflexed. *Petals* about half the length of the sepals and much narrower, lanceolate; the limb subulate; the claw ovoid, expanded, thick and densely tomentose at the margin. *Torus* cylindric, puberulous outside. *Ovary* ovoid, pointed, style long filiform, both puberulous. *Fruit* obovoid, compressed, 1·4 in. long and ·75 in. in diam.; pericarp membranous, glabrous, shining; mesocarp thick, pulpy and fibrous: pyrenes 3, of which one is 2-celled but contains only a single seed, the others abortive; the endocarp bony. Mast. in Hook. fil. Fl. Br. Ind. i. 392.

Nicobar Islands, Kurz: S. Andaman, Kurz, King.

This is very near *G. laurifolia*, Hook. but has very much larger fruit. A Malacca plant (Griffith, No. 630/2 Kew Distrib.) resembles this in leaves but not in flower. The only specimens which I have seen are too imperfect for determination.

9. GREWIA MIQUELIANA, KURZ, in Flora for 1872, p. 398. A tree 20 to 40 feet high: young branches at first very sparsely and minutely lepidote, afterwards glabrous, the bark dark brown. *Leaves* thinly coriaceous, glabrous, shining, ovate-lanceolate to lanceolate, shortly acuminate, entire, the base cuneate, faintly 3-nerved; both surfaces glabrescent soon becoming glabrous: main lateral nerves 5 or 6 pairs, not prominent; length 3 to 5 in., breadth 1 to 1.75 in.; petiole .2 to .3 in., scaly-tomentose; stipules oblong, blunt, oblique. *Panicles* axillary and terminal, lax, few-flowered, sparsely lepidote and puberulous, 1 to 2 in. long. *Flowers* .3 in. long, their pedicels very short. *Sepals* oblanceolate, acute, the edges inflexed, minutely tomentose. *Petals* much shorter than the sepals, the glabrescent linear acute limb shorter and narrower than the thickened rounded tomentose claw. *Torus* short, cylindric, puberulous with villous edges. *Ovary* globose-ovoid, tomentose, shorter than the cylindric glabrous style, 2-celled. *Drupe* pyriform, .75 in. long and .5 in. in diam., glabrous: pericarp smooth, glabrous, shining; mesocarp fibrous with a little pulp: pyrenes 2, each 1-celled, one 1-seeded, the other barren: the endocarp bony. *Inodaphnis lanceolata*, Miq. Fl. Ind. Bat. Suppl. 357; Ann. Mus. Lugd. Bat. iii. 89; Meisn. in DC. Prod. xv. 1, 265.

Malacca; Maingay, (Kew Distrib.) No. 244. Perak; Scortechini, King's Collector, at low elevations. Dindings; Curtis, No. 1613. Distrib. Sumatra.

There is an authentic fruiting specimen in the Calcutta Herbarium of Miquel's *Inodaphnis lanceolata* collected in Sumatra. And there is no doubt whatever that Kurz was right in referring the plant to *Grewia*. Miquel founded his genus on specimens without flowers; and, apparently from the structure of the fruit, he suggested its affinity to *Inocarpus*. Later on he suggested (Ann. Mus. Lugd. Bat. iii. 89) its affinity with the Rosaceous genera *Chrysobalanus*, *Parastemon* and *Diemenia* (= *Trichocarya*). Meissner in DC. Prod. (l. c.) briefly described the genus at the end of *Hernandiaceae*, but without indicating his opinion as to its proper place. Had these distinguished botanists had an opportunity of examining flowers, they would doubtless have referred it without hesitation to *Grewia*. The practice (fortunately confined to a few authors) of founding genera on specimens without flowers cannot be too strongly condemned.

6. TRIUMFETTA, Linn.

Herbs or undershrubs, generally more or less covered with stellate hairs. *Leaves* serrate or dentate, simple or lobed. *Flowers* yellowish, in dense cymes. *Sepals* 5, oblong, concave. *Petals* 5. *Stamens* 5-35,

springing from a fleshy, lobed, glandular torus. Ovary 2-5-celled, cells 2-ovuled; style filiform, stigma 5-toothed. Fruit globose or oblong, spiny or bristly, indehiscent or 3-6-valved. Seeds 1-2 in each cell, pendulous, albuminous embryo straight, cotyledons flat. Distrib. A genus of about 40 very variable species, mostly tropical weeds.

Fruit tomentose, bristles shorter than itself ...	1.	<i>T. rhomboidea</i> .
„ villous „ longer „ ...	2.	<i>T. pilosa</i> .
„ glabrous „ „ „ ...	3.	<i>T. annua</i> .

1. TRIUMFETTA RHOMBOIDEA, Jacq. DC. Prod. i. 507 Erect, herbaceous or shrubby, annual, glabrous or pubescent. Leaves polymorphous, but usually rhomboid, 3-lobed, coarsely and unequally serrate, the upper more or less lanceolate; length 1.75 to 3 in., breadth nearly as much in the rhomboid, much less in the lanceolate forms; petioles .25 to 1.25 in. Peduncles short, 4 to 6-flowered. Flowers about .15 in. long, the buds clavate. Sepals apiculate: petals oblong, ciliate at the base. Stamens 8 to 15. Fruit about .2 in. in diam., globose, tomentose, covered with short glabrous or pubescent hooked spines. Masters in Hook. fil. Fl. Br. Ind. i. 395. *T. angulata*, Lam. Dict. iii. 41; Wight Ic. t. 320; W. & A. Prodr. i. 74; Thwaites Enum. 31; Dalz. & Gibs. Bomb. Fl. 25; Wall. Cat. 1075, 2, C; Miq. Fl. Ind. Bat. pt. i. 197. *T. angulata*, β . *acuminata*, Wall. Cat. 1075 β . *T. Bartramia*, Roxb. Fl. Ind. ii. 463; Wall. Cat. 1075, D, E. *T. trilocularis*, Roxb. Fl. Ind. ii. 462; Wall. Cat. 1083. *T. vestita*, Wall. Cat. 1078, in part.

In all the provinces: a weed. Distrib. British India, Ceylon, Malacca, Archipelago, China, Africa.

2. TRIUMFETTA PILOSA, Roth Nov. Sp., 233. Erect, herbaceous or shrubby, annual; the whole plant, but especially the young branches and the under surface of the leaves, villous, stellate-tomentose. Leaves; the lower broadly ovate, sometimes 3-lobed; the upper ovate to ovate-lanceolate, acute or acuminate, unequally and rather coarsely serrate or dentate; length 2 to 4.5 in., breadth 1 to 1.75 in.; petiole .5 to 1 in. Stipules linear-subulate. Peduncles many-flowered, usually shorter than the petiole. Calyx .25 in. long, sparsely hairy. Petals spatulate-oblong, nearly as long as the calyx. Fruit globular, about .25 in. in diam., villous, densely covered with spines longer than itself which are hispid below, glabrous above, and usually hooked at the apex. W. & A. Prodr. i. 74; Hook. fil. Fl. Br. Ind. i. 394. *T. pilosa*, var. β , Thwaites Enum. 31; Dalz. & Gibs. Bomb. Fl. 25. *T. tomentosa*, Wall. Cat. 1078 C. *T. glandulosa*, Heyne Herb.; Wall. Cat. 1077, 5. *T. polycarpa*, Wall. Cat. 1079. *T. oblongata*, Link Enum. Pl. Hort. Ber. ii. 5; Wall.

Cat. 1077, 1, 2, 3. *T. ovata*, DC. Prodr. i. 507? *T. pilosa*, Wall. Cat. 1080. *T. pilosa*, var. *a*, Thwaites Enum. 31. *T. vestita*, Wall. Cat. 1078, 1, 2. *T. indica*, Ham. in Wall. Cat. 237, 1078 D; W. & A. Prodr. i. 74. *T. oblonga*, Wall. in Don. Prodr. 227.

Malacca, Singapore: Perak, King's Collector, No. 989; and probably in all the provinces. Distrib. British India, Ceylon, Africa.

A common and rather variable weed. The bristles of the fruit are usually hooked at the apex; but in some specimens they are quite straight. The species *T. tomentosa*, was founded by Bojer on specimens collected in Mombassa, having straight fruit-bristles and the lower leaves broadly oval or oblong and often 3-lobed. Many of the Indian forms have been referred to that, but I think they might very well be included in *T. pilosa*, and in the synonymy above quoted I have adopted this view.

3. TRIUMFETTA ANNUA, Linn. Mant. p. 73. Annual, shrubby, erect, 1 to 2 feet high; the whole plant with sparse pale straight hairs, the older parts glabrescent. *Leaves* thin, ovate-acuminate, coarsely dentate, 3-nerved, 3 to 5 in. long, by 1.5 to 2 broad: petioles nearly 1.5 in. *Stipules* subulate, minute. *Peduncles* axillary, 3-flowered. *Calyx* .25 in. long, nearly glabrous. *Petals* shorter than calyx. *Stamens* 10. *Fruit* globose, pitted, glabrous, .2 in. across, bearing numerous smooth glabrous thin hooked spines longer than the capsule. DC. Prodr. i. 507; Miq. Fl. Ind. Bat. i. pt. 2, 196; Hook. fil. Fl. Br. Ind. i. 396. *T. polycarpa*, Wall. Cat. 1079, partly. *T. trichoclada*, Link. ex DC. Prodr. i. 507; Wall. Cat. 1082. *T. indica*, Lam. Diet. iii. 420?

Perak: a weed. Distrib. British India, Malay Archipelago, Africa.

7. CORCHORUS, Linn.

Herbs or undershrubs, more or less covered with stellate pubescence, or glabrescent. *Leaves* simple. *Peduncles* axillary or opposite to the leaves, 1-2-flowered. *Flowers* small, yellow. *Sepals* 4-5. *Petals* 4-5, glandless. *Stamens* free, indefinite or rarely twice the number of the petals, springing from a short torus. *Ovary* 2-6-celled, style short, stigma cup-shaped. *Capsule* elongated, slender or subglobose, smooth or prickly, loculicidally 2-5-valved, sometimes with transverse partitions. *Seeds* numerous, albuminous, pendulous or horizontal; embryo curved. Distrib. 35 species, throughout the tropics.

Capsules globular	1. <i>C. capsularis</i> .
„ cylindric, 10-ridged	2. <i>C. olitorius</i> .
„ „ 6-winged	3. <i>C. acutangulus</i> .

1. *CORCHORUS CAPSULARIS*, L. sp. 746. Annual, shrubby, glabrescent. *Leaves* lanceolate, or oblong-lanceolate, acuminate, coarsely serrate, the base rounded and with 2 subulate appendages: length 2 to 4 in., breadth .75 to 1.5 in., petiole .5 in. or less; stipules linear-subulate .25 to .5 in. *Capsules* axillary, truncate-globose, ridged, wrinkled, sub-muricate, 5-celled. *Seeds* few in each cell. DC. Prodr. i. 505; Roxb. Fl. Ind. ii. 581; W. & A. Prodr. i. 73; Wall. Cat. 1071 A, B, C; Wight. Ic. t. 311; Thwaites Enum. 31; Dalz. & Gibs. Bomb. Fl. 25; Miq. Fl. Ind. Bat. i. pt. 2, 194; Hook. fil. Fl. Br. Ind. i. 397. *C. Marua*, Ham. in Wall. Cat. 6311.—Rumph. Amb. v. t. 78, f. 1.

Cultivated here and there in all the provinces for its fibre which is known in commerce as "Jute." Doubtfully wild.

2. *CORCHORUS OLITORIUS*, L. sp. 746. Annual, shrubby, glabrescent. *Leaves* ovate-lanceolate, serrate, the base rounded and with 2 subulate appendages: length 2 to 4 in., breadth .75 to 2 in., petiole .75 to 1.5 in.; stipules linear, .5 to 1 in. *Capsules* cylindric, 10 ribbed, 5-celled, 2 in. long. DC. Prodr. i. 504; Roxb. Fl. Ind. ii. 581; W. & A. Prodr. i. 73; Wall. Cat. 1072; 1, 2, 3, 4, D, E, F; Boiss. Fl. Orient. i. 845; Dalz. & Gibs. Bomb. Fl. 25; Miq. Fl. Ind. Bat. i. pt. 2, 195; Thwaites Enum. 31; Hook. fil. Fl. Br. Ind. i. 397. *C. decemangularis*, Roxb. Fl. Ind. ii. 582; Wall. Cat. p. 237, 1072 G.

Doubtfully wild: but occasionally cultivated in all the provinces under the name of "Jute."

3. *CORCHORUS ACUTANGULUS*, Lamk. Diet. ii. 104. Erect, herbaceous, the stems with a broad line of pubescence interrupted and varying in position at the nodes, otherwise glabrous. *Leaves* ovate to ovate-lanceolate, acute or acuminate, serrate, the base rounded, with or without subulate appendages, sparsely hairy on both surfaces; length 1.5 to 2 in., breadth .75 to 1.75 in.; petiole .25 to .75 in. slender, villous at the apex; stipules lanceolate, acuminate, .5 in. long. *Capsules* 1 to 1.5 in. long, cylindric, 6-winged, with 3 terminal bifid beaks, 3-celled. DC. Prodr. i. 505; W. & A. Prodr. i. 73; Wall. Cat. 1069, 1074 D, E; Wight Ic. t. 739; Thwaites Enum. 31; Dalz. & Gibs. Bomb. Fl. 25; Miq. Fl. Ind. Bat. i. pt. 2, 194; Hook. fil. Fl. Br. Ind. i. 398. *C. astuans*? Ham. in Wall. Cat. p. 237, 1074 C. *C. fuscus*, Roxb. Hort. Beng. 42; Fl. Ind. ii. 582; Ham. in Wall. Cat. 1069.

Johore: at the base of Gunong Pantii, King's Collector, No. 180. Distrib. India, Ceylon, Australia, Africa, W. Indies.

8. *TRICHOSPERMUM*, Blume.

Trees with penni-nerved, minutely stellate, puberulous leaves.

Flowers in axillary or terminal, umbellate, stalked cymes or panicles. *Sepals* 5 valvate, thick. *Petals* 5, membranous with a scale at the base. *Stamens* numerous, free, inserted on the inner surface of an annular marginally villous sub-crenate disk; anthers broad, short, versatile, the connective sub-orbicular. *Ovary* sessile, 2-celled, with numerous ovules on axile placentas: style short, stigma expanded, papillose. *Capsule* orbicular-reniform, much compressed at right angles to the dissepiments, loculicidally 2-valved, many-seeded. *Seeds* sub-lenticular, with a thin imperfect marginally villous arillus; albumen fleshy; embryo central the cotyledons orbicular, foliaceous; radicle straight. Distrib. 3 species 2 of which are Malayan and Polynesian.

1. TRICHOSPERMUM KURZII, King. A tree 40 to 60 feet high: bark of young branches very dark-coloured, sparsely and minutely stellate-pubescent when young, speedily glabrous. *Leaves* membranous, ovate-elliptic, shortly acuminate, minutely serrate-crenate especially near the apex; the base rounded, sub-truncate, sub-cordate, boldly 3-nerved: lateral nerves about 4 pairs: the transverse veins sub-horizontal, curved, bold: length 4 to 6 in., breadth 2 to 3 in., petiole about .5 in. *Panicles* axillary or terminal; stalked, cymose, 2-3-chotomous, much shorter than the leaves when in flower, nearly as long when in fruit, stellate-tomentose. *Sepals* oblong, acute, stellate-tomentose outside, glabrous inside except a tuft of hairs at the base. *Petals* about the size and shape of the sepals, glabrescent, with a fleshy scale at the base and a transverse belt of long hairs above it. *Ovary* sessile, densely villous; style shorter than the ovary, cylindric, expanding upwards into the broad papillose stigma. *Capsule* about .75 in. long and slightly wider, emarginate at the apex and crowned by the persistent style: pericarp leathery, villous and dark-coloured; inside white, shining and glabrous: placentas broad, seeds sessile or shortly stalked, sub-lenticular, the long hairs of the arillus forming a marginal ring. *Bixagrewia nicobarica*, Kurz, Trim. Journ. Bot. for 1875, p. 325, t. 169.

Nicobars: Kurz, Perak; King's Collector, Wray.

The genus *Trichospermum* was founded by Blume for his single species *T. Javanicum*. The generic definition which I have given above differs from that of Blume (Bijdr. 56), in these respects. Blume describes (1) the aestivation of the sepals as imbricate; (2) the style as absent; (3) the stigmas as two and emarginate. The definition also differs from that given by Benth. & Hook. (G. P. i. 236) inasmuch as these authors describe (1) the petals as naked at the base; (2) anthers oblong; (3) style almost none; (4) stigma sessile, retuse; (5) the apex of the capsule produced into a short thick leathery expansion; (6) leaves entire.

9. ELEOCARPUS, Linn.

Trees. *Leaves* simple. *Flowers* usually hermaphrodite, rarely polygamous, in axillary racemes. *Sepals* 5, distinct. *Petals* 5, usually lacinate at the apex, rarely entire, springing from the outside of a cushion-shaped, often 5-lobed torus. *Stamens* usually indefinite, never less than 10, arising from the inside of the torus, and more or less aggregated into groups opposite the petals and alternating with the glands of the torus; anthers innate, linear, opening by a terminal pore. *Ovary* sessile, 2-5-celled, cells 2-many-ovuled; style columnar. *Drupe* with a single bony stone which is 3-5 or, by abortion, 1-celled. *Seeds* pendulous, 1 in each cell, albumen fleshy; cotyledons flat. Distrib. About 50 species chiefly in the Indian Archipelago and India; a few in some of the South Sea Islands, New Zealand, and Australia.

Sect. I. *Ganitrus*. *Ovary* and *drupe* 5-celled, the latter globular.

Leaves glabrescent or glabrous, without stipules.

Leaves lanceolate 1. *E. Ganitrus*.

„ ovate-oblong 2. *E. parvifolia*.

Leaves softly rusty-pubescent or tomentose beneath, stipulate 3. *E. stipularis*.

Sect. II. *Eu-eleocarpus*. *Ovary* 3-celled: longer cell of anthers usually with an apical tuft of minute hair; petals cuneiform, fimbriate.

Leaves pubescent beneath, elliptic-oblong 4. *E. Scortechinii*.

Leaves glabrescent beneath; the midrib pubescent.

Leaves ovate to elliptic-ovate, with black dots beneath 5. *E. Wrayi*.

„ narrowly lanceolate, not dotted beneath... .. 6. *E. salicifolius*.

Leaves quite glabrous everywhere.

Leaves with rounded bases.

Petals glabrous 7. *E. robustus*.

„ glandular-pubescent 8. *E. nitidus*, var. *leptostachyus*.

Leaves with their bases much narrowed.

Petals glandular-pubescent; fruit ovoid or slightly obovoid, blunt 8. *E. nitidus*.

- Petals glabrous except on the edges: fruit ovoid-elliptic, slightly apiculate ... 9. *E. floribundus*.
- Sect. III. *Monocera*. Outer cell of anther produced into an awn. Ovary 2-celled. *Drupe* 1-celled, 1-seeded.
- Petals ovate-acuminate, entire ... 10. *E. paniculatus*.
- Petals about equally wide at base and apex; the apex toothed ... 11. *E. petiolatus*.
- Petals wider at the base than the apex, the edges much incurved below the middle, the apex irregularly toothed or fimbriate.
- Apex of leaves acuminate.
- Racemes longer than the leaves: stamens 35 to 40 ... 12. *E. Griffithii*.
- Racemes usually shorter than the leaves: stamens 20 ... 13. *E. Hullettii*.
- Apex of leaves obtuse: stamens about 15 ... 14. *E. pedunculatus*.
- Petals oblong, slightly obovate, apex obtuse with 6 to 8 broad teeth ... 15. *E. Kunstleri*.
- Petals cuneiform.
- Apex of petals with 8 to 10 rather broad teeth, sometimes 2-lobed: stamens 30 to 50 ... 16. *E. obtusus*.
- Petals oblong-cuneiform to cuneiform, with numerous fimbriae ... 17. *E. apiculatus*.
- Petals broadly cuneiform, lobed and fimbriate ... 18. *E. aristatus*.
- Sect. IV. *Acronodia*. Flowers 4-merous, polygamous; anthers not awned and usually not bearded (sometimes slightly bearded in *E. glabrescens*).
- Leaves sparsely and minutely pubescent or puberulous beneath, their edges serrulate; petals elliptic, the apex slightly lobed ... 19. *E. polystachyus*.
- Leaves rufous-tomentose beneath, subglabrescent only when very old, edges quite entire, recurved; petals oblong, obtuse, 8 to 10-toothed ... 20. *E. Jackianus*.

Leaves rufous-pubescent on lower surface
when young: ultimately glabrescent
or glabrous 21. *E. glabrescens*.

Leaves glabrous at all stages.

Leaves acute narrowed at the base
into the petiole: fruit oblong-
ovoid, .5 in. long 22. *E. punctatus*.

Leaves acuminate (often caudate)
base not passing into petiole:
fruit ovoid-globose, .35 in. long... 23. *E. Mastersii*.

1. *ELÆOCARPUS GANITRUS*, Roxb. Hort. Beng. 42: Fl. Ind. iii. 592. A tree 30 to 60 feet high: branchlets with dark bark, cinereously puberulous when quite young. *Leaves* membranous, lanceolate, acute at base and apex, obscurely serrulate, glabrescent or glabrous: main nerves 10 to 12 pairs, spreading, slender: length 3.5 to 5.5 in., breadth 1.25 to 2.25 in., petiole .3 to .5 in. *Racemes* from the branches below the leaves, drooping, shorter than the leaves, crowded, many-flowered. *Flowers* .35 in. long, narrow and pointed in bud; their pedicels rather longer, puberulous. *Sepals* lanceolate, shorter than the petals, puberulous outside, glabrescent inside. *Petals* obovoid, the base thickened, rounded and puberulous at the edge; the limb glabrous, lacinate for more than half its length. *Torus* short, fleshy, wrinkled, pubescent. *Anthers* about 30 to 35, sessile, slightly pubescent or glabrous; the cells slightly unequal, the longer with 1 (or sometimes 2) short white terminal hairs. *Ovary* ovoid-conic, with deep vertical grooves, minutely tomentose, 5-celled, each cell with about 4 ovules. *Style* much longer than the ovary, thin, fluted, puberulous or glabrescent, thickened towards the base. *Fruit* spherical, .75 to .9 in. in diam., glabrous, bluish-purple; the stone vertically 5-grooved, tubercled, 5-celled, often only one cell containing a ripe seed. Mast. in Hook. fil. Fl. Br. Ind. i. 400; Kurz Fl. Burm. i. 13; Wall. Cat. 2660 A to D; Dalz. & Gibs. Bomb. Fl. 27. *Ganitrus sphaerica*, Gærtn. Fruct. ii. 271, t. 139, f. 6; Wight Ic. i. 66.—Rumph. Amb. iii. t. 101. *E. cyanocarpa*, Maing. in Hook. fil. Fl. Br. Ind. i. 406.

Malacca; Maingay, No. 263. Penang; Curtis. Perak; King's Collector, Scortechini. Distrib. Java; British India, in damp tropical forests as far west as Nepal.

I have dissected flowers of the type specimen (Maingay No. 263) of *E. cyanocarpa*, Maingay, and I can find no difference in them from those of the type sheets of *E. Ganitrus* in Wall. Cat. Roxburgh's original drawing of *E. Ganitrus* in Herb. Calc. is wrong as regards the petals,

which it represents as too broad and with too many fimbriæ: otherwise it is an equally exact representation of the Indian plant described by him as *E. Ganitrus*, and of *E. cyanocarpa*, Maingay.

2. *ELEOCARPUS PARVIFOLIUS*, Wall. Cat, 2662 A & B. A tree 30 to 50 feet high: young branches at first minutely pubescent, ultimately glabrous greyish-brown and minutely lenticellate. *Leaves* membranous, ovate-oblong, rather bluntly acuminate, serrulate, the base cuneate: upper surface shining, glabrous; the lower dull of chocolate brown colour, glabrous or glabrescent, the midrib and 5 or 6 pairs of curved ascending nerves pubescent on both; length 2·5 to 4 in., breadth 1·1 to 1·4 in.; petiole ·6 to ·75 in., slender, puberulous. *Racemes* from the branches below the leaves, rather shorter than the leaves, the rachis, flower-pedicels and outside of calyx softly and shortly pubescent. *Flowers* ·3 in. in diam., their pedicels about ·1 in., recurved, buds conical. *Sepals* slightly shorter than the petals, lanceolate, puberulous within and 3-nerved. *Petals* cuneiform, slightly nerved, cut half-way down into numerous narrow lacinia, almost glabrous. *Torus* of 5 distinct, broad, shallow, fleshy, grooved, pale, velvety glands. *Stamens* 15, shorter than the petals, with short filaments; the anthers scaberulous, cells equal, obtuse, the outer sometimes with 2 or 3 minute pale apical hairs. *Ovary* globose, 5-grooved, 5-celled, sparsely pubescent. *Style* as long as the stamens, cylindric, faintly 5-grooved, glabrescent or glabrous. *Fruit* globose, sometimes ovoid-globose, ·75 to 1 in. in diam.: stone 5-celled, with fertile seeds in only 2 or 3 cells, ovoid, ·7 in. long, bluntly rugose, and with 5 very faint grooves from base to apex. C. Mull. Annot. de fam. Elæocarp. 24; Hook. fil. Fl. Br. Ind. i. 401.

Singapore; Ridley, King's Collector. Malacca; Griffith, (Kew Distrib.) 684, Maingay, 254. Penang and Singapore; Wallich, Curtis. Perak; King's Collector, Scortechini.

3. *ELEOCARPUS STIPULARIS*, Blume Bijdr. 121. A more or less rusty-pubescent tree 40 to 70 feet high: young branches thin, minutely tomentose. *Leaves* coriaceous, ovate to oblong-ovate, acute or acuminate: the edges usually entire, slightly recurved when dry, sometimes waved; the base slightly cuneate, or sometimes rounded: upper surface at first puberulous, ultimately glabrous, the midrib always pubescent: lower softly rusty-pubescent: main nerves 9 to 12 pairs, spreading, interarching close to the margin: length 3·6 to 6·5 in., breadth 1·75 to 2·5 in.; petiole ·5 to ·75 in., minutely tomentose, not conspicuously thickened at the apex; stipules halbert-shaped, tomentose, fugaceous. *Racemes* axillary and from the axils of fallen leaves, usually shorter than, but sometimes as long as the leaves; the rachises, pedicels

and outside of sepals minutely tomentose. *Flowers* .35 in. in diam., their pedicels .2 to .3 in. long; buds sub-globose, obtusely pointed. *Sepals* ovate-lanceolate, pubescent inside especially towards the base, the mid-rib thickened. *Petals* longer than the sepals, cuneiform, lobed and cut irregularly half-way into about 25 slightly unequal fimbriae, veined, glabrous, the edges villous. *Torus* of 5 distinct, fleshy, sub-globose, puberulous, transversely oblong, truncate, 2-grooved glands. *Stamens* 25, about half as long as the petals: filaments about half the length of the scaberulous anthers; cells unequal, the longer with (but sometimes without) an apical tuft of 4 or 5 stiff white hairs. *Ovary* ovoid-globose, vertically 5-furrowed, tomentose, 5-celled. *Style* twice as long as the ovary, conic-cylindric, pubescent at the thickened base, glabrescent above. *Fruit* globose, smooth; .8 to 1 in. in diam.; pulp thin: stone very hard, thick, 1-seeded. *Miq. Fl. Ind. Bat.*, i. pt. 2, p. 209; *Mast. in Hook. fil. Fl. Br. Ind.* i. 404; *Kurz Fl. Burm.* i. 170.

Malacca; Griffith, No. 683, Maingay, No. 255, (Kew Distrib.). Singapore, Malacca, Penang, Perak; very common at low elevations. Distrib. Java, Sumatra, Borneo, Burmah.

Var. *latifolia*, King. *Leaves* broadly elliptic to elliptic-oblong 5 to 7 in. long and 2.75 to 3.75 in. broad: petioles elongate, 1.5 to 2.75 in.; stipules lanceolate.

Perak; Scortechini No. 1991, King's Collector, Nos. 4412, 8176, 10786.

4. *ELEOCARPUS SCORTECHINII*, n. sp. King. A tree 30 to 50 feet high: young branches and stipules as in *E. stipularis*. *Leaves* elliptic-oblong otherwise as in *E. stipularis* except that the main nerves are only 8 to 10 pairs, and the under surface is only softly pubescent, not tomentose: length 5.5 to 7.5 in., breadth 2.25 to 3.25 in. *Flower pedicels* longer than in *E. stipularis*, and the flowers the same, except that the ovary is 3-furrowed and 3-celled. *Fruit* oval, 1 to 1.25 in. long and .5 to .75 in. in diam., glabrous and smooth when ripe, 1-celled, 1-seeded by abortion.

Perak; Scortechini, No. 1481; Wray, Nos. 1876, 1836, 2251; King's Collector, Nos. 3483, 10303.

This is one of the few plants to which the lamented Father Scortechini gave a manuscript name. He dedicated it to Jack: but as Wallich's species, dedicated to the same botanist, has long priority, I name this after my deceased friend. In everything but its 3-celled ovary and smooth oval fruit it agrees with *E. stipularis*, Bl.

5. *ELEOCARPUS WRAYI*, n. sp., King. A small tree: leaf-buds,

young branches and inflorescence pale tawny-pubescent. *Leaves* ovate to elliptic-ovate, shortly and bluntly acuminate, the margin cartilaginous, sometimes crenate-serrate, the base always entire and rounded; upper surface glabrous, shining: the lower dull, pale but not glaucous, with scattered black dots, glabrescent except the puberulous midrib and 6 or 7 pairs of rather prominent sub-ascending main nerves; the reticulations distinct, wide; length 2.25 to 3.75 in., breadth 1.25 to 1.75 in.; petiole .75 to 1.25 in., pubescent. *Racemes* mostly from the wood below the leaves (a few axillary) more than half as long as the leaves. *Flowers* .2 in. in diam., their pedicels .1 in. long or less: buds ovoid, blunt. *Sepals* lanceolate, sub-acute, outside tomentose, inside pubescent and the midrib thickened; the edges not incurved. *Petals* broadly cuneate, glabrous, cut for a third or a fourth of their length into about 25 narrow fimbriae; the base truncate. *Torus* of 5 distinct, fleshy, oblong, truncate, several-grooved, velvety glands. *Stamens* 20 to 25, shorter than the petals; filaments less than half as long as the minutely scaberulous anthers; cells sub-equal, the longer sometimes with 2 or 3 short white hairs. *Ovary* globose, pointed, grooved, tomentose, 3-celled. *Style* slightly longer than the ovary, conic-cylindric, pubescent at the base, glabrescent above. *Fruit* ovoid-globular, glabrous, slightly rugose, 1 to 1.25 in. long when ripe, and .8 to .9 in. in diam.: pulp rather thin: stone bluntly rugose: putamen very hard, thick: 1 cell with a solitary seed, the other 2 cells abortive.

Perak; on Gunong Bubu at 5000 feet elevation; Wray, No. 3857: Gunong Batu Patch, Wray, No. 1107; Scortechini, No. 400.

This resembles *E. parvifolius*, Wall. in some respects; but its leaves have more rounded bases, their nerves are rather more numerous and the petioles longer; the flower buds are blunt and not pointed as in that species, and they are tomentose rather than pubescent; also the stamens are more numerous and the ovary is 3-celled. This is found moreover at much higher elevations than *E. parvifolius* which is found at elevations under 1000 feet.

6. *ELEOCARPUS SALICIFOLIUS*, n. sp., King. A tree 30 feet high: young branches puberulous. *Leaves* thinly coriaceous, narrowly lanceolate, slightly oblique: acuminate, serrulate-crenulate except at the entire cuneate base; upper surface glabrous, shining, olivaceous when dry, the midrib puberulous; lower dull brown when dry, glabrescent, the midrib puberulous: main nerves about 8 pairs, rather straight, sub-ascending, slender; length 3 to 3.5 in., breadth .7 to .9 in.: petiole about .5 in., puberulous. *Racemes* from the lower axils and from the axils of fallen leaves, nearly as long as the leaves; the slender rachises,

and pedicels pubescent. *Flowers* .25 in. in diam., the pedicels .15 to .2 in. *Sepals* lanceolate, spreading, hoary adpressed-tomentose outside, pubescent inside, the midrib slightly thickened, the edges not incurved. *Petals* a little longer than the sepals, cuneiform, contracted into a rather narrow claw, divided more than half-way down into about eight 3-fimbriate lobes, glabrescent outside, pubescent inside. *Torus* of 5 distinct, subglobose, fleshy, externally grooved glands. *Stamens* 25, shorter than the petals: filaments half as long as the minutely scaberulous shining anthers: cells subequal, pointed, the upper with a minute apical tuft of short hairs. *Ovary* globose, pointed, tomentose, 3-celled. *Style* longer than the stamens, thick and tomentose at the base, cylindric and glabrous above. *Fruit* unknown.

Singapore; King's Collector, No. 1207.

This approaches *E. augustifolius*, Bl. but has smaller more pubescent petals, fewer stamens, and less glabrous leaves. It is also closely allied to *E. hypadenus*, Miq., but has not the characteristic rounded stipules of that species, and the leaf-venation is different. It is also allied to *E. parvifolius*, Wall. from which it differs in its narrower leaves with much more slender veins, and also by its 3-celled ovary.

7. *ELEOCARPUS ROBUSTUS*, Roxb. Hort. Beng. 42; Fl. Ind. ii. 598. A tree 40 to 60 feet high; young branches rather stout, at first puberulous; afterwards glabrous, cinereous, lenticellate. *Leaves* thinly coriaceous, ovate-lanceolate to ovate, acuminate or acute, serrate almost to the slightly narrowed rounded rarely cuneate base; both surfaces glabrous, the upper shining; the lower dull, slightly paler, the minute reticulations rather distinct and the 10 to 12 pairs of spreading curving nerves rather prominent: length 3.5 to 9 in., breadth 1.75 to 3.5 in.; petiole 1 to 2.25 in., thickened at the apex. *Racemes* from the branches beneath the leaves, and a few axillary, often nearly as long as the leaves: rachis, pedicels and outer surface of the sepals pubescent. *Flowers* .5 in. in diam., the pedicels slightly recurved and about .3 in. long. *Sepals* lanceolate, glabrous inside except the incurved pubescent edges, the midrib thick. *Petals* broadly cuneiform, much contracted in the lower half, the base acute, cut half-way down into about 30 narrow fimbriae, glabrous except the puberulous edges. *Torus* of 5 fleshy, truncate, cushion-like velvety glands. *Stamens* 30 to 50, shorter than the petals, scaberulous; the filaments curved, about one-fifth the length of the anthers; cells subequal, the longer with a small tuft of white hair at its apex. *Ovary* ovoid-globose, with about 6 shallow vertical grooves, tomentose, 3-celled. *Style* cylindric, longer than the ovary, shorter than the petals, pubescent in its lower, glabrous

in its upper half. *Fruit* ovoid-globose, 1 to 1.25 in. long: stone oblong-ovoid, rugose, slightly 3-grooved at base and apex, 3-celled. *Mast.* in Hook. fil. Fl. Br. Ind. i. 402; Kurz Fl. Burm. i. 169; Pierre Fl. For. Coch-Chine, t. 147; Wight Ic. t. 64; Wall. Cat. 2664. *E. ovalifolius*, Wall. Cat. 2665; C. Müll. Annot. de fam. Elæocarp. 21. *E. amygdalinus*, Wall. Cat. 6857. *E. serratus*, Wall. Cat. 2666 C. *E. oblonga*, Wall. Cat. 2677. *E. aristatus*, Wall. Cat. 2665 B. ? Wall. Cat. 9027. *E. Helferi*, Kurz MSS.; Hook. fil. Fl. Br. Ind. i. 402.

Penang; Curtis. Pahang; Ridley. Andaman Islands. Distrib. British India, from Burmah to the tropical forests of the E. Himalaya.

8. ELEOCARPUS NITIDUS, Jack Mal. Misc. Vol. i. No. 2, 41; Hook. Bot. Misc. ii. 84. A tree 25 to 35 feet high; young shoots deciduously pulverulent-pubescent, speedily glabrous as are all other parts except the inflorescence; young branches with blackish bark. *Leaves* thinly coriaceous, oblong-lanceolate to elliptic-oblong, acuminate, crenate-serulate, (sometimes obscurely so) the base cuneate (rounded in var. *leptostachya*); upper surface shining, the lower dull brown; main nerves 10 to 13 pairs, spreading, forming slender arches a little short of the margin: length 4.5 to 9 in., breadth 1.75 to 2.75 in.; petiole 1.25 to 2 in. thickened at the apex. *Racemes* crowded on the old wood below the leaves and rather more than half as long; rachis, flower-pedicels, and exterior of sepals sparsely puberulous. *Flowers* .35 in. in diam., their pedicels recurved and rather shorter. *Sepals* shorter than the petals, ovate-lanceolate, acute, puberulous and sometimes lenticellate outside, puberulous inside and the midrib very thick. *Petals* cuneiform, finely and irregularly lacinate for nearly half their length, the entire triangular part with thickened nerves and truncate base, glandular-pubescent especially at the edges. *Torus* of 5 truncate, sub-globular, fleshy, tomentose, cushion-like glands. *Stamens* 15 to 35; the filaments nearly as long as the scabrid obtuse anthers: cells sub-equal, awnless, but sometimes the longer with 2 or 3 small white hairs. *Ovary* globose, slightly thickened below and puberulous. *Fruit* ovoid, or slightly obovoid, smooth, 1.5 in. long, and 1 in. in diam. when quite ripe: stone 3-celled, only one cell bearing a perfect seed. Wall. Cat. 2670; Miq. Fl. Ind. Bat. i. pt. 2, p. 208; Mast. in Hook. fil. Fl. Br. Ind. i. 401; Wall. Cat., No. 2678 (*E. pedunculatus*) in part.

Penang; Jack, Curtis, No. 282, 463. Perak; King's Collector, No. 4926.

The anthers are sometimes without any terminal hairs: sometimes there are a few. I have seen no authentic specimen of Jack's naming,

and nothing that I have dissected quite fits his description of *E. nitidus*, of which he describes the stamens as 15: whereas in the plants which I refer to this species they vary from 15 to 35. Jack describes the putamen as 5-ridged and 5-celled: I do not find more than 3 cells in the ovary. In spite, however, of these discrepancies, I believe that Jack's specimen above cited belongs to the species which he named *E. nitidus*. Wallich's specimen No. 2679 has leaves which do not well answer to Jack's description "attenuate at the base." They are only slightly attenuate, and correspond rather with those of his own species *E. leptostachyus* which is sufficiently distinct as regards the shape of its leaves to be maintained as a variety, though not in my opinion entitled to specific rank.

Var. *leptostachya*. Leaves elliptic-oblong to elliptic-rotund, acute, the edge obscurely serrate-crenate, often sub-entire, the base rounded: length 6 to 9 in., breadth 2.75 to 4.5 in.; petiole 1 in. to 1.75 in., slightly thickened at the apex. *E. leptostachyus*, Wall. Cat. 2672; C. Müll. Annot. de fam. Elæocarp. 23; Mast. in Hook. fil. Fl. Br. Ind. i. 403.

Penang, Wallich; Perak; King's Collector, Nos. 409, 4905, 10105, 10240; Scortechini, Nos. 195, 1752; Wray, No. 2313.

9. *ELÆOCARPUS FLORIBUNDUS*, Blume Bijdr. 120. A tree 30 to 40 feet high: young shoots shortly silky; otherwise glabrous, except the inflorescence. Leaves thinly coriaceous ovate-elliptic to oblong-lanceolate or oblanceolate, shortly acuminate, coarsely crenate-serrate, the base much narrowed; both surfaces shining, with a blistered appearance when dry: main nerves 5 to 7 pairs; length 3 to 5.5 in., breadth 1.75 to 2.75 in., petiole 1 to 1.5 in., thickened at the apex. Racemes usually from below the leaves, sometimes axillary, usually shorter than, but sometimes nearly as long as the leaves; rachises, pedicels and outside of sepals puberulous. Flowers .4 in. in diam., their pedicels about .35 in. long. Sepals lanceolate, outside glabrescent and often pustulate; inside glabrous except the pubescent involute edge, the midrib prominent. Petals cuneiform, lobed irregularly half-way down, the lobes divided into about 25 fimbriae, glabrous except the pubescent edges, the lower half veined and thickened, often pustulate. Torus of 5 distinct, fleshy, oblong, sub-globular, truncate, tomentose glands. Stamens about 30, shorter than the petals, scaberulous, the filaments very short, the cells slightly unequal, the longer with a small apical tuft of white hair. Ovary ovoid-globose, tomentose, 3-celled. Style longer than the stamens, cylindric, puberulous in the lower, glabrous in the upper third. Fruit 1 in. long, ovoid-elliptic and slightly apiculate when ripe, oblong and much apiculate when young: stone narrowly ovoid tapering to each end, with 3 vertical grooves and many rather shallow large rugæ, 3-celled, one

or two of the cells sub-abortive, the walls thick. Mast. in Hook. fil. Fl. Br. Ind. i. 401; Miq. Fl. Ind. Bat. i. pt. 2, 210; Kurz Fl. Br. Burm. i. 167; Pierre Fl. Forest. Coch.-Chine, t. 143; Miq. Fl. Ind. Bat. i. pt. 2, 210. *E. serratus*, Roxb. (not of L.) Fl. Ind. ii. 596. *E. grossa*, Wall. Cat. 2661. *E. serratus*, Roxb. ex Wall. Cat. 2666 A, B. partly. *E. oblongus*, Wall. Cat. 2677; C. Müll. Annot. de fam. Elæocarp. 19, f. 30. *E. Lobbianus*, Turcz. in Mosc. Bull. 1858, 235.

The Nicobar Islands. Distrib. British India through Burmah to the E. Himalaya, in tropical forests.

There is no doubt that this is the plant which Roxburgh described as *E. serratus*, Willd.

10. *ELEOCARPUS PANICULATUS*, Wall. Cat. 2663. A tree 15 to 30 feet high: all parts glabrous except the inflorescence, young branches with dark polished bark. *Leaves* thinly coriaceous, lanceolate or oblanceolate-oblong to ovate-oblong, shortly acuminate; the edges entire, slightly wavy; base slightly cuneate, sometimes rounded: both surfaces glabrous, the upper shining; the lower paler and rather dull, the reticulations distinct; main nerves 5 to 7 pairs, sub-ascending, inter-arching freely within the margin: length 4.5 to 6.5 in., breadth 1.65 to 2.75 in.; petiole .8 to 2 in., glabrous. *Racemes* numerous, from the axils near the apices of the branches, longer than the leaves, erect, rachises puberulous, becoming glabrous: pedicels spreading, slender, minutely pubescent, .5 to .65 in. long. *Flowers* about .5 in. in diam.; buds ovoid with long narrow points. *Sepals* ovate, acuminate, adpressed-sericeous outside; glabrous inside except the pubescent infolded edges. *Petals* not longer than the sepals, ovate acuminate, entire, outside adpressed-sericeous, inside glabrous in the upper villous in the lower half and especially on the thickened midrib and infolded edges. *Torus* a shallow fleshy waved sericeous disk. *Stamens* 50, almost sessile, nearly as long as the petals; anthers sericeous, the cells sub-equal, the outer with a rather thick terminal awn. *Ovary* narrowly ovoid, sericeous, 2-celled. *Style* longer than the ovary, cylindric, glabrous. *Fruit* ellipsoid, blunt at each end, smooth, glabrous, bluish when ripe, .4 to .5 in. long and .25 to .35 in. in diam.; pulp rather thick, slightly fibrous; stone bony, minutely tuberculate, 1-celled, 1-seeded. C. Müll. Annot. de fam. Elæocarp. 12; Mast. in Hook. fil. Fl. Br. Ind. i. 407. *Monoceras leucobotryum*, Miq. Fl. Ind. Bat. Suppl. 409. *Monocera Griffithii*, Müll. l. c.

Singapore; Wallich, Anderson. Malacca; Griffith, Maingay (Kew Distrib.) No. 257. Perak; Scortechini, King's Collector; common at low elevations.

11. *ELEOCARPUS PETIOLATUS*, Wall. Cat. 2673. A tree 20 to 40 feet high; all parts glabrous except the inflorescence; young branches dark-coloured, about the thickness of a goose-quill. *Leaves* coriaceous, elliptic to elliptic-oblong, acute or shortly and bluntly acuminate; edges entire: base slightly cuneate or rounded; both surfaces shining, the lower slightly paler when dry, the reticulations sharply distinct on both surfaces: main nerves 7 or 8 pairs, sub-ascending, curving and interarching a little within the margin: length 4.5 to 6.5 in., breadth 2 to 2.75 in.; petiole 1.4 to 2.4 in. slender, dark-coloured, slightly thickened at the apex. *Racemes* numerous from the old wood just below the leaves, shorter than the leaves, rachises and pedicels deciduously puberulous. *Flowers* .5 in. in diam., their pedicels .35 in.; buds ovoid, rather abruptly pointed. *Sepals* lanceolate, acuminate, almost glabrous externally; quite glabrous internally, the infolded edges alone pubescent, the midrib thickened from base to apex. *Petals* about as long as the sepals, oblong, the apex cut into 10 to 13 narrow glabrous teeth, the lower two-thirds sericeous, cucullate at the base from the infolding of the edges, a large fleshy villous gland in the middle near the base with a quasi-cell at each side of it, the hairs on the inner surface retroversed. *Torus* a 10-lobed fleshy glabrescent disk. *Stamens* 18 to 25, shorter than the petals, with sericeous or glabrescent flat or sub-cylindric filaments much shorter than the shortly puberulous anthers: apex of anther deeply cleft, the outer cell with a sub-recurved thick awn shorter than the filament. *Ovary* ovoid, pointed, glabrous, 2-celled. *Style* as long as the stamens and much longer than the ovary, cylindrical, grooved, glabrous. *Fruit* elliptic, blunt at each end, smooth, .4 to .6 in. long, and .3 in. in diam.: the pulp thin, with very few fibres; stone very slightly rugose, 1-celled, 1-seeded. *Monocera petiolata*, Jack Mal. Misc. i. No. v, 43; ex Hook. Bot. Misc. ii. 86; Cum. et Zoll. in Bull. Mosc. xix, 495. *Monoceras petiolatum*, Miq. Fl. Ind. Bat. i. pt. 2, p. 212; Kurz Fl. Burm. i. 164; Pierre, Fl. Forest. Coch.-Chine, t. 140. *Eleocarpus integra*, Mast. (not of Wall.) in Hook. fil. Fl. Br. Ind. i. 408.

Malacca; Griffith No. 699; Maingay, No. 256, (Kew Distrib.); Derry. Singapore; Hullett, King. Penang; Curtis, No. 333. Perak; Scortechini, King's Collector, Wray, very common at low elevations. Distrib. Sumatra, Beccari, N. S. No. 668.

This is undoubtedly the *Monocera petiolata* of Jack; that it is the *Eleocarpus integra* of Wall. (Cat. No. 2668) I very much doubt. Wallich's No. 2668 was collected in Silhet from which no specimen anything like this has been collected since his day. In fact there is no evidence to show that this species is found in any part of British India (as distinguished from British Malaya), although Kurz includes it in his

Flora of Burmah. This species is a smaller tree than *E. pedunculatus*, which, however, it closely resembles, differing chiefly in the shape of the leaves, the nearly glabrous sepals and in the larger number of stamens. *E. ovalis*, Miq. (a species from Sumatra) must be very nearly allied to this. I have seen only a fruiting specimen of *E. ovalis*, but, except in having leaves of thicker texture and slightly larger fruit, I see little to prevent its being referred here.

12. *ELEOCARPUS GRIFFITHII*, Mast. in Hook. fil. Fl. Br. Ind. i. 408. A tree 30 to 40 or over 70 feet high, all parts glabrous except the inflorescence; young branches almost as thin as a crow-quill, dark-coloured. Leaves thinly coriaceous, ovate-lanceolate to lanceolate, acuminate, the edges cartilaginous with shallow mucronate crenulations, or sub-entire with remote marginal black points, the base sub-cuneate or rounded: both surfaces shining, the reticulations minute and distinct: main nerves 5 or 6 pairs spreading, forking and interarching at some distance from the margin, not prominent: length of blade 2.5 to 3.75 in., breadth .9 to 1.5 in., petiole .5 to 1 in. Racemes from the upper axils, longer than the leaves, rachises and pedicels softly and minutely pubescent. Flowers .5 in. in diam.; pedicels thickened at the apex, .6 to .8 in. long: buds ovoid-conic. Sepals lanceolate, acuminate, finely adpressed sericeous externally, glabrous internally except the pubescent infolded edges and the thickened sometimes sericeous midrib. Petals about as long as sepals, ovate, acuminate, the apex irregularly 2 or 3-toothed with 2 or 3 lateral fimbriæ, outside minutely adpressed-sericeous, inside retroversed hirsute especially on the large gland near the base; edges in the lower two-thirds much infolded so as to form with the gland 2 quasi-cells. Torus a shallow, acutely 10-lobed, fleshy disk. Stamens 35 to 40, shorter than the petals: filaments short, sericeous as are the unequally 2-celled anthers: outer cell with a tapering awn $\frac{1}{4}$ to $\frac{1}{3}$ of its own length, the inner with a few apical hairs. Ovary narrowly ellipsoid, tapering, glabrous except a few silky hairs, 2-celled. Style cylindric, grooved, glabrous, longer than the ovary. Fruit ellipsoid, blunt at both ends, smooth, .5 in. long and .3 in. in diam.: pulp thin with a few fibres; stone slightly rugose, 1-celled, 1-seeded. Kurz in Journ. As. Soc. Beng. pt. 2, for 1870, p. 68; for 1874, pt. 2, 123; For. Flora Burm. i. 164. *Monocera tricanthera*, Griff. Not. pt. 4, 518, t. 619, fig. 3. *Monocera Griffithii*, Wight Ill. i. 84, (not of Müll.). *Monocera holopetala*, Zoll. et Cum. Bull. Mosc. xix, 496. *Monoceras odontopetalum*, Miq. Fl. Ind. Bat. Supp. 409.

Malacca; Griffith, Maingay, No. 257/2 (Kew Distrib.). Perak, at low elevations; King's Collector, Wray. Penang; King's Collector. Distrib. Tenasserim, Helfer, No. 714, Kew Distrib.

13. *ELEOCARPUS HULLETTII*, n. sp., King. A tree 30 to 40 feet high: young branches very slender, dark-coloured; all parts glabrous except the inflorescence. *Leaves* thinly coriaceous, lanceolate to ovate-lanceolate, acuminate; edges slightly cartilaginous, entire or remotely and obscurely serrate; the base cuneate or rounded: both surfaces shining, the reticulations minute, elongate and rather distinct on the lower; main nerves 7 or 8 pairs curving, interarching within the edge, rather faint; length of blade 2·5 to 3 in., breadth ·75 to 1·4 in.; petiole ·65 to ·9 in., slender. *Racemes* from the leaf-axils below the apex, crowded, usually shorter than, but sometimes as long as, the leaves, the rachises glabrescent or puberulous, the pedicels silky puberulous. *Flowers* ·3 in. in diam., their pedicels ·35 in. long. *Sepals* linear-lanceolate, acuminate; externally adpressed-pubescent; internally glabrous below, puberulous near the apex and on the infolded edges. *Petals* ovate, concave at the base, narrowed to the 10 to 12-fimbriate apex; outside glabrous, inside villous on the much-thickened base of the midrib, otherwise puberulous. *Torus* very shallow, deeply 10-lobed, sericeous. *Stamens* 20, slightly shorter than the petals: filaments nearly as long as the minutely scaberulous anthers, outer cell with tapering awn nearly as long as itself. *Ovary* ovoid, pubescent, 2-celled. *Style* as long as the petals and much longer than the ovary, subulate, puberulous below, glabrous above. *Fruit* ellipsoid, blunt at each end, smooth, ·6 in. long, ·35 in. in diam.; pulp thin, very slightly fibrous: stone rugulose, rather thick, bony, 1-celled, 1-seeded.

Singapore; Hullett, No. 132. Penang; Curtis, No. 1091, King's Collector, No. 1475. Perak, on low hills; Scortechini, King's Collector.

A species not unlike *E. Griffithii*, Wall. but with smaller flowers and much shorter racemes.

14. *ELEOCARPUS PEDUNCULATUS*, Wall. Cat. 2678 in part. A tree 40 to 80 feet high: glabrous except the inflorescence: young branches nearly as thin as a crow-quill, polished, dark-coloured; their apices and the older branchlets rough and thickened. *Leaves* coriaceous, oblanceolate or narrowly elliptic-oblong, obtuse or slightly narrowed at the apex, the base very cuneate; the edges cartilaginous, remotely-mucronate crenate-waved, slightly recurved when dry: both surfaces shining: the lower very slightly the paler and with the minute reticulations distinct; main nerves 5 to 7 pairs, interarching at some distance from the edge; length of blade 3 to 4·5 in., breadth 1·25 to 1·8 in., petiole ·75 to 1·2 in. *Racemes* axillary but mostly from axils of fallen leaves, 3 or 4 in. long, rachises and pedicels hoary-pubescent. *Flowers* ·3 in. in diam., buds narrowly ovoid, sub-acute; pedicels recurved, slightly longer than the

flowers. *Sepals* lanceolate, sub-acute, minutely adpressed-sericeous externally, almost glabrous internally except the pubescent inverted edges, the midrib equally thickened from base to apex. *Petals* slightly longer than the sepals, oblong, expanded at the base, the apex broad, cut into 10 to 15 cylindric filiform glabrous fimbriæ one-fourth of the length of the petals: lower part sericeous on both surfaces but especially on the inner (where the hairs are reversed), cucullate with the edges much infolded, and with a large basal gland at each side of which is an imperfect cell. *Torus* a fleshy deeply 10-lobed glabrescent disc. *Stamens* about 15; slightly shorter than the petals, filaments pubescent, less than half the length of the puberulous sub-equal anthers: outer cell with a short sub-recurved awn. *Ovary* ovoid, pointed, smooth, glabrous, 2-celled. *Style* as long as the stamens, cylindric, grooved, glabrous. *Fruit* ellipsoid, blunt at each end, .5 in. long and .3 in. in diam., 1-celled, 1-seeded; pulp rather thin, slightly fibrous, stone minutely rugulose, 1-celled, 1-seeded. *Mast.* in Hook. fil. Fl. Br. Ind. i. 408.

Singapore; Wallich, Ridley. Malacca; Griffith, No. 698, Maingay No. 258 (Kew. Distrib.). Penang; Curtis, No. 256. Perak; Scortechinii, King's Collector, Nos. 269, 6907, 10831.

Miquel's *Monocera Palembangica*, from Sumatra, judging from the only authentic specimen which I have seen (and which has no flowers), if not identical with this must be a very closed allied species. Under his Catalogue, No. 2678, Wallich issued two species, the above described as *E. pedunculatus*, and another which is clearly *E. nitidus*, Jack.

15. *ELEOCARPUS KUNSTLERI*, n. sp., King. A tree 50 to 70 feet high: young branches as thick as a goose-quill, polished, thickened and rough at the apex: all parts glabrous except the inflorescence. *Leaves* coriaceous, rotund-obovate, the apex broadly obtuse, sometimes with a short broad apiculus, rather abruptly narrowed from below the middle to the acuminate base; both surfaces, shining, glabrous; main nerves about 10 pairs, ascending, interarching freely inside the entire or crenate-serrate edge: prominent beneath; the reticulations rather faint; length 5 to 8 in., breadth 2.75 to 3.75 in., petiole .5 to .7 in., pubescent. *Racemes* crowded from the axils of fallen leaves and a few axillary, less than half as long as the leaves, 6 to 9-flowered: rachises and pedicels slender, puberulous, glabrous when old. *Flowers* .6 in. in diam., their pedicels .5 in. or more long. *Sepals* lanceolate, sub-acute, pubescent on both surfaces, the midrib thickened and villous at the base inside. *Petals* about as long as the sepals, oblong slightly obovate, obtuse, thickened in the lower half, the apex with 6 to 8 rather broad teeth, adpressed-sericeous outside, densely villous inside. *Torus* a shallow

toothed villous cup. *Stamens* 28 to 30, shorter than the petals; filaments nearly as long as the minutely scaberulous anthers, swollen in the lower half, the apex of the outer anther-cell with a short recurved awn. *Ovary* ovoid, pointed, tomentose, 2-celled. *Style* cylindric, as long as the petals, puberulous below, glabrous at the apex. *Fruit* unknown.

Perak; at elevations under 1000 feet, King's Collector, No. 8323.

A species near *E. apiculatus*, Mast. but with broader, blunter leaves more abruptly attenuated to the base and quite glabrous, also with smaller flowers.

16. *ELEOCARPUS OBTUSUS*, Blume Bijdr. 125. A tree 30 or 40 feet high: young shoots minutely pale pubescent, ultimately glabrous. *Leaves* coriaceous, oblong-obovate, the apex rounded or retuse, gradually narrowed from above or below the middle into the acute or acuminate base; the edges sub-entire or with shallow mucronate crenations; upper surface glabrous, shining; the lower minutely puberulous at first, ultimately glabrous, the reticulations very minute and rather distinct; main nerves 6 to 8 pairs, sub-ascending, not much curved, rather prominent below, scrobiculate at their origin from the midrib; length 4.5 to 6.5 in., breadth 2.25 to 2.75 in., petiole .75 to 1.1 in. *Racemes* axillary, sometimes from the axils of fallen leaves, less than half as long as the leaves, few-flowered; rachises and pedicels puberulous when young, often nearly glabrous when old. *Flowers* .9 in. in diam; their pedicels slender, .75 to 1 in. long. *Sepals* oblong-lanceolate, outside minutely pubescent, inside sparsely adpressed-sericeous, the midrib thickened. *Petals* longer than the sepals, cuneiform, the base rather broad; the lower third thickened and its edges infolded, the apex with 8 to 10 rather broad teeth sometimes 2-lobed, sericeous on both surfaces but especially on the thickened lower third. *Torus* a wavy, sub-10-toothed, fleshy, sericeous cup. *Stamens* 30 to 50, shorter than the petals: the filaments slender, slightly swollen in the lower half, as long as the pubescent anthers; outer anther cell with a thin tapering awn about as long as itself. *Ovary* ovoid, sericeous or pubescent, 2-celled. *Style* tapering, cylindric, nearly as long as the petals, slightly grooved, puberulous. *Fruit* ovoid, oblong, not pointed, 1.5 in. long, and .9 in. in diam., smooth: stone boldly tuberculate, 1-celled, 1-seeded. *Monoceras obtusum*, Hassk. Tijds. Nat. Gesch. xii. 136; Miq. Fl. Ind. Bat. i. pt. 2, p. 212. *E. Monoceras*, Cav. (fide Mast. in Hook. fil. Fl. Br. Ind. i. 405). *E. littoralis*, Kurz (not of Teyssm. and Binn.) in Journ. As. Soc. Beng. 1874, pp. 132, 182; For. Fl. Burm. i. 167.

Malacca; Griffith, (Kew Distrib.) No. 700. Perak, at low elevations; King's Collector, Nos. 1096, 4671; Scortechini 1396. Pahang; Ridley, 1312. Distrib. Java, Borneo, Sumatra, Burmah.

E. Monoceras, Cav. to which Dr. Masters reduces this, was founded by its author on specimens from the island of Luzon. The species, however, is not given in the latest Flora of the Philippines (that of Sig. Vidal); and, as the original description of Cavanilles does not quite agree with the flowers of the Perak specimens, I think it safer not to go farther back than Blume's name, leaving it to be settled hereafter whether *E. obtusus*, Bl. is really the same plant as the Philippine *E. Monoceras*. The Perak plant is closely allied to *E. littoralis*, T. B. (for which Kurz mistook it); and also to the smaller-flowered Sumatran *E. cuneifolius*, Miq.

17. *ELEOCARPUS APICULATUS*, Mast. in Hook. fil. Fl. Br. Ind. i. 407. A tree 50 to 60 feet high: young branches glabrous, their apices much thickened, rough and puberulous. *Leaves* coriaceous, obovate or oblanceolate-oblong, slightly narrowed to the obtuse, sub-acute, or shortly apiculate apex, and much narrowed to the base, the edges sub-entire or with coarse shallow crenations; both surfaces glabrous, shining, the midrib on the lower glabrescent when young; under surface pale, the reticulations minute, rather distinct; main nerves 12 to 14 pairs, slightly prominent beneath and interarching freely within the margin, not scrobiculate; length 7 to 10 in.; breadth 2.5 to 3.75 in., petiole .3 to 1 in., thickened at the apex. *Racemes* few, mostly from the axils of fallen leaves, usually about a fourth but sometimes half the length of the leaves; the rachises and pedicels softly pubescent. *Flowers* .9 in. in diam.; buds oblong, sub-obtuse or pointed, their pedicels .75 to 1.25 in. *Sepals* oblong-lanceolate, rufous-pubescent outside, glabrous or glabrescent inside, the edge infolded and pubescent, the midrib thickened from base to apex. *Petals* slightly longer than the sepals, oblong-cuneiform to cuneiform, cut from one-fourth to one-fifth of their length into numerous rather broad fimbriæ; externally adpressed-sericeous in the lower half, glabrous in the upper; internally thickened and villous in the lower, glabrous in the upper, half. *Torus* a shallow fleshy puberulous cup. *Stamens* 30 to 40, half as long as the petals; filaments shorter than the minutely scaberulous anthers, bulbous at the base: outer anther-cell with short or long apical recurved awn. *Ovary* ovoid, rufous-tomentose, pointed, 2-celled. *Style* as long as the petals, conic-cylindric and pubescent in the lower half, filiform and glabrous in the upper. *Fruit* (fide Masters) "1 in. long, resembling the fruit of a *Diospyros*." *Terminalia moluccana*, Wall. (not of Lamk.) Cat. 3969.

Penang; Wallich. Malacca; Griffith, Maingay, No. 262 (Kew. Distrib.). Perak; Scortechini, King's Collector; common at low elevations.

Allied to *E. Kunstleri*, King and to *E. rugosus*, Roxb. In fact I am inclined to believe that it is merely a form of the latter, from which it should not be separated specifically. Dr. Prain has called my attention to Wallich's sheet No. 3969, which is unmistakably this species and has nothing to do with *Terminalia moluccana*, Lamk. which is *T. Catappa*, Linn.

18. *ELEOCARPUS ARISTATUS*, Roxb. Hort. Beng.: Fl. Ind. ii. 599. A tree 30 to 60 feet high: young branches of about the thickness of a swan's quill, smooth, thickened and rough towards the apex. Leaves thinly coriaceous, obovate, shortly and bluntly apiculate, remotely crenate-serrate, narrowed to the base, glabrous on both surfaces; main nerves 7 to 10 pairs, slender, curving, scrobiculate at the origin from the midrib; length 6 to 8.5 in., breadth 2.75 to 3.75 in., petiole .5 to .7 in. Racemes axillary and from the axils of fallen leaves, often nearly as long as the leaves, 3 to 5-flowered, rachises and pedicels puberulous or glabrous. Flowers nearly 1 in. in diam.; buds cylindric, pointed; pedicels .8 to 1.25, or longer in fruit. Sepals as in *E. apiculatus*. Petals also as in *E. apiculatus* but broadly cuneiform, and lobed as well as fimbriate. Stamens 50, otherwise as in *E. apiculatus*. Ovary less velvety, but otherwise as in *E. apiculatus*. Fruit ovoid, smooth, 1.25 to 1.4 in. long and .8 to .9 in. in diam., pulp rather thick; stone oblong, flattened, pointed at each end, rugose, slightly ridged in the middle of each side, 1 in. long, 1-celled, 1-seeded. Mast. in Hook. fil. Fl. Br. Ind. i. 405. *E. rugosus*, Wall. Cat. No. 2659 (not of Roxb.).

Andaman Islands; King's Collector. Distrib. Brit. India in Burmah, Chittagong, Sylhet, Assam, Khasia Hills and base of Eastern Himalaya.

This is very closely allied to *E. rugosus*, Roxb.—a species originally discovered by Roxburgh in Chittagong, but specimens of which from that province are very rare in collections. The plants distributed under this name by Wallich as No. 2659 of his Catalogue were not collected there but in Sylhet, while some of them were taken from trees cultivated in the Botanic Garden, Calcutta. They are not *E. rugosus* at all, but *E. aristatus*, Roxb.; and they differ from true *E. rugosus* in having their young branches thinner and smoother; and in leaves which are always glabrous, not so gradually narrowed to the base and with much longer petioles. Their racemes are also more numerous, the petals more broadly cuneiform and the stamens more numerous, (50 as against 30 to 40). The pulp of the fruit is thicker in Andamans specimens of this than in those from Sylhet and Assam; and the stone is proportionately smaller. There is in Assam and Burmah a plant closely allied to this which has

smaller leaves with very large scrobiculae on the lower surface at the junction of the petioles with the midrib. This has been named *E. simplex* by Kurz, (Fl. Burm. i. 165.) A similar form occurs in Travancore and has been named *E. venustus* by Beddome (Flora Sylvatica, t. 574).

19. *ELEOCARPUS POLYSTACHYUS*, Wall. Cat. 2671. A small tree: young shoots rather stout, minutely tawny-tomentose. *Leaves* coriaceous, pale when dry, elliptic to elliptic-oblong, abruptly and shortly acuminate; the edges rather remotely serrulate except at the base, sub-entire when old; the base broad, rounded; upper surface glabrous; the lower sparsely and minutely sub-adpressed puberulous, the midrib pubescent. *Main nerves* 7 to 10 pairs, ascending, curving, prominent beneath, the reticulations minute, faint: length 5.5 to 7.5 in., breadth 2.25 to 3.5 in.; *petioles* 2.5 to 4 in., minutely tawny-tomentose, slightly thickened at the apex. *Racemes* slightly longer than the petioles; the rachises, pedicels and outside of sepals densely minutely tawny-tomentose. *Flowers* .35 in. in diam., their pedicels .4 in., recurved, buds sub-globose. *Sepals* ovate, acute; inner surface glabrous, except the pubescent edges, the midrib thickened. *Petals* elliptic, little longer than the sepals, the apex obtuse, sometimes slightly lobed, not fimbriate: villous on both surfaces, the hairs on the inner reversed. *Torus* of 5 retuse thin densely villous glands. *Stamens* half as long as the petals: filaments nearly as long as the hispid-pubescent anthers; cells subequal, awnless, beardless. *Ovary* ovoid, blunt, densely villous, 2-celled. *Style* about as long as the ovary, puberulous. *Fruit* oblong, blunt, .6 in. long and .35 in. in diam., smooth, glabrous; stone minutely but sharply rugose, 1-celled, 1-seeded, pulp thin and slightly fibrous. C. Müll. Annot. de fam. Eleocarp. 20, f. 13; Mast. in Hook. fil. Fl. Br. Ind. i. 403.

Singapore; Wallich, Hullett, Ridley. Malacca; Maingay, Nos. 264, 266, (Kew Distrib.).

20. *ELEOCARPUS JACKIANUS*, Wall. Cat. 2679. A tree 40 to 80 feet high: young branches stout, densely rufous-tomentose. *Leaves* coriaceous, ovate-oblong to elliptic, rarely oblong-ovoid, shortly acuminate or acute, edges entire, recurved; the base rounded or slightly narrowed: upper surface rather dull and pale when dry; glabrous, the midrib alone sometimes pubescent, the lower softly rufous-tomentose, becoming sub-glabrescent when very old, the minute reticulations distinct; *main nerves* 8 to 10 pairs, sub-ascending, curving, prominent on the lower, impressed on the upper, surface: length 4 to 7.5 in., breadth 2.5 to 4.5 in.; *petiole* 1.75 to 3 in., stout, thickened at each end, tomentose. *Racemes* crowded on the branches below the leaves, sometimes

axillary, shorter than the petioles, the rachises pedicels and outside of sepals softly rufous-tomentose. *Flowers* .25 in. in diam.; their pedicels about .25 in. long, recurved. *Sepals* 4, ovate, acute; inside puberulous with infolded tomentose edges, the midrib thickened. *Petals* 4, very little longer than the sepals, oblong, slightly obovate, obtuse, shortly 8- to 10-toothed, villous outside, glabrescent inside, the edges villous. *Torus* a shallow rufous-villous cup. *Stamens* about 12, shorter than the petals, scaberulous, the filaments about half as long as the anthers; anther-cells slightly unequal, pointed, the longer sometimes with, but usually without, a minute tuft of white hair. *Ovary* (absent in most flowers) ovoid-oblong, glabrous, imperfectly 2-celled, one cell only perfect. *Fruit* ovoid, tapering at each end, smooth, shining; pulp thin, slightly fibrous: stone sharply rugulose, crustaceous, 1-celled. 1-seeded. *Monocera ferruginea*, Jack Mal. Misc. ex Hook. Bot. Misc. ii. 86.

Singapore; Jack, Kurz. Penang; Curtis, No. 465. Malacca; Griffith, No. 693; Maingay, No. 259, (Kew Distrib.). Perak; King's Collector; common at low elevations.

This species approaches *E. glabrescens*, Mast. but is larger in all its parts and much more persistently tomentose. This is unrepresented by any Wallichian specimen at Kew, and is therefore referred to by Masters in the *Flora of Brit. India* only in a note (i. 409).

21. *ELEOCARPUS GLABRESCENS*, Mast. in Hook. fil. Fl. Br. Ind. i. 403. A tree: young branches and petioles densely rufous-tomentose. *Leaves* coriaceous, ovate to ovate-lanceolate, acuminate; edges entire, slightly revolute; base rounded or slightly narrowed; upper surface glabrous, the midrib pubescent; lower surface at first rufous-pubescent ultimately glabrescent or glabrous, the reticulations minute but distinct: main nerves 5 or 6 pairs, prominent beneath, spreading, curved: length 2.5 to 3.5 in., breadth 1.2 to 1.6 in.; petiole 1.25 to 1.75 in., slightly thickened at the apex, glabrescent when old. *Racemes* rather longer than the petioles, axillary and from the axils of fallen leaves; rachises and pedicels sparsely pubescent. *Flowers* .25 in. in diam., the pedicels about .2 in. long, recurved. *Sepals* 4, sub-erect, ovate-lanceolate, thickened at the base, pubescent outside, glabrescent inside with puberulous edges, the midrib thickened. *Petals* 4, slightly longer than the sepals, oblong, the apex obtuse and with 6 to 12 short unequal teeth: pubescent outside, glabrescent inside, the edges shortly villous. *Torus* a very shallow villous cup. *Stamens* 10 or 12, shorter than the petals: filaments short: anthers scaberulous, the cells slightly unequal, slightly pointed, usually without small apical tufts of minute hair. *Ovary* (absent in many flowers) ovoid, glabrous, 1-celled. *Style* short, conic, glabrous. *Fruit* (fide Masters) the size of a cherry, 1-celled, 1-seeded.

Malacca; Maingay No. 256 (Kew Distrib.). Penang; Stoliczka; on Government Hill at 2,500 feet, Curtis, No. 1092.

The Malacca and Penang specimens agree with a specimen at Kew which Miquel has named *E. tomentosus*, Bl. The two species are no doubt close together: but Blume describes the leaves of his *E. tomentosus*, as "setaceous-denticulate" which is not the case here. This tree appears to be uncommon, for Herbarium specimens of it are very few.

22. *ELMOCARPUS PUNCTATUS*, King, n. sp. A small tree; all parts glabrous except the puberulous inflorescence; young branches thicker than a crow-quill, rough. *Leaves* coriaceous, oblong-lanceolate, acute, the edges cartilaginous, crenate or serrate, sometimes with a short seta on each tooth, the base much narrowed into the petiole, entire: both surfaces shining; main nerves 8 to 10 pairs, slender but distinct beneath as are the reticulations: length 1.75 to 3 in., breadth .5 to 1.1 in.; petiole .25 to .3 in., channelled in front. *Racemes* axillary and from the axils of fallen leaves, much shorter than the leaves; rachises and pedicels puberulous, becoming glabrescent. *Flowers* .25 in. in diam., their pedicels .2 in., recurved. *Sepals* 4, oblong-lanceolate, sub-acute, puberulous on both surfaces, the edges thickened and pubescent but not recurved; midrib thickened inside. *Petals* 4, obovoid-oblong, apex obtuse with 5 to 7 short broad teeth, glabrous. *Torus* a shallow wavy pubescent cup. *Stamens* 8 to 12; filaments less than half as long as the scaberulous obtuse beardless awnless anthers: the cells sub-equal. *Ovary* ovoid, glabrous, slightly grooved, 2-celled. *Style* about as long as the ovary, cylindrical, grooved, puberulous. *Fruit* oblong-ovoid, pointed, narrowed to both ends, glabrous, shining, pale, .5 in. long and .25 in. in diam.; pulp thin, and slightly fibrous; stone crustaceous, sharply rugose, 1-celled by abortion, 1-seeded. *Elmocarpus Acronodia*, Mast. in Hook. fil. Fl. Br. Ind. i. 408 *in part*. *Acronodia punctata*, Bl. Bijdr. 123; Miq. Fl. Ind. i. pt. 2, p. 213.

Perak; on Ulu Batang Padang, at 5000 feet, Wray. Malacca. Distrib. Java, Sumatra.

The leaves of specimens from Java and Sumatra are larger than those from Perak and have numerous black dots on the lower surface, whereas those from Perak have no such dots. In other respects the specimens agree: but the Perak material which I have as yet seen is scanty. The plant issued by Wallich as *E. punctatus*, (No. 2676 of his Catalogue) is not the *Acronodia punctata* of Blume, but an altogether different plant. Wallich's specimens are very bad, and Dr. Masters (Fl. Br. Ind. i. 406) suggests that perhaps the leaves are those of a *Pterospermum*; in reality they belong to a species of *Parinarium*.

23. *ELEOCARPUS MASTERSII*, King. A tree 30 to 50 feet high: young branches as thin as a crow-quill, smooth, puberulous; otherwise glabrous except the inflorescence. *Leaves* thinly coriaceous, oblong-lanceolate to ovate-lanceolate, acuminate, often caudate; the edge slightly cartilaginous, remotely and faintly serrate, the base cuneate; both surfaces shining and with the rather transverse reticulations distinct; main nerves 6 to 8 pairs, faint, spreading, interarching within the edge: length of blade 2.75 to 4.5 in., breadth .8 to 1.4 in.; petiole .5 to .75 in., slender. *Racemes* few-flowered, less than half as long as the leaves, from the axils under the apex; rachises and pedicels puberulous, becoming glabrous. *Flowers* .2 in. in diam.; buds narrowly ovoid, pointed. *Sepals* 4, ovate-lanceolate, subacute, puberulous or glabrescent outside: glabrous inside on the lower, often puberulous in the upper half and slightly on the infolded edges. *Petals* 4, oblanceolate or narrowly cuneate, the rounded apex with about 15 short teeth, thickened towards the base, veined, glabrous. *Torus* a very shallow wavy pubescent disk. *Stamens* 8 or 9, shorter than the petals, filaments nearly as long as the sub-scaberulous anthers; the cells blunt at the apex, awnless. *Ovary* (absent in many flowers), ovoid, blunt, glabrous, 2-celled. *Style* about as long as the ovary, thick, cylindrical, grooved, glabrous. *Fruit* ovoid-globose, the apex slightly pointed, smooth, .35 in. long and .25 in. in diam.; pulp thin and without fibres: stone smooth, cartilaginous, 1-celled, 1-seeded. *Elaeocarpus Acronodia*, Mast. in Hook. fil. Fl. Br. Ind. i. 401, in part (excl. syn. *Acronodia punctata*, Bl.).

Malacca; Griffith, No. 681; Maingay, No. 261, (Kew Distrib.). Singapore; Hullett, Ridley. Perak; common at low elevations, King's Collector, Scortechini, Wray.

This is a true *Acronodia* allied to *A. punctata*, Bl. (= *Elaeocarpus punctatus*, King, not of Wall.) but is distinguished by its less acuminate longer petiolate leaves, slightly different flowers and smaller more globose fruit. This occurs at low elevations and is a tree whereas the other is a shrub and is found as high as 7000 feet.

EXCLUDED SPECIES.

ELEOCARPUS PUNCTATUS, Wall. Cat. 2676 is, (as Kurz pointed out) no *Elaeocarpus* but a *Parinarium*. Maingay's Nos. 621 and 621/2 (Kew Distribution) seem to be conspecific with it.

JOURNAL
OF THE
ASIATIC SOCIETY OF BENGAL.



Part II.—NATURAL SCIENCE.

~~~~~  
No. II.—1891.  
~~~~~

V.—*The Butterflies of Sumba and Sumbawa, with some account of the Island of Sumba.*—By WILLIAM DOHERTY, Cincinnati, U. S. A. Communicated by the NATURAL HISTORY SECRETARY.

[Received April 9th :—Read May 6th, 1891.]

(With Plate II.)

The chain of the Lesser Sunda Islands, extending from Java eastwards to Timor Laut and New Guinea, is of great interest from many points of view, but especially from the ethnologist's. For, whereas a slight tincture of Muhammadan civilization, leading to the entire loss of the native product, has made the people of the Malay Peninsula, Sumatra, and Borneo the most uninteresting of all the sons of men, and only the minutest differences distinguish the natives of Penang from those of Macassar, fifteen hundred miles away, every little island east of Java has an astonishing wealth of peculiarities.

Taking the question of religion and government, Bali, the first, is a densely inhabited island, the home of an ancient civilization. The people are of the Hindu faith, the four original castes still prevail there as they did in India in the time of Manu, and suttee, extinct everywhere else, still flourishes. In Lombok, a Hindu aristocracy rules a Muhammadan proletariat of a more recent and less pronounced

civilization. In Sambawa* there are four Muhammadan kingdoms of considerable age, while a few tributary heathen tribes, but little inferior to them in refinement, inhabit the mountains. In part of Flores, the governing race is a tribe of Muhammadan slave-traders, the curse of all that region, and the pagan mountaineers are in a more or less savage state. In Sumba and Timor there are independent heathen tribes. In Solor, Savu (better written Sau or Sawu), and Roti a large part of the population is Christian. The Dutch have no possessions in these islands, except the town of Boeling in Bali, the fort at Bima in Sambawa, and the neighbourhood of Kupang in Timor. But owing to their command of the sea, they have a considerable, and I believe an increasing influence with the trading community, and with many of the native princes. At the present moment they are trying to get possession of Middle Flores, where tin has been discovered, and, if successful, the occupation must have the happiest effect on all the surrounding islands.

From the philologist's standpoint, all the dialects from Bali to Kupang belong to the Javanese branch of the Polynesian family. In Eastern Timor and the islands beyond, some of the languages are of a totally different type, probably that of the original Negrito inhabitants.

As regards race, no part of the world excels these islands in interest. In and east of Timor, the prevalence of wavy or frizzly-haired tribes, generally of low stature, indicates the Negrito as the first occupant. In Sumba the Polynesians are still numerous, and form the ruling race, while traces of them occur in Sambawa,† and even in the mountains of the Celebes. The people of Roti are a strikingly handsome tribe looking somewhat like the better class of Tamils or Telugus, and their origin is certainly a puzzle. Mr. Wallace suggests that they may be of Portuguese blood, introduced by some unrecorded shipwreck. But they themselves say they came from Serang (Ceram). A somewhat similar race occurs at Melolo in Eastern Sumba, and, I hear, in Flores. In Savu, the people have an obvious strain of Negrito blood, but some resemble the Rotinese, while universal tradition ascribes their

* So pronounced, also sometimes pronounced Sembawa, or, if written in the Hunterian manner, Sambawá. The Dutch call it Soembawa, which is not only incorrect, but confuses it, with Sumba (Soemba). It is remarkable that the inhabitants of the island have no name for it, Sambawa being simply the name of the western sultanate. Nothing could more surprise a native of Bima, than to be told that Bima and Sambawa are on the same island. The same is true of Flores, for that pretty word is purely European, and there is no native name for the whole island. I see that the island Dutch propose to call it "Soenda" so that these three great neighbouring islands are to stand as Soenda, Soemba, and Soembawa!

† Lengota, the *glarang* or headman of Kala in the mountains of Sambawa, is a fine example of a Polynesian.

origin to Sumba, and their language scarcely differs from Sumbanese. In Sambawa, Lombok, and Bali, the flood of Mongolian immigrants has swept away nearly all traces of the original inhabitants, and the people are indistinguishable from the Malays or Siamese. The same race has entered all the islands, I do not think there is a single island in the Archipelago or the Pacific where the Mongolians have not profoundly modified the original population, whether Polynesian or Negrito. In Sumba the mixture is of great interest, because it presents the same features as in New Zealand and among the eastern and higher tribes of American Indians, namely, a race chiefly or largely Mongolian in blood, but Polynesian in language and manners, and ruled by a princely caste of genuine Polynesian blood. Till I visited Sumba I had no idea of the possibility of this state of affairs so far west. But since then I have been struck with the prevalence of Polynesian features, and even to a certain extent of Polynesian manners* among the higher tribes between Assam and Burma, namely, some of the Naga tribes—the Angamis, Lhotas and Kachhas—the Chins and the Lushais. This country may well have been the starting point of this fine race, whence they have extended their conquests eastwards to New York and Yucatan and westwards to Madagascar, and where, judging from what I saw, they may yet survive after their extermination, now so rapidly going on, is complete everywhere else in the world.

Mr. Alfred Russell Wallace has formed all the Lesser Sunda islands, except Bali, separated from Java only by a narrow strait, into his Timorian division of the Austro-Malayan region. So far as the birds are concerned, he seems to have had good reason for this, for out of 160 land-birds known from the group, just half are found nowhere else, a larger proportion than exists even in the peculiar Celebesian fauna. On examination, however, it does not appear that the group is a zoological province in the same sense as is the Celebes. In that island, a great number of peculiar species, and a certain number of peculiar genera, range over the whole island from Menado to Macassar. But the Timor group contains hardly any peculiar genera,† and the peculiar species are generally confined to one or two of its component parts.‡

* As regards language, the euphonic and structural rules are remarkably alike, but the roots of Naga words are generally as wholly different from those of the Pacific islanders, as theirs are again from those of the American Polynesians.

† Two genera of butterflies, *Ancistroides* and *Jatana*, have been described from Timor only. But I must confess that I cannot find in either any generic character separating it from its allies.

‡ On examining the British Museum Catalogue of Birds as far as completed—*Passeres* and *Picidæ*—I find their distribution as follows. No genera are mentioned as peculiar to the Timor Group, or to any part of it. Only two species are men-

It is simply a long string of islands* which has received waifs and strays from various quarters, the eastern ones, Timor, Timor Laut and presumably Wetter, chiefly from the Moluccas, New Guinea, and Australia, the western chiefly from Java. As the stream of Javanese immigrants, crossing narrow seas, is regular and unceasing, the species from that quarter have had fewer opportunities of differentiation, while the visitors from the eastward have for the opposite reason generally become distinct. This renders the fauna of the eastern islands more interesting to the naturalist, and Timor Laut, Wetter,* Sumba, and the high country of Timor, offer a field of unusual interest.

But Lombok Strait, now known in science as Wallace's Line, after the great naturalist who discovered its faunal importance, is nevertheless an important frontier, cutting off a host of Indo-Malayan forms† from Lombok and the islands eastward, and a few Austro-Malayan forms, such as the cockatoos, from Bali and Java. However, it seems hardly so deep as Mr. Wallace supposed,‡ and it is not impassable even to mammals, seeing that the tiger has of late years crossed it, and is now,

tioned as common to the Indian and Australian regions, passing through these islands, but no doubt a few additional wide-ranging forms could be added.

Timor.

Peculiar species,	29
Extending to Flores only,	5
" " Lombok only,	2
" " Bali, "	2
Indo-Malayan,	13
Austro-Malayan,	3
					—
					54

Lombok, Sumbawa and Flores.

Peculiar species,	...	15	(Flores 4, Lombok 4, in com-
Extending to Timor only,...	...	7	mon 7).
Indo-Malayan,	...	28	
Austro-Malayan,	...	0	
		—	

50

So that only seven species (or nine including the two extending to Bali) are confined to the group, as a group, and no Austro-Malayan species extends west of Timor, while the Indo-Malayan species are numerous.

Nothing whatever is known of the birds of Sumba.

* The island of Wetter seems to be wholly surrounded by deep sea, and merits examination.

† Such as the *Cyprinidæ*.

‡ The depth of Lombok Strait, as now given, seems hardly over fifty fathoms at the deepest part of the shortest line across it. There are several islands in the Strait.

I hear, making fearful ravages among the herds of ponies for which Lombok was once celebrated. As the dividing line between homologous species, Lombok Strait is probably less important than Ombai or even Sumba Straits. I should rather call it the boundary between the Indian Region and the neutral zone beyond, than that between the Indian and the Australian regions.

Sumba is one of the largest of the Lesser Sunda islands, having an area probably exceeding six thousand square miles, for the unexplored southern coast-line, drawn on the maps as concave, is really convex, giving great breadth to the island.* It is called Sumba or Humba (the S and H being interchangeable here and in Savu, which is generally called Hau by the natives) by all the tribes inhabiting it, but on the maps the more usual names are Chendana (Tjendana), Sandelhout and Sandalwood, names of the same significance, given not because, as has been stated, sandalwood is exported, but because that tree is said to be *tabu* (or *palili* as the Sumbanese say) to the inhabitants, so that if any one chances to break a twig of it, he is cut into small pieces, and scattered about under the sacred branches. At least, that is what the Malays say, but the Sumbanese, both the mountaineers and the coast-dwellers, entirely deny the existence of the tree on their island.

Deep sea separates Sumba from Flores, the high peaks of which are distinctly visible from Nangamesi Bay, but a bank covered by 50-80 fathoms of water, connects it with Eastern Sambawa, while on the side of Savu and Roti there is apparently deep sea again. No part of the coast has been surveyed even in the most cursory manner, but on account of the development of the horse-trade, the north-east coast from Laura to Réndi has become pretty well known to Arab and Bugis skippers. Except Tarimbang, which has not, I believe, been visited for generations, there is no harbour anywhere in the island. The roadstead of Waigapu or Wayapu, the chief port, is difficult of access, lying between two long coral reefs laid bare at low tide.

The aspect of the north coast of Sumba is most forbidding. Long naked headlands—Sasa, Ngarulubu, Mandolu, famous for their horses—extend far into the sea, marked with the lines of raised beaches. All this side of the island, for as much as forty miles inland and up to a height of two thousand feet, is covered with a sheet of coral overlying sandstone.† The coral must be of considerable age, and is often extraordinarily hard, reminding one of the ancient metamorphic lime-

* The southern coast of Sambawa is set down quite wrongly on the maps, as I could see from the top of Haruhasa.

† Near Kawangu the sandstone is uncovered, forming hills curiously carved and water-worn.

stones of Greece, in Bœotia and Arædia. Its surface is infinitely rough and broken, capable of destroying the stoutest boots in a few days. It is owing to this that the Sandalwood ponies develop such hard hoofs that they rarely require to be shod. Fortunately, wherever the ground is level, the coral is hidden by a coating of indurated clay like laterite, and the native paths keep to this as much as possible. A scanty growth of grass, especially the horrible spear-grass, which renders travelling almost unendurable, covers the coral. Wherever the surface consists of irregular piles of jagged fragments, bristling with needle-like points, and full of deep rifts and well-like cavities, a dry, thorny jungle grows, since horses cannot find foothold there, nor fire reach it. The grass is burnt every May or June, and for some months later, the country is as black as a coal, but travelling is easier and is usually done at this season. In some places the soil is exceedingly rich, and the population dense, especially in Melolo and Laura; but the country is everywhere dreary, and is far from green even just after the rains. Nevertheless this region, the north-east coast from Laura to Réndi, is the civilized part of the island, and the seat of all the larger states. The coast itself is generally uninhabited for several miles inland, owing to the depredations of the Endinese pirates. The heat is terrible, but the coast seems singularly healthy, and the climate is more like that of Northern Australia than of the Indian Archipelago.

Till I came to Sumba, no European had ever visited the interior. Learning from the natives that a well-wooded and watered tract existed inland, I pushed across forty miles of a desolate coral wilderness and reached a wholly different country. At Pada Dalung, and thence to Mandas* (south-west) and Karita (south-east), and, I was told, to Tarimbang on the south coast, the rock is stratified and calcareous, apparently a soft decomposed chalk, and in one deep ravine I saw some huge round boulders which may have been granitic. The interior of the island is a great plateau, somewhat hollowed out in the middle by the river Kambéra, which rises in the forests around Léwa, and in that called Kótikujara or the Horse's Head, west of Mandas, flows eastward, and near Mandas is a considerable river in deep jungle, difficult to ford, haunted by crocodiles, and much larger in volume than at its mouth seventy or eighty miles below. Indeed most rivers of northern Sumba tend to disappear on approaching the coast. The table-land is flat in general outline, but deeply cut by an infinity of exceedingly steep ravines each with a clear swift stream. Flat or steep it is everywhere the richest possible meadow land. The forests lie in great masses, and, except

* Or Mandasu; spelt Maanala in Mr. Roos's map of Sumba, which, except over a part of the north coast, seems to have been compiled wholly from hearsay.

at Tabundung and one or two other exceptional places, they are wholly trackless and serve as the boundaries of hostile tribes. West of Mandas, the country appears to descend steeply into the Indian Ocean. This slope was described to me as covered with high forest, with a heavy rainfall* and a coast so stormy as to be inaccessible during the greater part of the year. The height of the tableland of the Kambéra is usually about 1500-2000 feet; the hill at Pada Dalung must be about 2500 feet above the sea. The climate of this region is delicious. Southeast and north-west the country rises, and by its upward trend conceals whatever high mountains may be in that direction. The great isolated *massif* of Tabundung, covered with high forest, lies south of Pada Dalung, and must be about 4000 feet high. East of this is the unknown *tana maringu* (cold country) of Masu, which lies back of Melolo, and is sacred ground. No war may be fought there, and the buffalo and horse have run wild, since those that escaped thither might not be caught and brought back. Masu is the Olympus of the Sumbanese, regarded as the original home of their ancestors, and the place whither their own souls shall go after death.†

West of Pada Dalung the country again rises, and beyond Léwa Paku (*Old Léwa*, the original demesne of the present king of Léwa, who now owns all the middle part of the interior plateau), and the sources of the Kambéra, lies another "cold country," probably of considerable height and extent. This is inhabited by rude mountain tribes, not yet visited either by Europeans or by the Arab and Bugis traders. West of Perwatana and Anakala, on the border of this region, which is called by the general name of Wayéwa, lies a great forest, and then comes Kodi, beyond which the land sinks precipitously into the sea near Gaura or Garu.

A volcano has been said to exist near Tarimbang on the south-western coast. But some people of that state told me this was quite untrue. However, the mountain of Tabundung, which I did not succeed in reaching, may possibly be of volcanic origin. This district, though rather out of the way, seems to be the best accessible collecting-ground on the island.

* In Java and all the islands to the east of it, with the possible exception of Timor, the rainfall is far greatest on the southern and south-western sides. Thus at Tjilatjap (south coast of Java) the rainfall is 170 inches, at Surabaya 65. At Bima in Sambawa it is 38 inches. At Waingapu in Sumba it can hardly be more than 20, while at Pada Dalung it must be fully 100 inches.

† The Muhammadans of Sambawa, call their Heaven by the Sanskrit name *Sorga* (Swarga), and, I believe, say it lies in some distant mountains to the westward, perhaps a idea derived from the Hindus of Java. Hell is called Anaraka.

The upland forests of Sumba are less luxuriant than in Java or Sumatra, and are singularly free from thorns and underbrush, but many of the trees reach the height of a hundred feet, and some of the figs are of enormous girth. The only bamboos on the island occur in the dry valleys near the coast. Palms, except the *lontar* or palmyra, and a few arecas, are exceedingly scarce. The Endinese, who import cocoa-nuts, always destroy the germ of each nut, which perhaps accounts for the absence of this useful tree.

Of the animals of Sumba I can say but little. The natives think there are three kinds of monkeys, but I saw only the *Macacus cynomolgus*, which is very common and tame. A deer like the *Cervus muntjac* is said to be common, as well as another with large branching horns, which they call by the Malay name of *rusa*. Wild pigs abound, and a wild cat. Among birds, cockatoos are so numerous that I have seen the trees white with them; the species is the common lemon-crested one.

Among domestic animals there are pigs (*wei* or *wawi*), goats, fowls (*manu*, a Javanese word), a few buffaloes (*kalambua*, a softened form of the Malay *kribau*), cats (*kamembu*), dogs (*ashu*), and pigeons. Buffaloes are used chiefly for ploughing and for funeral sacrifices. They are the largest animals of which the Sumbanese have any conception, and a huge, ferocious kangaroo-hound, who goes about with the king of Léwa as a very efficient body-guard, has been called by the awe-struck natives the "Roaring Buffalo." Fowls are used chiefly in taking auspices, and pigs and mares are the animals generally employed for food.

Horses are the most valuable product of the island, and "Sandalwood ponies" are perhaps the best in the world, and well known as far as Rangoon and Hong Kong. They are called *jara*, a word which, like the Malay *kuda*, is derived from the Sanskrit *ghora*. The horses live unguarded in troops of twenty or thirty, each having its own range of pasture, the limits of which are carefully respected. Being very curious, they used to follow me for miles over all obstacles, but never dared to cross the ravine which bounded their beat. The colts generally follow the leading stallion (and not their dams) in a long string, which has a most absurd appearance. The mares are rarely ridden, and as in Sambawa are kept for breeding and for food.* Only stallions are exported. The trade is wholly in the hands of the Arabs and Bugis, who carry the horses to Surabaya in their own vessels at a fixed time every spring. The Sumbanese are the best rough-country riders I have

* In Sambawa, though the people are Muhammadans, a man is allowed to kill a mare on his birthday and make a feast for his friends. This is also done at the end of Ramazan; and even the Imams do it, though they may have made the Mecca pilgrimage. The Do Donggo sacrifice mares at the time of the rice-harvest.

ever seen, (and I have lived among the Turkman, Bedawin and Hiats), galloping bare back down the steepest slopes. On foot they are a singularly helpless people, and would rather ride twenty miles than walk one. They are fond of their horses and give them the most ornate names, those of mine being interpreted to me as "Beautiful Flower," "Wind in the Grass," and "Lightning." No woman is allowed to mount a horse, and I have seen a princess on foot while her attendant slaves were mounted.

The staple food in Sumba is millet (*usukannu* or *uhukannu*) and maize,* generally planted alternately, and rice (*usuberesu* or white grain), which is hard to obtain except on the coast. The wet cultivation of paddy is unknown,† though the late king of Taimanu tried to introduce it at Yawahapi-Lukukatoba. Maize is usually eaten parched. Meat is only eaten on great occasions, and there are scarcely any vegetables. Curiously enough, the use of toddy (palm-wine) is unknown, though so common in Flores, Savu, and Roti, and even in the Muhammadan parts of the Celebes. Considering the wealth of the people, and the cheapness of Java rum, the Sumbanese are a sober people, and most of the mountaineers have never tasted spirits. The use of betel is universal. Salt is very scarce and dear.

The people of Sumba do not probably number less than 100,000, and perhaps much more if Laura and Melolo are really as populous as they are said to be. A small colony of curly-haired Savu people are settled at Waingapu and Kabaniru, and a similar race at Memboro. Some of the Melolo people are said to resemble the Rotinese in feature. Otherwise, the bulk of the people may be said to be Mongolians resembling the Javanese, with a Polynesian aristocracy.‡ The former are

* Maize is probably a recent introduction, but I could hear of no tradition on the subject. A common species of sorghum growing in marshes is called "wild maize." In many of the islands, the word *jawa* or Javanese is applied to maize, showing whence it came. In Sumba the word is *water*, but in Savu *water-jawa*, in Roti *mtela*, in Timor *pela*, in the Moluccas *milu*, in Ende (Flores) simply *jawa*, in Roka (Flores) *hai*, in Sambawa *baso*. The word *jawa* is applied to anything foreign. Europeans are called "white Javanese," and I was generally known in Sumba as *umbu maremba jawa* or the King's son from Java.

† The Do Donggo of the mountains of Sambawa have some of the finest wet paddy fields I have ever seen. Yet they are far inferior in capacity to the Sumbanese, and preserve a curious memento of their recent savage state in an annual three days' pilgrimage to the mountain-tops, where they sleep in the open and live wholly on what game they kill, leaving the villages guarded by the dogs tied up in the houses.

‡ Some of the western hill-tribes may belong to a lower race. The Kodi people are said to be of short stature, and to turn the toes inwards in walking, especially the women. To "walk like a Kodi woman" is a staple joke, appealing strongly to the Sumbanese sense of humour.

the same as everywhere. The latter are tall, light-brown men, of somewhat slender, but graceful, and manly proportions. The face is rather long, with a Roman nose and a finely-moulded chin; the hair is straight, rather dry and stiff, and a beard is not generally worn. The women are often of a refined and high-bred, though somewhat grave and melancholy beauty, contrasting strangely with their barbarous condition. The quiet dignity of the men is in striking opposition to the innate vulgarity of all Mongolians from Turcomania to Malayana.

The Sumbanese, both men and women, wear a large loose mantle of Manchester cotton dyed black in the mud of the rivers. The women wear also a short black skirt, and on gala occasions a black jacket tastefully embroidered with beads and small cowries. The men wear a waistcloth, a turban, a huge ivory armband, and a heavy belt like that worn by the Greeks and Albanians, containing their krisses and *parang*. They always go about with a square mat-work satchel, generally of very pretty design, containing betel. They exchange betel with everyone they meet outside their village, as a sign of peaceful intentions. I had to carry a supply about also, and never dared to decline it, though it is not at all nice, for the Sumbanese could only explain a refusal as a sign of hostility, just as Bedawin would excite at the refusal of salt. Bows are unknown in Sumba, and so are fire-arms, but a man goes nowhere without two spears, which are never laid aside for an instant. In the remote district of Mandas, I was amused to see that my visitors had covered their spearheads with sheaths tied on with thongs, as if to reassure me, reminding me of the old Norse custom.

“Thereat was the Wrath of Sigurd laid fast in a silver sheath,
And the peace-strings knit about it, for the blade was fain of death,
And 'tis ill to show such edges to the broad blue light of day,
Or to let the hall-glare light them, if ye list not play the play.”

The Kaumbéra language is understood over the greater part of the island, but Gaura and Laura in the west have languages of their own, and the Memboro dialect is very distinct. All these are closely allied to the Javanese and the languages of Sambawa and Flores. I have taken vocabularies of a number of these, which I hope to publish some day.

Strange to say, Sumba has a currency of its own in the shape of fine copper wire very intricately plaited and cut into lengths of two feet, worth half a rupee each. The ugly, fish-shaped earrings of gold beaten out thin, are always of the same size and value (about a dollar), and are likewise used for money.

The women have spinning-wheels and weaving-frames, and make cloth, especially blankets, generally white with curious figures of fish,

tortoises, prawns, ships, men, deer, etc., but all so conventional in form and so harmoniously arranged that the effect is good. The men also make nets and ropes, both of excellent quality and largely exported, and at Kadungu (Memboro) good pottery is made. The chief exports are horses, slaves and edible birds-nests.

There are three castes of Sumbanese, the *maremba* or lords, the *kabisu* or freeborn citizens, and the *towata* or slaves, the latter being the most numerous.

The ruling classes marry chiefly among themselves, and are inter-related in a most puzzling fashion. Marriages are arranged by the parents, and are of two kinds. If the wife is bought, whether with money or with service, she enters her husband's tribe. In this case she is his property, and he can kill her if he likes. If he pays nothing, he enters her tribe; but this is less usual. Polygamy is not common, but if a man's sisters-in-law remain unmarried, I believe they are after a time considered as his wives. The Sumba women make faithful wives, but before marriage incontinence is universal, and every girl, slave or princess, has her price. Infanticide and abortion are very common, and it is probably largely for this reason that the population is not increasing. Islam always, and Christianity often check this evil, so that the population is large and increasing in Muhammadan Ende and Sambawa, and again in Christian Roti and Solor. The old are treated with great respect. The Sumbanese struck me as a brave, honest and truthful people. But they are too proud to work for others, and will never become a thriving agricultural race like the Javanese.

Exogamy is usual, and the rules of intermarriage are often inconveniently complicated. For instance, I hear that Kanata men can marry only Lakoka women, and Lakoka men only Soru women. Now Lakoka* and Soru are small independent states in the interior, while Kanata (or Lubu) is fifty miles away on the coast of the Taimanu state.

Apart from the wars of extermination waged now and then by the great chiefs, disputes are continually going on between neighbouring tribes, generally concerning boundaries, horses, or women. They are usually settled without much bloodshed in the following manner. The men meet in a meadow, and form two lines on horseback. Then the chiefs recite war-songs, and make speeches, and the two sides exchange

* In 1886, shortly before my visit, the king of Léwa sacked Lakoka, in alliance with the Ende slavers. The men were killed, the king took the horses, and the women and children were carried off to Flores as slaves. This is the usual way in which the Endinese do business. It is to be hoped that the Dutch troops now in Flores will put an end to this murderous little state. Its supremacy in this region is owing to its possession of ships and rifles, of which the Sumbanese have none.

abuse and defiance in the Homeric fashion, till the proper degree of excitement is reached, upon which they charge, fighting with spears and shields. As soon as anyone gets speared, his side acknowledges itself beaten and pays a fine, while the others celebrate their victory with much noise and feasting. The horses on these occasions are decorated with collars of white horse-hair, and immense frontal tufts, giving them a most ferocious look, and are said to enjoy the fighting thoroughly. Sham fights, very similar to the real ones, and quite as dangerous, are often held. But horse-fights are the characteristic amusement of Sumba. Two stallions and a mare are placed in a little enclosure, and the former fight till one is dead. On great occasions there is dancing, generally performed by women, and sometimes a poet will sing the praise of his forefathers, exhibiting the skulls of their conquered enemies which have descended to him. The musical instruments in use are drums, gongs, and a guitar with two copper strings.

The dead are buried,* household articles being broken and thrown into the grave as in the Nicobars. A large oval horizontal slab of stone surrounded by small upright ones, marks the grave. The bodies of chiefs are exposed on the mountains for months after their death. When a propitious time for the funeral comes, a great feast is held, many buffaloes, pigs and mares are killed and eaten, and a number of slaves, both men and women, are strangled and thrown into the grave.† When I was in Sumba, the body of the late king of Taimanu had been lying exposed at Semparingu for more than a year.

I cannot say much about the religion of Sumba. The island presents a remarkable contrast to Sambawa in this respect. In Sumba, though there are a few *ratus* or professional magicians of little influence, the chiefs are the real religious leaders, and it seems to me that the union of church and state in the hands of practical men managing large temporal affairs has kept superstition in bounds. In Sambawa, both in the heathen and in the Muhammadan parts, the *juhís* or sorcerers are the descendants of the old local chiefs, now replaced by a centralized bureaucracy. Reduced to mere tricksters and jugglers dependent for their food on the popular faith in their magic powers, they have made the people as superstitious as any in the world. The same is the case

* The Do Donggo in Sambawa are buried sitting, but I can find no note of the Sumbanese custom.

† On the death of a Sultan of Mbojo (Bima) in Sambawa, 199 buffaloes are sacrificed. A new flagstaff is raised by his successor, and a slave is said to be strangled and buried beneath it. This, if true, illustrates the extreme conservatism of the East, for the people of Bima may be almost called a civilized race, and have been Muhammadans for some centuries.

with the heathen Do Donggo. Here the sorcerers hold a higher rank as *jubi Perafu* or priests of the god *Perafu*. But all temporal power is in the hands of the *glarangs* or headmen. The result is, that the *mori*, or ancestral spirits, and the *héncha*, or demons, are never out of peoples' minds, everything seen or done has some good or evil significance, there are sacred trees, mountains, springs, stones, and animals, while every spot is the scene of some absurd legend. Each village has its priest's house, priestess's house, and its *uma Perafu* or house of *Perafu*, closed, empty and of very archaic make. Belief in the evil eye, in the unluckiness of a thousand acts and signs, in the constant presence of evil demons, and in the disastrous effect of anything unusual or uncustomary, make these people the timid, unhappy race they are.*

The Sumbanese are said to worship one greater god, described as *umbu walu mendoku* or *he who makes all*, who owns all the sandalwood. Also two deities called *umbu awan*, lord of heaven, and *umbu tana*, lord of earth, to whom worship is paid at harvest time, and rice, pigs, horses and buffaloes sacrificed. They also believe in evil spirits, and the huge fig trees in some of the villages are apparently held in veneration. Certain things also are sacred, and hence *tabu* or forbidden (*palili* in Sumba, *leo* or *ponali* in Timor, *perafu* among the heathen of Sambawa). Though the crocodiles receive no regular ceremonial worship as in Roti,† the Sumbanese nevertheless throw them meat, saying, "Don't eat

* For instance, when I was at Kala in the Donggo country, the *jubis* kept praying and sacrificing all night to prevent evil resulting from my stay there. At Pelunto the people threatened to abandon their homes when they heard I was going to climb Harahasa, the chief mountain in those parts. And when I returned and nothing happened, they said I had not really done it, just as when the alpinist climbed Ararat, the Armenians would not believe it, because he had not seen the Ark standing intact on the summit, as St. Mesrob had seen it in his dream. At Oo, the *jubi* declared that my visit had caused the terrible rains we had then. Out of revenge I rolled my eyes at him tragically, and repeated the first stanza of "Simple Simon met a Pieman" once or twice when I met him, upon which he fled the country. Such things are quite impossible in Sumba, and their own little devices for keeping off the evil spirits are performed in rather a sceptical mood. As in India they snap their fingers when some one sneezes. If a young man hiccups, they box his ears, if an old man, they ask him respectfully why he did it, to which he calmly replies that he never did, and there is a general smile.

† The following story was told me by Mijneer Teffer, whose wife, a remarkably beautiful woman, now a Christian, was the daughter of the king of Hai in Roti. There is a caste of priests of the crocodile there. When they want to travel by water they call the crocodile, and he carries them wherever they wish on his back. On a certain day they go down to the bank, and call the crocodile, describing to him their rank and duties. When he comes out, they take him up and carry him, with a band of music and an applauding crowd, to his temple. There they give him rice and sweetmeats, put a robe on him, and begin praying over him. He dislikes the praying

me, but eat such a man, my enemy." But the chief Sumbanese deity is Merapu (the name is obviously equivalent to the Sambawan Perafu), who is the hearth-god, a kind of aggregation, I imagine, of the ancestral spirits. But some say he is a man who lives in Masu, and is a kind of intercessor with the great gods for men, and especially for kings. He is described as black in hue, for when I asked why the Sumbanese dressed wholly in black, they replied that it was Merapu's colour. The largest insect in the island, the black butterfly I have named *Papilio merapu*, is sacred to him. They pray to him in the forest, placing betel, siri, and a bit of gold or silver on a leaf, and setting it on the ground say "Merapu, give me this and that, pasturage for my horses, rain for my maize, and vengeance for my wrongs." Slaves pray to Merapu that the king may live a hundred (*ngasu*) years, for they are afraid of being sacrificed at his death.

The houses are large, with a thatched roof pointed at the top, and a floor raised five or six feet above the ground. Inside, the fireplace is always surrounded by four posts. That on the right hand on entering is called Merapu's post, and the enclosure is sacred to the god. Oaths are taken by laying the hand on this post, and no one is allowed to sing or play the guitar indoors when a fire is on the hearth.

Houses are gathered in a *paraing* (generally called *paré*) or village, or in a *negeri* (Sanskrit, through the Javanese) or town. The political unit, at least in the interior, is a group of open villages, protected by a fortress (*kota*, Sanskrit through the Javanese). Thus Watupéli is the central fortress of Melolo, Kamanu of Mandas, and Lambanapu of Kambéra. On the plains, these fortresses are defended by intricate cactus hedges, but I was told that in Laura (I think) towns are strongly walled with stone, as is the case with some of the hill forts elsewhere. These last are often very striking. Lateng in the Taimanu state, is built on a sharp spur of the mountains, the *col* connecting it with the main mass fortified by wall after wall. On the other side, the hill descends at a very steep angle to the river a thousand feet below, and this almost inaccessible gorge is so industriously cultivated as to be a perfect nest of verdure in this dreary country. In times of peace these forts are often left almost unguarded. Once I lost my way at nightfall in the Kiritana district, but when the moon rose I struck a path, and rode through a country of alternate thorny jungle, and meadows studded with great upright blocks of coral-like tombstones,

and struggles, so that it is necessary to quiet him with more food, and begin the prayers again. When the prayers are said, they carry him back to the river with music and dancing, and when he enters the water all the other crocodiles rise up and pay homage to him as their king.

worn into a thousand fantastic shapes like Gothic gargoyles. The impressive uncanniness of this place I cannot describe; my horse was in an agony of terror. Finally I came to a citadel on a steep crag, and climbing the wall in constant expectation of attack, I found a mass of huge fortress-like houses of stone, bigger than any built nowadays. After I had gone over most of them and found them full of grain and household utensils, but without inhabitants, I finally stumbled on three very old men, who were speechless with amazement at seeing me. They were in charge of the place and had not had a visitor for months.

The little district with its sheltering citadel was probably till modern times the only kind of state in Sumba. The recent evolution of governments like Léwa and Melolo has as yet had but little influence on the people of the interior.

On the coast, one can now ride from Waingapu to Melolo without receiving anything from the men he meets but polite salutations. In the interior, even in the middle of the Léwa dominions, I never met a native not belonging to the village where I was staying, but we both prepared for battle, and spear and revolver were held in readiness till we had exchanged betel. Twice I was within an ace of being speared, because I came on men suddenly in the forest. When two parties meet, they halt when yet a long way apart, dismount, and drive their spears deep into the earth as a sign of peace, then exchange a "cooey" (the well-known Australian cry, much used in Sumba), and yell out a question or two. Then two men advance, one from each party, and exchange betel, after which the others come forward warily, keeping a good grip on spear and shield. In spite of the tyranny of the kings over their subjects, and their occasional ferocity to conquered enemies, centralized government of any kind is better than this constant distrust of one's neighbours. The northern kings of Sumba have greatly strengthened their power by making it hereditary. The *umbu maremba*, or heir-apparent, is a power even in his father's lifetime. Whereas in Ende, Roti, Savu, and in the less advanced states of Sumba, the king's successor is elected by the nobles from the royal house.

The most powerful of the Sumba kings are those of Léwa (who holds *Kambéra* by right of conquest), and Melolo (who ruled half the island a generation ago), whose son rules at Petawang. West of Léwa come Taimanu, Kapundu, Palmédo, Kadungu (or Memboro), and finally Laura, which is said to be of great interest, but which has not been visited by any European. The Dutch claim allegiance from the Savu and Timorese immigrants at Waingapu and Kabaniru as well as over a few Arabs, Bugis, and Chinese who trade at Waingapu. But these all pay tribute to the king of Léwa, and the only time they

refused it, he plundered the village, and drove the Dutch agent out of Sumba. When the unhappy Achehese war is over, it is to be hoped that the Dutch will pay some attention to this fine island, hitherto neglected. Owing to the absence of fire-arms, it could be subjugated by an insignificant force; the horse-trade properly developed would prove a mine of wealth; and under settled government the island would be as prosperous as Roti or the Minahasa. However it may have been in the last century, no people in this can rule semibarbarous races better than the Dutch—when they think it worth their while.

Some idea of a Sumbanese king may be gathered from an account of my visit to Tunggu, king of Léwa, which I made with my kind friend Mr. K. H. de Roo van Alderwereld. The king was then at Kawangu near the coast.

We rode from Waingapu across the Matawai, past the Savu settlement of Kabairu, and reached the Kambéra river, where women were dyeing cloth in the black mud, and a frizzly-haired Timorese was fishing in a canoe. Forging the Kambéra and the Palamenjéli with much difficulty, we reached Kawangu, a village of thirty large houses. A narrow path wound zigzag past three great concentric hedges of cactus guarding the place, and brought us to the king's house. He was an ugly old man, well over six feet high, wearing nothing but a dirty waist-cloth, his skinny limbs uncovered. His long hair was white and knotted over the nape of his neck, his eyebrows were black and stood out from his head, the hairs more than an inch long, shading a pair of singularly bright, unsteady eyes, and giving him an extraordinary appearance. He shook hands with us feebly with his paralyzed left hand, holding his spear in his right all ready for action, for he is forever suspecting some treachery. He had two mares led up, and drove his spear into the throat of each with a wild shout; then he killed a pig for us, saying, politely, "Pork is for kings' sons, but mares' meat is good enough for soldiers." For the mares' meat was for his body-guard, a number of handsome and splendidly-formed young men, with whom he was at that time hoping to conquer the whole island. Later on, we saw him standing among them ladling the boiling meat out of a huge pot, and saying, according to our interpreter, "Eat, my children; he who fights shall eat meat; let slaves stick to millet."

He took us to see his elder brother, who was slowly dying of cancer, and had therefore given up the throne to him many years before. When my trip into the interior was suggested, he coolly made a counter-proposal that we should both go and help him in his campaign against Anakala. For he has no fire-arms, and no doubt thought a rifle or two would have great effect on those sturdy mountaineers. On our

declining this, he invited the leading nobles into the council-house and consulted the auguries about my journey, examining the liver of one fowl after another till one suited him. Every stain or flaw in the liver has its own meaning, and I was surprised at the acuteness of his inferences regarding them. Sometimes he would consult some of the old men, who seemed quite familiar with the science, and evidently considered it as reasonable and reliable as the multiplication table. The king is held the best haruspex in the country, and is also proud of his skill in causing rain or drought. He remarked incidentally that he would give me good weather for my trip. We sat up till late listening to a minstrel singing a song of the king's composition, twangling a guitar the while. The king sat glancing sharply at us alternately, some nervous affection keeping his head and hands in constant motion. Meanwhile his followers sat in a circle round him, looking singularly dignified and austere. They wore black mantles and turbans, their belts were full of handsomely-mounted weapons, their fine Roman faces perfectly quiet, in striking contrast to their master's, whom they so surpassed in dress and bearing. Still I think I understood then somewhat of the power which made this ignorant savage a ruler of men, personally more revered than any sovereign in civilized countries.

The butterflies mentioned in the following list were taken in 1887. The Sambawan specimens are in the hands of Mr. B. Neumoegen. The Sumba ones have remained four years in my possession, during the course of which most of them have suffered greatly, and many of the best have disappeared or been destroyed, including several uniques, of which I have only descriptions made at the time. As, however, no one may visit the interior of Sumba again for many years to come, I shall include these descriptions here. Owing to my small knowledge of Moluccan and Timorese butterflies, and the absence of specimens for comparison, my work is necessarily imperfect. When described, the Sumba butterflies will be sent to M. Charles Oberthür. Of the Sambawa butterflies I cannot give a complete list, and I am obliged to omit most of the *Euploas* and some others. At the time I was there, I counted about 140 species taken in each island. In this list I number only the Sumba species, those from Sambawa being introduced chiefly to illustrate the geographical distribution of the species, and to describe a few novelties.

There is a considerable Austro-Malayan element in the Sumba butterflies, but very few of these forms reappear in Sambawa, *Ornithoptera naias* and *Danais (Nasuma) haruhasa* being the most remarkable of those that do. The list scarcely does justice to this element in

Sumba, since a *Doleschallia* and a *Charaxes* probably of Moluccan type escaped me, and of a *Hypolimnias* apparently near *H. pandarus* I made no description, and the specimens are lost.

A few Papuan or Timorese forms occurring in Sumba do not extend to Sambawa, such as *Radena oberthurii*, and the above-mentioned *Doleschallia* and *Hypolimnias*. In one or two cases a species occurring with little change from Java to Timor has a wholly different representative in Sumba, as in the case of *Papilio marenba*. Ten Sumbanese forms are represented in Sambawa by other allied species, namely, six *Tanaïde*, three *Pieridæ* and a *Papilio*. Nine species of *Danaïds* occur in Sumba and nine in Sambawa, and of these six are the same, and three different. Ten species of *Euplœa* (of nine different groups) were taken in Sambawa, and only six in Sumba (perhaps on account of the continual rain in the interior), of these only one, a large *Salpinx*, was common to the two islands. The dominant *Euplœa* of Sumba was apparently *E. lewa*, that of Sambawa seemed to be the Javanese *E. (Selinda) eleusina*, and both have their mimics. *Trepsichrois*, of which a species is peculiar to each island, appears as a mimic, and rare, whereas further west it is usually a dominant genus.

Information regarding the seasonal forms of the *Satyridæ*, will be found under the head of that family. It will be seen that in these islands the non-ocellate brood appears when the ocellate brood appears in India, as might be expected, the seasons there being similarly reversed. In Sambawa I reared both forms of *Melanitis leda* from the wet-season one, by keeping a wet sponge along with the chrysalids in one box, whence only the ocellate brood was obtained.

My discovery of the dimorphism of these insects, made in 1882-3, and my theory regarding its cause, have now received confirmation from all sides, and may be regarded as proven.

My collecting in Sambawa was in the eastern part of the island in the sultanate of Bima or Mbojo, and was unfortunate, owing to the heavy and unseasonable rains. Owing to the assistance of Mijuheer A. C. de Heer, Controleur of Bima, for whose kindness I offer my best thanks, I was enabled to visit the mountains west of Bima, the Sultan sending his brothers to arrange matters for me. These mountains, which are of some height—two of the peaks, Haruhasa and Ndindi exceeding five thousand feet in elevation—are in the district of Bolo, the capital of which is Sila, and are inhabited by a timid race called the Do Donggo Bolo to distinguish them from the Do Donggo Kai near Prado. The higher parts of the mountains have a very wet climate, and are partly meadow and partly forest. The latter is exceedingly rich and luxuriant, resembling that of Sumatra or Borneo, but is of no great height on account of the violence of the wind.

My impression is, though I can scarcely prove it by lists of species, that the insects of this mountain region, are almost purely Indo-Malayan, or at least more so than those of the coasts. If this is true, it does not at all agree with Mr. Wallace's belief that the Indo-Malayan element is of recent introduction. As these mountains are very easy of access from Bima, where steamers stop every month, and as travelling in the island of Sambawa is safe and pleasant, it seems a pity that some competent ornithologist does not investigate the birds of this district, which ought to afford many novelties.

Family DANAIDÆ.

1. *SALPINX MEIZON*, n. sp.

Male, above, forewing rich brown with blue reflections, a short slender pale lilac spot in the interno-median space, a costal spot and seven large inner-submarginal ones, light blue with purple reflections, generally pointed outwardly and inwardly, the second largest, separated only by a vein from the first, which is prolonged costally, the last with an obscure streak below it. Hindwing with the blue gloss much less conspicuous, the velvety patch pale ochreous externally, darker internally; two or three small subapical lilac spots. *Below* dark brown, both wings with the cell and the spaces just beyond it much paler than the outer part. Forewing with a costal lilac dot, and sometimes one or two subapical, a larger one in the lower median space; below the lower median vein a large ochreous area, pale brown in the middle, extending below the submedian vein, enclosing a short sericeous band. Hindwing with a varying number of minute inner-submarginal lilac spots subapically, and sometimes two or three still smaller outer-submarginal ones subanally.

The basal tuft of the male is very large, light reddish at base, fuscous outwardly; the outer tuft white, very short.

Expanse of male over four inches, the female still larger. Type from Sumba, where it is scarce, also occurring in Sambawa, apparently unchanged. It differs from *S. leucostictos* and *pasithea* in the pale internal areas of the underside, and in the absence of most of the submarginal spots. It somewhat resembles *S. viola*, Butler, from the Celebes, but lacks the blue spots on the hindwing above.

SALPINX (*SELINDA*) *ELEUSINA*, Cramer.

Sambawa, very common. An undescribed *Isamia* occurs in Sambawa.

2. *SALPINX* (*CALLIPLÆA*) *SUMBANA*, n. sp.

Above, forewing brown, slightly glossed with blue, the outer margin paler; a costal white spot, and a submarginal row of nine others, the

first six more or less fused into a single mass, the first and sixth sometimes separate, the veins dark, the fourth (above the upper radial vein) much the largest, the seventh and eighth (between the median branches) minute. Hindwing whitish costally, unmarked. *Below* uniform brown, forewing with the spots reduced, those between the median veins generally absent. Hindwing with 4-6 small inner-submarginal spots subapically, the last minute.

Sumba, coast and interior. It seems quite distinct.

SALPINX (CALLIPLÆA) SAMBAVANA, n. sp.

Above, forewing with seven large lilac spots centred with white, the second and last largest, the upper ones sometimes slightly connected, Hindwing with several lilac spots subapically. *Below* both wings with two nearly complete submarginal series of small and delicate white spots, the outer ones minute and not extending to the apex.

Sambawa, one of the numerous local forms of this group. A very distinct species, *C. hyems*, Butler, occurs in Timor.

I have noticed that both *C. sumbana* and *C. sambavana* are occasionally found with the first subcostal vein united to the costal one, showing at the same time the relations this group has with *Hestia*, and the small value of classifications based wholly in venation.

3. STICTOPLEA MELOLO, n. sp.

Male, above rich dark brown, with blue reflections over all the forewing except the extreme outer margin; four lilac spots, sometimes centred with white, form a narrow subapical mass, the fourth well separated, generally a fifth below it, and occasionally a sixth. In the male the sex marks vary; in the specimen before me, the upper one is shorter than the lower, and only half as broad. Hindwing with two, sometimes three, subapical spots. *Below* rich brown, darker at the end of the cell and on the disc beyond it; a distinct white spot bordered with lilac at the end of the cell, and two near it in the median spaces; generally traces of a few other dots, especially subapically on the hindwing. The semicircle of spots beyond the cell of the hindwing is represented by obscure darker touches. The female is much paler.

Sumba, common; one of the numerous local forms of this genus.

4. STICTOPLEA LACORDAIREI, MOORE.

Sumba, common. The species was described from Java.

EUPLEA (TREPESICHOIS) DONGO, n. sp.

Male, above, forewing outwardly shining blue, basally blue-black, with two rows of pale blue submarginal spots, the outer of about eight

or nine dots, not extending above the radial veins, the inner of seven large spots placed irregularly, the first four and the last three in line, the first minute. Hindwing bronzy brown with a slight bluish lustre subapically, the velvety area extending below the upper median vein. *Below* chocolate-brown, forewing with a purple cell-spot, a costal and usually three or four minute discal spots, and one or two submarginal dots near the lower angle. Hindwing with a large paler area subapically around the subcostal branches; usually a few submarginal bluish dots. The wings are short and broad, quite unlike those of *T. midamus*.

Female with no trace of blue. Above, forewing with whitish markings, one geminate in the cell, (besides a basal pale streak there), one costal, two approximate beyond the cell, three discal, two outerdiscal spots, besides a few pale ones subapically and an interno-median pale streak. Hindwing with the white rays larger and more distinct than in *T. midamus*, the submarginal spots obscure. *Below*, all the markings are white and well-defined.

Nearest *T. mindanaensis*, Semper, from the Philippines, but the female and the underside of the male are very different.

Taken sparsely in the mountains of Sambawa, in the Donggo country.

5. *EUPLEA* (*TREPSICHOIS*) *ELWESII*, n. sp. Pl. II, fig. 1.

Female, above brown without any blue reflections, a pale longitudinal streak in the cell, a small round whitish spot at its end, two similar discal spots between the median branches, a long bent pale streak in the upper part of the interno-median space, a subapical area of five large white spots, separated by veins only, from the costa to the upper median vein, the lower spot large and quadrate. Hindwing with four white streaks occupying the greater part of the cell, two good-sized elongate quadrate markings at the base of the spaces between the upper subcostal and the radial vein, small spots beyond the cell below the radial vein, and above the lower median, narrow whitish streaks in the submedian space, and two in the internal space. *Below*, there are obscure subapical dots on both wings, and a few marginal ones nearly obsolete. The white masses of the forewing and the hindwing are unchanged.

This species, the most aberrant of the genus, obviously mimics *Radena oberthurii*, a dominant species in Sumba. No male was seen, and only two females, both now in bad condition, were taken at Koloki and Mandas, Central Sumba, 2—3000 ft.

I name this butterfly after Mr. H. J. Elwes, the well-known lepidopterist and ornithologist.

6. *EUPLEA* (*RASUMA* ?) *LEWA*, n. sp. Pl. II, fig. 2.

Male, above, forewing dark brown, the outer part paler, especially near the lower angle; a broad sericeous streak in the interno-median space, nearly half an inch long; a subapical mass of four blue-bordered white spots, with a minute one above them, the first two small, the third large and quadrate, the fourth smaller, pointed inwardly; a whitish point on the costa, another obsolescent discally in the upper median space. Hindwing unmarked, much paler than the forewing, especially outwardly. *Below*, forewing darkest on the disc and in the cell, the subapical band somewhat reduced in size, one bluish spot in the cell, one near the costa, and three on the disc, the lower one yellowish. Hindwing, with a space above the terminal part of the cell much darker than the rest, a pale band round the disc, one bluish spot in the cell, five or six dots beyond it, and eight rosy ones in an irregular line across the disc, two in each median and in the lower radial space, and one in each of the two next spaces. Expanse over three inches.

Sumba, apparently a dominant species. Like the next species it is separated from all allies by its white subapical band.

I did not find any species resembling this in Sambawa, though a species of *Penoa* occurs there having a somewhat similar sericeous band above. I took only a single male at 2000 feet. A wholly different species, *G. baudiniana*, Godart (*orope*, Boisduval), occurs in Timor, having the hindwing broadly whitish. *E. lewa* is apparently of Papuan affinities.

7. *EUPLEA* (*CRASTIA* or *VADEBRA*) *PALMEDO*, n. sp. Pl. II, fig. 3.

Closely resembling the preceding species. Male, forewing dark brown above, paler outwardly; a white, rather quadrate, subapical mass diffused at the edges, broken by three slender dark veins. Hindwing nearly white above the upper subcostal vein, the rest brown, the outer discal and subanal area much paler. *Below*, the pale areas are more obvious than in *Eupleia lewa*. Forewing with a bluish-white spot in the cell, and two in the disc beyond, besides traces of two streaks in the interno-median space. Hindwing with one spot in the cell, a semicircle of six minute ones beyond it, and a row of seven or eight larger ones in the yellowish discal area, all but one arranged linearly; only one or two submarginal dots visible.

Sumba, coast and interior. It is much less common than *E. lewa*, but as the *climena* group to which it belongs is in most places a dominant one, I am unwilling to believe it a mimic of that species. The species is a very distinct one.

An allied form occurs in Sambawa, with the margins broadly

whitish as in *E. clinena*, and without the conspicuous subapical white band of the forewing.

EUPLOEA (*CRASTIA* ?) *DEHEERII*, n. sp.

Male, above dark brown, with a slight violet gloss, the outer margin broadly paler, not glossed. Forewing with an irregular series of seven small white outer-discal spots, the first three subapical, cordate, separated by veins, the fourth and fifth beyond the line of the others, the fifth minute, the sixth and seventh in the median spaces, distinct, equal, the upper elongate; an obscure dot near the base of the upper median space. Hindwing with three good-sized white subapical inner-submarginal spots, and eight or nine obscure outer-submarginal dots, which do not reach the apex. *Below*, the cell and inner part of the disc of both wings dark brown, the rest paler, with a bronzy gloss. Forewing with a spot in the cell, one costal, four or five discal violet-white spots, seven inner-submarginal white ones arranged as above, and eight or nine outer-submarginal ones minute. Hindwing with a cell-spot, a semicircle of seven inner-discal violet-white spots, and one of nine or ten outer-discal ones, mostly white, some lilac; about twelve small submarginal white spots, larger than those on the forewing.

Like my *Euploea oceanis* from Engano, this species has a large, somewhat velvety, pale brown patch on the underside of the forewing. This lies along the internal vein for more than half its length, about three quarters of it lying above that vein. At the base of this there is an obscure whitish patch, chiefly below the internal vein, while parallel with it is an obscure longitudinal sericeous streak placed below the lower median vein. The hindwing is whitish apically and costally, with a pale brown area surrounding the subcostal veins, entering the cell and the space below the costal vein, extending narrowly along the subcostal veins three-quarters towards the margin.

The species perhaps belongs to Mr. Moore's genus *Gamatoba*. I took it in the mountains of Sumbawa, and name it in honour of my friend Heer A. C. de Heer, Controleur of Bima.

Another species from Sumbawa, belonged, I think, to the subgenus *Tronga*, making ten *Euploea*s in all from that island, some very rare. Only six were taken in Sumba. At a favourable season, I believe Sumba will yield a far larger number of species than I obtained there.

8. *DANAIS* (*LIMNAS*) *CHREYSIPPUS*, Linn.

Sumba, Sumbawa. Somewhat intermediate between typical *chrysippus* and *D. bataviana*; colour bright red as in *chrysippus*, the white

spot at the end of the cell absent, the black border of the hindwing broader than in Indian specimens, and inwardly diffused.

9. DANAIIS (SALATURA) GENUTIA, Cramer.

Sumba, Sambawa. My Sumba specimens are somewhat intermediate between *genutia* and the Javanese *D. intensa*. There is only one submarginal line of spots on the hindwing above, the small subapical spots are nearly obsolete on the forewing above, and below, the red area in the upper median space is present or absent. The general colour is not so dark as in *intensa* and the species larger.

10. DANAIIS (SALATURA) LITORALIS, n. sp. Pl. II, fig. 4, *underside*.

Male, above black, a narrow pale ferruginous band in the cell, a much larger one occupying most of the interno-median space, and another in the lower median space, extending much further outwardly, slightly irrorated with white scales in the middle; a narrow oblique white subapical macular band from the costa; the spot above the upper median vein much beyond the line of those above it; one below it, large; three costal marks, a dot beyond the end of the cell, five marginal and three submarginal spots in the median spaces, one apical and one at the lower angle, all white. Hindwing black, a broad quadrate white band across the disc, and the end of the cell as far as the lower subcostal vein, scarcely reaching the submedian scent-gland, continued outwardly by obscure ferruginous rays, the veins there widely black, an outer row of submarginal white spots, with two inner ones subapically. *Below*, forewing with the ferruginous cell-striga obsolescent, two rows of minute submarginal spots subapically. Hindwing with two complete rows of submarginal white spots, and a few costal ones, including one basally along the lower side of the costal vein, the white area more broken, its discal spots outwardly incised, the cell-spot occupying two-fifths of the cell; most of the disc, including the base of the cell and the costal and subcostal spaces, ferruginous, edged with black; the veins all dark, the submedian and internal veins black bordered with white for most of their length; the outer black border glossed with chocolate-brown.

Nearest *Danaïis abigar (chionippe)* from the Philippines, figured by Mr. Distant from Province Wellesley, Malay Peninsula, though that locality seems to me rather dubious. It differs in the smaller ferruginous area on the forewing and smaller white area on the hindwing. From *D. fulgurata*, *afinis*, *aruana*, etc., it differs in the absence of white in the interno-median space of the forewing.

Sumba, scarce. An apparently identical form is common on the dry-coast of Sambawa.

In the figure the forewing has been drawn much too short.

DANAIS (NASUMA) HARUHASA, n. sp.

Male, forewing extremely long and falcate, deep fuscous above; a long obscure reddish streak extends along the lower part of the cell, another more distinct, lighter in colour, and enlarged outwardly, in the interno-median space; a third, obscure, between the lower median veins; a small round discal whitish spot in each of the spaces below the upper radial vein, the second a little nearer the base than the others are; two small whitish spots, one on each side of the lower radial vein, are sometimes present (especially in the female) just beyond the cell. Hindwing with discal streak of pale brown, slender and rather obscure, a larger one in the cell: two rows of white submarginal spots, the outer subanal only, minute, the inner obsolescent near the median veins. *Below* dark brown, the apex of the forewing rufous, the pale reddish markings of the forewing somewhat larger, those of the hindwing larger and dull leaden-white in colour, reddish only at their truncate tips, the submarginal series complete and nearly equal.

Forewing with the whitish discal spots larger, those just beyond the cell. An additional white spot is present near the apex, and a series of outer-submarginal dots increasing towards the lower angle, an inner-submarginal series confined to the apex. In the female three or four of the outer-submarginal dots are sometimes visible above. In the male the sex-mark is somewhat less prominent than in *D. genutia*.

Sambawa, 1000—2500 feet, scarce. When on the wing it somewhat resembles an undescribed *Euploea* found there.

Nearest *Danais ismare* from the Moluccas, but having the markings of the upperside reddish instead of white, and much reduced in size and number.

11. DANAIS (NASUMA) TAIMAND, n. sp.

Female. It obviously differs from the preceding species in the presence of a broad quadrate white discal band on the forewing, between the first subcostal and the upper median vein, in six pieces separated only by slender dark veins; below this there is one or sometimes two very small white spots. The submarginal spots are all obsolescent except one or two at the apex of the hindwing. The basal marks on the forewing are very indistinct, and merely paler not reddish, that in the cell absent. The discal marks on the hindwing are wholly undefined, resembling a large pale area, broken by dark veins. *Below*, there are generally one or two dots beyond the end of the cell, and also a few minute ones at the apex, one between the lower subcostal veins

more distinct, more or less bifid. The hindwing has the leaden-whitish markings rather broader than in *D. haruhasa*, but those in the median spaces are much shorter, leaving the dark outer border very wide there. The submarginal dots are in one specimen wholly absent, in the other partly present, but very small.

The male is unknown. I took one female at Lateng (1000 feet) in Taimanu, Sumba, and another at Mandas, Sumba. I fear that both are now lost. When flying it somewhat resembles *Euploea lewa*, and no doubt its mimicry of that species accounts for the presence of the broad white band, absent in *D. haruhasa* and *D. ismare*.

The subgenus *Nasuma*, as far as known, inhabits only the Moluccas, Sumba, and Sambawa, but no doubt a species will be found in Timor, while none is known from Java. It is distinguished by its elongate wings, and, at least in the two species described here, the flight is swifter than in *Danais genutia*, etc. It seems likely that these insects have lost some of the protective qualities of their allies, and have acquired a swifter flight and become mimics of other butterflies, the Moluccan form resembling a *Radena*, while the Sumba and Sambawa species look like *Euploea*s when flying.

12. DANAIS (TIRUMALA) LIMNIACE, Cram.

13. DANAIS (TIRUMALA) MELISSA, Cram.

Following Herr Semper's instructions, I easily separated these two species, which are extremely alike in general appearance. The *melissa*-form somewhat resembled the figure of *D. australis*, Hombrom and Jacquinet. I also recorded a form of *D. gautama* in Sumba, but no specimens have turned up. *D. limniace* and *melissa* are both common in Sumba and Sambawa.

14. DANAIS (CHITTIRA) ORIENTIS, n. sp. Pl. II, fig. 5.

Near *D. nilgiriensis*. Cell-mark of forewing with all three rays distinct, though slender, in the females, the upper two obsolescent in the male, the interno-median marks broadly divided, the mark at the base of the lower median space wanting (present in *nilgiriensis*), that in the upper median space small and diffused (large and conspicuous in allied species), the streak above the radial vein much longer than the one above it (as in *D. larissa*), five or six submarginal dots. Hindwing with the cell-spot broad in the middle (narrow in *nilgiriensis*), divided longitudinally by a slender dark line (absent in *D. larissa* and *lazonensis*), a line of six outer-discal spots in the male, nine or ten in the female, the submarginal line of spots incomplete. *Below*, both lines of spots are complete.

Pada Dalung, Central Sumba: a very dull-coloured species. It appears to belong to Mr. Moore's newly-described genus *Badacara*, along with *B. nilgiriensis*.

A single male from Sambawa agrees in the main with those from Sumba. But the whitish markings are better defined and more transparent, the outer submarginal spots of the forewing extend on the underside to the apex, the elongate discal streak between the radial veins is shorter, and all the discal and submarginal marks of the hindwing are somewhat larger and more distinct. The specimen is not now in my possession, and I am unable to compare it with *D. larissa*.

15. *RADENA OBERTHÜRII*, n. sp. Pl. II. fig. 6.

Male, above dark brown, the markings yellowish, somewhat translucent; a pale streak along the costal vein, the tip clavate and more distinct; the basal cell-streak bifid, its upper ray very slender, short; terminal cell-spot narrow, obliquely transverse; interno-median space with two strong white bands scarcely convergent, a broad dark space between them; a large, elongate spot in the lower median space; a broad obliquely-transverse discal band of four large white spots separated by veins, one on the costa somewhat apart from the others, the fourth largest, ovate, the third incised outwardly; another small spot beyond these on the costa, and six small inner-submarginal spots, the upper three in a line across the apex, the others small, transverse, between the upper median and internal veins; no outer-submarginal spots are present. Hindwing paler brown than the forewing, the cell all white, a large spot in each space beyond it, making, besides the long submedian streaks, four in all, the second (above the upper median vein) incised outwardly, and projecting beyond the others, the first and fourth elongate, the third small, triangular; a line of about nine inner-submarginal dots, placed rather irregularly. *Below*, similar, five or six obscure outer-submarginal dots on the hindwing only. The tufts are long, as in *R. juvena*.

Nearest *Radena purpurata*, Butler, from New Guinea, from which it obviously differs in the two interno-median streaks, and the broad oblique discal band on the forewing.

I name this fine species in honour of the distinguished entomologist, M. Charles Oberthür, of Rennes. It is a dominant species in Sumba, occurring both on the coast and inland, and seems to be the most western representative of the Papuan group to which it belongs.

16. *RADENA KAMBERA*, n. sp. Pl. II. fig. 7.

Allied to *R. juvena*, from Java. The wings are shorter, and

most of the white markings larger. The basal cell-streak is short and dusky, the outer one large, upright, the upper part projecting like the lower, a slender streak above it. The two series of subapical streaks of *juventa* are in *kambera* united into three very long white strigæ, that between the radial veins being nearly half an inch in length; all are incised outwardly; the inner-submarginal spots are large. Hindwing with the dark streak in the cell-spot continued to the end of the cell, slightly forked in the middle. *Below*, the light markings are not yellowish and greenish as in *R. juventa*, but pure white with a slight lilac gloss.

A very distinct species, not very common in Sumba.

In Sambawa there are two species of *Radena*, both I think distinct local forms. One, which appears to be the representative of *R. vulgaris*, is common everywhere; the other is very close to the Javanese *R. juventa*, and is confined to the higher country, though I have taken it as low as 1500 feet. I have now no specimens of either species, and am unable to compare them with their allies.

Family SATYRIDÆ.

17. LETHE EUROPA, Fab.

A female, Sumba, 2000 feet; another, Sambawa, 4000 feet, both resembling Java specimens.

18. MYCALESIS (ORSOTRIENA) MEDUS, Fab.

Sumba, Sambawa, common in meadows.

19. MYCALESIS (CALYSISME) PERSEUS, Fab.

The wet-season, ocellate brood prevailed in Sumba till the middle of March, when the non-ocellate form (*blasius*) took their place. In Sambawa, the latter brood had already begun to appear in the middle of April, but a long succession of heavy rains exterminated them, and the ocellate form reappeared and continued in exclusive possession till the latter part of May.

20. MYCALESIS (JATANA) WAYEWA, n. sp.

Female, above dark brown, a small ocellus on the forewing between the lower median branches. Hindwing outwardly whitish, gradually darkening inwardly, with two submarginal dark lines, a small ocellus between the lower median branches. *Below*, dark brown, slightly rufous, not perceptibly striate; a median transverse whitish line, angled at the upper median and above the submedian vein; beyond this the wing is much paler, with a rather small ocellus above the upper radial

vein, a large one above and a small one below the lower median vein; a marginal line and two wavy, dark, submarginal ones. Hindwing with the outer half whitish, the dark area very sharply outlined, projecting outwardly above the upper median vein; seven ocelli nearly in line, the first, fourth, and fifth large, subequal, the second and seventh minute; beyond this are two wavy submarginal and marginal dark lines.

The male is darker and more uniform, the ocelli less marked, with a golden-brown sex-mark (as in *M. mineus*) on the submedian vein of the forewing, and a large subcostal ochreous tuft, the subcostal vein and its upper branch, slightly swollen around its bifurcation. The prehensors and sex-marks agree with those of *Calysime* and so does its venation, except in the point mentioned.

This species occurs both in Sumba and in Sambawa, confined in both to the higher and damper regions. It is apparently the local representative of *Mycalesis mynois*, Hewitson from Timor, but lacks the conspicuous white band of that species. I also suspect it to be the local representative of *M. mineus*, but without a more elaborate study of the prehensors I cannot well prove it. *Mycalesis mynois* is the type of Mr. Moore's genus *Jatana*, the *raison d'être* of which I have not been able to discover.

21. *YPHTHIMA ASTEROPE*, Klug.

Sumba only, found in the driest plains. It is worth remarking that this species has prehensors precisely like those of *Y. pandocus*, the size and habits of which are so different, while the markings are nearly identical.

22. *YPHTHIMA LEUCE*, n. sp.

A local form of *Y. philomela*. The forewing has a large, conspicuous, whitish sex-mark, and an ocellus as in *philomela*; the hindwing has the cilia white, and the outer and abdominal region grey, with two large blue-pupilled ocelli and two minute anal ones. *Below*; the striæ are very irregular; the forewing has a whitish discal band partly inclosing the large ocellus, and extending to the hind-margin; there is a submarginal dark band, the apex and costa are dark. Hindwing chiefly white, the disc being free from striæ over a considerable area, and elsewhere they are very delicate and irregular, forming an obscure transverse fascia, crossing the cell near its end, and a continuous, slender submarginal line. The six ocelli are in pairs, as in *Y. philomela* or *sepyra*, all black, pupilled with blue, and with large ochreous irides.

This species is found in Sumba and Sambawa. It may also be allied to *Y. aphnius* (Timor) of which only the dry-season form is known.

In that case it bears much the same relationship to *aphnius* as *Mycalis wayewa* does to the Timorese *M. myzoides*, the orange band of the hindwing of *T. aphnius* being absent.

Mr. de Nicéville has shown me that the true *Ypthima philomela* of Johanssen has no sex-mark. Since that is the case, the Indian form will stand, I suppose, as *Y. baldus*.

23. MELANITIS CONSTANTIA, Cram.

Sumba, Sambawa, not uncommon on the dry coast. A Sumba female before me has the ochreous band narrow and irregular, almost attaining the costa and the lower angle, its outer border undefined, with three obscure ocelli (the first chiefly white, the last nearly all dark), the outer border with three indentations, and projecting acutely inwardly along the lower median vein. Hindwing with two ocelli, the apical border narrowly ochreous. *Below*, the ocelli are rather small, the aspect that of *M. leda*, the wet-season form.

24. MELANITIS LEDA, Linnæus.

The dry-season brood (*ismene*) appeared in Sumba in the middle of March, in Sambawa in the middle of May; till that date the ocellate brood held the field.

Family ELYMNIADÆ.

ELYMNIA UNDPULARIS, Drury.

Sambawa, low country, not differing from Java specimens. No *Elymnias* is known from Sumba or Timor.

Family MORPHIDÆ.

A *Discophora (timora)*, Wallace has been found in Timor, but I saw none of the family either in Sumba or Sambawa.

Family BYBLIADÆ.

25. ERGOLIS ARIADNE, Linn.

Sumba, Sambawa, not differing from the Indian form, hardly so small and dark as the Javanese. I believe *E. merione* also occurs in Sambawa.

Family APATURIDÆ.

26. CYNTHIA DEIONE, Erich.

A single male, Sumba, interior. Common in Sambawa, where the females vary to a remarkable extent, some being as red as the male,

while others are dark green insects like *parthenos*. Intermediate forms are common.

27. *CETHOSIA PENTHESILEA*, Cram.

Sumba and Sambawa, common, even on the coast. A single female taken in the interior of Sumba was very large and richly coloured, resembling the female of the Javanese *C. hypsea*, Doub., but I am not sure that it was distinct from *C. penthesilea*.

CETHOSIA TAMBORA, n. sp

Black, the usual wavy submarginal lines absent on both wings. Forewing with an irregular ochreous subapical band of six marks, the first two slender, minute, the third elongate-quadrate, the fourth very small, triangular, the fifth long and narrow, the sixth large, triangular, paler; a dull reddish area on the hind margin and basally in the internommedian space, two or three reddish touches basally in the cell. Hindwing, basal half red, a small dark spot discally between the costal and the subcostal veins, and one between the subcostal branches. *Below*, all blue-black and ochreous-white, except a reddish area on the hind margin of the forewing.

The description, which is apparently that of a female, is a poor one, but I believe several specimens of both sexes are in Mr. Neumoen's possession. It is very unlike any known species.

Sambawa, mountains, scarce. I have named it after the great Sambawan volcano, celebrated for its eruption in 1815.

I have dubiously recorded a similar species in Sumba, not taken, the underside more variegated.

Two very beautiful species, *C. lamarckii* and *leschenaultii*, were taken by Mr. Wallace in Timor, but neither seem to extend further westwards.

28. *CUPHA ERYMANTHIS*, Drury.

Sumba, Sambawa, normal. No *Cirrochroa* was seen in either island, nor has any been recorded from Timor.

29. *AELLA PHALANTA*, Drury.

Sumba, Sambawa. Sumba specimens are richly marked with purple below.

30. *AELLA SINHA*, Kollar.

Sumba, Sambawa, normal.

31. *CYRESTIS NAIS*, Wallace.

Two tattered specimens from Pada Dalung in Sumba seem to be

almost precisely intermediate between *C. nivea* and *C. thyodamas*. To indicate their affinities would require a long description. They seem near Mr. Wallace's *C. nais* from Timor, but without better specimens I cannot be sure of their position.

SYMBRENTHIA HIPPOCLUS,* Cram.

Sambawa, none taken.

32. PYRAMEIS CARDUI, Linn.

Dry meadows, Sumba.

33. JUNONIA ATLITES, Joh.

Sumba, Sambawa.

34. JUNONIA ASTERIE, Linn. var., SUMBAE.

Above, the subapical ocellus is indistinct, merged in the black band from the costa; the lower ocellus is large and set in a black patch. On the hindwing the lower ocellus is much larger than in Indian specimens of *asterie*, and is marked like the upper one. *Below*, the forewing has only two ocelli, the upper pair on the hindwing are more perfectly merged into one, the black transverse lines are replaced by diffused pale reddish ones; the hindwing has three pale bands across it.

Sumba, Sambawa, common. It is merely an extreme form of the Java variety (*J. javana*, Felder), which connects it with the typical Indian one, differing chiefly on the underside.

The non-ocellate form, *J. almana*, probably conspecific with *asterie*, was not taken.

35. JUNONIA VELLIDA, Fab.

The upperside agrees with Godart's description. The underside is rather brilliantly marked with black or fuscous wavy lines on a pale grey ground, a reddish submarginal band, the hindwing with five ocelli, of which only the second and fifth are distinct, pupilled with bluish.

This pretty little species occurred only on the desert plains of Sumba, and seemed to be rare.

36. JUNONIA AONIS, Linn.

The markings of the forewing are rather fuscous than fulvous except the ocelli, of which only two are distinct, the lower small and attached to the upper. On the hindwing the second of the five ocelli

* In Eastern Java I found that the female of this butterfly was dimorphic, one form having the yellow spots replaced by white ones, so that it resembled a white *Neptis* instead of a yellow one. No intermediate forms were seen.

is larger than the others. Otherwise the specimens agree with Godart's description. In the female there are distinct, reddish-bordered ocelli on the forewing, and the sordid spots are larger. The underside is sometimes reddish, as is often the case with the female of *J. lemonias*. There seem to be two perfectly distinct seasonal forms, that of the dry-season resembling a dead leaf below.

Sumba, Sambawa, those from the latter island not examined. The species is certainly very close to the Javanese *J. erigone*.

37. JUNONIA TIMORENSIS, Wallace.

In the male there is no distinctly rufous area except a submarginal band on both wings, evanescent apically on the forewing. In some females nearly the whole upper surface is more or less rufous, outwardly brighter, a dark area over the apex and disc of the forewing, the subapical spots united, a small distinct ocellus (obscure in the male) between the lower median branches. *Below*, the male is dark with the ocelli distinct, while the above-mentioned females are glossed with silvery-grey scales over both wings, obscuring the subapical band and the ocelli, the median transverse line distinct, angled at the upper median vein. These females evidently belong to the dry-season brood, just appearing at the time I left Sumba, and resembling the other form much as *J. almana* resembles *asteria*. I have not observed any male of this brood.

Sumba. This very distinct species, as Mr. Wallace justly calls it, was previously known only from Timor,

38. JUNONIA ORITHYIA, Linn.

Sumba, Sambawa.

39. PRECIS IDA, Cram.

Sumba, Sambawa.

40. PRECIS IPHITA, Cram.

Sumba, Sambawa.

41. YOMA SABINA, Cram.

Sumba, Sambawa, low country.

I see that Doubleday, followed by M. Oberthür and Dr. Semper, places this species in the very heterogeneous African genus *Salamis*, the type of which somewhat resembles a *Doleschallia*. The only species of *Salamis* at all like *Yoma* are *S. anacardi* and *anteva*, forming Wallengren's genus *Protogoniomorpha*, a name which ought to be barred on account of its enormous length. But in these species the cell of the

hindwing terminates opposite the second forking of the median vein, in Yoma at some distance before it. The relations of homologous genera in different zoological regions are as yet very little known, and will no doubt greatly exercise the minds of naturalists in the next century. But for the present I think my genus *Yoma* may be allowed to stand.

42. *HYPOLIMNAS MISIPPUS*, Linn.

Sumba, Sambawa. The female mimics *Danais chrysippus* as usual.

43. *HYPOLIMNAS BOLINA*, Linn.

Sumba, Sambawa. The female sometimes has a red area near the hind margin of the forewing and over the disc of the hindwing.

44. *HYPOLIMNAS SAUNDERSII*, Wallace, (?).

Several broken specimens of what appeared to be a dwarf form of *H. pandarus* were taken in Sumba, but none have survived.

HYPOLIMNAS ANOMALA, Wallace.

One male, taken in the mountains of Sambawa. I am unable to say whether it was identical with Javanese specimens or not.

45. *DOLESCHALLIA* sp.

At least one species occurs in the dry coast region of Sumba, but no specimen was taken.

46. *HELCYRA CHIONIPPE*, Felder.

Several specimens seen at Pada Dalung in Sumba, but none taken.

47. *CHARAXES ATHAMAS*, Drury.

Sumba, Sambawa.

48. *CHARAXES* sp.

A very large *Charaxes* apparently of the *eudamippus* group was several times seen in the mountains of Sumba, and again in those of Sambawa. Unlike *C. eudamippus*, which is a ground butterfly, it always alighted high up on trees, so that I could never catch it. Another species, something like *C. pyrrhus*, was once seen in Sumba.

Family NYMPHALIDÆ.

49. *PHÆDYMA COLUMELLA*, Cr.

Sumba, Sambawa. The upper band of the hindwing above is much broader than in Indian specimens.

50. NEPTIS HORDONIA, Stoll.

Sumba, Sambawa. The yellow markings above are all much smaller than in Indian specimens.

51. NEPTIS NANDINA, var. SUMBA.

This form greatly resembles the Javanese *N. leucothoë*, a near ally of the Indian *N. varmona*, but the triangular white spot beyond the cell is narrow and elongate, sometimes extending on the underside two-thirds towards the margin. The subapical white band is somewhat less massive, and the four large discal spots are separated by veins, the first elongate, as large as any of the others. The upper band of the hindwing is narrower, the submarginal lines remoter from the margin. The general colour of the underside is dark red, the white bands very slightly outlined with dark. Though the markings resemble those of *N. leucothoë*, the species is probably more allied to *N. nandina*. From this it may easily be distinguished by the brilliant white markings of the upperside. The discal spots are large, that between the upper two median veins large and elongate, the cell-streak is well separated from the triangular spot beyond the cell. The upper band of the hindwing narrower, especially apically.

Sumba; a somewhat different form occurs in Sambawa, which I am now unable to examine. I did not observe any representative of the *varmona* group in Sumba.

52. ATHYMA PERIUS, Linn.

Sumba, Sambawa. As in India it generally occurs in open meadows, unlike all its allies.

53. ATHYMA KARITA, n. sp.

Male, above deep brown, variegated with darker areas, forewing with an outer-submarginal pale line, and an inner-submarginal series of slender obsolescent greenish streaks. An oblique subapical band of three greenish-yellow spots, the second largest, adjacent to the first, the third as large as the first, round, separate; cell unmarked; a broad discal macular band from the hind-margin to the second median vein, with a small triangular spot above it, that between the two lower median veins rounded-quadrate, as large as the one below it, separated from it only by a vein, and projecting beyond it. Hindwing with two greenish-yellow bands, the upper unbroken, very wide on the costa, extending to the submedian vein, where it tapers to a point; the lower band composed of six transverse spots cut by dark veins, their lower edges incised, the outer ones slender, lunular. Below the markings are similar, but whiter and more united, placed on a dull brown ground, a

darker chocolate area discally on the forewing, a row of obscure darker submarginal spots on the forewing, and discally on the hindwing; the abdominal margin of the hindwing is broadly suffused with bluish.

Sumba, a single male taken by the river Waibaku near Pada Dalung, at 1,500 feet. The species seems intermediate between *A. venilia* and *A. amhara*.

ATHYMA NEPTE, Cram.

One male taken near Ndindi, Sambawa, at 3000 feet elevation. It is one of the apparently numerous Indo-Malayan species inhabiting the mountainous interior of this island.

54. LIMENITIS* PROCRIS, Cram.

Sumba, Sambawa.

* I append the description of a very rare species from Perak, Malay Peninsula, allied to *L. daraxa*.

LIMENITIS AGNEYA, n. sp.

Male. Above very deep fuscous, a pea-green band across both wings. Basal forewing somewhat paler, with obliquely transverse darker markings in and below the cell: a submarginal pale line, an outer-discal row of six dark spots set in square paler spaces. The green band extends from the hind margin to the upper median vein in four spots separated by dark veins only, the upper spot a little out of line and smaller. Beyond these a line of three smaller spots runs to meet them obliquely, placed at right angles with the costa, extending from the upper median to the subcostal, the upper smaller than the others and whitish, placed above the inner half of the middle and largest one. More than halfway between these and the apex are two other spots also placed at right angles with the costa, and between the third and fifth subcostal veins, the lower small, greenish, the upper very small, whitish. On the hindwing the green band extends from the costa, where it is widest, to the submedian vein, tapering to a point. Its inner margin is convex, its outer straight; it is whitish at both ends and cut by slender black veins. The submarginal pale line and the outer-discal line of spots are much as on the forewing; cilia white, dark at the ends of the veins. Below light chestnut-brown, the band as above, but paler green and not cut by dark veins, the basal marks on both wings outlined with lilac, the outer-discal line of spots set in lilac areas, the submarginal line lilac, some darker red discal markings beyond the green band.

Obviously differs from *Limenitis daraxa* in the bifid green band of the forewing; agrees with it in the closed cells of both wings, which make the positions of the species in this genus rather dubious. My single male of *agneya* was taken on Larat Hill, Perak, at about two thousand feet elevation. *L. daraxa*, not apparently differing in any respect from the Himalayan form, was not uncommon at the summit of the same hill, nearly three thousand feet higher.

Prehensors, seen from the side. In *agneya* the uncus is long, sinuous, hooked at the tip, without branches, the clasp simple, straight, tapering gradually to the

LIMENITIS HOLLANDI, n. sp.

Male, above, black; cilia alternately black and white, a double undulating submarginal pale line, touched with whitish apically on the forewing; two united white spots placed obliquely subapically above the radial veins, with or without a smaller one below them. A broad common white band, very slenderly cut by dark veins, edged outwardly with purple, across both wings, extending on the forewing nearly to the upper median vein, the upper (fourth) piece small, the inner edge of the band straight, the outer irregular. Hindwing with the white band extending to the submedian vein, where it is tapering, both its edges nearly straight. *Below* chestnut-red of two different shades, the basal two-thirds of the cell mostly white, with a crooked dark mark, an irregular white spot at the end of the cell, its lower side tapering; the white band extends nearly to the lower radial vein; two submarginal pale lines, the inner mostly bluish-white with three white subapical spots; a chestnut longitudinal streak in the internommedian space. Hindwing with the two submarginal lines regular, grey, undulating, base chiefly white with transverse streaks of chestnut, namely, two in the cell, two between the costal and the first subcostal vein, one along the precostal vein; and a long one from the costa tapering to the submedian vein; the white band broad. It is rather a small species.

Several males taken in the Do Donggo country, Sambawa, all above 2000 feet. The species is nearest *L. lysanias* from the Celebes, but is without the rufous bands which that species has on the upperside.

I name the butterfly in honour of the Rev. W. J. Holland, of Pittsburgh, U. S. A., well-known as a lepidopterist.

55. SYMPHEDRA EGLE, n. sp.

Male, above black, with some obscure ochreous-greenish spots near the costa, and three in the cell (two at the end), two in the internommedian space, one basal, the other further out, geminate; a row of five white subapical spots in a semicircle above the middle median vein, the second largest, then the first, the third diffused; also a macular submarginal band of dull bluish-whitish spots (the lower two sagittate) extending from the hind margin nearly to the apex, separated by dull

tip. In *L. populi* the tip is abruptly bent downwards. In *daraxa* the tip is blunter than in *ageaya*, and there is a long process arising from its upper edge, longitudinal and slightly ascending, set with denticles on both sides. The uncus is also shorter. The lower uncus, absent in most butterflies, is well developed in this genus, strongly hooked at the tip, its point opposed to that of the true or upper uncus, which can be brought into contact with it by muscular action.

bluish-green spaces. Hindwing with a broad bluish-green band from the abdominal margin (where it is whitish) to the costal vein (where it is blue), consisting of spots twice as long as wide, separated by black veins, their inner border diffused, the outer pointed, enclosing a large oval or cordate black spot, slenderly surrounded with blue. *Below*, forewing nearly black, slightly olivaceous apically, the markings white, more or less tinged with greenish or ochreous, arranged in two series, a submarginal one of five or six spots, and a discal one of seven, of which the third is evanescent, the fifth and sixth obliquely elongate, the seventh geminate and purplish; there are also spots on the inner disc, between the median veins and above each radial; cell crossed by two bluish-white bands, one in the middle, one at the end, with several additional spots at the base. Hindwing purplish-brown with a slight bronzy lustre; a submarginal band of dark spots in paler spaces; an irregular discal series of pale spots, three spots in the cell. Eyes dark, proboscis scarlet.

The colour of the underside, the narrow and dull blue band, and the numerous spots on the upperside distinguish it from *S. dirtea*.

A male, Pada Dalung, a female, Mandas, both in Sumba. I have no description of the female.

An *Euthalia*, dark like *E. garuda*, seems also to inhabit Sumba, but none were taken.

Family LIBYTHEIDÆ.

56. LIBYTHEA GEOFFROYI, Godart.

Two males, interior of Sumba. The blue of the forewing almost obliterates the subapical spots, which are barely traceable above; the luteous band of the hindwing is very obscure.

57. LIBYTHEA NARINA, Godart.

One female, interior of Sambawa, another seen in Sumba, not taken. The Sambawan specimen agreed well with Moluccan examples; The Philippine form has the white band of the hindwing reduced, while the Assam variety (*L. rohini*, Marshall), has it enlarged.

Family NEMEOBIADÆ.

ZEMEROS PHEGYAS, Cram.

Sambawa, scarce. No species of this family was observed in Sumba.

Family LYCÆNIDÆ.

Subfamily THECLINÆ.

58. ARHOPALA ARAXES, Feld.

Sumba, coast. A local Celebesian form of *A. amantes*, but apparently distinct.

FLOS APIDANUS, Hew., var.

Sambawa, 3000 feet.

SURENDRA QUERCETORUM, Moore.

Sambawa, coast.

Subfamily APHNEINÆ.

IRAOTA TIMOLEON, Stoll.

Sambawa. I am almost inclined to think this species a mimic of the white species of *Neptis*; the resemblance is sometimes quite striking, and in any rate the *varmona* group of *Neptis* is partially protected.

59. CURETIS MALAYICA, Feld., var. KIRITANA.

Male black, the red area scarcely extending above the middle median vein, the hind margin dark. Hindwing with the end of the cell and the disc from the subcostal vein to about the lower median, red or reddish, the abdominal and costal margins very widely, the outer margin more narrowly black.

Sumba, Sambawa, scarce. This is, I think, the darkest *Curetis* known, Dr. Felder's *C. obscura*, described as a male, being really a female.

60. HYPOLYCENA SIPYLUS, Feld.

Sumba, Sambawa. This is near Dr. Felder's *H. thecloides*, but the lower part of the hindwing above is bluish, not ochreous.

61. LOXURA ATYMNUS, Linn.

Sumba, Sambawa. Two or three more species of the *Aphneinae* were taken in Sumba, but the specimens are now lost.

Subfamily DEUDORIGINÆ.

62. RAPALA IARBAS, Fab.

Sumba, Sambawa, mountains.

RAPALA ORSEIS, Hew.

Sambawa, 1,500 ft., a male and a female.

63. *RAPALA VARUNA*, Horsf.

Sumba, Sambawa, mountains. I also observed a species of *Sinthusa** in Sambawa, but no specimens were taken.

Subfamily *LYCENINÆ*.

CYANIRIS AKASA, Horsf.

Sambawa, 4-5000 feet, rare.

64. *CYANIRIS PUSPA*, Horsf.

Sumba, Sambawa, mountains. The white area is larger than in Indian specimens, extending over three or four spaces and into the cell. The female has the white areas very large and not marked with blue.

65. *ZIZERA PYGMEA*, Snellen.

Sumba, Sambawa.

66. *ZIZERA LYSIZONE*, Snellen.

Sumba, Sambawa. Another *Zizera* occurred in Sumba.

* I append the description of a rare Javanese species of this genus.

SINTHUSA ASPRA, n. sp.

Male, above, forewing with the costa and the outer margin narrowly black, the base as far as the end of the cell, and more especially the basal half of the intermedian space to the hind margin, light violet-blue; a diffused submarginal macular band of the same colour; the disc and the outer half of the hind-margin black, deep blue in some lights, with a few scattered light-blue scales. Hindwing violet-blue, (much richer than on the forewing) from the lower subcostal to the submedian vein; above the lower subcostal vein a line of pale blue scales; the abdominal border widely silvery-whitish. Below, white, the spots chiefly very small, black, not annular as in all the allied species: a broad dark fascia across the end of the cell of the forewing; six discal spots, the upper three small, in an oblique line, the lower three larger, in a transverse line nearer the base; a slender marginal dark line, cilia dark. Hindwing with the cell-fascia double; eight discal spots, the fifth evanescent and nearer the base, the eighth elongate and conspicuous; a large black spot in the lower median space; in the next a blue area adjoining the black lobe; beyond this a short black and blue marginal line; tail chiefly black. The prehensors resemble those of *Deudorix*.

The hindwing is angled at the end of the middle median vein; there is a short tail and a very small but distinct lobe. The venation and sex-marks are as in other species of *Sinthusa*. The species has no near allies.

Rare on Arjuno, Eastern Java, taken at 5000 feet in a flock of *Cyaniris akasa* from which it was indistinguishable when settled. The genus is usually mimetic. *S. nasaka*, Moore, strongly resembles *Hypolycaena erylus*, and I have several times mistaken *S. virgo*, Elwes, for a *Cyaniris*.

I have given Mr. de Nicéville the type of this species. The specific name means white in modern Greek.

67. *Polyommatus beticus*, Linn.
Sumba, Sambawa, high country.
68. *Chilades trochilus*, Freyer (*pulli*).
Sumba, Sambawa.
69. *Everes parrhasius*, Fab.
Sumba, Sambawa.
70. *Tarucus theophrastus*, Fab.
Sumba.
71. *Tarucus plinius*, Fab.
Sumba, Sambawa.
72. *Castalius ethion*, Doub.
Sumba, Sambawa.
73. *Castalius rosimon*, Fab.
Sumba, Sambawa.
74. *Castalius roxus*, Godt.
Sumba, Sambawa.
75. *Catochrysops cneiis*, Fab.
Sumba.
76. *Catochrysops pandava*, Fab.
Sumba.
77. *Catochrysops strabo*, Fab.
Sumba, Sambawa.
78. *Nacaduea gaura*, n. sp. Pl. II, fig. 8, *magnified two diameters.*
Male, above, pale violet, a narrow black border widening apically, the veins brown. Hindwing with a row of six black spots in whitish rings, a marginal dark line, cilia chiefly white. *Below* pale brown, heavily marked with white. Forewing with a serrate, catenulated line of dark spots, surmounted by a row of lunules outwardly whitish, inwardly dark; three transverse quadrate dark bands, paler in the middle, and bordered with white, namely, two in the cell, the basal one continued into the interno-median space, and a broader one across the

wing discally to the submedian vein, slightly dislocated inwardly at the upper median vein, below which it is narrow and broken. Hindwing with a row of serrate ocelli as on the forewing, a very large black ocellus in the lower median space, slenderly edged with dark orange, and then a darker ring; a black anal spot; the three transverse bands much broken, the discal one forming two dark quadrate areas, the upper from the costa to the lower subcostal vein, the lower lying further out, extending thence to the upper median vein; the disc is chiefly white.

This curious species* is something like Dr. Felder's *Lycæna palmyra* from Amboina. The venation places it in *Nacaduba*, but it has little resemblance to the usual forms of the genus, and looks more like a *Catochrysops*.

Sumba, rare.

79. *NACADUBA LAURA*, n. sp, Pl. II, fig. 9, ♀, *magnified two diameters.*

Female, above, forewing about three-fifths dark brown; a large white area, sprinkled, especially at the base, with blue scales, extending from the cell to the hind-margin. Hindwing with the cell and the two spaces beyond it discally white, the rest of the disc paler brown; two submarginal rows of black spots, the outer round, the inner lunular. *Below*, forewing with a large white area occupying the end of the cell and the disc to the hind-margin; a basal transverse dark band across the cell and the interno-median space; the discal transverse bands are obliterated, except costally, one being represented by two brown lines extending from the hind margin into the white area as far as the middle median vein. Hindwing with the white area much smaller and duller-coloured, the transverse bands confused. Both wings have two submarginal rows of conspicuous dark spots, the inner large, black, semicircular on the forewing, lanceolate on the hindwing, the outer paler, transverse. The hindwing has two slender ocelli with slender irides of orange, touched with metallic, the outer one with the black area very large.

* An allied species, of which *N. gaura* is presumably a local form, occurs in south-eastern Borneo and Java. I also took a male in Engano (unluckily omitted in my list), and I think I found it in the Nicobars. A Bornean specimen before me has the black marginal band above very narrow and equal, the black spots on the hindwing obsolete, except that in the lower median space, which is large. *Below* there are no distinct white areas, but all the markings are conspicuously edged with white. The discal band is bent outwardly and very irregular as far as the middle median vein, below which it is broad, straight and quadrate, the subanal ocellus is broadly bordered with orange.

I propose the name of *N. pseustis* for this species.

I took several females in Sumba, both on the coast and in the interior, and also a single male probably of this species. It was violet blue above, the margin very slenderly dark. Below the white areas were nearly obsolete on the forewing, and on the hindwing reduced a white bar in the band across the end of the cell, and a border one on the disc beyond the cell between the lower subcostal and upper median veins. Expanse over an inch and a quarter.

The species is near *N. perusia*, Felder, from Amboina, and *N. atrata*, Horsfield, from Java, but the extensive white areas of the female easily distinguish it.

80. *NACADUBA HERMUS*, Feld. (*viola*, Moore).

Sumba, Sambawa.

81. *NACADUBA ARDATES*, Moore.

Sumba, Sambawa.

82. *NACADUBA MACROPHALMA*, Feld.

Sumba. The female is marked almost as in that curious little species, *N. kerriana*, Distant, which I have taken at 4000 feet elevation in Karenni east of Burma.

83. *NACADUBA DANA*, de Nicéville.

Sumba, Sambawa. This is probably Mr. Druce's *N. almora*, but his figure and description are so bad that certainty is impossible.

84. *LAMPIDES BOCHUS*, Cram. (*Jamides bochus*.)

Sumba, Sambawa, rare. The only Sumba specimen I have examined has the blue area above very large, and not at all brilliant. It may be a distinct local form of this species, intermediate between *L. bochus*, Cram. and *L. astraptus*, Feld.

A *Lampides* which occurs in Borneo, Sumatra and the Malay Peninsula but has apparently escaped the notice of naturalists, has the underside like that of *L. bochus*, while the upperside has only a slender marginal dark line and is of a rich blue, darker than that of *L. elpis*. I have given Mr. de Nicéville a Bornean example for description.

85. *LAMPIDES ANOPS*, n. sp. Pl. II, fig. 10.

Male, above, rich azure-blue, virescent in some lights, darkening outwardly, translucent, a slender marginal dark line somewhat broader apically on the forewing, the cilia of the hindwing white. Below, ground-colour basally grey, darkening outwardly, but without the slightest

rufous tinge. The markings consist of slightly darker transverse bands, bordered by straight white lines, which are broad and very conspicuous. Forewing with one of these across the cell, with no markings above it costally, one discal from the costa to the middle median, then dislocated inwardly and continued obliquely to the lower median, below which the wing is white with a single dark streak in it. Hindwing with the bands confused and broken. Both wings have the cilia whitish, a marginal dark line, a catenulated line of dark streaks in a white ground, and behind this a line of very conspicuous black lunules, large and lanceolate on the hindwing; behind these are white lunules which extend far into the disc. There is no trace of ocelli, or of metallic scales.

The female is also blue, and has the outer two-thirds of the forewing black above.

Sumba, confined to the mountain-forests above 2000 feet. A beautiful and conspicuous species. I have not examined the prehensors, but the species is so unlike all others that its identification must be easy. In the figure the white markings of the underside have been made too narrow and inconspicuous.

86. *LAMPIDES MASU*, n. sp. Pl. II, fig. 11.

Male, above, bluish-white, whiter than *L. alianus*, a very slender marginal black line nearly obsolete apically; hindwing with this line more distinct; a broken, catenulated, submarginal dark fascia, double at the anal angle, obsolete apically, with a good-sized dark spot in the lower-median space. *Below* pale brown, the bands scarcely perceptibly darker, bordered by white lines, of which the basal pair on the hindwing are slender; on the forewing one band crosses the cell, one is beyond it from the costa to the middle median vein; these two are continued in common by another nearly to the hind-margin. On the hindwing the bands are irregular, extending further outwardly than in *L. anops* (in which the submarginal lunules greatly encroach on the disc), acutely angled in the interno-median space. Both wings have three conspicuous white submarginal lines enclosing two lines of spots, the outer linear, catenulated, slender, the inner large, black and conspicuous, transverse and wholly surrounded with white on the forewing, lanceolate and irregular on the hindwing. Hindwing with a large subanal ocellus with a narrow orange iris, surmounted by a black and a white lunule; a small similar anal ocellus; both are touched with metallic.

This species is very like the Amboina female figured by Cramer as *aratus*, and is probably a local variety of that species. The female of *L. masu* has the black border of the forewing broad and serrate apically, the inner cordate spots of the hindwing are large and black. The

male differs from *L. subditus*, Moore, in the whitish upperside, and in the pale underside, heavily marked with white, with the orange area smaller; it resembles it in the submarginal band of black spots. *L. subditus* is very close to *L. amphissa*, Felder, from Amboina.

87. LAMPIDES CELENO, Gr.
Sumba, Sambawa.

88. LAMPIDES ELPIS, Godt.
Sumba, Sambawa.

89. SPALGIS EPIUS, West.
Sumba, Sambawa, found on the acacias growing along the dry shore, the specimens normal.

90. MEGISEA MALAYA, Horsf.
Sumba, Sambawa.

91. NEOPITHECOPS ZALMORA, Butler.
Sumba, Sambawa.

Subfamily GERYDINÆ.

92. GERYDUS TEOS, n. sp.

A local form of *G. symethus*. The white area above is large, extending furthest below the middle median vein; there are no bluish scales; the upper median vein is swollen at its base. Hindwing all dark, not bluish-grey as in *symethus*. Below nearly uniform pale brown, a dark area crossing the end of the cell, surrounding the median vein and its branches discally; beyond this there is a whitish area from the middle median vein to the hind-margin; the transverse discal lunular band only extends down to the middle median vein. Hindwing with the discal lunules nearly joined, very distinctly marked, scarcely darker than the ground-colour, edged with paler. Both wings have a submarginal dark line edged inwardly with whitish, and containing a black dot in each space. The underside is wholly without the sordid irrorations found in *G. symethus*.

The female has a little less white on the forewing above, and a little more white below. The hindwing is acutely angled in the middle and is unmarked above.

Sumba, Sambawa. The Javan form (*G. pandu*, Horsf.) may also be distinct from *G. symethus*.

93. *GERYDUS BOISDUVALII*, Moore, var. *ACRAGAS*, nov.

Male, above, forewing with the base of the upper median vein swollen beyond the cell and placed in a small longitudinal pale space, no other markings above. Female with a narrow straight white band beyond the cell, extending obliquely to the lower median vein, broken by the dark middle median vein. *Below*, forewing with a white area over the disc to the hind margin, the transverse discal band formed of joined incomplete lunules in both sexes. The female has the hindwing slightly angled.

Sumba, Sambawa. I cannot compare this with the Javanese form, the male of which is still undescribed. But the shape of the band in the female of *G. acragas*, which resembles that of the male of *G. biggsii*, the absence of white or pale markings above in the male, and of sordid irrorations on the underside in both sexes, easily distinguish it from the Indian form of *G. boisduvalii*. It is obviously distinct from a form mentioned by Mr. Distant, who says "in an Amboinese species *G. boisduvalii*, Butler, the distinctive colouring of the anterior wing is reversed, the male having the largest white area to the anterior wings."

The male of *acragas* must certainly be very much like that of *G. irroratus*. Dr. Semper doubtfully identifies with that species a Philippine form having a large round median white patch in the female. I think this very unlikely to prove correct. *G. irroratus* is certainly very close to *G. boisduvalii*, but till the female of Mr. Druce's Siamese form is discovered, the question must be left open.

Family PIERIDÆ.

94. *NYCHITONA XIPHIA*, Fab.

Sumba, Sambawa, varying greatly in size and markings.

No *Elodina* was seen in either island, though *E. egnatia* is known from Timor.

95. *TERIAS HECABE*, Linn.

Sumba, Sambawa, several varieties.

96. *TERIAS SARI*, Horsf.

Sumba, Sambawa.

TERIAS DRONA, Horsf.

Sambawa, 2-4000 feet. *T. candida*, found by Mr Wallace in Timor, was not seen.

97. *TERIAS HARINA*, Horsf.

Sumba, Sambawa.

HUPHINA TEMENA, Hew.

Sambawa. A very beautiful species. .

98. HUPHINA JULIA, n. sp. Pl. II, fig. 12.

Male, above, creamy-white, a black marginal fascia, scarcely wider on the forewing than on the hindwing, its inner border diffused; the submarginal dark bands of the underside visible through the wings; all the veins of the forewing dark, as well as those of the hindwing outwardly. *Below*, forewing white, a submarginal diffused brown band, connected along the two upper median veins with a dark longitudinal band in the lower part of the cell; apex diffused ochreous-yellow, the yellow area just crossing the submarginal band costally, the outer margin narrowly brown. Hindwing bright yellow; a narrow dark marginal band, a broad, irregular, serrate dark submarginal band enclosing seven large orange-scarlet spots, obsoletely edged with yellow; the subcostal space nearly all scarlet, without any black border inwardly or outwardly. Expanse two and two-third inches.

Female, above, forewing with all the veins heavily outlined with dark, the spaces more or less white, a united subapical white band, and a row of submarginal spots. Hindwing pale yellow, the veins slightly darkened, a broad outer dark band enclosing white spots. *Below*, duller than the male, the radial and upper median veins of the hindwing marked with lines of blackish scales.

Nearest *H. lata*, Hewitson, from Timor, which has the forewing broadly and the hindwing slenderly black above, while below, the disc of the hindwing is uniform black, the submarginal dark band of the hindwing is obsolescent, and the scarlet costal stripe bordered on both sides with black. It is a much smaller butterfly than *H. julia*.

Sumba, interior.

This insect, the most beautiful Oriental species of *Pieris* known, has when flying none of the air of a protected butterfly. If it stood alone, I should certainly suppose it to be a mimic of some form of *Delias hyparete* yet undiscovered in the island. But both *H. lata* and *H. temena* require to be accounted for in the same way, and while it is possible that some Timorese *Delias* may resemble *H. lata*, I feel sure that *H. temena* can have no such original. It must then be assumed that this group is less pressed by its enemies in the Timorian Islands, and has therefore been able to acquire more brilliant colours than its allies.

Huphina lata, julia, temena and *tamar* form the nearest approach to a peculiar group of butterflies which these islands possess. But *H. tamar* is from Bali, beyond Wallace's Line, and no doubt extends into Eastern Java.

HUPHINA NAOMI, Wallace.
Sambawa.

99. HUPHINA EIRENE, n. sp.

Male. It differs from *H. naomi* in the colour of the hindwing below, and that of the subapical spots of the forewing, being lemon, instead of rich orange-ochreous. Above, the median vein and its upper branch are more widely marked with black, and the white of the cell and the hind margin of the forewing is clearer and purer.

In these points it agrees with *H. judith*, Fab. (Java), but differs in the narrow dark markings of the forewing above, the long white discal markings being slenderly continued nearly to the margin. *Below*, the forewing has three subapical lemon spots, a large white spot between the upper median branches, and two white spots beyond the cell, the one above these being obsolete. The black border of the hindwing is narrower than in *H. judith*, enclosing a lemon spot between the subcostals, one (obsolescent) above the radial veins, and one, large and diffused, between the upper median branches; an ochreous spot partly enclosed between the lower median and submedian veins; the anal angle slenderly edged with orange-ochreous.

Sumba, interior, rare. I cannot find the type, and the description is not very good. The species, however, which I compared with good series of *H. judith* and *naomi*, is distinct. It is curious that it should more approach the Javanese species than that of the neighbouring islands.

The species figured by Hombron and Jacquinot in the "Voyage au Pole Sud," as "*Pieris judith*, var.," and so placed by Kirby, is certainly distinct. I suggest for it the name of *Huphina imogene*. It is nearest my *H. ethel* from Engano.

HUPHINA VASO, n. sp.

A local form of the Javanese *H. corva*. Male, above, darker than in that species, all the veins heavily outlined with black. Forewing with an almost complete, irregular, outer-discal dark band extending obliquely from the costa to the hind-margin, and continued along the latter to its base; the white submarginal spots beyond this are large. Hindwing with the cell, the upper median, and both the subcostal veins clouded with black, the black border deeply dentate, with a line of diffused black spots submarginally in the middle of the spaces. *Below*, paler than in *corva*, the white markings, especially the submarginal ones, all larger and clearer, the veins all slenderly outlined with ochreous.

Sambawa, coast.

BELENOIS JAVA, Sparr. (*coronea*, Cr.).

Coast of Sambawa, common, not seen in Sumba. Cramer records this species from Borneo, and in the Singapore Museum there is a specimen labelled Jelebu (not far from Malacca). These localities are certainly doubtful, as the butterfly inhabits dry, sterile coasts, and would be quite out of place in forest countries like Malacca or Borneo. On the other hand a coast-butterfly of exceedingly weak flight, but able to float in the air for an indefinite time, would be more apt to be blown out to sea than other insects, and more likely to survive till its arrival in another island. So that stragglers may really have been taken remote from the true habitat of the species.

This butterfly flies like a *Hestia*, and seems to be the most perfectly protected of Eastern *Pierida*. It is hard to believe that it has anything to do with the Indian *Belenois mesentina*, the type of the genus, which is wholly different in appearance and in habits. Like *Delias* and *Prioneris* this genus has the claws bifid.

DELIAS PASITHOË, Linn., var.

I saw two specimens of this on Mt. Haruhasa in Sambawa at nearly 5000 feet elevation, but as well as I can remember neither were taken. An undescribed *Agarista* mimicking it was caught at the same place. No species of *Delias* was observed in Sumba.

DELIAS ORAIA, n. sp.

A local form of the Indian *Delias descombesii*, and greatly resembling it. It lacks, however, the black marginal band of *descombesii*, the costa and outer margin of both wings being slenderly grey, especially at the ends of the veins, the cilia lemon. Below the five subapical lunules on the forewing are yellow, not white. The female is generally brighter coloured than that of *descombesii*. The hindwing, however, is much darker over the base and disc, but below the submarginal spots are bright lemon, and the hind-margin rich ochreous.

Sambawa, 2—5000 feet, scarce. The specific name means *beautiful* in modern Greek. I thought it unnecessary to give my detailed description of the species.

The reappearance in Sambawa of a local form of an Indo-Malayan butterfly unknown to Java, is remarkable.*

* I take the opportunity to describe a new Javanese species of this genus.

DELIAS AURANTIA, n. sp.

Near *D. belisama*, Cr. Above orange over the basal half of the wings, including all the cell, the disc as far as the upper radial vein, far beyond the cell, and the hind

100. *APPIAS** *PAULINA*, Cr.

Two forms of this very puzzling group occurred both in Sumba and in Sambawa. One was all white, with only a slender dark marginal line, resembling *A. albina*. The other had the hindwing and the apex of the forewing bright ochreous-yellow below, resembling *A. lankapura*, but without the dark apex. It generally had a black or gray discal spot on the forewing.

I have dubiously recorded *Appias lyncida* from Sambawa, and from Sumba a female which I supposed to be that of *A. (Saletara) nathalia*.

101. *NEPHERONIA VALERIA*, Cr.

Sumba, Sambawa. The submarginal spots are wholly absent. No yellow female was taken. A different species is, I think, also found in Sumba.

margin to the lower angle; the rest black, its inner border serrate. Hindwing with the black border rather wide and equal. Below, forewing black with four (five in *belisama*) orange subapical streaks, the lowest obsolescent; a broad oblique orange streak borders the disco-cellular veins, the cell and median veins are outlined with pale orange, the rest of the cell irrorated with black scales, the hind-margin broadly whitish. Hindwing orange like the upperside, the red area darker than in *belisama* and much larger; the inner lunular band more continuous, a marginal yellow line (nearly obsolete in *belisama*), the inner half of the wing, including most of the cell, and on the disc from the hind-margin to beyond the middle median vein, densely irrorated with black scales. Expanse $3\frac{1}{4}$ inches. From *D. belisama* this may be distinguished by the much smaller black area of the upperside of the forewing, and the large orange areas of the underside of the same wing. The male of *D. belisama* is also usually white, or white tinged with lemon, or yellow with a slight ochreous tinge. From *D. nakula*, recently described from Java by Mr. H. Grose Smith, *D. aurantia* differs in its larger size and rich orange colour.

This fine butterfly is not uncommon on Mount Arjuno, Eastern Java; I did not take it below 2,500 feet elevation, and it is found at any rate up to 5000 feet. *D. belisama* is also common there, ranging from the low country up to about 3000 feet, so that there is a zone where both are found. When flying together *D. belisama* could always be distinguished by its smaller size, East Java specimens being apparently smaller than West Java ones, and under three inches in expanse. On visiting the great volcano of Sméra further east at a different season, I found *belisama* common, but did not see *aurantia* at all.

At 5000 feet on Arjuno, I took a single faded specimen of a *Delias* only $1\frac{1}{4}$ inch in expanse. It was white, the outer half dark, the forewing with a subapical bar and a row of five subapical spots. The hindwing had the margin broadly black with five yellow (?) spots in it below only. This species does not resemble anything known to me.

* This is one of Hübner's silly genera, grounded on nothing whatever. I use it most unwillingly as the equivalent of Mr. Wallace's *Tachyris*. Mr. Distant includes the species of *Huphina* (Mr. Wallace's *Pieris*) under *Appias*, for which I can see no reason. *Catopha* necessarily falls before *Appias*.

102. *HEBOMIA GLAUCIPPE*, Linn.
Sumba, Sambawa.

IXIAS REINWARDII, Voll.
Sambawa, a beautiful species, confined to the dry coast.

103. *IXIAS* near *PIRENE*, Linn. (*pyrene*).
Sumba, coast, several times seen, but no specimen taken.

CATOPSILIA PYRANTHE, Linn. (*evangelina*, Butler).
Sambawa, coast, not observed in Sumba.

104. *CATOPSILIA CATILLA*, Cr.
Sumba, Sambawa.

105. *CATOPSILIA CROCALE*, Cr.
Sumba, Sambawa.

106. *CATOPSILIA SCYLLA*, Boisd.
Sumba, Sambawa.

Family PAPILIONIDÆ.

107. *PAPILIO* (*ORPHEIDES*) *ERICHTHONIUS*, Cram. (*erithonius*).
Sumba, Sambawa.

108. *PAPILIO* (*ILIADÆS*) *MERAPU*, n. sp.

Male, above like *P. memnon*, the pale rays gray, and not very distinct on the forewing. Below the basal crimson areas are larger, the longitudinal streaks in the cell of the forewing obsolescent. Hindwing with the outer pale area partly gray, partly orange-ochreous, narrow, the outer line of spots many times larger than in *memnon*, the inner lying outside of the band, only touched outwardly with ochreous; a separate oblique orange stripe on the abdominal border. The species is the largest of the group, much larger than *memnon*, its expanse being six and a half inches.

Koloki, Pada Dalung, Sumba, 2—3000 feet.

This fine butterfly is called by the Sumbanese after their god Merapu, and I was told that it was after a fashion held sacred, though they raised no objection to my catching specimens. As an instance of the sober tastes of this people, I may remark that they consider this butterfly far handsomer than *Ornithoptera naias* or *Papilio maremba*.

A form of *Papilio memnon* occurs in Sambawa, but I have no specimens.

109. PAPILIO (MENEALIDES) OREON, n. sp.

A local form of *P. liris*, Godart, from Timor and Australia, differing from de Haan's figure of that species in the following particulars. Female, forewing with the dark border narrower than in *liris*, the pale area not tinged with yellow, extending nearly to the base and apparently more marked with black scales. On the hindwing below, the whitish area is less yellow, extends nearer to the base, (occupying two-thirds of the cell), and somewhat further discally, its outer margin scalloped. The white area between the costal and subcostal veins is nearer the base of that space, and is much narrower, being not so long as broad, deeply concave outwardly. There is a submarginal row of seven crimson spots (five in *liris*), that in the interno-median space joining the white area so as to enclose an oval black spot. On the upperside, the hindwing is obscurely marked with red as in *liris*. Expanse four inches and three quarters.

Of the male I have only one very bad specimen, not perceptibly different from the female.

Sumba, confined to the mountain-forests from which I have named it.

PAPILIO ARISTOLOCHIE, Linn.

Sambawa, normal.

110. PAPILIO (CHARUS) HELENUS, Linn.

Sumba, Sambawa, confined to the mountains, scarce.

111. PAPILIO (HARIMALA) MAREMBA, n. sp.

Male, above black, the markings metallic golden-green, greenish-blue in some lights. Forewing with the base, all the cell and two-thirds of the hind-margin uniformly powdered with green scales; just beyond the cell and forming a crescent round it, is a rather narrow band of rich green from the upper median vein to well above the subcostal, the veins black; beyond this a broad black band, but slightly irrorated with green; after which there is a broad subapical and submarginal area of diffused green, with separate green spots between the median veins; the outer and costal margins dark; the hind-margin rich green near the lower angle as far as the middle of the interno-median space. A large unbroken dark discal patch covered with a cottony mass of odoriferous hairs, extending from the internal to the upper median vein, reaching inwardly almost to the cell, and outwardly along the veins almost to the outer margin. Hindwing with the base powdered with green, a broad green discal area, scarcely entering the cell, extending from the upper subcostal vein to the hind margin, broadest discally;

beyond it are four green lunules; the outer part, including the tails, black. *Below*, the basal half of the wings deep brown, sparsely sprinkled with whitish scales; forewing with a pale transverse band (much more remote from the base than in *P. pericles*), broad subapically, deeply indented outwardly, below the two lower subcostal branches, abruptly narrowed below the lower radial vein. Hindwing with the disc somewhat whitish subabdominally, a row of seven narrow ocelli, outwardly silvery blue, inwardly dull reddish. The ocelli are much further from the outer margin than in the allied species.

This splendid species has no near allies. Apart from the sex-mark, it is nearest *Papilio brama*, Guérin, from Borneo, Sumatra, and the Malay Peninsula.

Sumba, rare near the coast, commoner in the remote interior.

PAPILIO (HARIMALA) PERANTHUS, var.

Sambawa, mountains, scarce. It seems to be intermediate between the Javanese *P. peranthus* and the Timorese *P. pericles*. I have unluckily neither specimens nor description.

112. PAPILIO (ZETIDES) SARPEDON, Linn.

Sumba, Sambawa. In Sumba the red markings of the underside are sometimes replaced by ochreous-yellow.

113. PAPILIO (ZETIDES) EURYPYLUS, Linn.

Interior of Sumba and Sambawa, scarce.

114. PAPILIO (ZETIDES) AGAMEMNON, Linn.

Sumba, Sambawa.

PAPILIO (PATHYSA) ANTIPHATES, Linn.

Sambawa.

115. PAPILIO (LAERTIAS) POLITES, Linn.

Sumba, Sambawa. In Sambawa one female imitates *P. aristolochia*, while another is similar, but with the white discal area replaced by orange. I have unluckily not recorded any female from Sumba.

116. ORNITHOPTERA NALAS, n. sp.

Male, above black, all the veins except at the extreme base and the internal vein, bordered with paler brown, a longitudinal pale streak generally in the middle of the cell. Hindwing with a small translucent golden patch occupying less than half of the cell longitudinally (its lower and basal part black); four discal spaces from the costal to the upper median vein occupied by quadrate golden areas, which are not incised outwardly as in *O. plata*; that between the costal and subcostal

veins is largest and broadest; a slight diffused golden touch between the upper and middle median veins, more distinct below. *Below* forewing with the pale streaks whitish. Collar and patches at the base of the wings red, abdomen mostly black except around the base of the valves.

Female, above dark brown, the pale streaks conspicuous, whitish, occupying two-thirds of the cell, where they are broad with a conspicuous streak in the middle. Hindwing with a golden area, slightly duller than that of the male, occupying two-thirds of the cell transversely, and six discal spaces from the subcostal to the fold above the submedian vein; a minute spot above the subcostal, the next area small, all of them deeply incised outwardly; four pairs of triangular yellow spots are usually on the outer disc, sometimes connected with the inner golden area, by rays of the same colour; there are a few marginal internervular ochreous touches. Collar and base of wings red, abdomen laterally and ventrally yellow with black spots.

The outer margin of the hindwing of the male is remarkably straight; its expanse is nearly five and a half inches.

The golden area of the male differs considerably from that of *O. criton* (Moluccas) and *O. plato* (Timor). In *plato* the red patches are wanting; the female is unknown. In *criton* the female is altogether different. The golden area is probably smaller in *naias* than in any other species of *Ornithoptera*, except the Celebesian *O. haliphron*, in which the cell is all black.

Common in Sumba, both on the coast and in the interior.

In Sambawa occurs a variety (*O. naias*, var. *sambavana*), which is considerably larger. The female has a large golden spot between the costal and subcostal veins of the hindwing; the outer discal pairs of spots are always connected with the central golden area, more or less enclosing large black spots. In the male a golden spot (generally present, though small in the Sumba form) between the upper and middle median veins, is always absent, while in *O. criton*, and presumably in *O. plato*, it is the largest of all.

Family HESPERIADÆ.

Ismene Group.

117. PARATA MALAYANA, Feld.

Sumba. Another species occurred both in Sumba and Sambawa.

118. HASORA BADRA, Moore.

Sumba, Sambawa. Both this and the Engano form resemble the Javanese, which may and may not be the same as the typical Indian one.

119. *BADAMIA EXCLAMATIONIS*, Fab.
Sumba, Sambawa.

Tagiadas Group.

120. *TAGIADAS BRASIDAS*, n. sp.

Male, above dark brown, forewing with three subapical hyaline spots, the first two approximate, the third well beyond the others, small; two small discal spots, and two terminally in the cell, hyaline. Hindwing, dark brown, the disc irrorated with gray scales, and bearing three dark spots; the abdominal angle widely white enclosing a line of marginal dark spots, the cilia long, white. *Below*, forewing touched with whitish below the lower median vein. Hindwing two-thirds white, with three subapical dark discal spots (the first united with the apical dark area), the second and third unequal, united, the apical border widely dark brown, tapering to below the lower median vein, twice interrupted by white. Female, like the male the hyaline spots in the cell of the forewing united, the discal ones larger, the brown spots bordering the hindwing smaller above and below.

Sumba, and Sambawa, but those from the latter island may not be quite the same. The species is nearest *T. helferii* from the Nicobars, differing in the white border on the hindwing above, and the dark spots on it below. The hyaline marks are also more conspicuous. This also distinguishes it from *T. alicia*, which is also much less white below, and has no distinct marginal spots.

121. *ABARATHA SYEICHTHUS*, Feld.

Above, the discal yellowish bands of the hindwing are much narrower and more obscure than in Indian specimens. Below, the forewing is altogether less white; the white mark beyond the cell, which in the Himalayan form extends far outwardly, is reduced to a slender, transverse crescent; the costal streaks above it are obsolete, the submarginal line of quadrate spots much smaller, and the fifth hyaline spot absent.

Sumba, rare.

ABARATHA HYPERIDES, n. sp.

Very near *Pterygospidea helias*, Felder, from the Celebes, but the bands and the discal hyaline spots of the forewing are absent, and the apex of the hindwing is broadly dark. From *A. sura* it differs in the forewing, which is almost uniform dark brown above and below. The hindwing is also less variegated above and below, the white area is larger and more uniform, the inner line of spots is obsolete, the outer

united, and dark apically, nearly obliterated by white scales in the median spaces.

Sambawa. Another species, more like *A. angulatus*, was found in Sumba, but no specimens have survived.

Gehlota Group.

122. *COLADENIA DAN*, Fab.

Sumba, Sambawa. Two species are confused under this name, and occur together in Sumba, Borneo, the Malay Peninsula, and perhaps elsewhere. They differ obviously in flight and in prehensors, but I cannot at present point out any difference in the markings. One has an egg with numerous ribs (over forty) as in *Gehlota*, the other with few (seventeen) as in *Tagiades*.

Pamphila Group.

123. *TILICOTA MESOIDES*, Moore.

Sumba, Sambawa. The orange bands are smaller and narrower than in Indian specimens, and the ground-colour dark below.

124. *TILICOTA NIGROLIMBATA*, Snellen.

Sumba, Sambawa. This is the species figured by Mr. Distant; I am not quite sure of its identity with Heer Snellen's species.

125. *TILICOTA GOLA*, Moore.

Sumba, Sambawa.

126. *AMPITTIA MARO*, Fab.

Sumba, Sambawa. I am doubtful of its identity with the Indian form.

Baoris Group.

127. *CHAPRA MATHIAS*, Fab.

Sumba, Sambawa.

128. *PAENARA NAROOA*, Moore.

Sumba, Sambawa. I am not sure of its identity with the Ceylon form.

Suastus Group.

129. *SUASTUS CHILON*, n. sp.

Above, male all dark brown, no hyaline markings nor patches of lighter-brown scales. *Below*, forewing with a minute white dot dis-cally in the lower median space, the subapical hyaline spots represented by two slight dark streaks, the lower (in one specimen) containing a

lighter dot. Hindwing nearly white (not gray as in *S. gremius*), the borders dark, a conspicuous black cell-spot, and a row of black discal spots, six in one specimen, four in the other.

The absence of hyaline spots distinguishes it from all others.

Two males, Sumba coast.

Kerana Group.

130. PLESIONEURA RESTRICTA, Moore.

Sumba, Sumbawa, mountains. My Sumbanese *Hesperidae* have suffered more than any other family, and I have been compelled to omit a number of species, a *Halpe*, two *Parnaras*, a *Parata*, etc.

EXPLANATION OF PLATE II.

- | | |
|--|---|
| Fig. 1. <i>Euplua clivesii</i> , n. sp. ♀. | 7. <i>Radena kambara</i> , n. sp. |
| 2. <i>Euplua lewa</i> , n. sp. | 8. <i>Nacaduba gaura</i> , n. sp., × 2. |
| 3. <i>Euplua palmedo</i> , n. sp. | 9. <i>Nacaduba laura</i> , n. sp. ♀, × 2. |
| 4. <i>Danais litoralis</i> , n. sp. | 10. <i>Lampides anops</i> , n. sp. |
| 5. <i>Danais orientis</i> , n. sp. | 11. <i>Lampides masu</i> , n. sp. |
| 6. <i>Radena oberthurii</i> , n. sp. | 12. <i>Huphina julia</i> , n. sp. |

VI.—*Natural History Notes from H. M. Indian Marine Survey Steamer 'Investigator,' Commander R. F. Hoskyn, R. N., commanding. No. 24. List of Deep-sea Holothurians collected during seasons 1887 to 1891, with descriptions of new species.—By Dr. J. H. TULL WALSH. Communicated by the SUPERINTENDENT OF THE INDIAN MUSEUM.*

[Received 6th April, 1891;—Read 7th May, 1891.]

Order ELASIPODA, Théel, Chall. Rep. vol. iv, Hol., p. 9.

Family Elpididæ, Théel, l. c., p. 10.

1. PENIAGONE WYVILLII, Théel, Chall. Rep. vol. iv, Hol., p. 42.

One specimen.

Station 118, 15th December, 1890, Bay of Bengal, lat. 12° 20' N., long. 85° 8' E., 1803 fathoms, globigerina ooze, bot. temp. 35° Fahr. (*Alcock*).

Family Deimatidæ.

2. ONEIROPHANTA MUTABILIS, Théel, l. c., p. 62.

One specimen.

April 12th to 13th, 1888, Andaman Sea, N. Sentinel Island bearing N. 15°, W. 18 miles, 250 fathoms (*Wood-Mason*).

3. *DEIMA VALIDUM*, Théel, l. c., p. 68.

"Transparent light orange-red" (*Alcock*).

Two specimens.

April 16th, 1888; Bay of Bengal lat. 11° 15' N., long. 91° 16' E., 1840 fathoms, globigerina ooze (*Wood-Mason*).

A fine specimen; length 130 mm.; length of lateral processes 55 mm.

Station 117, Bay of Bengal, lat. 11° 58' N., long. 88° 52' 17" E., 1748 fathoms, globigerina ooze with pieces of pumice, bot. temp. 35.3° (*Alcock*).

4. *DEIMA FASTOSUM*, Théel, l. c., p. 71.

Two specimens.

Station 118 (*Alcock*).

April 20th, 1888, Bay of Bengal, lat. 6° 18' N., long. 90° 40' E., 1520 fathoms (*Wood-Mason*).

5. *ORPHNURGUS ASPER*, Théel, l. c., p. 82, var. *GLABER*, nov.

Length 150 mm. Body smooth and leathery; skin thin and somewhat transparent; colour (in spirit), body greyish white, feet and processes yellowish-white. Tentacles 20, long, extremities brownish and dendritic. Ventral feet 24 on each side, longer and larger near the centre of the body than at the extremities; longest 20-25 mm. Lateral processes generally longer than the pedicles, longest 30 mm. Two rows of processes on each dorsal ambulacrum; these vary considerably in size and short and long ones seem to alternate. Mouth slightly ventral; anus terminal and patulous. Calcareous bodies, spinous rods or smooth rods with branched extremities. Polian vesicle single, 40 mm., wider in the middle than at the extremities.

Colour in the fresh state "salmon-pink" (*Alcock*).

One specimen.

Station 112, 7th November, 1890, Bay of Bengal, lat. 13° 47' 30" N., long. 92° 36' E., 561 fathoms, grey mud, bot. temp. 44.9° (*Alcock*).

6. *PANNYCHIA WOOD-MASONI*, n. sp.

Length 300 mm. Body long, flat and of equal width throughout; skin calcareous but flexible; colour yellowish-white (in spirit). Tentacles 20, round, with stumpy digits; mouth ventral; anus terminal and without teeth. The middle ambulacrum of the trivium bears a double row of feet; outside these the skin is thrown into longitudinal and transverse folds and ridges; the skin on the dorsum is also ridged.

The lateral ambulacra possess a single row of feet much longer than those on the middle ambulacrum. Just above this row of feet there is a row of long, pointed processes; these average 15 mm. in length. The dorsal ambulacra are furnished with a double row of processes; these are shorter than the lateral ones. One polian vesicle; one stone canal; one bundle of genital tubes which are branched and furnished with irregular dilatations. Calcareous ring small; muscles poorly developed relatively to the size of the animal.

The calcareous bodies consist of large, round, many-holed plates, the centre holes being fairly large, the marginal ones small. Besides these plates, a few branched spicules are present.

This species is very closely allied to *P. moseleyi*, Théel.

Numerous specimens.

January 2nd, 1888, off Port Blair, 271 fathoms, bottom green mud (Giles).

April 12th, 1888, 7½ miles E. of N. Cinque Island, Andaman Sea, 490 fathoms, bottom green mud (Wood-Mason).

Station 115, December 9th, 1890, Andaman Sea, lat. 11° 31' 40" N., long. 92° 46' 40" E., off Dyer Point and N. of Cinque island, 188—220 fathoms, green mud, bot. temp. 56° (Alcock).

AMPHIGYMNAS, nov. gen.

Body ovoid with narrow tail-like extremities; soft and appears to have been surrounded by a jelly-like material when fresh. Feet very numerous over the whole of the trivium and placed more or less irregularly. Lateral margins with two or three rows of long processes. Back covered with processes except near the mouth and anus where the body tapers and where the dorsal surface is naked; mouth terminal, small; tentacles 15, very small and retracted; anus terminal, small. Calcareous bodies moderate sized, irregularly rounded, many-holed plates somewhat like those of *Pannychia*. Calcareous ring of 5 small pieces loosely connected.

7. AMPHIGYMNAS MULTIPES, n. sp.

Length 80 mm., width near middle of body 22 mm.; at oral and anal end about 9 mm.

Colour (in spirit) pale purple; the long lateral processes measure about 15 mm., those on the back are shorter.

One specimen.

Station 115, December 9th, 1890, Andaman Sea, lat. 11° 31' 40" N., long. 92° 46' 40" E., off Dyer Point and N. of Cinque island, 188—220 fathoms, green mud, bot. temp. 56° (Alcock).

8. *LETMOGENE SPONGIOSA*, Théel, l. c., p. 80.

Crydora spongiosa, Théel, Prelim. Rep. Hol., p. 9.

One specimen.

April 17th, 1888, Bay of Bengal, lat. 10° N., long. $91^{\circ} 7'$ E., 1924 fathoms (*Wood-Mason*).

Family Psychropotidæ.

9. *EUPHRONIDES DEPRESSA*, Théel, Chall. Rep. vol. iv, Hol., p. 93.

Two specimens.

Station 118, December 15th, 1890, Bay of Bengal, lat. $12^{\circ} 20'$ N., long. $85^{\circ} 8'$ E., 1803 fathoms, globigerina ooze with pieces of pumice, bot. temp 35° (*Alcock*).

10. *BENTHODYTES PAPILLIFERA*, Théel, l. c., p. 102.

"Colour purple, tentacles violet black" (*Alcock*).

Two specimens 80 mm. and 68 mm. The fringe is damaged and the dorsal papillæ are not very apparent, but there is, I think, no doubt that these specimens belong to Théel's species.

Station 117 (*Alcock*).

11. *BENTHODYTES SANGUINOLENTA*, Théel, l. c., p. 104.

Four specimens; the row of transverse papillæ below the mouth cannot be made out.

Station 118 (*Alcock*).

12. *BENTHODYTES OVALIS*, n. sp.

Length 65 mm.; breadth 35 mm. including fringe. Body oval, flat, about twice as long as broad and of a deep purple colour throughout. A double row of feet on the middle ambulacrum of the trivium and no transverse line of feet below the mouth; lateral ambulacra with a single row of feet. On the dorsal surface there are three processes on each ambulacrum; the anterior ones are 15 mm. long and not retractile (?); the posterior one in each row is much smaller. Mouth ventral; anus terminal. Tentacles retracted. Calcareous bodies small 3-armed spicules with the end of each arm clubbed and perforated.

One specimen.

April 12th, 1888, $7\frac{1}{2}$ miles E. of N. Cinque island, Andaman Sea, 490 fathoms, bottom green mud (*Wood-Mason*).

13. *BENTHODYTES GELATINOSA*, n. sp.

Before describing the specimens as they appear in spirit it will be better to quote the graphic account of the fresh creature given by Dr. Giles, Ad. Rep. Marine Survey of India, 1877-88, p. 15:—

"I must first notice a very peculiar holothurian, several specimens of which were included in the catch..... When fresh, the animal consists of a tough, muscular sac of a yellowish pink colour, enclosed in a thick coating of perfectly transparent, deep violet jelly. Though not sticky or glairy, this jelly is of so delicate a consistence that it was almost impossible to clean the mud from the animal without stripping off the coating. After a short exposure to the action of spirit, this jelly, previously fully 15 mm. thick, shrinks to a thickness of less than 5 mm. and becomes comparatively dense. The violet colouring matter dissolves out into the spirit and exhibits a curious affinity for vegetable tissues, deeply staining a paper label which had been placed within the jar. Its attraction for animal tissues though less marked was still very noticeable, the nuclei shewing it best, so that, on microscopical examination, the animal was found to have become stained in a solution of its own pigment. After hardening in spirit it is sufficiently evident that this jelly-like coating cannot be considered as a secretion, but as an integral part of the tissues of the body wall, as it consists of a plexus of stellate and spindle-shaped cells, enclosing within their meshes many nuclei-form bodies and much hyaline connective substance. The body wall contains but few calcareous bodies. Those present consist of a circular plate, having articulated to it a fan-shaped body so shaped that the whole apparatus is not unlike the badge of a grenadier's cap. As far as could be made out, these bodies appertain to the jelly-like layer and not to the dense inner portion of the body wall, which latter appears to be purely muscular. When laid open it is seen that the lungs are very complex and racemose, and that the ambulacral tubes which are very small and straight, give off a number of minute branches to the pedicels. The polian vesicle is of moderate size and simple. It appears to belong to, or be near, the genus *Benthodytes* of the *Elasipod* family *Psychropotidae*, and is doubtless new as it is identifiable with neither of the species in the Challenger monograph. In all probability, many of its allies must share with this species the peculiarity of a jelly-like coating but no mention of any such appearance is made in the above report."

Length of specimens 100 to 140 mm.—Body soft; extremities and ventral feet are still of a beautiful purple, the body generally is a dark lavender colour (in spirit). The anterior end of the body has a wide lappet-like fringe furnished with papillæ and the border is continued down each side of the flatly cylindrical body and expanded in a less pronounced form as an anal lappet. Along the lateral border there are numerous feet in a single row. Mouth small and ventral; anus large, patulous and somewhat dorsal, situated just above the anal lappet.

Tentacles 15, very small and of a deep purple colour. One, or two, fairly large Polian vesicles; calcareous ring very minute, a double row of sucker feet on the middle ambulacrum of the trivium; these are more numerous and closer together near the anal end while they are entirely absent from the surface just below the mouth. A single row of fairly long, slender retractile papillæ are seen on each dorsal ambulacrum. Two small bundles of genital tubes. The longitudinal muscles are well-developed, broad, orange-yellow and visible through the skin—only a few calcareous spicules and granules were found in the muscular substance of the body.

Many specimens.

January 2nd, 1888, Andaman Sea off Port Blair, 271 fathoms, (*Giles*), two specimens.

April 12th, 1888, Andaman Sea $7\frac{1}{2}$ miles E. of N. Cinque island, 490 fathoms, (*Wood-Mason*). Two specimens.

Station 115, December 9th, 1890, Andaman Sea, lat. $11^{\circ} 31' 40''$ N., long. $92^{\circ} 46' 40''$ E., off Dyer Point and N. of Cinque island, 188—220 fathoms, green mud, bot. temp. 56° (*Alcock*). Eleven specimens.

APODOGASTER, nov. gen.

Body long flat and worm-like; slightly wider anteriorly than posteriorly. Down both sides there is a fringe, like that of *Benthodytes*, in which are seen the long tubes of numerous sucker feet. The odd ambulacrum is naked while the lateral ambulacra have one row of small sucker feet situated just below the lateral fringe. A single row of papillæ on each dorsal ambulacrum. Tentacles 15, mouth ventral. Anus terminal.

Calcareous bodies small granules and small wheel-like plates.

14. APODOGASTER ALCOCKI, n. sp.

Length 80 mm.

"Light pink" (*Alcock*).

One specimen.

Station 112, November 7th, 1890, Bay of Bengal, lat. $13^{\circ} 47' 30''$ N., long. $92^{\circ} 36' E.$, 561 fathoms, grey mud, bot. temp. 44.9° (*Alcock*).

Order APODA.

Sub-order PNEUMONOPHORA.

Family Molpadidæ.

15. ANKYRODERMA DANIELSSENI, Théel, Chall. Rep. vol. xiv, Hol. p. 39.

One specimen.

April 11th, 1888, 7 miles S. E. by S. of Ross Island, Andaman Sea; 265 fathoms, green mud (*Wood-Mason*).

16. *A. MARENZELLERI*, Théel, l. c., p. 41.

One specimen.

Station 55, 13th—14th April, 1889, Bay of Bengal, 30 miles W. of Middle Andaman Island (Cape Bluff), 480—500 fathoms, globigerina ooze (*Alcock*).

17. *EUPYRGUS SCABER*, Lütken, Videnskab. Meddel. Kjöbenhavn 1857, p. 23.

After some indecision as to the correct title of these specimens I have thought it better to name them as above. In general form and appearance they all agree with *Echinostoma hispidum*, Semper, Reis. in Philipp. 1867, p. 44, but differ from *Eupyrgus hispidus*, Barrett, Ann. Mag. Nat. Hist. xx, p. 46, which Semper gives as a synonym of his *Echinostoma*, in having no sucker feet. In the Challenger Report, vol. xiv, Hol. p. 49, *Echinostoma hispidum*, Semp. is given as a synonym of *Eupyrgus scaber*, Lütken, and the *Eupyrgus* of Barrett is not mentioned.

Three specimens.

January 15th, 1888, off Cinque island, Andaman Sea, 650 fathoms (*Giles*.)

Station 107, October 23rd, 1889, Laccadive Sea, lat. 8° 23' N., long. 75° 47' E., 738 fathoms, green mud, bot. temp. 41.9° (*Alcock*).

Station 116, December 9th, 1890, Bay of Bengal lat. 11° 25' 5" N., long. 92° 47' 6" E. off Cinque and Rutland islands, 405 fathoms, green mud, bot. temp. 47° (*Alcock*).

18. *TROCHOSTOMA ANDAMANENSE*, n. sp.

This species is very closely allied to *T. antarcticum*, Théel, Chall. Rep. vol. xiv, Hol. p. 44.

Length 90 mm.

Body cylindrical and narrowed behind into a tail-like extremity. Skin rough but thin. In spirit the ground colour has become greenish-grey and the spots are more or less blood-red. The tentacles fifteen in number, are yellow and between every two there is, near the base, a blue-black triangular mark. Each tentacle has three very small digits at the free end. Mouth and anus terminal, the latter without teeth, but in one specimen with numerous fine papillæ. One Polian vesicle, one free stone canal; 2 (3?) lungs, not much branched; one bundle of yellow genital tubes. The radii of the calcareous ring are produced backwards as spines. Calcareous bodies not very numerous consisting of a few wide-holed plates and tower like rods which are perforated with 4 to 6

holes. Under the microscope the "chocolate spots" are seen to be composed of rust-red ovoid bodies with a concentric arrangement. They are insoluble in caustic potash.

Colour when fresh "dirty flesh-colour with closely placed deep chocolate spots; the crown (tentacles) being a sort of raw-meat-colour." (*Giles*).

Three specimens.

December 8th, 1887, S. E. of Cinque island, Andaman Sea, 500 fathoms, green mud (*Giles*).

VII.—*On an undescribed Oriental species of Nepeta*.—By D. PRAIN.

[Received 2nd March 1891;—Read 6th May 1891.]

(With Plate III).

In the Calcutta herbarium there is an example of a very distinct species of *Nepeta* which appears to be as yet undescribed. As it has been collected beyond the frontiers of India, it could not with propriety be dealt with in the paper on *Indian Labiatae* recently read before the Society. But as it possesses rather more interest than isolated new species usually do, its position in the arrangement of Oriental species elaborated by the late M. Boissier in his work (*Flora Orientalis*, iv, 637-670) dealing with the area in which it occurs, as well as a description and figure of the specimen, are now presented.

NEPETA LINN.

SECT. I. EUNEPETA. SERIES I. Perennes. *Sub-ser.* 2. *Nuculae tuberculatae*.

§ MACROSTEGLE *Boiss.*, *Flor. Orient.* iv, 638 (1879) *ampl.*—Verticillata densiflora vel laxiuscula remota. Bractea ovata vel oblonga. Calyx fauce pilosus vel glaber.

1. Calyx ore obliquus.

a. Calyx fauce pilosus.

N. Bellevii.

b. Calyx fauce glaber.

N. glomerulosa, *N. juncea*.

2. Calyx ore rectus, fauce glaber.

N. Scordotis, *N. Sibthorpii*, *N. leucostegia*.

34 b. NEPETA BELLEVII *Prain*; robusta, laxe lanata nivea, caulibus elongatis ramosis puberulis obtuse 4-gonis superne subundis, foliis masculis sursum decrescentibus oblongo-lanceolatis crenato-dentatis

subrugosis apice acutis basi truncatis, utrinque parce lanatis, inferioribus breve petiolatis cæteris subsessilibus ramealibus ovato-lanceolatis; *verticillastris* multifloris laxiusculis remotis, *bracteis* membranaceis ovato-lanceolatis acuminatis cucullatis calyces æquantibus, *calycis* longe hirsuti membranacei striati ore intus piloso obliquo, dentibus e basi angustiore subulatis summo tubum subæquante cæteris eo brevioribus, *corolla* extus puberula tubo calycis fere 2-plo longiore, staminibus exsertis, *nuculis* oblongis apice obtusis ruguloso-tuberculatis areolâ ferri equini calcaribus sursum directis ad instar affixis.

AFGHANIA AUSTRALI; inter Kandahar et Kekat-i-Ghilzai, *Bellew!*

Rhizoma deest, caulis p. q. s. plus quam 40 cm. basi fere 1 cm. crassa, foliis inferioribus 7—9 cm. longis his 3—4 cm. latis petiolis 1 cm. longis, ramealibus 2—3.5 cm. longis his 1—1.5 cm. latis sessilibus, *spicâ* 12 cm. longâ hæc 2.5 cm. latâ verticillastris (10—12) 12—16-floris imis 2 cm. summis vix 1 cm. remotis, bracteis 12 mm. longis 4—7 mm. latis 3—5-nervis coloratis supra glabris subtus nervis sub lente puberulis margine prope basin parce pilis albis divergentibus orato prope apicem acuminatum et tandem subulatum sub lente puberulo, *calyce* tandem subulato pedicello 1—2 mm. longo tubo 6 mm. longo dentibus summo 6 mm., lateralibus 5 mm., imis 4.5 mm. longis extus æqualiter pilis albis divergentibus parce hirsuto intus tubi ore et dentium basibus pilis simillimis densissime piloso, *corolla* tubo 11 mm. longo extus puberulo colore forsan rosea, *nuculis* 3 mm. longis his 1.5 mm. latis brunneis areolâ albâ calcaribus 1 mm. longis exceptâ manifeste rugulosis.

This very distinct species bears a considerable resemblance to the European *N. tuberosa* but the whorls are much more lax, the calyx mouth is oblique and no secondary reticulation of the membranous bracts is visible. The outlets of this species also closely resemble those of *N. tuberosa* and *N. Sibthorpii*, but in these European species there is a much smaller areola of the usual type. Among Oriental species it most resembles *N. glomerulosa*, next to which it is now placed, but it differs greatly in having much larger bracts, much longer spikes and many-flowered lax whorls with longer corollas. The outlets too are different—the horse-shoe shaped areola of the outlets being perhaps the most remarkable character of this species; its pilose calyx-mouth also is a character which connects it with § *Psilonepeta* Benth. to which group this feature has hitherto been supposed to be confined. Except for this character, however, it is much more obviously related to the other *Macrostegia* than to any *Psilonepeta*. Perhaps a preferable view would be to look on this species as the type of a new sub-section intermediate between *Psilonepeta* and *Macrostegia* though to be

placed in the same subseries as the latter. This subsection would be characterised as follows:—

§ *PSILOSTEGIA* (*Sub-sect. nov.*),—*Verticillastra laxiuscula remota*. Bractea ovata. Calyx fauce pilosus. (Nuculae tuberculatae).—Inter *PSILONEPETAS Benth.* et *MACROSTEGIAS Boiss.* quasi medianas et proximae *MACROSTEGIAS (Boiss., Flor. Orient. iv, 638 et 651) antepenultimae.*

EXPLANATION OF PLATE III.

NEPETA BELLEVII, Prain.

- | | |
|---|-----------------------------|
| Fig. 1. Bract. | Fig. 4. Corolla tube. |
| 2. Calyx. | 5. Upper lip of Corolla. |
| 3. The same, laid open. | 6. Nutlet, external aspect. |
| 7. The same, showing inner face with characteristic horse-shoe shaped <i>areola</i> . | |

VIII.—*Noviciae Indicae.* IV. *Two additional species of GLYPTOPETALUM.*—
By D. PRAIN.

[Received 20th April, 1891;—Read 6th May, 1891.]

In a collection of plants from Great Coco, an island thirty miles north of N. Andaman, are complete examples of a species of *Glyptopetalum* nearly related to *G. zeylanicum*, Thw. This plant was previously collected, but only in fruit, either in Tenasserim or the Andamans by Helfer* and an example of the gathering (Helfer n. 1973) was described by Kurz in this *Journal* (vol. xli, [1872], pt. 2, p. 299) as *Euonymus calocarpus*, Kurz. The same plant (specimens also incomplete) has been reported from Kondil (Nicobars) by Calcutta garden collectors, and this gathering, along with a plant from Great Nicobar (Novara 188, Jelinek 245; not represented at Calcutta), has been

* This gentleman, as members are aware, was assassinated in North Andaman; his collections were disposed of in Europe and distributed thence; the circumstances of his decease rendered it impossible to differentiate positively his Tenasserim from his Andamans specimens. Kurz believed this gathering (n. 1973) to be from Tenasserim, and Lawson (*F. B. I.* i, 612) gives the same locality; I am not aware, however, what authority these authors had for this definite decision, the *Herb. Calcutta* specimen on which Kurz based his description yields none.

referred by Kurz in this *Journal* (vol. xlv, [1876], pt. 2, p. 123) to *Euonymus javanicus* Blume.*

As it is necessary to formally remove the plant from the genus to which it has hitherto been referred† the opportunity has been taken of providing a full description; at the same time a description is given of a second species collected—in fruit—in the Mishmi Mts. by Griffith; a synopsis of all the species now known precedes these descriptions.

In the four species of which the fruit is known, the dorsal raphe does not terminate at the organic base of the seed but there divides into 3-4 lacinate segments of the same appearance and consistence as the raphe itself; from the raphe they only differ in being slightly branched and in not quite reaching the hilum. They form a closely adherent

* No opinion can be expressed here regarding the Novara expedition specimens; no example of *E. javanicus* has hitherto been obtained in the Nicobars by Calcutta collectors.

† It is true that Kurz did not think *Glyptopetalum* Thw. generically separable from *Euonymus* Linn. for in this *Journal* (vol. xlv, [1875], pt. 2, p. 259) he formally relegates it to *Euonymus* (as a section) and in the *Forest Flora of British Burma* (vol. i, [1877], p. 249) he does not accord *Glyptopetalum* even sectional rank. It must also be pointed out that Bentham and Hooker had already (*Gen. Pl.* i, [1862], p. 361) pointed out how slight are the characters—the principal one being the solitary pendulous ovules—that separate *Glyptopetalum* from *Euonymus*; Baillon also (*Hist. des Plantes*, vi, [1875], p. 1, footnote) takes the same view as Kurz. If therefore the views of Kurz and Baillon ultimately prevail this plant will again be known as *Euonymus calocarpus* Kurz.

But while this is the case it will be seen on referring to the place of its publication that Kurz did not recognise in this species an example of his own section *Glyptopetalum*. In the *Flora of British Burma* too the generic description of *Euonymus* given by Kurz implies that the cells of the ovary are at least 2-ovuled—an implication opposed to his own statement (*J. A. S. B.* xlv, pt. 2, 159) as regards *Glyptopetalum sclerocarpum* and, as regards the species under review, incorrect. Kurz's views regarding the generic position of *Glyptopetalum* may therefore, I think, be ignored, and Baillon's authority can hardly be quoted in Kurz's support since that author takes so comprehensive a view of *Euonymus* that he is prepared to merge in it not merely *Glyptopetalum* Thw. but also *Lophopetalum* Wight, a step which Kurz has nowhere proposed. Moreover the genus *Glyptopetalum*, as founded by Thwaites (*Hook., Jour. Bot.* viii, [1856], p. 267), is sustained by Bentham and Hooker (*Gen. Pl.* i, [1862], p. 361), by Hooker and Thwaites (*Enum. Pl. Zeylan.* [1864], p. 73), by Bédome (*Flor. Sylvest.* i, [1874], t. 162), by Lawson (*Flor. Brit. Ind.* i, [1875], p. 612), by Trimen (*Cat. Ceylon Pl.* [1885], p. 18) and by Durand (*Index Gen. Phaner.* [1888], p. 66); considering too the large number of species of *Euonymus* proper already described and the ease with which species of *Glyptopetalum* are distinguished from these, it appears inadvisable at present to follow Kurz and Baillon in suppressing the latter genus. The present retention of *Glyptopetalum* moreover disturbs the synonymy of only one species instead of changing that of several.

arillar structure with meridional segments differing in colour from the testa that it overlies. The presence of this arillus proves that the coloured "aril" (which in turn loosely overlies it) is not a true arillus but, as Planchon has pointed out as regards the aril of *Euonymus*, an arillode.

GLYPTOPETALUM THWAITES.

Characters of *Euonymus* but ovules solitary and pendulous from the apex of the cell.

Cymes shorter than leaves, flowers under 20 mm. diam. :—

Fruit quite smooth :—

Peduncles $\frac{1}{2}$ as long as leaves, *cymes lax*
2-3 times divided; leaves thinly coriaceous oblong-lanceolate serrate, petals oblong, flowers 12 mm. diam., arillode covering $\frac{2}{3}$ of seed and lobed at the margin ... 1. *G. zeylanicum*.

Peduncles very short, cymes dense 1-2 times divided :—

Cymes longer than petioles, peduncles longer than pedicels; leaves coriaceous ovate-oblong entire, nerves indistinct, petals oblong, flowers 8 mm. diam., arillode covering $\frac{2}{3}$ of seed and lobed at the margin ... 2. *G. calocarpum*.

Cymes not exceeding petioles, peduncles shorter than pedicels, leaves membranous, ovate-oblong serrate, nerves prominent beneath, arillode hardly covering $\frac{1}{3}$ of seed, margin entire... 3. *G. Griffithii*.

Fruit rough tubercular; peduncles $\frac{1}{2}$ as long as leaves, *cymes lax* 2-3 times divided, leaves thickly coriaceous, oblong-lanceolate serrate, petals suborbicular, flowers 8 mm. diam., arillode covering $\frac{1}{2}$ of seed, margin sinuate... 4. *G. sclerocarpum*.

Cymes longer than leaves, flowers over 20 mm. diam.; *cymes lax* once divided, leaves thinly coriaceous, oblong-elliptic entire, petals obovate, flowers 30 mm. diam.; fruit unknown ... 5. *G. grandiflorum*.

1. *GLYPTOPETALUM ZEYLANICUM* Thwaites. (*F. B. I.* i, 612).
INDIA AUSTRALIS; ZEYLANIA.

2. *GLYPTOPETALUM CALOCARPUM*.—(*Euonymus calocarpus* Kurz, *Jour. As. Soc. Beng.* xli, ii, 299 (1872) et xliv, ii, 159 (1875); *Laws. in Hook. f., Flor. Brit. Ind.* i, 609 (1875); Kurz, *For. Flor. Brit. Burma* i, 249 (1877).—*E. javanicus* Kurz, *Jour. As. Soc. Beng.* xlv, ii, 123 [*Veg. Nicobar.* 19] (1876), *saltem in parte, haudquaquam E. javanicus Blume.*)

TENASSERIM?; *Helper* (n. 1973). N. ANDAMANS; Great Coco, Little Coco, Table island, Prain. NICOBARS; Kondil, *Calcutta garden collectors.*

Arbuscula glabra ramulis glabris teretibus, foliis coriaceis supra viridibus subtus prasinis elliptico-oblongis utrinque cuneatis apice obtusis margine integris nervis 5—6-paribus indistinctis, petiolis brevibus; cymis axillaribus vel parum extra-axillaribus paucifloris densioribus pedunculis semel rarius iterumque divisis; pedicellis brevibus; floribus parvis albo-viridescens, calyce 4-lobo lobis omnibus rotundatis exterioribus interioribus 2-plo minoribus, corollae petalis 4 oblongis obtusis planis prope basin 2-foveolatis; staminibus 4 filamentis erectis in disci angulis obtusis insertis, loculis antherarum basi divergentibus; ovario pyramidato-tetragono disco affixo stigmate minuto, 4-loculare; ovulis in loculis singulis pendulis anatropis, raphi extrorsa; capsula 1—4-sperma rotundata loculicidim dehiscens, seminibus oblongis pendulis, arillo laciniato in testa membranacea adherente e raphi orto ornatis et arillodio carnoso rubro obtectis; embryo intra albumen carnosum orthotropo, cotyledonibus oblongis foliaceis, radícula brevi.

Arbuscula 4—12 metr. alta, foliis oppositis 9—16 cm. longis his 6—9 cm. latis, petiolis .75 cm., cymis 3—9-floris pedunculis 20—30 mm. longis, pedicellis 4 mm. longis, sepalis exterioribus 1.5 mm. interioribus 3 mm. diam., petalis 3 mm. longis latisque, capsulis extus viridibus intus pallidis 1 cm. longis his 1—2 cm. latis seminibus 8 mm. longis his 7 mm. latis testa aurantiaca, arillo pallido, arillodio rubro 6 mm. alto.

Differs from *G. zeylanicum* by its rounded branches, thicker broader leaves with margins entire, by its much smaller flowers with petals flat and not reflexed at the margin, and by its rather shallower arillode. The true arillus is identical in both and the margin of the arillode is similarly lobed.

3. *GLYPTOPETALUM GRIFFITHII*.

In montibus MISHMI; *Griffith* (n. 1996).

Arbuscula? glabra ramulis glabris sub-4-gonis, foliis membranaceis late-virentibus, elliptico-oblongis apice acuminatis basi cuneatis margine parce et minute dentatis, nervis 6—7-paribus subtus prominentibus,

petiolis distinctis; *cymis* axillaribus paucifloris densis pedunculis semel divisis, pedicellis elongatis; *floribus*; *capsula* 1—4-sperma rotundata loculicidim dehiscente; *seminibus* oblongis pendulis basi tantum arillodio carnoso obtectis; *embryone* intra albumen carnosum orthotropo, cotyledonibus oblongis foliaceis, radícula brevi.

Arbuscula? *foliis* oppositis 10—15 cm. longis his 6—9 cm. latis, petiolis 1 cm. longis, *cymis* 3-floris pedunculis 2—4 mm. longis pedicellis 8—9 mm. longis, *sepalis* omnibus 2.5 mm. diam., capsulis extus læte-viridibus intus pallidis 1 cm. longis his 1—2 cm. latis seminibus 8 mm. longis his 7 mm. latis, arillodio 2.5 mm. alto.

This species is well distinguished by its very short cymes with peduncles shorter than the pedicels and by its much shallower arillode. The sepals are all of equal size, the capsules externally are not distinguishable from those of *G. zeylanicum*.

4. *GLYPTOPETALUM SCLEROCARPUM* Kurz. (*F. B. I.* i, 613).
PEGU.

5. *GLYPTOPETALUM GRANDIFLORUM* Beddome. (*F. B. I.* i, 613).
INDIA AUSTRALIS.

JOURNAL
OF THE
ASIATIC SOCIETY OF BENGAL.

Part II.—NATURAL SCIENCE.

No. III.—1891.

IX.—*Additional Uredineae from the Neighbourhood of Simla.*
By A. BARCLAY, M. B., Bengal Medical Service.

[Received 12th June, 1891; Read 1st July, 1891.]

(With Plates IV and V.)

Since the completion of my Descriptive List of the Simla *Uredineae*, published in former volumes of the Journal of this Society I have found, or have been supplied with, 32 other species, which are described in the following pages. Of these 32 species, 4 are *Uromyces* (2 new), 11 *Puccinia* (5 new), 3 *Phragmidia* (all new), 1 *Xenodochus* (new), 2 *Melampsora* (1 new), 5 isolated Accidial Forms (3 probably new), and 6 isolated *Uredo* Forms (5 probably new). Of the 32 species described, therefore, probably 20 are new.

HEMIUROMYCES.

UROMYCES VIGNAE, n. s.

On *Vigna vexillata*, Benth.

I found this plant (on Tara Devi) towards the end of August immensely attacked by a species of *Uromyces*. The leaves were sometimes almost blackened with pustules. These were irregularly scattered over the leaf blade on both surfaces, but more freely over the lower

surface; and it appeared to me probable that they first emerged from this surface, and later from the upper surface, at a more advanced stage. Each pustule was round and convex, and without any paling of tissue around. On a few of the younger leaves I found some paler brown uredo pustules, on both surfaces of the leaf. Except in colour they were similar to the teleutospore pustules.

The *uredospores* are brownish red, oval to round, thin walled, very spiny, and with a few immature teleutospores among them (Pl. V, fig. 19). They measured when fresh and just wetted $26-19 \times 19-18\mu$.

The *teleutospores* are deep brown, oval, very deciduous, with a portion of colourless stalk adherent, with a pale brown shallow mammilla at the free end, a clear nuclear vesicle, a germ pore just under the mammilla, and a smooth surface (Pl. V, fig. 19). The fresh spores just wetted measured $35-27 \times 22-20\mu$. The portion of adherent stalk measured up to 40μ in length. These spores refused to germinate, probably requiring a period of rest. But some uredospores germinated throwing out simple long unbranched tubes.

As this host is nearly related to *Lathyrus sativus*, I at first thought the fungus must be the same as that I have described elsewhere* on that plant, namely, *Uromyces Pisi*, Pers.; but the uredospores of the fungus on *Vigna* are considerably smaller, whilst the teleutospores are larger. It is therefore probable that the species I have here described is different, and I have named it *Uromyces Vignae*.

UROMYCES AGROPYRI, n. s.

On *Agropyrum*, sp.

This grass was collected also by Mr. Lacey at Ralli (Bashahr), 7,000 feet, in October.

On some leaves there were light brown pustules on both leaf surfaces, and these contained uredospores. Other pustules were long, linear and black on the under leaf surface, somewhat resembling pustules of *P. graminis*.

The *uredospores* are round to oval, pale brown, densely warted, with several germ pores (4 to 5), and measured when just wetted $23-20 \times 21-19\mu$.

The *teleutospores* are lightly adherent to their beds, coming off with a small piece of stalk attached. They are light yellowish brown, much thickened at the apex, smooth on the surface, with a clear nuclear vesicle, measuring when just wetted $37-28 \times 18-16\mu$.

Hitherto, so far as I am aware, this genus of grasses was known to

* Journal of Botany, Sept. 1890.

harbour only two species of *Puccinia* (*P. graminis* and *P. coronata*). This is therefore probably a new species.

UROMYCES PULVINATUS, Kalchb. et Cooke.?

On *Euphorbia hypericifolia*, Linn. var. *indica*.

Mr. Lace collected a *Puccinia* on this species of *Euphorbia* in October in a low valley of Bashahr, about 6,000 feet above the sea level. The under-surfaces of the leaves were profusely covered with dark brown circular pustules, usually distinct from one another, but sometimes coalescing. The spores are readily scraped off. These consist of:—*uredospores* which are yellow, thin-walled, spiny, and $20 \times 19\mu$; and *teleutospores*. These are small, brown, thick-walled, single-celled, with a small piece of stalk adhering (sometimes none), warted on the surface, and generally with a clear nuclear vesicle. When just wetted they measure $21-17 \times 18-16\mu$.

Of the several species of *Uromyces* which inhabit species of *Euphorbia* the one I have just described comes nearest *U. pulvinatus*, Kalchb. et Cooke, and *U. Myristica*, B. et B. The former has been described from South Africa, and the latter from North America. In size of spore the Himalayan fungus resembles the former more than the latter; but the spores of *U. pulvinatus* are smooth, whilst those of *U. Myristica* are punctated like mine.

MICRUROMYCES or LEPTUROMYCES.

UROMYCES AMBIENS, Cooke.

On *Buxus sempervirens*, Linn.

Specimens of this fungus were gathered by Mr. Lace in the Bhabar Valley, Bashahr (6,000 feet) in October. The leaves were studded with circular patches 1—4 mm. in diameter. Some of these were orange yellow, and others pale in the centre, but surrounded by a black circle. The centres of all patches contained very prominent papillae (spermatogonia?). All the pustules were hypophyllous. In the case of patches which were not black the central papillae were surrounded by a circular mound covered with epidermis. This mound was a circular covered spore pustule. The spores were orange red, but as they were thick walled and otherwise like immature teleutospores I do not regard them as uredospores. The black circles surrounding the other patches were naked teleutospore beds. The spores from such parts are large, yellowish brown, round to oval, fairly firmly adherent, coming off usually with a short stalk or none, thick walled, showing a distinct pore at the

summit which is not thickened. Their surfaces were smooth, and they measured when just wetted $44-40 \times 36-34\mu$.

This is doubtless *Uromyces andiensis*, but I have no access to the measurements of the spores. Cooke's specimens apparently came from the Himalayas.

HEMIPUCCINIA.

PUCCINIA SORGHI, Schw.

On *Zea Mays*, Linn.

I had long searched in fields of maize for a Uredine, but without success until 1890 when I found it in some fields at Mashobra. Up to this time I was acquainted with the fungus only on *Sorghum vulgare*, on specimens of the plant sent to me for examination from the Poona district. But this is absolutely the first record of its existence on maize in India, so far as I am aware. My specimens were gathered early in October. Pustules were found abundantly on both leaf surfaces, some covered entirely with a scale of epidermis, whilst others were more or less naked. The covered pustules contained uredospores, the naked ones mostly teleutospores, and the perfectly matured open ones, which are inky black, only teleutospores. Some pustules were minute and circular, others long and even linear.

The uredospores are pale brownish red, round to oval, beset with shallow warts or short spines, and measure when just wetted $30-26 \times 26-20\mu$. By applying sulphuric acid I detected 3 germ pores in each spore, arranged around the short equator.

The teleutospores are firmly attached, breaking off with a portion of stalk adhering. They are reddish brown, rounded at both ends, thickened at the free end, constricted at the septum, and apparently smooth on the surface. When just wetted they measured $42-32 \times 18-16\mu$. They refused to germinate immediately after ripening. There were no paraphyses.

This fungus is much more like *Pucc. Sorghi* than that which grows on *Sorghum vulgare*, and which I have described elsewhere.* In the first place the fungus on *Zea* has no paraphyses, and in the second the measurements of both uredo- and teleutospores approximate those given for *Pucc. Sorghi* much more closely. In the publication above alluded to I referred the parasite on *Sorghum* with hesitation to *P. Sorghi*. I am now inclined to think that the latter is a different species. Lastly, even later I found a *Puccinia* on *Pennisetum typhoideum*, Rich. (Bajra)

* Journal of Botany, Sept. 1890.

at Erode in the Madras Presidency, which is undoubtedly the same as that on *Sorghum*. As neither of these fungi (on *Sorghum* and *Pennisetum*) are Himalayan I shall not describe them in detail here; but in order to enforce my argument that the fungus on *Zea* is *P. Sorghi*, whilst that on *Sorghum* and *Pennisetum* is a different species, which I shall call *Puccinia Penniseti*, I subjoin in tabular form their salient characters.

Host.	Uredospores.	Teleutospores.	Paraphyses.	Apical thickening.	Germ pores in uredospores.
<i>Pennisetum</i>	34-30 × 24-21	18-44 × 29-26	present.	none.	2
<i>Sorghum</i> ...	34-30 × 22-20	50-41 × 29-22	present.	none.	4-5
<i>Zea</i> ...	30-26 × 26-20	12-32 × 18-16	none.	present.	3

PUCCINIA ELLISII, De-Toni ?

On *Angelica glauca*, Edgw.

This plant was found by Dr. G. Watt at Fagoo in August, bearing uredo- and teleutospore pustules, both minute, discrete, and hypophyllous; but whilst the former are pale yellow, the latter are dark brown to black. Both kinds of pustule occurred together on the same leaf.

The *uredospores* are round to oval, very pale yellow, very spiny, and when just wetted 30 - 24 × 25 - 20 μ . (Pl. IV, fig. 9).

The *teleutospores* are deep brown, rounded at both ends, slightly constricted at the septum, coming off with little or no stalk adhering, not thickened as a rule at the free end, and tuberculated over both cells (Pl. IV, fig. 9). They are sometimes irregular in shape. They measured when just wetted 38 - 22 × 22 μ .

This is possibly *Pucc. Ellisii*, though both the uredo- and teleutospores of this last mentioned species are larger, the former measuring 35 - 30 × 30 - 24, and the latter 40 - 35 × 25 - 20 μ . It is certainly not *Pucc. Angelicae*, Schum. which shows a tendency to erupt along the nerves, among other differing characters.

PUCCINIA CASTAGNEI, Thüm ?

On *Apium graveolens*, Linn.

I have hitherto confused a fungus on this host with *Pucc. Pimpinellae*, Strauss; but it is certainly distinct.

The *uredospores* are very pale brown, covered sparsely with spines, and with three germ pores, each covered with a hyaline semilunar

thickening, like that described in *Pucc. Prenanthis*. Through each of these a commencing germ tube protrudes, but one only develops fully. They are in pustules mostly hypophyllous, but some few epiphyllous. These uredospores afford another instance of extremely long retained power to germinate. I put some spores scraped off from leaves gathered on the 31st October into water on the 13th June, and found on the following day that many had germinated most freely, although the accompanying teleutospores remained ungerminated.

The *teleutospores* are brownish yellow, very irregular in shape, the septum often oblique, and even perpendicular, slightly constricted at the septum, generally not thickened at the free end, though sometimes slightly so, mostly rounded at both ends, but sometimes with the lower cell narrowing towards the stalk (Pl. V, fig. 18). After lying 24 hours in water these spores measured $41 - 26 \times 24 - 18\mu$.

This is possibly *Pucc. Castagnei*, whose teleutospores are said to be very irregular and to measure $46 - 36 \times 24 - 18\mu$.

Puccinia Eulaliae, n. s.

On *Pollinia japonica*, Haeck.

The leaves of this grass presented dark reddish brown linear pustules, mostly on their under-surfaces. These pustules contained uredo- and teleutospores with numerous capitate paraphyses.

The *uredospores* are oval or pyriform, pale brown, and spiny, and measure when just wetted $30 - 28 \times 21 - 20\mu$.

The *teleutospores* are reddish brown: the upper cell is rounded and not thickened specially anywhere; the lower is broadly wedge-shaped. The surface of the spores is smooth, and they usually have a short piece of stalk adhering. They measure when just wetted $38 - 34 \times 20 - 18\mu$. The paraphyses are numerous, reddish brown, capitate, the heads measuring about 16μ in diameter.

This is probably a new species.

MICROPUCCINIA.

Puccinia excelsa, n. s.

On *Phlomis lamiifolia*, Royle.

I found this plant early in September on the summit of the Huttoo peak bearing *Puccinia* pustules. I next found it, about the same time at Mahasu, a hill close to Simla. The leaves were densely bespattered on the under surfaces with fairly large, round to oval, dark brown, almost black, circular hemispherical pustules, with a smaller less fre-

quent epiphyllous eruption opposite the beds below. The upper surface of attacked leaves is rendered conspicuous by the yellow zones surrounding the pustules. Pustules were also found on the petiole and smaller stems, though not so frequently.

The spores are fairly easily detached, coming off as a rule with no portion of stalk adhering. They are dark brown, somewhat irregular in size and shape, mostly distinctly constricted at the septum, and mostly narrowing towards the free end, where there is a slight pale mammilla or conical thickening (Pl. V, fig. 12). At the base the spore is sometimes rounded, and sometimes narrowed. It is smooth on the surface. The fresh spores examined at once in water measure $40 - 28 \times 18 - 14\mu$.

No uredospores are apparently formed by this species, since I got specimens of teleutospores from the earliest stages of development. The teleutospores would not germinate immediately after ripening.

There is no doubt, I think, that this fungus is distinct from *P. Phlomidis*, Thüm.

LEPTOPUCCINIA.

PUCCINIA USTALIS, Berk. ?

On *Ranunculus hirtellus*, Royle.

I found seedlings of this plant bearing teleutospore pustules on the Mattiana hill on the 4th September. The pustules were quite young, and there was no trace of uredospore. The teleutospore pustules were small, dark, circular, and hypophyllous, with spots of paling on the upper leaf surface. These pustules were confined to the youngest leaves, and were never found on the upper ones. The pustules had a distinct though not pronounced circinate arrangement. The spores are very firmly adherent; they are long, more or less spindle-shaped, pale yellowish brown, much thickened and conical at the free end, well constricted at the septum, and narrowing towards the stalk. The surface is smooth (Pl. V, fig. 13). In scrapings I noticed many empty spore cases, and I presume therefore that the species is a *Leptopuccinia*. When just wetted the spores measured $61 - 46 \times 23 - 8\mu$, the apical thickening being $10 - 4\mu$ in depth. I placed some spores in water on the 10th September, and on the following day found some had germinated. The sporidia are oval or somewhat semilunar, colourless, and $13 \times 6\mu$, in measurement. The sterigmata are four in number, short, conical, and pointed, and together with the whole premycelium colourless.

This is possibly *P. ustalis*, Berk.; but I have no access to the spore measurements.

MICRO or LEPTOPUCCINIA.

PUCCINIA DOLORIS, Speg. ?

On *Erigeron alpinus*, var. *multicaulis*, Wall.

This plant, harbouring a species of *Puccinia*, was gathered by Mr. Lace on the 27th August near the banks of the Sutlej in Bashahr, at an elevation of about 7,000 feet. The under surfaces of the leaves bore numerous large black pustules, irregularly scattered as a rule, but sometimes with several smaller pustules in a circlet around a larger central one. Most pustules were naked, but some were covered over with a scale of epidermis. Though most pustules are hypophyllous some few are on the upper leaf surface.

The spores are readily detached, coming off with only a very small fragment of stalk adhering. They are yellowish brown, rounded at both ends, well constricted at the septum, thickened and broadly conical at the free end, beset externally over both cells with shallow short ridges and tubercles, and measuring when just wetted $42-34 \times 18-16\mu$. The thickening at the free end is usually 6μ in depth. I placed them in water in October, but they did not germinate.

This fungus resembles *Pucc. doloris* as described by De-Toni* in many respects, and I have accordingly named it so. But as *P. doloris* is known only, so far as I am aware, from the Argentine Republic, it is quite possible that the Himalayan species is distinct.

PUCCINIA SAXIFRAGAE-MICRANTHAE, n. s.

On *Saxifraga micrantha*, Edgw.

Mr. Lace collected this plant, bearing a *Puccinia*, in Bashahr at an elevation of 9,500 feet. On the under leaf surfaces were a number of minute, circular, discrete, brown pustules, sometimes very numerous.

The telentospores are readily detached, and many were found to be empty. The species is probably therefore a *Leptopuccinia*. They are pale brown, not thickened anywhere, with a slight apical mammilla, usually slightly constricted at the septum, but sometimes considerably so, and somewhat irregular in size and shape. When just wetted they measure $35-26 \times 14-12\mu$. I placed these spores in water, but none germinated. There were no uredospores in the specimens sent to me.

This species is quite distinct from *Pucc. Saxifragae-ciliatae* mihi. It is also evidently distinct from *P. Saxifragae*, Schlect.

* Saccardo, Sylloge Fungorum.

PUCCINIA CAUDATA, n. s.

On *Stellaria pycnulata*, Edgw.

Dr. G. Watt collected this plant in Narkanda, bearing a *Puccinia*. The teleutospore beds are dark brown, well raised with a circinate tendency, and hypophyllous, with paled patches on the upper leaf surface. The spores are very adherent, more or less spindle-shaped, pale brown, well constricted at the septum, usually much thickened at the apex, with a long piece of adherent stalk (often twice, or a little more, than the whole length of the spore) and measuring $37 - 28 \times 16 - 13\mu$. (Pl. V, fig. 17.).

This is evidently not *P. Arenariae*, Schum. and I have regarded it as a new species.

PUCCINIA CRASSA, n. s.

On *Pimpinella Griffithiana*, Boiss.

This was gathered by Mr. Lace in Ziarat in Afghanistan at 8,000 feet. The teleutospore pustules are dark brown and hypophyllous. The spores are brown, almost rounded at both ends, but diminishing somewhat towards the stalk, of which a small fragment remains adherent. They are slightly constricted at the septum, and slightly thickened at the apex (Pl. V, fig. 16). When just moistened the spores measure $50 - 41 \times 24 - 22\mu$. The epispore has shallow tubercles over both cells. They would not germinate.

This fungus is quite distinct from *P. Pimpinellae*, Strauss which is one of the most common of the *Uredineae* in Simla. Neither are the spores like any of the other species inhabiting *Pimpinella* described by De-Toni* (except perhaps *P. Pimpinella*, St var. *Eryngii*, D. C.) It is I think a new species.

PUCCINIA PULVINATA Rabenh.?

On *Echinops niveus*, Wall.

A *Puccinia* on this host was gathered for me by Dr. G. Watt in Simla. The leaves had numerous minute well raised black pustules all epiphylous. The spores are readily detached from their beds. The plant was in full flower inclining to seed. The spores come off sometimes with a long piece of stalk attached, sometimes with a short. The spore surface is studded with shallow tubercles. They are brown, slightly constricted at the septum, mostly rounded at both ends, the free end slightly thickened. Most spores were found empty, so they

* Saccardo, loc. cit.

presumably germinate at once. They measure when just wetted $52 - 45 \times 24 - 22\mu$.

This is possibly *Puccinia pulvinata*, though the measurements given by De-Toni are greater than mine ($68 - 51 \times 38 - 35\mu$).

PHRAGMIDIUM.

PHRAGMIDIUM LACEIANUM, n. s.

On *Potentilla argyrophylla*, Wall.

I first received specimens of this plant harbouring a parasite from Mr. Lace, who gathered them in Bashahr. Subsequently I found specimens myself at Narkanda, and Dr. J. Murray procured specimens at Sarhan in Kulu.

The *uredo pustules* are hypophyllous, brilliantly orange red, discrete, but often so numerous as to give an appearance of coalescence. Each individual pustule is circular and minute. The *spores* are round to oval or pyriform, bright orange red, spiny, and measuring when just wetted $24 - 20 \times 18 - 16\mu$ (Pl. IV, fig. 3).

The *teleutospore pustules* are also hypophyllous, black, circular, discrete. The *spores* are deep brown, on stalks which narrow gradually away from the spore. They are mostly five-celled; but some are four and some six-celled. The surface is apparently smooth, the free end rounded and slightly thickened, with usually a minute colourless papilla. When just wetted they measure $132 - 94 \times 50 - 41\mu$. I could not detect any germ pores; and they would not germinate immediately after ripening.

The specimens sent by Mr. Lace, and those collected by myself were on the red flowered variety, whilst those collected by Dr. Murray were on the yellow flowered variety. This fact may tend to confirm the view taken in Hooker's Flora of British India that these two are varieties of the same species.

This fungus is, I think, distinct from any of the three noted in Saccardo's Sylloge as inhabiting species of *Potentilla*. It is not *Phr. Fragariastris*, D. C., which has warty teleutospores, $75 - 45\mu$ long and 3-5 celled. It is not *Phr. Potentillae*, Pers. which has smooth teleutospores, $90 \times 26\mu$, and 3-7 celled. And it is not *Phr. Tormentillae*, Fuck. which has spores often bent, $115 \times 28\mu$, 3-8 celled, and light brown.

PHRAGMIDIUM NEPALENSE, n. s.

On *Potentilla nepalensis*, Hook.

I found this host in September attacked with a *Phragmidium* at Mattiana, some miles towards the interior from Simla. The host at

that time was in flower and held both uredo- and teleutospore pustules. Both pustules were mainly hypophyllous, but a few (especially uredo pustules) were epiphyllous.

The uredo pustules are very brilliantly orange red and circular, but were often so closely aggregated that they ran together. The spores are round or oval, or more or less irregular in shape, and bright orange red (Pl. IV, fig. 2). They varied much in size, $28 - 22 \times 22 - 17\mu$. There were no paraphyses.

The teleutospore pustules are much smaller and appear like minute black dots scattered about irregularly. The spores are very dark brown, generally constricted at the septa, apparently with 2-4 pores to each cell, 2-5 celled, but usually 4 celled, with a small and inconspicuous mammilla at the free end, looking more like a slight general thickening of the epispore (Pl. IV, fig. 2). The stalk is long, thin, and inflated at some distance from the attachment to the spore, the inflation containing orange red matter. Normal 4 celled spores measured when just wetted $68 - 66 \times 26\mu$ and a 3 celled spore $54 \times 24\mu$.

Of the three species of *Phragmidium* on species of *Potentilla* described by De-Toni* only one, namely, *Phr. Fragariastris* has so few cells to each spore, and the one I have just described is, I think, not identical with it. I have therefore named it *Phragmidium nepalense*.

PHRAGMIDIUM OCTOLOCLARE, n. s.

On *Rubus rosaefolius*, Smith.

The general appearance of the teleutospore stage of this fungus (the only one I know it in) is very like that of *Phr. Barclayi*, Dietel,† with somewhat large circular pulverulent hypophyllous pustules.

The teleutospores are dark brown 7-9 celled usually, but mostly 8-celled. The surface is distinctly and coarsely tuberculated, and at the free end there is sometimes a minute colourless papilla, but often none. The stalk is long, swells up somewhat in water, is inflated away from the spore, and this lower part of the stalk swells, more than the upper part adjoining the spore (Pl. IV, fig. 1). The stalk is not unlike that of *Phr. Barclayi*. After lying 24 hours in water the gelatinous sheath shrinks upwards towards the spore, leaving a central axis with orange red swellings at the ends. The spore is constricted at each septum to a slight degree. Each cell of the spore is more flattened from above downwards than in the last mentioned species, the vertical depth of each cell being 10μ against 14μ in *Phr. Barclayi*. The spores (taking

* Saccardo, *loc. cit.*

† I incorrectly regarded it as *Phr. Rubi*, Pers., in my Descriptive List.

those which have 7-9 cells) measure when recently wetted $130 - 94 \times 23 - 25\mu$ but exceptionally a spore of only two cells may be found, measuring $44 \times 24\mu$. I have not observed the germination of these spores; but some spores collected in autumn and put into water did not germinate, and from this I conclude that they must rest. On the other hand some of these spores showed a cell here and there empty; so apparently under special conditions they may also germinate at once.

I am inclined to regard this fungus as distinct from *Phr. Barclayi* and *Phr. quinqueloculare*, mili. If it be identical with either, it is with *Phr. Barclayi*; but the spores of the latter are usually 6-celled with smooth surface (or with very shallow inconspicuous warts) somewhat less in length and greater in diameter. I am also unable to match it with any of the six species described by De-Toni* on species of *Rubus*. I propose naming it *Phragmidium octoloculare*.

XENODOCHUS.

XENODOCHUS CLARKIANUS, n. s.

On *Astilbe rivularis*, Ham.

This fungus was collected by Dr. Clark in the Cheog forest about the beginning of August. On one specimen there were large irregular aecidial patches, especially on the stem, accompanied by hypertrophy, just like that caused by *Phragmidium subcorticium* on *Rosa moschata*. These aecidia were exceedingly brilliantly orange red: they occurred also on leaves. The spores are very bright orange red, in rows, squarish or oblong, densely warted, thickened a little at one end (seen best in empty spores) (Pl. IV, fig. 5). They measure $31 - 26 \times 26 - 22\mu$. There were no paraphyses.

But much more numerous were orange waxy looking beds, which to the naked eye resembled *Coleosporium* beds. These in many places simply covered the under leaf surface. I noticed that in some parts these orange waxy beds were gradually changing, and at others had changed into black beds, naked, and under a field lens looking like *Puccinia* beds. The spores from the latter are scraped off with difficulty, and when examined under the microscope presented characters most like those I have read described as *Xenodochus* spores. They have characters, as far as I am able to judge, intermediate between this genus and *Phragmidium*. The spores were in rows, usually 5 to 6 in each row, pale brown, rows being sometimes transversally septate, at others irregularly in various obliquities. The rows of spores were some-

* Saccardo, loc. cit.

times closely amalgamated with rows on either side. Each spore of a regular row measured about $16 \times 13\mu$ (Pl. IV, fig. 4). I placed these spores in water, but they refused to germinate.

This is apparently a new species of *Xenodochus*. It differs considerably from *X. carbonarius*, Schlect. in which the aecidiospores are $28 - 16 \times 20 - 15\mu$ and among which there are paraphyses. Moreover, in the last named species the teleutospore rows have 10 to 20 loculi.

MELAMPSORA.

MELAMPSORA CILIATA, n. s.

On *Populus ciliata*, Wall.

This host is abundant in Simla, and many are occasionally attacked by a species of *Melampsora*. The uredo stage makes its appearance in August or September.

The uredo pustules are minute and discrete, but often very abundant; they are light yellow, and entirely hypophyllous, with scattered yellow dots on the upper leaf surface. The spores are pale orange red, mostly oval, pretty densely covered with spines, thick walled, and measure when fresh $30 - 21 \times 22 - 20\mu$ (Pl. V, fig. 15). They are borne singly on stalks, and among them are some capitate paraphyses, with heads very distinctly, and often greatly thickened at the free end (Pl. V, fig. 15).

The teleutospore beds are at first orange red, but become brown later. They are entirely hypophyllous. The spores are in compact beds and each spore measures $34 - 30 \times 9 - 8\mu$.

This fungus is, I think, different from any of the three described by De-Toni. This will be apparent when the characters are shown tabularly.

	Uredospores.	Teleutospores.	Paraphyses.	Remarks.
<i>M. Tremulae</i> , Tul.	24 - 15 × 18 - 13	55 - 45 × 12 - 11	Clavate, 9 - 15.	
<i>M. aecidioides</i> , D. C.	24 - 17 × 17 - 15	?	Clavate.	
<i>M. populina</i> , Jacq.	38 - 28 × 20 - 15	45 - 40 × 13	Capitate, 20-17.	epiphyllous.
<i>M. ciliata</i> , n. s.	30 - 21 × 22 - 20	34 - 30 × 9 - 8	Capitate.	hypophyllous.

MELAMPSORA AECIDIODES, D. C. ?

On *Populus alba*, Lin.

Mr. Lace sent me specimens of the leaves of this plant collected at an elevation of 8,500 feet on the 30th August. The under surfaces

were almost wholly covered with brilliant orange red pustules, minute and discrete, though from their enormous numbers they at first sight appeared to be coalescing. The upper leaf surface is profusely flecked with yellow discoloured spots. The spores are orange red, thick walled, spiny, and measure when just wetted $24 - 21 \times 18 - 16\mu$. There were no teliospores.

It is impossible from the uredo spores alone to determine the exact position of this fungus. Possibly it is *M. accidioides* which occurs also on *Populus alba*. The uredospore measurements coincide very closely with those I have just described, but *M. accidioides* has paraphyses whilst the Himalayan form has not.

ISOLATED AECIDIA.

AECIDIUM CUNNINGHAMIANUM, n. s.

On *Cotoneaster bacillaris*, Wall.

I found the leaves of this plant bearing several characteristic *Roestelia* patches first on the Mahasu hill by the road side; then fairly abundantly in Narkanda; and lastly, scarcely in Mashobra. These were found at the end of August. In all these localities, especially Narkanda and Mashobra, I did not see a single *Cupressus* tree, and this inclines me to regard the fungus as distinct from *Gymnosporangium Cunninghamianum*, mihi. The leaf patches were red above, with long filiform peridia on the lower surface, about 3-5 mm. in length. From one to five such patches were found on a single leaf. On superficial examination this fungus looks different from *Gymnosporangium Cunninghamianum*, although the peridium bursts in the same way, namely, by irregular slits on the tube sides. With a field lens numerous spermogonia could be seen on the upper leaf surface.

The aecidiospores are yellowish brown, round or oval, densely beset with minute and very shallow warts, and with apparently numerous germ pores (the addition of sulphuric acid discloses eight pores). These spores measure $28 - 26 \times 28 - 24\mu$ (Pl. IV, fig. 7). The peridial cells are paler in colour than the aecidiospores, elongated, separating readily from one another laterally, very spiny (not ridgy), and measure from $60 - 58 \times 26 - 24\mu$ (Pl. IV, fig. 7). The aecidiospores would not germinate in water.

Had it not been for the absence, as far as I could see, of *Cupressus* trees in the neighbourhood of these aecidial patches, I should have been disposed on the whole to regard this fungus as *G. Cunninghamianum*; and, indeed, this identity is still quite possible, since it is by no means easy to be certain that no *Cypress* tree exists in the forests in those

regions. In support of the view of identity are the manner of dehiscence of the peridium, and the closely corresponding sizes of the accidiospores and the peridial cells (the former in the case of *G. Cunninghamianum* being on an average $28.6 \times 24.6\mu$, and the latter $70 \times 22\mu$). I should note also that the peridial tubes of the accidium on *Cotoneaster* are somewhat longer than those on *Pyrus* (in the latter they are 1 to 2 mm.)

The only *Aecidium* known on *Cotoneaster* is *Aec. Mespili*, D. C.; but the accidiospores of this species are $24 - 19\mu$ in diameter, and I do not think the Simla species can be the same. Until more is known of its life history I propose naming it *Aec. Cunninghamianum*, believing that it will probably prove to be identical with *Gymnosporangium Cunninghamianum*, mihi.

ÆCIDIDIUM MORI, n. s.

On *Ficus palmata*, Forsk.

I found this plant first attacked in a valley to the north of Mashobra, and subsequently in Simla itself during October. The leaves are often densely covered with an orange red eruption. In some cases the whole of the lower leaf surface was a mass of these bright pustules. But although the eruption is mainly hypophyllous it is also largely epiphyllous. With a field lens it is difficult to determine the nature of the fungus. Each spore heap is very minute, but shreds of white tissue are seen about them. Under the microscope however, all doubt ceases, for there are very characteristic peridial cells, and the fungus is consequently an *Aecidium*. But it is a very remarkable one in having so small and inconspicuous a peridium. To the naked eye the fungus resembles a *Uredo* form. The spore beds are scattered irregularly all over the leaf surface; but sometimes on somewhat swollen weals on the petiole and midrib. The accidiospores are very brilliant orange red bodies, round to oval, and apparently smooth on the surface. When just wetted they measure $17 - 14 \times 16 - 14\mu$. The peridial cells are colourless and more delicate than usual. They are mostly six-sided and spiny or tuberculated. They measure $22 \times 18 - 13\mu$. I placed some of the accidiospores in water on the 17th of October, and a few were found to have germinated on the following day. The germ tube is long, simple, and unbranched.

Saccardo mentions a *Puccinia sepulta*, B. et C. on the leaves of a species of *Ficus* from Nicaragua and two *Uredo* forms (*U. Fici*, Cast. and *U. ficicola*, Speg.); but even supposing these *Uredo* forms are really *Aecidia*, the spores of the former are much too large, and those of the latter considerably larger than the Simla fungus to permit of their being considered identical. The Simla species is probably therefore new.

This fungus is identical with that I have described as *Caeoma Mori*; and as I have subsequently found that the peridium is distinctly present though incoherent, I feel disposed to relinquish the name *Caeoma Mori* and to substitute *Aecidium Mori*.

AECIDIUM FLAVESCENS, n. s.

On *Senecio rufo-nervis*, D. C.

I found this plant in August bearing an *Aecidium* in the Mashobra woods. The aecidial patches are indicated conspicuously by brown patches with yellow irregular halos around them on the upper leaf surface. On the under surface the peridia are indistinctly seen against the white tomented natural leaf surface, as they are pale in colour. The peridia are densely aggregated together, and always on the under leaf surface. The tubes are short and open stellately. I counted from one to thirty-two aecidial patches on single leaves. The patches varied from a half to 2 c.m. in diameter. The aecidiospores are oval or round, pale orange red, measuring when well moistened $22 - 20 \times 16 - 14\mu$. The peridial cells are almost isodiametric, very rugose on the surface, with short ridges and spines, and measuring $28 - 22 \times 22 - 20\mu$. (Pl. IV, fig. 8).

Later in the season leaves in the same locality bore uredospores, and still later teleutospores in the form of *Coleosporium*. I have unfortunately had no opportunity of examining these forms.

Two species of *Aecidia* on species of *Senecio* are noted by Saccardo, *Aec. Huallatinum*, Speg., and *Aec. sclerothecium*, Speg.; but the spores of the Simla species are much smaller than those of either of these two species.

AECIDIUM AQUILEGIE, Pers. ?

On *Aquilegia vulgaris*, Linn.

Mr. Lace collected this plant on the 9th August, 1890 at an elevation of 10,000 feet bearing an *Aecidium*. The peridia are hypophyllous, forming a patch of yellow below with a paled area above on which spermogonia may be seen with a field lens. The peridia open by a clean regular margin. The peridial cells are angular, almost isodiametric, five to six-sided, spiny and ridgy, and about $26 \times 20\mu$. The aecidiospores are round or faceted when just scraped off, tuberculated, and $24 - 18 \times 18 - 12\mu$.

This fungus is very possibly *Aec. Aquilegiae*, Pers. though the spores in the latter are said to be larger ($30 - 16 \times 20 - 14\mu$.) and perhaps more spiny.

AECIDIUM ORBICULARE, n. s.

On *Clematis grata*, Wall." *orientalis*, Linn." *puberula*, H. f. and T.

This *Aecidium* was sent to me first by Mr. Lacey and Dr. Watt, and subsequently I found it myself at Mattiaiu on *Clematis grata*.

Accidial patches were very numerous on stems, petioles, and leaves, causing considerable hypertrophy of stems and petioles, especially on *Clematis grata*. On leaves the peridia were markedly circinate in arrangement, and all were hypophyllous. The peridial tubes were long and cylindrical, and opened at the summit with an almost clean margin, *i. e.*, very minutely serrated. They measured about 0.75 m.m. in length. The spores are bright orange red, densely beset with shallow warts, round to oval or angular, 25 - 20 × 20 - 14 (*Clematis grata*) 21 - 16 × 16 - 15 (*C. orientale*) 22 - 19 × 19 - 15 μ . (*C. puberula*).

The peridial cells were in all cases about 26 - 20 μ . in diameter, four to six-sided, with bright orange red matter in their centres, and spiny, (Pl IV, fig. 6).

This may possibly be the New Zealand *Aec. otagense*, Linds.; but the only description available to me is the very brief one by De-Toni, from which it is impossible to decide, since no spore measurements are given. The only other *Aecidium* described by De-Toni on *Clematis* is *Aec. Clematidis*; but the description of the peridium is unlike that I have described. The Himalayan species is therefore, I think, distinct. It is possible that this *Aecidium* is related to *Puccinia Wattiana*, which occurs plentifully on *Clematis Gauriana* and *C. grata*; but in the absence of experimental proof it is impossible to determine.

ISOLATED UREDO FORMS.

UREDO COLEBROOKIAE, n. s.

On *Colebrookia oppositifolia*, Smith.

This fungus was collected by Dr. Watt near Suni in October. The under surfaces of the leaves were almost uniformly orange red in colour from innumerable orange red pustules. These pustules are really distinct, but appear to be coalescing from their great numbers, and from the hairy nature of the leaf surface, in which the spores get entangled. Dr. Watt informed me that showers of red dust fell from the leaves as he picked them. The spores are pale orange red, oval, very spiny, measuring when just wetted 28 - 20 × 20 - 17 μ . These spores were put into water some few days after collection, but they did not germinate freely; one or two did so, however, throwing out long simple sinuous tubes.

UREDO ICHNOCARPI, n. s.

On *Ichnocarpus frutescens*, Br.

This fungus was also collected by Dr. Watt in the same neighbourhood, and at the same time. The leaves bore a few scattered isolated, bright orange red pustules on the under leaf surface. The spores were readily scraped off. They are bright orange red, oval, studded with large coarse warts or spines, and measure when just wetted $24 - 20 \times 16 - 21\mu$. Even in empty spores I could not detect any germ pores. Among the scraped off uredospores I saw some immature single celled stalked spores which I imagine are teleutospores (*Uromyces*); but they were too immature to allow of decision. They were colourless at this early stage.

An *Aecidium Apocyni*, Schwein. is known to occur in Carolina, Kansas, and Illinois; but it is impossible as yet to say whether it is in any way related to the Himalayan fungus: probably it is not.

UREDO IPOMAEAE, n. s.

On *Ipomaea hederacea*, Jacq.

This plant was collected by Dr. Watt near Sairi in September with numerous, white, irregularly shaped covered pustules on the under leaf surface. On examining the pustules they were found to contain an abundance of white powdery spores. The spores are colourless, round or squarish, apparently quite smooth on the surface, and measuring when just wetted $20 - 17 \times 16 - 14\mu$. Although I have placed this fungus here among *Uredo* forms I should note that it looks much like the *Aecidium* of a *Phragmidium*, differing only in not being coloured.

UREDO PILEAE, n. s.

On *Pilea trinervia*, Wt.

I found a few of these plants in the woods at Mashobra early in October, attacked by a *Uredo*. The pustules were minute, pale yellow, and scattered on the under leaf surface. The spores are oval, very pale yellow, studded with prominent spines, and measured when just wetted $25 - 20 \times 20 - 14\mu$. I could find no trace of any teleutospore form up to the middle of October.

UREDO EHRETIAE, n. s.

On *Ehretia serrata*, Roxb.

This fungus was collected by Mr. J. S. Gamble on the banks of the Tous river near Chakrata. The leaves are attacked by a Uredinous

fungus of somewhat uncertain nature. Some leaves had circular spots varying in diameter from 1 to 8 m.m., whilst others had large hypertrophies of the petiole as it enters the lamina, and these were uniformly covered with bright orange red pulverulent spores. Transverse sections through the fungus and leaf show that the spores are not borne separately on stalks, but from a cup-like depression like the pit of an aecidium without any peridium. There were also numerous superficial spermogonia. The spores are orange red, oval or pear-shaped, very spiny, decidedly thickened at the free end (remining one of the aecidio- and uredospores of *Puccinia Prainiana*). When just wetted they measure $38 - 30 \times 22 - 20\mu$. They become detached without any portion of the stalk adhering. The spores are given off from both surfaces of the leaves. By applying nitric acid I saw that each spore had two germ pores.

This is the first member of the *Boragineae* which I have seen attacked in India by a Uredine. As *Puccinia Rubigo-vera*, or some variety or allied species, is undoubtedly the most prevalent and destructive rust on wheat, barley, and oats in India, I had long looked for some associated form on a Boragineous host, and this not only by personal search, but also by correspondence. I am afraid, however, that this particular Uredine cannot be the associated form I have been looking for, although it is just possible that it is. For although the spores are given off like uredospores, the cup-shaped depressions in which they are formed, the presence of spermogonia, and the hypertrophy of the host's tissue all render it possible that we have here an anomalous *Aecidium*. This question will have to be tested by experiment.

UREDIO AGRIMONIAE, D. C.

On *Agrimonia Eupatorium*, Linn.

This plant is frequently found attacked with a Uredo in these regions; but I have never seen a teleutosporic stage. The Uredo pustules are hypophyllous, very numerous, and minute. The spores are orange yellow, oval, warty or spiny, and measure $22 - 17 \times 14 - 13\mu$, when just wetted.

This is no doubt the widely distributed *Uredo Agrimoniae*.

EXPLANATION OF PLATES.

Plate IV.

1. Telentosporic stage of *Phr. octolocularis*, $\times 220$.
2. Telentosporic and uredosporic stages of *Phr. nepalense*, former $\times 220$, and latter $\times 350$.

3. Telento- and uredospores of *Phr. Laccianum*, $\times 220$ and $\times 350$.
4. Telentosporo of *Xenodochus Clarkianum*, $\times 350$.
5. Accidiospore of the same $\times 350$.
6. Peridial cells of *Accidium orbiculare*, $\times 350$.
7. Peridial cells and accidiospores of *Acc. Cunninghamianum*, a $\times 350$,
b $\times 220$.
8. Peridial cells of *Acc. flavescens*, $\times 350$.
9. Telento- and uredospores of *Pucc. Ellisii*, $\times 350$.
- *10. Telentosporo of *Pucc. Wattiana*, $\times 350$.

Plate V.

11. Telento- and accidiospores and promycelium of *Phr. quinqueloculare*,
 $\times 350$.
12. Telentosporos of *Pucc. excelsa*, $\times 350$.
13. Telentosporos of *Pucc. occulta*, $\times 350$.
- *14. Telentosporos of *Pucc. McIntirianus*, $\times 350$.
15. Uredospores and paraphyses of *Melampsora ciliata*, $\times 350$.
16. Telentosporos of *Pucc. crassa*, $\times 350$.
17. Telentosporos of *Pucc. caudata*, $\times 350$.
18. Telentosporos of *Pucc. Castagnei*, $\times 350$.
19. Telento- and uredospores of *Uromyces Vignae*, $\times 350$.

X.—Notes on the Collection of Snakes in the Indian Museum with descriptions of several new species.—By W. L. SCLATER, M. A., Deputy Superintendent of the Indian Museum.

[Received 1st August, 1891; read 5th August, 1891.]

(With Plate VI.)

The following notes were drawn up while critically examining and rearranging the collection of Snakes in the Indian Museum; the collection is a fairly large one though there are many Indian species still unrepresented, and I should be very glad if any one would assist me to fill up the gaps especially in the Southern Indian and Ceylonese forms in which the Museum is specially deficient.

The number of Snakes described by Mr. Boulenger in his work on the Reptiles of the Indian Empire and Ceylon amounts in all to 264, of this number the Indian Museum possesses examples of 196, so that no less than 68 are still wanting to complete the Indian Museum Collection; of the 68 deficiencies, however, 22 at least of the species have only been got once and are represented in the British Museum or elsewhere usually by a single specimen.

* For descriptions of these, see J. A. S. B., Vol. LIX, Pt. II, No. 2, 1890.

To the 264 snakes given in Mr. Boulenger's work I have now eleven to add, of which five are new, and described here for the first time and six are exotic species now recorded from the Indian Empire for the first time, so that the total number of Indian Snakes is now raised to 275.

The Indian Museum possesses a fair number of types, but unfortunately several which I believe should be here are no longer to be found in the collection, such is the case with *Typhlops axlamaniensis*, Stoliczka, *Calamaria catenata*, Blyth, and several others.

The following is a list of the species, twenty-one in number, of which the types are still in the Museum.

<i>Typhlops theobaldianus</i> , Stol.	<i>Coluber prasinus</i> , Bly.
<i>Trachischium fuscum</i> , (Bly.)	<i>Pseudoxenodon macrops</i> , (Bly.)
" <i>tenuiceps</i> , (Bly.)	<i>Tropidonotus platyceps</i> , Bly.
<i>Blythia reticulata</i> , (Bly.)	" <i>nigrocinctus</i> , Bly.
<i>Lycodon gammiei</i> , (Blanf.)	<i>Dipsas multifasciatus</i> , Bly.
<i>Hydrophobus davisoni</i> , (Blanf.)	<i>Hypsirhina blanfordi</i> , Boul.*
<i>Pseudocyclophis bicolor</i> , (Bly.)	<i>Distira tuberculata</i> , (Anders.)
<i>Ablabes scriptus</i> , Theob.	<i>Amblycephalus modestus</i> , (Theob.)
" <i>nicobarensis</i> , Stol.	" <i>macularius</i> , (Bly.)
<i>Zamenis ladaccensis</i> , Anders.	<i>Trimeresurus cantor</i> , Bly.
<i>Zaocys nigromarginatus</i> , (Bly.)	

To these must be added the types of the five new species described below.

<i>Ablabes stoliczkae</i> .	<i>Tropidonotus pealii</i> .
<i>Simotes woodmasoni</i> .	<i>Tropidonotus nicobaricus</i> .
<i>Zaocys tenasserimensis</i> .	

There are also in the Indian Museum the types of the following species found on the borders of the Indian Empire but not yet recorded from within its limits.

<i>Typhlops persicus</i> , Blanf.	<i>Pseudocyclophis persicus</i> , (Anders.)
<i>Calamaria stalknechti</i> , Stol.	<i>Dipsas rhinopoma</i> , Blanf.

The total number of types therefore in the Indian Museum is thirty.

It is only due to Mr. Boulenger to add that it is entirely owing to his useful and excellent manual on Indian Reptiles in the Indian Fauna series that I have been able to work out and rearrange the Snakes in the Indian Museum, so much has been done since Günther's Reptiles of British India appeared and that scattered over so many various separate works and periodicals.

The Snakes in the Museum are now arranged in accordance with

* Originally described by Blanford as *H. maculosa*, name changed by Boulenger.

Mr. Boulenger's work, and a complete list of them will shortly be published, in the meantime the following notes may be found useful.

TYPHLOPS DIARDI, Schleg., Boulenger, p. 238.

TYPHLOPS BOTHRIORHYNCHUS, Günth., Boulenger, p. 239.

I have found it difficult to separate these two species, in fact the only distinction pointed out by Boulenger is the presence of the pits below the nostrils, these I have found vary considerably in depth and conspicuousness; all the specimens of *T. bothriorhynchus* in the Indian Museum were procured in Assam where *T. diardi* also seems to be most abundant; it would therefore seem possible that this pit might be a sexual or seasonal mark, and that these two species are really one and the same.

TYPHLOPS THEOBALDIANUS, Stol., Boulenger, p. 240.

Of this species, which seems to be unrepresented in the British Museum Collection, the Indian Museum possesses two examples, one the type, the history of which is unknown, and a second procured by Capt. J. Butler at Samagooting in the Naga hills.

TYPHLOPS ACUTUS, (Dum. et Bibr.), Boulenger, p. 241.

This species is not confined to Southern India, it extends to Bengal, and the Indian Museum contains specimens from Sipri in Gwalior, Chybassa in Chota Nagpur, Calcutta and Krishnaghur.

PYTHON MOLURUS, (Linn.), Boulenger, p. 246.

Boulenger does not mention whether the present species extends to China, Günther also expresses a doubt on the subject, the matter may now be settled as the Indian Museum possesses an undoubted specimen of *P. molurus* procured by Surgeon-Major Hungerford at Haiphong (Hiaofung?) near Shanghai.

SILYBURA NIGRA, Beddome, Boulenger, p. 263.

Silybura wood-masoni, Theobald (Cat. Rept. Brit. Ind., p. 135) the type of which is in the Indian Museum, seem rather referable to *Silybura nigra* than to *S. pulneyensis*.

SILYBURA BREVIS, Günth., Boulenger, p. 268.

This species extends northwards to the Ganjam hills, whence the Museum possesses a specimen presented by Col. R. H. Beddome.

CALAMARIA CATENATA, Blyth, Boulenger, p. 282.

The type of this species is no longer in the Museum, it had apparently disappeared before the collections of the Asiatic Society were transferred to the Indian Museum, as is mentioned by Theobald in his Catalogue of Reptiles in the Asiatic Society's Museum so that unless the species is rediscovered in Assam, nothing further can be known about it beyond what is contained in Blyth's rather meagre description.

CALAMARIA SUMATRANA, Edeling, Natur. Tijdsch. Ned. Indië xxxi, p. 379, (1870).

A specimen of this apparently rather rare snake from Singapore presented to the Museum by Mr. Davison, was kindly identified for me by Mr. Boulenger; it has hitherto been known from Sumatra only.

TRACHISCHIUM GUENTHERI, Boulenger, p. 285.

There are three snakes in the Indian Museum which seem referable to this species, one from Katmandu in Nepal and two labeled "Allahabad J. Cockburn," the latter probably came from Naini Tal as there are other Himalayan snakes in the collection presented by Mr. Cockburn labelled Allahabad; it is probable therefore that this is the more western representative of *Trachischium fuscum* from which it differs merely in colouration and the number of ventral shields.

TRACHISCHIUM RUBRIVENTER, (Jerdon), Boulenger, p. 286.

The type of this species does not seem to be in the British Museum as the species is "unknown to Mr. Boulenger," neither have I been able to find it among the specimens in the Indian Museum, unless therefore the species is rediscovered, nothing further can be known about it.

LYCODON STRIATUS, (Shaw), Boulenger, p. 292.

LYCODON TRAVANCORICUS, (Bedd.), Boulenger, p. 293.

There are no specimens of *L. striatus* in the Museum from Southern India, the localities from which there are specimens are the hills below Simla, Jemper in Sind, Lahore, Agra, Ajmere, Rajputana, and the Ganjam district; the species is recorded from the Anamalai hills by Günther, but this was before *Lycodon travancoricus* which resembles *Lycodon striatus* very strongly had been discriminated; of this species (*S. travancoricus*), the Museum possesses examples from the Nilgiri hills, the South Arcot district and Tinnevely hills, it therefore appears probable that it entirely replaces *S. striatus* in the southern part of the Indian Peninsula.

LYCODON FASCIATUS, (Anders.), Boulenger, p. 295.

The Indian Museum possesses two specimens of this rather rare species, one from Shillong in the Khasia hills collected by the late Major Cock, and one from Tezpur obtained by Col. Godwin-Austen; the species has been hitherto recorded only from the Kakhyen hills and Western Yunnan.

POLYDONTOPHIS BISTRIGATUS, (Günth.), Boulenger, p. 304.

The geographical area of this species can be extended to the Nicobars whence there is an example procured by Mr. de Roepstorff.

ABLABES STOLICZKAE, sp. nov. Pl. VI, fig. 1.

Rostral shield broader than deep, the part visible from above about two-thirds the length of the internasals; internasals shorter than the prefrontals and somewhat triangular, frontal hardly as long as its distance from the end of the snout, and a good deal shorter than the parietals; nostrils rounded, about the middle of the length of an undivided shield which is quite three times as large as the loreal; this latter is very small and squarish; one preocular not reaching the upper part of the head, two postocular both in contact with the parietals; eye of moderate size, about half the length of the snout; temporals 1 + 2; upper labials eight, fourth and fifth entering the eye; four lower labials in contact with the anterior chin shields which are about as long as the posterior. Scales smooth, in fifteen rows. Ventrals 153-4, subcaudals 116-9. Anal divided. Colour light olive-brown above, lighter olive yellow below, the two colours separated in front by a conspicuous broad longitudinal black streak, extending from just in front of the eye back along the neck for an inch or so.

There are two specimens of this species in the Museum. One procured at Samagooting in the Naga hills of Assam by Capt. J Butler, the other to which no locality is attached but which was received from Dr. F. Stoliczka of the Geological Survey.

This species differs from all the Indian species of *Ablabes* described by Boulenger, except *Ablabes calamaria*, in having a single nasal shield; in all the other species of the genus the nasal shield is divided or semi-divided.

From *A. calamaria* to which it is most nearly allied, it differs in having a loreal distinct from the nasal shield, in having both postoculars in contact with the parietal and in the much larger number of subcaudals, 116-9 against 64-76 in *A. calamaria*.

ABLABES DORIAE, (Boul.), Boulenger, p. 306.

The Indian Museum is indebted to Mr. R. D. Oldham of the Geological Survey for a specimen of this rather rare snake which he procured in Manipur.

ABLABES COLLARIS, (Ménétries), Blanford, Persia, p. 405.

Two snakes collected by Dr. Anderson in Palestine near Lake Galilee seem referable to this species; they agree very well with the description given by Blanford (*l. c.*) of a Persian specimen in the Genoa Museum except that in the Palestine specimens, the posterior chin shields are slightly smaller than the anterior ones, whereas in the Persian specimen the chin shields are said to be equal in size; the allied species *Ablabes modestus* has been recorded from Palestine, but not so far as I am aware the present one.

SIMOTES CYCLURUS, (Cantor), Boulenger, p. 311.

The type specimens of *Simotes obscurus* and *Simotes crassus* are undoubtedly both faded specimens of this species, as was surmised by Boulenger (*l. c.*); with regard to some of the specimens in the Museum, it is difficult to say, whether they should be referred to this species or *Simotes albocinctus*, since the number of anterior temporals and labials entering the eye seem to vary somewhat, so that some of the examples are coloured like one species, but scaled like the other, possibly this may be due to hybridism.

Judging from the specimens of these two snakes in the Indian Museum, it would appear that *S. cyclurus* was an inhabitant of higher lands than *S. albocinctus*.

SIMOTES PURPURASCENS, (Schleg.), Boulenger, P. Z. S. 1890, p. 34.

This species with which according to Boulenger (*l. c.*) *Simotes trinotatus*, Dum. et Bibr., *S. labuanensis*, Günth., *S. catenifer*, Stol. and *S. dennysi*, Blanf., are conspecific, may now be included among the snakes of the Indian Empire, since a specimen was procured by Dr. Anderson from Tavoy in Tenasserim and is now in the Museum.

SIMOTES WOOD-MASONI, sp. nov. (Plate VI, fig. 2.)

Nasal divided; portion of the rostral seen from above a little shorter than its distance from the frontal; suture between the internasals shorter than that between the prefrontals; frontal longer than its distance from the end of the snout, as long as the parietals; loreal small, longer than deep; one preocular, one subocular separating the third labial from the

eye margin, two postoculars; temporals 1 + 2; six upper labials, the fourth alone entering the eye, the fifth the largest; four lower labials in contact with the anterior chin shields; posterior chin shields small, less than half the size of the anterior. Scales in 17 rows. Ventrals strongly angulate, 180-6. Anal undivided. Subcaudals 57.

Colour; adult, above brick reddish with traces of a dorsal and three lateral narrow lighter longitudinal lines; below dusky reddish with a light longitudinal line on either side at the angle of the ventrals; head with a dark median longitudinal mark extending back from the anterior end of the frontal to the nape where it bifurcates into the ground colour of the back, an oblique dark streak across the anterior nasal and the three anterior labials; a second oblique streak from the 4th and 5th labials through the eye above which it bifurcates and meets its fellow from the opposite side, a third oblique streak across the parietals and the sides of the neck. In the young the markings are much more conspicuous, the lighter colour being yellow and contrasting strongly with the darker brown of the markings.

This species is most nearly allied to *Simotes violaceus* and *S. octolineatus*; from the former species it differs in the reduced number of its labials, of which only the fourth enters the eye, and also in colour.

From *S. octolineatus* it differs in having a subocular which enters between the third labial from the eye and in possessing only a single anterior temporal; in colouration, however, especially of the young, the two species much resemble one another.

There are two examples of this species in the Indian Museum, from which this description was drawn up, one from the Andamans collected and presented by Mr. Wood-Mason, and one from the Nicobars where it was procured by the late Mr. F. A. de Roepstorff.

SIMOTES THEOBALDI, Günth., Boulenger, p. 318.

The Indian Museum contains four examples of this rather uncommon species, from Mandalay (Anderson), from Meiktalla (Collett), and from Mergui.

SIMOTES PLANICEPS, Boulenger, p. 316.

An example of this snake was procured by Dr. Anderson during one of his two expeditions to Yunnan and Upper Burma, which had apparently never been named or described; unfortunately the specimen is without locality, though it no doubt comes from Upper Burma; the species has hitherto been known from a single specimen only, procured by Fea at Minbla in Burma.

OLIGODON DORSALIS, (Gray), Boulenger, p. 318.

The range of this species may be extended from the Khasia hills south to the Naga and Chittagong hills whence the Indian Museum possesses examples procured by Capt. J. Butler and Mr. Bruce respectively.

OLIGODON SUBLINEATUS, Dum. & Bibr., Boulenger, p. 320.

Two examples of this species indistinguishable from the typical Ceylon specimens were got by Mr. de Roepstorff in the Nicobars and are now in the Museum.

OLIGODON SUBGRISEUS, Dum. & Bibr., Boulenger, p. 321.

This typically Indian species has spread over the natural boundaries of India as far as Killa Abdulla near the Khojak Pass in British Baluchistan whence the Museum possesses a specimen presented by Mr. J. A. Murray.

OLIGODON MELANOCEPHALUS, (Günth.), Boulenger, p. 317.

Gunther (P. Z. S. 1864, p. 491) and Jan. (Icon. Ophid. livr. xiii, pl. iii, fig. 4, Oct. 1865), seem to have described and figured the same snake independently under the same specific name, the former making a new genus for its reception, the latter including it in the genus *Homolosoma*.

The examination of a specimen brought by Dr. Anderson from Sebastiyeh (Samaria) in Palestine confirms Boulenger in placing the species in the genus *Oligodon*.

RHAGERRHIS PRODUCTA, (Gerv.), Peters Monatsb. Akad. Berlin, 1862, p. 275; Murray, Ann. Mag. N. H. (5) xiv, p. 104.

This rather curious snake was described by Peters (*l. c.*) from Senaar in N. E. Africa, and has since been recorded by Murray from Tanjistan and Bushire in Persia; one of the Tanjistan specimens is now in the Indian Museum and agrees with Peters' description in every respect.

ZAMENIS KORROS, (Schleg.), Boulenger, p. 324.

There is an undoubted example of this species in the Indian Museum said to have been procured by Dr. E. F. Keleart in Ceylon; Anderson quoting from Ferguson's "Reptile Fauna of Ceylon" states that this species is not found in Ceylon; it is possible therefore that the specimen in question may be wrongly labelled though there does not seem to be any particular reason for this being the case.

ZAMENIS VENTRIMACULATUS, (Gray), Boulenger, p. 325.

ZAMENIS LADACCENSIS, Anderson, Boulenger, p. 326.

There are no specimens of *Z. ventrimaculatus* in the Indian Museum from Persia or anywhere outside the Indian Empire. The Museum possesses examples from the following localities. Below Simla, Sabathu, Rajanpur in the Punjab, Jeypore Rjpt., and Karachi. Of *Z. ladaccensis* the Indian Museum possesses specimens from Shiraz (including the type of *Gonyosoma dorsalis*, Anders) Bushire, Karman and Regan in Persia; Askan, Zamran, Hung, and Quetta in Baluchistan; Gilgit and Ladak.

If this really represents the true distribution of the two species their geographical areas are quite separate and the two species may be considered quite distinct.

ZAMENIS DIADEMA, (Schleg.), Boulenger, p. 328.

The distributional area of this snake may be extended eastwards as far as Allahabad perhaps as far as Purneah, as there is a specimen probably from the latter place in the Indian Museum.

ZAOCCYS TENASSERIMENSIS, sp. nov. (Plate VI, fig. 3.)

Rostral as broad as deep, just visible from above; suture between the internasals two-thirds of that between the pre-frontals, frontal longer than its distance to the end of the snout, shorter than the parietals; three loreals, one larger anterior, and two smaller posterior; one long and narrow preocular reaching the top of the head but not touching the frontal; one subocular below it wedged in between the fourth and fifth labials; two postoculars; a single pair of long temporals on either side, the posterior temporals not larger than the ordinary scales; upper labials 7 to 8 in number, one very large labial alone entering the eye; in the specimen described this is on one side the fourth, on the other the fifth; five lower labials in contact with the anterior chin shields which are equal to the posterior; scales in sixteen rows all smooth, the two median dorsal rows and the two outer rows adjoining the ventrals on either side are more or less broad and quadrangular, whereas the five intermediate rows on either side are narrow and oblique. Ventrals not angulate, 201; Subcaudals 123; anal divided.

Colour, black above to greenish olive on the head, anteriorly traces of a white vertebral line, in the middle part of the body a series of ill defined white transverse bands bordered with black posteriorly, on the hinder part of the body, the bands are gradually transformed into seven longitudinal series of white spots separated by a black network, of the series of spots the outer series on either side are on the ventrals; below

yellowish the outer edges of the ventrals dusky, tail with a median dusky streak.

This snake is quite different from the only other Indian species of the genus, *Z. nigromarginatus*; in fact it belongs to the other section of the genus characterized by Günther by the possession of three loreals and named by him *Zapyrus*.

It seems to most resemble *Zaocys fuscus* from Borneo, but differs from this snake in colouration and also in the number and position of upper labials.

COLUBER HELENA, Daud., Boulenger, p. 331.

The range of this snake extends somewhat beyond the limits imposed by Boulenger; the Indian Museum possesses examples from the Purneah district and Mutlah in Bengal and from Samagooting in Assam.

COLUBER RETICULARIS, Cantor, Boulenger, p. 332.

The range of this species too may be extended from Sikkim and Assam southwards to Arakan and Pegu whence the Museum possesses specimens.

COLUBER TAENIURUS, (Cope), Boulenger, p. 333.

Coluber nuthalli, Theobald (Cat. Rept. As. Soc. Mus., p. 51) the type of which is in the Indian Museum, is obviously a young specimen of *C. taeniurus* and is not identical with *Coluber helena* as suggested by Boulenger.

COLUBER RADIATUS, Schleg., Boulenger, p. 333.

The Indian Museum possesses examples of this snake from Backergunge in Lower Bengal (E. Taylor) and from Hong Kong, from neither of which localities is this species recorded by Boulenger.

COLUBER PRASINUS, Bly., Boulenger, p. 334.

COLUBER OXYCEPHALUS, Boie, Boulenger, p. 335.

There are in the Museum examples of both these snakes from Darjeeling (Gammie) whence they are not recorded by Boulenger.

TROPIDONOTUS CHRYSARGUS, Schleg., Boulenger, p. 345.

TROPIDONOTUS NIGROCINCTUS, Bly., Boulenger, p. 346.

The distinction given by Boulenger in his key between these two species I find to be by no means a constant one; several of the specimens of *Trop. nigrocinctus* in the Museum possess two anterior temporals; the

colouration, however, of the two species is very distinct, the three oblique black streaks, below the eye, behind the eye and on the neck at once distinguishing *Tropidonotus nigrocinctus* from its ally.

TROPIDONOTUS HIMALAYANUS, Günth., Boulenger, p. 347.

This species is much more distinct from *T. subminiatus* than would be gathered from a perusal of Boulenger's description; *T. himalayanus* is much darker and very nearly always retains traces of the dorso-lateral series of white and black spots which are always found in the young, whereas in *T. subminiatus* the ground colour above is much lighter, almost blue, and very seldom retains any traces of the spots; in *T. himalayanus* the labials are all light coloured, edged with black and the oblique dark streak, under the eye so conspicuous a feature in *T. subminiatus* is altogether absent; the ventral surface in *T. himalayanus* is very dark, in some specimens almost black, whereas in *T. subminiatus* it is never dusky.

In none of the large number of specimens of *T. subminiatus* which I have examined, is there any trace of the keels on the outer row of scales, in *T. himalayanus*, however, the outer row of scales is as often keeled as not, and sometimes almost as strongly as in *T. chrysargus* and *T. nigrocinctus*.

The Indian Museum possesses examples of *T. subminiatus* from Sikkim, the Garo, Khasia and Naga hills, Manipur, Yunnan, Burma and from throughout Tenasserim; of *T. himalayanus* from Darjeeling (3000 to 4000 feet), the hills of Assam and Moulmein in Burma.

TROPIDONOTUS PLUMBICOLOR, Cantor, Boulenger, p. 351.

As is so often the case with Southern Indian forms, this species extends its range northwards to Mt. Aboo in Rajpootana; the Indian Museum also possesses examples from Nowgong and the Upper Godavery district in the Central Provinces, from the Nilgiri, Anamalai and Tinnevely hills in S. India and from Galle in Ceylon.

TROPIDONOTUS ANGUSTICEPS, Blyth, J. A. S. B. xxiii, p. 295; Boulenger, p. 352.

Three bottles containing four snakes were found amongst the collection of the Indian Museum labelled *Tropidonotus angusticeps*, of these snakes one was obviously *T. piscator* and does not seem to have been one of Blyth's original specimens; of the others, two with no history attached are without doubt examples of *Tropidonotus hydrus*, and the fourth, which was said to be the actual type of *T. angusticeps* collected by Capt. Abbott

in Ramri Island on the Arakau coast is a very faded specimen of *Pseudoxenodon macrops*.

The description of *T. angusticeps* seems to refer to both the latter species, *T. hydrus* and to *Pseudoxenodon macrops*.

TROPIDONOTUS PEALII, sp. nov. (Plate VI, fig. 4.)

Eye moderate, its diameter hardly equal to its distance from the nostril; rostral just visible from above; internasals broadly truncated anteriorly, suture between them shorter than that between the prefrontals; frontal longer than its distance to the end of the snout, shorter than the parietals; loreal nearly square; 1-2 preoculars; 2-3 postoculars; temporals 2 + 2; upper labials nine, fourth and fifth entering the eye; five pairs of lower labials in contact with the anterior chin shields, which are shorter than the posterior. Scales in 19 rows, strongly keeled, outer row also keeled but not so strongly as the rest; ventrals 142-144, subcaudals 75-77, anal entire.

Colour in spirit dark brown above, with a narrow light longitudinal line on either side, edged rather darker reaching the length of the body, below on either side occupying the lateral scales bordering the ventrals another light and much broader band two scales wide; head dark brown above, the upper and lower labials and rostral yellow, edged and blotched with brown, ventrals very dark brown each tipped laterally with light yellow, the longitudinal band so formed enlarges anteriorly to form a large white mark under the posterior lower labials; an indistinct yellow line along the middle of the ventral shields rather more conspicuous posteriorly.

Total length, 20 in.; tail 5 in.

This species is a very well marked one in every way; it differs from all the other Indian species (except *T. plumbeicolor*?) in having an undivided anal shield; apart from this it is perhaps somewhat allied to *Tropidonotus parallelus* with which it agrees in having a small eye and the outer row of scales keeled.

There are in the Indian Museum two examples of this snake, both collected in the Sibsagar district of Assam by Mr. S. E. Peal, who has contributed very largely to our collection of snakes and after whom I have much pleasure in naming this fine new species.

TROPIDONOTUS NICOBARENSIS, sp. nov. (Plate VI, fig. 5.)

Eye large, its diameter exceeding its distance from the nostril; nostril just visible from above; internasals truncated, suture between them and between the prefrontals about equal in length; frontal longer than its distance to the end of the snout, very nearly as long as the parietals;

loreal squarish; one large preocular reaching the top of the head, but not touching the vertical; three postoculars; temporals 1 + 2, the posterior pair very much smaller than the anterior and hardly larger than the scales around; upper labials 7 or 8, the third and fourth, or fourth and fifth entering the eye; five pairs of lower labials in contact with the anterior chin shields which are much shorter than the posterior ones. Scales in 19 rows strongly keeled including the outer row; ventrals 162; subcaudals 119; anal divided.

Colour, above bluish olive with a longitudinal dorsal band from the nape to the tail about two scales wide bordered on either side by a narrow black line, another indistinct white line laterally on either side, head uniform, a black streak behind the eye; below lighter than above, uniform; chin, upper labials and snout yellowish without the bluish tinge.

The only example of this new species is one from Camorta in the Nicobars, procured there by Mr. de Roepstorff.

I have referred this snake to *Tropidonotus*, but I am by no means certain that it is properly there located; the only example in the Museum is a small and obviously young one, and I cannot make out that there is very much difference in the size of the maxillary teeth, and the number of subcaudals is very high for this genus; the only species which it seems to resemble at all is *Prymniodon chalcus*, Cope (Günther, Reptiles Brit. Ind. p. 274) which is said to have come from Siam; with the description of this species in Günther, it agrees admirably except in the two important characters of the dentition and the anal shield.

The maxillary teeth of *Prymniodon* are said to be very considerably larger anteriorly than posteriorly, and the anal shield is entire; in the species before me the maxillary teeth appear to be of equal length throughout, and the anal is divided; it is therefore impossible to identify the Nicobar species with *Prymniodon* and I have thought it best, until more specimens are forthcoming to leave it in the genus *Tropidonotus*.

TROPIDONOTUS RHODOMELAS, Boie, Blanford, P. Z. S. 1881, p. 221.

Tropidonotus mortoni, Theobald, (Cat. Rept. As. Soc. Mus. p. 57) is referable to this species of which we have examples from Singapore and Sinkip Island, Sumatra.

TROPIDONOTUS TRIANGULIGERUS, Schleg., Anderson, Journ. Lin. Soc. xxi, p. 345.

This snake was obtained by Dr. Anderson in Mergui and the Museum also contains examples from North Tenasserim, the Malay Peninsula and from Sinkip Island, Sumatra; it is not mentioned by Boulenger, and must be added to the Indian Fauna.

DIPSAS MULTIFASCIATA, Blyth, J. A. S. B. xxix, p. 114.

This species, the type of which was described by Blyth, from Subathu near Simla in the North-West Himalayas, was afterwards identified by Stoliczka (J. A. S. B. xxxix, p. 199) with *D. ceylonensis* a species which is otherwise apparently confined to Southern India and Ceylon.

I cannot agree with Stoliczka with regard to this identification, in the first place the preocular shield which extends nearly to the vertical in the case of *D. ceylonensis*, does not or barely reaches the top of the head in *D. multifasciata*; again in none of the specimens of *D. ceylonensis* which I have been able to examine is there the slightest trace of the narrowing of the preocular shield below, so that the lower corner of the loreal enters the eye, this arrangement is found more or less markedly in all the specimens of *D. multifasciata*; finally the colouration of the two species is very different, though perhaps difficult to describe.

The following shows at a glance the difference between the two species, the characters being taken from an examination of the specimens in the Museum:—

Dipsas multifasciata.

Loreal entering the eye below the preocular except in two cases where the lower corner of the loreal is prolonged in that direction, but does not quite reach it.

Preocular barely reaching the top of the head in some cases.

Temporals 1+2 or 2+1.

Scales in 21 rows.

Ventrals 231-248. Subcaudals 96-109.

Head with a well marked lateral black band from the prefrontals to the end of the parietals.

A median black streak on the nape.

The labials edged with black.

A well marked streak from the eye to the gape.

Ventrals marked with lateral square blotches throughout.

Dipsas ceylonensis.

No approach of the loreal to the eye, preocular of equal width throughout.

Preocular reaching the top of the head and sometimes touching the vertical.

Temporals 2+3.

Scales in 19 rows.

Ventrals 221-3. Subcaudals 90.

All the upper head shields black blotched, no definite longitudinal band.

Three ill-defined longitudinal black bands on the nape generally joined by a transverse band behind and forming a trident.

Labials not edged with black.

Streak from eye to gape ill-defined and interrupted.

Ventrals irregularly and very lightly spotted.

Of *Dipsas ceylonensis*, the Indian Museum possesses examples from the Anamalai hills only; of *D. multifasciata* besides the type from Subathu near Simla examples from Mussooree, Naini Tal and Darjeeling.

DIPSAS CYANEA, (Dum. & Bibr.), Boulenger, p. 361.

This species occurs in Tenasserim; there is an example in the Indian Museum from Tavoy.

DIPSAS CYNODON, Cuv., Günther Reptiles Brit. Ind., p. 368.

This species must be added to the fauna of the Indian Empire; there are in the Indian Museum undoubted examples from the Garo hills (Capt. Williamson), Samagooting (Capt. Butler) and Cachar (Museum Collector) in Assam; and from Thyetmyo (W. T. Blanford) the Burma-Siam hills (Museum Collector), and Mergui (W. Theobald) in Burma; this species is recorded from two of the above localities by Theobald in his Catalogue of the Snakes of the Asiatic Society, and it seems curious that this should have been overlooked by Boulenger.

DRYOPHIS FRONTICINCTUS, Günth., Boulenger, p. 368.

There is a snake in the Indian Museum from Sibsagar in Assam presented by Mr. S. E. Peal which I am unable to identify with any other species; if this is the case it cannot be that this snake is confined to the neighbourhood of brackish water as is suggested by Stoliczka.

DRYOPHIS MYCTERIZANS, (Daud.), Boulenger, p. 370.

This snake like many other Southern Indian species extends north-westwards as far as Mt. Aboo in Rajputana.

DRYOPHIS PULVERULENTUS, (Dum. & Bibr.), Boulenger, p. 371.

This species is apparently generally distributed throughout Peninsular India, there are six undoubted examples in the Indian Museum collected by Mr. V. Ball in Maunbhoom.

CERBERUS RHYNCHOPS, (Schneid.), Boulenger, p. 374.

This species appears to be common on the Andamans and Nicobars.

HYPsirrhina BLANFORDII, Boulenger, p. 377.

This species is founded on a single specimen described by Blanford as probably from the neighbourhood of Bassein; this unique specimen does not seem to be in the British Museum as Mr. Boulenger distinctly says that he has not been able to examine it; there is, however, an ex-

ample of this species in the Museum with no recorded history, which had been identified by Dr. Anderson as *Gerarda bicolor*, and it is possible that this is the missing type which has lost its label.

HYPSSIRHINA SIEBOLDII, (Schleg.), Boulenger, p. 377.

This species occurs in Assam; the Indian Museum contains a specimen from Samagooting in the Naga hills.

FORDONIA LEUCOBALIA, (Schleg.), Boulenger, p. 378.

This species occurs in the Sunderbunds, there is an example thence in the Museum, presented by the Rev. H. J. Harrison.

CALLOPHIS NIGRESCENS, Günth., Boulenger, p. 384.

This species varies considerably in colour, of the specimens in the Indian Museum, there is one referable to the spotted variety (var. A. of Boulenger) from the Wynaad; three of the variety with three longitudinal white-edged bands (var. B. of Boulenger) from Malabar, the Anamalai and the Shevaroy hills; and finally two of the variety with five longitudinal bands from Malabar and Ganjam.

MEGAEROPHIS FLAVICEPS, Reinh., Günther Rept. Brit. Ind. p. 346.

This species, which is not mentioned by Boulenger, should be included among the Snakes of the Indian Empire; there is an example of it in the Museum from Mergui in Tenasserim, presented by Mr. W. Theobald.

BURGARUS CAERULEUS, (Schneid.), Boulenger, p. 388.

This snake, which is found all over India proper, appears to be very rare on the eastern side of the Bay of Bengal, in fact the Indian Museum possesses what I believe to be the only specimen recorded thence, one procured by Col. Nuthall from Rangoon, and this might have easily reached Burma by ship. Lately, however, the Museum has received a pair of "Kraits" from Meiktalla in Upper Burma where they were found by General H. Collett which differ in some respects from the ordinary Indian form of this snake, although hardly perhaps enough to merit specific distinction.

The following are the points in which this variety differs from the typical Indian form.

The rostral is deeper than broad and extends back to about two-thirds or three-fourths of its distance from the frontal; the ventral shields number 223 and 228 respectively and the subcaudals 49.

The colour above is a dark slaty blue, the dorsal scales are alternately yellow and slate coloured, the yellow extending for about 10 or 12 scales and reappearing again after a somewhat longer interval; the scales on the sides of the yellow dorsal scales also exhibit conspicuous pale edgings so that the whole snake has the appearance of about 12 light coloured bands encircling the body.

BUNGARUS BUNGAROIDES, (Cantor), Boulenger, p. 389.

An example of this snake from Darjeeling presented by Mr. J. Gammie has a well marked loreal shield present on either side of the head.

BUNGARUS LIVIDUS, Cantor, Boulenger, p. 389.

The range of this species extends as far as Lower Bengal as is shown by an example procured at Saidpur in the Dinajpur district presented by Mr. W. de W. Peal.

The vertical scales of this snake are said by Mr. Boulenger to be "but feebly enlarged and not broader than long," in two, however, out of the three examples of this species in the Museum, the vertebral scales are certainly broader than long in the posterior part of the body.

NAIA TRIPUDIANS, Merr., Boulenger, p. 391.

There are in the Indian Museum a very large number of Cobras from different parts of India, and it seemed worth while to try and make out how far the various colour varieties of the Cobra were constant to fixed geographical areas.

The specimens in the Museum are all spirit-preserved ones and not very large, and although a great deal more has still to be done before an accurate knowledge of the geographical distribution of the varieties can be made out, the following seems to roughly indicate the truth.

a. Hood with the well known spectacle-marking on it; above and below, stone coloured, with the characteristic dark pectoral band across the chest.

In the Indian Museum examples from Banda, N.-W. P., Ganjam district, Calcutta, and the Krishnagar district.

This is "var. a" of Günther's Reptiles and the "Gokurrah" of Fayrer, and is probably found throughout the peninsula of India.

b. Hood with a white circular marking, edged with black, behind the hood-marking is a dark, followed by a light ring encircling the body, rest of the body dark, mottled lighter.

In the Indian Museum examples from Calcutta only where it ap-

pears to be the commonest variety, this is the "Keantiah" of Fayer and "var. θ " of Günther who also records it from Assam and Sikkim.

c. Hood as above with the circular marking; uniform brown above, below, anteriorly light coloured, posteriorly darker.

In the Indian Museum examples from Assansole, Bengal, Calcutta, Sibsagar and Samagoting, Assam, Chittagong and the Andamans.

This is the variety most commonly met with, I believe, throughout Burma, it appears to correspond to "var. ξ " of Günther which he records from Siam.

d. Hood with no marking, light coloured above and below with darker pectoral bands.

The Indian Museum contains examples of this species from the Rungpur district in Bengal, from Assam, the Chittagong hills, Mandalay and Mergui.

e. Hood with no marking, blue black above and below except slight traces of lighter colour on each side of the throat.

The Indian Museum contains examples of this well-marked variety from the Andamans (?) and Singapore, where it appears to be very abundant.

It seems to correspond to part of "var. ϵ " of Günther.

f. No marking on the hood; very dark brown, almost black above and below.

The Indian Museum contains examples of this variety from the Punjab and Rajputana, and this is probably the same as part of Günther's "var. ϵ " which came from the Deccan.

g. No marking on the head, colour a light sandy with bluish tinge; two dark bands on the neck extending all round, behind these two anterior bands a series of chevron-shaped forwardly directed bands, which become fainter towards the tail.

There are examples of this curiously coloured variety in the Indian Museum from Khojak in British Baluchistan and from the Punjab.

This variety was first described by Eichwald as *Tomyris oxiana* and has been shown by Boulenger and Boettger to be conspecific with the typical *N. tripudians*.

DISTIRA CYANOCINCTA, (Daud.), Boulenger, p. 410.

The types of *Hydrophis trachyceps*, Theobald, (Cat. Rept. As. Soc. Mus. p. 70) and *Hydrophis crassicollis*, (Anderson, J. A. S. B. xl, p. 19) may both be referred to this species, the former was got at Mergui, the latter in the Hooghly below Calcutta.

AMBLYCEPHALUS MONTICOLA, (Cantor), Boulenger, p. 415.

There is a snake in the Indian Museum from Camorta in the Nico-

bars presented by Mr. F. A. de Roepstorff which seems quite identical with undoubted examples of *A. monticola* from Assam.

AMBLYCEPHALUS MACULARIUS, (Blyth ?), Boulenger, p. 416.

AMBLYCEPHALUS CARINATUS, (Reinw.), Günther, Reptiles Brit. Ind., p. 326.

A good deal of confusion in the synonymy of these snakes has been caused by Mr. Theobald, who when examining and cataloguing the Asiatic Society's collection of Snakes identified as the adult and young of the same species certain snakes which had been referred by Blyth to two different species, *i. e.*, *Aplopeltura boa*, Schleg. (Theobald's adult) and *Pareas macularius*, Blyth (Theobald's young).

I have been quite unable to find any published description by Blyth of the latter species and so I must conclude that *Pareas macularius* is a manuscript name.

These snakes which were five in number Theobald first of all (J. Linn. Soc. x, p. 54), referred to *Pareas macularius*, the manuscript name given by Blyth to the three smaller specimens only; afterward (Cat. Rept. Mus. As. Soc., p. 63) he gave a new name, *Pareas berdmorei*, to these same five snakes, and finally (Cat. Rept. Brit. Ind., p. 203) he identified them with *Pareas margaritophorus*, Jan.

On examining the five snakes in question, it was at once evident that the two larger examples were quite distinct from the three smaller ones in colour and disposition of the head-shields and in fact in every way.

The larger snakes have internasals nearly as large as the prefrontals which latter shields are excluded from the eye, and in every other respect resemble *A. carinatus*; while the smaller snakes in which the prefrontal enters the eye, agree with the description of *A. macularius* as given in Boulenger's Reptiles.

Pareas berdmorei is therefore only in part a synonym of *Amblycephalus macularius* as described by Boulenger, and also in part a synonym of *Amblycephalus carinatus*.

There are in the Indian Museum only the three original specimens of *A. macularius*, which were procured by Major Berdmore at Martaban in Burma; of the other species, *A. carinatus*, which is an addition to the Fauna of the Indian Empire, besides the two specimens procured by Major Berdmore in Tenasserim, there are two from Tavoy (Museum Collector), one from the Burma-Siam hills and one from Mergui (Anderson).

TRIMERESURUS GRAMINEUS, (Shaw), Boulenger, p. 429.

TRIMERESURUS PURPUREOMACULATUS, (Gray), Boulenger, p. 429.

Of these two species there is a very large series in the Museum; and if the insular and Malayan forms be excluded, the two species are fairly

constant in scaling and colour. The points of distinction between the two species are as follows :

In *T. purpureomaculatus* the head scales are juxtaposed and convex or keeled, the temporal scales are strongly keeled, the scales are in 25-27 rows and the colour is in spirit a porphyraceous blue; in *T. gramineus* the head scales are small, smooth, and imbricate, the temporal scales are also smooth, the scales are in 19 to 23 rows and the colour is a uniform green.

It is in most cases perfectly easy to distinguish these two snakes, but there are in the Indian Museum some specimens which are intermediate in character, whether they are hybrids or not it is difficult to say, but as the two snakes inhabit approximately the same geographical area, it is quite possible that this may be the case.

Among the intermediate forms may be mentioned nine snakes from Sibsagar in Assam (register no. 4015-23) which have the juxtaposed convex head shields of *T. purpureomaculatus*, but the smooth temporals and 21 rows of scales of *T. gramineus*; again a snake (register no. 4109) from Moulmein has the convex head shields and keeled temporals but only twenty-one rows of scales.

Of *T. purpureomaculatus* the Museum contains examples from Lower Bengal, and from various localities in Assam and Burma; and of *T. gramineus*, examples from Simla, Sikhim, Assam, Burma and Hongkong.

In the Islands of the Bay of Bengal, Preparis Isle, Cocos Isles, the Andamans and Nicobars there are several different coloured varieties of Pit vipers which, in my opinion with one exception only, are better referred to *Tr. gramineus* than to *Tr. purpureomaculatus*; besides these varieties there is a closely allied form, which is recognised by Boulenger as a separate and distinct species; this is *Tr. cantoris* of Blyth, of which the Museum possesses a fair series, all from the Nicobars with one exception, which is from the Andamans.

The following is a list with brief descriptions of the various varieties above mentioned.

a. Preparis Island variety.

Light brown above with darker dorsal and lateral spots; ventrals light coloured, marbled with brown, scales in 25 rows; this form has the characteristic juxtaposed convex head scales and keeled temporals, and I have considered it to be an insular variety of *T. purpureomaculatus*.

b. Mottled variety.

Brown above blotched with blue, below bluish with brown blotches, scales in 23-25 rows, head scales as in *T. gramineus*. In the Indian Museum there are a good number of snakes of this variety from the Andaman Islands alone.

c. Brown variety.

This resembles the last, but there is little or no trace of the blue mottling on the ventrals which are almost uniform brown. Two examples from the Andamans in the Indian Museum.

d. Uniform or banded variety.

Colour uniform light reddish or dusky or with regular white transverse bands; scales in 21 rows.

This is the variety described by Stoliczka under the name of *T. mutabilis*, and is found on both the Andamans and Nicobars.

e. Green above, lighter below, resembling the typical variety found in Assam and Burma, scales in 21-25 rows.

This variety occurs on the Andamans, Cocos and Nicobars.

All the above varieties with the exception of the first seem to be referable to *T. gramineus* rather than to *T. purpureomaculatus*.

EXPLANATION OF PLATE VI.

- Fig. 1. *Ablabes stoliczkae*, sp. nov. × 2.
 2. *Simotes woodmasoni*, sp. nov.
 3. *Zaocys tenasserimensis*, sp. nov.
 4. *Tropidonotus pealii*, sp. nov.
 5. *Tropidonotus nicobaricus*, sp. nov. × 2.

XI.—*Catalogue of the Diptera of the Oriental region* by MONS. J. M. F. BIGOT. PART I. Communicated by the SUPERINTENDENT OF THE INDIAN MUSEUM.*

[Received Feb. 26th, 1891. Read March 3rd, 1891.]

Order DIPTERA.

Suborder HOMALOCERATI.

J. Bigot, *adhuc ined.*

Division NEMATOCERATE.

J. Bigot, *adhuc ined.*: Nematocera, Meigen, *Syst. Besch.*, i, 1818.

Family CULICIDÆ.

J. Bigot, *adhuc ined.*: Culiciformes, Meigen *Syst. Besch.* i, 1818; Culicidæ, Latr. *Fam. Natur.* 1825; Macquart, *S. d. Buff. Dipt.* i, 1834; Culicinæ, Zetterst., *Ins. Japonica*: Culicinæ Rondaui, *Prodr.* i, 1856; Culicina, Schiner, *Faun. Austr. Fliegen* 1864.

* This Catalogue was drawn up by M. Bigot at the suggestion of the late Mr. Atkinson and was to have formed part of the series of Catalogues of the Insecta of the Oriental Region, which have been discontinued since Mr. Atkinson's death.

• Genus ANOPHELES.

- Meigen, *Syst. Besch.* i, 1818, p. 10: *Culex*, pt., Linn.
annularis, V. der Wulp, *Notes Leyden Mus.*, vi, 1884, p. 249.
 Hab. Java.
barbirostris, id. *ibid.*, p. 248.
 Hab. Java.
sinensis, Wiedem., *Auss. Europ. Zweifl. Ins. Hamm.* 1828, i, p. 47.
 Hab. China.

Genus MEGARHINUS.

- Rob.-Desvoidy, (*alias Megarhina*) *Essai sur les Tipul.*: *Culex*, pt.
splendens, (*Culex*) Wiedem., *Zool. Magaz.* iii, p. 2.
 Hab. Java.

Genus CULEX.

- Linn., *Faun. Suec.* 1761: Meigen, *Illig. Magaz.* ii, p. 260, (auct).
laniger, Wiedem., *Auss. Europ. Zweifl. Ins. Hamm.* 1828, p. 5.
 Hab. Java.
fuscatus, id. *ibid.*, p. 6.
 Hab. India.
fatigans, id. *ibid.*, p. 10.
 Hab. India.
molestus, id. *ibid.*, p. 542.
 Hab. Sumatra.
sitiens, id. *ibid.*, p. 543.
 Hab. Sumatra.
vagans, id. *ibid.*, p. 545.
 Hab. China.
amboinensis, (*Melius Megarhina*) Doleschall *Natuur. Tijdschr. Nederl. Indië.* xiv,
 1857, p. 381.
 Hab. Amboina.
aureostriatus, id. *ibid.*, p. 385.
 Hab. Amboina.
cingulatus, id. *Natuur. Tijdschr. Nederl. Indië, Batavia.* x, 1856, p. 405.
 Hab. Java.
subulifer, id. *ibid.*, xiv, 1857, p. 382.
 Hab. Amboina.
nero, id. *ibid.*, xiv, 1857, p. 383.
 Hab. Java.

setulosus, id. *ibid.*, xiv, 1857, p. 384.

Hab. Java.

luridus, id. *ibid.*, xiv, 1857, p. 384.

Hab. Java.

variegatus, id. *ibid.*, xvii, 1858, p. 77.

Hab. Amboina.

annulipes, Walker, *Journ. Proceed. Linn. Soc. London*, i, 1857, p. 6.

Hab. Singapore.

imprimens, id. *ibid.*, v, 1861, p. 144.

Hab. Amboina.

ventralis, id. *ibid.*, v, 1861, p. 144.

Hab. Amboina.

dives, Schiner *Novara Reise* 1868, p. 31.

Hab. Batavia.

longipalpis, V. der Wulp, *Tijdschr. Ent.* xxii, blz. lxxvii, p. 9.

Hab. Sumatra.

crassipes, id. *ibid.*, p. 9.

Hab. Sumatra.

Family CHIRONOMYDI.

J. Bigot, *adhuc ined.*: Chironomydæ, Schiner *Faun. Austr. Flieg.*, ii, 1864, p. 574; Chironomides Westw., *Introd. Entom.* 1840: Tipul. Culiciform Macquart. *S. à Buff. Dipt.* i, 1834: Chironomides, Macquart, *Dipt. Exot.*, 1838, p. 36.

GENUS CERATOPOGON.

Meigen, *Illig. Magaz.* ii, 1803, p. 261: Culex pt. Linn.: Chironomus pt. Fabr. Fallen: Culicides, Latr.; Palpomyda, Prionomyda, Ceratopogon pt., Rossi.

trichopus, Thomson, *Eugenia Resa*, 1868, p. 444.

Hab. China.

agas, Rondani, *Ann. Mus. Civ. Genova*, vii, 1875, p. 462.

Hab. Borneo.

GENUS CHIRONOMUS.

Meigen, *Illig. Magaz.* ii, 1803, p. 260; Thalassomyia, pt. Schiner: Tipula, pt. Linn.

venustus, Wiedem., *Aus. Europ. Ziccifl. Ins. Hamm.* 1828, i, p. 547.

Hab. China.

vicarius, Walker, *Ins. Saundés. Dipt. London*, 1856, p. 423.
Hab. India.

cubiculorum, Doleschall, *Natuur. Tijdschr. Nederl. Indie, Batavia*, x, 1856, p. 405.
Hab. Java.

pictus, id. *ibid.*, xiv, 1857, p. 386.
Hab. Java.

socius, Walker, *List Dipt. Ins. Brit. Museum London*, 1848, pt. 1, p. 16.
Hab. India.

Genus TANYPUS.

Meigen, *Illig. Magaz.*, ii, 1803, p. 261: *Tipala* pt. Linn., Degeer: *Chironomus*
pt. Fabr.

pardalis, Doleschall, *Naturk. Tijdschr. Nederl. Indie, Batavia*, v, 1856, p. 405.
Hab. Java.

melanurus, id. *ibid.*, p. 405.
Hab. Java.

cyanomaculatus, id. *ibid.*, p. 406.
Hab. Java.

nigrocinctus, id. *ibid.*, p. 406.
Hab. Java.

ornatus, id. *ibid.*, xiv, 1857, p. 385.
Hab. Java.

crux, Wiedem. *Anal. Entom.* 10, id., *Ausser Europ. Zweifl. Ins.* i, Hamm, 1828,
p. 10.
Hab. India.

Genus MACROPEZA.

Meigen, *Syst., Besch.*, i, 1818, p. 87: Wiedem., Macq., Walker, Schiner,
Rondani, (et auct.).

gibbosa, Wiedem., *Ausser Europ. Zweifl. Ins.* pt. i, Hamm, 1828, p. 20.
Hab. India.

Family CECIDOMYDI.

Bigot, *adhuc ined.* 1891: *Tipulariae Gallicoloe*, Gallmücken, Meigen, *Syst.*
Beschr. 1818, i, p. xxxiv: *Cecidomyida*, Westw. *Introd. Entom.* 1840:
Cecidomyidae, Schiner, *Faun. Austr. Fliegen*, ii, Wien, 1864, p. xviii.

Genus CECIDOMYIA.

Meigen, *Illig. Magaz.* ii, 1803, p. 261, Latr., Wiedem., Macq., Schiner,
Rondani, V. de Wulp, Walker, (et auct.): *Tipala*, pt. Degeer: *Dasyneura*
pt. Rondani: *Oligotrophus*, pt. Latr.

deferenda, Walker, *Journ. Proceed. Linn. Soc. London*, 1, 1857, p. 105.
Hab. Sarawak.

oryzae, Wood-Mason, *Indian Museum News*, i, p. 103, pl. vi, fig. 6.
Hab. Bengal.

Genus LASIOPTERA.

Meigen, *Syst. Besch.* i, p. 88, 1818: *Cecidomyia* pt. Meigen: *Tipula* pt.
Linn., Degeer.

bryonicæ, Schiner, *Novara Reise* 1868, p. 5.
Hab. Madras.

Family DIXADI.

Bigot, *adhuc ined.* 1891.

Genus DIXA.

Meigen, *Syst. Besch.* i, 1818, p. 216.

guttipennis, Thomson, *Eugenie Resa* 1868 p. 448.
Hab. China.

Family TIPULIDI.

Tipulidæ, Leach, in *Sam. Comp.* 1819: *Tipulidæ*, Schiner, *Faun. Austr.*
Fliegen ii, 1864, p. 495: *Tipulariæ*, Latr. *Hist. Nat. Ins.* 1802: *Tipulari-*
des, Leach *Edinb. Encycl.* 1815: *Tipulides*, Westw. *Introd. Mod. Class.*
Insects, 1840.

Genus CTENOPHORA.

Meigen, *Illig. Magaz.* ii, 1803, p. 263: Wiedem., Macq., Schiner, Zetterst.,
Rondani, Ost.-Sacken pt.: *Pseliophora*, pt. Ost.-Sacken. *Berlin Ent.*
Zeitschr. 1886, xxxi. p. 168: *Tanyptera*, Latr.: *Dictenidia* et *Xyphura*,
pt. Brullé.: *Ceroctena* et *Xyphura* pt. Rondani.

melanura, Walker, *List Dipt. Ins. Brit. Museum London*, i, 1848, p. 78.
Hab. Nepal.

xanthomelana, id. *ibid.*, p. 77.
Hab. India.

melanura, id. *ibid.*, p. 78.
Hab. Nepal.

Genus PSELLIOPHORA.

Ost-Sacken, *Berlin Entom. Zeitschr* 1886, xxx, p. 168: *Ctenophora* pt. Meigen
læta (*Ctenophora*) Wiedem., *Ausser. Europ. Zweifl. Ins.* i, p. 40: (*Tipula* id. Fabr.)
Hab. India.

- ardens, (*Ctenophora*), id. *ibid.*, p. 39.
Hab. Java.
- compedita, (*Ctenophora*) id. *ibid.*, p. 39.
Hab. Java.
- taprobanes, (*Ctenophora*) Walker *List. Dipt. Ins. Brit. Mus. London*, 1848, i, p. 77.
Hab. Ceylon.
- fumiplena, (*Ctenophora*) Walker *Ins. Saunders. Dipt. i, London*, 1856, p. 449.
Hab. China.
- javanica, (*Ctenophora*) Doleschall, *Natuurk. Tijdsch. Nederl. Indie, Batavia*, x,
1856, p. 406.
Hab. Borneo, Java.
- chrysophila, (*Ctenophora*) Walker *Journ. Proceed. Linn. Soc. Lond.* i, 1857, p. 6.
Hab. Singapore.
- curvipes, (*Ctenophora*) V. der Wulp, *Notes Leyden Mus.* vi, 1884, p. 254.
Hab. Gorontalo, Java?
- annulosa, (*Ctenophora*) id. *ibid.*, vii, p. 1.
Hab. Java.
- rubra, Ost.-Sacken, *Berlin. Entom. Zeitschr.* xxx, 1886, p. 171.
Hab. Laos.

Genus PRIONOTA.

- V. de Wulp, *Notes Leyd. Mus.* vii, 1885 p. 1.
- nigriceps, V. der Wulp, id. *ibid.*, p. 2.
Hab. Java.

Genus OLIGOMEA.

- Doleschall, *Natuurk. Tijdschr. Nederl. Indie, Batavia* xiv, 1857, p. 387,
Eriocera, pt. Maquart.
- javensis, Doleschall, *ibid.*, p. 387: Eriocera acrostacta, V. der Wulp.
Hab. Java.

Genus PACHYRHINA.

- Macquart, *S. à. Buff. Dipt.* i, 1834, p. 88: Tipula pt.: Nephrotoma, pt.
Olivier, *Encycl. Method.*
- bombayensis, Macquart *Dipt. Exot. Paris*, 1855, 5th *Suppl.* p. 15.
Hab. Bombay.
- delecta, Walker, *Ins. Saunders. Dipt. Lond.* 1856, p. 445, Tipula id. Macq.
Hab. India.

quadrivittata, V. der Wulp, *Notes Leyden Mus.* vi, 1885, p. 9.

Hab. Java.

triplasia, id. *ibid.*, p. 10.

Hab. Java.

fasciata, Macquart, *S. à Buff. Dipt.*, i, Paris, 1834, p. 90.

Hab. Java.

doleschalli, (*Nomen Novum*), *Tipula javensis*, *Doleschall Natuur. Tijdschr. Nederl. Indie, Batavia*, x, 1856, p. 406; *Ost.-Sacken, Ann. Mus. Civ. Genova*, xvi, 1881, p. 399: *P. fasciata*? Macq. (*Vid. Suppl.*).

Hab. Java.

Genus TIPULA.

Linn., *Faun. Suec.* 1761, p. 430 (et auct).

praepotens, Wiedem., *Auss. Europ. Zweifl. Ins.* i, *Hann.* 1828, p. 40.

Hab. Java.

monochroa, id. *ibid.*, p. 41.

Hab. Java.

pedata, Wiedem., *Dipt. Exot.* i, p. 23.

Hab. Java.

umbrina, Wiedem., *Auss. Europ. Zweifl. Ins.* i, *Hann.* 1828, p. 49: *T. congrua*, Walker.

Hab. Java.

javana, id. *Dipt. Exot.* i, 1821, p. 27.

Hab. Java.

castanea, Macq. *Dipt. Exot.* i, 1838, p. 54.

Hab. Java.

venusta, Walker, *List Dipt. Ins. British Museum*, i, 1848, p. 64.

Hab. Sylhet.

fulvipennis, id. *ibid.*, p. 67.

Hab. Nepal.

reposita, id. *ibid.*, p. 67.

Hab. Nepal.

melanomera, id. *ibid.*, p. 68.

Hab. Nepal.

nova, id. *ibid.*, p. 71.

Hab. Hong-Kong.

vicaria, id. *Ins. Saunders. Dipt. Lond.* 1856, pt. i, p. 445.

Hab. India.

javensis, *Doleschall, Natuur. Tijdschr. Nederl. Indie, Batavia*, x, 1856, p. 406.

Hab. Java.

- longicornis*, id. *ibid.* xvii, 1858, p. 79.
Hab. Amboina.
- vittis*, Walker, *Journ. Proceed. Linn. Soc. London*, i, 1857, p. 108.
Hab. Borneo.
- fumifinis*, id. *ibid.*, v, 1861, p. 145.
Hab. Amboina.
- sorrata*, V. der Wulp, *Notes Leyden Mus.* vii, 1885, p. 5.
Hab. Serahan?
- pilosula*, id. *ibid.*, p. 5.
Hab. Java.
- leucopyga*, id. *ibid.*, p. 6.
Hab. Java.
- brobdignagia*, Westwood, *Trans. Ent. Soc. Lond.* 1876, p. 504.
Hab. N. China.

Genus CONOSIA.

- V. der Wulp, *Tijdschr. Entom.* 1880, p. 159 : *Limnobia*, pt. Wiedem.
- irrorata*, V. der Wulp, *loc. cit.* p. 161 : *Limnobia* id. Wiedem. *Ausser Europ. Zweifl. Ins.* i, Hamm, 1828, p. 574.
Hab. Java.
- crux*, id. *ibid.*, p. 161 : (*Limnophila*) Doleschall, *Natuur. Tijdschr. Nederl. Indie, Batavia*, xiv, 1858, p. 388.
Hab. Java.

Genus LIMNOPHILA.

- Macquart, *S. á Buff. Dipt.* i, 1834, 95 : *Limnobia* pt. Meigen (et auctor).
Limnomyia, Rondani *Prodr.* 1861, vi, p. 11.
- basilaris*, Macquart, *Dipt. Exot.* i, Paris, 1838, p. 66 : *Limnobia* id. Wiedem. *Auss. Europ. Zweifl. Ins.* i, Hamm, 1828, p. 27.
Hab. Java.
- bicolor*, id. *ibid.*, p. 66 : *Eriocera* id. Ost.-Sacken.
Hab. Bengal.

Genus ERIOCERA.

- Macquart, *Dipt. Exot.* pt. v, Paris, 1838, p. 74, *Pterocosmus* pt. Walker :
Allarithmia, Loew, *Berust. Faun.* 1850, p. 38 : *Physecerania*? pt. Bigot,
Ann. Soc. Ent. France, (3) vii, 1859, p. 123 : *Arrhenica*, pt. Ost.-
Sacken, *Proceed. Acad. N. Sci. Philadelphia* 1859, p. 243.
- selene*, Ost.-Sacken *Ann. Mus. Civ. Genova*, xvi, 1881, p. 406.
Hab. Sumatra.

humberti, Ost.-Sacken, *Berlin. Ent. Zeitschr.*, xxxi, 1887, p. 221.
Hab. Ceylon.

meleagris, id. *ibid.*, p. 222.
Hab. Ceylon.

pachyrrhina, id. *ibid.*, p. 222.
Hab. Ceylon.

crystalloptera, id. *ibid.*, p. 222.
Hab. Ceylon.

albonotata, id. *ibid.*, p. 223 : *Limnobia*, id. Loew, *Peter's Reise*, p. 1.
Hab. Ceylon.

acrostacta, V. der Wulp, *Notes Leyden Mus.* vi, 1884 p. 11 : *Limnobia*, id. Wiedem.
Dipt. Exot. p. 1 : *Cylindrotoma*, id. Macquart, *Dipt. Exot.*, i, p. 168, *Suppl.* iii,
p. 7 : *Oligoneura javensis*, Doleschall.
Hab. Java.

albipunctata, V. der Wulp, *Tijdschr. Entom.* xxiii, p. 158.
Hab. Java.

ferruginosa, id. *Notes Leyden Mus.* vii, 1885, p. 13.
Hab. Java.

lunata, Westwood, *Trans. Ent. Soc. London*, 1881, pt. iii, p. 367.
Hab. Sarawak.

GENUS PTEROCOSMUS.

Walker, *List Dipt. Ins. Brit. Mus.* pt. 1, Lond. 1848, p. 78 : *Eriocera*, pt.
V der Wulp.

velutinus, Walker, *loc. cit.* p. 79.
Hab. Nepal.

hilpa, id. *ibid.*, p. 79 : *Eriocera* id., V. der Wulp, *Notes Leyden Mus.* vii, 1885
p. 12.
Hab. Hong-kong.

lunigerus, Walker, *Journ. Proceed. Linn. Soc. London*, i, 1857, p. 107.
Hab. Borneo.

infixus, id. *ibid.*, p. 107.
Hab. Borneo.

optabilis, id. *ibid.*, p. 107.
Hab. Borneo.

combinatus, id. *ibid.*, p. 107.
Hab. Borneo.

dilutus, id. *ibid.*, p. 108.
Hab. Borneo.

Genus LIMNOBIA.

Meigen, *Syst. Besch.* 1818, p. 116: Tipula, pt. Linn., Fabr.: Limonia,
Meigen, *Illig. Magaz.*, 1803: Dicranomyia, pt. et Limnobia, pt. Ost-
Sacken: Limnomyza, Rondani, *Prodr.* i, 1856: Glochina, pt. Meigen,
Staeger.

diana, Macq., *S. à Buff. Dipt. Paris*, 1834, p. 107.

Hab. Bengal.

sorbillans, Wiedem., *Auss. Europ. Zweifl. Ins. Hamm*, 1828, p. 551.

Hab. Sumatra.

trentepohlii, id. *ibid.*, p. 551.

Hab. Sumatra.

apicalis, id. *ibid.*, p. 551.

Hab. Sumatra.

bibula, id. *ibid.*, p. 552.

Hab. China.

mesopyrrha, id. *ibid.*, p. 26.

Hab. Java.

costalis, id. *ibid.*, p. 37.

Hab. India.

sumatrensis, Macquart, *Dipt. Exot. Paris, Suppl.*, iv, 1850, p. 16.

Hab. Sumatra.

aterrima, Walker, *Ins. Saunders, Dipt. London*, 1856, p. 434.

Hab. India.

substituta, id., *List Dipt. Ins. Brit. Mus. London*, i, 1848, p. 39.

Hab. China.

sanguinea, Doleschall, *Natuur. Tijdschr. Nederl. Indie*, xiv, *Batavia*, 1857, p. 391.

Hab. Java.

impressa, Walker, *Journ. Proceed. Linn. Soc. London*, i, 1857, p. 106.

Hab. Borneo.

rubescens, id. *ibid.*, p. 106.

Hab. Borneo.

pyrrhochroma, id. *ibid.*, p. 106.

Hab. Borneo.

argentocincta, id. *ibid.*, p. 107.

Hab. Borneo.

leucotelus, id. *ibid.*, p. 6.

Hab. Singapore.

plecioides, id. *ibid.*, p. 6.
Hab. Singapore.

vittifrons, id. *ibid.*, v, 1861, p. 144.
Hab. Amboina.

Genus CYLINDROTOMA.

Macquart, *S. à Buff. Dipt. Paris*, 1834, p. 107: *Limnobia*, pt. (anctor.)
albitarsis, Doleschall, *Naturk. Tijdschr. Nederl. Indie* xiv, *Batavia*, 1857, p. 391.
Hab. Java.

Genus PÆCILOSTOLA.

Loew, *Wien. Entom. Monatschr.*, vii, 1863, p. 222.
pallens, V. der Wulp, *Notes Leyden Mus.* vii, 1885, p. 13.
Hab. Java.

Genus TANYDERUS.

Philippi, *Verh. K. K. Z. B. Gesellsch. Wien*, 1865, p. 780.
ornatissimus, Ost.-Sacken, *Berlin. Ent. Zeitschr.* xxxi, 1887, p. 228: *Cylindrotoma*
id., Doleschall, *Natuur. Tijdschr. Nederl. Indie*, xvii, *Batavia*, 1858, p. 80.
Hab. Amboina.

Genus MEGISTOCERA.

Wiedem., *Auss. Europ. Zweifl. Ins.* i, *Hamm*, 1828, p. 55: *Nematocera*, *olim*
Wiedem., *Dipt. Exot.* i, 1821, p. 29: Macquart, Schiner, Ost.-Sacken,
etc.
fuscana, Wiedem. *loc. cit.* p. 55: *Nematocera*, id., Wiedem., *Dipt. Exot.* i, 1821.
Hab. Java.
verticalis, id. *ibid.*, p. 56.
Hab. Java
atra, Doleschall, *Naturk. Tijdschr. Nederl. Indie*, xvii, *Batavia*, 1858, p. 80.
Hab. Amboina.

Genus TRICHOCERA.

Meigen, *Illig. Magaz.* ii, 1803, p. 262: *Tipula*, pt. (anctor.): *Limonia*,
Latr., Macq., Walker, Schiner, Rondani, Ost.-Sacken, etc.
ocellata, Walker, *Ins. Saunders. Dipt. London*, 1856, p. 433.
Hab. India.

• Genus DICRANOMYIA.

Stephens *Catal. Brit. Ins.* 1829.

saltens, Ost.-Sacken, *Berlin. Ent. Zeitschr.* xxxi, 1887, p. 172: *Limnobia* id.
Doleschall, *Naturk. Tijdschr. Nederl. Indie*, xiv, *Batavia*, 1857, p. 390.
Hab. Java.

• Genus LIBNOTES.

Westwood, *Trans. Ent. Soc. London*, 1876, p. 505.

thwaitesiana, Westw. *loc. cit.* p. 505.
Hab. Ceylon.

Genus MONGOMA.

Westwood, *Trans. Ent. Soc. London*, 1881, p. 364.

simplex, Ost.-Sacken, *Ann. Mus. Civ. Genova*, xvi, 1881, p. 402.
Hab. Ternate.

pœcilopectera, id. *ibid.*, p. 403.
Hab. Sumatra.

aurantiaca, id. *Berlin. Ent. Zeitschr.* xxxi, 1887, p. 181: *Limnobia* id. Doleschall,
Naturk. Tijdschr. Nederl. Indie, Batavia, xvii, 1859, p. 78.
Hab. Java.

Genus TEUCHOLABIS.

Osten-Sacken, *Proceed. Acad. Nat. Sci. Philadelphia*, 1859, p. 222.

bicolor, Ost.-Sacken, *Ann. Mus. Civ. Genova*, xvi, 1881, p. 404.
Hab. Sumatra.

fenestrata, id. *Berlin. Ent. Zeitschr.* xxxi, 1887, p. 188.
Hab. Ceylon.

determinata, id. *ibid.*, p. 188.
Hab. Sula.

Family MYCETOPHILIDI.

J. Bigot, *ad hoc in ed.* 1891: *Mycetophilidæ*, Schiner, *Faun. Austr. Dipt.* ii,
Wien, 1864, p. 416: *Mycetophilides*, Westwood, *Introd. Modern Class.*
Ins. 1840: *Tipul. Fungicolæ*, Latr.: *Tipul. Fungivoræ*, Meigen.

Genus MYCETOPHILA.

Meigen, *Illig. Magaz.* ii, 1803, p. 263: *Sciara* pt. Fabr.

bimaculata, Walker, *List Dipt. Ins. Brit. Museum*, i, London, 1818, p. 100.
Hab. India.

pennipes, Osten-Sacken, *Berlin. Entom. Zeitschr.* xxxi, 1887, p. 204.
Hab. Borneo.

Genus PLATYURA.

Meigen, *Illig. Magaz.* ii, 1803, p. 264: *Sciophila*, pt. Meigen: *Sciara*, pt.
Fabr.: *Ceroptatus*, pt. Fabr., Latr.: *Asindulum*, pt. Latr.

aenusta, Walker, *Ins. Saunders. Dipt.* i, London, 1856, p. 421.
Hab. India.

Genus GLAPHYROPTERA.

Winnertz, *Verhandl. K. K. z. b. Gesellsch. Wien*, xiii, 1863, p. 781: *Leia* pt.
winthemi, Lehmann, V. der Wulp, *Reisen in Midden Sumatra* iv, *Natuurlijke
Histoire, Diptera*, p. 7.
Hab. Sumatra. Borneo?

Genus SCIOPHILA.

Meigen, *Syst. Besch.* i, 1818, p. 245: *Platyura*, pt. Fabr., Meig.: *Asindu-
lum*, pt. Latr.

tropica, Doleschall, *Naturk. Tijdschr. Nederl. Indie*, xiv, *Batavia*, 1857, p. 392.
Hab. Java.

Genus SCIARA.

Meigen, *Illig. Magaz.* ii, 1803, p. 263: *Tipula*, pt. Linn., Fabr.: *Hirtea*,
Fabr.: *Molobrus*, Latr.: *Planetes*, pt. Walker: *Planetella*, pt. Westwood.

femoralis, Doleschall, *Naturk. Tijdschr. Nederl. Indie*, xvii, *Batavia*, 1858, p. 78.
Hab. Amboina.

indica, Walker, *Ins. Saunders. Dipt.* i, London, 1856, p. 418.
Hab. India.

laticornis, id. *Journ. Proceed. Linn. Soc. London*, i, 1857, p. 105.
Hab. Borneo.

solita, id. *ibid.*, p. 105.
Hab. Borneo.

rufithorax, V. der Wulp, *Reisen in Midden Sumatra*, iv, *Natuurlijke Histoire, Dip-
tera*, p. 7.
Hab. Sumatra.

thomæ, id. *ibid.*, p. 7.
Hab. Sumatra.

sulcata, id. *Tijdschr. Entom.* xxx, 1886-87, p. 177.
Hab. Java.

• Family RHYPHIDI.

J. Bigot, *adhuc in ed.* 1891: Rhyphidae Schiner, *Faun. Austr. Flieg.* ii, Wien., 1864, p. xxiv.

• Genus RHYPHUS.

Latr. *Diction.* iv, 1804: Tipala pt. Scopoli: Sciara, Rhagio, pt. Fabr.: Anisopus (*olim*) Meigen.

maculipennis, V. der Wulp, *Notes Leyden Mus.* vii, 1885, p. 14.
Hab. Java.

• Family BIBIONIDI.

J. Bigot, *adhuc. in ed.* 1891: Bibionidae, Schiner, *Faun. Austr. Flieg.*, 1864, p. xvi: Bibionides, Westwood, *Introd. Mod. Class. Ins.* 1840: Tipulariae florales, Latr.: Tip. Musciformes et Latipennes Meig.

• Genus SCATHOPSE.

Geoffroy, *Hist. d. Ins.* ii, 1764, p. 545: Tipala pt. Linn., De Geer, Fabr.: Hirtea pt. Fabr. (*Meig. olim*): Ceria, Scopoli: Aspistes, Rnth. pt.

pusilla, Doleschall, *Naturk. Tijdschr. Nederl. Indie, x, Batavia*, 1859, p. 407.
Hab. Java.

• Genus PLECIA.

Wiedem., *Auss. Europ. Zweifl. Ins.*, i, *Hamm.* 1828, p. 72: Crapitula, pt. Gimmerth. *Bull. Moscou*, 1845: Penthetria, pt. Meig. *Illig. Magaz.* ii, 1803, p. 264.

melanaspis, Ost.-Sacken, *Ann. Mus. Civ. Genova*, xvi, 1881, p. 307: Penthetria, id. Wiedem., *Auss. Europ. Zweifl. Ins.* 1828, i, p. 72: Penth. japonica, id. *ibid.*, p. 618: Plecia ignicollis, Walker, *List. Dipt. Ins. Brit. Mus. London*, 1848, p. 116.
Hab. Nepal.

fulvicollis, V. der Wulp, *Tijdschr. Entom.* xxiii, p. 156: Hirtea id. Fabr., *Syst. Antl.* p. 53: Plecia id., Wiedem., *Auss. Europ. Zweifl. Ins.* p. 73: Penthetria id. Wiedem. *Dipt. Exot.* i, p. 31: Plecia id. Macquart, *S. à Buff. Dipt.* i, p. 176: Plecia id. Doleschall *Natuur. Tijdschr. Nederl. Indie, x, Batavia*, 1839, p. 407: Plecia, id. Rondani, *Ann. Mus. Civ. Genova*, vii, p. 462: Plecia dorsalis, Walker, *Journ. Proceed. Linn. Soc. London*, i, 1857, p. 5.
Hab. Sumatra and Malay Peninsula.

tergorata, Rondani, *Ann. Mus. Civ. Genova* vii, 1875, p. 462.
Hab. Borneo.

foreipata, Ost.-Sacken *Ann. Mus. Civ. Genova*, xvi, 1881, p. 397.
Hab. Sumatra.

subvarians, Walker, *Journ. Proceed. Linn. Soc. London*, i, 1857, p. 105.
Hab. Sarawak.

tristis, V. der Wulp, *Notes Leyden Mus.* vi, 1884, p. 251.
Hab. Java.

Genus BIBIO.

Geoffroy, *Hist. Ins.* ii, 1764, p. 571: Tipula pt. De Geer, Linn., Schin.
Hirtea, pt. Fabr., Panzer., Meig., (*olim*), Zetterst., etc.

bicolor, Walker, *List. Dipt. Ins. Brit. Mus.* pt. i, London, 1848, p. 121.
Hab. India.

rubicundus, V. der Wulp, *Notes Leyden Mus.* vi, 1884, p. 251.
Hab. Java.

Genus SIMULIUM.

Latreille, Buffon. *Hist. Nat. Crust. et Ins.* xiv, 1805, p. 294.

indicum, Becher *Journ. Asiat. Soc. Bengal*, liii, 1884, p. 199, pl. xiv.
Hab. Assam.

Division BRACHYCERATÆ.

J. Bigot, *adhuc ined.* 1891.

Sub-division EMPODIATA.

J. Bigot, *adhuc ined.* 1891.

Family TABANIDI.

J. Bigot, *adhuc ined.* 1891: Tabanii, Latr., 1802, Meigen, Macquart,
Zetterst., Loew: Tabanidæ, Leach, 1819, Schiner, 1862: Tabanides,
Latr. 1829: Tabanina, Tabaninæ, Rondani, 1846: Tabanidæ, West-
wood, Walker: Tabanidii, J. Bigot (*olim*): Anthracina, Rafinesque:
Sclerostoma, Dumèril: Tabanica, Burmeister: Tabanites, Newman.

Genus CHRYSOPS.

Meigen, *Illig. Magaz.*, ii, 1803, p. 267: Tabanus pt.

ligatus, Walker, *List. Dipt. Ins. Brit. Museum, Lond.*, i, 1848, p. 195.
Hab. Bengal.

terminalis, id. *ibid.*, p. 195.
Hab. India.

semicirculus, id. *ibid.*, p. 196.
Hab. India.

stimulans, id., *Ins. Saunders. Dipt. London*, i, 1856, p. 73.
Hab. India.

pellucidus, Fabr. *Syst. Antl.*, p. 113.
Hab. Tranquebar.

dispar, Fabr., *Entom. System. Suppl.*, p. 567.
Hab. India.

fasciatus, Wiedem., *Dipt. Exot.*, i, p. 103.
Hab. Java.

translucens, Macquart, *Dipt. Exot.* pt. i, 1838, p. 158.
Hab. Java.

flaviventris, id. *ibid.*, *Suppl.* i, 1846, p. 44.
Hab. India.

rufitarsis, id. *ibid.*, *Suppl.*, ii, 1847, p. 14.
Hab. Java.

fixissimus, Walker, *Journ. Proceed. Linn. Soc. London*, i, 1857, p. 112.
Hab. Borneo.

unizonatus, Rondani, *Ann. Mus. Civ. Genova*, vii, 1875, p. 459.
Hab. Borneo.

albicoinctus, V. der Wulp, *Tijdschr. Entom.*, xi, 1868, p. 103.
Hab. Salawatti.

striatus, id., *Notes Leyden Mus.*, vii, 1885, p. 79.
Hab. Amoy, China.

clavierus, Thomson, *Eugenie Resa, Stockholm*, 1858-68, p. 452.
Hab. Malacca.

Genus PANGONIA.

Latr., *Hist. Nat. Crust. et Ins.*, iii, 1794, p. 437: *Tanyglossa*, Meigen
Klassif. p. 174: *Tabanus* pt. Linn., Rossi, Fabr.: *Bombylinas*, pt. Olivier,
Encycl. Method.

amboinensis, Fabr., *Syst. Antl.*, p. 91.
Hab. Amboina, Himalayas.

longirostris, Walker, *List Dipt. Ins. Brit. Museum, London*, 1851. pt. v, *Suppl.*
p. 139.
Hab. India.

taprobanea, id. *ibid.*, *Addenda*, 1854, p. 324.
Hab. Ceylon.

rufa, Macquart, *Dipt. Exot. Suppl.* pt. iv, Paris, 1850, p. 18.
Hab. Bombay.

Genus SILVIUS.

Meigen, *Syst. Besch.* ii, 1820, p. 27, Wiedem., Macq.: *Tabanus* pt.

dimidiatus, V. der Wulp, *Tijdschr. Entom.*, xi, 1868, p. 102.
Hab. Salawatti.

Genus HÆMATOPOTA.

Meigen, *Illig. Magaz.* ii, 1803, p. 267: *Tabanus* pt.

cana, Walker, *List Dipt. Ins. Brit. Mus. London*, pt. i, 1848, p. 207.
Hab. Bengal.

atomaria, id. *Journ. Proceed. Linn. Soc. London*, i, 1857, p. 112.
Hab. Borneo.

irrorata, Macquart, *Dipt. Exot.* pt. i, 1838, p. 163.
Hab. Java.

lunulata, id. *ibid.*, *Suppl.* ii, *Suites*, 1847, p. 15.
Hab. Java.

roralis, Fabr. *Syst. Antl.*, p. 107.
Hab. Tranquebar, Madras Pr.

javana, Wiedem. *Ausser. Europ. Zweifl. Ins.* i, Hamm, 1828, p. 218.
Hab. Java.

cingulata, id. *ibid.*, p. 216.
Hab. Java.

borneana, Rondani, *Ann. Mus. Civ. Genova*, vii, 1875, p. 461.
Hab. Borneo.

pungens, Doleschall, *Natuur. Tijdschr. Nederl. Indie*, x, Batavia, 1856, p. 407.
Hab. Java.

oilipes, J. Bigot, *Nouv. Archives Mus. Paris*, (3) ii, 1890, p. 205.
Hab. Laos.

pachycera, id. *ibid.*, p. 206.
Hab. Laos.

macrocera, id. *ibid.*, p. 207.
Hab. Laos.

Genus RHINOMYZA.

Wiedemann, *Nov. Dipt. Gener.*, 1820, p. 8.*fusca*, Wiedem. *ibid.* p. 8.

Hab. Java.

Genus DITYLOMYIA.

J. Bigot, *Rev. et Mag. Zool. Guérin*, (2) xi, 1859, p. 305.*ornata*, J. Bigot, *ibid.* p. 306.

Hab. Ceylon.

Genus TABANUS.

Linn., *Faun. Suec.* ii, 1761, p. 462, et auct.: *Theriopectes* pt. Zeller, *Isis Agelanius*, pt. Rondani, 1863: *Atylotus*, pt. Ost.-Sacken, 1876.*auricinctus*, Macquart, *Dipt. Exot.* pt. i, Paris, 1838, p. 130.

Hab. Indian Archipelago.

pusillus, *id. ibid.*, p. 127.

Hab. China.

albilateralis, *id. ibid.*, p. 129.

Hab. Java.

cœrulescens, *id. ibid.*, p. 128.

Hab. Java.

basalis, *id. ibid.*, p. 126.

Hab. India.

brunneus, *id.*, S. à Buff. *Dipt.* i, Paris, 1834, p. 203.

Hab. Java.

sumatrensis, *id. ibid.*, p. 201.

Hab. Sumatra.

consanguineus, *id.*, *Dipt. Exot.* pt. i, 1838, p. 127.

Hab. Malabar.

servillei, *id. ibid.*, p. 128.

Hab. India.

rubicundus, *id. ibid.*, *Supplem.*, 1846, p. 32.

Hab. India.

dorsolinea, Wiedem., *Analect. Entomol.* p. 22: *T. striatus* ? Fabr.

Hab. India.

virgo, *id. ibid.*, p. 22.

Hab. India.

orientalis, *id. ibid.*, p. 21.

Hab. India.

- rubidus*, id., *Dipt. Exot.*, i, p. 69.
 Hab. Bengal.
- ardens*, id. *ibid.*, p. 78.
 Hab. Java.
- immanis*, id. *Ausser Europ. Zweifl. Ins.*, i, *Hamm*, 1828, p. 123.
 Hab. Java.
- hybridus*, id. *ibid.*, p. 557.
 Hab. Macao, China.
- striatus*, Fabr., *Entom. System.*, iv, p. 371: *T. dorsolinea*? *Wiedem.*
 Hab. China; Java.
- rufiventris*, id. *ibid.*, p. 96.
 Hab. India.
- javanus*, id. *Syst. Antl.*, p. 103.
 Hab. Java.
- tenebrosus*, Walker, *List Dipt. Ins. Brit. Mus. London*, v, *Suppl.* p. 242.
 Hab. Malabar.
- pyrausta*, id., *Zoologist*, vii, 1849, *Append.* p. lxxv.
 Hab. Java.
- vagus*, id. *ibid.*, *Append.* p. lxxviii.
 Hab. Hong-Kong.
- megalops*, id. *List Dipt. Ins. Brit. Mus. London*, v, *Suppl.* i, 1854, p. 247.
 Hab. Java.
- aurotestaceus*, id. *ibid.*, p. 247.
 Hab. Shanghai.
- orientis*, id., *List Dipt. Ins. Brit. Mus.*, pt. i, 1848, p. 152, and v, *Suppl.* i, 1854
Addenda, p. 328.
 Hab. Sikkim and Nepal.
- explicatus*, id., *ibid.* v, *Suppl.* i. 1854 *Addenda*, p. 328.
 Hab. Sikkim.
- univentris*, id. *ibid.*, i, 1848, p. 151.
 Hab. Borneo.
- auriflamma*, id. *ibid.*, p. 155.
 Hab. Silhet, Assam.
- inscitus*, id. *ibid.*, p. 161.
 Hab. Silhet, Assam.
- mentitus*, id. *ibid.*, p. 162.
 Hab. China.
- sinicus*, id. *ibid.*, p. 163.
 Hab. Hong-Kong.

- amœnus, id. *ibid.*, p. 163.
Hab. Hong-Kong.
- internus, id. *ibid.*, p. 164.
Hab. Silhet.
- jucundus, id. *ibid.*, p. 187.
Hab. Hong-Kong.
- albulus, id. *Ins. Saunders. Dipt. i, London, 1856, p. 46.*
Hab. India.
- pyrrhus, id. *ibid.*, p. 47.
Hab. India.
- albimediis, id. *ibid.*, p. 48.
Hab. India.
- tenens, id. *ibid.*, p. 49.
Hab. India.
- hilaris, id. *ibid.*, p. 49.
Hab. India.
- crassus, id. *ibid.*, p. 50.
Hab. India.
- vagus, id. *ibid.*, p. 50.
Hab. India, Java.
- rubiginosus, id. *ibid.*, p. 51.
Hab. India.
- umbrosus, id. *ibid.*, p. 52.
Hab. India.
- hirtus, id. *ibid.*, p. 52.
Hab. India.
- puella, id. *ibid.*, p. 53.
Hab. India.
- sanguineus, id. *ibid.*, p. 54.
Hab. Java.
- obconicus, id. *ibid.*, p. 54.
Hab. India.
- consocius, id. *ibid.*, p. 56.
Hab. India.
- perlina, id. *ibid.*, p. 56.
Hab. India.
- nigropictus, Macquart, *Dipt. Exot. Suppl., v, Paris, 1855, p. 24.*
Hab. India.
- yao, id. *ibid.*, p. 24.
Hab. North China.

clausicella, (*Bellardia*, Rondani), id. *ibid.*, p. 25.

Hab. China.

confucius, id. *ibid.*, p. 26.

Hab. China.

hoang, id. *ibid.*, p. 27.

Hab. China.

bubali, Doleschall, *Natuur. Tijdschr. Nederl. Indie, Batavia*, x, 1856, p. 407.

Hab. Java.

monoculus, id. *ibid.*, xvii, 1858, p. 85.

Hab. Java.

cinnamomeus, id. *ibid.*, xvii, 1858, p. 84.

Hab. Amboina.

furunculigenus, id. *ibid.*, xvii, 1858, p. 84.

Hab. Amboina.

abscondens, Walker, *Trans. Ent. Soc. London*, v, 1860, p. 275.

Hab. Burmah.

univentris, (*nomen bis lectum*) id. *ibid.*, i, 1857, p. 9.

Hab. Mt. Ophir; Borneo.

partitus, id. *ibid.*, i, 1857, p. 9.

Hab. Singapore.

nexus, id. *ibid.*, i, 1857, p. 110.

Hab. Borneo.

fumifer, id. *ibid.*, i, 1857, p. 110.

Hab. Borneo.

optatus, id. *ibid.*, i, 1857, p. 111.

Hab. Borneo.

simplicissimus, id. *ibid.*, i, 1857, p. 111.

Hab. Borneo.

serus, id. *ibid.*, vi, 1862, p. 20.

Hab. Ceram.

justorius, Rondani, *Ann. Mus. Civ. Genova*, 1875, p. 455.

Hab. Sarawak.

alboscutatus, id. *ibid.*, p. 456.

Hab. Sarawak.

pauper, id. *ibid.*, p. 456.

Hab. Sarawak.

ignobilis, id. *ibid.*, p. 457.

Hab. Sarawak.

- dives*, id. *ibid.*, p. 457.
Hab. Sarawak.
- fulvissimus*, id. *ibid.*, p. 458.
Hab. Sarawak.
- variegatus*, id. *ibid.*, p. 458.
Hab. Sarawak.
- apicalis*, id. *ibid.*, p. 459.
Hab. Sarawak.
- goniculatus*, V. der Wulp, *Midden Sumatra*, iv, *Natuurlijke Histoire, Diptera*, p. 16.
Hab. Sumatra.
- incultus*, id. *ibid.*, p. 17.
Hab. Sumatra.
- tristis*, id. *ibid.*, p. 17.
Hab. Sumatra.
- fumipennis*, id. *ibid.*, p. 18.
Hab. Sumatra.
- minimus*, id. *ibid.*, p. 18.
Hab. Sumatra.
- equestris*, id., *Notes Leyden Mus.*, vii, 1885, p. 77.
Hab. Sumatra, Java and Borneo.
- felderi*, id. *ibid.*, p. 78.
Hab. Ning-po, China.
- bucolicus*, Schiner, *Novara Reise*, 1868, p. 81.
Hab. Hong-Kong.
- nicobarensis*, id. *ibid.*, p. 81.
Hab. Nicobar Islands.
- mandarinus*, id. *ibid.*, p. 83.
Hab. Hong-Kong.
- administrans*, id. *ibid.*, p. 83.
Hab. Hong-Kong.
- ceylanicus*, id. *ibid.*, p. 93.
Hab. Ceylon.
- leucosparsus*, J. Bigot *N. Archives Mus. Paris*, (3) ii, 1890, p. 203.
Hab. Laos.
- nigrotectus*, (*Bellardia*, Rondani), id. *ibid.*, p. 204.
Hab. Laos.

melanognathus, (*Atylotus*, Ost.-Sacken), id. *ibid.*, p. 204.
Hab. Laos.

laotianus, (*Atylotus*, Ost.-Sacken), id. *ibid.*, p. 205.
Hab. Laos.

Family STRATIOMYDI.

J. Bigot, *adhuc in ed.*, 1891: Stratiomydæ, Latr. 1802: Xylophagei, Fall. 1810: Notachanta, Latr. 1817: id. Macquart, 1834: Notachanten, Brauer 1882: Stratiomydæ, Leach, 1819: id. Rondani, 1856: Stratiomydinæ, Rondani, 1856: Stratiomydes Latr. 1802: Xylophagi, Meigen, 1820: Xylophagidæ Steph., 1829: id. Rondani, 1856: id. Bigot: Xylophagina, Rondani, 1836: Xylophagi, Walker, 1848: Beridæ, Westwood, 1840: Beridina, Rondani, 1836: Cœnomydæ, Westwood, 1840: Cœnomyna, Rondani, 1836: Sicarii, Latr., 1825: id. Macq. 1834: Polychæta, Zetterstedt.

Genus XYLOPHAGUS.

Meigen, *Illig. Magaz*, ii, 1803, p. 266.

brunneus, Wiedem., *Auss. Europ. Zweifl. Ins.* pt. iv, Hamm, 1828, p. 85.
Hab. India.

Genus ANTIDOXION.

Sneller V. Vollenhoven, *Versl. en Med. K. Akad. Wetensch.* xv, 1863.

flavicorne, id. *ibid.*
Hab. Java.

Genus ACRASPIDEA.

Schiner *in lit.*, V. Brauer *Zweifl. K. Mus. Wien.*, 1882, p. 75-19.

felder, id. *ibid.*, p. 76-20.
Hab. Ceylon.

Genus SUBULA.

Meigen, *Syst. Besch.* ii, 1820, p. 15: Xylophagns, pt.

calopodata, J. Bigot, *Ann. Soc. Ent. France*, (5) ix, 1879, p. 195.
Hab. Ternate.

flavipes, Doleschall, *Natuur. Tijdschr. Nederl. Indie*, xvii, 1858, p. 85.
Hab. Amboina.

vittata, id. *ibid.*, p. 86.
Hab. Amboina.

inamcena, (*Solva*) Walker, *Journ. Proceed. Linn. Soc. London* iv, 1860, p. 98: Ost.-Sacken, *Ann. Mus. Genova*, xvi, p. 407.

Hab. Java; Celebes.

Genus THYLACOSOMA.

Schiner in lit., V. Brauer, *Zweifl. Mus. Wien*, 1882, p. 77: ? Ruba, pt. Walker,

amboinense, Schiner in lit., V. Brauer, *loc. cit.*, p. 77.

Hab. Amboina.

Genus ENGONIA.

Schiner, in lit., V. Brauer, *Zweifl. Mus. Wien*, 1882, p. 76: Negritomyia? J. Bigot, *Ann. Soc. Ent. France*, (5) ix, 1877, *Bulletin*, p. lxxiv.

aurata, Schiner, in lit., v. Brauer, *loc. cit.*, p. 76.

Hab. Amboina.

Genus NEGRITOMYIA.

J. Bigot, *Ann. Soc. Ent. France*, 1877, *Bulletin*, p. lxxiv.

bilineata, (*Stratiomys*) Fabr. *Syst. Antl.* p. 79: V. der Wulp, *Notes Leyd. Mus.* vii. p. 59.

Hab. Java.

Genus CLITELLARIA.

Meigen, *Illig. Magaz.*, ii, 1803, p. 265: Ehippium, Latr., 1802: Rhapsiocera pt. Macq.: Stratiomys, pt. Fabr. *Syst. Antl.*

heminopla, Wiedem., *Auss. Europ. Zweifl. Ins.* ii, *Hamm*, 1830, p. 48: id., *Zool. Mag.* iii, p. 30.

Hab. Tranquebar, Madras Pr.

varia, Walker, *List Dipt. Ins. Brit. Mus.*, pt. v, *Suppl. i*, London, 1854, p. 63.

Hab. Java.

angusta, id. *loc. cit.*, p. 62: Ehippium id., Macq. *S. à Buñ. Dipt.* i, 1834, p. 252.

Hab. Java.

tonebrica, id., *loc. cit.*, iii, 1849, p. 522.

Hab. Java.

notabilis, Walker, *Journ. Proceed. Linn. Soc. London*, i, 1857, p. 108.

Hab. Borneo.

bivittata, Wiedem., *Auss. Europ. Zweifl. Ins.*, ii, *Hamm*, 1830, p. 46: Stratiomys bilineata, Fabr. *Syst. Antl.*: Ehippium spinigerum? Dolesch.: Rhapsiocera spinithorax Macq. *Dipt. Ez.*

Hab. Java.

spinigerum, (*Ephippium*) Doleschall, *Natuur. Tijdschr. Nederl. Indie*, x, 1856,
p. 407: Engonia, id. Schin., Brauer.

Hab. Java.

nigerrimum, (*Ephippium*) Doleschall, *loc. cit.*, xvii, 1858, p. 81.

Hab. Java.

flaviceps, Walker, *Journ. Proceed. Linn. Soc. London*, i, 1857, p. 7.

Hab. Singapore.

Genus BERIS.

Latr. *Hist. Nat. Crust. et Ins.* iii, 1802, p. 340: Stratiomys, pt. Fabr., Fall. :
Actina, pt. Meig.

javana, Macquart, *Dipt. Exot.* i, pt. ii, Paris, 1838, p. 188.

Hab. Java.

Genus CULCUA.

Walker, *Journ. Proceed. Linn. Soc. London*, i, 1857, p. 109.

simulans, Walker, *ibid.*, p. 109.

Hab. Borneo, Malacca.

Genus CYCLOGASTER.

Macquart, *S. & Buff. Dipt.* i, Paris, 1834, p. 256.

radians, Walker, *Journ. Proceed. Linn., Soc. London*, i, 1857, p. 7.

Hab. Singapore.

Genus HERMETIA.

Latr. *Dict. Hist. Nat.* xxiv, p. 192, Fabr., Wied., Macq. : *Nemotelus*, pt.
Degeer. : *Bibio*, *Syrphus*, *Mydas*, pt. auctor. : *Xylophagus*, pt. Lamarek.

melanæsiæ, Bigot, *Ann. Soc. Ent. France*, (5) ix, 1879, p. 202: *H. batjanensis*,
V. der Wulp, *Tijdschr. Entom.*, xxiii, p. 161: *Massicyta corioides* Walker. ?

Hab. Indian Archipelago.

rufiventris, (*Nomen bislectum*) Walker, *Journ. Proceed. Linn. Soc. London*, v, 1861,
p. 145.

Hab. Amboina.

Genus PHYLLOPHORA.

Macquart, *Dipt. Exot.* i, pt. i, Paris, 1838, p. 178.

angusta, Walker, *Journ. Proceed. Linn. Soc. London*, i, 1857, p. 7.

Hab. Singapore.

Genus TINDA.

Walker, *Journ. Proceed. Linn. Soc. London*, iv, 1860, p. 101.

indica, V. der Wulp, *Notes Leyd. Mus.*, vi, 1885, p. 57: *Biartes* id. Walker, *Ins. Saunders. Dipt.*, i, London, 1856, p. 87: *Tinda modifera*, Walker, *Journ. Proceed. Linn. Soc. London*, iv, 1860, p. 101: *Phyllophora bispinosa*, Thomson, *Eugenie Resa*, p. 454.

Hab. Java, Manilla.

Genus EVAZA.

Walker, *Journ. Proceed. Linn. Soc. London*, i, 1857, p. 109.

bipars, Walker, *ibid.*, p. 110.

Hab. Borneo.

flavipes, J. Bigot, *Ann. Soc. Ent. France*, (5) ix, 1879, p. 219.

Hab. India.

argyroceps, id. *ibid.*, p. 219.

Hab. The Moluccas.

fulviventris, id. *ibid.*, p. 220.

Hab. The Moluccas.

Genus NERNA.

(*nec Nerua*) Walker, *Journ. Proceed. Linn. Soc. London*, iii, 1859, p. 81.

mollis, Oat.-Sacken, *Ann. Mus. Civ. Genova*, xvi, 1881, p. 415.

Hab. Sumatra.

Genus PTILOCERA.

Wiedem., *Auss. Europ. Zweifl. Ins.* ii, Hamm, 1830, p. 58: *Stratiomys*, pt.

quadridentata, Wiedem., *loc. cit.*, p. 59.

Hab. Java.

continua, Walker, *Ins. Saunders. Dipt. London*, 1856, pt. i, p. 84.

Hab. Java.

smaragdina, Snellen v. Vollenhoven, *Mem. Entomol. Pays-Bas*, i, 1853, p. 92.

Hab. Celebes.

amethystina, id. *ibid.*, p. 92.

Hab. Amboina, Java and Celebes.

fastuosa, Gerstaecker, *Linn. Ent.* xi, 1857, p. 333.

Hab. Ceylon.

Genus WALLACEA. •

Doleschall, *Naturk. Tijdschr. Nederl. Indie*, xvii, Batavia, 1858, p. 82.

argentea, Doleschall, *loc. cit.*, p. 82. •

Hab. Amboina.

Genus ACRASPIDEA.

Brauer, *Zweift. Kais. Mus. Wien*, 1882, p. 19.

felderi, Brauer, *loc. cit.*, p. 20.

Hab. Ceylon.

Genus ACANTHINA.

Wiedem., *Auss. Europ. Zweift. Ins. ii, Hamm*, 1830, p. 50.

azurea, Gerstaecker, *Linn. Ent.*, xi, 1857, p. 335.

Hab. Ceylon.

Genus ELASMA.

Jaenicke, *Neu. Exot. Dipter. Frankfurt*, 1867, p. 14.

acanthinoidea, Jaenicke, *loc. cit.*, p. 15.

Hab. Java.

Genus PACHYGASTER.

Meigen, *Illig. Magaz.*, ii, 1803, p. 266: Nematelus, pt. Panzer: Sargus, pt.

Fall.: Vappo, pt. Latr.

rustitarsis, Macquart, *Dipt. Exot. Paris, Suppl.* 1846, p. 57.

Hab. Pondicherry.

Genus TOXOCERA.

Macquart, *Dipt. Exot. Suppl. iv, Paris*, 1850, p. 44.

limbiventris, Macq., *loc. cit.*, p. 45.

Hab. Java.

Genus MASSICYTA.

Walker, *Journ. Proceed. Linn. Soc. London*, i, 1857, p. 8.

bicolor, Walker, *loc. cit.*, p. 8.

Hab. Singapore.

Genus RHACHICERUS.

Haliday, *List Dipt. Ins.* 1848, p. 154.

zonotus, Ost. Sacken, *Ann. Mus. Genova*, xvi 1881, p. 408.

Hab. Sumatra.

Genus STRATIOMYS.

Geoffr. *Hist. Nat. Ins.* ii, 1764: Hirtea, pt. Scopoli: *Hoplomyia*, Zeller, Loew.

barca, Walker, *List Dipt. Ins. Brit. Museum, London*, 1849, p. 530.
Hab. China.

garatas, id. *ibid.*, p. 532.
Hab. China.

apicalis, id. *ibid.*, Part v, 1854, p. 53.
Hab. North China.

tutatus, id. *ibid.*, 1849, p. 532.
Hab. Malacca.

solennis, id., *Ins. Saunders. Diptor. London*, i, 1856, p. 79.
Hab. India.

inanimis, id., *Trans., Ent. Soc. London*, iv, 1857, p. 121.
Hab. China.

viridana, Wiedem., *Auss. Europ. Zweifl. Ins.*, ii, *Hamm*, 1830, p. 66.
Hab. Bengal.

minuta, Fabr., *Entom. System.*, iv, p. 268.
Hab. Tranquebar, Madras Pr.

pusilla, id. *ibid.*, p. 271
Hab. Tranquebar, Madras Pr.

rufipennis, Macquart, *Dipt. Exot. Suppl. v. Paris*, 1855, p. 42.
Hab. South China.

flavoscutellata, V. der Wulp, *Notes Leyd. Mus.* vii, 1885, p. 60.
Hab. Java.

Genus ODONTOMYIA.

Meigen, *Klassific.* i, 1804, p. 123: *Stratiomys*, pt. apud V. der Wulp.

viridana, Wiedem., *Analect. Entom.*: Macquart *Dipt. Exot. Suppl.*, iv, 1850, p. 48.
Hab. Bengal.

consobrins, Macquart, *Dipt. Exot., Suppl.*, ii, *Paris*, 1847, p. 16.
Hab. Java.

diffusa (*Stratiomys*) Walker, *List Dipt. Ins. Brit. Mus.*, part v, *London*, 1854, p. 53.
Hab. Java.

mutica, V. der Wulp, *Notes Leyd. Mus.* vii, 1885, p. 62.
Hab. Ternate.

Genus CAMPEPROSOPA.

- . Macquart, *Dipt. Exot. Suppl.*, iv, 1850, p. 46.
flavipes, Macquart, *loc. cit.*, p. 46.
Hab. Java.
munda, Osten-Sacken, *Ann. Mus. Civ. Genova*, 1881, p. 409.
Hab. Sumatra.

Genus CHRYSOCHLORA.

- Latr. *Fam. Natur.*, 1825, Macquart: Sargus, pt.
baccoides, Rondani, *Ann. Mus. Civ. Genova*, vii, 1875, p. 454.
Hab. Sarawak.
vitripennis, Doleschall, *Natuur. Tijdschr. Nederl. Indie.*, x, 1856, p. 408.
Hab. Java.

Genus MICROCHRYZA.

- Loew, *Verhandl. K. K. z. b. Gesellsch. Wien*, 1855, p. 146: Sargus, pt.
gemma, J. Bigot, *Ann. Soc. Ent. France*, (5) ix, 1879, p. 231.
Hab. Ceylon.

Genus TRICHOCHETA.

- J. Bigot, *Ann. Soc. Ent. France*, (5) ix, *Bullet.* 1878 p. xxii, et errata.
nemoteloides, *id. ibid.*, p. xxii.
Hab. Ternate.

Genus RAPHIOCERA.

- Macquart, *S. à Buff. Dipt.*, i, 1834, p. 253.
spinithorax, Macquart, *Dipt. Exot. Suppl.* ii, Paris, 1847, p. 17: Stratiomys bilineata, Fabr.: Clitellaria bivittata Wiedem.: Ehippium spinigeram, Doleschall:
Hab. Java.

Genus PTECTICUS.

- Loew, *Verhandl. K. K. z. b. Gesellsch. Wien.*, v, 1855, p. 142.
apicalis, Loew, *loc. cit.*, p. 142.
Hab. Pinang.
cingulatus, *id. ibid.*, p. 143.
Hab. Pinang.

australis, Schiner, *Novara Reise*, 1868, p. 65.
Hab. Nicobar Islands.

illustris, id. *ibid.*, p. 65.
Hab. Nicobar Islands.

apicalis, (nom. bisectum), V. der Wulp, *Notes Leyden Museum*, vii, 1885, p. 62.
Hab. Sumatra, Borneo.

brevipennis, V. der Wulp, *loc. cit.*, p. 63: Sargus ♂. Rondani, *Ann. Mus. Civ. Genova*, vii, 1875, p. 454.
Hab. Borneo, Java.

infascia, V. der Wulp, *loc. cit.*, p. 64: Sargus, id. Walker *Journ. Proceed. Linn. Soc. London*, i, 1857, p. 110.
Hab. Borneo, Java.

Genus SARGUS.

Fabr. *Entom. System. Suppl.*, 1798, p. 566: Rhagio pt. Schranck: Nematolus pt. Degeer.

gemmifer, Walker, *List Dipt. Ins. Brit. Mus.* iii, London, 1849, p. 516.
Hab. Sylhet, Assam.

tenebrifer, id. *ibid.*, p. 517.
Hab. China.

aurifer, id. *ibid.*, v. *Suppl.* i, London, 1854, p. 96.
Hab. India, China.

luridus, id. *Journ. Proceed. Linn. Soc. London*, i, 1857, p. 8.
Hab. Singapore.

quadrifasciatus, id. *ibid.*, v, 1861, p. 146.
Hab. Amboina.

metallinus, Fabr., *Syst. Antl.*, p. 258: V. der Wulp, *Notes Leyd. Mus.* vii, 1885, p. 65.
Hab. Bengal; Java.

longipennis, Wiedem., *Analect. Entomol.*, p. 31.
Hab. Java.

insignis, Macquart, *Dipt. Esot. Suppl.*, v, Paris, 1855, p. 16.
Hab. China.

viridiceps, id. *ibid.*, 1856, p. 110: Ptecticus id. V. der Wulp.
Hab. China.

pallipes, Bigot, *Ann. Soc. Ent. France*, (5) xi, 1879, p. 222.
Hab. Ceylon.

magnificus, id. *ibid.*, p. 222.
Hab. Assam.

rufus, Doleschall, *Naturk. Tijdschr. Nederl. Indie*, xvii, Batavia, 1858, p. 83.
Hab. Amboina.

ferrugineus, id. *ibid.*, p. 83.
Hab. Amboina.

formicæformis, id. *ibid.*, xiv, 1857, p. 403 : Sarg. metallinus, V. der Wulp.
Hab. Amboina.

leoninus, Rondani, *Ann. Mus. Civ. Genova*, vii, 1875, p. 454.
Hab. Sarawak.

brevipennis, id. *ibid.*, p. 454 : Ptecticus id., V. der Wulp.
Hab. Sarawak.

lætus, V. der Wulp, *Notes Leyden Mus.* vii, 1885, p. 66.
Hab. Sumatra.

rubescens, id. *ibid.*, p. 67.
Hab. Gorontalo.

Genus CHRYSOMYIA.

Macquart, *S. à Buff. Dipt. Paris*, 1834, p. 262 : Sargus, pt. et Nemotelus,
pt. Degeer.

flaviventris, Wiedem., *Analect. Entom.*, p. 31 : Sargus id., Macquart, *Dipt. Exot.* :
Microchryza id., Ost.-Sacken, 1881.
Hab. India, Java.

affinis, id. *ibid.*, p. 31 : Sargus id., Macquart, *Dipt. Exot.*
Hab. India.

Genus EUDMETA.

Wiedem., *Auss. Europ. Zweifl. Ins. Hamm*, 1830, p. 43 : Hermetia, pt.

marginata, Wiedem., *loc. cit.*, p. 45 : Hermetia, id. Fabr., *Syst. Antl.* p. 63 : Hermetia
cingulata (vel cingulalis,) Guerin, *Icon. Regn. Anim.*, p. 543.
Hab. Amboina, Java.

Family NEMESTRINIDI.

J. Bigot, *adhuc inæd.* : Nemestrinidæ, Macquart, *S. à Buff. Dipt.*, i, Paris,
1834, p. 370 : Anthracii et Bombyliarii, Latr. : Nemestrinidæ, Schiner
Faun. Austr., Flieg., i, 1862, Wien., p. 44.

Genus NEMESTRINA.

Latr. *Hist. Nat. Crust. et Ins.* xiv, 1804, p. 319.

javana, Macquart *Dipt. Exot.* ii, pt. i, Paris, 1840, p. 17.
Hab. Java.

Genus COLAX.

Wiedem., *Analecta Entomol.*, 1824.

javanus, Wiedem., *loc. cit.*, 1824, p. 18.
Hab. Java.

variegatus, Westwood, *Cabinet of Orient. Entomol.*, London, p. 38.
Hab. China.

Family LEPTIDI.

J. Bigot, *adhuc ined.*: Leptides, Meigen, *Syst. Besch.* ii, 1820: Leptidæ, Westw., *Introd. Entom.*, 1840: Schiner, *Faun. Austr. Flieg.* i, Wien, 1862, p. 170.

Genus LEPTIS.

Fabr., *Syst. Antl.*, 1806, p. 9: Nemetelus, pt. Degeer: Rhagio, pt. Latr., Olivier, Walker.

decisa, Walker, *Journ. Proceed. Linn. Soc. London*, i, 1857, p. 15.
Hab. Malacca.

uniguttata, Ost.-Sacken, *Ann. Mus. Civ. Genova*, xvi, 1881, p. 422.
Hab. Sumatra.

Genus HELIOMYIA.

Doleschall, *Naturk. Tijdschr. Nederl. Indië*, xiv, Batavia, 1857, p. 402, Chrysopyla? pt.

ferruginea, Doleschall, *loc. cit.*, p. 402.
Hab. Amboina.

Genus ATHERIX.

Meigen, *Illig. Magaz.*, 1803, p. 271: Rhagio pt. Fabr.: Bibio, pt. Fabr.: Leptis, pt. Fabr., Fall.: Ibisia, pt. Rondani.

nigritarsis, Doleschall, *Naturk. Tijdschr. Nederl. Indië*, xvii, Batavia, 1858, p. 92.
Hab. Amboina.

labiatus, Bigot, *Bull. Soc. Zool. France*, xii, Paris, 1887, p. 21.
Hab. Ceylon.

Genus CHRYSOPILA.

Macquart, *Dipt. Nord. France*, i, 1827: Leptis, pt. Meigen: Rhagio, pt. Latr., Walker: Atherix, pt. Fabr.

maculipennis, Walker, *Journ. Proceed. Linn. Soc. London*, i, 1857, p. 118.
Hab. Borneo.

lupina, Ost.-Sacken, *Ann. Mus. Civ. Genova*, 1881, p. 420.
Hab. Sumatra.

uniguttata, id. *ibid.*, p. 422.
Hab. Sumatra.

ferruginosa, id. *ibid.*, p. 419; *Leptis* id., *Wiedem. Zool. Magaz.*, iii, p. 4.
Hab. Java. and N. Ceram.

insularis, Schiner, *Novara Reise*, 1868, p. 119.
Hab. Nicobar Islands.

Genus SURAGINA.

Walker, *Journ. Proceed. Linn. Soc. London*, iv, 1860, p. 110.
signipennis, id. *ibid.*, vi, 1862, p. 8.
Hab. Ternate.

Family CYRTIDI.

J. Bigot, *adhuc ined.*: *Vesiculosidæ*, Bigot (*olim*): *Acrocera*, Meigen, 1803:
Acroceridæ, Leach, 1819: *Acrocerinæ*, Zetterst., 1842: *Inflatæ*, Latr.,
1809: *Cyrtidæ*, *Oncodinæ*, Rondani, 1846.

Genus ONCODES.

Latr. *Precis*, 1796, p. 154: *Henops*, Illiger, 1798, (*alias Oggeodes*)
costalis, (*Henops*), Walker, *Ins. Saunders. Dipt. London*, i, 1856, p. 203.
Hab. India.

INDEX.

Names of New Genera and Species have an asterisk (*) prefixed.

- Abaratha angulatus*, 196
 * " *hypobides*, 195
 " *sura*, 195
 " *syrichthus*, 195
Abelmoschus moschatus, 45
Ablabes, 234
 " *calamaria*, 234
 " *collaris*, 235
 " *doriae*, 235
 " *modestus*, 235
 " *nicobarensis*, 231
 * " *Stoliczka*, 231, 234, 250
 " *scriptus*, 231
Abroma, 58, 89
 " *angulata*, 89
 " *angusta*, 89
 " *fastuosum*, 89
 " *Wheeleri*, 89
Abrus precatorius, 294, 296, 310, 353, 401, 402
 " *pulchellus*, 296, 310, 353, 401
Abutilon, 39, 42
 " *asiaticum*, 42
 " *indicum*, 42
 " *populifolia*, 43
Acacia, 347
 " *concinna*, 312, 353, 403
 " *horrida*, 3
 " *pennata*, 312, 353, 401, 402
 " *rubricaulis*, 296
 ACANTHACEÆ, 322, 345, 349
Acanthina, 276
 " *azurea*, 276
Acanthophora Thierii, 341, 359, 378
Acesina, 33
 " *ammon*, 34
 * " *ammonides*, 34
 * " *ariel*, 33, 34
 " *paraganesa*, 33
 * " *zephyretta*, 33
Achyranthes aspera, 355, 371, 377
 " *var. porphyristachya*, 324, 356, 371
 " *aspera*, *var. typica*, 324
 " *porphyristachya*, 381, 383
 ACREIDÆ, 11, 13
Acraspidea, 272, 276
 " *felderi*, 272, 276
- ACROCERA, 282
 ACROCERIDÆ, 282
 ACROCERINÆ, 282
Acronodia, 121, 140
 " *punctata*, 139, 140
Acrostichum appendiculatum, 294, 316, 358, 388, 389, 394
 " *appendiculatum*, *var. setosa*, 337
 " *scandens*, 293, 337, 343, 345, 358, 389
Actina, 274
Adenanthera pavonina, 312, 353, 403
Adenostemma viscosum, 317, 354, 369, 372, 377, 380, 383
Adiantum lunulatum, 294, 336, 358, 389
Adolias, 9
Aecidium, 224, 225, 226, 227, 228, 229
 " *Apocyni*, 228
 " *Aquilegia*, 226
 " *clematidis*, 227
 * " *Cunninghamianum*, 224, 225, 230
 * " *flavescens*, 226, 230
 " *Hualtatinum*, 226
 " *Mespili*, 225
 * " *mori*, 225, 226
 * " *orbiculare*, 227, 230
 " *otagensis*, 227
 " *sclerothecium*, 226
Aegiceras, 293, 379
 " *majus*, 318, 354, 381
Acridis multiflorum, 328, 356, 380
Afghania australi, 205
Aganosma, 390
Agarista, 189
 AGARISTIDÆ, 14
Agelanius, 267
Ageratum conyzoides, 317, 354, 369, 377
Aglais andamanica, 295, 306, 352, 397
Agrimonia Eupatorium, 229
Agropyrum, 212
Alangium, 396
Albizzia Lobbek, 312, 353, 401, 402
 " *procera*, 295, 312, 353, 401, 402
 ALGÆ, 287, 300, 341, 343, 344, 348, 350, 378, 406
Allarithmia, 257

- Allophylus* Cobbe, 292, 308, 352, 382, 397
Allotinus, 29
 " *aphacha*, 29
 " *Horsfieldii*, 29
Allocasia, 296
 " *foenicata*, 294, 333, 346, 357, 404
Alsodicia bengalensis, 294, 296, 302, 340, 346, 351, 403
Alysicarpus vaginalis, 310, 353, 368, 377
 AMARANTACEÆ, 324, 345, 350
 AMARYLLIDACEÆ, 329, 345, 350
Amathusia amythaon, 8, 25
 " " *var. insularis*, 25
 " *dilucida*, 25
 " *potheus*, 25
 " *Westwoodii*, 25
Amblycephalus carinatus, 248
 " " *macularius*, 231, 248
 " " *modestus*, 231
 " " *monticola*, 247, 248
Amora Rohituka, 306, 352, 403
Anorphophallus, 297, 298, 333, 346, 357, 396, 398
 " *bulbifer*, 298, 333
 " *tuberculiger*, 333
 AMPELIDÆ, 307, 345, 349
 **Amphigynas*, 199
 " " *multipes*, 199
Ampittia mara, 196
 ANACARDIACEÆ, 308, 345, 349
Ancistroides, 143
Andropogon contortus, 295, 300, 336, 345, 358, 389, 394
Anilema, 396
 " *ovatum*, 330, 357, 401
Angelica glauca, 215
 ANGIOSPERMÆ, 344
Anisomeles ovata, 297, 323, 355, 370, 372, 377
Anisopus, 263
Ankyroderma Danielsseni, 202
 " *Marszelleri*, 203
Anodendron, 347
 " *paniculatum*, 296, 319, 354, 387
 ANONACEÆ, 38, 301, 345, 349
Anopheles, 251
 " *annularis*, 251
 " *barbistrotris*, 251
 " *sincensis*, 251
Anosia, 9
 ANTHRACI, 280
 ANTHRACINA, 264
Antiaris toxicaria, 328, 356, 398
Antidoxion, 272
 " *flavicorne*, 272
Antitaxis calocarpa, 293, 296, 301, 351, 403
Apatura, 9
 APATURIDÆ, 11, 12, 13, 25, 170
 APHNEINÆ, 27, 34, 179
Apium graveolens, 215
Aplopeltura lca, 248
 APOCYNÆÆ, 318, 344, 345, 349
 APODA, 202
 **Apodogaster*, 202
 " *Alcocki*, 202
Aporosa villosula, 326, 356, 404
Appias, 190
 " *albina*, 190
 " *lippo*, 29
 " *lankapura*, 190
Appias lycida, 29, 190
 " *paulina*, 190
 " (*Saletara*) *nathalia*, 190
Aquilegia vulgaris, 226
Ardisia, 288
 " *humilis*, 296, 297, 318, 354, 381, 382, 397
Argyranis, 13
 " *Childrenii*, 34
Argyria Hookeri, 297, 320, 355, 403
 " *lanccolata*, 320, 355, 403
 " *titiaefolia*, 297, 320, 355, 379, 381
Arhopala ainea, 32
 " *amantes*, 34, 179
 " *anarte*, 34
 " *araxes*, 179
 " *bazalus*, 34
 " *belphæbe*, 34
 " *camdeo*, 34
 " *centaurus*, 34
 " *khamti*, 32, 38
 " *ramo*, 34
 " *singla*, 34
 " *teesta*, 34
Aristolochia tagala, 325, 356, 88, 389, 390
 ARISTALOCHIACEÆ, 324, 345, 350
 AROIDEÆ, 333, 345, 350
Arrhenica, 257
Arrhenothrix, 35
Artocarpus Gomeziana, 296, 328, 356, 398
 ASCLEPIADACEÆ, 345, 349
 ASCLEPIADEÆ, 319
Asindulum, 262
Asparagus racemosus, 330, 357, 393
Aspistes, 263
Astilbe rivularis, 222
Atella alcippe, 26
 " *phalanta*, 26, 171
 " *sinha*, 171
Aterica, 9
Atheriz, 281
 " *labiatus*, 281
 " *nigritarsis*, 281
Athyma, 14
 " *amhara*, 176
 " *karita*, 175
 " *nefte*, 176
 " *perius*, 175
 " *venilia*, 176
Atolophlebia fasciatus, 409

- ATTIDE, 2, 4**
Atylotus, 267
 " *laetianus*, 272
 " *melanognathus*, 272
Aticennia, 288, 293, 299, 301, 313, 379
 " *officinalis*, 293, 307, 323, 355, 381
Azanza acuminata, 48
Badacara, 167
 " *nilgiriensis*, 167
Badamia exclamationis, 195
 " *Bahora chrysea*, 23
Balanopteris Tothila, 80
Bambusa gigantea, 374
Bamia betulifolia, 45
 " *chinensis*, 45
 " *multiformis*, 45
Baoris, 196
Barclaya longifolia, 301
Barringtonia racemosa, 314, 354, 380
 " *speciosa*, 292, 298, 314, 353, 380
Belenois coronca, 189
 " *java*, 189
 " *mesentina*, 189
Bellardia clausiella, 270
 " *nigrotectus*, 271
Benthodytes, 201, 202
 " *gelatinosa*, 200
 " *ocalis*, 200
 " *papillifera*, 200
 " *sanguinolenta*, 200
BERIDE, 272
BERIDINA, 272
Beris, 274
 " *javana*, 274
Berrya, 96, 108
 " *Ammonilla*, 108, 305, 352, 403
Biartes indica, 275
Bibio, 264, 274, 281
 " *bicolor*, 264
 " *rubicundus*, 264
BIBIONIDE, 263
BIBIONIDES, 263
BIBIONIDI, 263
BIGNONIACEÆ, 322, 345, 349
Bindohara phocides, 27
 " *sugriva*, 27
Binnindykia trichostylis, 95
Bisagrewia nicobarica, 119
Blachia andamanica, 297, 326, 340, 356, 384, 404
Blatogona, 12
Blumea virens, 297, 317, 345, 354
Blythia reticulata, 231
Barhaaria repens, 297, 324, 345, 355, 370, 372, 377, 378, 381, 383, 393, 394
BOMBACEÆ, 39
Bombax, 39, 48, 295, 303, 304, 390, 395
 " *ceiba*, 49
 " *festivum*, 48
 " *heptaphylla*, 49
 " *heterophyllum*, 304
 " *insigne*, 48, 49, 295, 304, 340, 351, 388
 " *insigne*, var. 303
 " *malabaricum*, 48, 49, 295, 304
 " " var. *albiflora*, 49
 " " " *albiflorum*, 304
 " *orientale*, 50
 " *pentandrum*, 50
BOMBYLIARFI, 280
Bombylius, 265
BORAGINEÆ, 229, 319, 345, 349
Boschia, 39, 54
 " *Griffithii*, 55
Bovista lilacina, 339, 358, 389
BRACHYGERATE, 264
Bragantia tomentosa, 324, 376, 404
Briedelia, 292
 " *Kurzii*, 326, 356, 397
 " *tomentosa*, 326, 356, 397
Brownlowia, 96, 97
 " *clata*, 98, 99
 " *Kleinhovioidea*, 97
 " *lanceolata*, 97
 " *macrophylla*, 97, 99.
 " *peltata*, 99
 " *Scortechinii*, 97, 98
BROWNLOWIÆ, 96
Bruguiera, 293, 299, 301
 " *gymnorhiza*, 293, 313, 353, 380
Bryum coronatum, 338, 358, 389
Buettneria, 59, 89
 " *andamanensis*, 90, 93, 305, 352, 393, 394, 403
 " *aspera*, 90, 92
 " *Curtisii*, 89, 90
 " *elliptica*, 90, 91
 " *grandifolia*, 93
 " *hypoleuca*, 94
 " *Jackiana*, 90, 92
 " *lanceifolia*, 90
 " *Maingayi*, 90, 91
 " *nepalensis*, 93
 " *uncinata*, 90, 91
BUETTNERIÆ, 58
Buuca virens, 388
Bungarus bungaroides, 246
 " *caeruleus*, 246
 " *lividus*, 246
BURSERACEÆ, 306, 345, 349
Buxus sempervirens, 213
BYBLIADÆ, 170
Cacoma Mori, 226
Cassalpinia Bonducella, 292, 297, 311, 353, 380
 " *Nuga*, 293, 296, 312, 353, 380
Calamaria catenata, 231, 233
 " *stalknechti*, 231
 " *sumatrana*, 233
Calamus, 294, 296, 347

- Calamus andamanicus*, 332, 357, 404
 " *tigrinus*, 332, 357, 404
Calanthe, 294, 388, 394
 " *veratrifolia*, 294, 356, 388, 389, 394
Caleosporium, 222
Calladium, 6
Calliana pieridoides, 17
 CALLIDULIDÆ, 15
Calliptax, 15, 16, 17, 22
 " *hyems*, 160
 " *mazares*, 22
 * " *sambavana*, 160
 * " *sumbana*, 159, 160
Callophis nigrescens, 245
Calanus nicobarica, 394, 399
Calophyllum inophyllum, 302, 351, 380
Calothrix pulvinata, 342, 359, 378
 CALYCIFLORE, 309, 344
Calympetes Dozyanum, 338, 358, 389
Calysisme, 169
 " *perseus*, 168
Camellia theifera, 340
Campeprosopa, 278
 " *flavipes*, 278
 " *munda*, 278
Camponotus micans, 3
Canarium, 382, 396
 " *commune*, 295
 " *euphyllum*, 306, 352, 403
Canavalia obtusifolia, 292, 311, 353, 379, 380
Cansjera Rheedii, 306, 352, 397
 CAPPARIDÆ, 302, 345, 349
Capparis, 347
 " *oxyphylla*, 294, 302
 " *scpiaria*, 293, 294, 296, 397
 " " *var. grandifolia*, 302, 351
 " *tenera*, 296, 397
 " " *var.*, 294
 " " *latifolia*, 302, 351
Capsicum minimum, 321, 355, 370, 372, 377
Carapa, 379
 " *moluccensis*, 298, 306, 352, 380
 " *obovata*, 306
Cardisoma, 292
Carica, 345
 " *Papaya*, 315, 321, 354, 369, 372, 377
Carpophaga bicolor, 327, 394, 399, 402
 " *insularis*, 399
Caryota sobolifera, 296, 331, 357, 381, 384
 CASSIDA, 7
Cassytha, 290, 292
 " *filiformis*, 325, 356, 381
Castalius ethion, 181
 " *rosimon*, 181
 " *roxus*, 181
Casuarina, 374
Casuarina cuissetifolia, 373, 374
Catochryps, 182
 " *cucius*, 27, 28, 181
 " *kandarpa*, 28
 " *padava*, 27, 181
 " *strabs*, 27, 28, 181
 " " *var. litharyyria*, 27
Catopha, 190
Catopsis, *Catilla*, 191
 " *crocale*, 191
 " *evangelina*, 191
 " *pyranthe*, 191
 " *scylla*, 191
Caulerpa clavifera, 342, 359, 378
 " *plumaris*, 342, 359, 378
 CECIDOMYDÆ, 253
 CECIDOMYDI, 253
Cecidomyia, 253, 254
 " *deferenda*, 254
 " *oryzæ*, 254
 CECIDOMYIDA, 253
Ceiba pentandra, 50
Celanthe veratrifolia, 328
 CELASTRINÆ, 306, 345, 349
 CELLULARES, 338, 344
Celosia cristata, 324, 355, 371, 372, 377
 CENOMYNA, 272
Ceratopogon, 252
 " *agas*, 252
 " *trichopus*, 252
Ceratopteris thalictroides, 294, 337, 346, 358, 388, 392
Cerbera Odollam, 318, 354, 381
Cerberus rhynchops, 244
Cervus muntjac, 148
Ceria, 263
Ceriops, 293, 299, 301
 " *Candolleana*, 313, 353, 380
 " *Roxburghiana*, 313, 353, 380
Ceroctena, 254
Ceroplastus, 262
Cethosia, 11, 12, 13
 " *hypsæ*, 171
 " *hypsina*, 14
 " *Lamarckii*, 171
 " *Leschenaultii*, 171
 " *penthesilea*, 171
 * *tambora*, 171
Champercia Griffithiana, 298, 325, 356
Chapra mathias, 32, 196
Chara, 299, 388
 " *fatida*, 334, 337, 358, 392
 CHARACEÆ, 337, 344
Charaxes, 158, 174
 " *athamas*, 174
 " *endamippus*, 174
 " *pyrrhus*, 174
Chartacalyx, 106
 " *acrescens*, 106
Charus helenus, 192
 " " *var. enganius*, 31

- Chickrassia tabularis*, 306, 352, 403
Chilades putli, 181
 " *trochilus*, 181
 CHIRONOMIDES, 252
Chironomus, 252, 253
 " *cubiculorum*, 253
 " *pictus*, 253
 " *socius*, 253
 " *venustus*, 252
 " *vicarius*, 253
 CHIRONOMIDÆ, 252
 CHIRONOMIDI, 252
 **Chittira orientis*, 166
Chonemorpha, 347
 " *macrophylla*, 319, 354, 389,
 391
Chrysobalanus, 115
Chrysochlora, 278
 " *baccoides*, 278
 " *vitripennis*, 278
Chrysomyia, 280
 " *affinis*, 280
 " *flaviventris*, 280
Chrysopila, 281
 " *ferruginosa*, 282
 " *insularis*, 282
 " *lupina*, 282
 " *maculipennis*, 281
 " *uniguttata*, 282
Chrysops, 264
 " *albicinctus*, 265
 " *clavicus*, 265
 " *dispar*, 265
 " *fasciatus*, 265
 " *fixissimus*, 265
 " *flaviventris*, 265
 " *ligatus*, 264
 " *pellucidus*, 265
 " *rufitarsis*, 265
 " *semicirculus*, 265
 " *stimulans*, 265
 " *striatus*, 265
 " *terminalis*, 265
 " *translucens*, 265
 " *unizonatus*, 265
Cirrochroa, 171
Claoxylon, 297
 " *longifolium*, 326, 356, 404
Clematis, 227
 " *Gauriana*, 227
 " *grata*, 227
 " *orientale*, 227
 " *orientalis*, 227
 " *puberula*, 227
Clerodendron, 341
 " *inermis*, 292, 298, 323, 341,
 355, 379, 381
Clitellaria, 273
 " *angusta*, 273
 " *bivittata*, 273, 278
 " *flaviceps*, 274
Clitellaria heminopia, 273
 " *nigerrimum*, 274
 " *notabilis*, 273
 " *spinigerum*, 274
 " *tenebrica*, 273
 " *varia*, 273
Cnesmone javanica, 297, 327, 356, 404
Cocos, 374
 " *nucifera*, 292, 296, 297, 332, 339,
 346, 357, 371, 372, 373, 375, 377, 378,
 381, 384, 385
Calostegia, 39, 56
 " *bornensis*, 57
 " *Griffithii*, 57
 " *sumatrana*, 57
 CÆNOMYDÆ, 272
Coladenia dan, 196
Colax, 281
 " *javanus*, 281
 " *variegatus*, 281
Colebrookia oppositifolia, 227
Colcosporium, 226
Collema nigrescens, 338, 358, 389
Coluber helena, 239
 " *Nuthalli*, 239
 " *oxycephalus*, 239
 " *prasinus*, 231, 239
 " *radiatus*, 239
 " *reticularis*, 239
 " *tæniurus*, 239
Colubrina asiatica, 292, 307, 352, 380
 COMBRETACEÆ, 313, 345, 349
Commelina obliqua, 330, 357, 401
 COMMELYNACEÆ, 345, 350
 COMMELYNEÆ, 330
Commersonia, 59, 93
 " *echinata*, 94
 " " *var.*, 94
 " *javensis*, 94
 " *platyphylla*, 93
 COMPOSITE, 317, 345, 349
 CONNARCEÆ, 309, 345, 349
Connarus gibbosus, 309, 353, 403
Conosia, 257
 " *cruz*, 257
 " *irrorata*, 257
 CONVULVULACEÆ, 320, 344, 345, 349, 383,
 396
Convolvulus parviflorus, 294, 321, 355,
 381, 383
Corchorus, 96, 117
 " *acutangulus*, 117, 118
 " *æstuans*, 118
 " *capsularis*, 117, 118
 " *decemangularis*, 118
 " *fuscus*, 118
 " *Marua*, 118
 " *olitorius*, 117, 118
Cordia subcordata, 298, 319, 355, 381
 COROLLIFLOREÆ, 316
Corypha, 297, 331

- Corypha clata*, 298, 331, 357, 404
 " *Gebanga*, 331
 " *macropoda*, 331
Costus speciosus, 329, 346, 356, 398
Cotoneaster, 225
 " *bacillaris*, 224
Crapitula, 263
Crastia, 9, 11, 21
 " *climena*, 10, 22
 " *Deheerii*, 163
 " *enganensis*, 20, 22
 " *oceanis*, 21, 22
 " *palmedo*, 162
Crinum, 345
 " *asiaticum*, 292, 329, 357, 381
Crotalaria retusa, 309
 " *sericea*, 309, 353, 368, 372, 377
Croton sublyratus, 296, 326, 356, 404
Crydora spongiosa, 200
CRYPTOGAME, 336, 344
Ctenophora 254
 " *annulosa*, 255
 " *ardens*, 255
 " *chrysophila*, 255
 " *compedita*, 255
 " *curvipes*, 255
 " *fumiplena*, 255
 " *javanica*, 255
 " *lata*, 254
 " *melanura*, 254
 " *taprobanes*, 255
 " *xanthomelana*, 254
CUCURBITACEÆ, 315, 345, 349
Calua, 274
 " *simulans*, 274
Culex, 251, 252
 " *amboinensis*, 251
 " *annulipes*, 252
 " *aurcostriatus*, 251
 " *cingulatus*, 251
 " *crassipes*, 252
 " *dives*, 252
 " *fatigans*, 251
 " *fuscus*, 251
 " *imprimens*, 252
 " *laniger*, 251
 " *longipalpis*, 252
 " *luridus*, 252
 " *molestus*, 251
 " *nero*, 251
 " *setulosus*, 252
 " *sitrens*, 251
 " *splendens*, 251
 " *subulifer*, 251
 " *vagans*, 251
 " *variegatus*, 252
 " *ventralis*, 252
CULICIDÆ, 250
Culicidæ, 252
CULICIFORMES, 250
CULICINA, 250
CULICINÆ, 350
Cupha erymanthis, 25, 171
Cupressus, 224
Curetis, 179
 " *malayica*, var. *kiritana*, 179
 " *obscura*, 179
Cyaniris, 29, 180
 " *akasa*, 180
 " *danis*, 36
 " *puspa*, 29, 180
 " *puspinus*, 29
CYCADACEÆ, 328, 345, 350
Cycas, 288, 379
 " *Rumphii*, 292, 328, 338, 339, 343
 " 346, 356, 381
Cyclea peltata, 301, 351, 401, 402, 403
Cyclogaster, 274
 " *radians*, 274
Cyclostemon assamicus, 294, 296, 326, 346,
 356, 404
Cylindrotoma, 260
 " *acrostacta*, 258
 " *albitarsis*, 260
 " *ornatissimus*, 260
Cymodocea, 288, 379
 " *ciliata*, 287, 334, 357, 378
Cynometra, 295, 298
 " *ramiflora*, 293, 296, 312, 346,
 353, 379, 380
Cynthia, 11, 12, 13
 " *deione*, 170
 " *parthenos*, 171
CYPERACEÆ, 334, 344, 345, 350, 372
Cyperus, *dilutus*, 334, 357, 392
 " *elegans*, 294, 334, 337, 346, 357,
 392
 " *pennatus*, 297, 334, 345, 357, 381
 " *polystachyus*, 295, 299, 334, 345,
 357, 392
CYPRINIDÆ, 144
Cyrestis nais, 171, 172
 " *nivea*, 172
 " *periander*, 8, 26
 " *themire*, 26
 " *thyodamas*, 172
CYRTIDÆ, 282
CYRTIDI, 282
Dacalana, 35
Daldinia vernicosa, 340, 358, 389
Danaida, 9
DANAIDÆ, 4, 8, 9, 11, 12, 13, 14, 15, 17,
 19, 158, 159
DANAINE, 9
Danais, 9, 10, 11, 12, 13, 15, 16, 17, 18,
 24, 158
 " *abigar*, 164
 " *agnis*, 164
 " *aglaia*, 12, 14
 " *aglaioides*, 12, 14
 " *archippus*, 9
 " *aruana*, 164

- Danais australis*, 166
 „ *bataviana*, 163
 „ *chionippe*, 164
 „ *chrysea*, 32
 „ *chrysippus*, 13, 14, 163, 174
 „ *crocea*, 14, 23
 „ *eurydice*, 23
 „ *fulgurata*, 164
 „ *Gautama*, 166
 „ *Gautamoides*, 14
 „ *genutia*, 13, 14, 164, 165, 166
 „ *haruhasa*, 166
 „ *hegesippus*, 14
 „ *intensa*, 164
 „ *ismare*, 165, 166
 „ *larissa*, 14, 166, 167
 „ *limniace*, 13, 14, 15, 17, 166
 „ *litoralis*, 197
 „ *luzonensis*, 166
 „ *melaneus*, 14
 „ *melanoleuca*, 14
 „ *melissa*, 166
 „ *nesippus*, 14
 „ *nilgiriensis*, 166, 167
 „ *orientis*, 197
 „ *philomela*, 23
 „ *Pietersii*, 23, 26, 32
 „ *plexippus*, 9
 „ *septentrionis*, 14
 „ *tytia*, 13, 14
 „ *vulgaris*, 14
 „ (*Bahora*) *chrysea*, 23
 „ (*Chiffra*) *orientis*, 166
 „ (*Limnas*) *chrysippus*, 163
 „ (*Nasuma*) *haruhasa*, 157, 165
 „ „ *taimanu*, 165
 „ (*Salatura*) *genutia*, 164
 „ „ *litoralis*, 164
 „ (*Tirumala*) *limniace*, 166
 „ „ *melissa*, 166
 DANAUS, 9
Danisepa, 16
Darasana paramuta, 34
 „ *perimuta*, 34
Dasyncura, 253
Datura, 396
Davallia solida, 295, 336, 337, 358, 389
Dedalea concentrica, 339, 358, 389
 „ *flabellum*, 339, 358, 363, 389
 „ *quercina*, 339, 358, 389
 „ *sanguinea*, 339, 358, 389
Dehaasia Kurzii, 325, 356, 397
Deima fastosum, 198
 „ *validum*, 198
 DEIMATIDÆ, 197
Delias, 187, 189, 190
 „ *aurantia*, 189, 190
 „ *belisama*, 189, 190
 „ *Descombasi*, 189
 „ *hyparete*, 187
 „ *nakula*, 190
 „ *Delias orata*, 189
 „ *pasithoë* var., 189
Dendrobium, 390
 „ *secundum*, 295, 328, 356, 389
Dendrocalamus, 294
 „ *strictus*, 294, 336, 346,
 358, 363, 364, 401, 402,
 404, 405
 „ „ var., 300
Derris, 347
 „ *scanflens*, 296, 311, 353, 403
 „ *sinuata*, 311, 353, 380, 383
 „ *uliginosa*, 292, 296, 311, 353, 380
Desmodia, 383, 393
Desmodium laxiflorum, 294, 309, 346, 353,
 393, 394
 „ *polycarpum*, 292, 297, 309,
 345, 353, 380, 382, 384, 396,
 394
 „ *triflorum*, 310, 353, 368, 377
 „ *triquetrum*, 292, 297, 309, 345,
 353, 380, 382, 393, 394
 „ *umbellatum*, 292, 298, 309,
 353, 379, 380, 384
Dendrorix, 180
 „ *epiarbas*, 27
 DEUDORIGINÆ, 27, 179
 DICOTYLEDONES, 344
Dicranomyia, 259, 261
 „ *saltens*, 261
Dictenidia, 254
Dictyota dichotoma, 341, 359, 378, 379
Diemenia, 115
Dioscorea, 294, 296, 347, 390
 „ *bulbifera*, 390
 „ *glabra*, 329, 357, 389
 „ *pentaphylla*, 330, 357, 389
 DIOSCOREACEÆ, 329, 345, 350
Diospyros, 135
 „ *Kurzii*, 300, 318, 354
Diplospora singularis, 294, 316, 340, 354,
 403
Dipsas ceylonensis, 243, 244
 „ *cyanea*, 244
 „ *cynodon*, 244
 „ *multifasciata*, 243, 244
 „ *multifasciatus*, 231
 „ *rhinopoma*, 231
 DIPTERA, 250
 DIPTEROCARPEÆ, 38, 302, 345, 349
Dipterocarpus, 290, 292, 295, 297, 300,
 384, 388
 „ *alatus*, 303, 351, 403
 „ *Griffithii*, 302
 „ *pilosus*, 302, 351, 403
Dischidia nummularia, 319, 354, 389
 DISCIPLOREÆ, 305, 341
Discophora, 170
 „ *timora*, 170
Distira cyanocincta, 247
 „ *tuberculata*, 231

- Ditylomyia*, 267
 " *ornata*, 267
Diza, 254
 " *guttipennis*, 254
 DIXADI, 254
Dodonaea viscosa, 297, 308, 352, 380, 385, 388
Dolenschallia, 158, 173, 174
 " *bisaltide*, 26
 " *niatica*, 26
 " *polibete*, 26
 " *pratipa*, 26
 DOLICHODERIDE, 1
Dolichos lineatus, 311
 Dorites, 295
 " *Wightii*, 295, 296, 356, 389
 " var., 328
Dracena angustifolia, 330, 357, 381
 " *spicata*, 294, 296, 330, 357, 398
Dracontomelum, 382, 396
 " *mangiferum*, 308, 352, 380, 382, 403
 " *sylvestre*, 295
Drina donina, 34
 " *mancia*, 34
Dryophis fronticinctus, 244
 " *mycterizans*, 244
 " *pulchralentus*, 244
Durio, 39, 50, 54
 " *affinis*, 52
 " *Loisianus*, 51
 " *malaccensis*, 51, 53
 " *Ozleyanus*, 50, 54
 " *perakensis*, 52
 " *testitudinarum*, 52
 " " var., *macrophylla*, 53
 " " var., *Pinangiana*, 53
 * " *Wrayi*, 53
 " *tibethicus*, 50, 51
 EBENACEÆ, 318, 345, 349
Echinops niveus, 219
Echinosoma, 203
 " *hispidum*, 203
Ehretia serrata, 228
Eleocarpus, 96, 120, 140
 " *Acronodia*, 139, 140
 " *amygdalinus*, 127
 " *angustifolius*, 126
 " *apiculatus*, 121, 134, 135, 136
 " *aristatus*, 121, 127, 136
 " *cuneifolius*, 135
 " *cyanocarpa*, 122, 123
 " *floribundus*, 121, 128
 " *Ganitrus*, 120, 122, 123
 " *glabrescens*, 121, 122, 138
 " *Griffithii*, 121, 131, 132
 " *grossa*, 129
 " *Helferi*, 127
 " *Hullettii*, 121, 132
Eleocarpus hypadenus, 126
 " *integra*, 130
 " *Jackianus*, 121, 137
 " *Kunstleri*, 121, 133, 136
 " *leptostachyus*, 120, 128
 " *littoralis*, 134, 135
 " *Lobbianus*, 129
 " *Mastersii*, 122, 140
 " *Monoceras*, 134, 135
 " *nitidus*, 120, 127, 128, 133
 " " var., *leptostachya*, 128
 " *oblonga*, 127
 " *oblongus*, 129
 " *obtusus*, 121, 134, 135
 " *ovalifolius*, 127
 " *ovalis*, 131
 " *paniculatus*, 121, 129
 " *parvifolius*, 120, 123, 125, 126
 " *pedunculatus*, 121, 127, 131, 132, 133
 " *petiolatus*, 121, 130
 " *polystachyus*, 121, 137
 " *punctatus*, 122, 139, 140
 " *robustus*, 120, 126
 " *rugosus*, 136
 " *salicifolius*, 120, 125
 " *Scortechinii*, 120, 124
 " *serratus*, 127, 129
 " *simplex*, 137
 " *stipularis*, 120, 123, 124
 " " var., *latifolia*, 124
 " *tomentosus*, 139
 " *venustus*, 137
 * " *Wrayi*, 120, 124
 ELASIPODA, 197
Elasma, 276
 " *acanthinoidea*, 276
 ELATERIDÆ, 7
Eleusine ægyptiaca, 295, 336, 358, 372, 377
 " *indica*, 295, 336, 358, 372, 377, 401
Elodina, 186
 " *egnatia*, 186
 ELPIDIDÆ, 197
 ELYMNIADÆ, 24, 170
Elymnias, 170
 " *dolorosa*, 25
 " " var., *enganica*, 24
 " *panthera*, 25
 " *undularis*, 170
 ENPODIATA, 264
Engonia, 273
 " *aurata*, 273
 " *spinigerum*, 274
Enhalus accroides, 334
Entada scandens, 298, [312, 353, 378, 380, 383
Eoözyllides tharis, 8, 27, 35
Epeorus, 406, 413
 " *psi*, 413

- Ephemerica*, 406, 409, 410, 411 •
 * " *consors*, 412
 " *expectans*, 409
 " *faciata*, 410
 " *immaculata*, 409
 * " *remensa*, 410, 412
 " *supposita*, 409, 411
Ephemerella, 406, 413
 EPHEMERIDÆ, 403
Ephippium, 273
 " *angusta*, 273
 " *nigerrimum*, 274
 " *spinigerum*, 273, 274, 278
Eranthemum, 292
 " *album*, 322, 355, 404
 " *cinnabarinum*, 322
 " " *var.*, *succisifolia*, 322
 " *succifolium*, 296, 322, 355, 381
Eryolis ariadne, 170
 " *merione*, 170
Erigeron alpinus, *var.*, *multicaulis*, 218
Eriocera, 255, 257, 258
 " *acrostacta*, 255, 258
 " *albipunctata*, 258
 " *albonotata*, 258
 " *bicolor*, 257
 " *crystalloptera*, 258
 " *ferruginosa*, 258
 " *hilpa*, 258
 " *Humberti*, 258
 " *lunata*, 258
 " *melantria*, 258
 " *pachyrrhina*, 258
 " *selene*, 257
Eriodendron, 39, 49, 390
 " *anfractuosum*, 49, 296, 304, 352, 388
 " *orientale*, 50
Erioglossum edule, 308, 352, 397
Erycibe paniculata, 297, 320, 355, 397
Erycina, 10
 ERYCINIDÆ, 10
Erythrina, 288, 298, 386
 " *indica*, 292, 296, 310, 353, 379, 380
Erythropsis Roxburghiana, 72
Eudmeta, 280
 " *marginata*, 280
Eu-elyocarpus, 120
Eugenia, 396
Eunepeta, 204
Euonymus, 207, 208
 " *calcarpus*, 206, 207, 209, 306
 " *javanicus*, 207, 209
Euphorbia, 213
 " *Atoto*, 288, 292, 326, 356, 379, 381
 " *hypericifolia*, *var.*, *indica*, 213
 " *pilulifera*, 326, 356, 371, 377
 EUPHORBACEÆ, 326, 344, 350, 384
Euphoroides depressa, 200
Euphoca, 10, 11, 13, 15, 16, 17, 18, 22, 157, 158, 163, 165, 166
 " *alcatheë*, 13, 18
 " *baudiniana*, 162
 " *camorta*, 13, 22
 " *climena*, 21, 162, 163
 " *core*, 9, 10, 11, 13, 22
 * " *elevsii*, 197
 " *espeii*, 13
 * " *leuca*, 158, 162, 166, 197
 " *midamus*, 13, 14, 20
 " *mulciber*, 20
 " *oceanis*, 15, 163
 " *orope*, 162
 * " *pahakela*, 22
 * " *palmedo*, 197
 " *pinwillii*, 13
 " *rhadamanthus*, 13, 14
 " *sepulchralis*, 21
 " *simulatrix*, 9, 13, 21
 * " (*Crastia*) *cheerri*, 163
 * " " *enganensis*, 20, 22
 * " " *oceanis*, 21, 22
 * " " *palmedo*, 162
 * " (*Pecoa*) *menetricii*, 22
 * " (*Rasuma*) *leuca*, 162
 * " (*Sclinda*) *elcusina*, 158
 * " (*Trepsichrois*) *dongo*, 160
 * " " *elevsii*, 161
 * " " *malakoni*, 20, 22
 " " *midamus*, 16
 " " *Verhuelii*, 22
 * " (*Tronga*) *niasica*, 22
 * " (*Vadebra*) *palmedo*, 162
 EUPLOINA, 9
 EUPLOINÆ, 9
Eupyrus, 203
 " *hispidus*, 203
 " *scaber*, 203
 EUSTERCULIA, 59
Euthalia, 9, 178
 " *aconthea*, 9
 " *adonia*, 9
 " *evelina*, 9
 " *garuda*, 178
 " *labentina*, 9
 Evaza, 275
 " *argyroceps*, 275
 " *bipars*, 275
 " *flavipes*, 275
 " *fulvicentris*, 275
Everes argiades, 28
 " *parhasius*, 28, 181
 FICOIDÆ, 315, 345, 349
Ficus, 225, 294
 " *Benjamina*, 327, 356, 397
 " *brevicuspis*, 296, 327, 340, 356, 398
 " *callosa*, 327, 356, 398
 " *comosa*, 327

- Ficus*, *Dæmonum*, 296
 " *grisea*, 328, 356, 398
 " *hispida*, 356, 398
 " " *var.*, *Dæmonum*, 327
 " *nitida*, 340
 " *palmata*, 225
 " *retusa*, 397
 " " *var.*, *nitida*, 327, 356
 " *Rumphii*, 290, 292, 327, 356, 397
FILICES, 336, 343, 344, 345, 350
Fimbristylis, 297, 299, 345
 " *diphylla*, 334, 357, 371, 377
 " *ferruginea*, 335, 357, 381
 " *miliacea*, 335, 357, 392
 " *quincangularis*, 335, 357, 392
Fimiana, 60, 74
 " *colorata*, 72
 " " *var.*, 72
Flagellaria indica, 293, 330, 357, 379, 381
FLAGELLARIÆ, 330, 345, 350
 * *Flos ahanus*, 33, 38
 " *apidanus*, *var.*, 179
 " *asaka*, 33
 " *fulgidus*, 33
 " *mullerii*, 34
Flueggia microcarpa, 326, 356, 397
Fordonia leucobalia, 245
FORNICIDÆ, 1
FROHLICHE, 9
FUNGI, 338, 344, 348, 350, 385, 390, 404
Gamboba, 21, 163
GAMOPETALE, 343, 344
Ganitrus, 120
 " *sphærica*, 122
Garcinia, 302, 351
Gardenia, 340
Garuga pinnata, 306, 352, 397
Gchlota, 196
Gelidium, 342
 " *cornutum*, 342, 359, 378
GENTIANACEÆ, 319, 345, 349
Gerarda bicolor, 245
GERYDINÆ, 29, 36, 185
Gerydus, 37
 " *acragas*, 186
 " *ancoa*, 37
 " *Biggsii*, 36, 37, 186
 " *Boisdavalii*, 37, 186
 " " *var.*, *acragas*, 186
 " *croton*, 36, 37
 " *gopara*, 36
 " *heracleion*, 36, 37
 " *irroratus*, 37, 186
 " " *var.*, *assamensis*, 37, 38
 " *melanion*, 37
 " *pandu*, 185
 " *symcthus*, 36, 37, 185
 " *teos*, 185
GRACIÆ, 350
Glaphyoptera, 262
 " *Winthemi*, 262
Glochina, 259
Gloriosa, 347
 " *superba*, 330, 357, 378, 381, 385, 388, 389, 390
Glossospermum cordatum, 88
 " *velutinum*, 88
Glycosmis pentaphylla, 296, 305, 346, 352, 397
Glyptopetalum, 206, 207, 208
 " *calocarpum*, 208, 209, 294, 296, 306, 346, 352, 403
 " *grandiflorum*, 208, 210
 " *Griffithii*, 208, 209
 " *sclerocarpum*, 207, 208, 210
 " *zeylanicum*, 206, 208, 209, 210, 307
Gomphrena, *globosa*, 324, 356, 371, 372, 377
Gonyosoma dorsalis, 238
GOODENOVIEÆ, 317, 345, 349
Gossampinus rubra, 49
Gracilaria crassa, 342, 359, 378
GRAMINEÆ, 335, 344, 350, 372
Grapsus, 393
Grewia, 96, 108, 109, 110, 115, 294, 306
 " *affinis*, 111
 " *antidesma folia*, 109, 113
 " " *var.*, *hirsuta*, 113
 " *Blumei*, 111
 " *calophylla*, 109, 114, 305, 352, 403
 " *caudata*, 95
 " *fibrocarpa*, 109, 111, 112
 " *globulifera*, 109, 112
 " *heteroclita*, 95
 " *laevigata*, 305, 352, 397
 " *latifolia*, 109, 112
 " *laurifolia*, 109, 114
 " *Microcos*, 305, 352, 397
 " *Miqueliana*, 109, 115
 " *paniculata*, 109, 110, 111
 " *pedicellata*, 110
 " *umbellata*, 109
GREWICÆ, 96
Guettarda speciosa, 292, 297, 298, 316, 354, 380
GUTTIFERÆ, 302, 345, 349
GYNOSPERMÆ, 328, 344
Gymnosporangium Cunninghamianum, 224, 225
Gyrocarpus, 288, 294, 298, 388
 " *Jacquinii*, 292, 295, 296, 297, 314, 318, 346, 353, 366, 380
Hæmatopota, 266
 " *atomaria*, 266
 " *borneana*, 266
 " *cava*, 266
 " *cilipes*, 266

- Hamatopota, cingulata*, 266
 " *irrorata*, 266
 " *juvana*, 266
 " *lunulata*, 266
 " *macrocera*, 266
 " *pachcera*, 266
 " *pungens*, 266
 " *roralis*, 266
Halimeda Opuntia, 342, 359, 378
Halophila ovalis, 288, 334
Halpe, 197
Hamadryas, 10
 **Harimata maremba*, 192
 " *peranthus, var.*, 193
Hasora badra, 32, 194
Hebomoia glaucippe, 191
Helcyra chionippe, 174
 HELICTEREE, 58
Helicteres, 58, 81
 " *angustifolia*, 81, 83
 " " *var., obtusa*, 83
 " *chrysocalyx*, 82
 " *hirsuta*, 81, 82, 83
 " " *var. oblonga*, 82
 " " *vestita*, 82
 " *Isora*, 81, 82, 83
 " *lanccolata*, 83
 " *spicata*, 82
 " *virgata*, 83
Heliomyia, 281
 " *ferruginea*, 281
Hemipuccinia, 214
Hemiuromyces, 211
Henops, 282
 " *costalis*, 282
Hepatica, 338
 HEPATICE, 338, 344, 350
Heritiera, 58, 79, 300, 356
 " *acuminata*, 80
 " *Fomes*, 80
 " *littoralis*, 79, 80, 295, 305, 352,
 350
 " *macrophylla*, 80
 HERMANNIE, 58
Hermetia, 274, 280
 " *batjanensis*, 274
 " *bislectum*, 274
 " *cingulata*, 280
 " *cingulalis*, 280
 " *marginata*, 280
 " *melanoxis*, 274
 " *rußcentris*, 274
Hernandia, 288
 " *peltata*, 292, 296, 297, 325,
 356, 379, 381
 HERNANDIACEE, 115
 HESPERIADÆ, 32, 194, 197
Hestia, 9, 10, 12, 13, 16, 17, 18, 160, 189
 " *Cadellii*, 13, 18
 " *Hadenii*, 13
 " *lynceus*, 13
- HETEROPETALE, 96
Heterophragma adenophyllum, 297, 322,
 355, 389
Heteropyxis, 55
Hexagonia, 406, 409
Hexagona pergamenca, 339, 358, 389
 " *sericeo-hirsuta*, 358, 389
 " *sericeo-hirsutus*, 339
 " *tenuis*, 339, 358, 389
Hidari irava, 32
 HIBISCEE, 39
Hibiscus, 39, 44
 " *Abelmoschus*, 44, 303, 351, 368,
 372, 377
 " *clatum*, 46
 " *flavescens*, 45
 " *floccosus*, 44, 46
 " *furcatus*, 45
 " *longifolius*, 45
 " *macrophyllus*, 44, 45
 " *populneoides*, 48
 " *populneus*, 48
 " *pseudo-abelmoschus*, 45
 " *ricinifolius*, 45
 " *rosa-sinensis*, 340
 " *sabdariffa*, 303, 351, 368, 372,
 377
 " *sagittifolius*, 45
 " *setosus*, 46
 " *spathaceus*, 45
 " *surattensis*, 44, 45
 " *tiliaceus*, 44, 46, 290, 292, 293,
 298, 303, 351, 379, 380
 " *tortuosus*, 46
 " *vestitus*, 46
Hirnicola polytricha, 340, 358, 389
Hirtea, 262, 263, 264, 277
 " *fulvicollis*, 263
 HOLOPETALE, 96
 HOMALOCERATI, 250
Homolosoma, 237
Hoplomyia, 277
Hoya, 295, 390
 " *diversifolia*, 319, 354, 389
 " *parasitica*, 319, 354, 389
Huphina, 190
 " *corva*, 188
 * " *cirene*, 188
 * " *ethel*, 29, 188
 " *imogene*, 188
 " *judith*, 30, 188
 * " *julia*, 187, 197
 " *leta*, 187
 " *lea*, 30
 " *naomi*, 188
 " *tamar*, 187
 " *temena*, 187
 * " *vaso*, 188
Hydrophis crassicollis, 247
 " *trachiceps*, 247
Hydrophobus davisoni, 231

- Hydrosaurus*, 402
 " *salvator*, 402
Hygrophila, 300
 " *quadriculvis*, 299, 322, 324,
 355, 392
Hypolimnas, 158
 " *anomala*, 26, 174
 " *bolina*, 26, 174
 " *missippus*, 174
 " *pandarus*, 158, 174
 " *saundersii*, 174
Hypolycaena *erylus*, 180
 " *sipylus*, 179
 " *thecloides*, 27, 179
Hypsirhina *blanfordii*, 231, 244
 " *maculosa*, 231
 " *sieboldii*, 245
Irisia, 281
Ichuocarpus frutescens, 228
Ideopsis, 10, 13, 17
 " *daos*, 14
**Iliaes merapu*, 191
 " *oceani*, 31
Illigera Conyzadenia, 314, 353, 388, 403
 INCOMPLETE, 324, 344
 INFLORE, 282
Inocarpus, 115
Inodaphnis lanceolata, 115
Ipomoea, 288, 292, 297
 " *Batatas*, 320, 355, 369, 372, 377
 " *biloba*, 288, 290, 292, 296, 297,
 298, 321, 355, 379, 381
 " *coccinea*, 320, 355, 369, 372, 377
 " *denticulata*, 292, 297, 320, 355,
 381, 386
 " *digitata*, 320, 355, 379, 381
 " *grandiflora*, 294, 320, 355, 381
 " *hederacea*, 228
 " *palmata*, 294
 " *Turpethum*, 321, 355, 384, 401,
 402
Iraota timoleon, 179
Isamia, 16, 22, 159
 " *lowii*, 22
Isamia standingerii, 22
Ischemum ciliare, 295, 336, 358, 389, 401
 " *muticum*, 295, 336, 345, 358,
 381
Ismene, 194
Isora corylifolia, 82
 ITHOMIADÆ, 11
Leias pyrene, 191
 " *pyrene*, 191
 " *reinwardtii*, 191
Isora brunnescens, 292, 297, 316, 354, 389
 " *cuneifolia*, 316, 354, 403
 " *grandiflora* var. *Kurzeana*, 316, 354
 " *grandifolia*, 403
 " *Kurzeana*, 316
Jamides bochus, 28, 183
Jania tenella, 341, 359, 378
Jatana, 143, 169
 " *wayaca*, 168
Junonia almana, 172, 173
 " *anonis*, 172
 " *asterie*, 172, 173
 " *atrites*, 172
 " *erigone*, 173
 " *javana*, 172
 " *semonias*, 173
 " *orithyia*, 173
 " *timorensis*, 173
 " *vellida*, 172
Kerana, 197
Kleinhovia, 58, 80
 " *Hospita*, 80, 98
Kyllinga, 299
 " *brevifolia*, 334, 357, 371, 377
 LABIATE, 204, 297, 323, 345, 350
Laertias polites, 193
Laetmogene spongiosa, 200
Lagerstræmia, 315, 354, 403
 " *calyculata*, 315
 " *hypoleuca*, 300, 314, 354,
 403
Lampides, 12, 28, 183
 " *alianus*, 14, 184
 " *amphiasa*, 185
 " *anops*, 183, 184, 197
 " *aratus*, 184
 " *astraptes*, 183
 " *bochus*, 28, 183
 " *celeno*, 28, 185
 " *elpidion*, 28
 " *elpis*, 28, 183, 185
 " *masu*, 184, 197
 " *nicobaricus*, 28
 " *subditus*, 28, 185
 " var. *telanjang*, 28
Lasioptera, 254
 " *bryonica*, 254
Lathyrus satirus, 212
 LAURINEÆ, 325, 345, 350
Leca, 290, 382
 " *hirta*, 307, 352, 380, 382, 327
 " *sambucina*, 293, 295, 307, 346, 352,
 379, 380, 382
 LEGUMINOSÆ, 309, 344, 349, 383
Leia, 262
Lejeunia, 338, 358
Lentianus leucochrous, 338, 358, 389
Leuzites deplanata, 338, 358, 389
 " *subferruginea*, 338, 358, 389
Lepraria, 338, 355
 LEPTIDÆ, 281
 LEPTIDES, 281
 LEPTIDI, 281
Leptis, 281
 " *decisa*, 281
 " *ferruginosa*, 282
 " *unguttata*, 281
Leptonychia, 59, 94, 95

- Leptonychia, acuminata*, 95
 " *glabra*, 94
 " *var. Mastersiana*, 95
 " *heteroclita*, 95
 " *macarroides*, 95
Leptopuccinia, 217, 218
Lepturomyces, 213
Lethe europa, 24, 168
Lettonia peguensis, 320, 355, 403
Libnotes, 261
 " *thwaitesiana*, 261
Libythea, 13
 " *geoffroyi*, 178
 " *narina*, 178
 " *rohini*, 178
 LIBYTHEIDÆ, 178
 LICHENES, 338, 344, 350
 LILIACEÆ, 330, 345, 350
 * *Limenitis agpeya*, 176, 177
 " *daraza*, 176, 177
 * " *hollandii*, 177
 " *lysianis*, 177
 " *populi*, 177
 " *procris*, 176
 LIMNAINA, 9
Linnanthemum, 299
 " *indicum*, 299, 319, 355, 392
Linnaea, 9
 " *chrysippus*, 163
Limnobia, 257, 259, 260, 261
 " *acrostacta*, 258
 " *albonotata*, 258
 " *apicalis*, 259
 " *argentocincta*, 259
 " *aterrima*, 259
 " *aurantiaca*, 261
 " *basilaris*, 257
 " *bibula*, 259
 " *costalis*, 259
 " *diana*, 259
 " *impressa*, 259
 " *irrorata*, 257
 " *leucotelus*, 259
 " *mesopyrrha*, 259
 " *plecioides*, 260
 " *pyrrhochroma*, 259
 " *rubescens*, 259
 " *sanguinea*, 259
 " *strbillans*, 259
 " *substituta*, 259
 " *sumatrensis*, 259
 " *trentepohlii*, 259
 " *vittifrons*, 260
Limnomyia, 257
Limnomyza, 259
Limnophila, 257
 " *basilaris*, 257
 " *bicolor*, 257
 " *crux*, 257
Limonia, 259, 260
Lippia nodiflora, 323, 355, 381, 383, 392
Lithothamnium polymorphum, 341, 359, 378
Livistona, 296, 331, 357, 404
 " *Jenkinsiana*, 331, 332
 " *speciosa*, 331, 332
Legunia, 29, 37
 " *marmorata*, 37
 * " *massalia*, 37, 38
 " *erica*, 29
Lophopetalum, 207
 LORANTHACEÆ, 325, 345, 350
Loranthus longiflorus, 325, 356, 394, 404
Lorura atymnus, 14, 179
Lumnitzera, 299, 301
 " *racemosa*, 314, 353, 380
Lycæna, 36
 " *atroguttata*, 36
 " *palmyra*, 182
Lycænaesthes bengalensis, 28
 LYCÆNIDÆ, 12, 27, 32, 179
 LYCÆNINÆ, 27, 36, 180
Lycodon fasciatus, 234
 " *gammiei*, 231
 " *striatus*, 233
 " *travancoricus*, 233
Lygodium flexuosum, 294, 337, 358, 389
 " *scandens*, 345
 LYTHRACEÆ, 314, 345, 349
Macacus cynomolgus, 148
Macaranga Tanarius, 296, 297, 327, 346, 356, 381, 384
Macropeza, 253
 " *gibbosa*, 253
Macroplæa, 10, 11, 13, 16, 17
 " *corus*, var. *micronesia*, 19
 " *elisa*, 10, 19
 " *micronesia*, 22
 " *phæretena*, 20, 22
 " *phæbus*, 19, 20, 22
 " *semicirculus*, 20
Macrostigma, 204, 205, 206
Mahintha, 9
Malais, 29
Malesia, 56
Mallotus acuminatus, 326, 356, 404
 " *andamanicus*, 296, 326, 356, 404
 " *Helferianus*, 326
 MALVACEÆ, 38, 303, 345, 349
Malvariscus populneus, 42
 MALVEÆ, 39
Massaga, 35
 " *pediada*, 35
 " *pharyge*, 35
 " *potina*, 35
Massicyta, 276
 " *bicolor*, 276
 " *cerioides*, 274
Megaerophis flaviceps, 245
Megarhina, 251
Megarhinus, 251
 " *splendens*, 251

- Meyisba malaya*, 185
Meyistocera, 260
 " *atra*, 260
 " *fuscana*, 260
 " *verticalis*, 260
Melampsora, 211, 223
 " *acidoides*, 223, 224
 " *ciliata*, 223, 230
 " *populina*, 223
 " *Tremulae*, 223
Melanitis, 12
Melanitis constantia, 170
 " *ismenc*, 24, 170
 " *leda*, 24, 158, 170
Melanocyna faunula, 18
 MELASTOMACEÆ, 314, 345, 349
 MELIACEÆ, 292, 306, 345, 349
Melius Megathina, 251
Melochia, 58, 87
 " *affinis*, 87
 " *concatenata*, 87
 " *corchorifolia*, 87
 " *pauciflora*, 87
 " *supina*, 87
 " *truncata*, 87
 " *velutina*, 88
Memecylon edule, 314, 354, 397
Menana, 9, 21
Menelaides aristolochia, 31
 " *oreon*, 192
 MENISPERMACEÆ, 301, 345, 349
Mezoneuron enneaphyllum, 294, 311, 353, 403
Microchryza, 278
 " *flaviventris*, 280
 " *gemma*, 278
Microcos, 103
 " *tomentosa*, 111
Micropuccinia, 216, 218
Microromyces, 213
Milusa, 292, 293, 295, 301, 346, 351, 396, 397, 403
 " *macrocarpa*, 301
 " *Rozburghiana*, 301
Mimusops, 288, 290, 294, 295, 297, 338, 341, 379, 386
 " *littoralis*, 292, 295, 297, 300, 318, 336, 338, 340, 341, 346, 354, 381, 385
Modecca, 296, 315, 347*
 " *cordifolia*, 294, 315, 354, 397
 " *cardiophylla*, 315
Molobrus, 262
 MOLPADIDÆ, 202
Mongoma, 261
 " *aurantiaca*, 261
 " *paciloptera*, 261
 " *simplex*, 261
Monocera, 121
 " *ferruginca*, 138
 " *Griethii*, 129, 131
Monocera holopetala, 131
 " *Pulembanica*, 133
 " *petiolata*, 130
 " *tricanthera*, 131
Monoceras leucobotryum, 129
 " *obtusum*, 134
 " *odontopetalum*, 131
 " *petiolatum*, 130
 MONOCOTYLEDONES, 328, 844
Morinda bracteata, 380
 " *citrifolia*, var. *bracteata*, 316, 354
Moringa pterygosperma, 309, 353, 368, 372, 377
 MORINGEÆ, 309, 345, 349
 MORPHIDÆ, 11, 12, 25, 170
Mucuna, 298, 383
 " *gigantea*, 293, 296, 310, 353, 380, 383
 " *pruriens*, 294, 310, 353, 401
 MUNTERE, 9
Musa, 345
 " *sapientum*, 329, 357, 371, 372, 377
 MUSCI, 337, 344, 350
Mussaenda calycina, 316, 354, 397
 " *macrophylla*, 316
Mycalexis blasius, 168
 " *mincus*, 8, 24, 169
 " *mynois*, 169, 170
 " *wayewa*, 170
 " (*Calysisme*) *perseus*, 168
 " (*Jatana*) *wayewa*, 168
 " (*Orotriena*) *medus*, 168
Mycetophila, 261
 " *bimaculata*, 261
 " *pcunipes*, 262
 MYCETOPHILIDÆ, 261
 MYCETOPHILIDES, 261
 MYCETOPHILIDI, 261
Mydas, 274
Myristeca glauca, 325, 356, 397
 " *Irya*, 325, 356, 397
 MYRISTICÆ, 325, 345, 350
 MYRSINÆ, 318, 345, 349
 MYRTACEÆ, 314, 345, 349
Nacaduba, 182
 " *almora*, 183
 " *ardates*, 27, 183
 " *atrata*, 183
 " *dana*, 183
 " *gaura*, 181, 182, 197
 " *hermus*, 183
 " *kerriana*, 183
 " *laura*, 182, 197
 " *macrothalma*, 27, 183
 " *perusia*, 183
 " *prominens*, 27
 " *psustis*, 182
 " *viola*, 27, 183
Naia tripudians, 246, 247
 NAIADACEÆ, 334, 350
 NAIADÆ, 345

- Nasuma*, 166
 * " *haruhasa*, 157, 165
 * " *tainanu*, 165
Nectaria, 13, 16, 17
 " *clara*, 19
 " *leucocœ*, 19
 " " *var. engania*, 19
Neesia, 39, 55
 " *altissima*, 56
Neesia Griffithii, 54
 " *synandra*, 56
Negritomyia, 273
 " *bilineata*, 278
Nematocera, 250, 260
 " *fuscana*, 260
 NEMATOCERATÆ, 250
 NEMEBIADÆ, 11, 12, 178
Nemestrina, 280
 " *jaana*, 280
 NEMESTRIDÆ, 280
 NEMESTRIDIDÆ, 280
Nemotelus, 274, 276, 279, 280, 281
Neocheritra, 35
 " *gama*, 35
Neopitheops zalmora, 185
 NEOTROPIDÆ, 10, 11, 12
Nepeta, 204
 " *Belleii*, 204, 206
 " *glomerulosa*, 204, 205
 " *juacca*, 204
 " *leucostegia*, 204
 " *Scordatis*, 204
 " *Sibthorpii*, 204, 205
 " *tuberosa*, 205
Nepheronia valeria, 190
Nephotoma, 255
Neptis, 14, 172, 179
 " *hordonia*, 175
 " *leucothœ*, 175
 " *nandina*, 175
 " " *var. sumba*, 175
 " *ombalata*, *var. engano*, 27
 " *soma*, *var. mecidiei*, 26
 " *varmona*, 14, 175, 179
Nerna, 275
 " *mollis*, 275
Nipa fruticans, 374
Niphobolus adnascens, 295
Nostocaceus Alga, A, 342
 NOTACHANTA, 272
 NOTACHANTEN, 272
Nychitona riphia, 186
 NYCTAGINÆ, 324, 345, 350
Nymphæa, 299
 " *Lotus*, 299, 300, 301, 351, 368, 392
 " *rubra*, 299, 301, 367, 372, 377
 NYMPHEACÆ, 301, 345, 349
 NYMPHALIDÆ, 11, 12, 26, 174
Ochna, 349
 " *squarrosa*, 340
Ochrosia borbonica, 318, 354, 381
Odina Wodier, 308, 352, 397
Odontomyia, 277
 " *consobrina*, 277
 " *diffusa*, 277
 " *mutica*, 277
 " *viridana*, 277
Oecophylla smaragdina, 2, 4
Oxygeodes, 282
 OLACINÆ, 306, 345, 349
Oligodon, 237
 " *dorsalis*, 237
 " *melanocephalus*, 237
 " *subgriseus*, 237
 " *sublineatus*, 237
Oligomera, 255
 " *javensis*, 255, 258
Oligotrophus, 253
Omphacarpus, 109
Oncodes, 282
 " *costalis*, 282
 ONCODINÆ, 282
Onciophanta mutabilis, 197
Oplismenus, 297, 396
 " *compositus*, 294, 335, 346, 357, 394
 ORCHIDACÆ, 328, 345, 350
Ornithoptera, 18, 194
 " *amphrysus*, 31
 " *criton*, 194
 " *haliphron*, 194
 " *minos*, 30
 * " *naias*, 157, 191, 193, 194
 * " *var. sambarana*, 194
 * " *nercis*, 30, 31
 " *plata*, 193, 194
 " *pompeius*, 30
 " *ruficollis*, 14
Orocyllum indicum, 296, 297, 322, 355, 381, 385, 389
Orpheides erichthonius, 191
 " *erithonius*, 191
 * *Orpheurgus asper var. glaber*, 198
Orsotricena medus, 168
Orthothecium hirsutum, 82
 " *javense*, 83
Oudemansia hirsuta, 82
 " *integerrima*, 83
 " *javensis*, 83
Pachygaster, 376
 " *rufitarsis*, 276
Pachyrhina, 255
 " *bombayensis*, 255
 " *delecta*, 255
 * " *daleschalli*, 256
 " *fasciata*, 256
 " *quadrivittata*, 255, 256
 " *triplasia*, 256
Pademna, 16
Padina paronia, 288, 341, 359, 378
Padraona palmarum, *var. kayapu*, 32

- Pedleria fetida*, 294, 296, 317, 354, 397, 399
Palmingia, 406
 " *ampla*, 408
 " *lata*, 407
 " *robusta*, 407
 " *minor*, 408
 PALMEE, 331, 345, 350
Palpomyda, 252
Pamphila, 196
 PANDANACEE, 345, 350
 PANDANEE, 333
Pandanus, 285, 286, 288, 290, 292, 294, 296, 298, 300, 371, 382, 403
 " *odoratissimus*, 297, 333, 357, 371, 379, 381
Pangonia, 265
 " *amboinensis*, 265
 " *longirostris*, 265
 " *rufa*, 266
 " *taprobanes*, 266
Panicum, 396
 " *ciliare*, 335, 357, 372, 377, 401
 " *colonum*, 295, 335, 357, 372, 377, 401
 " *Helopus*, 295, 335, 357, 372, 377, 401
 " *javanicum*, 335, 357, 401, 402
 " *montanum*, 335, 346, 357
 " *Myurus*, 299, 324, 335, 357, 392
Pannychia, 199
 " *moscleyi*, 199
 " *wood-masoni*, 198
Papilio, 11, 158
 " *antiphus*, 31
 " *aristolochiae*, 192, 193
 " *brama*, 193
 " *ganesa*, 34
 " *liris*, 192
 " *maremba*, 158, 191
 " *Memaon*, 31, 191
 " *merapu*, 154
 " *peranthus*, 193
 " *pericles*, 193
 " *presaspes*, 31
 " (*Charus*) *helenus*, 192
 " " " *var. enganius*, 31
 " (*Harimata*) *maremba*, 192
 " " *peranthus var.*, 193
 " (*Hiades*) *merapu*, 191
 " " *oceanii*, 31
 " (*Lavitas*) *polites*, 193
 " (*Menclaidis*) *aristolochiae*, 31
 " " *oreon*, 192
 " (*Orpheides*) *erichthonius*, 191
 " " *erithonius*, 191
 " (*Pathysa*) *antiphates*, 193
 " (*Zetides*) *agamemnon*, 31, 193
 " " *curypylus*, 193
 " " *sarpedon*, 31, 193
 PAPILIONIDÆ, 30, 191
Paradoxurus, 7
Paragerydus, 29
 " *horsfieldii*, 29
 " *unicolor*, 8, 29
Parantica, 17
 " *aglaia*, 23
Parastemon, 115
Parata, 197
 " *malayana*, 194
Parcas bermorci, 248
 " *macularius*, 248
 " *margaritophorus*, 248
Parinarium, 139, 140
Parishia insignis, 308, 352, 403
Paritium tiliaceum, 46
Parnara, 197
 " *naroca*, 196
Parthenos gambrisius, 26
 " *ilacinus*, 26
Paspalum acrobiculatum, 299, 335, 357, 392
 PASSERES, 143
 PASSIFLORACEÆ, 315
 PASSIFLOREÆ, 315, 349
Pathysa antiphates, 193
Pavetta indica, 316, 354, 397
Pemphis, 313
 " *acidula*, 292, 297, 298, 314, 354, 380
Pentagone wyvillii, 197
Pennisetum, 215
 " *typhoideum*, 214
Penca, 162
 " *menetricisii*, 22
Pentace, 96, 99
 " *Curtisii*, 100, 103, 104
 " *eximia*, 100, 103
 " *floribunda*, 100, 102
 " *Griffithii*, 100, 104
 " *Hookeriana*, 100, 101
 " *Kunstleri*, 100, 101
 " *macrophylla*, 100, 102
 " *perakensis*, 100, 101
 " *polyantha*, 101
 " *Scortechinii*, 100, 104
 " *strychnoidea*, 100, 105
 " *triptera*, 100
Pentagonia, 409
Penthetria fulvicollis, 263
 " *japonica*, 263
 " *melanaspis*, 263
Peristrophe acuminata, 323, 355, 381, 384, 404
Phedyma columella, 174
Phalacroopsis, 340
 PHANEROGAMÆ, 301, 344
Phaseolus, 310, 353, 368, 372, 377
 " *Phengaris*, 36
Phlebocalymna Lobbiana, 306, 352, 403
Phlomis lamifolia, 216
Pholidota imbricata, 329, 356, 389, 390

- Phragmidia*, 211
Phragmidium, 220, 221, 222, 228
 " *Barclayi*, 221, 222
 " *Fragariastris*, 220, 221
 " *Laccianum*, 220, 230
 " *nepalense*, 220, 221, 229
 " *octoloculare*, 221, 222, 229
 " *Potentillæ*, 220
 " *quinqueloculare*, 222, 230
 " *Rubi*, 221
 " *subcorticium*, 222
 " *Tormentillæ*, 220
Phyllanthus columnaris, 326, 356, 404
Phyllocladus spinosa, 327, 356, 397
 PHYLLOPHAGA, 7
Phyllophora, 274
 " *angusta*, 274
 " *bispinosa*, 275
Physalis minima, 297, 321, 355, 381, 383,
 396, 397
Physcia, 338, 358
 " *obscura*, 338, 358, 389
Physecrania, 257
 PICIDÆ, 143
 PIERIDÆ, 9, 12, 13, 29, 158, 186, 189
Pieris, 187, 190
 " *judith var.*, 188
Pilea trinervia, 228
Pimpinella, 219
 " *Grißthiana*, 219
Piper caninum, 325, 356
 PIPERACEÆ, 325, 345, 350
Pisonia, 288, 379, 384, 393, 394
 " *aculeata*, 293, 296, 324, 347, 355,
 366, 381, 393, 394
 " *excelsa*, 324, 355, 366, 381, 384,
 393, 394
Pithecopus hylax, 29
Planetella, 262
Planetes, 262
Platyura, 262
 " *venusta*, 262
Plebeius balliston, 28
 " *kupu*, 27
 " *lysimon*, 29
 " *polysperchianus*, 28
 " *siraha*, 28
 " *talinga*, 28
Pleria, 263
 " *dorsalis*, 263
 " *forcipata*, 263
 " *fulvicollis*, 263
 " *ignicollis*, 263
 " *melanaspis*, 263
 " *subvarians*, 264
 " *tergorata*, 263
 " *tristis*, 264
Plecospermum, 347
 " *andamanicum*, 293, 327,
 356, 397
Plesioneura restricta, 197
Pluchea indica, 292, 297, 317, 354, 380,
 384
 PNEUMONOPHORA 202
Pacilostola, 260
 " *pollens*, 260
Pollia sorzogonensis, 330, 357
 " *var.*, 401, 402
Pollinia japonica, 216
Polyalthia, 396
 " *longifolia*, 396
 POLYCHÆTA, 272
 POLYGONACEÆ, 324, 345, 350
Polygonum, 299, 300, 322
 " *barbatum*, 324, 356, 392
Polymitarceus, 406, 408
 " *indicus*, 408
Polyodontophis bistrigatus, 294
Polyommatus laticus, 181
 POLYPETALÆ, 343, 344
Polypodium adnascens, 337, 358, 389
 " *irioides*, 337, 358, 389
 " *quercifolium*, 295, 337, 358,
 389
 " (*Niphobolus adnascens*, 295
Polyporus australis, 339, 358, 389
 " *fulvus*, 339, 358, 389
 " *grammatocephalus*, 339, 358,
 389
 " *sanguinalis*, 385
 " *sanguineus*, 339, 358, 389
 " *xanthopus*, 339, 358, 389
Pometia tomentosa, 308, 352, 397
Pongamia, 288
 " *glabra*, 292, 311, 339, 340, 346,
 353, 380
Populus alba, 223, 224
 " *ciliata*, 223
Porana spectabilis, 296, 321, 355, 388, 403
Poritia, 36
 PORITINÆ, 35
Potamanthus (Atolophlebia) fasciatus, 409
Potentilla, 220, 221
 " *argyrophylla*, 220
 " *nepalensis*, 220
Pothos scandens, 334, 357, 398
Precis ida, 26, 173
 " *iphita*, 173
Premna, 323, 355, 381
 " *integrifolia*, 292, 323, 355, 381
 " *obtusifolia*, 323, 355
Prioneris, 189
Prionomyda, 252
Prionota, 255
 " *nigriceps*, 255
Protogonionomorpha, 173
Prymniodon, 242
 " *chalcus*, 242
Pselliophora, 254
 " *annulosa*, 255
 " *ardens*, 255
 " *chrysophila*, 255

- Pselliophora*, *compedita*, 255
 " *curvipes*, 255
 " *funiplena*, 255
 " *javanica*, 255
 " *lacta*, 254
 " *rubra*, 255
 " *taprobanes*, 255
Pseudergolis, 12
Pseudocyclophis bicolor, 231
 " *persicus*, 231
Pseudoxenodon macrops, 231, 241
Psilonepeta, 205
 PSILONEPETE, 205, 206
 *PSILOSTEGIA, 206
Psychotria adenophylla, 317, 354, 397
 PSYCHROPODIDÆ, 200, 201
Ptecticus, 273
 " *apicalis*, 278, 279
 " *australis*, 279
 " *brevipennis*, 279, 280
 " *cingulatus*, 278,
 " *illustris*, 279
 " *latifascia*, 279
 " *viridiceps*, 279
Pterocarpus indicus, 300, 311, 353, 388,
 403
Pterocosmus, 257, 258
 " *combinatus*, 258
 " *dilutus*, 258
 " *hilpa*, 258
 " *infixus*, 258
 " *lunigerus*, 258
 " *optabilis*, 258
 " *velutinus*, 258
Pterocymbium, 61
 " *campanulatum*, 76
 " *javanicum*, 76
Pterospermum, 58, 83, 139
 " *accrifolium*, 84, 86, 87
 " *aceroides*, 84, 86, 87
 " *Blumeanum*, 84, 86
 " *cinnamomeum*, 85
 " *diversifolium*, 84
 " *fuscum*, 85
 " *Jackianum*, 84, 85
 " *javanicum*, 85
 " *lanceifolium*, 85
 " *oblongum*, 86
 " *rubiginosum*, 85
Pterygospidea helias, 15
 PTERYGOTA, 60
 " *alata*, 74
 " *Roxburghii*, 74
Ptilocera, 275
 " *amethystina*, 275
 " *continua*, 275
 " *fastuosa*, 275
 " *quadridentata*, 275
 " *smaragdina*, 275
Puccinia, 211, 213, 214, 216, 218, 219, 222
 " *Angelica*, 215
Puccinia, *Arenaria*, 219
 " *Castagnei*, 215, 216, 230
 " *caudata*, 219, 230
 " *coronata*, 213
 " *crassa*, 219, 230
 " *doloris*, 218
 " *Ellisii*, 215, 230
 " *Eulalie*, 216
 " *excelsa*, 216, 230
 " *graminis*, 212, 213
 " *McIntirianus*, 230
 " *occulta*, 230
 " *Penniseti*, 215
 " *Phlomidis*, 217
 " *Pimpinella*, 215, 219
 " *var. Eryngii*, 219
 " *Prainiana*, 229
 " *Prenanthis*, 216
 " *pulvinata*, 219, 220
 " *Rubigo-vera*, 229
 " *Saxifrage*, 218
 " *Saxifrage-ciliatæ*, 218
 " *Saxifrage-micranthæ*, 218
 " *sepulta*, 225
 " *Sorghii*, 214, 215
 " *ustalis*, 217
 " *Wattiana*, 227, 230
Pueraria Candollei, 311, 353, 403
 " *phascoloides*, 311, 353, 403
Pyrameis cardui, 172
Pyrus, 225
Python molurus, 232
Quercus, 341
Radena, 13, 14, 17, 18, 24, 166, 168
 " *juventa*, 16, 24, 167, 168
 " *kamera*, 167, 168, 197
 " *longa*, 24, 32
 " *macra*, 24, 32
 " *nicozarica*, 14
 " *oberthurii*, 158, 161, 167, 197
 " *purpurata*, 167
 " *similis*, 24
 " *volgaris*, 14, 16, 24, 168
Randia longiflora, 316, 354, 403
Ranunculus hirtellus, 217
Rapala tarbas, 179
 " *orscis*, 179
 " *varuna*, 180
Raphiocera, 278
 " *spinithorax*, 278
 **Rasuna leuca*, 162
Rauwolfia serpentina, 318, 354, 397
Rhachicercus, 276
 " *zonotus*, 276
Rhagerrhis producta, 237
Rhagio, 268, 279, 281
 RHAMNEÆ, 307, 345, 349
Rhaphiocera, 273
 " *spinithorax*, 273
Rhinomyza, 267
 " *fusca*, 267

- Rhizophora*, 293
 " *conjugata*, 313, 353, 380
 " *mucronata*, 313, 353, 380
RHIZOPHOREÆ, 313, 345, 349
RHYTHIDEÆ, 263
RUYFRIDI, 263
Rhyphus, 263
 " *maculipennis*, 263
Rhytisma, 340, 358, 389
Riedleia concatenata, 87
 " *corchorifolia*, 87
 " *supina*, 87
 " *truncata*, 87
Rocstelia, 224
Rosa moschata, 222
Ruba, 273
RUBIACEÆ, 316, 344, 345, 349, 396
Rubus, 222
 " *rosafolius*, 221
Rungia parviflora, var. *pectinata*, 322, 355
 " *pectinata*, 370, 377
RTIACEÆ, 305, 345, 349
Salacia, 347
 " *prinoides*, 293, 307, 346, 352, 397
Salamis, 173
 " *anacardii*, 173
 " *anteca*, 173
Salotura genutia, 164
 * " *litoralis*, 164
Salctara nathalia, 190
Salmalia malabarica, 49
Salpinx, 15, 16, 17, 18, 21, 22, 158
 " *kollarii*, 13
 " *leucostictos*, 20, 22, 159
 * " *meizon*, 159
 " *novaræ*, 13, 20
 " *pasithea*, 159
 * " *phane*, 20, 22
 " *splendens*, 13
 " *vestigata*, 20
 " *viola*, 159
 " (*Calliptera*), 22
 * " " *sambavana*, 160
 * " " *sumbana*, 159
 " (*Isamia*) *locii*, 22
 " " *standingerii*, 22
 " (*Selinda*) *eleusina*, 159
 " (*Tiruna*) *ochsenheimeri*, 22
SALTICIDÆ, 3
Salticus, 2, 3
 " *formicarius*, 2, 3
SALTIGRADÆ, 2
SANTALACEÆ, 325, 345, 350
SAPINDACEÆ, 308, 345, 349
Sapindus Danura, 308, 352, 397
SAPOTACEÆ, 318, 345, 349
Sarcotobus globosus, 319, 354, 381, 386
Sarcostigma, 347
 " *edule*, 293, 296, 306, 397
 " *Wallichii*, 306, 352 [379.
Sargassum ilicifolium, 288, 341, 358, 378,
Sargus, 276, 278, 279, 280
 " *afinis*, 280
 " *aurifer*, 279
 " *brevipennis*, 279, 280
 " *ferrugineus*, 280
 " *flaviventris*, 280
 " *formiciformis*, 280
 " *gemmifer*, 279
 " *insignis*, 279
 " *latus*, 280
 " *latifascia*, 279
 " *leoninus*, 280
 " *longipennis*, 279
 " *luridus*, 279
 " *magnificus*, 279
 " *metallinus*, 279, 280
 " *pallipes*, 279
 " *quadrifasciatus*, 279*
 " *rubescens*, 280
 " *rufus*, 280
 " *tenebrifer*, 279
 " *viridiceps*, 279
Satanga, 16
SATYRIDÆ, 11, 12, 24, 158, 168
Saxifraga micrantha, 218
Scævola, 345
 " *Königii*, 297, 317, 354, 380
SCAPHIUM, 60, 74
 " *affine*, 75
 " *Wallichii*, 75
Scathopse, 263
 " *pusilla*, 263
Schoutenia, 96, 105, 106, 107
 * " *glomerata*, 106, 107
 " *hypoleuca*, 107
 * " *Kunsteri*, 106, 107
 " *Mastersii*, 106
 " *ovata*, 107
Sciara, 261, 262, 263
 " *femoralis*, 262
 " *indica*, 262
 " *laticornis*, 262
 " *rufithorax*, 262
 " *solita*, 262
 " *sulcata*, 262
 " *thoma*, 262
Sciudapsus officinalis, 295, 334, 357, 404
Sciophila, 262
 " *tropica*, 262
Scirpus subulatus, 300, 335, 357, 392, 402
SCITAMINEÆ, 329, 345, 350
SCLEROSTOMA, 264
Scoparia dulcis, 321, 370, 355, 377
SCROPHULARINEÆ, 321, 345, 349
Selinda, 16
 " *eleusina*, 158, 159
Semecarpus heterophylla, 295, 397
 " *heterophyllus*, 308, 352
 " *subpanduriformis*, 308, 352, 397
Senecio, 226
 " *rufinervis*, 226

- Sesuvium*, 288
 Portulacastrum, 292, 315, 354,
 379, 380
- SICARIÆ**, 272
- Sida*, 39
- acuta*, 40, 303, 351
 - althæifolia*, 42
 - asiatica*, 42
 - Belare*, 43
 - canariensis*, 41
 - carpinifolia*, 40, 368, 377
 - chinensis*, 41
 - compressa*, 41
 - cordifolia*, 41
 - corynocarpa*, 41
 - cuneifolia*, 87
 - Eteroomischos*, 43
 - fasciculiflora*, 40
- Sida glutinosa*, 40
- herbacea*, 42
 - hirta*, 40
 - indica*, 43
 - lanceolata*, 40
 - micans*, 42
 - mysorensis*, 40
 - nervosa*, 40
 - olens*, 40
 - orientalis*, 41
 - philippica*, 41
 - populifolia*, 43
 - radicans*, 40
 - rhubifolia*, 40, 41
 - " *var. microphylla*, 41
 - " " *obovata*, 41
 - " " *retusa*, 41
 - " " *rhomboidca*, 41
 - " " *scrabrida*, 41
 - rotundifolia*, 42
 - scoparia*, 40
 - Stauntoniana*, 40
 - stipulata*, 40
 - tenax*, 41
 - urticifolia*, 40
- Silvius*, 266
- dimidiatus*, 266
- Silybura brevis*, 232
- nigra*, 232
 - pulneyensis*, 232
 - wood-masoni*, 232
- Sima rufa-nigra*, 2
- Simotes albocinctus*, 235
- catenifer*, 235
 - crassus*, 235
 - cyclurus*, 235
 - dennysi*, 235
 - labuanensis*, 235
 - obscurus*, 235
 - octolineatus*, 236
 - planiceps*, 236
 - purpurascens*, 235
 - theobaldi*, 236
- Simotes trinotatus*, 235
- violaceus*, 236
 - " *wood-masoni*, 231, 235, 250
- Simulium*, 264
- " *indicum*, 264
- Sinthusa*, 180
- " *aspra*, 180
 - " *nasaka*, 180
 - " *virgo*, 180
- Siphonocladus filiformis*, 342, 359, 378
- Siphonodon celastrincus*, 296, 298, 307,
 352, 403
- Smilax macrophylla*, 330, 357
- " *macrophyllus*, 398
- SOLANACEÆ**, 321, 345, 349
- Solanum Melongena*, 321, 355, 370, 372,
 377
- Solva inamona*, 273
- Sophora tomentosa*, 292, 297, 311, 353,
 379, 380
- Sorghum*, 214, 215
- " *vulgare*, 214
- Spalgis epius*, 185
- Spondias*, 382, 396
- " *mangifera*, 295, 308, 352, 403
- Stellaria paniculata*, 219
- Stephegyne*, 288, 386
- " *diversifolia*, 292, 296, 316, 354,
 379, 380
- Sterculia*, 58, 59, 69, 382
- " *affinis*, 61, 75
 - " *alata*, 60, 73, 295, 305, 352, 388,
 403
 - " *angustifolia*, 60, 68, 69, 70, 71
 - " *armata*, 62
 - " *Balanghas*, *var. glabrescens*, 70
 - " " *mollis*, 69
 - " *bicolor*, 60, 67, 68
 - " *campanulata*, 61, 76, 305, 352,
 388, 403
 - " *campanulatum*, 76
 - " *coccinea*, 64, 74
 - " *colorata*, 60, 71, 73, 305, 352,
 403
 - " " *var.*, 73
 - " *ensifolia*, 60, 70
 - " *fulgens*, 60, 72, 73
 - " *Heymii*, 74
 - " *hyposticta*, 59, 64
 - " *Jackiana*, 70
 - " *javanicum*, 76
 - " *Kunsteri*, 60, 65
 - " *laevis*, 59, 63, 65
 - " *linearicarpa*, 60, 74
 - " *macrophylla*, 59, 63
 - " *Maingayi*, 67
 - " *mollis*, 69, 70
 - " *ornata*, 59, 62
 - " *pallens*, 73
 - " *parviflora*, 66, 67, 70, 305, 352,
 403

- Sterculia, parvifolia*, 60, 64
 " *pubescens*, 60, 71
 " *rubiginosa*, 60, 69, 296, 382
 " " *var.* 380, 403
 " " *glabrescens*, 70, 305, 352
 " *scaphigera*, 60, 74
 * " *Scortechinii*, 60, 67
 " *tubulata*, 61, 76
 " *villosa*, 59, 61, 62, 296, 304, 352, 403, 404, 405
 STERCULIACEÆ, 57, 304, 345, 349
 STERCULIÆ, 58
Stichophthalma, 18
Stictoplea, 13, 16, 17
 " *lacordairei*, 13, 160
 * " *melolo*, 160
 STRATIOMYDÆ, 272
 STRATIOMYDÆ, 272
 STRATIOMYDI, 272
 STRATIOMYDINÆ, 272
Stratiomys, 273, 274, 275, 277
 " *apicalis*, 277
 " *barca*, 277
 " *bilineata*, 273, 278
 " *difusa*, 277
 " *flavoscutellata*, 277
 " *garutæ*, 277
 " *inanimis*, 277
 " *lutatus*, 277
 " *minuta*, 277
 " *pusilla*, 277
 " *rufipennis*, 277
 " *solenis*, 277
 " *viridana*, 277
Strobilanthes phyllostachyus, 297, 322, 355, 404
Strophanthus, 390
 " *Wallichii*, 319, 354, 388, 403
Strychnos, 396
Suaetus, 196
 * " *chilon*, 196
 " *gremius*, 197
Subula, 272
 " *calopodata*, 272
 " *flavipes*, 272
 " *inamena*, 273
 " *vittata*, 272
Suragina, 282
 " *signipennis*, 282
Surendra quercetorum, 179
Sus andamanensis, 308
Symbrenthia hippoclus, 172
 * *Symphædra ngle*, 177
 " *dirtea*, 178
Synageles picata, 3, 4
Synemosyna formica, 4
Syrphus, 274
 TABANICA, 264
 TABANIDÆ, 264
 TABANIDES, 264
 TABANIDI, 264
 TABANIDII, 264
 TABANII, 264
 TABANINA, 264
 TABANINÆ, 264
 TABANITES, 264
Tabanus, 264, 265, 266, 267
 " *abscondens*, 270
 " *administrans*, 271
 " *albilateralis*, 267
 " *albimediæ*, 269
 " *alboscutatus*, 270
 " *albulus*, 269
 " *amænus*, 269
 " *apicalis*, 271
 " *ardens*, 268
 " *auricinctus*, 267
 " *auriglanma*, 268
 " *avrotestaccus*, 268
 " *basalis*, 267
 " *brunneus*, 267
 " *bubali*, 270
 " *bucolicus*, 271
 " *ceylanicus*, 271
 " *cinnamomeus*, 270
 " *clausicella*, 270
 " *cœrulescens*, 267
 " *confucius*, 270
 " *consanguineus*, 267
 " *consocius*, 269
 " *crassus*, 269
 " *dives*, 271
 " *dorsolinea*, 267, 268
 " *equestris*, 271
 " *explicatus*, 268
 " *felderi*, 271
 " *fulvissimus*, 271
 " *fumifer*, 270
 " *fumipennis*, 271
 " *furunculigenus*, 270
 " *geniculatus*, 271
 " *hilaris*, 269
 " *hirtus*, 269
 " *hoang*, 270
 " *hybridus*, 268
 " *ignobilis*, 270
 " *immanis*, 268
 " *incultus*, 271
 " *inscitus*, 268
 " *internus*, 269
 " *javanus*, 268
 " *juvencus*, 269
 " *justorius*, 270
 " *lastianus*, 272
 " *lectum*, 270
 " *leucosparsus*, 271
 " *mandarinus*, 271
 " *megalops*, 268
 " *melanognathus*, 272
 " *mentitus*, 268

- Tabanus, minimus*, 271
 " *monoculus*, 270
 " *navus*, 270
 " *nicobarensis*, 271
 " *nigropictus*, 269
 " *nigrolectus*, 271
 " *obconicus*, 269
 " *optatus*, 270
 " *orientalis*, 267
 " *orientis*, 268
 " *partitus*, 270
 " *pauper*, 270
 " *perlina*, 269
 " *puella*, 269
 " *pusillus*, 267
 " *pyrausta*, 268
 " *pyrchus*, 269
 " *rubicundus*, 267
 " *rubidus*, 268
 " *rubiginosus*, 269
 " *rufiventris*, 268
 " *sanguineus*, 269
 " *serus*, 270
 " *servillei*, 267
 " *simplicissimus*, 270
 " *sinicus*, 268
 " *striatus*, 267, 268.
 " *sumatrensis*, 267
 " *tenebrosus*, 268
 " *tensus*, 269
 " *tristis*, 271
 " *umbrosus*, 269
 " *unicentris*, 268, 270
 " *vagus*, 268, 269
 " *variegatus*, 271
 " *virgo*, 267
 " *yao*, 269
Tabernaemontana crispa, 292, 319, 354, 381.
Tacca pinnatifida, 292, 329, 357, 381, 384.
 TACCACEÆ, 329, 345, 350
Tachyris, 190
Taxaris, 18
Togiades, 195, 196
 " *alica*, 195
 " *atticus*, 32
 * " *brasidas*, 195
 " *helferii*, 195
Tajuria maculata, 36
Tamarindus indica, 312, 353, 369, 372, 377.
Tanyderus, 260
 " *ornatissimus*, 260
Tanyglossa, 265
Tanyptera, 254
Tanypus, 253
 " *crux*, 253
 " *cyanomaculatus*, 253
 " *melanurus*, 253
 " *nigrocinctus*, 253
 " *ornatus*, 253
 " *pardalis*, 253
Taraka hamada, 36
Tarrietia, 58, 77
 * " *Curtisii*, 78
 " *javamica*, 78
 * " *Kunstleri*, 77, 79
 " *penangiana*, 77
 * " *perakensis*, 77, 78
 " *simplicifolia*, 77, 78, 79
Tarucus plinius, 181
 " *theophrastus*, 181
Telicota gola, 196
 " *nigrolimbata*, 196
Terias candida, 186
 " *drona*, 186
 " *harina*, 29, 186
 " *hecabe*, 29, 186
 " *sari*, 29, 186
Terminalia, 288
 " *biolata*, 296, 314, 353, 388,
 403
 " *Catappa*, 136, 292, 296, 297,
 313, 353, 366, 379, 380,
 382
 " *moluccana*, 135, 136
 " *procera*, 314
Tetrapoda, 11
T eucholabis, 261
 " *bicolor*, 261
 " *determinata*, 261
 " *fenestrata*, 261
 THALAMIFLORE, 301, 344
Thalassomyia, 252
 THECLINE, 32, 179
Thelephora incrustans, 339, 358, 389
Theriopectes, 267
Thespesia, 39, 47, 288, 298
 " *populnea*, 47, 292, 298, 303,
 351, 380
 **Thriz*, 35
 " *gama*, 35
Thuarea sarmentosa, 292, 295, 296, 336,
 345, 357, 379, 381
Thumbergia, 347
 " *laurifolia*, 294, 296, 322, 355,
 403
Thylacosoma, 273
 " *amboinense*, 273
 TILIACEÆ, 95, 305, 345, 349
Tilicota mexoides, 196
 TILIÆ, 96
Tinda, 275
 " *indica*, 275
 " *modifera*, 275
Tipula, 252, 253, 254, 255, 256, 259, 260,
 262, 263, 264
 " *broddignagia*, 257
 " *castanea*, 256
 " *congrua*, 256
 " *fulvipennis*, 256
 " *sumifensis*, 257
 " *javana*, 256
 " *javensis*, 256

- Tipula, leucopyga*, 257
 " *longicornis*, 257
 " *melanomera*, 256
 " *monochroa*, 256
 " *nora*, 256
 " *pedata*, 256
 " *pilosula*, 257
 " *praeputens*, 256
 " *reposita*, 256
 " *serrata*, 257
 " *umbrina*, 256
 " *venusta*, 256
 " *vicaria*, 256
 " *vilis*, 257
Tipularia, 254
 " *florales*, 263
 " *Fungicola*, 261
 " *Fungitorae*, 261
 " *Gaëlicolae*, 253
 " *Latipennes*, 263
 " *Muscaformis*, 263
 TIPULARIDES, 254
 TIPULIDE, 252, 254
 TIPULIDES, 254
Tirumala, 16
 " *linniace*, 166
 " *melissa*, 166
Tiruna ochsenheimieri, 22
Zomyris oxiata, 247
Tournefortia argentea, 292, 297, 320, 346,
 355, 379, 381
Tozocera, 276
 " *limbiventris*, 276
Trachischium fuscum, 231, 233
 " *guentheri*, 233
 " *rubricenter*, 233
 " *tenuiceps*, 231
Trepsichrois, 15, 18, 158
 * " *dongo*, 160
 * " *elwesii*, 161
 * " *malakoni*, 20, 22
 " *midamus*, 15, 16, 161
 " *mindanensis*, 161
 " *verhuelli*, 22
Trichocarya, 115
Trichocera, 260
 " *ocellata*, 260
Trichochæta, 278
 " *nemoteloides*, 278
Trichosanthes, 347
 " *palmata*, 296, 315, 354, 397
Trichospermum, 96, 118, 119
 " *javanicum*, 119
 " *Kurzii*, 119
Tremeresurus cantoris, 231, 249
 " *gramineus*, 248, 249, 250
 " *mutabilis*, 250
 " *purpureomaculatus*, 248, 249,
 250.
Triumfetta, 96, 115
 " *angulata*, 116
Triumfetta, angulata var. *acuminata*, 116
 " *annua*, 116, 117
 " *Bartramia*, 116
 " *glandulosa*, 116
 " *indica*, 117
 " *oblonga*, 417
 " *oblongata*, 116
 " *ovata*, 117
 " *pilosa*, 116, 117
 " *pilosa*, var. 116, 117
 " *polycarpa*, 116, 117
 " *rhomboidea*, 116
 " *tomentosa*, 116, 117
 " *trichoclada*, 117
 " *trilocularis*, 116
 " *vestita*, 116, 117
Trochetia contracta, 80
 * *Trochostoma andamanense*, 203
 " *antarcticum*, 203
Tronga, 22, 163
 " *niasica*, 22
Tropidonotus, 242
 " *angusticeps*, 240, 241
 " *chrysargus*, 239, 240
 " *himalayanus*, 240
 " *hydrus*, 240, 241
 " *mortoni*, 242
 * " *nicobarensis*, 241
 * " *nicobaricus*, 231, 250
 " *nigrocinctus*, 231, 239, 240
 " *parallelus*, 241
 * " *pealii*, 231, 241, 250
 " *piscator*, 240
 " *platyceps*, 231
 " *plumbicolor*, 240, 241
 " *rhodomelas*, 242
 " *subminiatus*, 240
 " *trianguligerus*, 242
Turbinaria, 288
 " *ornata*, 288, 341, 358, 378
Turraa trichostylis, 95
Typhlops acutus, 232
 " *andamanensis*, 231
 " *bothriorhynchus*, 232
 " *diardi*, 232
 " *persicus*, 231
 " *theobaldianus*, 231, 232
Udaspes folus, 32
 UREDINEÆ, 211, 219
Uredo Agrimoniar, 229
 * " *Colebrookiae*, 227
 * " *Ehretiae*, 228
 * " *Fici*, 225
 " *ficiola*, 225
 * " *Ichnocarpi*, 228
 * " *Ipomæae*, 228
 * " *Pilcae*, 228
Urena, 39, 43
 " *cana*, 43
 " *heterophylla*, 44
 " *Lappago*, 44

- Urena, lobata*, 43, 294, 303, 346, 351, 368,
 372, 377, 393, 394
 " " var. *scabriuscula*, 44
 " " " *sinuata*, 43
 " *morifolia*, 44
 " *muricata*, 44
 " *palmata*, 43
 " *scabriuscula*, 44
 " *sinuata*, 43
 " *tomentosa*, 43, 44
 URENEE, 39
Uromyces, 211, 213, 228
 * " *Agrappri*, 212
 " *ambiens*, 213, 214
 " *Myristica*, 213
 " *Pisi*, 212
 " *puberulus*, 213
 * " *Vigna*, 211, 212, 330
 URTICACEE, 327, 344, 345, 350
Fadebra, 9, 11, 20, 21
 * " *palmedo*, 162
Valonia, 342
 " *confervoides*, 342, 359, 378
 " *fastigiata*, 342, 359, 378
Vappo, 276
 VASCULARIS, 336, 344
Vaucleria, 342, 359, 378
Ventilago calyculata, 307, 352, 388, 403
 VERBENACEE, 323, 345, 350
Vernonia cinerea, 297, 317, 345, 354, 369,
 372, 377, 388
 " *divergens*, 297, 317, 354, 388
 VESICULOSIDE, 282
Vigna, 212
 " *lutea*, 290, 292, 311, 353, 379, 380
 " *vesillata*, 211
 VISLACEE, 302, 345, 349
Viscacia concatenana, 87
 " *corchorifolia*, 87
 " *indica*, 88
 " *javanica*, 88
 " *supina*, 87
 " *umbellata*, 88
 VITEE, 290, 384
 " *Negundo*, 293, 298, 323, 355, 379,
 381, 384
 " *pubescens*, 323, 355, 381
 " *sumatrana*, 355
 " *Wimberleyi*, 323, 355, 281
Vitis caruosa, 307, 352, 397
 " *pedata*, 296, 307, 352, 397
 " *pentagona*, 307, 352, 397
Vittaria elongata, 337, 358, 389
 WALLACEA, 276
 " *argentea*, 276
Waltheria, 58, 88
 " *americana*, 88
 " *elliptica*, 88
 " *indica*, 88
 * *Weberia Kurzii*, 316, 354, 403
Wedelia scandens, 317, 354, 380
Xenodochus, 211, 222, 223
 " *carbonarius*, 225
 " * *Clarkianum*, 230
 * " *Clarkianus*, 222
Xylaria, 300, 340
 * " *clararioides*, 340, 358, 404
Xylocoryne, 310
 XYLOPHAGEI, 272
 XYLOPHAGI, 272
 XYLOPHAGIDÆ, 272
 XYLOPHAGINA, 272
Xylophagus, 272, 274
 " *brunneus*, 272
Xyphura, 254
Yoma, 73, 174
 " *sabina*, 173
Ypithima aphnius, 169, 170
 " *asterope*, 169
 " *baldus*, 170
 * " *leuce*, 169
 " *panocus*, 169
 " *philomela*, 169, 170
 " *scypra*, 169
Zamenis diadema, 238
 " *korros*, 237
 " *ladaccensis*, 231, 238
 " *ventrimaculatus*, 238
Zanichellia, 299
 " *palustris*, 334, 357, 392
Zaocys fuscus, 239
 * " *nigromarginatus*, 231, 239
 * " *tenasserimensis*, 231, 238, 250.
Zopyrus, 239
Zea, 214, 215
 " *Mayo*, 214
Zemeros phlegyas, 178
Zephyrus distortus, 34
Zelides agamemnon, 31, 193
 " *euryppylus*, 193
 " *sarpedon*, 31, 193
Zeylania, 209
Zinzypa distorta, 34
Zingiber, 296, 329, 346, 357, 398
 " *corollianum*, 329
Zizera, 180
 " *karsandra*, 29
 " *lysiocoe*, 180
 " *pygmaea*, 180
Zizyphus (Euphria), 307, 352, 397



H. L. D. and G. C. C. del.

NEW ORIENTAL BUTTERFLIES.






K.P. Das del.


NEPETA BELLEVII P. Prain

A. D. Mollis.



UREDINEAE FROM SIMLA

Illustrated by G. S. S. S. S.



CILIA FROM SIMLA

Copyright 1891

ZOOLOG

BO