



LIBRARY OF THE UNIVERSITY OF TORONTO

UNIVERSITY OF TORONTO LIBRARY

VOLUME XVII

UNIVERSITY OF TORONTO LIBRARY

11

UNIVERSITY OF TORONTO LIBRARY

VOLUME XVII

20

UNIVERSITY OF TORONTO LIBRARY

UNIVERSITY OF TORONTO LIBRARY

UNIVERSITY OF TORONTO LIBRARY

UNIVERSITY OF TORONTO LIBRARY

UNIVERSITY OF TORONTO LIBRARY

UNIVERSITY OF TORONTO LIBRARY

UNIVERSITY OF TORONTO LIBRARY

KNO.

11

PHYSICAL SCIENCE DEPARTMENT

UNIVERSITY OF TORONTO LIBRARY



RECORDS
of the
INDIAN MUSEUM

(Vol. XVII.)

CATALOGUE
OF
ORIENTAL AND SOUTH ASIATIC
NEMOCERA.

By
E. BRUNETTI.

Calcutta :

PUBLISHED BY THE DIRECTOR, ZOOLOGICAL SURVEY OF INDIA,
PRINTED AT THE BAPTIST MISSION PRESS.

JUNE (1920.)



Acc. no. P707
Dt. 15/5/07

591-5405
ANN

AUTHOR'S PREFACE.

—◇—

The preparation of the present Catalogue has extended over several years, and it has been laid aside frequently in favour of more urgent work; this fact, in conjunction with the wholesale revisions in the classification of the Culicidae, rendered necessary in its progress by the publication of numerous papers by various authors, and the large number of additions and alterations in Cecidomyiidae, Chironomidae and Tipulidae combining to render the task one of considerable difficulty.

It follows that omissions and inaccuracies must have occurred, and, as a matter of fact, they are continually coming to light. The catalogue is not made up to any particular date, all available names being incorporated up to about the middle of 1919, but the printing has been a matter of many months and it has therefore been impossible to include, even in the "Errata and Addenda," the very latest accessions to the Oriental Fauna. The Author trusts that these facts may be taken into consideration. Readers will very greatly assist him by notifying errors as they are discovered.

E. BRUNETTI.

Calcutta, May 6th, 1920.

I. CATALOGUE OF ORIENTAL AND SOUTH ASIATIC NEMOCERA.

By E. BRUNETTI.

INTRODUCTION.

Van Der Wulp's "Catalogue of the Diptera of South Asia" (1896) has been the only one available since its publication, and the want of a more up-to-date work has probably been felt by all workers in this somewhat neglected order of insects. The present effort is, so far as the first suborder goes, an endeavour to fill the breach *pro tempore*, for the study of oriental diptera is very obviously still in its infancy.

The arrangement of families and subfamilies is the one that appears to me most natural, and the nomenclature adopted is that which has, with few exceptions, been used by the leading dipterologists of the past three quarters of a century. The opportunity may here be taken to state my unshakeable adherence to both the present classification and nomenclature. The continual shuffling of names under the guise of priority is a wholly unnecessary handicap to effective study, and the only way to establish finality is to retain the nomenclature which has been found satisfactory by practically all the greatest writers on diptera. Osten Sacken and Verrall openly confessed to "continuity before priority" and I cannot but whole-heartedly agree with them in the best interests of the science itself.

As has been pointed out by others besides myself, Meigen's paper of 1800 (which has been the chief cause of the suggested general overturning of the principal generic and family names throughout the order by the strict priorist) was not binomial, no species being given as representative of the genera described. This fact, therefore, in itself entirely invalidates its acceptance under the customary laws of zoological literature, and the adoption of the names contained in it seems to suggest merely a morbid craving for continual change.

It is not out of place to recall Verrall's view that it is incumbent upon us to accept the methods of the writers of old and the etiquette of their times. Meigen's "1800 paper" as it has been called was wholly preparatory; the author himself almost entirely ignored it in his later works, and his right to rename his genera was never questioned by his *confidères* at the time he wrote.

My own fixed policy is to retain the nomenclature as employed by the principal authors of the last century, Schiner, Loew, Meigen, Macquart, Walker, Osten Sacken, Verrall and their contemporaries, and family and subfamily names should be held in

Acc. No. P707
Dt. 15/5/07



violable for ever. Generic and specific names of long standing should be similarly regarded and the strict priorist discouraged in every possible way. Within the last few years I read somewhere that the great majority of American and German zoologists were opposed to the rigid application of the law of priority yet the International Commission appear to be mainly composed of those of the opposite way of thinking.

Some endeavour has been made to ascertain the location of types but for various reasons the results have not been over-satisfactory. My best thanks are due and tendered to those who have assisted me in this matter, notably Mr. Edwards of the British Museum and Mr. F. V. Theobald respecting many of his types in Culicidae.

A number of types (though of which species is by no means certain) of Walker and of Doleschall described from the Malay Peninsula and the East Indian Islands are definitely known to be lost and a certain number of those of Osten Sacken have been for many years, presumed lost also. In cases where the location of the type is not stated in the present Catalogue it simply means that no information has been obtained about it. All types of my own species, unless otherwise stated, were in good condition at the time of description.

REFERENCE LIST OF LOCALITIES.

- Albay, a Province in Luzon.
 Angeles, Pampanga, Philippine Is.
 Armageon, east coast, India.
 Aru (Aroe), one of the East Indian Islands.
 Astrolabe Bay, Papua.
 Aurangabad, Hyderabad State.
- Badrinath, 10,200 ft., Garhwal Dist., United Provinces, India.
 Bago, 450 ft., Negros Occidental, Philippine Is.
 Bakloh, 4,500—5,000 ft., Punjab.
 Balighai, near Puri, Orissa.
 Bandaong (Bandong), Java.
 Bangalore, 3,000 ft., Mysore State, S. India.
 Barogh, 5,000 ft., Simla Dist., Western Himalayas.
 Bataan, one of the Philippines.
 Batticaloa, district on E. Coast of Ceylon.
 Bayembang, Pangasinan, Philippine Is.
 Benguet, Province in Luzon.
 Berars, Province in the Deccan, India.
 Berhampur, Murshidabad Dist., Bengal.
 Berlinhafen, Papua.
 Bhim Tal, 4,500 ft., Kumaon Div., W. Himalayas.
 Bhogaon, Purneah Dist., Bihar, India.
 Bhowali, 5,700 ft., Kumaon Div., W. Himalayas.
 Buitenzorg, Java.

Bukit Kutu, 3,400 ft., Federated Malay States.
Bulaco, Mindoro Is., Philippines.

Camerines Sur, Province in Luzon.

Canara, see North Kanara.

Canlaon Volcano, 2,300 ft., Negros Is., Philippines.

Castle Rock, N. Kanara District, Western India.

Cavite, near Manila (Luzon).

Cebu, one of the Philippines.

Chilka Lake, east coast of India.

Chitlong, Nepal.

Chittagong, Eastern Bengal.

Cochin State, South India.

Coconada, East Coast, Madras Presidency.

Coonor, 6,000 ft., Nilgiri Hills, Madras Presidency.

Coromandel, 2,500 ft., east coast of India.

Cottabato, Mindanao, Philippine Is.

Dacca, district in Eastern Bengal.

Damukdia, Damukdia Ghat, on River Ganges (Calcutta-Darjiling route), N. Bengal.

Darjiling, Eastern Himalayas: most of the localities given in this catalogue are from the town and its immediate neighbourhood, about 6,900 ft. The altitude of the district ranges from about 1,000 to 10,000 ft.

Dawna Hills, 2,000—3,000 ft., Amherst Dist., Burma.

Deccan, a general term for Hyderabad State and adjacent districts, India.

Deesa, Cantonment in Bombay Presidency.

Dehra Dun, foot of Mussoorie Hills, W. Himalayas.

Depok, Java.

Deslac Island, off coast of Papua.

Dharampur, 5,000 ft., Simla District.

Dharamsala, West Himalayas.

Dindings, Straits Settlement.

Djokjokarta, Java.

Dondra, extreme south of Ceylon.

Dorey, a small island off Papua.

Dum Dum, near (and N. of) Calcutta.

Ellichpur, Berar, Central India.

Ennur, Madras Coast.

Erima, Astrolabe Bay, Papua.

Ernakulam, Cochin.

Etawah, district in United Provinces, India.

Etna Bai, South Papua.

Ferozepore, district and city in the Punjab.

Foo Chow, South China.

Fort McKinley, Luzon, Philippine Is.

Friedrich Wilhelmshafen, Papua.

Galle, south coast of Ceylon.
 Gangtok, 6,000 ft., Sikkim State.
 Garhwal Dist., base of Western Himalayas.
 Ghoom, 7,500—8,000 ft., above Darjiling.
 Goa, district on west coast of India (Portuguese).
 Gombong, Mid-Java.
 Gonda, N. India, south of Nepal.
 Gopkuda Island, Chilka Lake, east coast of India.
 Gorontalo, Malay Archipelago.
 Guimaris Is., Philippines.
 Gunung Ungaren, Java.

Hakgala, 7,000 ft., Central Ceylon.
 Hajonoy, Luzon.
 Hoshangabad, Central Provinces, India.
 Huon Gulf, Papua.

Jalpaiguri, N. Bengal plains, south of Darjiling.
 Jeypore, State in Madras Presidency.
 Jhansi, district in United Provinces, India.
 Jhaunsa, Tons Valley, Western Himalayas.
 Jolo, Island and City, Philippines.
 Jubbelpore (Jubbulpur, etc.), Central Provinces, India.
 Jugra, 1,000 ft., on solitary hill on Langat Riv., Federated Malay States.

Kandy, 1,600—1,700 ft., Central Ceylon.
 Kangra Valley, 4,500—4,800 ft., Punjab.
 Karachi, city on west coast of India, Sind.
 Karwar, Coast Town, Bombay Presidency.
 Kalimpong, 2,000—5,500 ft., Darjiling District.
 Kasauli, 6,300 ft., Simla District.
 Katihar, Purnea Dist., Bihar, India.
 Katmandu, Nepal.
 Kawkareik, base of Dawna Hills, Tenasserim.
 Key Islands, Eastern Malay Archipelago.
 Khasi Hills, Assam, S. of Brahmaputra R.
 Klang, on Klang Riv., Federated Malay States.
 Kuala Lumpur, Selangor State, Federated Malay States.
 Kuching, British N. Borneo.
 Kufri, 8,000 ft., Simla District, Western Himalayas.
 Kumaon, Area in Western Himalayas, including Naini Tal and Bhim Tal.
 Kurseong, about 5,000 ft., below Darjiling.

Laguna, Province in Luzon.
 Lahago, Central Lias Island.
 Lahore, Punjab.
 Lake Chilka, east coast of India.
 Laos, Siam.

Leyte, one of the Philippines.
 Lushai Hills, Assam, on N. E. Frontier of India.
 Luzon, the principal island in the Philippines, in which is
 Manila.

Macassar (Makassar), Celebes.
 Maddathorai, Travancore State, S. India.
 Madhupur, Bengal.
 Madulsima, 2,500-3,000 ft., Central Ceylon.
 Makerian, Punjab.
 Mandalay, district and city in Upper Burma.
 Manila, Luzon.
 Manipur, State in Assam.
 Margherita, Assam.
 Marinduque, one of the Philippines.
 Masbata, one of the Philippines.
 Maskeliya, 4,000 ft., Central Ceylon.
 Matheran, Bombay Presidency, India.
 Matiana, 8,000 ft., Simla District, Western Himalayas.
 Mazbat, Mangaldai Dist., Assam.
 Medan, Sumatra.
 Meenglass, Dooars, Jalpaiguri, Bengal.
 Merauke, South Papua.
 Mergue, Assam.
 Mergui, Tenasserim, Lower Burma.
 Mian Mir, near Lahore (*q.v.*).
 Mindanao, one of the Philippines.
 Mindoro, one of the Philippines.
 Moerah Tewah, Borneo.
 Moradabad, United Provinces, India.
 Moulmein, Amherst District, Tenasserim, Lower Burma.
 Mount Hauseman, Astrolabe Bay, Papua.
 Mount Singalang, Sumatra.
 Mount Siya-Siya; *v.* Canlaon Volcano.
 Mozufferpore, Bihar, India.
 Muina, Papua.
 Mungphu, 5,000 ft., Darjiling District, Eastern Himalayas.
 Murree Hills, W. Himalayas, Punjab.
 Mussoorie, 6,000-7,000 ft., Western Himalayas.
 Mysore, State in S. India.

Nagpur, town and district in Central Provinces, India.
 Naini Tal, 6,500-7,000 ft., Kumaon, W. Himalayas.
 Nara Ghat, Nepal, near Indian Frontier.
 Narcondam Island, Bay of Bengal, 80 miles East of North
 Andaman Island.
 Nedumangad, 10 miles N.E. of Trivandrum (*q.v.*).
 Negros (Occidental), one of the Philippines.
 Nicobar Islands, south of the Andamans.
 Nilgiri Hills, about 3,000-8,000 ft., Madras Presidency.

Nongpoh, Assam.

North Kanara Dist., on Goa Frontier, Western Coast of
Bombay Presidency.

N.B.—South Kanara Dist. is in Madras Presidency.

Ootacamund, Nilgiri Hills, South India, alt. 7,000--8,000 ft.
Orissa, East Coast of India.

Padang, W. Sumatra.

Pahang, Malay States.

Palawan, one of the Philippines.

Pallode, Travancore State, S. India.

Pampanga, Province in Luzon.

Pangasinan, Province in Luzon.

Pangerango, Java.

Pangkor Haut, Malay Peninsula.

Parang, Mindanao, Philippine Is.

Paresnath Hill (4,800 ft.), Bihar.

Paumonu River, Papua.

Peradeniya, Ceylon, 4 miles from Kandy.

Perak, Federated Malay States.

Phagu, 8,700—9,000 ft., Simla District, Western Himalayas.

Phrapatoon, Siam.

Port Canning, 30 miles S. of Calcutta, on Matla River.

Port Swettenham, Selangor, Federated Malay States.

Preanger, Borneo.

Pundaluoya, 5,000 ft., Mid-Ceylon.

Puri, Orissa Coast.

Purneah, district in Bihar, India.

Pusa (Agricultural Research Institute), Bihar, India.

Quilon, Coast Town, Travancore, S. India.

Rajmahal, N. Bengal.

Rajpore, base of Mussoorie Hills, Western Himalayas.

Rajshahi, North Bengal.

Ranikhet (Ranaghat, Reneghat), 4,000 ft., Kumaon, Western
Himalayas.

Raub, Pahang, Federated Malay States.

Rawas, Mid-Sumatra.

Rio Baco, Chicago, Mindoro, Philippine Is.

Rizal, Province in Luzon.

Samar, one of the Philippines.

Sambalpur, Central Provinces, India.

Selangor, Federated Malay States.

Seleo Berlinhafen, Papua.

Selve, Papua.

Semarang, Java.

Senchal, 8,000 ft., Darjiling District, E. Himalayas.

- Serang Tjimahi, Java.
 Shahjahanpur, district in United Provinces, India.
 Shamnagar, Bengal.
 Shan Hills, Upper Burma.
 Shaohyling, S. China.
 Shasthancotta, Travancore State, S. India.
 Shencotta, Travancore State, S. India.
 Shillong, 4,900—6,400 ft., Assam.
 Sibsagar, Assam.
 Sikkim (Sikkim), State in Eastern Himalayas, often confused
 with Darjiling District.
 Siliguri, base of Darjiling Hills, 200 ft.
 Simbang, Huon Gulf, Papua. [alayas.
 Simla, 7,000—8,000 ft., town in Simla District, Western Him-
 Singla, 1,500 ft., Darjiling District.
 Soekaboemi, City on E. extremity of Java.
 Soerabaya, East Java.
 Soeroelangoen, Sumatra.
 Soondrijal, Nepal.
 Stephansort, Papua.
 Sukna, 500 ft., base of Darjiling Hills.
 Sukwani, Nepal, near Indian Frontier.
 Sula, Indo-Malay Archipelago.
 Sureil, 5,000 ft., Darjiling District.
 Sylhet, district in Assam.
- Taiping, Perak, Federated Malay States. [dency.
 Talewadi, near Castle Rock, N. Kanara District, Bombay Presi-
 Tamsui, Formosa.
 Tandjong, N.E. Sumatra.
 Tandjong Priok, near Batavia, Java.
 Tandong, 4,000 ft., Tenasserim.
 Tayabas, Province in Luzon.
 Tellschong, Nicobar Islands.
 Tenmalai, Western Ghats, Travancore State.
 Thaumaspur, Lower Nepalese Himalayas.
 Thayetmyo, Upper Burma.
 "The Gap," Pass between Selangor and Pahang, Federated
 Malay States.
 Theog, 8,000 ft., Simla District, Western Himalayas.
 Tonglu, 10,000 ft., Darjiling Distr.
 Tosari, E. Java.
 Travancore, State in S. India.
 Trivandrum, Capital of Travancore State, S. India.
 Trincomalee (Hot Wells), 8 miles inland, east coast of Ceylon.
- Ukhrul, 6,400 ft., Manipur State, Assam.
 Ulu Gombok, 13 miles from Kuala Lumpur, Selangor.
 Ulu Klang, near Kuala Lumpur.
 Umling, Assam.

Waigiou, one of the East Indian Islands.
 Waria River, British Papua.
 Weligama, south coast of Ceylon.
 West Lake, Hangchow (Hankow), South China.
 Wonosobo, Java.

Yomba, Papua.
 Yunnan, Province in South China on Burmese Frontier.

Family CECIDOMYIIDAE.

It should be remarked respecting the nomenclature of this family that the alterations suggested by Dr. Karsch, and accepted by Rübsaamen, Keiffer and others, have been vigorously opposed by both Aldrich and Osten Sacken. Aldrich says (Cat. N. Amer. Dipt., 152): "The practical effect of the changes adopted by Karsch, Rübsaamen and Kieffer is to apply the name *Cecidomyia* to the *Diplosis* of Osten Sacken's catalogue, and to divide up his *Cecidomyia* between *Rhabdophaga* and *Dasyneura*. the adoption of the changed nomenclature will leave few members of the family in the same genera to which Osten Sacken referred them."

Osten Sacken had expounded and, I think, fully substantiated his views (Ent. Month. Mag., 1901, p. 40) and concluded as follows: "What I would propose now would be to restore as far as possible the original concept of *Cecidomyia* of Meigen and Latreille, and to consider the very numerous new genera, since formed at its expense, as *subgenera*."

My own view is identical, but having no practical experience of the family I refrain from inaugurating the change in the present catalogue.

It must also be noted that species of which the adult form is not known are not included as an integral part of the catalogue, but the *names* of such species are given, with a brief reference to the works in which they are described.

Subfamily CECIDOMYINAE.

Sect. LASIOPTERARIAE.

Kieff., Gen. Ins., Fasc. 152, 18, tab. gen.

Lasioptera, Mg.

Syst. Besch. I, 88 (1818).

Sch., F. Austr. II, 406: Kieff., Gen. Ins., Fasc. 152, 30: Auctt. var.

GENOTYPE, *Cecidomyia albipennis* Mg., according to Coquillett's designation in 1910.

Diomyza, Megerle in Mg., Syst. Besch., I, 89 (1818).

GENOTYPE, *Lasioptera picta* Mg., by original designation.

bryoniae Sch., Nov. Reise, Dipt., 5, ♂ ♀, pl. i, 2, wing, 3, ant. (1868).

Madras. Type in Vienna Museum.

textor Kieff., Ann. Soc. Sci. Brux. XXIX, 155, ♀ (1905).

Bengal. Type in Kieffer's collection. Kieffer erects two species, *longispatha* and *trilobata* in Marcellia VII, but only the gall and larva of each is described.

fluitans Felt, Ent. News, XXVIII, 73, ♂ ♀ (1917).

Coimbatore, 3-xii-15 [Y. R. Rao]. From galls in *Panicum fluitans*. "Type Cecid. 2779."

Prolasioptera, Kieff.

Bull. Soc. Hist. Nat. Metz XXVIII, 47 (1913).

Kieff., Gen. Ins., Fasc. 152, 33.

GENOTYPE, *Lasioptera niveocincta* Kieff., by original designation.

javanica Kieff. and Leeuwen-Reijnvaan, Marcellia IX, 123, ♂ ♀ (*Lasioptera*) (1909).

Java. Type in Kieffer's collection.

Sect. OLIGOTROPHARIAE.

Kieff., Gen. Ins., Fasc. 152, 34, tab. gen.

Microperrisia, Kieff.

Neue Gallmucken Gattungen, Bitsch, 2 (1912).

GENOTYPE unknown to present author.

pulvinariae Felt, Jour. N. Yk. Ent. Soc. XXIII, 174 (1915).

Philippine Is.

Panteliola, Kieff.

Bull. Soc. Hist. Nat. Metz XXVIII, 49 (1913).

Kieff., Gen. Ins., Fasc. 152, 46.

GENOTYPE, presumably *P. haasi*, sp. nov., *loc. cit.*

One species only (*P. haasi*, Kieff., *loc. cit.*) has been described of this genus, but the imago is not known. It is from Bengal.

Oligotrophus, Latr.

Hist. Nat. Crust. Ins. XIV, 288 (1805).

Kieff., Gen. Ins., Fasc. 152, 49.

GENOTYPE, *Tipula juniperinus* Linn., by original designation.

No species in the adult has been described from the Orient. The following have been set up on the gall, larva or pupa. *O. indicus* (*nom. nov.* for *indianus* preocc.), *ischaemi*, *mangiferae*, *quadrilobatus*, *saligneus* and *tenuispatha*, all by Kieffer except

saligneus for which no author is quoted by that author in the Genera Insectorum.

Trichoperrisia, Kieff.

Rec. Ind. Mus. IX, 199 (1913).

Kieff., Gen. Ins., Fasc. 152, 65.

GENOTYPE, *T. pipericola* Kieff., by original designation.

pipericola Kieff., Rec. Ind. Mus. IX, 199, ♂ (1913).

Peradeniya. The *type* should be in the Indian Museum, but has apparently not been returned by the author.

Harpomyia, Felt.

Can. Ent. XLVIII, 401 (1916).

GENOTYPE, *H. indica* Felt, sp. nov., loc. cit.

indica Felt, Can. Ent. XLVIII, 401, ♂ ♀ (1916).

Pusa (presumably), 19-viii-15 [*Bahadur*]; reared from larvae found under the lining of a felt cap.

Dasyneura, Rond.

Sopra alcuni nuovi generi d'insetti Ditteri,

Parma, p. 17 (*Dasineura*) (1840).

Kieff., Gen. Ins., Fasc. 152, 86.

GENOTYPE, *D. obscura* Rond., by original designation.

gossypii Felt, Can. Ent. XLVIII, 29, ♀ (1916).

"*Type* Cecid. a2678." Sent by Mr. T. B. Fletcher, from S. India; ♂ ♀ figured (unnamed) in his "South Indian Insects."

Sect. ASPHONDYLARIAE.

Kieff., Gen. Ins., Fasc. 152, 87, tab. gen.

Schizomyia, Kieff.

Ent. Nach. XIV, 183 (1889).

Kieff., Gen. Ins., Fasc. 152, 88.

GENOTYPE, *S. galii* Kieff.

Only two species *S. indica*, Kieff. and *S. incerta*, Kieff. (*Marcellia* VII, 154, 1908) have been described from the Orient, but the imagos are not known.

Asphondylia, Loew.

Dipt. Beitr. IV, 21 (1850).

Sch., F. Austr. II, 395: Kieff., Gen. Ins., Fasc. 152, 91: *Auctt.*

GENOTYPE, *Cecidomyia sarothamni* Lw., according to Coquillett.

Phyllophaga, Rond., Dipt. Ital. Prod. I, 199 (1856).

GENOTYPE, *Cecidomyia fusca* Mg., by original designation.

Cylindrocera, Lioy, Atti. Istit. Veneto (3) IX, 503 (1863).

viticola Kieff. and Leeuwen, Marcellia VIII, 124, ♂ ♀ (1909).

Java. *Type* in Kieffer's collection.

sesami Felt, Can. Ent. XLVIII, 31, ♂ ♀ (1916).

South India, vi-12, reared from galls on the "gingelly" plant (*Sesamum indicum*). Adult (unnamed) and gall figured in Fletcher's "South Indian Insects," p. 364.

Daphnephila, Kieff.

Ann. Soc. Sci. Brux. XXIX, 144 (1905).

Kieff., Gen. Ins., Fasc. 152, 96.

GENOTYPE, *D. haasii* Kieff., by original designation.

haasi Kieff., Ann. Soc. Sci. Brux. XXIX, 144, ♂ ♀ (1905).

Bengal. *Types* in Kieffer's collection.

glandifex, *id.*, *loc. cit.*, 148, ♀ (1905).

Bengal. *Type* in Kieffer's collection.

linderae, *id.*, *loc. cit.*, 149, ♂ (1905).

Bengal. *Type* in Kieffer's collection.

Sect. BRACHYNEURARIAE.

Kieff., Gen. Ins., Fasc. 152, 101, tab. gen.

Hallomyia, Kieff.

Spol. Zeyl. VIII, 25 (1912).

Kieff., Gen. Ins., Fasc. 152, 106.

GENOTYPE, *H. iris* Kieff., by original designation.

iris Kieff., Spol. Zeyl. VIII, 25, ♀, figs. 1, 2 (1912).

Peradeniya, 17 and 19-v-10. Three *paratypes* in Indian Museum.

Sect. DIPLOSARIAE.

Eumerosema, Kieff.

Marcellia, XII, 42 (1913).

Kieff., Gen. Ins., Fasc. 152, 140.

GENOTYPE, *E. dispar* Kieff., by original designation.

dispar Kieff., Marcellia, XII, 42, ♂ ♀ (1913).

Formosa. *Type* in Entomological Museum, Berlin-Dahlem.

Chrysodiplosis, Kieff.

Trans. Linn. Soc. Lond. XIV, 318 (1911).

Kieff., Gen. Ins., Fasc. 152, 145.

GENOTYPE, *C. pulchricornis* Kieff., sp. nov., *loc. cit.*, by present designation.

squamatipes Kieff., Spol. Zeyl. VIII, 28, ♂ (1912).
Peradeniya, 20-v-10. Type in Indian Museum, unique.

Coelodiplosis, Kieff.

Marcellia XII, 43 (1913).

Kieff., Gen. Ins., Fasc. 152, 141.

GENOTYPE, *C. magnipennis* Kieff., by original designation.

magnipennis Kieff., Marcellia XII, 43, ♂ ♀ (1913).
Formosa. Type in the Entomological Museum, Berlin-Dahlem.

Courteia, Kieff.

Marcellia VIII, 131 (1909).

Kieff., Gen. Ins., Fasc. 152, 153.

GENOTYPE, *C. graminis* Kieff., by original designation.

graminis Kieff., Marcellia VIII, 131, ♀ (1909).
Java. Type in Kieffer's collection.

Feltodiplosis, Kieff.

Gen. Ins., Fasc. 152, 154 (1913).

Endaphis, Felt, Ent. News XXII, 128-224 (1911).

GENOTYPE, *F. abdominalis* Felt, by original designation.

hirta Felt, Ent. News XXII, 224 (1911).
Ceylon. Type in Kieffer's collection.

Arthrocnodax, Rübs.

Wien. Ent. Zeitg. XIV, 189 (1895).

Kieff., Gen. Ins., Fasc. 152, 155.

GENOTYPE, *A. vitis* Rübs., sp. nov.

rutherfordi Felt, Jour. N. Yk. Ent. Soc. XXIII, 180 (1915).
Ceylon.

walkeriana, *id.*, *loc. cit.*, 181 (1915).
Ceylon.

Plutodiplosis, Kieff.

Spol. Zeyl. VIII, 27 (1912).

Kieff., Gen. Ins., Fasc. 152, 166.

GENOTYPE, *P. magnifica* Kieff., by original designation.

magnifica Kieff., Spol. Zeyl. VIII, 27, ♀ (1912).
Peradeniya, 15-vii-10. The unique type accidentally destroyed.

Indodiplosis, Felt.

Can. Ent. XLVIII, 403 (1916).

GENOTYPE, *I. mangiferae*, *id.*, *loc. cit.*, sp. nov.**mangiferae** Felt, *loc. cit.*, 403, ♂ ♀ (1916).

♂ Bred from galls in mango leaf, Pusa, 13-iii-14, "Case No. 1023"; ♀ from same source, Pusa 26-ix-15 [T. Ram], "Case No. 100."

Streptodiplosis, Felt.

Can. Ent. XLVIII, 405 (1916).

GENOTYPE, *S. indica*, *id.*, *loc. cit.*, sp. nov.**indica** Felt, *loc. cit.*, 405 ♂ (1916).Kusti, North Wynaad, S. India, 16-ii [Bahadur], on leaves infested by the scale insect *Mytilaspis piperis*, Green.**Dentifibula**, Felt.

Bull. N. Yk. State Mus. CXXIV, 389 (1908).

Kieff., Gen. Ins., Fasc. 152, 174.

GENOTYPE, *Contarinia viburni* Felt, by original designation.**ceylanica** Felt, Jour. N. Yk. Ent. Soc. XXIII, 175 (1915).
Ceylon.**obtusilobae**, *id.*, *loc. cit.*, 176 (1915).
Ceylon.**Stictodiplosis**, Kieff.

Bull. Soc. Ent. Fran. LXIII, 28 (1894).

Kieff., Gen. Ins., Fasc. 152, 183.

GENOTYPE, *S. nubilipennis* Kieff., by original designation.**pulcherrima** Kieff., Rec. Ind. Mus. III, 130, fig. 2, wing (*Contarinia*) (1909).

Kurseong [Haas]. Type in Kieffer's collection.

Procontarinia, Kieff. & Cecconi.

Marcellia V, 135 (1906).

Kieff., Gen. Ins., Fasc. 152, 183.

GENOTYPE, *P. matteiana* Kieff. & Cecconi, by original designation.**matteiana** Kieff. & Cecconi, Marcellia, V, 135 ♂ ♀ (1906).
India. Type in Kieffer's collection.**Hypodiplosis**, Kieff.

Gen. Ins., Fasc. 152, 187 (1913).

Clinodiplosis partim., Kieff. & Leeuwen-Reijnvaan,
Marcellia VIII, 129 (1909).GENOTYPE, *Clinodiplosis paederiae* Kieff. & Leeuwen, by original designation.

paederiae Kieff & Leeuwen, Marcellia, VIII, 129, ♂ ♀ (1909).
Java. Type in Kieffer's collection.

Oribremia, Kieff.

Rec. Ind. Mus. IX, 199 (1913).

Kieff., Gen. Ins., Fasc. 152, 194.

GENOTYPE, *O. multifida* Kieff., by original designation.

multifida Kieff., Rec. Ind. Mus. IX, 199, ♂ (1913).
Bhowali, Kumaon, W. Himalayas, 5700 ft. [Imms]. Type apparently mislaid, not in Indian Museum.

Lestodiplosis, Kieff.

Bull. Soc. Ent. Fran. LXIII, 28 (1894).

Kieff., Gen. Ins., Fasc. 152, 195.

GENOTYPE, *L. alternans* Kieff., by his designation.

Coprodiplosis, Kieff., Feuille Jeune Natur. XXIV, 84 (1894).

GENOTYPE, *Diplosis polypori* Lw., by original designation.

ceylanicus Kieff., Spol. Zeyl. VIII, 28, ♀ (1912).
Peradeniya. Type in Indian Museum.

Diplecus, Kieff.

Bull. Soc. Hist. Nat. Metz. XXVIII, 55 (1913).

Kieff., Gen. Ins., Fasc. 152, 201.

GENOTYPE, *Coprodiplosis inconspicuus* Kieff., by original designation.

inconspicuus Kieff., Spol. Zeyl. VIII, 29, ♀ (*Coprodiplosis*) (1912).
Peradeniya, 11 and 21-vi-10. No definite type set up. Described from five specimens, said by Kieffer to have been taken in May, but the two in the Indian Museum forming part of the set of five are dated June. These become *paratypes*.

Diadiplosis, Felt.

Jour. N. Yk. Ent. Soc. XIX, 54 (1911).

GENOTYPE, *D. coccidivora*, *id.*, by original designation.

coccidivora Felt, Entom. XLVII, 86 (1914).
Ceylon.

smithii Felt, Jour. N. Yk. Ent. Soc. XXIII, 178 (1915).
Philippine Is.

Dyodiplosis, Rubs.

Zeits. wiss. Insektenbiol. VIII, 49 (1912).

Kieff., Gen. Ins., Fasc. 152, 205.

GENOTYPE, *D. arenariae* Rubs., by original designation.

generosi Felt, Jour. N. Yk. Ent. Soc. XXIII, 183 (1915).
Ceylon.

Pseudhormomyia, Kieff.

Synop. d. Cecid. d'Eur. et d'Alg. 57 (1898).

GENOTYPE, *P. granifex* Kieff., sp. nov., loc. cit.

Amaurosiphon, Rubs., Zeits. wiss. Insek., VII, 391 (1911).

GENOTYPE, *A. caricis* Rubs., sp. nov., by original designation.

fluviatilis Felt, Ent. News, XXVIII, 74, ♂ ♀ (1917).

Coimbatore, 29-vii-16. From galls on *Panicum fluitans* [Y. R. Rao]. "Type Cecid. a2777."

cornea Felt, loc. cit., 76, ♂ ♀ (1917).

Coimbatore, 2-viii-16 [Rao]. From galls on *Ischaemum ciliare*. "Type Cecid. a2778."

Lowiola, Kieff.

Miscell. Ent. V, 5 (1896).

Kieff., Gen. Ins., Fasc. 152, 206.

GENOTYPE, *L. centaureae*, according to Kieffer.

costata Felt, Jour. N. Yk. Ent. Soc. XXIII, 182 (1915).
Ceylon.

Orseoliella, Kieff.

Marcellia XI, 231 (1912).

Kieff., Gen. Ins., Fasc. 152, 152.

GENOTYPE, *O. javanica* Kieff., by original designation.

javanica Kieff., Marcellia VIII, 125, ♂ ♀ (*Orseolia*) (1909).

Java. Type in Kieffer's collection.

Cecidomyia, Mg.

Illig. Mag. II, 261 (1803).

Sch., F. Austr. II, 369 et Auct.: Kieff., Gen. Ins., Fasc. 152, 211.

GENOTYPE, *Tipula pini* DeGeer, according to Coquillett, and Kieffer.

Diplosis, Loew, Dipt. Beitr. IV, 20 (1850).

Cryptodiplosis, Kieff., Bull. Soc. Ent. Fran. LXIV, 194 (1895).

GENOTYPE, *Tipula pini*, DeGeer, by original designation.

deferenda Walk., Proc. Linn. Soc. Lond. I, 105, ♀ (1857).

Sarawak. Type apparently lost; not in British Museum.

Pachydiplosis, Kieff.

Bull. Soc. Hist. Nat. Metz, XXVIII, 108 (1913).

Kieff., Gen. Ins., Fasc. 152, 224.

GENOTYPE, *P. apricans* Kieff., by original designation.

- oryzae* Wood-Mason, Amer. Naturalist, p. 149 (*Cecidomyia*) (1881):
 Y. R. Rao, J. Asiat. Soc. Beng. n.s. XIII, p. clxxxiv (1917).
 Bengal; living in "paddy," causing the disease known as
 "silvershoots." Type untraceable, not in Indian Museum.
- graminicola* Kieff., Marcellia, VIII, 127, ♂ ♀ (*Clinodiplosis*) (1909).
 Java. Type in Kieffer's collection.
- ceylanica* Kieff., Spol. Zeyl. VIII, 26, ♀, fig. 3, tip of antenna
 (*Clinodiplosis*) (1912).
 Peradeniya, 13-v-10; 2, 9, and 25-vi-10; Pusa, 19-vi-11, at
 light [Howlett]. Paratypes in Indian Museum; no type erected.

Clinodiplosis, Kieff.

Feuille Jeunes Natur. Paris, XXIV, 121 (1894).
 Kieff., Gen. Ins., Fasc. 152, 237.
 GENOTYPE, *C. cilicrus* Kieff., by original designation.

No species in the adult stage is recorded from the Orient,
 but Kieffer has described the gall and the larva of three,
artemisiarum, *cellularis* and *nodifex*, in Marcellia VII, from
 the East Indies.

Androdiplosis, Felt.¹

- Jour. N. Yk. Ent. Soc. XXIII, 182 (1915).
 GENOTYPE, *F. coccidivora*, *id.*, *loc. cit.*, sp. nov.
- coccidivora* Felt, *loc. cit.*, 183 (1915).
 Ceylon.

Mycodiplosis, Rubs.

- Ent. Nachr. XXI, 186 (1895).
 GENOTYPE, *Diplosis coniophaga* Winn., by original designation.
- simulacri* Felt, Jour. N. Yk. Ent. Soc. XXIII, 177 (1915).
 Philippine Is.

Xiphodiplosis, Felt.¹

- Jour. N. Yk. Ent. Soc. XXIII, 180 (1915).
 GENOTYPE, *X. fulva*, *id.*, *loc. cit.*, sp. nov.
- fulva* Felt, *loc. cit.*, 180 (1915).
 Ceylon.

Sect. PORRICONDYLARIAE.

Kieff., Gen. Ins., Fasc. 152, 258, tab. gen.

Dicroneurus, Kieff.

Ent. Nachr. XXI, 122 (1895).
 Kieff., Gen. Ins., Fasc. 152, 270.

¹ The work in which these genera are described is inaccessible in India, and it is therefore impossible to place them in their exact systematic position.

GENOTYPE, *Epidosis lineatus* Kieff., by original designation.

indicus Kieff., Rec. Ind. Mus. IX, 200, ♂ (*Epidosis*) (1913).

Kalighat, 6000 ft. Type, apparently mislaid, not in Indian Museum.

Misocosmus, Kieff.

Bull. Soc. Hist. Nat. Metz, XXVIII, 55 (1913).

Kieff., Gen. Ins., Fasc. 152, 276.

GENOTYPE, *Epidosis ceylanicus* Kieff., by original designation.

ceylanicus Kieff., Spol. Zeyl. VIII, 29, ♀ (*Epidosis*) (1912).

Peradeniya, 19-v-10. Type in Indian Museum.

Colpodia, Winn.

Linn. Ent. VIII, 188 (1853).

Kieff., Gen. Ins., Fasc. 152, 268.

GENOTYPE, *C. angustipennis* Winn., sp. nov., loc. cit.

fletcheri Felt, Can. Ent. XLVIII, 400, ♂ (1916).

Pusa, a unique ♂, 1-i-16. [Bahadur]. "Type Cecid. 1696." No location of type is stated.

Epidosis, Lw.

Dipt. Beit. IV, 21 (1850).

GENOTYPE, *E. fragilis* Lw., sp. nov., loc. cit., p. 38, the 1st species by present designation.

indicus Kieff., Rec. Ind. Mus., IX, 200 (1913).

India.

Subfamily **LESTREMINAE**.¹

Peromyia, Kieff.

Bull. Soc. Ent. Fran. 63 (1894).

Kieff., Gen. Ins., Fasc. 152, 292.

GENOTYPE, *P. leveillei* Kieff., by original designation.

bengalensis Kieff., Ann. Soc. Sci. Brux. XXIX, 158, ♂, fig. 4, wing (1905).

Bengal. Type in Kieffer's collection.

Lestremia, Macq.

Recueil Soc. Agric. Lille, 173 (1826).

Sch., F. Austr. II, 413 et Auctt.

Kieff., Gen. Ins., Fasc. 152, 307.

GENOTYPE, *Lestremia cinerea* Mcq., according to Coquillett.

Cecidogona, Lw., Stett. Ent. Zeit. V, 324 (1844).

GENOTYPE, *Lestremia carnea* Lw., by original designation.

¹ For table of genera, see Kieff., Gen. Ins., Fasc. 152, 305.

- Furcinerva*, Rond., Nuovi Ann. Soc. Nat. Bologna (2) VI, 369 (1846).¹
Mimosciara, *id.*, Mem. Ia. ditt. Ital. 25 (1840).
Molobraea, *id.*, Atti. Soc. Ital. Sci. Nat. Milano, II, 287 (1860).
Yposataea, *id.*, Dipt. Ital. Prod. I, 198 (1856).
indica Kieff., Rec. Ind. Mus. III, 129, fig. 1, wing (1909).
 Calcutta [*Annandale*]; Kurseong, 26-iii-10 [*Gravely*].
ceylanica Kieff., Spol. Zeyl. VIII, 29, ♀ (1912).
 Peradeniya, 13-v-10, at light. *Type* in Indian Museum, unique.

Family MYCETOPHILIDAE.

Subfamily SCIARINAE.

Sciara, Mg.

Illig. Mag. II, 263 (1803).

Sch., Fauna Austr. II, 417: Auctt. var.

Brun., Fauna Brit. Ind., Dipt., 120, figs. (p. 121), pl. ii, 15-17, wings.

GENOTYPE, *S. thomae* L.²*Molobrus*, Latr., Hist. Nat. Crust. Ins. XIV, 288 (1805).GENOTYPE, *Tipula thomae* L., as the "Tipule de Thomas."*Planctes*, Walk., Ent. Month. Mag. III, 178 (1836).GENOTYPE, *P. extremus*, *id.*, *loc. cit.*, sp. nov.*Planatella*, Westw., Synop. gen. Brit. Ins., 126 (1840).*indica* Walk.,³ Ins. Saunds. Dipt. V, 419 (1856).

Brun., Fauna Brit. Ind. I, 130, ♀.

East India; Darjiling, 5-viii-09 [*Paiva*]; Kurseong, 13-16-vii-07; common, 14-viii-09 [*Paiva*]; Siliguri, 18-20-vii-07; Phagu, 21-v-16; Bhim Tal, 19-22-ix-06 [*Annandale*]; Senchal, v-13. *Type* in British Museum in good condition.*pusilla* Dol., Nat. Tijd. Ned. Ind. X, 407, pl. iv, 4 (*Scatopse*) (1856).Java. *Type*, location unknown, apparently lost.*laticornis* Walk., Proc. Linn. Soc. Lond. I, 105, ♂ ♀ (1857).Sarawak. *Type* in British Museum, damaged but recognizable.

¹ *Furcinerva* was founded on the union of *Zygoneura* Mg., *Lestremia* Macq., *Mimosciara* Rond. and *Catocha* Hall. In his Prodrôme vol. I, Rondani admits *Catocha* as valid, ignores *Furcinerva* and erects *Yposataea* for the three other genera without any type species. Coquillett proposed *Mimosciara molobrina* Rond., which was the genotype of this latter genus.

² *S. thomae* L., the common European species, was recorded from Sumatra by Van der Wulp (Dipt. Sum. Exped. 7, and Cat. Dipt. S. Asia, 1), but in the latter work he says that Rübsaamen (Berl. Ent. Zeits. XXXIX, 32) doubts the identity of the species. I therefore refrain from including it in the present catalogue.

³ *S. indica* Walk. is a good and conspicuous species, but his others are mostly unrecognizable from the descriptions, except perhaps *filipes*.

- solita* Walk., *loc. cit.*, I, 105, ♀ (1857).
Sarawak. Type in British Museum, in fair condition.
- femoralis* Dol., Nat. Tijd. Ned. Ind. XVII, 78 (1858).
Amboina. Type, location unknown, apparently lost.
- selecta* Walk., Proc. Linn. Soc. Lond. III, 77, ♂ (1859).
Aroe. Type in British Museum, in good condition.
- longipes*, *id.*, *loc. cit.*, V, 263 (1861).
Tondano. Type in British Museum, in fair condition (abdomen missing).
- ponderosa*, *id.*, *loc. cit.*, VII, 230, ♀ (1864).
North Ceram; Mimika River, Dutch Papua, viii-1910. Type in British Museum, in good condition.
- varipes*, *id.*, *loc. cit.*, VIII, 102, ♀ (1865).
Papua. Type in British Museum (no abdomen).
- filipes*, *id.*, *loc. cit.*, VIII, 102 (1865).
Papua. Type in British Museum, in fair condition but without antennae.
- Both Walker and Van der Wulp state that this species represents a new genus, but the type (and only) specimen is too damaged to diagnose its characters.
- rufithorax* Wulp, Mid. Sum. Exped., Dipt. 6, pl. i, 1, full insect (1881): Rübs., Berl. Ent. Zeits. XXXIX, 39: Meij., Tijd. v Ent. L. 200, pl. v, 1: Brun., Fauna Brit. Ind., Dipt. I, 128, ♂ ♀, pl. iii, 15, ♂ genitalia.
Probably widely distributed in the East. Sumatra, Java (var. locs.), Ceylon, India (many localities, *vide* Brun., Fauna), Assam, Burma, and probably South China and other parts. Recorded during the greater part of the year. Occurs as far north as Simla, x-08 [Howlett]. Type in Leyden Museum, *cotype* in Amsterdam Museum.
- sulcata* Wulp, Tijd. v. Ent. XXX, 177, pl. xi, 12 (1887).
Java. Type probably lost, originally in Van der Wulp's collection.
- pruinosa* Rübs., Berl. Ent. Zeits. XXXIX, 33, pl. i, 2 (1894).
Borneo. Type in Berlin Museum.
- rotunda*, *id.*, *loc. cit.*, 34 (1894).
Himalayas. Type in Berlin Museum.
- singhalensis*, *id.*, *loc. cit.*, 35 (1894).
Ceylon. Type in Berlin Museum.
- trilineata* Brun., Fauna Brit. Ind., 126, ♀ (1912).
Darjiling, 8-viii-09, *type*; 6-viii-09 [both Paiva]. Type in Indian Museum.
- trifasciata*, *id.*, *loc. cit.*, 126, ♀ (1912).
Darjiling, 10-16-x-05 [Brunetti]. Type, unique, in Brunetti collection.
- opposita*, *id.*, *loc. cit.*, 127, ♂ (1912).
Kurseong, 3-vii-08. Type, unique, in Indian Museum.

- distinguenda*, *id.*, *loc. cit.*, 127, ♂ (1912).
Darjiling (7000 ft.), 12-viii-09 [Jenkins]. Type, unique, in Indian Museum.
- rufoabdominalis*, *id.*, *loc. cit.*, 129, ♀ (1912).
Ukhrul, Manipur (6400 ft.) [Pettigrew]. Type, unique, in Indian Museum.
- luteiventris*, *id.*, *loc. cit.*, 129, ♀ (1912).
Phagu, 12-v-09 [Annandale]. Type, unique, in Indian Museum.
- flammiventris*, *id.*, *loc. cit.*, 129, ♀ (1912).
Darjiling, 6-viii-09 [Paiva]. Type, unique, in Indian Museum.
- flavofemorata*, *id.*, *loc. cit.*, 130, ♂ (1912).
Kurseong, 8-vii-08; Simla, x-08 [Howlett]. Type in Indian Museum.
- nigripennis*, *id.*, *loc. cit.*, 131, ♂ ♀ (1912).
Darjiling, 26-v-10, type ♂; 29-ix-08; 28-v-10 [all Brunetti]; Kurseong, 13-viii-09 [Paiva]; Bhim Tal, 19-22-ix-06 [Annandale]; Simla, x-08 [Howlett]; Pashok, Sikkim (2000 ft.), 5-ix-09; Ukhrul, Manipur; Shillong; Pattipola, Ceylon. Types in Indian Museum.
Some ♀ ♀ taken by Mr. C. W. Beebe at Kuching, Sarawak, 23-vi-10 and 4-vii-10 are probably a small variety of this species.
- diversipes*, *id.*, *loc. cit.*, 131, ♂ ♀ (1912).
Darjiling, 10-viii-09 [Paiva]; Ghoom (above Darjiling, 7500 ft.), 18-ix-08; Kurseong, 13-16-vii-07; Siliguri, 18-25-vii-07. Type ♂ ♀ in Indian Museum.
- fratercula*, *id.*, *loc. cit.*, 132, ♀ (1912).
Kurseong, 7-ix-09 [Annandale]; Bhim Tal, 22-27-ix-06 [Annandale]. Type in Indian Museum.
- exacta*, *id.*, *loc. cit.*, 132, ♀ (1912).
Darjiling, 4-7-viii-09, type [Paiva]; Kurseong, 7-ix-09; 4-8-vii-08; Ghoom (above Darjiling), 19-ix-08; Siliguri, 18-20-vii-07; Bhowali, Kumaon, vii-09 [Imms]; Ukhrul, Manipur; all in the Indian Museum. In my collection from Darjiling, 10-16-x-05 and Shanghai, 9-v-06 [both Brunetti], and from Ohiya, Ceylon [Green]. Type in Indian Museum.
- longinervis*, *id.*, *loc. cit.*, 133, ♀ (1912).
Siliguri, 18-20-vii-07, type; Kurseong, x-1910 [D'Abreu]; in the Pusa collection from Mussoorie, viii-1906. Type in Indian Museum.
- flavicollis*, *id.*, *loc. cit.*, 134, ♂ ♀ (1912).
Siliguri, 18-20-vii-07, type ♂; Darjiling, 1-x-08, type ♀ [Brunetti]; 26-29-vi-10; Kurseong, 29-vi-10 [both Annandale]. Types in Indian Museum.
- orientalis*, *id.*, *loc. cit.*, 135, ♂ ♀ (1912).

Darjiling; Kurseong; Siliguri; Bhim Tal; Mandalay; Ukhul; Dibrugarh, 17-19-xi-11; Rotung, N.E. Ind. Frontier, 24-25-xii-11 [both *Kemp*]; Calcutta (common Jan. to May and July to Sept.); Lucknow; Bijnor District; Ganges Delta; Travancore State (var. locs.); Ceylon; Dawna Hills; Lower Burma. Occurs, speaking generally, nearly all the year round in many localities. *Types* in Indian Museum, *cotypes* in my collection.

fascipennis, *id.*, *loc. cit.*, 136, ♀ (1912).

Darjiling, xi-1910 [*D'Abreu*]. *Type*, unique, in Indian Museum.

sexsetosa, *id.*, *loc. cit.*, 137 ♀ (1912).

Darjiling, 20-ix-08, *type* [*Brunetti*]; 6-viii-09 [*Paiva*]; Pashok, Sikkim (2000 ft.), 5-ix-09. *Type* in Indian Museum.

quadrisetosa, *id.*, *loc. cit.*, 137, ♂ (1912).

Darjiling, 6-viii-09 [*Paiva*]. *Type* in Indian Museum.

pallescens, *id.*, *loc. cit.*, 137 (sex uncertain) (1912).

Khayon (or "Farm") caves, 9 miles from Moulmein, Tenasserim, 7-iii-08, taken in the darkest part of the cavern by Dr. Annandale, in company with *Crysithyris speloca*, the only known moth adapted for a cavernicolous existence. *Type*, unique, in Indian Museum.

fulvescens, *id.*, *loc. cit.*, 138, ♀ (1912).

Rajshai, East Bengal, 1-6-ii-07 [*Annandale*]. *Type*, unique, in Indian Museum, possibly identical with *pallescens*.

setilineata, *id.*, *loc. cit.*, 138, ♂ ♀ (1912).

Simla, 10-v-09, *type* ♂ [*Annandale*]; Darjiling, 6-viii-09, *type* ♀ [*Paiva*]. *Types* in Indian Museum.

latelineata, *id.*, *loc. cit.*, 139, ♂ (1912).

Dawna Hills, Tenasserim (on western slopes, 2000-3000 ft.), 2-3-iii-08 [*Annandale*]. *Type*, unique, in Indian Museum.

radicum, *id.*, *loc. cit.*, 139, ♂ ♀ (1912).

Calcutta, 21-i-10 to 24-ii-10 [*Annandale*]. *Types* in Indian Museum; *cotypes* in my collection.

The larva lives on lily bulbs and kindred plants.

impostor, *id.*, *loc. cit.*, 140, ♂ (1912).

Darjiling, 1-x-08, *type*, 2-x-08, 10-16-x-05 [all *Brunetti*]. *Type* in Indian Museum, *cotype* in my collection.

segmenticornis, *id.*, *loc. cit.*, 141, ♂ (1912).

Bindukhera (Naini Tal Distr., plains), 13-iv-09. *Type* in Indian Museum, unique.

compacta, *id.*, *loc. cit.*, 141, ♀ (1912).

Gangtok, Sikkim (6150 ft.), 8-ix-09, *type*; Butal, Nepal, 12-ii-08. *Type* in Indian Museum.

inconspicua, *id.*, *loc. cit.*, 142, ♀ (1912).

Sylhet, 14-i-05 [*Hall*]. *Type*, unique, in Indian Museum.

hirtilineata, *id.*, *loc. cit.*, 142, ♂ ♀ (1912).

- Phagu, 12-v-09, *type* ♂, 12-v-09; Simla, 10-v-09; Darjiling, 6-viii-09; Kurseong, 4-vii-08 [Annandale and others]. *Types* in Indian Museum.
- niveiapicalis*, *id.*, *loc. cit.*, 143, ♀ (1912).
Ukhrul, Manipur (6,400 ft.) [Pettigrew]. *Type* in Indian Museum.
- longipennis*, *id.*, *loc. cit.*, 143, ♂ ♀ (1912).
Below Simla, 6-v-10, *type* ♂ [Annandale]; Darjiling, 25-v-10, *type* ♀ [Brunetti]; Kurseong, 4-vii-08; 26—27-vi-10 [Annandale]; Tonglu, Darjiling (10,000 ft.), 22-iv-10 [Beebe]. *Type* in Indian Museum.
- flaviseta*, *id.*, *loc. cit.*, 144, ♀ (1912).
Simla, 10-v-09 [Annandale]. *Type*, unique, in Indian Museum.
- nitidithorax*, *id.*, *loc. cit.*, 145, ♂ (1912).
Kalighat (Garhwal Distr., 6000 ft., India), 4-vi-10 [Imms]. *Type*, unique, in Indian Museum.
- longitudinalis*, *id.*, *loc. cit.*, 145, ♂ ♀ (1912).
Kurseong, 8-vii-08, *type* ♂ [Annandale]; Darjiling, 28-v-10, *type* ♀ [Brunetti]; Tonglu, Darjiling, 10,000 ft., 22-iv-10 [Beebe]. *Types* in Indian Museum.
- ruficoxa*, *id.*, *loc. cit.*, 146, ♀ (1912).
Kurseong, 25-vi-10 [Annandale]. *Type*, unique, in Indian Museum.
- flavipleura*, *id.*, *loc. cit.*, 146, ♂ ♀ (1912).
Darjiling, 2-x-08, *type* ♂ [Brunetti]; 6-viii-09, *type* ♀ [Paiva]; Kurseong, 3-ix-09 [Annandale]. *Types* in Indian Museum.
- evanescens*, *id.*, *loc. cit.*, 147, ♂ ♀ (1912).
Simla, 9-v-09, *type* ♂; Calcutta, 25-ii-09 [both Annandale]; Darjiling, 6-viii-09 [Paiva]; Peradeniya, 14-vii-10, 7-viii-10. *Types* in Indian Museum.
- parallela*, *id.*, *loc. cit.*, 147, ♂ ♀ (1912).
Simla, 9-v-09, *type* ♂ [Annandale]; Darjiling, 2-x-08, *type* ♀, and 26-v-10 [Brunetti]. *Types* in Indian Museum.
- turrita* de Meij., Bijdr. tot Dierk, XIX, 45, ♀, pl. i, 1, head, 2, wing, (1913).
Ceram, ii. *Type*, unique, in (?) Amsterdam Museum.
- heteroptera* de Meij., Tijds. v. Ent. LVI, 318, ♂ ♀; pl. xv, 1, wing; 1a, genitalia ♂ (1913).
Gedeh crater, Java [Roepke]. *Type* in Amsterdam Museum.
- fuscolimbata* Ender., Arch. Naturg. LXXVII, Supp. 3, 150 [*Lycoria*] (1912).
Sumatra.
- sumatrana*, *id.*, *loc. cit.*, 151 [*Lycoria*] (1912).
- bispinosa* Petzey, Phil. J. Sci. XIII, D, 215, ♂, pl. i, 2, ♂ clasper, 4, wing (1918).

Los Banos, Luzon, 3-iii-17. Unique *type* in Cornell University collection.

uichancoi, *id.*, *loc. cit.*, 215, ♂ ♀, pl. i, 1, ♂ clasper, 3, wing (1918).

Mt. Maquiling, Luzon, 28-viii-17. *Types* in Cornell University collection.

Phorodonta, Coq.

Proc. U.S. Mus. XXXVIII, 589, nom. nov. for *Odontonyx*, Rübs., preocc. (1910).

Odontonyx, Rübs., Berl. Ent. Zeits. XXXIX, 19 and 25 (1894).

GENOTYPE, *O. fruhstorferi* Rübs., by original designation.

Odontonyx preoccupied by Stephens in Coleoptera (1828), and used again by Guerin in 1843.

fruhstorferi Rübs., Berl. Ent. Zeits. XXXIX, 25, fig. 3, pl. iii, 13, 23 (1894).

Java. *Type* in Berlin Museum.

Psilomegalosphys, Enderl.

Archiv f. Naturges. LXXVII, 137 (1911).

GENOTYPE, *P. xanthogaster* Enderl., by original designation.

xanthogaster Enderl., *loc. cit.*, 137, ♀, pl. i, 6, wing (1911).

Sumatra. *Type* in Stettin Zoolog. Museum.

Trichomegalosphys, Enderl.

Archiv f. Naturges. LXXVII, 136 (1911).

GENOTYPE, *T. funesta* Enderl., by original designation.

funesta Enderl., *loc. cit.*, 136, ♀ (1911).

Soekaranda (Sumatra). *Type* in Stettin Zoolog. Museum.

Scythropochroa, Enderl.

Archiv f. Naturges. LXXVII, 138 (1911).

GENOTYPE, *S. latefurcata* Enderl., by original designation.

velata Enderl., *loc. cit.*, 139, ♀ (1911).

Soekaranda (Sumatra). *Type*, unique, in Stettin Zoological Museum.

Zygoneura, Mg.

Syst. Besch. VI, 304 (1830).

GENOTYPE, *Z. sciarina* Mg.

remota deMeij., Nova Guinea Res. IX, 305, ♂ (1913).

Riverkamp, ii. *Type* in Amsterdam Museum.

Subfamily *MACROCERINAE*.*Macrocera*, Mg.

Illig Mag. II, 261 (1803).

Sch., F. Austr. II, 430 : *Auctt. var.* : Johan., Gen. Ins., Fasc. 93, 28 : Brun., Fauna Brit. Ind., 49.GENOTYPE, *M. lutea* Mg., by designation of Curtis, Brit. Entom., 637 (1837).*Tipula longicornis* Mg., has been proposed as the type-species, but the identification seems doubtful.*Geneja*, Lioy, Atti Istit. Ven. (3) IX, 229 (1863).¹GENOTYPE, *Macrocera fasciata* Mg., by Coquillett's designation.? *Macrourea*, Berendt, Organ. Reste Bernstein I, 51, nom. nud. (1845).*ornata* Brun., Fauna Brit. Ind., 51, ♂ ♀, pl. i, 1, wing; pl. iii, ♂ genitalia (1912).Darjiling, 28-v-10, type ♂ [*Brunetti*]; Kurseong, 15-viii-09, type ♀ [*Jenkins*]. Types in Indian Museum.*alternata*, *id.*, *loc. cit.*, 52, ♂ ♀, pl. i, 2, wing (1912): *Id.*, Rec. Ind. Mus. XIII, 61.Naini Tal, type ♂, 10-vi-09; Mussoorie, vi-09, type ♀ [*Brunetti*]; Simla, x-08 [*Howlett*]. Type ♂ in Indian Museum, type ♀ in my collection.*flavicosta*, *id.*, *loc. cit.*, 53, ♀, pl. i, 3, wing (1912).

Exact locality uncertain, probably India, but certainly the Orient. Type in my collection.

brunnea, *id.*, *loc. cit.*, 53, ♀, pl. i, 5, wing (1912).Phagu, 12-v-09 [*Annandale*]. Type in Indian Museum.*elegans*, *id.*, *loc. cit.*, 54, ♂, pl. i, 4, wing (1912).Darjiling, 26-v-10 [*Brunetti*]. Type in Indian Museum, unique.*inconspicua*, *id.*, *loc. cit.*, 54, ♂ (1912).Naini Tal Distr., 28-v-09; Simla, x-08 [*Howlett*]. Type in Indian Museum, unique.*ferruginea*, *id.*, *loc. cit.*, 55, ♀ (1912).Bhim Tal, 22—27-ix-06 [*Annandale*]. Type in Indian Museum, unique.*fryeri* Edw., Ann. Mag. Nat. Hist. (8) XII, 200, ♂ (1913).

Peradeniya, ix-1911.

Subfamily *CEROPLATINAE*.*Ceroplatus*, Bosc.

Act. Soc. Hist. Nat. Paris I, i, 42 (1792).

Sch., F. Austr. II, 433 : *Auctt. var.* : Johan., Gen.

Ins., Fasc. 93, 18 : Brun., Fauna Brit. Ind., 56.

¹ A *nom. nov.* for *Macrocera* Mg., under the plea of the latter being preoccupied by Spinola in Hymenoptera. I cannot trace any such preoccupation.

GENOTYPE, *Tipula lineata* F., designated by Rondani as *Platyura laticornis* Mg.

quadripunctatus Brun., Fauna Brit. Ind. 57, ♀, pl. i, 6, wing; pl. iii, 2, abdomen (1912).

Calcutta, 27-vii-09 [Annandale]. Type, unique, in Ind. Mus.

If *Cerotelion* Rond., Dipt. Ital. Prod. I, 191 (1856), is admitted as generically distinct from *Ceroplatus*, this species will come in it.

Rhynchoplatyura, de Meij.

Tijd. v. Ent. LVIII, Supp. 3 (1915).

GENOTYPE, *R. longirostris*, *id.*, sp. nov., by original designation.

longirostris de Meij., *loc. cit.*, 3 (1915).

Simalur Is., Sumatra.

Platyura, Mg.

Illig. Mag. II, 264 (no species) (1803).

Mg., Klass, 101 (1804).

Sch., F. Aust. II, 435: *Auctt.*: Johan., Gen. Ins., Fasc. 93, 20: Brun., Fauna Brit. Ind., 58.

GENOTYPE, *Platyura fasciata* Mg., by Zetterstedt's designation.

Orfelia, A. Costa, Il Giambatt. Vico, II, 448 (1857).

GENOTYPE, unknown to present author.

Isoneuromyia, Brun., Fauna Brit. Ind., 66 (1912).¹

GENOTYPE, *I. annandalei*, *id.*, sp. nov., *loc. cit.*

venusta Walk., Ins. Saund. Dipt. pt. V, 421 (1856).

Brun., Fauna Brit. Ind., 63, ♀.

Bareilly, 15—22-iii-07; Pusa, 16-iv-07, iv-08, v-07; Chapra, Bengal; Chittagong, 14-vii-08 [Hall]. Type, location unknown.

grandis Brun., Fauna Brit. Ind., 60, ♂ ♀, pl. i, 7, wing (1912).

Mussoorie, 22-vi-05, type ♂ ♀, in cop., 26-vi-05 [both Brunetti].

Types in my collection.

suffusinervis, *id.*, *loc. cit.*, 60, ♂ ♀, pl. i, 9, wing (1912).

Sylhet, 10-iv-05, type ♂ ♀, in cop. [Hall]. Types, unique, in Indian Museum.

apicipennis, *id.*, *loc. cit.*, 61, ♀, pl. i, 8, wing, pl. iii, 4, head (1912).

Naini Tal, 2-vi-09. Type, unique, in Indian Museum.

affinis, *id.*, *loc. cit.*, 62, ♀ (1912).

Darjiling, 25-v-10 [Brunetti]. Type, unique, in Ind. Mus.

flaviventris, *id.*, *loc. cit.*, 62, ♀ (1912).

Darjiling, 23-ix-08 [Brunetti]. Type, unique, in Ind. Mus.

vicina, *id.*, *loc. cit.*, 63, ♀ (1912).

Naini Tal, 3-vi-09. Type in Indian Museum.

¹ I have accepted Edwards' opinion (in litt.) as to the synonymy: Rec. Ind. Mus. XIII, 61.

- octosegmentata*, *id.*, *loc. cit.*, 64, ♂ (1912).
Allahabad, 9-ix-09 [Imms]. *Type*, unique, in Ind. Mus.
- ruficornis*, *id.*, *loc. cit.*, 64, ♀ (1912).
Sylhet, 2-ii-45, 30-ii-04; Naini Tal, 10-vi-09. *Type* in Indian Museum.
- indistincta*, *id.*, *loc. cit.*, 65, ♂ (1912).
Kurseong, 20-vi-10 [Annandale] *Type*, unique, in Ind. Mus.
- longifurcata*, *id.*, *loc. cit.*, 65, ♂ ♀ (1912).
Kurseong, 24—26-iii-10 [Annandale]. *Types* in Ind. Mus.
- funerea*, *id.*, *loc. cit.*, 65, ♀ (1912).
Gangtok, 6150 ft., Sikkim, 9-ix-09. *Type*, unique, in Indian Museum.
- flavomarginata*, *id.*, *loc. cit.*, 66, ♂ (1912).
Naini Tal, 3-vi-09. *Type*, unique, in Indian Museum.
- fumipes*, *id.*, *loc. cit.*, 66, ♂ (1912).
Peradeniya, 22-vii-10. *Type*, unique, in Indian Museum.
- annandalei* Brun., *loc. cit.*, 67, ♀, pl. iii, 5, full insect (*Isoneuromyia*) (1912).
Sylhet, 27-v-05, *type* [Hall]; Kandy, v and viii-09 [Green];
Cochin State, 28—29-xi-14 [Gravelly]. *Type* in Indian Museum,
cotypes in my collection.
- rufescens*, *id.*, *loc. cit.*, app. 559, ♀ (*Isoneuromyia*) (1912).
Simla, 20-vii-11 [Annandale]. *Type*, unique, in Indian Museum.
- limbata*, *nom. nov.*, Brun., Rec. Ind. Mus. XIII, 61 (1917).
P. marginata, Brun., *loc. cit.*, 62, ♂ ♀, pl. iii, 3, genitalia ♂ (1912), *preocc.* Mg.
Naini Tal, 3 and 10-vi-09. *Types* in Indian Museum.

Subfamily SCIOPHILINAE.¹*Sciophila*, Mg.

Syst. Besch. I, 245 (1818).

Sch., F. Austr. II, 441: *Auctt. var.*: Johan., Gen.

Ins., Fasc. 93, 36: Brun., Fauna Brit. Ind., 68.

GENOTYPE, *S. hirta* Mg., designated by Curtis,

Brit. Ent. 641 (1837).

Lasiosoma, Winn., Verh. z.b. Ges. Wien XIII, 748 (1863).GENOTYPE, *Sciophila pilosa* Mg., by Coquillett's designation.*bicolor* Brun., Fauna Brit. Ind., 70, ♀, pl. i, 10, wing (1912).Darjiling, 7—8-viii-09, *type* [Paiva]; 28-v-10 [Brunetti].
Type in Indian Museum.¹ For table of genera see Kieff., Gen. Ins., Fasc. 93, 31.

Mycomyia, Rond.

Dipt. Ital. Prod. I, 194 (*Mycomyia*) (1856).

Johan., Gen. Ins., Fasc. 93, 45: Brun., Fauna Brit. Ind., 70.

GENOTYPE, *M. marginata* Mg., designated by Rondani, Dipt. Ital. Prod. I.

Sciophila, Mg., pt.

Sciophila, Winn., Verh. z.b. Ges. Wien XIII, 707 (1863).

GENOTYPE, apparently no species ever selected.

? *Sciobia*, Lw., Bernst. u. Bernst. Fauna, 33 (1850).

? *Cnephocophila*, Phil., Verh. z.b. Ges. Wien XV, 618 (1869).

flaviventris Brun., Fauna Brit. Ind., 72, ♂ ♀ (1912).

Kurseong, 3-vii-08, type ♂ [*Annandale*]; Ohiya, Ceylon, ii-07, type ♀ [*Green*]. Type ♂ in Indian Museum, type ♀ in my collection.

bifascipennis, *id.*, *loc. cit.*, 72, ♂, pl. iii, 6, ♂ genitalia (1912).

Brun., Rec. Ind. Mus. XIII, 61, note.

Bhowali, 5700 ft., vii-09, type [*Imms*]; Simla, x-08 [*Howlett*]; Maskeliya, Ceylon [*Green*]. Type in Indian Museum, cotype in my collection.

basalis Brun., Fauna Brit. Ind., 73, ♂, pl. i, 13, wing (1912).

Bhowali, 5700 ft., vii-09 [*Imms*]. Type in Indian Museum.

The abdomen of the unique type has since been lost.

tinctipennis, *id.*, *loc. cit.*, 74, pl. i, 11, wing (1912).

Kurseong, 3-vii-08 [*Annandale*]. Type, unique, sex uncertain, in Indian Museum.

ferruginea, *id.*, *loc. cit.*, 74, ♂ ♀, pl. i, 12, wing (1912).

Kurseong, 3-vii-08, type ♂ [*Annandale*]; Darjiling, 29-v-10, type ♀ [*Brunetti*]; Calcutta, 11-ii-90; Peradeniya, xii [*Gravelly*]. Types in Indian Museum.

trilineata, *id.*, *loc. cit.*, 75, ♂ (1912).

Simla, 10-v-09 [*Annandale*]; x-11, sweeping near the Dhobi Ghat [*Howlett*]. Type in Indian Museum.

indefinita, *id.*, *loc. cit.*, 76, ♂, and ? ♀ (1912).

Brun., Rec. Ind. Mus., XIII, 62.

Darjiling, 29-v-10 [*Brunetti*]; Simla, x-11. Type and second specimen in Indian Museum.

indica, *id.*, *loc. cit.*, 76, ♂ ♀ (1912).

Simla and Phagu, 10-12-v-09 [*Annandale*]. Type in Indian Museum.

flavithorax, *id.*, *loc. cit.*, 77, ♂ (1912).

Darjiling, 25-v-10 [*Annandale*]. Type, unique, in Ind. Mus.

curvilinea, *id.*, *loc. cit.*, 77, ♂ (1912).

Tonglu, Darjiling Distr., 10,000 ft., 22-iv-10, type [*Beebe*]; Darjiling, 29-viii-10 [*Brunetti*]. Type in Indian Museum.

Polylepta, Winn.

- Verh. z.b. Ges. Wien XIII, 745 (1863).
 Sch., F. Austr. II, 447: *Auctt. var.*: Johan., Gen.
 Ins., Fasc. 93, 43: Brun., Fauna Brit. Ind., 78,
 pl. i, 14.
 GENOTYPE, *P. undulata* Winn., by designation of
 Johannsen (Gen. Ins., Fasc. 93, 43, 1909).
dublosa Brun., Fauna Brit. Ind., 79, ♂ (1912).
 Mundali, Dehra Dun, 9000 ft., 10-v-10 [Beebe]. *Type*,
 unique, in Indian Museum.
incerta, *id.*, *loc. cit.*, 79 (1912).
 Mundali, 12-v-10 [Beebe]. *Type*, a unique specimen, appar-
 ently a ♀, in Indian Museum.

Neoempheria, Ost. Sack.¹

- Cat. N. Amer. Dipt., Ed. II, 9 and 16, *nom. nov.* (1878).
Empheria, Winn., Verh. z.b. Ges. Wien XIII, 738
 (1863), preocc. Hagen 1856 in Psocidae; Sch., F.
 Austr. II, 446.
 GENOTYPE, *Sciophila striata* Mg., by Coquillett's
 designation.
 * *Mycomyia*, Johann., pt., Gen. Ins., Fasc. 93, 45 (1909).
 GENOTYPE, *Sciophila striata* Mg., according to
 Coquillett.
tropcia Dol., Nat. Tijd. Ned. Ind. XIV, 392, pl. iv, 2 (*Sciophila*)
 (1857).
 Java. *Type* possibly lost; location unknown to present
 author. Van der Wulp removes the species from *Sciophila* to
 this genus on the strength of Doleschall's figure.
indivisa Walk., Proc. Linn. Soc. Lond. VII, 233 (*Leia*) (1864).
 Waigou. *Type* in good condition in British Museum.
 Mr. Edwards informs me the species belongs here.
propinqua de Meij., Tijd. v. Ent., L, 203, ♂ ♀, pl. v, 4, wing, pt.
 (*Empheria*) (1907).
 Semarang, iii [Jacobson]. *Types* in Amsterdam Museum,
 including pairs *in cop.*
apicalis, *id.*, *loc. cit.*, 204, ♂, pl. v, 5, wing, pt. (*Empheria*) (1907).
 Semarang, i [Jacobson]. *Unique type* in Amsterdam Museum.

Subfamily DIADOCIDIINAE.

Chiasmoneura, de Meij.

- Tijd. v. Ent., LVI, 323 (1913).
 GENOTYPE, *C. anthracina*, *id.*, *loc. cit.*, sp. nov.
anthracina de Meij., *loc. cit.*, 324, ♀, pl. xv, 1b, wing (1913).

¹ I followed Johannsen in uniting this genus with *Mycomyia* in my Fauna volume, but both Coquillett and de Meijere keep them separate.

Wonosobo, Java [*Jacobson*]. *Type*, unique, in Amsterdam Museum.

Subfamily MYCETOBIIINAE.

Mesochria, Enderl.

Tr. Linn. Soc. Lond. XIV, 65 (1910).

GENOTYPE, *M. scoffiana*, sp. nov., *loc. cit.*

cinctipes de Meij., Tijds. v. Ent. LVI, 322, ♂ (1913).

Batavia, viii [*Jacobson*]. *Type* in Amsterdam Museum.

Subfamily MYCETOPHILINAE.¹

(For tab. orient. gen. v. Brun., Fauna Brit. Ind., 80).

Boletina, Staeg.

Naturh. Tidsskr. III, 233 (1840).

Johan., Gen. Ins., Fasc. 93, 73.

GENOTYPE, *Leia trivittata* Mg., by designation of Johannsen.

brevirostris Brun., Fauna Brit. Ind., 83, ♂, pl. iii, 7, head (*Gnoriste*) (1912).

Mundali, Dehra Dun, 9000 ft., 12-v-10 [*Beebe*]. *Type*, unique, in Indian Museum.

The name of this species is misleading now that it is referred here, but was applicable when placed in *Gnoriste*. My error was kindly pointed out to me by Mr. Edwards (*in litt.*).

Leptomorphus, Curt.

Brit. Entom. 365 (1831).

Sch., F. Austr. II, 453: Johan., Gen. Ins., Fasc.

93, 71: Brun., Fauna Brit. Ind., 84, pl. ii, 1, wing.

GENOTYPE, *L. walkeri* Curt., by original designation.

ornatus Brun., Fauna Brit. Ind., 85, ♀ (1912).

Sylhet, 3-ii-05 [*Hall*]. *Type*, unique, in Indian Museum.

Palacoanaclinia, Meun.

Monog. Mycetoph., 143 (1904).

Johan., Gen. Ins., Fasc. 93, 85, pl. v, 22, 23,

wing (1909); Brun., Fauna Brit. Ind., 85, pl.

ii, 7, wing (1912).

GENOTYPE, *P. distincta* Meun., by original designation.

spathulata Brun., Fauna Brit. Ind., 84, ♂, pl. iii, 8, head (*Gnoriste*) (1912).

Darjiling, 29-v-10 [*Brunetti*]. *Type*, unique, in Indian Museum.

¹ For table of genera see Kieff., Gen. Ins., Fasc. 93, 52.

flavohirta Brun., *loc. cit.*, 86, ♂ (1912).

Darjiling, 11-viii-09 [*Jenkins*]. *Type*, unique, in Indian Museum.

Allactoneura, de Meij.

Tijd. v. Ent. L, 201 (1917).

Johan., Gen. Ins., Fasc., 93, 70: Brun., Fauna Brit. Ind., 88.

GENOTYPE, *A. cincta* de Meij., the original species.

Scottella, Enderl., Trans. Linn. Soc. Lond. XIV, 60 (1910).

GENOTYPE, *S. argenteosquamosa*, sp. nov., *loc. cit.*

obscurata Walk., Proc. Linn. Soc. Lond. VIII, 130 (*Mycetophila*) (1865).

Salawatti. *Type* in British Museum, in fair condition, abdomen missing.

cincta de Meij., Tijd. v. Ent. L, 202, ♂ ♀, pl. v, 2, 3, wing (1907).

Brun., Fauna Brit. Ind., 88 ♂ ♀, pl. i, 17, wing: Meij., Tijd. v. Ent. LVI, 321.

Scottella formosana, Enderl., Trans. Linn. Soc. Lond. XIV, 63, ♀, fig. 3, wing (1910).

Allactoneura formosana, de Meij., Tijd. v. Ent. LVI, 320.

Scottella argenteosquamosa, Enderl., Trans. Linn. Soc. Zool. (2) XIV, 60 (1910).

Sylhet, 1-i-05 [*Hall*]; Thaumaspur, Nepal, 18-20-ii-08; Kobo, N.E. Ind. Frontier, 2-xii-11 [*Kemp*]; Peradeniya, ix, xii-09 [*Green*]; 10-30-vi-10 and 15-viii-10 [both *Gravelly*]; Semarang, ii, iii-06; Buitenzorg, ix; Nongkodjadjar, i; Wonosobo, iv [all Java and *Jacobson*]; Kyukokado, Formosa, 6-i-08 [*Sauter*]. *Types* of *cincta* in Amsterdam Museum, of *S. formosana* in Stettin Zoological Museum.

nigrofemorata de Meij., Tijd. v. Ent. LVI, 322, ♀.

Nongkodjadjar, Java, i [*Jacobson*]. *Type*, unique, in Amsterdam Museum.

Odontopoda, Aldr.

Rep. Geol. Indiana, XXI, 187 (1898).

Johann., Gen. Ins., Fasc. 93, 70: Brun., Fauna Brit. Ind., 89, pl. i, 16, wing.

GENOTYPE, *O. sayi* Aldr., by original designation.

Proanaclinia, Meun., Monog. Myceto., 145 (1904).¹

indica Brun., Fauna Brit. Ind., 90, ♂ (1912).

Darjiling, 1-x-08 [*Brunetti*]. *Unique type* in Indian Museum.

Anomalomyia, Hutton.

Index Fauna Nov. Zeal., 134 (1904).

Johann., Gen. Ins., Fasc. 93, 83: Brun., Fauna Brit. Ind. 90, pl. ii, 6, wing.

¹ I cannot obtain access to this work.

Anomala, Marshall (*nec* Stephens), Trans. N. Zeal. Inst. XXVIII, 293 (1896).

GENOTYPE, *Mycetophila guttata* Hutton, by original designation.

indica Brun., Fauna Brit. Ind., 91, ♂ (1912).

Darjiling, 7-viii-09 [*Paiva*]. Unique *type* in Indian Museum.

Acrodicrania, Skuse.

Proc. Linn. Soc. N. S. Wales (2) III, 1194 (1888).

Johann., Gen. Ins., Fasc. 93, 81: Brun., Fauna Brit. Ind. 92.

GENOTYPE, *A. atricauda* Skuse, by original designation.

ferruginea Brun., Fauna Brit. Ind., 93, ? ♀ (1912).

Khasi Hills, 1000-3000 ft., x-06. Unique *type* in Pusa coll.

incerta, *id.*, *loc. cit.*, 93, ♀ (1912).

Kurseong, 20-vi-10 [*Annandale*]. Unique *type* in Ind. Mus.

Clastobasis, Skuse.

Proc. Linn. Soc. N.S. Wales (2) V, 617 (1890).

Johann., Gen. Ins., Fasc. 93, 80: Brun., Fauna Brit. Ind., 94, pl. ii, 5, wing.

GENOTYPE, *C. tryoni* Skuse, by original designation.

vittigera Brun., Fauna Brit. Ind., 95, ♀ (1912).

Sadiya, Assam. Unique *type* in Indian Museum.

Leia, Mg.¹

Syst. Besch. I, 253 (1818).

Sch., F. Austr. II, 459; Johann., Gen. Ins., Fasc. 93, 76: Brun., Fauna Brit. Ind., 96: *Auctt.*

GENOTYPE, *L. fascipennis* Mg., by Curtis's designation, 1837.

Lejomya, Rond., Dipt. Ital. Prod. I, 195 (1856).

Lejosoma, *id.*, *loc. cit.*, I, Corrigenda (1856).

Glaphyroptera, Winn., Verh. z.b. Ges. Wien XIII, 781 (1863).

GENOTYPE, *Leia fascipennis* Mg., by Coquillett's designation.

Neoglaphyroptera, Ost. Sack., Cat. N. Amer. Dipt., 10, *nom. nov.* (1878).

Greenomyia, Brun., Fauna Brit. Ind., 87 (1912).

GENOTYPE, *G. nigricoxa*, *id.*, *loc. cit.*, sp. nov.

winthemi Lehm., Ins. spec., etc., 39 (1822): Meig., Syst. Besch.

VII, 48: Winn., Verh. zool.-bot. Ges. Wien XIII, 789

(*Glaphyroptera*): Wulp, Dipt. Sum. Exped., 7 (*Glaphyroptera*):

Auctt. var.: Brun., Fauna Brit. Ind., 97, ♂ ♀, pl. ii, 2 wing.

¹ Respecting *Leia* v. *Lejomya*, see Johann., Gen. Ins., Fasc. 93, 76, and Edw., Ann. Mag. Nat. Hist. (8) XII, 201 (1913).

Mycetophila maculipennis, Say, in Long's Exped. St. Peter's Riv., App. 365 (1824).

Leia trifasciata, Walk., List Dipt. Brit. Mus. I, 93 (1848).

Simla, 25-iv-07, 5-v-07, 10-v-09; Matiana, Simla District, 28—30-iv-07 [all *Annandale*]; Naini Tal, 2-vi-09; Darjiling, 29-v-10 [*Brunetti*, *Paiva*]; Ukhrul, Manipur [*Pettigrew*]; Sumatra. Also occurs in Europe and North America. *Type*: the location of this unknown to me.

bimaculata Walk., List Dipt. Brit. Mus. I, 100 (*Mycetophila*) (1848).

East Indies. *Type* in British Museum in fair condition (no abdomen). Mr. Edwards informs me it belongs to this genus.

maculicoxa Enderl., Trans. Linn. Soc. Lond. XIV, 69 (1910).

Edw., Ann. Mag. Nat. Hist. (8) XII, 200, note.

Rhymosia fascipes, Brun., Fauna Brit. Ind., 106, ♂, pl. iii, 12, ♂ genitalia (1912).

Peradeniya, x, 1912 [*Edw.*], 7-viii-10 [*Gravelly*]. *Type* of *fascipes* in Indian Museum, location of that of *maculicoxa* unknown to me.

nigricoxa Brun., Fauna Brit. Ind., Dipt., 87, ♂, pl. ii, 8, wing; pl. iii, 9, ♂ genitalia (*Greenomyia*) (1912).

Leia spathulata, *id.*, *loc. cit.*, 101, ♂ (1912).

Phagu, 3-v-07, *type* of *nigricoxa* [*Annandale*]; Darjiling, 25-v-10, *type* of *spathulata* [*Brunetti*]. The identity of these two species was pointed out to me by Mr. Edwards (*in litt.*).

flavobrunnea Brun., Fauna Brit. Ind., 99, ♂ (1912).

Darjiling, 29-v-10 [*Brunetti*]. *Type*, a unique specimen, in Indian Museum.

arcuata, *id.*, *loc. cit.*, 99, ♂ ♀, pl. ii, 3, wing (1912).

Brun., Rec. Ind. Mus. XIII, 62.

Kurseong, 9-vii-08, *type* ♂, 5-vii-08, *type* ♀, 3-vii-08, 15-iv-11, 26-vi-10; Darjiling; Naini Tal, 10-vi-09; Simla, x-08 [*Howlett*]; Bhowali, Kumaon, 28-vi-10 and vii-09 [*Imms*]; Khasi Hills (1000—3000 ft.), iii-07; Komsing, N.E. Ind. Frontier, 1-iii-12 [*Kemp*]; Ceylon. *Type* in Indian Museum, *co-type* in Pusa collection.

bicolor, *id.*, Fauna Brit. Ind., 100, ♀ (1912).

Kurseong, 7-ix-09 [*Annandale*]. *Type*, a unique specimen, in Indian Museum.

insignis, *id.*, *loc. cit.*, 101, ♀ (1912).

Ukhrul, Manipur (6400 ft.). *Type*, a unique specimen, in Indian Museum.

nigra, *id.*, *loc. cit.*, 101, ♀ (1912).

Naini Tal, 2-vi-09, *type*; Simla, 9-v-09 [*Annandale*]; Senchal, v-13 [*Lord Carmichael* coll.]. *Type* in Indian Museum.

demeijerei *nom. nov.*

- L. nigra*, de Meij., Bijd. tot. Dierk. XIX, 46, pl. i, 3, wing (1913).
Ceram, ii. Type in Amsterdam Museum
The name *nigra* is preoccupied by me in 1912.

Rhymosia, Winn.

- Verh. z.b. Ges. Wien XIII, 810 (1863).
Sch., F. Austr. II, 463: *Auctt. var.*: Johan.,
Gen. Ins., Fasc. 93, 102: Brun., Fauna Brit.
Ind., 102.
GENOTYPE, *R. fasciata* Mg., by designation of
Johannsen (Gen. Ins., Fasc. 93, 102, 1909).
- flavolimbata* Brun., Fauna Brit. Ind., 103, ♂ ♀ (1912).
Kurseong, 4-vii-08, type, 21-vi-10 [both *Annandale*]; Simla,
x-03 [*Howlett*]; Kumaon, vii-09 [*Imms*]. Type in Ind. Mus.
- genitalis*, *id.*, *loc. cit.*, 104, ♂ (1912).
Darjiling, 29-v-10 [*Brunetti*]. Type, a unique specimen, in
Indian Museum.
- albolateralis*, *id.*, *loc. cit.*, 104, ♂ (1912).
Naini Tal, 10-vi-09. Type in Indian Museum, a unique
specimen.
- annulicornis*, *id.*, *loc. cit.*, 105, ♂ ♀, pl. ii, 10, wing; pl. iii, 13, 14,
genitalia and antennae (1912).
Madhupur, Bengal, 13-x-09, type ♂ ♀, 15-x-09 [*Paiva*]; Cal-
cutta, 18-vii-08 [*Annandale*]; Peradeniya, 14-16-vii-10 [*Gravelly*].
Types in Indian Museum.
- humeralis*, *id.*, *loc. cit.*, 106, ♂ (1912).
Sylhet, 3-ii-05 [*Hall*]. Type, a unique specimen, in Indian
Museum.

Allodia, Winn.

- Verh. zool.-bot. Ges. Wien XIII, 826 (1863).
Sch., F. Austr. II, 466: *Auctt. var.*: Johan.,
Gen. Ins., Fasc. 93, 103: Brun., Fauna Brit.
Ind., 107.
GENOTYPE, *Mycetophila lugens* Wied., by Johann-
sen's designation.
- Brachycampta*, Winn., Verh. z.b. Ges. Wien XIII,
833 (1863).
GENOTYPE, *Mycetophila alternans* Zett., by Co-
quillett's designation.
- nigrofasciata* Brun., Fauna Brit. Ind., 108, ♂ ♀, pl. ii, 11, wing
(1912).
Simla, 10-v-09, type ♂ ♀ [*Annandale*]; Kufri, v; Mundali,
Dehra Dun, 9000 ft., 10-12-v-10 [*Beebe*]. Types in Indian
Museum.

Macrobrachius, Dzied.

- Hor. Soc. Ent. Ross. XXIII, 520 (1889).
Johan., Gen. Ins., Fasc. 93, 98: Brun., Fauna
Brit. Ind., 108.

GENOTYPE, *M. kowarzii* Dzied., by original designation.

longicosta Brun., Fauna Brit. Ind., 109, ♂, pl. iii, 11, ♂ genitalia (1912).

Kurseong, 25-vi-10 [Annandale]. Type, a unique specimen, in Indian Museum.

Phronia, Winn.

Verh. zool.-bot. Ges. Wien XIII, 857 (1863).

Sch., F. Austr. II, 471: *Auctt. var.*: Johan., Gen.

Ins., Fasc. 93, 95: Brun., Fauna Brit. Ind., 110, pl. ii, 9, wing.

GENOTYPE, *P. rustica* Winn., by designation of Johannsen.

simplex Brun., Fauna Brit. Ind., 111, ♀ (1912).

Assam. Type, unique, in Indian Museum.

semifumata, *id.*, *loc. cit.*, 111, ♂, pl. iii, 10, ♂ genitalia (1912).

Tonglu, Darjiling, 21-iv-10 [Beebe]. Type in Indian Museum, a unique specimen.

Exechia, Winn.

Verh. zool.-bot. Ges. Wien XIII, 879 (1863).

Sch., F. Austr. II, 475: *Auctt. var.*: Johan.,

Gen. Ins., Fasc. 93, 106: Brun., Fauna Brit. Ind., 112.

GENOTYPE, *E. fungorum* DeGeer, by Johannsen's designation.

Parexechia, Bech., Ins. v. Jan Mayen, 62 (1886).

GENOTYPE, *P. concolor*, *id.*, sp. nov., *loc. cit.*, pl.

v., fig. 3, 3a—3c, full ins. var. chars.

Brachydicerania, Skuse, Proc. Linn. Soc. N.S. Wales (2) III, 1215 (1888).

GENOTYPE, *B. pullicauda*, *id.*, sp. nov., *loc. cit.*

basilinea Brun., Fauna Brit. Ind., 113, ♂ ♀, pl. ii, 12, wing (1912).

Kurseong, 3—4-vii-08 [Annandale]; Simla, x-08 [Howlett]. Types, a unique pair, in Indian Museum.

Euryschalis, Brun.

Fauna Brit. Ind., App., 559 (1912).

GENOTYPE, *E. spectralis*, the original species.

spectralis Brun., *loc. cit.*, 560, ♂ (1912).

Kurseong, 16-iv-11 [Annandale]. Type, a unique specimen, in Indian Museum.

Mr. Edwards informs me (*in litt.*) that *Euryschalis* may be synonymous with *Coelosia* Winn., but as there are no specimens at hand of the latter genus for comparison, I leave the question open for the present. The costa attains the wing tip in *Euryschalis*; the genital organs are very different to those figured in the Genera Insectorum; the fore metatarsi are about equal to the corresponding tibiae.

Mycetophila, Mg.

Illig. Mag. II, 263 (1803).

Sch., F. Austr. II, 484: *Auctt.*: Johan., Gen. Ins., Fasc. 93, 115: Brun., Fauna Brit. Ind., 113.GENOTYPE, *M. agarici* Oliv., proposed by Johannsen (Gen. Ins., Fasc., 93, 106, 1909).*Mycetina*, Rond., Dipt. Ital. Prod. I, 195 (1856).GENOTYPE, *Mycetophila flavipes* Macq., by original designation.*Mycozetoea*, *id.*, *loc. cit.*, IV, Corrigenda, 12 (1861).*cinctiventris* Brun., Fauna Brit. Ind., 115, ? ♂ (1912).Simla, 10-vii-09 [Annandale], x-08 [Howlett]. *Type*, unique, in Indian Museum.*quadrifasciata*, *id.*, *loc. cit.*, 115, ♂, pl. ii, 13, wing (1912).Simla, 10-v-09 [Annandale]. *Type*, unique, in Ind. Mus.*curvilinea*, *id.*, *loc. cit.*, 116, ♀, pl. ii, 14, wing (1912).Darjiling, 10-16-x-05 [Brunetti]. *Type*, unique, in my collection.*suffusa*, *id.*, *loc. cit.*, 117, ♀ (1912).Simla, 10-v-09 [Annandale]. *Type*, unique, in Ind. Mus.*himalayensis*, *id.*, *loc. cit.*, 117, ♂ (1912).Simla, 10-v-09 [Annandale]; Naini Tal, 10-v-09. *Type* in Indian Museum.*binotata* *id.*, *loc. cit.*, 118, ♂ ♀ (1912).*Id.*, Rec. Ind. Mus., XIII, 63.Darjiling, 20-ix-08, *type*, ♂, 26-ix-08, 29-v-10 [all Brunetti]; Simla, 10-v-09, *type* ♀ [Annandale], 8-x-11 [Howlett]; Ukhrul, Manipur. *Types*, ♂ ♀, in Indian Museum.Mr. Edwards calls my attention to the name *M. binotata*, Halid., which, as he states, is a synonym of *Zygomia pictipennis*. The name of my species need not therefore be changed.**Delopsis, Skuse.**

Proc. Linn. Soc. N.S. Wales (2) v, 623 (1890).

Brun., Fauna Brit. Ind., 118: Johan., Gen. Ins., Fasc. 93, 115.

GENOTYPE, *D. flavipennis* Skuse, by original designation.*griseolateralis*, *id.*, *loc. cit.*, 116, ♂ (*Mycetophila*) (1912).Tonglu, Darjiling, 10,000 ft., 21-iv-10 [Beebe]. *Type*, unique, in Indian Museum.*brunettii* Edwards, Ann. Mag. Nat. Hist. (8) XII, 55, *nom. nov.* (1913).*collaris*, Brun., Fauna Brit. Ind., 119, ♂ (1912).Maddathorai (Travancore), 17-xi-08 [Annandale]. *Type* in Indian Museum, unique.This species is renamed by Edwards who finds that *Mycetophila collaris*, Enderl., from the Seychelles, is a *Delopsis*.

Family BLEPHAROCERIDAE.

Hammatorhina, Loew.

Bull. Soc. Ent. Ital. I, 4 (1869).

Brun., Fauna Brit. Ind., 152.

Kellogg, Gen. Ins., Fasc. 56, 9.

GENOTYPE, *H. bella*, the original and only species.*bella* Loew, Bull. Soc. Ent. Ital. I, 96, pl. ii, 4-6 (1869).

Kellogg, Gen. Ins., 9, pl. ii, 17, antenna, 27, wing: Brun.,

Fauna Brit. Ind., 152, fig. 14, wing, antenna.

Ceylon. *Type*, location uncertain.

Apistomyia, Big.

Ann. Soc. Ent. Fran. (4) II, 109 (1862).

Kellogg, Gen. Ins., Fasc. 56, 8.

Brun., Fauna Brit. Ind., 153.

GENOTYPE, *A. elegans* Big.*trilineata* Brun., Rec. Ind. Mus. IV, 315, ♂ (1911).*Id.*, Fauna Brit. Ind., 154, ♀.Kurseong, 4-ix-09 [Annandale]. *Type*, a unique ♂, in Indian Museum.

Blapharocera, Macq.

Ann. Soc. Ent. Fran. (2) I, 61 (1843) (*Blepharicera*).Sch., F. Austr. II, 637 (*Blepharicera*): Kellogg,

Gen. Ins., Fasc. 56, 11: Brun., Fauna Brit. Ind.,

155 (*Blepharocera*).GENOTYPE, *Asthenia fasciata* Westw. (as *Blepharicera limbipennis*, sp. nov.).*Asthenia*, Westwood, Guer. Mag. d. Zool. (2) IV,

pl. xciv (1842); preocc. Hubn. 1816.

GENOTYPE, *Asthenia fasciata* Westw., by original designation.*indica* Brun., Rec. Ind. Mus. IV, 316, ♂ ♀ (1911).*Id.*, Fauna Brit. Ind., 156, ♂ ♀, fig. 15, wing.Phagu, 12-15-v-00 [Annandale]. *Types* in Indian Museum.

Philorus, Kellogg.

Proc. Acad. Sci. Calif. (3) Zool. 3, 199 (1903).

Kellogg, Gen. Ins., Fasc. 56, 14.

GENOTYPE, *Liponeura bilobata* Loew.¹*bionis* Agharkar, Rec. Ind. Mus. X, 160, ♂ ♀, pl. xvi, figs. of

various larvae, pl. xvii, full figs. ♂ ♀, head ♂ ♀, genitalia

♂ ♀, wing (1914).

Nagaberan, Kashmir. *Types* in Indian Museum.¹ Apparently this species was selected by Kellogg as the type.

Family BIBIONIDAE.

Subfamily SCATOPSINAE.

Scatopse, Geoff.¹

Hist. Nat. II, 545 (1764).

Sch., F. Austr. II, 348: *Auctt. var.*: Brun.,
Fauna Brit. Ind., 179.*Scathopse*, Geoff., Hist. abregée Ins. II, 544 (1762).GENOTYPE, *Tipula notata* L., by designation of
Latreille, as *Tipula albipennis* F.**brunnescens** Brun., Rec. Ind. Mus. IV, 281, ♀ (1911).*Id.*, Fauna Brit. Ind., 180, ♀, fig. 20, wing.Peradeniya, 9-vii-10 [Gravelly]. *Type*, unique, in Ind. Mus.**nigronitida** Brun., Rec. Ind. Mus. IV, 281, ♂ ♀ (1911).*Id.*, Fauna Brit. Ind., 181, ♂ ♀, fig. 21, antenna, wing, leg.Dharampur, 5000 ft., Simla Hills, 14-v-08, two pairs *in cop.*
[Annandale]. *Types* in Indian Museum.**crassiuscula** de Meij., Tijd. v. Ent. LVI, 325, pl. xv, 2, wing (1913).Batavia, x [Jacobson]. *Type*, unique, in Amsterdam
Museum.**palliditarsis**, *id.*, *loc. cit.*, 325, ♀, pl. xv, 3, wing (1913).Banjuwangi, E. Java [MacGillavry]. *Type*, unique, in
Amsterdam Museum.**pallidicornis**, *id.*, *loc. cit.*, 326, pl. xv, 4, wing (1913).Semarang, x [Jacobson]. *Type* in Amsterdam Museum.**crassicornis**, *id.*, *loc. cit.*, 326, pl. xv, 5, wing (1913).Batavia, x [Jacobson]. *Type* in Amsterdam Museum.

Subfamily BIBIONINAE.

Crapitula, Gimm.

Bull. Soc. Imp. Nat. Mosc. XVIII, 2, 330 (1845).

GENOTYPE, *Penthetria melanaspis* Wied., by original
designation.*Pleciomyia*, Brun., Rec. Ind. Mus. IV, 269 (1911):*id.*, Faun. Brit. Ind., 160.GENOTYPE, *Penthetria melanaspis* Wied., by original
designation.**melanaspis** Wied., Auss. Zweifl. I, 72 (*Penthetria*) (1828).Brun., Rec. Ind. Mus. IV, 270 (*Pleciomyia*): *id.*, Fauna Brit.Ind., 161, ♂ ♀, pl. xii, 11, wing, 14, antenna (*Pleciomyia*).*Penthetria japonica*, Wied., Auss. Zweifl. II, 618 (1830).*Crapitula motschulskii*, Gimm., Bull. Soc. Imp. Nat. Mosc. XVIII,
2, 330, fig. (1845).

¹ Coquillet upholds the original spelling, *Scathopse*, but as no named species was quoted at its erection, though two unnamed species were described, this generic name cannot be accepted.

Id., *id.*, Loew., Berl. Ent. Zeits. II, 106, pl. i, 12.

Crapitula melanaspis, Brun., Rec. Ind. Mus. IX, 256.

Plecia ignicollis, Walk., List Dipt. Brit. Mus. I, 116 (1848).

Plecia melanaspis, Auctores.

Himalayas (many locs.); Nepal; Assam; N.E. Ind. Frontier; China; Japan; Siberia. Probably common everywhere throughout the temperate regions of the Orient, and in Eastern Asia from about March to October. Not found in the plains in India. Occurs in Australia. *Types* in the Leyden and Vienna Museums.

Plecia, Wied.

Auss. Zweifl. I, 72 (1828).

Brun., Rec. Ind. Mus. IV, 270: *Id.*, Fauna Brit. Ind. 162: Auctores.

GENOTYPE, *Hirtea fulvicollis* F., by designation of Blanchard (Hist. Nat. Ins. III, 576, 1840).

Eupeitenus, Macq., Dipt. Exot. I, 88 (1838).

GENOTYPE *E. atra* Macq., by original designation.

Rhinoplecia, Bell., Ditt. Messico I, 16 (1859).

GENOTYPE, *Plecia rostellata* Lw., as *rostrata*, sp. nov., by original designation.

fulvicollis F., Syst. Antl., 53 (*Hirtea*) (1805).

Plecia fulvicollis, Wied., Auss. Zweifl. I, 73: Macq., Hist. Nat. Dipt. I, 176: Dol., Nat. Tijd. Ned. Ind. X, 407, pl. vi, 3: Ost. Sack., Ann. Mus. Gen. XVI, 397: *id.*, Berl. Ent. Zeits. XXVI, 96: de Meij., Tijd. v. Ent. LIII, 59, larva and pupa, pl. iv, 1, 2: Brun., Rec. Ind. Mus. IV, 271: *id.*, Fauna Brit. Ind., 163, ♂ ♀, pl. xii, 12, wing, 15, antenna.

Plecia dorsalis, Walk., Proc. Linn. Soc. Lond. I, 5 (1857), Singapore, Mount Ophir.

Plecia subvarians, *id.*, *loc. cit.*, I, 105 (1857), Sarawak.

Penthetria thoracica, Guer., Belanger's Voy. aux Indes Orient., Zool., 557, pl. iv, 9 (1833).

India; Ceylon; Burma; Assam; Java; Sumatra; Borneo; Ternate; Celebes; Dutch Papua; Philippines; probably common throughout the Oriental Region (though confined mainly to the plains), during the greater part of the year. *Type*. Probably no individual specimen was ever selected as such: the species is found in the Fabrician and other older collections.

tergorata Rond., Ann. Mus. Gen. VII, 462 (1875).

Ost. Sack., *loc. cit.*, XVI, 397: Brun., Rec. Ind. Mus. IV, 271, notes: *id.*, Fauna Brit. Ind. 164, pl. xii, 17, antenna.

Bhim Tal, 19-27-ix-06, including ♂ ♀, *in cop.*; Sukna, 1-vii-08 [both *Annandale*]; Kurseong, 6-9-viii-09 [*Paiva*]; Mussoorie, x-06; Bhowali, 13-vi-10 [*Imms*]; Shan Hills, Upper Burma, 26-vi-10 [*Annandale* and *J. C. Brown*]; Shillong, 5500-6400 ft., 29-viii-5-ix-15 [*Kemp*]; Khasi Hills, 17-iii-07; Malabar,

5-viii-07; Singapore; Java; Sarawak. Type in Genoa Museum.

forcipata Ost. Sack., Ann. Mus. Gen. XVI, 397 (1881).

de Meij., Tijd. v. Ent. LIII, 63.

Sumatra; Semarang, iii, viii, Wonosobo, iv [both Java and Jacobson]. Type in Genoa Museum.

tristis Wulp, Notes Leyden Mus. VI, 251, ♀ (1884).

de Meij., Tijd. v. Ent. LVI, 327, note.

Mount Ardjoeno, East Java; Gunung Gedeh, iii; Nongkodjadar [both Java and Jacobson]. Type in Leyden Museum.

indica Brun., Rec. Ind. Mus. IV, 271, ♂ ♀ (1911).

Id., Fauna Brit. Ind., 165, ♂ ♀.

Darjiling, type ♂ ♀, 2-x-08 and 10—16-x-05 [both Brunetti]; Theog., 27-iv-06; Kumaon, ix-06 [both Annandale]; Mussoorie, x-06; Soondrijal, Nepal; Kimoli, 24-x-07; Kangra Valley (4500 ft.); Ukhrul, Manipur [Pettigrew]; Khasi Hills, 17-iii-07. Types in Indian Museum; cotypes in Pusa coll. and my own.

atra Brun., Rec. Ind. Mus. IV, 272, ♀ (1911).

Id., Fauna Brit. Ind., 165, ♀, pl. xii, 13, wing, 16, antenna.

Bhim Tal (4500 ft.), including type, 19—22-ix-06 [Annandale]; Soondrijal, Nepal. Type in Indian Museum.

obscura Brun., Rec. Ind. Mus. IV, 272, ♂ ♀ (1911).

Id., Fauna Brit. Ind., 166, ♂ ♀.

Mussoorie, 24-v-05, types [Brunetti]. Types in my collection.

impostor, *id.*, Rec. Ind. Mus. VII, 446, ♂ ♀ (1912).

Airadeo, Kumaon Div., 6,880 ft., 31-v-12, type ♂ ♀; Bhowali, 25-vi-12 [both Imms]. Type ♂ ♀ in Indian Museum, cotype ♀ in Forest Zoology coll., Dehra Dun.

Parapleciomyia, Brun.

Rec. Ind. Mus. IX, 446 (1912).

GENOTYPE, *P. carbonaria* Brun., by original designation.

carbonaria Brun., *loc. cit.*, 447, ♂ ♀ (1912).

Bhowali, 11-v-12, type ♂; Sat Tal, near Kumaon, 9-v-12 [both Imms]. Type, a unique pair, in Indian Museum.

Bibio, Geoff.

Hist. Nat. Ins. II, 571 (1764).

Sch., F. Austr. II, 357; Brun., Rec. Ind. Mus.,

IV, 273, tab. orient. spp.: *id.*, Fauna Brit. Ind.,

166, fig. 17, head ♂ ♀: *Auctt. var.*

GENOTYPE, *B. hortulanus* L., by Latreille's designation.

Pullata, Harris, Expos. Engl. Ins., 77 (1776).

GENOTYPE, *Tipula pomonae* F., as *funestus*, sp. nov., by original designation.

Hirtea, Mg., Illig. Mag. II, 264 (1803).

GENOTYPE, no species selected.

- johannis** L., Syst. Nat. XII, 2, 976 (1767).
Sch., F., Austr. II, 361: Brun., Rec. Ind. Mus. IV, 278: *id.*,
Fauna Brit. Ind. 172, ♂ ♀: *Auctt. var.*
Bibio rufipes, Mg., Syst. Besch. VII, 55 (1838).
Hirtea praecox, Fab., Syst. Antl., 51 (1805).
Hirtea hyalina et praecox, Mg., Klass. I, 110, 111 (1804).
Hirtea johannis et pyri, *Auctt.*
Tipula pomonae, Schrk., Fauna Boica III, 79 (1803).
Tipula pyri, F., Ent. Syst. IV, 249 (1794).
Matiana, 28—30-iv-07: Theog., 2-v-07 [both *Annandale*].
Type, location unknown to present author.
A common European species.
- bicolor** Walk., List Dipt. Brit. Mus. I, 121, ♀ (1848).
East Indies. *Type* in British Museum in good condition.
Van der Wulp (Cat.) quotes East India as the locality, which
is quite different. This error is frequent in his Catalogue.
- obediens** Ost Sack., Ann. Mus. Gen. XVI, 395, ♀ (1881).
Hatam, Papua vii, 1875 [*Beccari*]; Heuvel-Biwak, xi, ♀.
Type, a unique ♀, in Genoa Museum.
- plecioides** Ost. Sack., *loc. cit.*, 396, ♀ (1881).
Hatam, Papua. vii, 1875 [*Beccari*]. *Type* in Genoa Museum.
- rubicundus** Wulp., Notes Leyd. Mus. VI, 251, ♀ (1884).
Java [*Muller*]. *Type*, a unique ♀, in Leyden Museum.
Possibly identical with *obediens* Ost. Sack.
- obscuripennis** de Meij., Bijd. tot Dierk. XVII, 86, ♂ ♀ (1904).
Brun., Rec. Ind. Mus. IV, 275, notes: *id.* Fauna Brit. Ind.,
170, ♂ ♀, fig. 18, wing.
Darjiling, 16-x-05 [*Brunetti*], 27-v-10 [*D'Abreu*]; Kurseong,
27—28-iv-10 [*D'Abreu*]; Chitlong, Nepal; Mt. Tahe-pum, 4000
—5000 ft., N.E. Burmese Frontier. xi-10 [*C. W. Bebe*]; Rotung,
Dihang Riv., 25-xii-11 [*Kemp*]; Matiana, 28—30-iv-07 [*Annandale*];
Naini Tal, 26-ix-07. *Type* in Amsterdam Museum.
- hortulanoides** Brun., Rec. Ind. Mus. IV, 274, ♂ ♀ (1911).
Id., Fauna Brit. Ind., 169, ♂ ♀.
India (probably Darjiling District), *type* ♂; Kurseong, 15-v-10,
type ♀; Darjiling District, vi-12. *Types* in Indian Museum.
- proximus** Brun., Rec. Ind. Mus. IV, 275, ♂ (1911).
Id., Fauna Brit. Ind., 171, ♂.
Darjiling, xi-1910 [*D'Abreu*]. *Type*, a unique ♂, in Ind. Mus.
- abdominalis** Brun., Rec. Ind. Mus. IV, 276, ♂ ♀ (1911).
Id., Fauna Brit. Ind., 172, ♂ ♀.
Western Himalayas; Badrinath, 10,200 ft., Garhwal Distr.,
27-v-10, ♂ ♀, *types in cop.* [*A. D. Imms*]; Phagu, 11-v-09
[*Annandale*]. *Types* in Indian Museum.
- approximatus** Brun., Rec. Ind. Mus. IV, 277, ♀ (1911).
Id., Faun. Brit. Ind., 173, ♀.
Kurseong, xi-10 [*D'Abreu*]. *Type*, a unique ♀, in Ind. Mus.
- discalis** Brun., Rec. Ind. Mus. IV, 278, ♀ (1911).

- Id.*, Fauna Brit. Ind., 174, ♀.
Phagu, 11-v-09 [Annandale]. *Type*, a unique ♀, in Ind. Mus.
- rufifemur** Brun., Rec. Ind. Mus. IV, 279, ♀ (1911).
Id., Fauna Brit. Ind. 175, ♀.
Darjiling, 20-x-05 [Brunetti]. *Type* and *cotype* in my collection.
- fuscitibia** Brun., Rec. Ind. Mus. IV, 279, ♂ (1911).
Id., Fauna Brit. Ind., 175, ♂.
Phakia, Kumaon Div., 10,700 ft., 3-vi-09 [Imms]. *Type* in Indian Museum.
- defectus** Brun., Rec. Ind. Mus. IV, 280, ♂ (1911).
Id., Fauna Brit. Ind., 176, ♂.
Kurseong, 13-viii-07 [Paiva]. *Type* in Indian Museum, a unique specimen.
- nigripennis**, *id.*, Rec. Ind. Mus. IX, 256, ♀ (1913).
Senchal, Darjiling, 8000 ft., v-13. *Type*, a unique ♀, in inferior condition, in Indian Museum.

Dilophus, Mg.

- Illig. Mag. II, 264 (1803).
Sch., F. Austr. II, 355: Brun., Fauna Brit. Ind., 176: *Auctt.*
GENOTYPE, *D. febrilis* L., designated by Latreille.
Acanthocnemis, Blanchard, Gay's Hist. fis. polit. Chile. Zool. VII, 355 (1852).
GENOTYPE, unknown to present author.
- gratiosus** Big., Jour. Asiat. Soc. Beng. LIX, 265, ♂ ♀ (1890).
Brun., Rec. Ind. Mus. IV, 280, redescr. ♂ ♀: *id.*, Fauna Brit. Ind., 178, ♂ ♀, fig. 19, var. chars.
Dharamsala, W. Himalayas, ix-1890, *type* ♂ ♀ *in cop.* [Major Sage]; Kurseong, 24-vi-10; Theog., 2-v-07; Phagu, 11-v-09 [both Annandale]; Bhowali, Kumaon, 5700 ft., 20-vi-10; vii-1909 [both Imms]; Darjiling, 21-iv-10 [Beebe]; Sadon, Upper Burma, 5,000 ft., 2-ii-11, at light; Amangarh, Central India, 24-ii-10. Also occurs in Yunnan, S. China. *Types* in Ind. Mus.
- nigriventris** de Meij., Tijd. v. Ent. LVI, 327, ♀ (1913).
Nongkodjadjar, Java, i [Jacobson]. *Type*, a unique ♀, in Amsterdam Museum.

Family SIMULIIDAE.¹

Simulium, Latr.

- Hist. Nat. Crust. Ins. III, 426 (1802); XIV, 294 (1805).
Sch., F. Austr. II, 363 (*Simulia*): Brun., Rec. Ind. Mus. IV, 282, tab. orient. spp.: *id.*, Fauna Brit. Ind., 182, fig. 23, early stages.

¹ For family characters v. Sch., F. Austr. II, xvii and Brun., Fauna, 182.

Simulium et *Simulia*, Auctt.

Atractocera, Mg., Klass. I, 94 (1804).

GENOTYPE, *S. reptans* Mg., as designated by Zetterstedt. Coquillett suggests *columbaczensis* Schoub. (*Culex id.*), but *reptans* is most generally accepted.

indicum Becher, Jour. Asiat. Soc. Beng. LIII, 199, ♀, pl. xiv, var. chars (1884).

Brun., Rec. Ind. Mus. IV, 286, notes and *neotype* ♂ described.

Id., Fauna Brit. Ind., 191, ♂ ♀, figs. 24, 25, var. chars.

Simla, 24-iv-07 [Annandale]; Darjiling, 12-viii-09 [Jenkins]; Kurseong, 10-26-ix-09 [Lynch]; Bamon and Thadier (3500 ft.); Tous Valley, N.W. Himalayas [C. W. Rogers]; Sylhet, 2-iii-05, 18-iv-05 [Hall]; Bombay, 25-xi-09 (*neotype*); Khasi Hills; Ukhul, Manipur [Pettigrew].

Neotype ♂ in Indian Museum. Location of original type unknown to present author, probably no particular specimen selected as such from the series, all of which were females. Specimens from Rotung taken on the Abor Expedition, xii-1911 and 6-i-12 are probably this species.

nobile de Meij., Tijd. v. Ent. I, 206, ♂ (1907).

Semarang, iii [Jacobson]. *Type*, a unique ♂, in Amsterdam Museum.

rufithorax Brun., Rec. Ind. Mus. IV, 282, ♂ ♀ (1911).

Id., Fauna Brit. Ind., 187, ♂ ♀.

Kanara, Bombay, viii-07 (*type* ♂); Karwar, Bombay, viii-07;

Kurseong, 10-26-ix-09 [Lynch]; 7-ix-09 (*type* ♀) [Annandale].

Type ♂ in Pusa collection, *type* ♀ in Indian Museum.

grisescens Brun., Rec. Ind. Mus. IV, 283, ♂ (1911).

Id., Fauna Brit. Ind., 188, ♂.

Kurseong, 10-26-ix-09 [Lynch]. *Type* in Indian Museum, a unique specimen.

metatarsale Brun., Rec. Ind. Mus. IV, 284, ♂ (*metatarsalis*) (1911).

Id., Fauna Brit. Ind., 189, ♂.

Kurseong, 28-iii-10 [Annandale]. *Type*, a unique ♂, in Indian Museum.

griseifrons Brun., Rec. Ind. Mus. IV, 285, ♀ (1911).

Id., Fauna Brit. Ind., 190, ♀.

Kalighat, Kumaon (6000 ft.), 4-vi-10 [Imms]. *Type*, a unique ♀, in Indian Museum.

rufibasis Brun., Rec. Ind. Mus. IV, 285, ♀ (1911).

Id., Fauna Brit. Ind., 190, ♀.

Kurseong, 18-vi-10 [Annandale]. *Type*, a unique ♀, in Indian Museum.

aureohirtum Brun., Rec. Ind. Mus. IV, 287, ♂ ♀ (1911).

Id., Fauna Brit. Ind., 194, ♂ ♀ : de Meij., Tijd. v. Ent. LVI, 331.

Umling, Assam, iii-07 (including *type* ♂); Kanara, Bombay.

- viii-07 (including *type* ♀); Kurseong, 6-ix-09; Gunung Gedeh, Java, iii [Jacobson]. *Types* in Pusa collection.
- senile** Brun., Rec. Ind. Mus. IV, 288, ♂ (*senilis*) (1911).
Id., Fauna Brit. Ind., 191, ♂.
 Phagu, 8-v-07 [Annandale]. *Type*, a unique ♂, in Indian Museum.
- striatum** Brun., Spol. Zeyl. VIII, 90, ♂, pl., var. chars. (1912).
Id., Fauna Brit. Ind., App. 561, ♂: Patton and Cragg, Text Bk. Med. Ent. 172, pl. xxxii, 1-4, var. chars.
 Peradeniya, vii-1911 (*type*), xiii-10, 2-vi-19 [Green and Gravely].
 Kodaikanal, S. India, 6000 ft., very common, breeding in hill streams.
- eximium** de Meij., Tijds. v. Ent. LVI, 330, ♂ ♀ (1913).
 Nongkodjadar, Java, i [Jacobson]. *Types* in Amsterdam Museum.
- atratum**, *id.*, loc. cit., 331, ♂ ♀ (1913).
 Gunung Gedeh, Java, iii [Jacobson]. *Types* in Amsterdam Museum.
- argyrocinctum**, *id.*, loc. cit., 332, ♂ ♀ (1913).
 Gunung Gedeh, Java, iii [Jacobson]. *Types* in Amsterdam Museum.
- iridescens**, *id.*, loc. cit., 333, ♂ ♀ (1913).
 Nongkodjadar, Java, i [Jacobson]. *Types* in Amsterdam Museum.

Family CHIRONOMIDAE.¹

In the Chironomidae, a family with which I am but poorly acquainted, it has been my endeavour to steer an intermediate course between the broader view as instanced by Dr. Kertész (Catalogue, 1902), and the narrower one followed by Dr. Kieffer. The former sinks in *Ceratopogon* many groups (*Culicoides*, *Palpomyia*, *Sphaeromyias*, etc.), that are regarded by the latter as generic, some of which have yet again been subdivided into further genera. The exact amount of pubescence on the wing, the degree of incrassation of the femora, also the presence thereon or absence of small spines or spinules, are all characters liable to vary much in degree, so that drawing a hard and fast generic line between the groups becomes increasingly difficult as a greater number of species are discovered.

In admitting *Sphaeromyias*, *Serromyia*, *Heteromyia* and *Bezzia* to generic rank I follow Coquillett (1910), but though that author admits *Culicoides* as generically distinct from *Ceratopogon*, I have been unable to sufficiently differentiate them to do more than list the species of the different groups herein regarded as subgeneric, according to the views of their authors.

Professor Kieffer, in describing many of his new species, has

¹ A number of Indian Museum *types* in this family are still with Prof. Kieffer, but communication with Germany being interrupted by the present war, these *types* are entered in this catalogue as "belonging to the Indian Museum."

not specified any individual as actual *type*, so that in cases where the species has been described from several specimens these are herein designated as paratypes, each being regarded as of equal value.

Subfamily CERATOPOGONINAE.¹

Calyptopogon, Kieff.

Mem. Ind. Mus. II, 209 (1910).

GENOTYPE, *C. albitarsis* Kieff., sp. nov.

*gibbosa*² Wied., Anal. entom., 10 (*Macropeza*) (1824).

Macropeza gibbosa, id., Auss. Zweifl. I, 20: Macq., Hist. Nat. Dipt. I, 69: de Meij., Tijd. v. Ent. L, 216; LVI, 337.

Macropeza javanensis, Kieff., Mem. Ind. Mus. II, 210, *nom. nov.* for *gibbosa* de Meij. (1910).

? *Macropeza javanensis*, Edwards, Ann. Mag. Nat. Hist. (8), XII, 202.

"Ostindien" [*Wied.*], Semarang, i [*Jacobson*]; Peradeniya, ii-1912. *Type* in Copenhagen Museum.

¹ For characters and table of genera *v.* Kieff., Gen. Ins., Fasc. 42, p. 44; also (for table of genera), id., Rec. Ind. Mus. IX, 161 (Culicoidinae).

² The synonymy of these two species is somewhat confused, owing to erroneous conceptions. Wiedemann described *gibbosa* as entirely black with the front part of the thorax projecting strongly over the head. He then says, "Vorderste Beine" yellow with black tips; "hintere" lengthened, whitish, etc. Did Wiedemann by "beine" mean *legs or feet* (i.e. *tarsi*)? "Legs" would be correct, and consistent with his general use of the term. De Meijere adopts this view in his redescription (Tijd. v. Ent. L, 216), giving the legs as reddish-yellow, with knees and extreme tips of tibiae, also last joint of all the tarsi, black.

In the common Indian species, of which the Indian Museum has a dozen specimens, and which I had always regarded (prior to Kieffer setting up *albitarsis*) as *gibbosa* (interpreting "beine" for once as "feet" [*tarsi*] and not "legs"), the femora and tarsi are black or dark brown—irrespective of a little paleness at the bases of one or more pairs in individual specimens; the front tarsi are normally yellow, but may occasionally vary from light brown to nearly white, with black tips; the posterior tarsi are white or whitish, the hind pair generally the whitest, all with narrowly black tips. This species has the conspicuous, conically produced shining black thorax and wholly black body attributable to *gibbosa*.

The erection of *Calyptopogon* and *albitarsis* first caused me to doubt this identification of *gibbosa*, as Kieffer described as *albitarsis* the common Indian species I had regarded as *gibbosa*. He mistranslates the older author's species thus: "anterior and middle feet yellow with black tips; posterior feet whitish." He wrongly employs the terms "anterior" and "posterior" for "fore" and "hind"; because whether Wiedemann meant legs or tarsi in this particular instance, he certainly by "vorderste" alluded to the front pair only. The colour of the middle tarsi may well vary from yellow to white. He concludes that Meijere's identification of *gibbosa* from Java is incorrect assuming that as he (Meijere) retained the species in *Macropeza*, the thorax could not be conically produced, but must be as in normal species of that genus. He renamed Meijere's species "*javanensis*." As I pointed out elsewhere (Rec. Ind. Mus. VIII, 150) Meijere could not have failed to note Wiedemann's very distinct description of the peculiarly shaped thorax, so that, whatever his species might be, Meijere's "*gibbosa*, Wied." must at least be a *Calyptopogon*.

Mr. Edwards notes (Ann. Mag. Nat. Hist. [8], XII, 202, 1913) a specimen each of two species from Ceylon: (1), "*javanensis* Kieff. (*gibbosa* Meij. *nec* W.)," with dull brown thorax only slightly and bluntly produced forwards, and with yellow femora and tibiae; and, (2), "*gibbosa*, W. (*albitarsis*, Kieff.)," with shining black, sharply and conically produced thorax, and black femora and tibiae.

albitarsis Kieff., Mem. Ind. Mus. II, 209, ♀, pl. viii, 2, hind claws, 6, full ins. ♀ (1910).

Kieff., Rec. Ind. Mus. VI, 124 ♂; IX, 165, note.

Brun., *loc. cit.*, VIII, 150 (note on synonymy).

Macropeza gibbosa, Edwards, Ann. Mag. Nat. Hist. (8), XII, 202.

Calcutta, II-viii, 27-viii, 25-ix, 3-x, 25-x, 11-18-xi-07; Katihar, Bengal; Madhupur, Bengal, x; Monghyr, Bengal, 22-ix [Jenkins]; Damukdia Ghat, Ganges Riv., E. Bengal, at light, 30-vi; Lake Chilka, 5-iii; Puri, at light; Dibrugarh, Assam, 17-19-xi-11 [Kemp]; Peradeniya, x-1911, ii-1912. *Paratypes* in Indian Museum.

Jenkinsia, Kieff.

Rec. Ind. Mus. IX, 165 (1913).

GENOTYPE, *J. setosipennis* Kieff., n. sp., by original designation.

setosipennis Kieff., *loc. cit.*, 165, ♀ (1913).

Calcutta, 17-ix-09 [Paiva]; Bhogaon [Paiva]. Two *paratypes* in Indian Museum.

Dasyhelina, Kieff.¹

Rec. Ind. Mus. IX, 179 (1913).

GENOTYPE, *D. halophila*, Kieff., by original designation.

His view of *gibbosa* is the same as my first impression, that it possesses black femora and tibiae.

More recently, Meijere (Tijd. v. Ent. LVI, 337) considers that both the species accepted by Edwards are distinct from the true *gibbosa*, W., and his other remarks are contingent on the view of *gibbosa* being a yellow-legged species. He notes that Kieffer's *albitarsis* must be different from *gibbosa*, as the former has wholly white tarsi; but though Kieffer actually states this, it is an error, as the paratypes returned by him to the Indian Museum all possess distinctly black tips to all the tarsi.

It is clear, therefore, that two species exist, one with yellow femora and tibiae represented by Meijere's Javan and Edwards' Ceylonese specimen (unless the brown blunt thorax in the latter should make it distinct, which I doubt, thinking it only a question of personal use of terms or perhaps of actual variation); and one with black femora and tibiae, that is Kieffer's *albitarsis*.

Therefore, if *gibbosa* has yellow legs the synonymy will be as given in this catalogue, if it has black legs and is identical with Kieffer's *albitarsis*, the synonymy will be as follows:—

1. *gibbosa*, W. (*Macropeza*).
Calyptopogon albitarsis, Kieff., Mem. Ind. Mus. II: Brun., Rec. Ind. Mus. VIII, 150.
Macropeza gibbosa, Edwards, Ann. Mag. Nat. Hist. (8), XII, 202.
2. *species sine nom.* (with yellow femora and tibiae).
Macropeza gibbosa, de Meij., Tijd. v. Ent. I, 216.
Macropeza javanensis, Edwards, Ann. Mag. Nat. Hist. (8), XII, 202.

Only a reference to the type of *gibbosa* can decide the exact identity of the species.

¹ In Dr. Kieffer's table of genera (Rec. Ind. Mus. IX, 161), *Dasyhelina* falls in two of his principal divisions; his subgenus *Kempia* in that of "empodium well developed"; the subgenus *Prokempia* in that of "empodium atrophied." All the species seem characterised by pubescent eyes.

ornaticornis Kieff., Rec. Ind. Mus. IX, 180, ♀ (subgen. *Prokempia*) (1913).

Puri, 13-ix [Kemp]. Three *paratypes* in alcohol in Ind. Mus.

longicornis, *id.*, *loc. cit.*, 180, ♀ (subgen. *Prokempia*) (1913).

Katihar, Bengal, 24-ix [Paiva]. Several *paratypes* in alcohol in Indian Museum.

minima, *id.*, *loc. cit.*, 181, ♂ (subgen. *Prokempia*) (1913).

Calcutta, 29-i-10 [Annandale]. Two *paratypes* in Ind. Mus.

calcuttensis, *id.*, *loc. cit.*, 181, ♀ (subgen. *Kempia*) (1913).

Calcutta, 20-viii-07 [Annandale]. *Type* in Indian Museum, much damaged, unique.

perplexa, *id.*, *loc. cit.*, 182, ♀ (subgen. *Kempia*) (1913).

Calcutta, 20-viii-07. *Type* in Indian Museum, unique.

Ceratopogon, Mg.

Illig. Mag. II, 261 (1803).

Sch., F. Aust. II, 574: Kieff., Gen. Ins., Fasc. 42, 48: *id.*, Mem. Ind. Mus. II, 182, tab. Ind. spp.: *id.*, Rec. Ind. Mus. VI, 321, tab. spp. nov.: GENOTYPE, *C. communis* Mg., as *Tipula barbicornis*, F.

Forcipomyia, Mg., Syst. Besch. 1st Ed. I, 73 (1818):

de Meij., Tijd. v. Ent. L, 208, tab. sp. nov.

GENOTYPE, *Ceratopogon ambiguus* Mg., by Coquillett's designation.

Apogon, Rond., Dipt. Ital. Prod. I, 175 (1856).

GENOTYPE, *Ceratopogon flavipes* Mg., as *hortulanus* Mg., by original designation.

Labidomyia, Steph., Cat. Brit. Ins. II, 239 (1829).

GENOTYPE, *Tipula bipunctata* L., by Westwood's designation.

Ceratopogon, sensu lato.

trichopus Thoms., Eugen. Resa, 444 (1869).¹

China. *Type*, location unknown to present author.

agas Rond., Ann. Mus. Gen. VII, 462 (1865).

Borneo. *Type* (presumably) in Genoa Museum.

lignicola Brun., Rec. Ind. Mus. VII, 445, ♂ (1912).

Thano, Dehra Dun, W. Himalayas, 1500 ft., 9-ii-12 [Inms].

Type in Indian Museum; *co-types* in Forest Zoology coll., Dehra Dun.

Sub-genus *Ceratopogon*, sensu stricto.

hirtipes de Meij., Tijd. v. Ent. L, 209, ♂ ♀ (1907).

Semarang [Jacobson]; Tosari [Kobus], both Java; Peradeniya, 17-xi-11; i-12. *Types* in Amsterdam Museum.

¹ This species is retained here because it figured in Van der Wulp's Catalogue, but it may more properly belong to the Palaearctic Fauna.

jacobsoni, *id.*, *loc. cit.*, L, 212, ♂ ♀; pl. v, 8, 9, wing, ♂ ♀ (1907).
Semarang, i [Jacobson]; Peradeniya i-12 "attracted by
alcohol." *Types* in Amsterdam Museum.

stimulans, *id.*, *loc. cit.*, LII, 197, ♀, pl. xii, 12 (1909): *id.*, *loc. cit.*,
LVI, 334.

Sumatra [Sahn]; Semarang, vi [Jacobson]. *Type* in Am-
sterdam Museum.

salmi, *id.*, *loc. cit.*, 197, ♂ ♀, pl. xii, 3-6 (1909).

Sumatra [Sahn]. *Type* in Amsterdam Museum.

auronitens Kieff., Mem. Ind. Mus., II, 183, ♂, pl. xi, 12, wing
(1910).

Base of Dawna Hills, 2-iii-08 [Annandale]. *Unique type*
in Indian Museum.

macrorhynchus, *id.*, *loc. cit.*, 183, ♂, pl. ix, 3, mouth and palpi,
8, antenna, pt. (1910).

Calcutta, vii-07 [Annandale]. *Type* belonging to Indian
Museum.

albonotatus, *id.*, *loc. cit.*, 185, ♂, pl. viii, 10, wing, (1910).

Calcutta, 3-vi-08 [Annandale]. *Unique type* in Indian
Museum.

Subgenus *Atrichopogon*, Kieff.

Gen. Ins., Fasc. 42, 53 (1906).

Kieff., Rec. Ind. Mus. IX, 174, notes, tab. Ind.
spp.

GENOTYPE, *Ceratopogon exilis* Coq., by designa-
tion of Coquillett, 1910.

indianus Kieff., Mem. Ind. Mus. II, 185, ♂ (1910).

Calcutta Zoolog. Gardens, viii [Annandale]. *Unique type*
in Indian Museum.

rivicola Kieff., Rec. Ind. Mus. VI, 324, ♀ (1911).

Barogh, Simla, 10-vi-10 [Annandale]; Darjiling, 24-ix
[Brunetti]. Three *paratypes* in Indian Museum.

flaviceps Kieff., Supp. Ent. Berl. I, 28 (1912).

Formosa.

xanthopus, *id.*, Rec. Ind. Mus. IX, 175, ♀ (1913).

Lake Chilka, 4-iii-10 [Annandale]. *Unique type* in Indian
Museum.

flavipalpis, *id.*, *loc. cit.*, 175, ♂ ♀ (1913).

Rajmahal, Bengal, 5-vii-09, *type* ♂ [Annandale]; Adra,
Manbhurn Dist., Chota Nagpur, 12-x-09, *type* ♀ [Jenkins].
Unique types in Indian Museum.

origenus, *id.*, *loc. cit.*, 176, ♀ (1913).

Kurseong, 9-ix-09 [Annandale]. *Unique type* in Indian
Museum.

- sublimatus**, *id.*, *loc. cit.*, 176, ♀ (1913).
Kurseong, 6-ix-09 [Annandale]. Unique type in Indian Museum.
- flavellus**, *id.*, *loc. cit.*, 177, ♂ ♀ (1913).
Maddathorai, Travancore, 18-xi; Bangalore, S. India, 3000 ft. [Annandale]; Dawna Hills, 1-iii-08 [Annandale]. Three paratypes in Indian Museum.
- spurius**, *id.*, *loc. cit.*, 177, ♀ (1913).
Madhupur, Bengal, 15-x-09 [Paiva]. Unique type in Indian Museum.
- aterrimus**, *id.*, *loc. cit.*, 178, ♀ (1913).
Darjiling, 8-viii-09 [Jenkins]. Unique type in Indian Museum.
- setosipennis**, *id.*, *loc. cit.*, 178, ♀ (1913).
Madhupur, Bengal, 16-x-09 [Paiva]. Unique type in Indian Museum.
- brevistilus**, *id.*, *loc. cit.*, 178, ♀ (1913).
Kurseong, vii [D'Abreu]. Unique type in alcohol in Indian Museum.

Subgenus **Prohelia**, Kieff.

Rec. Ind. Mus. VI, 319 (1911).¹

GENOTYPE, *Ceratopogon decipiens* Kieff., by present designation.

? *Oecacta* Poey., Mem. Hist. Nat. Cuba, I, 236 (1851).²

- decipiens** Kieff., Mem. Ind. Mus. II, 182, ♂, pl. xi, 10, wing (*Ceratopogon*) (1910).
Id., Rec. Ind. Mus. VI, 321 (*Prohelia*).
Simla, v [Annandale]. Unique type in Indian Museum.
- cubicularis** Kieff., Rec. Ind. Mus. VI, 320, ♂ (1911).
Kurseong, in bedroom, 23-vi-10 [Annandale]. Two paratypes in Indian Museum.

¹ A little confusion may occur here. In Rec. Ind. Mus. VI, 319 (December 1911), the genus is characterised, but no species is quoted as type. The first species mentioned is "*tetraclada* Kieff." (not described as new), and this is followed by two new species and by *decipiens* Kieff. described in 1910. Now in the *Spolia Zeylanica* VIII, 1, published January 1912, *Prohelea* is noted as a new subgenus, and *tetraclada* (described for the first time) is the second of five species described as new. On the face of it, it would appear as if the paper in the *Spolia Zeylanica* had been written first, but any way, that in the Rec. Ind. Mus. was undoubtedly published first, and therefore the correct original reference to the genus must be that of that journal. Following on this reasoning, the type must be either *decipiens* (a previously described species) or one of the two described as new (*cubicularis* and *himalayae*). I propose *decipiens* as genotype on account of being the first described of the species now referred to *Prohelea*. Of course for the two new species described in the *Spolia* paper the original references will be that journal.

² Coquillett gives this latter genus as synonymous with *Culicoides* Latr., with *O. furens* Poey, as genotype.

himalayae, *id.*, *loc. cit.*, 320, ♂ (1911).

Kurseong, 22-vi-10, in bedroom [*Annandale*]. Unique *type* in Indian Museum.

ornaticrus Kieff., Spol. Zeyl. VIII, 1, ♀ (1912).

Peradeniya, 9 and 11-viii-10. Two *paratypes* in Indian Museum.

calcarata Kieff., Spol. Zeyl. VIII, 3, ♀ (1912).

Peradeniya, 2-vi-10. Unique *type* in Indian Museum.

noctivaga, *id.*, *loc. cit.*, 3, ♀ (1912).

Peradeniya, 19-ix-10, at light. Two *paratypes* in Indian Museum.

theobromae, *id.*, *loc. cit.*, 3, ♀, fig. 1, antenna, pt. (1912).

Peradeniya, 6-vi-10, bred from larvae in decomposed fruit of *Theobroma cacao* L. [*Gravely*].

tetraclada, *id.*, *loc. cit.*, 2, ♀ (1912).

Id., Rec. Ind. Mus. VI, 320 (1911).¹

Peradeniya, 23-v-10, *type*; Kurseong, 23-vi-10 [*Annandale*]. *Type* in Indian Museum.

Subgenus *Forcipomyia*, Meg. in Mg.²

Syst. Besch., 1st Ed., I, 73 (1818).

Kieff., Gen. Ins., Fasc. 42, 52: *id.*, Rec. Ind.

Mus. VI, 321 and IX, 166, tab. Ind. spp.

GENOTYPE, *Ceratopogon ambiguus* Mg., by Coquillett's designation, 1910.

vexans de Meij., Tijds. v. Ent. LII, 198, ♂ ♀, pl. xii, 7-10 (1909).

Sumatra [*Dr. Salm*]. *Type* in Amsterdam Museum.

There is a *Ceratopogon vexans*, Staeg, 1839.

albosignata Kieff., Mem. Ind. Mus. II, 184, ♂ (*Ceratopogon*) (1910).

Rangoon, 24-ii-08, *type* [*Annandale*]. *Type* (pinned) in Indian Museum, also *coltypes* in alcohol.

var. *asticta* Kieff., Rec. Ind. Mus. IX, 172, ♀ (1913).

Kurseong, vii [*D'Abreu*]. *Type* in Indian Museum.

var. *variicrus*, *id.*, *loc. cit.*, 172, ♂ (1913).

Port Canning, 5-ii [*Annandale*]. Unique *type* in Indian Museum.

calotricha Kieff., Rec. Ind. Mus. VI, 322, ♀; pl. xiv, 1 wing, pt. (1911).

Kurseong, 22-vi-10, in bedroom [*Annandale*]. Unique *type* in Indian Museum.

¹ When mentioned in this periodical *tetraclada* was a *nomen nudum*: see footnote 1 on previous page.

² De Meijere regards *Forcipomyia* as a subgenus of *Ceratopogon*; Coquillett sinks it in *Culicoides*.

- montivaga**, *id.*, *loc. cit.*, 322, ♀ (1911).
Darjiling, 28-v-10 [Brunetti]. Unique type in Indian Museum.
- alticola**, *id.*, *loc. cit.*, 323, ♀ (1911).
Darjiling, 8-v-10 [Annandale]. Unique type in Indian Museum.
- oriphila**, *id.*, *loc. cit.*, 324, ♀ (1911).
Darjiling, 29-v-10 [Brunetti]. Type in Indian Museum.
- macrothrix**, *id.*, *loc. cit.*, 325, ♀ (1911).
Kurseong, 20-vi-10 [Annandale]. Type in Indian Museum.
- var. **flaviceps**, *id.*, *loc. cit.*, IX, 173, ♀ (1913).
Kurseong, vii [D'Abreu].
Kieffer also describes, (*l. c.* 172) a further, unnamed variety from Kurseong, 4-vii [Annandale]: two specimens being in the Indian Museum.
- urbicola** Kieff., Rec. Ind. Mus. VI, 325, ♂ (1911).
Kurseong, 22-vi-10 [Annandale]. Type (pinned) in Indian Museum, also *cotypes* in alcohol.
- xanthophila**, *id.*, *loc. cit.*, 326, ♀ (1911).
Kurseong, 13—16-viii-07 [Annandale]. Unique type in Indian Museum.
- longicalcar** Kieff., Spol. Zeyl. VIII, 4, ♀, fig. 2, wing (1912).
Peradeniya, 20-v-10. Unique type in Indian Museum.
- semipilosus** Kieff., *loc. cit.*, 5, ♀ (1912).
Peradeniya, 4-vii-10. Unique type in Indian Museum.
- sauteri** Kieff., Supp. Ent. Berl. I, 27 (1912).
Formosa.
- conifera** Kieff., Rec. Ind. Mus. IX, 168, ♂ (1913).
Giridih, Bengal, 7-xi [Kemp]. Unique type in alcohol in Indian Museum.
- trinotata**, *id.*, *loc. cit.*, 169, ♂ ♀ (1913).
Naini Tal [Lloyd]; Simla, 9-v [Annandale]; Kurseong, 4-vii; 5-ix [Annandale]. Three *paratypes* in Indian Museum.
- tristicta**, *id.*, *loc. cit.*, 169, ♂ (1913).
Kurseong, 6-iii [Annandale]. Type belonging to Indian Museum.
- psychasta**, *id.*, *loc. cit.*, 170, ♂ ♀ (1913).
Kurseong, 4-vii-08, type ♂ ♀ [Annandale]. Unique types in Indian Museum.
- confluens**, *id.*, *loc. cit.*, 170, ♀ (1913).
Kurseong, vii [D'Abreu]. Two *paratypes* in alcohol in Indian Museum.
- monticola**, *id.*, *loc. cit.*, 171, ♂ (1913).
Kurseong, 4-vii-08 [Annandale]. Type in Indian Museum.
- heterocera**, *id.*, *loc. cit.*, 173, ♀ (1913).
Darjiling, 8-viii-09 [Paiva]. Unique type in Indian Museum.

Subgenus *Culicoides*, Latr.

Gen. Crust. et Ins. IV, 251 (1809).

Kieff., Gen. Ins., Fasc. 42, 53: *id.*, Mem. Ind. Mus. II, 186, tab. Ind. spp.: Annand., Rec. Ind. Mus. IX, 246, notes on four Indian blood-sucking species.GENOTYPE, *Culex pulicarius* L. (as *punctata*, sp. nov.) by original designation.

- guttifer** de Meij., Tijds. v. Ent. L, 209, pl. v, 6, wing (1907).
Semarang, 1 [Jacobson]. Unique *type* in Amsterdam Museum.
- pungens**, *id.*, *loc. cit.*, LII, 200, pl. xii, 11, 12 (1909).
Sumatra. *Type* in Amsterdam Museum.
- brevimanus** Kieff., Mem. Ind. Mus. II, 187, ♀ (1910).
Calcutta, 20-viii-07 [Annandale]. Unique *type* in Indian Museum.
- macrostoma**, *id.*, *loc. cit.*, 188, ♂ ♀, pl. viii, 4, wing; pl. vi, 14, mouth parts (1910).
Calcutta, ii-08 [Annandale]. Unique *type* now completely destroyed.
- montivagus**, *id.*, *loc. cit.*, 188, ♂, pl. viii, 3, wing (1910).
Simla, 11-v-08 [Annandale]. Two *paratypes* in Indian Museum.
- opacus**, *id.*, *loc. cit.*, 189, ♂ ♀, pl. xi, 3, antenna ♀ (1910).
Calcutta, 6-7-ii-08 [Annandale]. Three *paratypes* in Indian Museum.
- scapularis**, *id.*, *loc. cit.*, 189, ♂ ♀ (1910).
Calcutta, var. dates in January [Annandale]. Several *paratypes* in Indian Museum.
- nitidulus**, *id.*, *loc. cit.*, 189, ♀ (1910).
Calcutta, ii, vi, vii [Annandale]; base of Dawna Hills, iii. Several *paratypes* in Indian Museum.
- quadrilobatus**, *id.*, *loc. cit.*, 189, ♂ ♀, pl. ix, 2, 4, 9, 12, pl. x, 8, 10, var. chars. (1910).
Calcutta Zoolog. Gardens, 25-viii-07 [Annandale]. *Types* in Indian Museum in Canada balsam.
- setiger**, *id.*, *loc. cit.*, 190, ♂ ♀, pl. x, 1, 2, 3, var. chars. (1910).
Calcutta, 16-vii-07. *Types* belonging to Indian Museum.
- peregrinus**, *id.*, *loc. cit.*, 191, ♀, pl. viii, 1, wing (1910).¹
Puri, 24-ii-08, 2-iii-08 [Paiva]. Three *paratypes* in Indian Museum.

¹ See Annandale, Rec. Ind. Mus. IX, 246, for blood-sucking notes.

- molestus*, *id.*, *loc. cit.*, 192, ♂ ♀, pl. viii, 9, wing (1910).
Id., Rec. Ind. Mus. IX, 183, ♀, notes.
 Calcutta Zoolog. Gardens, 24-viii-07 [*Annandale*]; Kurseong,
 vii [*D'Abreu*]. *Type* in alcohol in Indian Museum.
- paivai* Kieff., Mem. Ind. Mus. II, 193, ♀, pl. viii, 5 (1910).
 Puri, Orissa, 2-iii-08 [*Paiva*]. *Unique type* in Indian Museum.
- oxystoma*, *id.*, *loc. cit.*, 193, ♀, pl. ix, 1, mouth, palpi (1910).
 Calcutta Zoolog. Gardens, 25-viii-07 "attacks oxen and deer"
 [*Annandale*]. *Type* in Indian Museum in Canada balsam.
- himalayae* Kieff., Rec. Ind. Mus. VI, 326, ♂ ♀ (1911).
 Kurseong, 22-vi-10 [*Annandale*]. Three *paratypes* in Indian Museum.
- pictiventris*, *id.*, *loc. cit.*, 327, ♀ (1911).
 Tonglu, Darjiling District, 22-iv-10 [*Beebe*]. *Unique type* in Indian Museum.
- formosae* Kieff., Supp. Ent. Berl. I, 29 (1912).
 Formosa.
- indecora*, *id.*, *loc. cit.*, 29 (1912).
 Formosa.
- ceylanicus* Kieff., Spol. Zeyl. VII, 5, ♀ (1912).
 Peradeniya, 25-vi-10. *Unique type* in Indian Museum.
- bilobatus*, *id.*, *loc. cit.*, 6, ♂ (1912).
 Peradeniya, 14-v-10. *Unique type* in Indian Museum.
- xanthocoma* Kieff., Rec. Ind. Mus. IX, 182 ♂ (1913).
 Puri, Orissa, 13-xi [*Kemp*]. *Unique type* in alcohol in Indian Museum.
- kiefferi* Patton, Ind. Jour. Med. Res. I, 336, ♂ ♀, pl. xviii, var. chars. (1913).
 Madras, Nov. to March; Coimbatore [*Fletcher*].

Palpomyia, Meg. in Mg.

- Syst. Besch. 1st Ed. I, 82 (1818).
 Kieff., Gen. Ins., Facs. 42, 61: *id.*, Mem. Ind. Mus. II, 194 and Rec. Ind. Mus. VI, 113; IX, 187, tab. Ind. spp.
 GENOTYPE, *Ceratopogon flavipes* Mg.
- crassinervis* de Meij., Tijd. v. Ent. L, 213, pl. v, 10, wing (1907).
 Semarang, 1 [*Jacobson*]. *Type* in Amsterdam Museum.
- discolor*, *id.*, *loc. cit.*, 214 (1907).
 Semarang, 1 [*Jacobson*]. *Type* in Amsterdam Museum.
- luteinervis*, *id.*, *loc. cit.* 214, pl. v, 11, wing (1907).
 Semarang, 1 [*Jacobson*]. *Type* in Amsterdam Museum.

- burmae** Kieff., Mem. Ind. Mus. II, 196, ♀ (1910).
Base of Dawna Hills, 1-iii-08 [Annandale]. Type in Indian Museum.
- filicornis**, *id.*, *loc. cit.*, 196, ♀; pl. x, 4-7, 9, var. chars. (1910).
Calcutta, 16-vii-07 [Annandale]. Type belonging to Indian Museum.
- cinerea** *id.*, *loc. cit.*, 197, ♀ (1910).
Rajmahal, Bengal, 3-vii-07. Unique type in Indian Museum.
- pulchripes**, *id.*, *loc. cit.*, 198, ♀, pl. xi, 8, wing (1910).
Calcutta, 13-viii-07; Purneah, 6-viii-07. Paratypes in Indian Museum.
- pictipes**, *id.*, *loc. cit.*, 199, ♀ (1910).
Calcutta, 15-viii-07; 13-x [Annandale]; Katihar, 4-viii-07.
Two paratypes in Indian Museum.
- interrupta**, *id.*, *loc. cit.*, 199, ♀, pl. xi, 16, front tarsi claws (1910).
Calcutta, 10-27-vii-07; 27-viii-07. Several paratypes in Indian Museum.
- inaequalis**, *id.*, *loc. cit.*, 200, ♀ (1910).
Purneah, 5-viii-07. Unique type in Indian Museum.
- roseiventris** Kieff., Rec. Ind. Mus. VI, 114, ♀ (1911).
Calcutta, 28-viii-07. Unique type in Indian Museum.
- leucopogon**, *id.*, *loc. cit.*, 114, ♂ (1911).
Calcutta, 1-x-07. Unique type in Indian Museum.
- disticta**, *id.*, *loc. cit.*, 115, ♀, pl. vi, 1, wing (1911).
Calcutta, 3-ix, at light [Paiva]; 27-viii, 5-ix, 31-x-07; Dum Dum, near Calcutta, 28-vii [Lord]; Madhupur, Bengal, x, at light; Khustea, Bengal, x [Jenkins]; Jalpaiguri, N. Bengal, ix, at light; Bhogaon, Purneah, 2-x [Paiva]; Chennia, Sunderbunds, Bengal, xi, at light on board launch [Jenkins]; Allahabad, viii [Lord]. Several paratypes in Indian Museum.
- tetrasticta**, *id.*, *loc. cit.*, 116, ♀ (1911).
Calcutta, 7-ix-07. Unique type in Indian Museum.
- polysticta**, *id.*, *loc. cit.*, 117, ♀ (1911).
Calcutta, 7-ix-07. Unique type in Indian Museum.
- brevispina**, *id.*, *loc. cit.*, 117, ♀ (1911).
Calcutta, 27-vii-07. Unique type in Indian Museum.
- rivularis**, *id.*, *loc. cit.*, 328, ♀ (1911).
Valley of Sutlej Riv. below Simla, 6-v-10 [Annandale].
Unique type in Indian Museum.
- taitana** Kieff., Supp. Ent. Berl. I, 29 (1912).
Formosa.
- albinervis** Kieff., Rec. Ind. Mus. IX, 187, ♀ (1913).
Pallode, Travancore, 15-xi [Annandale]. Type in Indian Museum.

- photophila**, *id.*, *loc. cit.*, 188, ♀ (1913).
Madhupur, Bengal, at light, x [*Paiva*]; Calcutta, at light, vi.
Type in Indian Museum.
- abjuncta**, *id.*, *loc. cit.*, 189, ♀ (1913).
Calcutta, 14-ix, at light [*Paiva*]. *Type* belonging to Indian
Museum.
- disjuncta**, *id.*, *loc. cit.*, 189, ♀ (1913).
Bhogaon, N. Bengal, x [*Paiva*]. *Type* belonging to Indian
Museum.

Sphaeromyias, Curt.

Brit. Entom., 285, pl. (1829).

Nec Sphaeromyias, Walk.

Xylocrypta Kieff., Bull. Ent. Soc. Fr., 69 (1899).

GEONOTYPE of *Sphaeromyias* Curt., *Ceralopogon fasciatus*, Mg., as *albomarginatus*, sp. nov.

Kieffer's type of *Xylocrypta* is the same species.

- bisignata** Kieff., Mem. Ind. Mus. II, 200, ♀, pl. xi, 1, tarsus,
part (1910).
Purneah, 5-viii [*Paiva*]. *Type* belonging to Indian Mu-
seum.
- bimacula**, *id.*, *loc. cit.*, 201, ♂, pl. xi, 5, tarsus, 7, wing (1910).
Calcutta, 14-viii; Purneah, 5-viii [*Paiva*]. *Types* belonging
to Indian Museum.
- filitarsis**, *id.*, *loc. cit.*, 202, ♀, pl. xi, 9, claws (1910).
Calcutta, 3-13-viii-07. "Near *Macropeza*." *Paratypes* in
Indian Museum.
- albiditarsis**, *id.*, *loc. cit.*, 203, ♀ (1910).
Calcutta. *Unique type* in Indian Museum.
- viridiventris**, *id.*, *loc. cit.*, 203, ♀, pl. viii, 8, tarsus (1910).
Dawna Hills, 1000 ft., 2-3-iii-08 [*Annandale*]. *Type* in
Indian Museum, a unique specimen.
- stictonota** Kieff., Rec. Ind. Mus., VI, 118, ♂ (1911).
Calcutta (Zoolog. Gardens), 8-iii-01. *Type* belonging to
Indian Museum.
- inermicrus**, *id.*, *loc. cit.*, ix, 186, ♀ (1913).
Rajmahal, Bengal, 5-vii-09 [*Annandale*]. *Type*. a unique
specimen, in Indian Museum.

Stilobezzia, Kieff.

Rec. Ind. Mus. VI, 118 (1911).

Kieff., *loc. cit.*, IX, 184, notes.

GENOTYPE, *S. festiva* Kieff., by original designa-
tion.

- festiva** Kieff., Rec. Ind. Mus. VI, 118, ♂ ♀, pl. vi, 2-6, var.
chars. (1911).
Calcutta, 24-viii, 11-ix, 1-29-x. *Paratypes* in Ind. Mus.

var. *scutellaris* Kieff., Spol. Zeyl. VIII, 7, ♀ (1912).

Peradeniya, 24-v-10. Type in Indian Museum, a unique specimen.

inermipes Kieff., Spol. Zeyl. VIII, 8, ♀ (1911).

Peradeniya, 2-vi-10. Type in Indian Museum.

A second specimen from Ceylon (? Peradeniya) in same collection.

lineata Kieff., Rec. Ind. Mus. IX, 184, ♂ ♀ (1913).

Madhupur, Bengal, 13-x; 18-x [*Paiva*]; Giridih, Bengal, 7-xi [*Kemp*]; Karmatar, Bengal, 23-x [*Paiva*]. Paratypes in Indian Museum.

Serromyia, Meg. in Mg.

Syst. Besch. 1st Ed. I, 66 (1918).

GENOTYPE, *Ceratopogon femorata* Mg., the original species.

Prionomyia, Steph., Cat. Brit. Ins. II, 237 (1829).

GENOTYPE, *Ceratopogon femoratus* Mg., by Westwood's designation.

Ceratolophus, Kieff., Bull. Soc. Ent. Fran. LXVIII,

69 (1899): *id.*, Gen. Ins., Fasc. 42, 59.

GENOTYPE, *Ceratopogon femoratus* Mg., by original designation.¹

Johannseniella, Will., Jour. N.Yk. Ent. Soc. XV, 1

(1907), *nom. nov.* for *Ceratolophus* Kieff., *preocc.*

Bocoust, 1873.

notatus de Meij., Tijds. v. Ent. L. 210, ♂ ♀, pl. v, 7, wing (*Ceratolophus*) (1907).

Semarang, 1 [*Jacobson*] Type in Amsterdam Museum.

Coquillett considers *Johannseniella* Will., as synonymous with *Serromyia* Mg. and I follow him.

Heteromyia, Say.

Americ. Entom. II, 79 (1825).

Kieff., Gen. Ins., Fasc. 42, 64: *id.*, Rec. Ind.

Mus. IX, 183.

GENOTYPE, *H. fasciatus* Say, by original designation.

Pachyleptus, Walk., Ins. Saund. Dipt. pt. 5, 426 (1856).

GENOTYPE, *P. fasciatus* sp. nov., *loc. cit.*

indica Kieff., Rec. Ind. Mus. IX, 183, ♀ (*Pachyleptus indicus*) (1913).

Maddathorai, Travancore, 17-xi-08 [*Annandale*]. Type in Indian Museum.

Haasiella, Kieff.

Rec. Ind. Mus. IX, 190 (1913).

GENOTYPE, *H. semiflava*, by original designation.

¹ Kieffer overlooked the fact of *femoratus* having been employed as the type of *Serromyia*.

semiflava Kieff., *loc. cit.*, 190, ♀ (1913).

Tinpahar, Bengal, 7-vii-09 [*Annandale*]. *Type*, a unique ♀, in Indian Museum.

Bezzia, Kieff.¹

Bull. Soc. Ent. Fr. LXVIII, 69 (1899).

Kieff., Gen. Ins., Fasc. 42, 57: *id.*, Mem. Ind.

Mus. II, 204 (tab. Ind. spp.) (1910).

GENOTYPE, *Ceratopogon crnatus* Mg., by Kieffer's designation.

Probezzia, Kieff., Gen. Ins., Fasc. 42, 57 (1906).

GENOTYPE, *Ceratopogon venusta* Mg., by Coquillett's designation.

guttipennis de Meij., Nova Guin. Results, 69 (1906).

Papua. *Type* in Amsterdam Museum.

acanthopus de Meij., Tijds. v. Ent. L, 215, ♀, pl. v, 12, wing (1907).

Semarang, i, ii, iii [*Jacobson*]. *Type* in Amsterdam Museum.

flaviventris Kieff., Mem. Ind. Mus. II, 204, ♀ (1910).

Noalpur (Nepal), 22-ii-08. Unique *type* in Indian Museum.

gracilipes, *id.*, *loc. cit.*, 205, ♀ (1910).

Purneah, 4-viii [*Paiva*]. Two *paratypes* in Indian Museum.

var. *lineola*, *id.*, *loc. cit.*, 205, ♀.

Furi, January. *Type* in Ind. Mus., a unique specimen.

var. *analisis* Kieff., Rec. Ind. Mus. IX, 196, ♀ (1913).

Balighai, Orissa, 24-x [*Annandale*]. *Type*, a unique specimen, in Indian Museum.

var. *bengalensis*, *id.*, *loc. cit.*, 196, ♀ (1913).

Madhupur, Bengal, 16-x [*Paiva*]. *Type*, a unique specimen, in Indian Museum.

var. *flavescens*, *id.*, *loc. cit.*, 196, ♀ (1913).

Madhupur, Bengal, 13-x [*Paiva*]. *Type* in Indian Museum.

fascialis, *id.*, Mem. Ind. Mus. II, 206, ♂, pl. viii, 11, wing (1910).

Calcutta (Zoolog. Gardens), 25-vii-07 [*Annandale*]. *Type* in Indian Museum, a unique specimen.

lacteipennis, *id.*, *loc. cit.*, 206, ♀ (1910).

Calcutta, 15-viii-07, *type*; 11-viii-07, 12—13-ii-08. *Type* in Indian Museum.

An unnamed variety from Lake Chilka also described in same work. In Indian Museum.

¹ De Meijere ranks *Bezzia* as a subgenus of *Ceratopogon*: Coquillett (1910) admits it as a good genus.

In Kieffer's last paper (Rec. Ind. Mus. IX) he places *gracilipes*, *nigricans*, and *eucera*, with a number of new species, in the subgenus *Probezzia*, which differs from *Bezzia* in a minor and possibly variable character. All the species are incorporated herein under *Bezzia*. Even in the Genera Insectorum Kieffer lists all the species alphabetically under *Bezzia* without noting to which of the two subgenera *Probezzia* and *Bezzia*, *s. str.* each belongs.

nigricans, *id.*, *loc. cit.*, 207, ♀ (1910).

Id., Rec. Ind. Mus. IX, 193, ♀, notes (*Probezzia*) (1913).

Noalpur, Nepal, 22-ii-08; Calcutta, 27-vii-07 [*Paiva*]; Madhupur, Bengal, 14-x-09 [*Paiva*]; Balighai, Orissa, 24-x-08, *type*; Jalpaiguri, 10-ix-09 [*Annandale*]; Shencottah, 25-x-08 and Quilon, 9-xi-08 [both Travancore and *Annandale*].

armatipes Kieff., Mem. Ind. Mus. II, 207, ♀ (1910).

Puri, January. The unique *type* is now completely destroyed.

rufiventris, *id.*, *loc. cit.*, 208, ♀ (1910).

Calcutta, 10-viii-07 [*Annandale*]. *Type*, a unique specimen, in Indian Museum.

trispinosa Kieff., Rec. Ind. Mus. VI, 123, ♀, pl. vi, 9, antenna (1911).

Calcutta, 31-vii-07. *Type* belonging to Indian Museum.

eucera, *id.*, *loc. cit.*, 123, ♀, pl. vi, 10, antenna (1911).

Id., *loc. cit.*, IX, 196, note.

Calcutta, 30-x-07; Trivandrum, Travancore, 14-xi-08 [*Annandale*]. *Type* technically in Indian Museum.

Described originally from the Calcutta specimen, that should be the *type*, but it is not to be found in the Indian Museum, though the Travancore specimen is there.

albipennis Kieff., *loc. cit.*, VI, 124, ♀ (*B. nigricans*, var. *albipennis*) (1911).

Bezzia (*Probezzia*) *albipennis*, Kieff., *loc. cit.*, IX, 195, ♀ (1913).

Calcutta, 10-ix-07, 26-xi; Sara Ghat, Ganges Riv., Bengal, 11-ix [*Annandale*]; Madhupur, Bengal, 16-x [*Paiva*]; Balighai, Orissa, 25-x [*Annandale*]. *Paratypes* in Indian Museum.

vaga, *id.*, Rec. Ind. Mus. IV, 329, ♀ (1911).

Noalpur, Nepal, 22-ii-08, *type*; Madhupur, Bengal, 18-x-09 [*Paiva*]; Igatpuri, Western Ghats, Bombay Presidency, 20-xi-09 [*Annandale*]. *Type* in Indian Museum.

indecora, *id.*, Spol. Zeyl. VIII, 8, ♂ (1912).

Peradeniya, 14-v-10. *Type*, unique, in Indian Museum.

allotropica Kieff., Rec. Ind. Mus. IX, 193, ♀ (*Probezzia*) (1913).

Frasergunge, 12-xi; Basanti, 16-xi [both Sunderbunds and *Jenkins*]; at light on board launch, Bologhatta, Khulna, Ganges Delta, 28-viii, at light on launch; Besondhur, Khulna, 29-viii [both *Jenkins*]. *Paratypes* in Indian Museum.

hamifera, *id.*, *loc. cit.*, 194, ♀ (*Probezzia*) (1913).

Madhupur, Bengal, 15-16-x-09, at light [*Paiva*]; Balighai, Orissa, 25-x-08 [*Annandale*]. *Paratypes* in Indian Museum.

var. *arcuatipes* Kieff., *loc. cit.*, 195, ♀ (*Probezzia*) (1913).

Noalpur, Nepal, 22-ii; Madhupur, Bengal, 14, 16 and 18-x, at light [*Paiva*]; Jalpaiguri, N. Bengal, at light in railway carriage, 10-ix [*Annandale*]; Sara Ghat, Ganges Riv., E. Bengal, 11-ix [*Annandale*]; Balighai, Orissa, 24-25-x; Shencottah, 25-x; Qui-

lon, 9-xi; Kulattupuzha, 19-xi [all three Travancore and *Annandale*]; Igatpuri, Bombay Presid., 20-xi [*Annandale*].

calcuttensis, *id.*, *loc. cit.*, 195, ♀ (*Probezzia*) (1913).

Calcutta, 13-ix-07 [*Annandale*]. *Type* in Indian Museum, unique.

kempi, *id.*, *loc. cit.*, 197, ♀ (1913).

Puri, 13-x [*Kemp*]. *Type*, a unique specimen in alcohol, in Indian Museum.

formosana Kieff., Supp. Ent. Berl. I, 30 (1912).

Formosa.

chrysolopha, *id.*, *loc. cit.*, 31 (1912).

Formosa.

Subgenus *Dibezzia*, Kieff.

Rec. Ind. Mus. VI, 120 (1911).

GENOTYPE, *D. clavata* Kieff., sp. nov., by original designation.

clavata Kieff., Rec. Ind. Mus. VI, 120, ♀, pl. vi, 7, 8, tarsal joints (1911).

Calcutta, 5-9-ix-07. *Paratypes* in Indian Museum.

Kieffer (*l.c.*) also describes a variety from Calcutta.

longistila, *id.*, *loc. cit.*, 121, ♀ (1911).

Calcutta, 29-x-07. *Type*, a unique specimen, in Indian Museum.

brevistila, *id.*, *loc. cit.*, 122, ♀ (1911).

Sylhet, 3-iv-05 [*Hall*]. *Type*, a unique specimen, in Indian Museum.

himalayae, *id.*, *loc. cit.*, VI, 328, ♀ (1911).

Kurseong, 26-vi-10. *Type*, a unique specimen, in Indian Museum.

ceylanica Kieff., Spol. Zeyl. VIII, 6, ♀ (1912).

Peradeniya, 9-vii-10. Two *paratypes* in Indian Museum.

pictipennis Kieff., Rec. Ind. Mus. IX, 191, ♀ (1913).

Shasthancottah, Travancore 8-xi-08 [*Annandale*]. *Type*, a unique specimen, in Indian Museum.

spinigera, *id.*, *loc. cit.* x, 313, ♀ (1914).

Purneah, 5-viii-07. *Type* in Indian Museum.

Leptoconops, Skuse.

Proc. Linn. Soc. N.S. Wales (2) iv, 288 (1890).

De Meij., Tijd. v. Ent. LVIII, 98.

GENOTYPE, *L. stygius*, *id.*, *loc. cit.*, sp. nov.

albiventris de Meij., Tijd. v. Ent. LVIII, 98, ♀, pl. i, 6 (1915).

North Papua.

Subfamily TANYPINAE.¹

Tanypus, Mg.

Illig. Mag. II. 261 (1803).

Sch., F. Austr. II, 615; Kieff., Gen. Ins., Fasc.

42, 40: *id.*, Mem. Ind. Mus. II, 220, tab. Ind. spp.GENOTYPE, *Tipula cincta* F., by Latreille's designation, 1810.

melanurus Dol., Nat. Tijd. Ned. Ind. X, 405, pl. iv, 1, full insect and wing (1856).

Ambarawa, Java, in houses during the rains. *Type* probably lost.

cyaneomaculatus, *id.*, *loc. cit.*, 406, pl. xi, 2, full insect (1856).

Ambarawa, Java, abundant in bedrooms. *Type* probably lost.

nigrocinctus, *id.*, *loc. cit.*, 406, pl. xi, 3, full insect (1856).

Java. *Type* probably lost.

manillensis Sch., Reise Novara, Dipt., 26 (1868).

Manila. *Type* in Vienna Museum.

signifer de Meij., Nova Guin. Results, 71 (1906).

Papua. *Type* in Amsterdam Museum.

macrocerus Kieff., Mem. Ind. Mus. II, 212, ♀ (*Isoplastus*) (1910).

Id., Rec. Ind. Mus. VI, 333, note (*Pelopia*); *id.*, *loc. cit.*, IX, 160 (*Pelopia*).

Thamaspur, Nepal, xi and 18—20-ii-08; Puri, Orissa, 18-i [Annandale]. *Type* in Indian Museum.

A specimen in the same collection from Puri, 18—19-i-08, is noted by Kieffer as a variety.

paivai Kieff., Mem. Ind. Mus. II, 213, ♂ (*Isoplastus*) (1910).

Id., Rec. Ind. Mus. IX, 159, ♂ ♀ (*Pelopia*) (1913).

Bengal, between Manihari and Manshahi, on railway track, 4-viii-07, *type* ♂ [*Paiva*]; Madhupur, Bengal, 14-x-09, *type* ♀ [*Paiva*]; Maniharighat, Ganges Riv., on sand bank, 17-ii [Chaudhuri]. *Types* in Indian Museum.

ornatipes Kieff., Mem. Ind. Mus. II, 214, ♀ (1910).

Id., Rec. Ind. Mus., IX, 160, ♂ ♀ (1913).

Balighai, Orissa, 25-x [Annandale]; Madhupur, Bengal, 14-x, at light [*Paiva*]; Berhampore Court, Bengal, 12-iv [Kemp]; Calcutta, 16-ix; Sonadigee, Ganges Delta, 6-xii, at light [Jenkins]; Ernakulam, Cochin State, 4-xi [Annandale]. *Paratypes* in Indian Museum. Three *paratypes* of an unnamed variety also in the Museum.

pulchripes Kieff., Mem. Ind. Mus. II, 214, ♂ ♀ (1910).

Calcutta, ii, vii, at light; xii [Annandale]; Sonarpur, Bengal, at light, 24-xii; Puri, i, iii. *Paratypes* in Indian Museum.

¹ For characters and tab. gen. v. Kieff., Gen. Ins., Fasc. 42, 33—34 (1906), and Rec. Ind. Mus. IX, 153, under the inadmissible name of *Pelopiinae*.

- violaceipennis** Kieff., *loc. cit.*, II, 222, ♀ (1910).
Calcutta, 2-ii-08, *type*, 20-iii-07 [Annandale]. *Type* in Indian Museum.
- albolineatus**, *id.*, *loc. cit.*, 224, ♂ (1910).
Mandalay, 12-iii-08 [Annandale]. *Unique type* in Indian Museum.
- oriplanus** Kieff., *Rec. Ind. Mus.* vi, 125, ♂ (*Isoplastus*) (1911).
Id., *loc. cit.*, IX, 161, ♀ (*Pelopiea*) (1913).
Simla, 25-iv-07, *type* ♀ [Annandale]. *Type* ♂ apparently lost, *type* ♀ in Indian Museum.
- saltatrix**, *id.*, *Rec. Ind. Mus.* VI, 330, ♂ (*Pelopiea*) (1911).
Simla, 9-v-10 "in numbers at dusk, males aerial dancing" [Annandale]. *Paratypes* in Indian Museum.
- brunettii**, *id.*, *loc. cit.*, 332, ♂ (*Pelopiea*) (1911).
Darjiling, 28-v-10 [Brunetti]. *Type*, a unique specimen, in Indian Museum.
- riparia**, *id.*, *loc. cit.*, 332, ♂ (*Pelopiea*) (1911).
Barogh (Simla), 10-v-10 [Annandale]. *Type* in Indian Museum, a unique specimen.
- himalayae**, *id.*, *loc. cit.*, 333, ♂ (*Pelopiea*) (1911).
Barogh (Simla), 10-v-10 [Annandale]. *Two paratypes* in Indian Museum.
- annulatipes** Kieff., *Spol. Zeyl.* VIII, 8, ♀ (*Pelopiea*) (1912).
Peradeniya, 17-v-10. *Two paratypes* in Indian Museum.
- pallidipes**, *id.*, *loc. cit.*, 9, ♀ (*Pelopiea*) (1912).
Peradeniya, 29-v-10. *Type* in Indian Museum, a unique specimen.
- birmanensis** Kieff., *Rec. Ind. Mus.* IX, 159, ♂ (1913).
Mandalay, 11-iii-08 [Annandale]. *Type*, a unique specimen, in Indian Museum, reduced to a fragment.
- macrochaeta**, *id.*, *loc. cit.*, 159, ♀, pl. xii, 25, antenna ♀ (1913).
Kurseong, 27-iii [Annandale]. *Type* belonging to Indian Museum.

Clinotanypus, Kieff.

Rec. Ind. Mus. IX, 157 (1913).

GENOTYPE, *Procladius atratus* Kieff., the first species, by present designation.¹

- ornatissimus** Kieff., *Mem. Ind. Mus.* II., 216, ♂ (*Procladius*) (1910).
Id., *Rec. Ind. Mus.* IX, 158, note (*Clinotanypus*) (1913).
Calcutta, 21-vii, 19-viii [Annandale]; Kushtea, Bengal, 7-x [Jenkins]; Purnea, vii [Paiva]; Madhupur, Bengal, 16, 20-x,

¹ Kieffer in instituting this genus refers six of his species of *Procladius* to it without nominating any one as type.

at light [*Paiva*]; Nedumangad, Travancore, 14-xi [*Annandale*]; Victoria Gardens, Colombo, 26-iv [*Paiva*]; Mandalay, iii [*Annandale*].

var. described by Kieffer, Rec. Ind. Mus. VI, 126 (1911) from Calcutta, ix [*Lloyd*]; 2-i, at light [*Annandale*]. *Paratypes* of normal form and the variety in Indian Museum.

fumipennis, *id.*, Mem. Ind. Mus. II, 217, ♀ (*Procladius*) (1910).

Calcutta, iii, vi [*Annandale*]; Giridih, Bengal [*Kemp*]. *Type* belonging to Indian Museum. A variety (descr. Kieff., *loc. cit.*), from Puri, iii [*Paiva*], in Indian Museum.

paivai, *id.*, *loc. cit.*, 218, ♀ (*Procladius*) (1910).

Puri, 2-iii-08 [*Paiva*]. *Type* in Indian Museum, unique.

fuscusignatus, *id.*, *loc. cit.*, 218, ♂ ♀, pl. xi, 6, wing (*Procladius*) (1910).

Calcutta Zoolog. Gardens, vii [*Annandale*]; Sylhet, vii [*Hall*]; Purneah [*Paiva*]; Madhupur, x [*Paiva*]; Giridih, Bengal, [*Kemp*].

var. **aterrimus**, *id.*, *loc. cit.*, 219, ♀.

Puri, iii [*Paiva*]. Unique.

var. **flavidus**, *id.*, *loc. cit.*, 219, ♀

Bengal, between Bolpore and Rampore, at light in railway carriage; Calcutta, viii [*Annandale*]; Sylhet, vii. [*Lt-Col. Hall*].

var. **vulgaris**, *id.*, Rec. Ind. Mus., IX, 157, ♀ (1913).

Madhupur, Bengal, x [*Paiva*]; Tinpahar, Bengal, 7-vii; Bho-gaon, N. Bengal, 22-iii [*Paiva*]; Calcutta, 17-vii, 13-viii, 22-ix [*Annandale*]; Kankondigee, Sunderbunds, Ganges Delta, on board launch, at light, 14-xi [*Jenkins*], Bosondhur, Khulna Dist., Ganges Delta, on board launch at light, 29-viii [*Jenkins*]; on board ship 10 miles off Masulipatam, Madras Coast, 4-vi [*Paiva*]; Balighai, Orissa, 26-x [*Annandale*]. An additional form of this var. described (*l.c.*) by Kieffer from Madhupur, Bengal.

var. **jenkinsi**, *id.*, *loc. cit.*, 157, ♀ (1913).

Calcutta, 18-ix; Bosondhur, Khulna Dist., Ganges Delta, on board launch at light, 29-viii [*Jenkins*]; Balighai, Orissa, 25-x [*Annandale*]; Shasthancottah, Travancore, 6-xi [*Annandale*].

var. A further form, unnamed, described by Kieffer (*loc. cit.*, 158) from on board ship 10 miles off Masulipatam, Madras Coast, 4-vi [*Paiva*]. *Paratypes* of all the above varieties and the normal form in the Indian Museum. No *type* set up.

atratus, *id.*, Mem. Ind. Mus. II, 219, ♀ (*Procladius*) (1910).

Sylhet, 22-vii-05 [*Hall*]. *Type*, unique, in Indian Museum.

9-punctatus, *id.*, *loc. cit.*, 219, ♂ (*Procladius*) (1910).

Calcutta Zoological Gardens, iii, vii [*Annandale*]; at light, 2 and 14-viii; 12-ix [all *Annandale*]; Rajmahal, Bengal, vii; between Bolpore and Rampore, at light in railway carriage; Bettiah, Champaran, iii (both Bengal); Mandalay, iii [*Annandale*]; Puri, iii [*Paiva*]. Several *paratypes* in Indian Museum.

Trichotanypus, Kieff.

Ann. Soc. Sci. Brux. XXX, 319 (1906): no species quoted,
but one mentioned during same year.

Kieff., Gen. Ins., Fasc 42, 42.

GENOTYPE, *Tanypus posticalis* Lundb. accord-
ing to Coquillett (1910).

Subgenus Protenthes, Johan.

Ent. News Philad. XVIII, 400 (1908).

GENOTYPE, *Tipula cincta* F., as *punctipennis*
Mg.

bilobatus Kieff., Mem. Ind. Mus II, 221, ♂ ♀ (*Tanypus*) (1910).
Id., Rec. Ind. Mus. VI, 330 (*Trichotanypus*, subgenus *Pro-*
tenthes).

Sylhet, ii [Hall]; Calcutta, v, vi, vii, xi [Annandale]; Raj-
shahi, E. Bengal, ii.

microcerus Kieff., Mem. Ind. Mus. II, 224, ♀ (1910).

Puri, 2-iii-08 [Paiva]. *Type* in Indian Museum, a unique
specimen.

Kieffer describes (Rec. Ind. Mus. IX, 155) a variety, also
a ♀ from Naini Tal [Lloyd]. Three specimens in the Indian
Museum.

photophilus Kieff., Nova Guin. Results, 223, ♂ (1910).

Id., Rec. Ind. Mus. IX, 154, ♀ (1913). *Paratypes* in Indian
Museum.

Calcutta, i, at light [Annandale], 16-ix, at light, 18, 21-vi;
11-vii; 14-viii [Annandale]; Asansol, Bengal, 14-ii [Paiva,
Caunter]; Ernakulam, Cochin State, 4-xi [Annandale]; Shen-
cotta, Travancore, 25-xi [Annandale]; on ship 10 miles off
Masulipatam, Madras Coast, 5-vi [Paiva]. *Paratypes* in Indian
Museum.

Procladius, Skuse.

Proc. Linn. Soc. N.S. Wales (2), IV, 283 (1889).

Kieff., Gen. Ins., Fasc. 42, 38: *Id.*, Mem. Ind.

Mus. II, 216, tab. Ind. spp.

GENOTYPE, *P. paludicola* Skuse, by designation
of Coquillett (1910).

crux Wied., Analec. Entom., 10 (*Tanypus*) (1824).

Id., Auss. Zweifl. I, 19 (*Tanypus*): Wulp, Notes Leyd. Mus.
VI, 250 (*Tanypus*): de Meij., Tijd. v. Ent. LVI, 335
(*Procladius*).

pardalis Dol., Nat. Tijd. Ned. Ind. X, 405, pl. vi, 1 (1856).

ornatus, *id.*, loc. cit., XIV, 385, pl. vii, 1 (1897).

Java, Semarang, i, ii, vi, vii [Jacobson]. *Type* in Copen-
hagen Museum.

- formosanus** Kieff., Supp. Ent. Berl. I, 31 (1912).
bilobatus Kieff., Rec. Ind. Mus. IX, 155, ♂ ♀ (1913).
 Asansol, Bengal, 13-ii [Paiva, Caunter]. *Types*, a unique pair, in Indian Museum.
inconspicuus, *id.*, *loc. cit.*, 156, ♀, pl. xii, 23, antenna ♀ (1913).
 Moulmein, 26-ii [Annandale]. *Type* belonging to Indian Museum.
pallidus, *id.*, *loc. cit.*, 156, ♀ (1913).
 Mandalay, 10-iii, ♀ [Annandale]. *Type* belonging to Indian Museum.
obscuripes de Meij., Tijds. v. Ent. LVI, 336, ♀ (1913).
 Semarang, ii, iii [Jacobson]. *Type* in Amsterdam Museum.

Isoplastus, Skuse.

- Proc. Linn. Soc. N.S. Wales (2) IV, 279 (1889).
 Kieff., Gen. Ins., Fasc. 42, 43: *id.*, Mem. Ind. Mus. II, 210 (tab. Ind. spp.).
 GENOTYPE, *I. notabilis* Skuse, the first species, by designation of Coquillett (1910).
annandalei, Kieff., Mem. Ind. Mus. II, 211, ♂, pl. ix, 6, 7, 9, var. chars. (1910).
 Calcutta, vii 07 [Annandale]. *Type* belonging to Indian Museum.
indicus, *id.*, *loc. cit.*, 212, ♂ ♀ (1910).
 Calcutta, i, vii [Annandale]; Katihar, vii [Paiva]. *Paratypes* in Indian Museum.
setosicornis, *id.*, *loc. cit.*, 213, ♀ (1910).
 Bettiah, Champaran, Bengal, 6-iii-08. *Type*, a unique specimen, in Indian Museum.
variipes, *id.*, *loc. cit.*, 215, ♂ (1910).
 Mandalay, 11-iii-08 [Annandale]. *Type*, a unique specimen, in Indian Museum.
photophilus Kieff., Rec. Ind. Mus. VI, 126, ♀, pl. vi, 11, antenna (1911).
 Katihar, at light, 14-x-07 [Paiva]. *Type*, a unique specimen, in Indian Museum.

Subfamily *CHIRONOMINAE*.¹

Section *CLUNIONARIAE*.²

Nepalia, Kieff.

Rec. Ind. Mus. VI, 348 (1911).

GENOTYPE, *N. filipes*, the original species.

¹ For groups *v.* Kieff., Rec. Ind. Mus. IX, 120.

² For table of all genera *v.* Kieff., Rec. Ind. Mus. IX, 121.

filipes Kieff., Rec. Ind. Mus. VI, 338, ♂ (1911).

Noalpur, Nepal, 23-ii-08. The unique *type* is no longer in existence.

Section ORTHOCLADIARIAE.¹

Metriocnemus, Wulp.

Tijd. v. Ent. XVII, 136 (1874).

Kieff., Gen. Ins., Fasc. 42, 31.

GENOTYPE, *Chironomus albolineatus*, Mg., by Coquillett's designation (1910).

Wulpiella, Kieff., Bull. Soc. Ent. Fran. LXVIII, 66 (1899).

callinotus Kieff., Rec. Ind. Mus. VI, 175, ♂ (1911).

Simla Hills, 25-iv-07 [*Annandale*]. *Type*, a unique specimen, in Indian Museum.

fusiger, *id.*, *loc. cit.*, 348, ♂ ♀ (1911).

Simla, 12-v-08 [*Annandale*]. Four *paratypes* in Indian Museum.

Camptocladus, Wulp.

Tijd. v. Ent. XVII, 133 (1874).

Kieff., Gen. Ins., Fasc. 42, 24.

GENOTYPE, *Tipula byssina* Schrk., by Coquillett's designation (1910).

monticola Kieff., Rec. Ind. Mus. VI, 346, ♀, pl. xiv, 9, wing (1911).

Simla, 11-v-08. The unique *type* in the Indian Museum is a mere fragment.

Rhopalocladus, Kieff.

Rec. Ind. Mus. VI, 347 (1911).

GENOTYPE, *R. himalayae*, the original species.

himalayae Kieff., Rec. Ind. Mus. VI, 347, ♂.

Barogh, Simla, 10-v-10 [*Annandale*]. *Type* lost.

Conacladius, Kieff.

Mem. Ind. Mus. II, 242 (1910).

GENOTYPE, *C. flavus*, the original species.

flavus Kieff., Mem. Ind. Mus. II, 242, ♂, pl. xi, 15, head, 13, tip of genitalia (1910).

Calcutta. *Type* belonging to Indian Museum.

¹ For table of genera *v.* Kieff., Rec. Ind. Mus. IX, 122, and Spol. Zeyl. VIII, 23.

Dolichocladius, Kieff.

Rec. Ind. Mus. VI, 176 (1910).

GENOTYPE, *D. heteropterus* Kieff., the original species.**heteropterus** Kieff., Rec. Ind. Mus. VI, 176, ♂ (1910).Base of Dawna Hills, 4-iii-08 [*Annandale*]. Type in Indian Museum, a unique specimen.**Orthocladius**, Wulp.

Tijd. v. Ent. XVII, 132 (1874).

Kieff., Gen. Ins., Fasc. 42, 25.

GENOTYPE, *Tipula stercoraria*, De Geer, by Coquillett's designation (1910).Subgenus **Trichocladius**, Kieff.

Ann. Soc. Sci. Brux. XXX, 356 (1906).

Kieff., Gen. Ins., Fasc. 42, 25; *id.*, Rec. Ind. Mus. VI, 177 (1911).GENOTYPE, *T. fissicornis* Kieff., the original species.**spatulicornis** Kieff., Rec. Ind. Mus. VI, 177, ♂, pl. vii, 33, antenna, 34, genitalia, pt. (1911).Bengal, between Manihari and Manshahi, on railway track, 4-viii-07. Three *paratypes* in Indian Museum.**pentazonus**, *id.*, *loc. cit.*, VI, 345, ♂ (1911).Thaumaspur, Nepal, 18-ii-08; Tharbani, Nepal, 27-ii-08. Three *paratypes* in Indian Museum.**oriplanus**, *id.*, *loc. cit.*, 346, ♂ (1911).Tonglu, Darjiling, 10,000 ft., 21-iv-10 [*C. W. Beebe*]. Type, unique, but much damaged, in Indian Museum.**anomalous**, *id.*, *loc. cit.*, IX, 124, ♂ (1913).Valley of Sutlej River, below Simla, 6-v [*Annandale*]; Darjiling, 8-viii [*Jenkins*]. Type belonging to Indian Museum.Subgenus **Psectrocladius**, Kieff.

Ann. Soc. Sci. Brux. XXX, 356 (1906).

Kieff., Gen. Ins., Fasc. 42, 26; *id.*, Rec. Ind. Mus. VI, 176 (1911).GENOTYPE, *Orthocladius psilopterus*, Kieff., according to Coquillett (1910).**flavicollis** Kieff., Rec. Ind. Mus. VI, 176, ♀ (1911).Calcutta, 31-i-08 [*Annandale*]. Type in Indian Museum, unique.

Subgenus *Dactylocladius*, Kieff.

Ann. Soc. Sci. Brux. XXX, 356 (1906).

Kieff., Gen. Ins., Fasc. 42, 26: *id.*, Rec. Ind.

Mus. IV, 174 (1911).

GENOTYPE, *Orthocladius brevicornis*, Kieff., according to Coquillett (1910).*tenuicrus* Kieff., Rec. Ind. Mus. VI, 174, ♂ (1910).Rajshahi, E. Bengal, 6-ii-07 [*Annandale*]. *Type*, unique, in Indian Museum.*ceylanicus* Kieff., Spol. Zeyl. VIII, 21, ♂, fig. 8, genitalia tip (1912).Peradeniya, 14-vii-10. *Type*, unique, in Indian Museum.*Cardiocladius*, Kieff.

Spol. Zeyl. VIII, 22 (1912).

nom. nov. for *Thalassomyia* Johann., nec. Sch.*Thalassomyia*, Johan., Bull. N.Yk. State Mus.LXVIII, 437, ♂ ♀ (1903): nec *Thalassomyia*, Sch., 1856.Kieffer sets up *ceylanicus*, sp. nov. as type of his new genus, stating it to be synonymous with Johannsen's, and that the latter is different from Schiner's.¹*ceylanicus* Kieff., Spol. Zeyl. VIII, 22, ♀, fig. 9, flagellum (1912).Peradeniya, 2-vi-10. *Type* in Indian Museum.*Cricotopus*, Wulp.

Tijd. v. Ent. XVII, 132 (1874)

Kieff., Gen. Ins., Fasc. 42, 23.

GENOTYPE, *Chironomus tibialis* Mg., by designation of Coquillett (1910).*carneus* Kieff., Supp. Ent. Berl. I, 32 (1912).

Formosa.

brunetti Kieff., Rec. Ind. Mus. IX, 125, ♂ ♀ (1913).Kurseong, 6-ix-09, at light [*Annandale*]; Darjiling, 6000 ft., 28-ix-08 and 1-x-07 [*Brunetti*]. Several *paratypes* in Indian Museum.Section CHIRONOMARIAE.²*Tanytarsus*, Wulp.

Tijd. v. Ent. XVII, 134 (1874).

Kieff., Gen. Ins., Fasc. 42, 29: *id.*, Mem. Ind.Mus. II, 239, tab. Ind. spp.: *id.*, Rec. Ind. Mus.

VI, 166, tab. add. Ind. spp. nov.

¹ The work in which *Thalassomyia*, Johann., is described is inaccessible in India.² This name is formed to be in keeping with the names of the other two sections admitted by Kieffer, but the term used by that author is *Tendipedariae*, which is inadmissible, being founded on an inadmissible generic name, *Tendipes*. For table of genera in this section, see Kieffer, Rec. Ind. Mus. IX, 126.

GENOTYPE, *Chironomus punctipes* Wied., by Coquillett's designation (1910).

- clavatitarsis** Kieff., Mem. Ind. Mus. II, 240, ♀, pl. ix, 5, middle tarsus, part (1910).
Calcutta, vii. Type belonging to Indian Museum.
- setosipennis**, *id.*, *loc. cit.*, 240, ♂ (1910).
Calcutta, vii-07 [Annandale]. Type belonging to Indian Museum.
- breviventris**, *id.*, *loc. cit.*, 241, ♀ (1910).
Calcutta, vii-07 [Annandale]. Type belonging to Indian Museum.
- pilosipennis**, *id.*, *loc. cit.*, 241, ♀ (1910).
Calcutta, vii-07 [Annandale]. Type belonging to Indian Museum.
- viridis** Kieff., Rec. Ind. Mus. VI, 168, ♂ ♀, pl. vii, 25, wing, ♂ (1911).
Puri, 2-iii [Paiva]. Paratypes in Indian Museum.
Kieffer describes a variety, *loc. cit.*, ix, 152, from Puri, 19-i; 24-ii; 2-iii [Paiva]. Type in Indian Museum.
- hirtipes**, *id.*, *loc. cit.*, 168, ♂ (1911).
Sylhet, 27-i-05 [Hall]. Type in Indian Museum, a unique specimen.
- lasiopterus**, *id.*, *loc. cit.*, 169, ♂, pl. vii, 26, genitalia, part (1911).
Rajshahi, E. Bengal, 6-ii-07 [Annandale]. Type belonging to Indian Museum.
- confundendus**, *id.*, *loc. cit.*, 169, ♂, pl. vii, 27, genitalia (1911).
Port Canning, 21-vii [Annandale]; Calcutta, 23-vii [Lloyd].
Paratypes in Indian Museum.
- adjacens**, *id.*, *loc. cit.*, 170, ♀ (1911).
Rajshahi, E. Bengal, 6-ii-07 [Annandale]. Type, a unique specimen, in Indian Museum.
- semiglaber**, *id.*, *loc. cit.*, 170, ♀ (1911).
Calcutta, 30-vii-07, at light [Annandale]. Type in Indian Museum, a unique specimen.
- macrochirus**, *id.*, *loc. cit.*, 171, ♀ (1911).
Bhim Tal, 27-ix-06 [Annandale]. Type, a unique specimen, in Indian Museum.
- leptogastrus**, *id.*, *loc. cit.*, 172, ♂ (1911).
Port Canning, 21-vii-07 [Annandale]. Type, a unique specimen, in Indian Museum.
- lasiopus**, *id.*, *loc. cit.*, 172, ♀ (1911).
Port Canning, 21-vii-07 [Annandale]. Type, a unique specimen, in Indian Museum.
The month of capture is July, not August as quoted by Kieffer.

- formosanus** Kieff., Supp. Ent. Berl. I. 42 (1912).
Formosa.
- lobatus** Kieff., Spol. Zeyl. VIII, 18, ♀ (1912).
Peradeniya, 2-vi-10. Type in Indian Museum, unique,
damaged.
- transversalis**, *id.*, *loc. cit.*, 19, ♂ ♀, fig. 6, antenna tip (1912).
Peradeniya, 2 and 23-vi-10, type ♂ ♀. Types in Indian
Museum, unique.
- prasiogaster**, *id.*, *loc. cit.*, 20, ♂ ♀ (1912).
Peradeniya, 30-v-10. Paratypes in Indian Museum.
- ceylanicus**, *id.*, *loc. cit.*, 20, ♀, fig. 7, antenna tip (1912).
Peradeniya, 17 and 19-v-10. Paratypes in Indian Museum.
- poecilus**, *id.*, *loc. cit.*, 21 (1912).
Peradeniya, 13-v-10. Fragments of type in Indian Museum,
in Canada balsam.
- bicornutus** Kieff., Rec. Ind. Mus. IX, 151, ♀ (1913).
Giridih, Bengal, 2-xi [Kemp]. Two paratypes in Indian
Museum, in alcohol.
- pentatomus**, *id.*, *loc. cit.*, 152, ♀, pl. xii, 22, antenna (1913).
Puri, 13-xi [Kemp]. Paratypes in Indian Museum, in al-
cohol.

Subgenus *Micropsectra*, Kieff.¹

- indica** Kieff., Mem. Ind. Mus. II, 242, ♀ (*indicus*) (1910).
Calcutta, vii-07 [Annandale]. Type belonging to Indian Mu-
seum.

Polypedilum, Kieff.²

- tripartitum** Kieff., Mem. Ind. Mus. II, 231 (*Prochironomus tripar-
titus*) ♂ ♀ (1910).
Id., Rec. Ind. Mus. IX, 150, notes.
Madhupur, Bengal, 17-x, at light [Paiva]; Poradah, E. Bengal,
in railway station, 26-vi [Jenkins]; Bosondhur, Ganges Delta,
29-viii, on launch at light [Jenkins]; Puri Distr., 14-i and
8-xii [Caunter]; Kuching, Sarawak, 16-vi [Beebe]; on board
ship, Suez Canal, 9-x-07; type ♂ and ♀ [Annandale]. Types
in Indian Museum.
- annulatipes**, *id.*, Mem. Ind. Mus. II, 234, ♀ (1910).
Id., Rec. Ind. Mus. IX, 151, ♂ ♀ (1913).
Puri, iii [Paiva]; Lake Chilka, at light, iii [Annandale];
Khulna Distr., Ganges Delta, on board launch at light [Jenkins].
Paratypes in Indian Museum.

¹ I have been unable to trace the origin of this genus.

² I have been unable to trace the origin of this genus. Kieffer (Rec. Ind. Mus. IX, 146) gives a table of the Indian species.

fasciatipenne Kieff., Mem. Ind. Mus. II, 238, ♀, pl. xi, 4, antenna tip (*Prochironomus*) (1910).

Id., Rec. Ind. Mus. VI, 135 (*Chironomus*): *id.*, *loc. cit.*, IX, 150 (*Polypedilum*).

Calcutta, i, ii, vi, vii, viii, ix [mostly *Annandale*]. Larva feeds on *Hydra* (v. Journ. As. Soc. Bengal (n.s.), II, 1906, p. 112): Ann., Faun. Brit. Ind., Freshwater Sponges, 155; Bains Prashad, Rec. Ind. Mus. XV, 71, anatomy larva of this or an allied species, pl. xxiii, var. chars. *Paratypes* in Indian Museum.

pelostolum Kieff., Supp. Ent. Berl. I, 41 (1912).

Formosa.

crisatum Kieff., Rec. Ind. Mus. IX, 146, ♂ ♀ (1913).

Katihar, N. Bengal, ix [*Paiva*]; Port Canning, 6-xii; Lake Chilka, iii [*Annandale*]. *Paratypes* in Indian Museum.

kempi, *id.*, *loc. cit.*, 147, ♀, pl. xii, 18, antenna (1913).

Puri, 13-xi [*Kemp*]. *Type* in Indian Museum, in alcohol.

angustiforceps, *id.*, *loc. cit.*, 148, ♂, xii, 19, genitalia (1913).

Katihar, Bengal, ix [*Paiva*]. *Type* belonging to Indian Museum.

milnei, *id.*, *loc. cit.*, 148, ♂ (1913).

Berhampur, Murshidabad Distr., Bengal, xii [*Milne*]. *Type*, a unique specimen, in Indian Museum.

The only specimen in the Museum, which is marked "type," has no month given but is dated 1907 only.

purimanus, *id.*, *loc. cit.*, 149, ♂ (1913).

Calcutta, ix [*Lloyd*]. *Type* belonging to Indian Museum.

brumalei, *id.*, *loc. cit.*, 149, ♀, pl. xii, 20, antenna (1913).

Giridih, Bengal, 7-xi [*Kemp*]. Two *paratypes* in alcohol in Indian Museum.

Paratendipes, Kieff.¹

Rec. Ind. Mus. IX, 126 (1913).

GENOTYPE, unknown to present author.

diagraphis Kieff., Rec. Ind. Mus. VI, 335, ♂ ♀, pl. xiv, 3, genitalia, part (*Tendipes*) (1911).

Simla, 9-v-10; Mundali (Dehra Dun, W. Himalayas), 12-v-10. *Paratypes* in Indian Museum.

melanothorax, *id.*, *loc. cit.*, 335, ♂ ♀, pl. xiv, 2, genitalia, part (*Tendipes*) (1911).

Thamaspur, Nepal, 20-ii-08; Tharbani, 27-ii-08. *Type* belonging to Indian Museum.

¹ The only characterisation of this genus that I can find is its inclusion in a table of genera of this group, as quoted, but it is not therein given as new. Kieffer gives a table of four Indian species, p. 143.

pelargus, *id.*, *loc. cit.*, IX, 143, ♂ ♀, pl. xii, 16, last 2 joints ♀ antenna (1913).

Calcutta, 29-viii-07, *type*, ♂, 11-xii-07, *type* ♀. *Types* in Indian Museum, the only specimens.

ditaenius, *id.*, *loc. cit.*, 144, ♀ (1913).

Katihar, 14-x-07 [*Paiva*]. The unique *type* in the Indian Museum is a useless fragment.

caunteri, *id.*, *loc. cit.*, 144, ♀ (1913).

Rajmahal, Bengal, 5-vii, at light [*Annandale*]; on board launch at light, Borondhur, Khulna Distr., Ganges Delta, 29-viii [*Jenkins*]; Chennia, 24 Perghs., Sunderbunds, 10-xi [*Jenkins*]; Balighai, near Puri, 23-x [*Annandale*]. *Paratypes* in Indian Museum.

dolens, *id.*, Mem. Ind. Mus. II, 233, ♀ (*Prochironomus*) (1910).

Id., Rec. Ind. Mus. IX, 145, ♂ ♀, pl. xii, 17, genitalia tip (1913).

Calcutta, ii, viii, xi, xii; at light, i; Bettiah, Bengal, iii; Katihar, x [*Paiva*]; Sonapur, Bengal, 24-xii, at light in railway carriage [*Annandale*]; Puri, 2-iii [*Paiva*]. Several *paratypes* in Indian Museum.

A variety from the base of the Dawna Hills, taken in March by Dr. Annandale, noted by Kieffer (Mem. Ind. Mus. II, 234).

Glyptotendipes, Kieff.¹

GENOTYPE, *G. sigillatus* Kieff., by original designation.

verrucosus Kieff., Rec. Ind. Mus. VI, ♂ ♀ (*Chironomus*) (1911).

Id., *loc. cit.*, IX, 140, note, pl. xi, 11, genitalia tip (*Tendipes*).

Bhim Tal, 25, 27-ix-06 [*Annandale*]; Calcutta, 26-v; 14-vii; 28-xii-07. *Paratypes* in Indian Museum.

Prochironomus, Kieff.²

Mem. Ind. Mus. II, 225 (1910).

GENOTYPE unknown to present author.

prussicus Kieff., Mem. Ind. Mus. II, 229, ♂ (1910).

Simla, 12-v-08 [*Annandale*]. *Type*, a much damaged unique specimen, in Indian Museum.

pictiventris, *id.*, *loc. cit.*, 229, ♂ (1910).

Bettiah, Champaran, Bengal, 4-iii-08. *Type*, a unique specimen, in Indian Museum.

¹ The original description of this genus is unknown to me.

² Although Kieffer removes three species, *striatipennis*, *punctatipennis* and *fasciatipennis*, to *Chironomus*, he says nothing about the remainder from which I infer he still considers *Prochironomus* a valid genus.

annandalei, *id.*, *loc. cit.*, 230, ♀, pl. ix, 10, 11, var. chars. (1910).

Calcutta, vii-07 [*Annandale*]. *Type* technically in Indian Museum, but the only specimen there is labelled "Calcutta, ix-07, *typé*." An error seems to have crept in somewhere.

stripes, *id.*, *loc. cit.*, 231, ♀ (1910).

Katihar, 4-viii-07 [*Paiva*]. *Type*, a unique specimen, in Indian Museum.

At first considered possibly the ♀ of *albiforceps*, but later (Rec. Ind. Mus. VI, 143) Kieffer admits it as a good species.

clavatipes, *id.*, *loc. cit.*, 232, ♂ (1910).

Bengal, between Manihari and Maushali, on railway track, 4-viii-07 [*Paiva*]. *Type* in Indian Museum, a unique specimen.

graciliforceps, *id.*, *loc. cit.*, 235, ♂ (1910).

Bengal, between Bolpore and Rampore, at light in railway carriage, 3-viii-07 [*Paiva*]. *Type* in Indian Museum, a unique specimen.

curtimanus, *id.*, *loc. cit.*, 235, ♀ (1910).

Mandalay, iii [*Annandale*]. *Type* belonging to Indian Museum.

hallii, *id.*, *loc. cit.*, 237, ♀ (1910).

Sylhet, 9-ii [*Hall*]. *Type* belonging to Indian Museum.

tenuitarsis, *id.*, *loc. cit.*, 237, ♀ (1910).

Rangoon, 25-ii-08 [*Annandale*]; Mandalay, 11-iii-08 [*Annandale*]. Two *paratypes* in Indian Museum.

fimbriatus, *id.*, *loc. cit.*, 238, ♀ (1910).

Tharbani, Nepal, 27-ii-08. Kieffer erroneously quotes November. *Type*, a unique specimen, in Indian Museum.

Chironomus, Mg.

Illig. Mag. II, 260 (1803).

Sch., F. Austr. II, 595: *Auctt. var.*: Kieff., Gen. Ins., Fasc., 42, 14: *id.*, Mem. Ind. Mus. II, 225, tab. Ind. spp.: *loc. cit.*, VI, 127 and VI, 333, tab. addit. spp. nov. (as *Tendipes*): *loc. cit.*, IX, 129, tab. 37 Ind. spp. (as *Tendipes*): *id.*, Spol. Zeyl. VIII 10, tab. spp.

GENOTYPE, *Tipula plumosa* L., by Latreille's designation.

venustus W., Auss. Zweifl. I, 547 (1828).¹

China. *Types* in Wiedemann's and Trentepohl's collections.

socius Walk., List Dipt. Brit. Mus. I, 16 (1848).

East India. *Type* in British Museum.

¹ Probably should not be included in oriental lists.

- cubiculorum** Dol., Nat. Tijd. Ned. Ind. X, 405, pl. v, 2 (1856).
Java. Type probably lost.
- vicarius** Walk., Ins. Saund. Dipt., pt. V, 423, ♂ ♀ (1856).
East India. Type, location unknown to present author.
- pictus** Dol., Nat. Tijd. Ned. Ind. XIV, 386, pl. vii, 2 (1857).
Java. Type probably lost.
- instabilis** Walk., Proc. Linn. Soc. Lond. VIII, 103 (1865).
Papua. Type, location unknown to present author.
- trochanteratus** Thoms., Eugen. Resa, 445 (1869).
Manila. Type, location unknown to present author.
- novaeguineensis** de Meij., Nova Guinea Results, V, 69 (1906).
Papua. Type in Amsterdam Museum.
- campogaster** Kieff., Mem. Ind. Mus. II, 227, ♂, pl. xi, 2, abdomen tip (1910)
Calcutta. Type belonging to Indian Museum.
- longivalvis**, *id.*, *loc. cit.*, 228, ♂, pl. xi, 11, genitalia, pt. (1910).
Calcutta. Type belonging to Indian Museum.
- bipunctatus**, *id.*, *loc. cit.*, 228, ♂ (1910).
Id., Rec. Ind. Mus. IX, 139, ♀ (*Tendipes*).
Rangoon, 25-ii-08 [*Annandale*]. Two *paratypes* in Indian Museum.
- albiforceps**, *id.*, Mem. Ind. Mus. II, 231, ♂ (*Prochironomus*) (1910).
Calcutta, 5-i; ii, iii, 18-v; 2-vii; viii, ix, x, xi, xii; Dum Dum, 28-vii [*Lord*]; Bhogaon, Bengal, 2-x; Katihar, Bengal, 4-viii [*Paiva*]; on board launch at light, Port Canning [*Annandale*]; Bologhatta, near Khulna, Ganges Delta, 28-viii and Khulna, 27-viii [both *Jenkins*]; Katihar, viii [*Paiva*]; Mandalay, 12-iii [*Annandale*]; Peradeniya, 17-v-10, at light. *Paratypes* in Indian Museum.
- punctatipennis**, *id.*, Mem. Ind. Mus. 234, ♂ ♀ (*Prochironomus*) (1910).
Id., Rec. Ind. Mus. IX, 138, pl. xi, 10, genitalia (*Tendipes*).
Rajmahal, Bengal, 31-vii-07, *type* ♀, viii; Puri 2-iii-03 [*Paiva*]; Katihar, 4-viii-07, *type* ♂; ix [*Paiva*]; at light on launch, Bosphor, Khulna Distr., Ganges Delta, 29-viii [*Jenkins*]; Shencottah, Travancore, 25-xi-08 [*Annandale*]. *Types* in Indian Museum, the ♂ a mere fragment.
- striatipennis** Kieff., Mem. Ind. Mus. II, 326, ♀ (*Prochironomus*) (1910).
Id., Rec. Ind. Mus. VI, 134, ♂ ♀, pl. vi, 12, genitalia tip, (*Chironomus*): *id.*, *loc. cit.*, IX, 138, notes (*Tendipes*).
Bhim Tal, 27-ix-06 [*Annandale*]; Calcutta, 22-xii; Shencottah, Travancore, 25-xi [*Annandale*]; Victoria Gardens, Colombo, 26-iv [*Paiva*] Several *paratypes* in Indian Museum.
- speciosus** Kieff., Rec. Ind. Mus. VI, 133, ♂ (1911).
Matiana [*Annandale*]. Type, a unique specimen, in Indian Museum.

- pruinus**, *id.*, *loc. cit.*, 134, ♀ (1911).
Barrackpore, near Calcutta, 11-viii-07 [Annandale]. *Type*
in Indian Museum, a unique specimen.
- nigrosparus**, *id.*, *loc. cit.*, 135, ♀ (1911).
Mandalay, 11-iii-08 [Annandale]. *Type*, a unique specimen,
in Indian Museum.
- ceylanicus**, *id.*, *loc. cit.*, 136, 137, ♂ ♀ (1911).
Colombo, vii-1901. *Paratypes* in Indian Museum. Known
in Colombo as the "Lake Fly"; see Ind. Mus. Notes, V, 191
and 195.
- polystictus**, *id.*, *loc. cit.*, 137, ♂ (1911).
Bengal (between Bolpore and Rampore), 3-viii-07, at light in
railway carriage [Paiva]. *Type* in Indian Museum, a unique
specimen.
- lobaticeps**, *id.*, *loc. cit.*, 138, ♂ ♀, pl. vi, 13, genitalia, tip (1911).
Sylhet, ii [Hall]; Bijnor (United Prov., India), i; Calcutta, xi,
xii; Port Canning, Bengal, xii. *Paratypes* in Indian Museum.
var. **validus** Kieff., in H. Sauter's *Formosa Ausbeute*, *Tendi-*
pedidae (Supp. Ent. 1912, V, 1, p. 311, ♂ ♀) (*Tendipes*)?
Id., Rec. Ind. Mus. IX, 140, ♂ ♀, note (*Tendipes*).
Calcutta, 30-i; iii, iv, 5-xii; Madhupur, Bengal, x; Sonadigee,
Ganges Delta, 6-xii, at light; Formosa, ii, iii [Sauter]. *Para-*
types in Indian Museum.
- lobaticollis** Kieff., Rec. Ind. Mus. VI, 139, ♂ ♀, pl. vi, 14, geni-
talia tip (1911).
Sylhet, ii [Hall]. *Paratypes* in Indian Museum
- flaviventris**, *id.*, *loc. cit.*, 139, ♂ (1911).
Id., *loc. cit.*, IX, 141, note (*Tendipes*).
Puri, 20--21-i-08, *type*; Balighai, near Puri, 26-x-08 [Annandale].
Type in Indian Museum.
- grossipes**, *id.*, *loc. cit.*, VI, 141, ♀ (1911).
Calcutta, v, vii [Annandale]. Two *paratypes* in Indian
Museum.
- nigromarginatus**, *id.*, *loc. cit.*, 141, ♀ and 344, note (1911).
Id., Spol. Zeyl. VIII, 17, note (*Tendipes*).
"Probably the ♀ of *C. callithorax*" (Kieffer).
Lucknow, iv; Calcutta, i, iii, v, viii, also iii, at light; Port
Canning, 21-viii [Annandale]; Lake Chilka, iv; Chandan Chowki,
United Prov., India, 8-v-07; Kawkereik, Amherst Distr., Lower
Burma, 5-iii; Peradeniya, at light. 14-v; 17-v; 17-vi-10.
var. **macrogaster**, *id.*, Rec. Ind. Mus. VI, 142, ♀ (1911).
Maldhun, Naini Tal Distr., 14-ii, United Prov., India; Kuch-
ing, Borneo. *Paratypes* of normal form and variety in
Indian Museum. The var. *macrogaster* is represented by a unique
specimen.
- callithorax**, *id.*, *loc. cit.*, 142, ♂ (1911).
Lucknow 21-v-07: Bijnor Distr., United Prov., India, var.

- locs., xi-07; Sonarpur, Lower Bengal, 24-xii, at light in railway carriage [*Annandale*]; Calcutta, 30-iv-07, at light; 24-vi-07; Rajshahi, E. Bengal, 6-ii-07; Mandalay, 12-iii-08. Several *paratypes* in Indian Museum.
- melanostolus**, *id.*, *loc. cit.*, 143, ♀ (1911).
Lake Chilka, 7—15-viii-07. *Type* in Indian Museum, a unique specimen.
- nigriforceps**, *id.*, *loc. cit.*, 143 ♂ (1911).
Chandan Chowki, United Prov., India, 8-v-07. *Type*, a unique specimen, in Indian Museum.
- dolichogaster**, *id.*, *loc. cit.*, 144 ♂ (1911).
Id., *loc. cit.*, ix, 142, ♂ ♀, pl. xii, 15, genitalia tip (*Tendipes*) (1913)
Calcutta, ii-08 [*Annandale*]; viii. Three *paratypes* in Indian Museum.
- seminiger**, *id.*, *loc. cit.*, 144, ♂ (1911).
Id., *loc. cit.*, ix, 141, ♂, pl. xi, 13, genitalia (*Tendipes*).
Calcutta, 3—23-iii-07 [*Annandale*]; viii. Several *paratypes* in Indian Museum.
- lampronotus**, *id.*, *loc. cit.*, 144, ♀ (1911).
Sylhet, 14-iii-05 [*Hall*]. *Type* in Indian Museum, a unique specimen.
Kieffer quotes April (iv) as the date, but though it is not quite certain, it appears to be intended for iii.
- longicrus**, *id.*, *loc. cit.*, 145, ♂ (1911).
Matiana [*Annandale*]. *Type*, a unique ♂, in Indian Museum.
- fuscitarsis**, *id.*, *loc. cit.*, 145, ♀ (1911).
Id., Spol. Zeyl. VIII, 17, note (*Tendipes*).
Calcutta, 20-viii; Peradeniya, 2-vi-10; Mandalay, 12-iii-08 (*type*) [*Annandale*]. *Type* in Indian Museum.
Kieffer (Rec. Ind. Mus. IX, 142) queries this as a variety of *lamprothorax*.
- callicomus**, *id.*, Rec. Ind. Mus. VI, 146, ♂ (1911).
Matiana [*Annandale*]. *Type* in Indian Museum.
- lamprothorax**, *id.*, *loc. cit.*, 146, ♀ (1911).
Calcutta, vii, viii, ix, xi, xii; Port Canning, vii [*Annandale*]; Katihar, viii; Puri, at light, 24-ii [*Paiva*]; between Bolpore and Rampore, Bengal, at light in railway carriage, 3-viii-07, *type* [*Paiva*]. *Type* in Indian Museum.
- var. **radialis**, *id.*, *loc. cit.*, 147, ♀ (1911).
Calcutta, vi, vii, viii. *Paratypes* in Indian Museum.
- var. **conjungens**, *id.*, *loc. cit.*, X, 315, ♂ ♀ (1914).
Calcutta, Ind. Mus. gardens, 25-viii-10. *Type* in Indian Museum.
- tenerrimus**, *id.*, *loc. cit.*, VI, 147, ♂ ♀, pl. vi, 15, genitalia, pt. (1911).

- Calcutta, i, ii, iii, vi, viii, often at light [*Annandale*]. Several *paratypes* in Indian Museum.
- var. *leucotarsus*, *id.*, *loc. cit.*, 148, ♂ (1911).
Id., *loc. cit.*, VI, 158, var. ♂, pl. vii, 22, genitalia, pt. (1911).
 Calcutta, vi, at light, viii-07 [*Annandale*]. Three *paratypes* in Indian Museum.
- Another variety of this variety described (*loc. cit.*) from Calcutta, vi, viii, at light.
- dolichotomus*, *id.*, *loc. cit.*, 149, ♂, pl. vii, 17, genitalia (1911).
 Purneah Distr., 5-viii-07 [*Paiva*]. *Type* in Indian Museum, a unique ♂.
- callinotus*, *id.*, *loc. cit.*, 150, ♂, pl. vii, 18, genitalia, pt. (1911).
 Bijnor Distr., United Prov., India, 11-ii-07. *Type* in Indian Museum, a unique ♂.
- melanophorus*, *id.*, *loc. cit.*, 150, ♀ (1911).
 Calcutta, 13-vi-07. *Type*, a unique ♀, in Indian Museum.
 Kieffer, *loc. cit.*, IX, 143, describes a variety (♀) from Rajshai, East Bengal, ii [*Annandale*].
- planicollis*, *id.*, *loc. cit.*, 151, ♀ (1911).
 Sylhet, 27-i-03 [*Hall*]. Two *paratypes* in Indian Museum.
- breviforceps*, *id.*, *loc. cit.*, 151, ♂ ♀ : 342, pl. xiv, genitalia ♂, tip (1911).
 Theog, 2-v-07 [*Annandale*]; Dharampur (Simla) [*Annandale*].
 Two *paratypes* in Indian Museum.
- oriplanus*, *id.*, *loc. cit.*, 152, ♂ (1911).
 Theog, 2-v-07 [*Annandale*]. *Type* in Indian Museum, a unique specimen.
- forficularius*, *id.*, *loc. cit.*, 153, ♂, pl. vii, 19, genitalia (1911).
 Calcutta, 13-viii-07. *Type* in Indian Museum, a unique specimen.
- barbatitarsis*, *id.*, *loc. cit.*, 154, ♂ ♀, pl. vii, 20, antenna ♀, 21, genitalia ♂ (1911).
Id., *loc. cit.*, IX, 141 (*Tendipes*).
 Calcutta, iv, v, vii, viii, ix, xi, xii; Port Canning, viii, xii; Bettiah, Champaran, Bengal, 4-iii; Lake Chilka, viii; Mandalay, iii [both *Annandale*]. Several *paratypes* in Indian Museum.
- var. *semiflavus*, *id.*, *loc. cit.*, 154, ♂.
 Calcutta (Zoolog. Gdns.), 21--25-viii-07 [*Annandale*]. Two *paratypes* in Indian Museum.
- callichirus*, *id.*, *loc. cit.*, VI, 155, ♂ (1911).
 Calcutta (Zoolog. Gdns.), 4-v. *Type*, a unique specimen, in Indian Museum.
- melanochirus*, *id.*, *loc. cit.*, 155, ♂ ♀ (1911).
 Phagu, 3-v-07 [*Annandale*]. *Types* in Indian Museum, a unique pair.
 Kieffer quotes "October," but the inscription (badly written) is intended for May.

- melanostictus**, *id.*, *loc. cit.*, 156, ♂ (1911).
Calcutta, 1-viii-07 [*Annandale*]. *Type* in Indian Museum, a unique ♂.
- callisphyrus**, *id.*, *loc. cit.*, 156, ♂ ♀ (1911).
Sylhet, 1-ii; v, vi, xi [*Hall*]. Several *paratypes* in Indian Museum.
- var. **dolichomerus**, *id.*, *loc. cit.*, 157, ♀.
Sylhet, 29-i-05 [*Hall*]. *Type* in Indian Museum, a unique specimen.
Kieffer notes another variety in *Rec. Ind. Mus.* VI, 343. ♂, from Simla, 9-v-10 and Kurseong, 20-vi-10.
- aplochirus**, *id.*, *loc. cit.*, 157, ♂ ♀ (1911).
Sylhet, 6—8-i-05; 1—10-ii [*Hall*]. Four *paratypes* in Indian Museum.
- aploneurus**, *id.*, *loc. cit.*, 158, ♀ (1911).
Sylhet, 2-xii-04 [*Hall*]. *Type*, a unique specimen, in Indian Museum.
- leptochirus**, *id.*, *loc. cit.*, 158, ♂ ♀ (1911).
Id., *loc. cit.*, IX, 141, note.
Kumaon, 22-ix; Calcutta, vii, viii, ix, xii; Port Canning, 6—24-xii; Mandalay, 6—8-i. *Paratypes* in Indian Museum.
- psilochirus**, *id.*, *loc. cit.*, VI, 159, ♂ (1911).
Id., *loc. cit.*, IX, 141, pl. xii, 14, genitalia tip (*Tendipes*).
Calcutta, 27-xi-07; Port Canning, 21-vii-07, *type* [*Annandale*].
Type in Indian Museum.
- lasiochirus**, *id.*, *loc. cit.*, VI, 159, ♂ ♀ (1911).
Id., *loc. cit.*, IX, 141, note (*Tendipes*).
Port Canning, 21-vii-07 [*Annandale*]. *Paratypes* in Indian Museum.
- filitarsis**, *id.*, *loc. cit.*, VI, 160, ♂ (1911).
Id., *loc. cit.*, IX, 143, ♀ (*Tendipes*) (1913).
Calcutta, 26-vii-08, *type* ♂; 23—31-vii; 31-vii-07, *type* ♀ [*Annandale*]; 26-viii; Port Canning, vii [*Annandale*]; Purneah and Katihar [*Paiva*]; Moulmein, 28-ii [*Annandale*]. *Types* in Indian Museum.
- lucernarum**, *id.*, *loc. cit.*, VI, 160, ♀ (1911).
Calcutta, vi, viii, at light [*Annandale*]. Two *paratypes* only, both in Indian Museum; a variety determined by Kieffer is in the same collection from Calcutta, 20-viii-07.
- calligaster**, *id.*, *loc. cit.*, 160, ♂ (1911).
Id., *loc. cit.*, IX, 140, note, pl. xi, 12, genitalia tip (*Tendipes*).
Port Canning, 21-vii-07 (*type*); 6-xii [both *Annandale*]. *Type* in Indian Museum.
- var. **vilis**, *id.*, *loc. cit.*, 161.
Calcutta, 9-viii-07, at light [*Annandale*]. *Type*, unique, in Indian Museum.

- sexpunctatus**, *id.*, *loc. cit.*, 161, ♀ (1911).
Calcutta, 15-ii-18 [Annandale] Type, a unique ♀, in Indian Museum.
- callimorphus**, *id.*, *loc. cit.*, 162, ♂ (1911).
Dawna Hills, 2—3 iii 08 [Annandale] Type, a unique ♂, in Indian Museum.
- viridiventris**, *id.*, *loc. cit.*, 162, ♂ ♀ (1911).
Mandalay, iv; 11—12-iii-08 [both Annandale]. Paratypes in Indian Museum.
- atrosignatus**, *id.*, *loc. cit.*, 163, ♀ (1911).
Calcutta, 30-vii [Annandale]; ix [Lloyd]. Type in Indian Museum.
- nudipes**, *id.*, *loc. cit.*, 164, ♂ (1911).
Calcutta, 10-viii-07 [Annandale]. Type, a unique ♂, in Indian Museum.
- rostratus**, *id.*, *loc. cit.*, 164 ♂ (1911).
Sylhet, 3-i-05 [Hall]. Type, a unique ♂, in Indian Museum.
- prasiogaster**, *id.*, *loc. cit.*, 165, ♂ (1911).
Sylhet, 26-ii-04 [Hall]. Type, a unique ♂, in Indian Museum.
- chlorogaster (I)**, *id.*, *loc. cit.*, 165, ♂ (1911).
Mandalay, 12-iii-08 [Annandale]. Type, a single ♂, in Indian Museum.
N.B.—*Vide chlorogaster (II)*, *post.*, p. 78.
- semiviridis**, *id.*, *loc. cit.*, 166, ♂ ♀, pl. vii, 13, wing, 24, genitalia (1911).
Mandalay, iii; Moulmein, ii [Annandale]; Bettiah, Champaran, Bengal, ii; Noalpur, Nepal, ii; Puri, ii [Paiva]; iii. Types in Indian Museum.
- choricus**,¹ *id.*, *loc. cit.*, 336, ♂ ♀, pl. xiv, 4, tip of ♀ antenna, 5 tip of genitalia (1911).
Barogh (Simla), ♂ ♂ in air dance at dusk; Sukhwani, 15-ii-08; Dahawangahary Hill, 16-ii-08 (both Nepal); Bijnor Distr., United Prov., India, 24-ii-10. Paratypes in Indian Museum.
- himalayanus**, *id.*, *loc. cit.*, 337, ♂ (1911).
Mundali, Dehra Dun Distr., 10-v-10. Type, a unique specimen, in Indian Museum.
- fulvescens**, *id.*, *loc. cit.*, 338, ♀, pl. xiv, 6, tip of antenna (1911).
Butal, Nepal, 12-ii-08. Type belonging to Indian Museum.
- polius**, *id.*, *loc. cit.*, 339, ♂, pl. xiv, 7, ♂ genitalia (1911).
Kasauli, N.W. India, 16-v-08 [Annandale]. Type belonging to Indian Museum.

¹ All the species from *choricus* to *bipunctatus* are described by Kieffer under the inadmissible generic name *Tendipes*.

- nepalensis**, *id.*, *loc. cit.*, 339, ♂ ♀ (1911)
Noalpur, Nepal, 23-ii-08; Sicktan, 13-ii-08; Simla, 11-v-08
[Annandale]. *Types* belonging to Indian Museum.
- macroscelus**, *id.*, *loc. cit.*, 340, ♀ (1911).
Dahawangahary Hill, Nepal, 16-ii-08. *Type* belonging to
Indian Museum.
- stictogaster**, *id.*, *loc. cit.*, 341, ♂ (1911).
Simla, 12-v-08 [Annandale]. *Type* belonging to Indian
Museum.
- rufescens**, *id.*, *loc. cit.*, 341, ♂ (1911).
Chandan Chowki, United Prov., India, 8-v-07. *Type*
belonging to Indian Museum.
- sessilis**, *id.*, *loc. cit.*, 342, ♀ (1911).
Thamaspur, Nepal, 18-ii-08. *Type* belonging to Indian
Museum.
- pretiosus**, Kieff., Spol. Zeyl. VIII, 11, ♂ ♀, fig. 3, ♂ genitalia
(1912).
Peradeniya, 2-vi-10, *type* ♂; 31-v-10, *type* ♀; 21-v; 11-vi-10.
Types in Indian Museum.
- sumptuosus**, *id.*, *loc. cit.*, 12, ♀ (1912).
Peradeniya, 2-vi-10. *Type* in Indian Museum, a unique ♀.
- elatus**, *id.*, *loc. cit.*, 12, ♀ 1912.
Peradeniya, 2-vi-10. *Type* in Indian Museum, a unique
specimen.
- superbus**, *id.*, *loc. cit.*, 13, ♂ ♀, fig. 4, ♂ genitalia, part (1912).
Peradeniya, 14-29-v; 2-vi-10. *Paratypes* in Indian Mu-
seum.
- gloriosus**, *id.*, *loc. cit.*, 14, ♀ (1912).
Peradeniya, 20-v-10, at light. *Type* in Indian Museum,
a unique specimen.
- allothrix**, *id.*, *loc. cit.*, 15, ♀, fig. 5, antenna (1912).
Peradeniya, 25-v-10, at light. *Type* in Indian Museum, a
unique specimen.
- perichlorus**, *id.*, *loc. cit.*, 16, ♀ (1912).
Peradeniya, 17-iv-10. *Type* in Indian Museum, a unique ♀.
- chlorogaster** (II), *id.*, *loc. cit.*, 16, ♀ (1912).
Peradeniya, 17-v-10. *Type*, a unique ♀, in Indian Museum.
Kieffer has previously described a *Chironomus chlorogaster*
♂ (*vide ante*, p. 77). He calls the present form a new species.
Chlorogaster ♂ is too damaged to be of much use, but the
two specimens apparently represent different species.
- heptatomus**, *id.*, *loc. cit.*, 17, ♀ (1912).
Peradeniya, 2-vi-10. Two *paratypes* in Indian Museum.
- hexatomus**, *id.*, *loc. cit.*, 18, ♀ (1912).
Peradeniya, 2-vi-10. *Type* in Indian Museum.

- variicornis*, *id.*, *loc. cit.*, 18, ♀ (1912).
Peradeniya, 22-v-10. Type in Indian Museum, a unique specimen.
- formosae* Kieff., *Supp. Ent. Berl.* I, 35 (1912).
Formosa.
- sordidus*, *id.*, *loc. cit.*, 36 (1912).
Formosa.
- tainanus*, *id.*, *loc. cit.*, 36 (1912).
Formosa.
- formosicula*, *id.*, *loc. cit.*, 37 (1912).
Formosa.
- prasinellus*, *id.*, *loc. cit.*, 37 (1912).
Formosa.
- chlorophorus*, *id.*, *loc. cit.*, 38 (1912).
Formosa.
- chlorostolus*, *id.*, *loc. cit.*, 38 (1912).
Formosa.
- pelochloris*, *id.*, *loc. cit.*, 39 (1912).
Formosa.
- glauciventris*, *id.*, *loc. cit.*, 40 (1912).
Formosa.
- curtitarsis* Kieff., *Rec. Ind. Mus.* IX, 132, ♀ (1913).
Sara Ghat, Ganges Riv., N. Bengal, 23-iii-10, at light [*Annandale*]. Type in Indian Museum, a unique specimen.
- longipalpis*, *id.*, *loc. cit.*, 132, ♂, pl. xi, 4, genitalia, pt. (1913).
Port Canning, 24-xii-07 [*Annandale*]. Type in Indian Museum, a unique specimen.
- brunettii*, *id.*, *loc. cit.*, 133, ♀ (1913).
Madhupur, Bengal, 15—16-x-09 [*Paiva*]. Three paratypes in Indian Museum.
- ornatissimus*, *id.*, *loc. cit.*, 133, ♂ ♀ (1913).
Calcutta, 14-xii-07, type ♂: 11-i-08, type ♀ [both "at light," and *Annandale*]. Types in Indian Museum, a unique pair.
- anomalipes*, *id.*, *loc. cit.*, 134, ♀ (1913).
Bhogaon, 18-x-07 [*Paiva*]. Type in Indian Museum, a unique specimen.
- nitidus*, *id.*, *loc. cit.*, 134, ♂, pl. xi, 5, genitalia, pt. (1913).
Calcutta, 26-viii-07. Type in Indian Museum, a unique specimen.
There is a *C. nitidus* Mg., from Europe.
- heterocerus*, *id.*, *loc. cit.*, 135, ♀, pl. xi, 6, antenna (1913).
Bettiah, Champaran, Bengal, 4-iii; Calcutta, 13, 30-i; 4-ii; 4-ix. Paratypes in Indian Museum.
- glabripes*, *id.*, *loc. cit.*, 135, ♀ (1913).
Calcutta, 17-xii-07. Type in Indian Museum, a unique specimen.

- tenuiforceps*, *id.*, *loc. cit.*, 136, ♂ ♀, pl. xi, 7, genitalia, pt. (1913).
Calcutta, ix, x, xi. *Paratypes* in Indian Museum.
- atrophus*, *id.*, *loc. cit.*, 136, ♂, pl. xi, 8, genitalia, pt. (1913).
Calcutta, 11-viii-07. *Type* in Indian Museum, a unique specimen.
- orissae*, *id.*, *loc. cit.*, 137, ♂ (1913).
Puri, 2-iii-08 [*Paiva*]. *Type* in Indian Museum, a unique specimen.
- armatifrons*, *id.*, *loc. cit.*, 137, ♂ (1913).
Mandalay, 12-iii-08 [*Annandale*]. *Type* in Indian Museum, a unique specimen.
- conifer*, *id.*, *loc. cit.*, 137, ♂, pl. xi, 9, genitalia tip (1913).
Mandalay, 13-iii-08 [*Annandale*]. *Type* in Indian Museum, a unique ♂; abdomen now missing.
- erythropterus*, *id.*, *loc. cit.*, 138, ♀ (1913).
Between Bolpore and Rampore Haut, Bengal, at light, in railway carriage, 3-viii-07 [*Paiva*]. *Type* in Indian Museum, a unique specimen.
- glabrimanus*, *id.*, *loc. cit.*, 138, ♀ (1913).
In company with *erythropterus*. *Type*, a unique ♀, in Indian Museum.
- bipunctatus*, *id.*, *loc. cit.*, 139, ♀ (1913).
Rangoon, ii [*Annandale*]. *Type* in Indian Museum.
- perileucus*, *id.*, *loc. cit.*, x, 314, ♂ ♀ (1914).
Calcutta (Indian Museum gardens), 10-16-vii. *Type* in Indian Museum.

Tripelina, Kieff.

Rec. Ind. Mus. IX, 128 (1913).

GENOTYPE, *T. pallidum* Kieff., the original species.

- pallidum* Kieff., *loc. cit.*, 128, ♂ ♀ (1913).
Calcutta, 9-xi; 23-xi. at light [*Annandale*]. *Paratypes* in Indian Museum.

Family PSYCHODIDAE.¹

Subfamily PSYCHODINAE.

Psychoda, Latr.

Précis gen. Ins. 152 (1796).

Sch. Fauna Austr. II, 635: Brun., Rec. Ind. Mus. II, 369, notes; iv, 289 (tab. Ind. spp.); 291, notes: *id.*, Fauna Brit. Ind. Dipt. I, 218.

¹ For general family notes see Brun., Rec. Ind. Mus. IV, 289, and Fauna, 196.

Trichoptera Mg., Illig. Mag. II, 261 (1803).

Tinearia Schlübrig. Gen. des mouches (1803), pl.

40, fig. 1, a, b, c, fig. 2, b, d.

GENOTYPE, *Tipula phalaenoides*, L.

bengalensis Brun., Rec. Ind. Mus. II, 371, ♂ ♀, pl. xxiv, 1, wing, 2, antenna, 3, ♂ genitalia (1908).

Id., loc. cit., IV, 295 (localities): *id.*, Fauna Brit. Ind. 229, ♂ ♀, pl. iv, 6, antenna, 8, genitalia ♂.

India, common in many localities for the greater part of the year, E. and W. Himalayas up to 8000 ft.; Peradeniya, 8-vi-10; Moulmein, 25-ii-08; Tenasserim, iii; Rangoon iii. *Types* in Indian Museum.

distincta, *id.*, Rec. Ind. Mus. II, 372, ♀, pl. xxiv, 4, wing, 5, antenna (1908).

Id., loc. cit., IV, 298: *id.*, Fauna Brit. Ind. 225, ♀, pl. iv, 12, wing, 13, antenna.

Sylhet, 1-i-05 [Hall]. *Type* in Indian Museum.

albonotata, *id.*, Rec. Ind. Mus. II, 373, ♀, pl. xxiv, 7, wing (1908).

Id., loc. cit., IV, 298: *id.*, Fauna Brit. Ind. 224, ♀.

Calcutta, 5-6-i-08 [Annandale], vii; Sylhet, January [Lt.-Col. Hall]. *Type*, a unique specimen, in Indian Museum.

albonigra, *id.*, Rec. Ind. Mus. II, 374, ? ♀ (1908).

Id., loc. cit., iv, 295: *id.*, Fauna Brit. Ind. 228.

Calcutta [Annandale]. *Type* in Indian Museum.

nigripennis, *id.*, Rec. Ind. Mus. II, 376, ♂ ♀, pl. xxiv, 8, antenna (1908).

Id., loc. cit., IV, 297 (localities): *id.*, Fauna Brit. Ind., 232, ♂ ♀, pl. iv, 16, flagellum.

India, common from April to August in many Indian localities, hills and plains; Travancore, xi; E. and W. Himalayas up to 9000 ft., the next commonest species after *bengalensis*. *Types* in Indian Museum.

vittata, *id.*, Rec. Ind. Mus. II, 377, ♀ (1908).

Id., loc. cit., IV, 298, ♂: *id.*, Fauna Brit. Ind., 233, ♂ ♀.

Calcutta, 11-26-vii-08; 16-xi-08; 11-vi to 2-vii-08 [Annandale]; Maddathorai, Travancore, 18-xi-08, *type* ♂. *Type* in Indian Museum.

geniculata, *id.*, Rec. Ind. Mus. IV, 294, ♀ (1911).

Id., Fauna Brit. Ind., 227, ♀.

Peradeniya, x-10 [Green]. *Type*, a unique ♀, in Indian Museum.

distans, *id.*, Rec. Ind. Mus. IV, 296, ♀ (1911).

Id., Fauna Brit. Ind., 230, ♀.

Maddathorai, Travancore, 16-xi-08 [Annandale]. *Type*, a unique ♀, in Indian Museum.

albopicta, *id.*, Rec. Ind. Mus. IV, 296, ♀ (1911).

Id., Fauna Brit. Ind., 231, ♀.

Pusa, 8-i; Ceylon. *Type*, a unique ♀, in Pusa collection.

- fulvohirta*, *id.*, Rec. Ind. Mus. IV, 297, ♀ (1911).
Id., Fauna Brit. Ind., 233, ♀.
 Darjiling, 7-viii-09, *type* [*Paiva*]; Kurseong, 26-vi-10 [*Annandale*]. *Type* in Indian Museum.
- orbicularis*, *id.*, Rec. Ind. Mus. IV, 298, ♀ (1911).
Id., Fauna Brit. Ind., 231, ♀.
 Pusa, 21-xii-08. *Type*, a unique ♀, in Pusa collection.
- decora* Brun., Rec. Ind. Mus. IV, 299, ♀ (1911).
Id., Fauna Brit. Ind., 226, ♀.
 Tenmalai (Travancore), 22-xi-08 [*Annandale*]. *Type*, unique, in Indian Museum.
- maculipennis* Brun., Rec. Ind. Mus. IV, 299, ? ♂ (1911).
Id., Fauna Brit. Ind., 223, ? ♂.
 Peradeniya, 17-vi-10 [*Green*]. *Type*, a unique specimen, in Indian Museum.
- transversa* Brun., Rec. Ind. Mus. IV, 300, ? ♀ (1911).
Id., Fauna Brit. Ind., 226, ? ♀.
 Kurseong, 20-vi-10 [*Annandale*]. *Type* in Indian Museum, a unique specimen.
- hirtipennis* Brun., Rec. Ind. Mus. IV, 300, ♀ (1911).
Id., Fauna Brit. Ind., 227, ♀.
 Kurseong, 5-ix-09 (*type*); Bangalore, 15-x-10; Maddathorai, 18-xi-08; Trivandrum, 12—13-xi-08 [last three *Annandale*]; Calcutta Botanical Gardens, 27-xi-10 [*Gravelly*]. *Type* in Indian Museum.
- apicalis* Brun., Rec. Ind. Mus. IV, 301, ? ♀ (1911).
Id., Fauna Brit. Ind., 223, ? ♀.
 Maddathorai, 16-xi-08 [*Annandale*]. *Type*, apparently a female, in Indian Museum, a unique specimen.
- notatipennis* Brun., Rec. Ind. Mus. VIII, 151, pl. vi, 4, wing (1913).
 N.E. Indian Frontier. *Type*, sex uncertain, a unique specimen, in Indian Museum.

Pericoma, Walk.

- Ins. Brit. Dipt. 256 (1856).
 Sch. F. Austr. II, 632: Brun., Rec. Ind. Mus. IV, 301, tab. orient. spp.: *id.*, Fauna Brit. Ind. I, 234.
 GENOTYPE, *Trichoptera fasciata* Mg., by Coquillett's designation, 1910.
- spinicornis* Brun., Rec. Ind. Mus. II, 378, ♂; pl. xxiv, 9, antennal joint, 10, genitalia ♂, 11, wing (*appendiculata*) (1908).
Id., *loc. cit.*, iv, 304, locs.: *id.*, Fauna Brit. Ind., 238, ♂ ♀; pl. iv, 4, base of antenna, 15, wing.
appendiculata, Brun., Rec. Ind. Mus. II, 379 (1908).

E. and W. Himalayas, not uncommon from early May to early October in many localities. Siliguri, base of Darjiling Hills, 18—20-vii-07 [*Hodgart*]. *Types* in Indian Museum.

annandalei Brun., Rec. Ind. Mus. II, 380, ♀ (1908).

Id., loc. cit., 304: *id.*, Fauna Brit. Ind., 237, ♀.

Kurseong, 5-vii-08, *type*: 22—29-vi-10 [both *Annandale*]. *Type* in Indian Museum.

margininotata Brun., Rec. Ind. Mus. II, 381, ♂ ♀; pl. xxiv, 12, wing (1908).

Id., loc. cit., iv, 304: *id.*, Fauna Brit. Ind., 240, ♂ ♀.

P. bella, Brun., Rec. Ind. Mus. II, 383, ♀ (1908).

P. margininotata, var. *bella*: *id.*, Fauna Brit. Ind., 241, ♀.

Simla, 25-iv-07; 11-v-08; 9—10-v-09; Phagu, 11—15-v-09 [all *Annandale*]; Darjiling, 26-v-10, end of Sept. and Oct. 1, 08 (including *type* of *bella*) [both *Brunetti*]; Kurseong, 25—27-vi-10 [*Annandale*]; Pallode, Travancore, 15-xi-08 [*Annandale*]. *Types* of *margininotata* (typical form) and *bella* var. in Indian Museum.

lacteitarsis Brun., Rec. Ind. Mus. II, 382, ♀ (*margininotata*, var. *lacteitarsis*), (1908).

lacteitarsis, *id.*, loc. cit., IV, 307: *id.*, Fauna Brit. Ind., 245.

Kurseong, 4-vii-08, *type*; Quilon, Travancore, 9-xi-08 [both *Annandale*]. *Type* in Indian Museum.

gilvipes Brun., Rec. Ind. Mus. II, 382, ♀ (*margininotata*, var. *gilvipes*), (1908).

gilvipes, *id.*, loc. cit., iv, 308, notes: Fauna Brit. Ind., 246, ♀.

Calcutta, 28-vii-08, *type*, 2-viii-08, 17—18-vii-08, 9-ii-10, 8—23-viii-08, 1—26-ix-08 [all *Annandale*]; Madhupur, Bengal, 17-x-09, at light [*Paiva*]; Ernakulam, Cochin, 4-xi-08; Quilon, 9-xi-08 [both *Annandale*]. *Type* in Indian Museum.

squaminnervis Brun., Rec. Ind. Mus. IV, 303, ? ♀ (1911).

Id., Fauna Brit. Ind., 237, ? ♀.

Kurseong, 4-viii-08 [*Annandale*]. Unique *type* in Indian Museum.

metatarsalis Brun., Rec. Ind. Mus. IV, 305, ♀ (1911).

Id., Fauna Brit. Ind., 243, ♀.

Simla, 9-v-09, *type*; 12-v-09; Phagu, 11-v-09 [all *Annandale*]. *Type* in Indian Museum.

mixta Brun., Rec. Ind. Mus. IV, 306, ♀ (1911).

Id., Fauna Brit. Ind., 244, ♀.

Simla, 6-v-09 [*Annandale*]. Unique *type* in Indian Museum.

proxima Brun., Rec. Ind. Mus. IV, 308, ? ♂ (1911).

Id., Fauna Brit. Ind., 245, ? ♂.

Peradeniya, viii-1910 [*Green*]. Unique *type* in Indian Museum.

impunctata Brun., Rec. Ind. Mus. IV, 309 (1911).

- Id.*, Fauna Brit. Ind., 247.
 Tenmalai, Travancore, 22 xi-08 [Annandale]. The unique
type in Indian Museum, of uncertain sex.
- unicolor** Brun., Rec. Ind. Mus. IV, 309 (1911).
Id., Fauna Brit. Ind., 247.
 Kurseong, xi-1910 [D'Abreu]. Unique *type*, of uncertain
 sex, in Indian Museum.
- indica** Eaton., Trans. Linn. Soc. Lond. XV, 424, figs. (1913).
 India, also the Seychelles.

Brunettia, Annand.

- Rec. Ind. Mus. V, 141 (1910).
 Brun., Rec. Ind. Mus. IV, 310: *id.*, Fauna Brit.
 Ind., 248.
Diplonema, Annand. (*nec* Loew.), Rec. Ind. Mus.
 IV, 39 (1910).
- superstes** Annand., Jour. Asiat. Soc. Beng. (new ser.) IV, 353, ♂ ♀ ;
 pl. xii, var. chars. [*Diplonema*] (1908).
Id., Rec. Ind. Mus. V, 142, ♂ ♀ , redescr: fig. genitalia ♂ ;
 pl. xii, var. chars. [*Brunettia*] (1910).
 Brun. Fauna Brit. Ind., 249, ♂ ♀ , fig. 34 (♂ genitalia), 35
 venation, pl. iv, 1, full insect, 3 head.
 Kurseong, vi-vii [Annandale]. *Types* in Indian Museum.
- indica** Eaton, Trans. Linn. Soc. Lond. XV, 424, ♂ ♀ ; pl. xxvi,
 fig. 2, a-f; fig. 3, a-f (1913).
 Matheran, Western Ghats, Bombay, 800 metres, 8-vii-02 [Biro].

Parabrunettia, Brun.

- Rec. Ind. Mus. IV, 310, 311, tab. spp. (1911).
 Brun. Fauna Brit. Ind., 251.
 GENOTYPE, *Psychoda squamipennis*, Brun., by
 original designation.
- squamipennis** Brun., Rec. Ind. Mus. II, 375, ♀ (*Psychoda*), (1908).
Id., Fauna Brit. Ind., 252, ♀ .
 Calcutta, 5-viii-07 [Annandale]. Unique *type* in Indian
 Museum.
- argenteopunctata** Brun., Rec. Ind. Mus. II, 375, ♀ ; pl. xxiv, 6,
 wing (*Psychoda*), (1908).
Id., Fauna Brit. Ind., 254, ♀ .
 Calcutta, 27-viii-07. Unique *type* in Indian Museum.
- atrisquamis** Brun., Rec. Ind. Mus. II, 376, ♀ (*Psychoda*), (1908).
Parabrunettia atrisquamis, *id.*, *loc. cit.*, iv, 312: Fauna Brit. Ind.,
 253, fig. 36, wing, pl. iv, 2, full insect.
Brunettia travancorica, Annand., Rec. Ind. Mus. V, 144, ♀ ; pl.
 xii, var. chars. (1910).
 Calcutta, 22-vii-08, *type* ♀ [Annandale]; Peradeniya, 18-vii,
 14-viii and x-1910 [Green]; base of Western Ghats, Travancore,

xi-08 (*type of travancorica*), xi-08 [*Annandale*]. *Types* in Indian Museum.

albohumeralis Brun., Rec. Ind. Mus. IV, 312, ♀ (1911).

Id., Fauna Brit. Ind., 254.

Peradeniya, ix—x-10 [*Gravelly*]. *Type* in Indian Museum.

novemnotata Brun., Rec. Ind. Mus. IV, 313, ♀ (1911).

Id., Fauna Brit. Ind. 255, ♀.

Puri, Orissa, 12-xi-10 [*Annandale*]. *Unique type* in Indian Museum.

flavicollis Brun., Rec. Ind. Mus. IV, 314, ♂ (1911).

Id., Fauna Brit. Ind., 256, ♂.

Peradeniya, viii-10. *Unique type* in Indian Museum.

longichaeta Brun., Rec. Ind. Mus. IV, 314, ♂ (1911).

Id., Fauna Brit. Ind., 256, ♂.

Maddathorai (Travancore), 17-xi-08 [*Annandale*]; Bangalore, 3000 ft., x [*Annandale*]. *Type* in Indian Museum.

Referred doubtfully and provisionally to the present genus.

Subfamily PHLEBOTOMINAE.¹

Phlebotomus, Rond.

Mem. prima ditt. Ital., 12 (*Flebotomus*), (1840).

Phlebotomus, Sch. F. Austr. II, 630: Brun., Fauna

Brit. Ind., 199; tab. Ind. spp. 203: Annand.,

Rec. Ind. Mus. IV, 43, tab. Ind. spp.: Spol.

Zeyl. VII, 57, notes, tab. spp. (incorrect): *l.c.*,

VII, 159 (corrected table): Alcock, Entom. for

Med. Officers, 119 (brief description of oriental

species).

Cyniphes, G. Costa, Ann. Accad. Aspir. Natural

I, 4 (1843).

Hebotomus, Rond., Ann. So. Ent. Fr. (2), I, 264

(1843).

Haemasson, Loew, Stett. Ent. Zeit. V, 115 (1844).

GENOTYPE, *Bibio papatasi*, Scop.

papatasi Scop., Deliciae Faun. Flor. Insul. I, 55, pl. xxii, fig.

B (*Bibio*), (1786): Sch. F. Austr. II, 630: Grassi, Mem. So.

Ital. Sci. (3) XIV, 353 (copious descriptions and figs., also life-

history) (1907): Annand., Rec. Ind. Mus. IV, 51, ♂ ♀ redescr.,

pl. iv, 4, wing; pl. vi, 2, ♂ genitalia: Newstead, Bull. Ent.

Res. II, 76; Brun., Fauna Brit. Ind. 211, ♂ ♀, figs. 26—31,

var. chars.: Summers, Jour. Lond. Sch. Trop. Med. II, 105,

fig. ♂, genitalia.

Flebotomus papatasi, Rond., Mem. Ditt. Ital., 13, pl. (1840).

¹ For subfamily characters see Brun., Rec. Ind. Mus. IV, 289 and Fauna Brit. Ind. 199.

Hebotomus, *id.*, *id.*, Ann. So. Ent. Fran. (2) I, 265, pl. x, 1—4 (1843).

Cyniphes molestus, Costa, Ann. Accad. Aspir. Nat. I, 4 (1843).

Haemasson minutus, Loew (*nec* Rond.), Stett. Ent. Zeit. V, 115, pl. i, 1—5 (1844).

? *Phlebotomus angustipennis*, de Meij. Tijd. v. Ent., LII, 202, pl. xii, 14, wing (veins incorrectly numbered), (1909).

Phlebotomus sp., Howlett, in Lefroy's Indian Insect Life, 560, fig. 359, ♂ genitalia (1909).

Rawalpindi, Chitral, 6000 ft., Quetta, Miranshad, N.W. Frontier [*Walton*]; Punjab; Poona, Kirkee (near Bombay), viii-07; Lahore; North Bengal, common in April; Pusa. It also occurs in South Europe, Palestine, Egypt and elsewhere. *Type*, Prof. Bezzi informs me this is no longer in existence. That of *angustipennis* is in the Amsterdam Museum.

minutus Rond., Ann. So. Ent. Fran. (2), I, 263 (1843).

Brun., Fauna Brit. Ind., 206, ♂ ♀ (1912): Alcock, Jour. Lond.

Sch. Trop. Med. I, 205 (♀ dimorphism): Summers, *loc. cit.*

II, 106: Howlett, Ind. Jour. Med. Res. I, 34.

babu, Annand., Rec. Ind. Mus. IV, 49, ♂ ♀, pl. iv, 1, wing, pl. vi, 3, 3a, ♂, genitalia (1910).

babu var. *niger*, Annand., Rec. Ind. Mus. IV, 320 (1911): Summers, Jour. Lond. Sch. Trop. Med. II, 109.

Phlebotomus sp., Howlett, in Lefroy's Indian Insect Life, 559, fig. 358 (1909).

India, "probably all over the plains of India" [*Annandale*]; Rawalpindi, September [*C. W. Mason*]; Quetta, Chitral, Allahabad, October [*Imms*]; Rajmahal, on the Ganges, July; Asansol, Purneah, Pusa [*Howlett*]; Madhupur, Puri, Calcutta (all the year); Port Canning, Igatpuri, Bombay (Western Ghats), November; Poona, Kirkee, Rambha (N.E. Madras); Trivandrum, Pallode and Maddathorai (all in Travancore) in November; Peradeniya. It also occurs in South Europe. *Type* in Florence Museum; *types* of *babu*, ♂ ♀, in Indian Museum.

argentipes Annand. and Brun., Rec. Ind. Mus. II, 101, ♂ ♀, fig. wing (veins incorrectly numbered), (1908)

Annand., *loc. cit.*, IV, 44, ♂ ♀, redescr. fig. 4 (p. 42), ♂ genitalia, pt; pl. iv, 3, wing; pl. vi, 6, ♂ genitalia: Brun., Fauna Brit. Ind., 207, ♂ ♀, pl. iv, 5, ♂ genitalia: Summers, Jour. Lond. Sch. Trop. Med. II, 108.

P. marginatus, Annand., Spol. Zeyl. VII, pt. xxvi, 62, ♀, fig. 7, wing (1910).

P. argentipes var. *marginatus*, Brun., Fauna Brit. Ind., 209, ♀: Summers, Jour. Lond. Sch. Trop. Med. II, 109.

India, generally distributed, found in Calcutta at least all the year round; Peradeniya, iii, vi, vii. *Types* of *argentipes*, ♂ ♀ and *type* of *marginatus*, a unique ♀, in the Indian Museum.

- perturbans** de Meij., Tijds. v. Ent. LII, 201, pl. xii, 13, wing (veins incorrectly numbered), (1909).
 Annand., Rec. Ind. Mus. IV, 47, ♂ ♀, redescr. pl. v, 3, wing; pl. vi, 5, 5a, 5b, ♂ genitalia (1910): Brun., Fauna Brit. Ind., 205, ♂ ♀, pl. iv, 11, ♂ genitalia: Summers, Jour. Lond. Sch. Trop. Med. II, 112.
 Java; Sukna (base of Darjiling Hills), 1-3-vii, common "at light" [Annandale]. Type in Amsterdam Museum.
- major** Annand., Rec. Ind. Mus. IV, 46, ♂, pl. v, 4, wing; pl. vi, 4, ♂ genitalia (1910): Brun., Fauna Brit. Ind., 210, ♂, pl. iv, 7, ♂ genitalia: Summers, Jour. Lond. Sch. Trop. Med., II, 110.
Phlebotomus sp., Giles, Gnats or Mosquitoes (2nd ed.), 5, fig. 2 (1902).
 East and West Himalayas from base up to 8000 ft.; Naini Tal, vii; Simla vii; Kurseong, iv to vii, base of Nepalese Himalayas; Paresnath, W. Bengal, iv. Type in Indian Museum.
- var. *griseus*, Annand., Rec. Ind. Mus. IV, 320 (*grisea*), (1911).
 Brun., Fauna Brit. Ind., 211: Summers, Jour. Lond. Sch. Trop. Med. II, iii.
 Kurseong, vi-1910 [Annandale]. Type in Indian Museum.
- var. *perniciosus*, Newstead, Bull. Ent. Res. II, 70, ♂ ♀ (as a good species) (1911).
 Brun., Fauna Brit. Ind., 211, ♂ ♀: Summers, Jour. Lond. Sch. Trop. Med., II, 107. This variety is recorded from Malta.
- malabaricus** Annand., Rec. Ind. Mus. IV, 48, ♂ ♀, pl. v, 1, head, 2, wing; pl. v, 1, ♂ genitalia (1910).
 Brun., Fauna Brit. Ind., 214, ♂ ♀: Summers, Jour. Lond. Sch. Trop. Med. II, 111.
 Travancore (var. locs.), "at light", xi-08 [Annandale]. Types in Indian Museum.
- himalayensis** Annand., Rec. Ind. Mus. IV, 50, ♂ ♀, pl. iv, 2, wing; pl. vi, 7, ♂ genitalia (1910).
 Brun., Fauna Brit. Ind., 204, ♂ ♀, pl. iv, 10, ♂ genitalia: Summers, Jour. Lond. Sch. Trop. Med. II, 109.
 East and West Himalayas, 4000 to 7000 ft.; Naini Tal (Bhowali); Kurseong, common at both places, May to July.
- zeylanicus** Annand., Spol. Zeyl. VII, 60; ♂ ♀, fig. 4, full insect, ♀, 5, wing, 6, ♂ genitalia (1910).
 Brun., Fauna Brit. Ind., 215, ♂ ♀, fig. 32, full insect, pl. iv, 14; Summers, Jour. Lond. Sch. Trop. Med. II, 112: Newstead, Bull. Ent. Res. V, 187.
 Peradeniya, v, vi, vii, viii. Types in Indian Museum.
- stantoni** Newstead, Bull. Ent. Res. V, 109, ♀ (1914-1915).
 Kuala Lumpur, 15-vi-14 [Dr. Stanton], a unique ♀.

CULICIDAE.

The subdivision and arrangement of genera and species in this family has been a source of much difficulty. In the Anophelini there has been no hesitation in referring the bulk of the species to *Anopheles* itself, as is now done by most recent writers, the only other groups accorded generic rank here being *Calvertina* and *Bironella*. *Chagasia*, a non-oriental genus, was the only other one regarded as valid in the Anophelini in my "Critical Review." The classification, however, of the rest of the Culicinae is still in a very transitory state, and some attempt has been made to meet the views of others. A certain number of the recently established genera are evidently valid, others appear entitled to subgeneric rank, whilst the remainder must be regarded as of indefinite taxonomic value at present. When a genus with all its species has been bodily sunk in another by Mr. Edwards, the synonymy has been at once accepted, as the sooner invalid generic names are permanently shelved the better. No doubt a good many other genera will be weeded out as synonymous when the oriental species have been thoroughly overhauled by systematists.

In the present catalogue the subfamily Culicinae is divided merely into Anophelini and Culicini, as though the group Megarhini is apparently distinct enough, it is more than can be said for the groups or sections dominated by such genera as *Aedes*, *Culex*, *Sabethes*, etc., so that while the juggling of genera and higher groups still continues even amongst those best fitted to dictate,¹ no harm will be done by lumping them all together under the general term Culicini without assigning it any definite taxonomic value.

The accidental omission of the heading "Subfamily Megarhininae" in both my Catalogue of Culicidae and the Supplement gave the impression that the genera concerned were to be considered Anophelines, which of course was never intended.

Theobald's sequence of genera has been mainly followed because that author's work is known to all students in this family, and also because of personal inability to dogmatise.

As regards specific synonymy it has frequently been a case of "when doctors disagree" both metaphorically and actually, and in such instances I have usually followed the latest writer, or whichever has had any purely entomological support.

The wholesale overthrowing of a large proportion of the recently manufactured genera and species was early foretold by other systematists as well as by me, and the truth of the prophecy is visible in their continual reduction in almost every paper now published. It is to be hoped that the new species erected nowadays will prove to be founded on more reliable characters.

As regards location of types in this family considerable difficulty has been experienced. Respecting Leicester's species, Mr.

¹ There are, for instance, several genera which Mr. Edwards regards as anomalous.

Edwards wrote me that "types or typical specimens" of nearly all his species were in the collection that he presented to the British Museum, also that Dr. Leicester had not made it a rule to designate individual specimens as types. The types of Theobald's species, except when otherwise stated, are presumably in the British Museum also.

Subfamily CULICINAE.

GROUPS IN CULICIDAE HIGHER THAN GENERA.¹

Anophelinae.

Theob., Gen. Ins., Fasc., 26, 5: *id.*, Monog. Culic., IV, 18; tab. gen., 22; V, 1, tab. gen.: Blanch., Moust., 157, tab. gen., 159; Leices., Culic. Mal., 18, and tab. gen.: Brun., Rec. Ind. Mus. I, 302; IV, 414.

Anophelini.

Brun., Rec. Ind. Mus. X, 32, notes, tab. gen.

Anophelina.

Theob., Monog. Culic. I, 97; III, 9: Giles, Handbk., etc., 2nd ed., 280.

Anophelines.

How., Dyar and Knab, Mosq. N. Amer. III, pt. I, 193.

Epialurgi.

Alcock, Ent. Med. Off. Ind. 64.

Culicinae.

Theob., Gen. Ins., Fasc. 26, 26, 30: *id.*, Monog. Culic. IV, 19, 147 and tab. gen.; V, 112 and tab. gen.: Blanch., Moust., 231: tab. gen., 152, 390: Giles, Handbk., etc., 2nd ed., 334, tab. gen., Leices., Culic. Mal., 64: Alcock, Ent. Med. Off. Ind., 94: Brun., Rec. Ind. Mus. I, 326; IV, 439: Patton and Cragg, Text-bk. Med. Entom., 192: Edw., Bull. Ent. Res. II, 241, tab. African gen.

Culicini.

Brun., Rec. Ind. Mus. X, 35, notes and tab. gen.: How., Dyar and Knab, Mosq. N. Amer. III, pt. 1, 189, tab. N. Amer. gen. 194: IV, pt. 2, 525.

¹ For the convenience of those who may desire it, this short list is given of the various sections into which the mosquitoes have been divided by recent writers. Only the principal sections and references are given, as they form no integral part of the present catalogue. In Howard, Dyar and Knab's monumental work the fullest particulars can be obtained. No attempt at systematic sequence is made in listing them as the great majority of the groups will probably be entirely ignored in the early future.

Culicina.

Theob., Monog. Culic. I, 97, 247; III, 124; Giles, Handbk., etc., 2nd ed., 333, tab. gen., 334.

Culicines.

How., Dyar and Knab, Mosq. N. Amer. III, pt. I, 192.

Culicales.

Alcock, Ent. Med. Off. Ind., 94.

Megarhininae.

Theob., Gen. Ins., Fasc., 26, 12; *id.*, Monog. Culic. IV, 19, 126 (*Ankylorhynchae*, Lutz. and *Lynchiellina*, Lahille, given as synonymous); V, 88; Blanch., Moust., 218, tab. gen. 221; Leices., Culic. Mal., 48; Brun., Rec. Ind. Mus. IV, 434.

Megarhinina.

Theob., Monog. Culic. I, 97, 215; III, 113; Giles, Handbk., etc., 2nd ed., 265.

Megarhini.

Brun., Rec. Ind. Mus. X, 34, notes and tab. gen.

Megarhines.

How., Dyar and Knab, Mosq. N. Amer. III, pt. I, 193.

Megalorrhini.

Alcock, Ent. Med. Off. Ind., 113.

Toxorhynchitinae.

Theob., Gen. Ins., Fasc., 26, 13.

Joblotinae.

Theob., Gen. Ins., Fasc., 26, 33; Blanch., Moust. 428 and tab. gen.

Joblotina.

Theob., Monog. Culic., III, 334.

Aedeomyinae.

Theob., Gen. Ins., Fasc. 26, 33, tab. gen., 34; Giles, Handbk., etc., 2nd ed., 475; Blanch., Moust., 153, tab. gen.; Leices., Culic. Mal., 180 and tab. Malay gen.; Brun., Rec. Ind. Mus. I, 362; IV, 487.

Aedeomyina.

Theob., Monog. Culic. I, 98; II, 218; III, 275.

Aedeinae.

Blanch., Moust., 399, tab. gen.

Aedinae.

Theob., Monog. Culic. IV, 19, 537; V, 482.

Aedomina.

Giles, Handbk., etc., 2nd ed., 475.

Uranotaeniae.¹

Theob., Monog. Culic. IV, 19, 556; V, 497.

Sabethini.

How., Dyar and Knab, Mosq. N. Amer. IV, pt. 1, 19: Brun., Rec. Ind. Mus. X, 48, notes and tab. gen.

Sabettinae, Blanch.

Theob., Monog. Culic. V, 574: Blanch., Moust., 420 and tab. gen.

Wyeomyinae.

Leices., Culic. Mal., 250.

Metanototricha.

Alcock, Ent. Med. Off. Ind., 114.

Trichoprosoponina.Theob., Monog. Culic. I, 98; II, 283; IV, 585 (*Trichoprosoponinae*); V, 554.**Dendromyinae, Lutz.**

Theob., Monog. Culic. IV, 20, 594; V, 574.

Limatinae.

Theob., Monog. Culic. IV, 627; V, 595.

Deinoceratinae.²

Theob., Monog. Culic. V, 553.

Deinoceratines.

How., Dyar and Knab, Mosq. N. Amer. III, pt. 1, 191.

Heptaphloeomyinae (*Heptaphloeomyina*).Theob., Gen. Ins., Fasc., 26, 41: *id.*, Monog. Culic. III, 336; IV, 531; V, 481: Blanch., Moust., 398.¹ This genus is attributed to Mitchell by Theobald, but he gives no reference.² The name *Deinoceratinae* is attributed by Theobald to Mitchell without giving any reference.

Sect. ANOPHELINI.

Anopheles, Mg.¹

Mg., Syst. Besch. I, 10 (1818).

Sch. F. Austr. II.

GENOTYPE, *A. maculipennis* Mg., by general European acceptance.*Conchyliastes*, Theob., in Howard's "Mosquitoes," 235 (1901).GENOTYPE, *Culex posticatus* W., as *musicus* Say, by Coquillett's designation.*Cycloleppteron*, Theob., Monog. Culic. II, 312 (1901).*Id.*, *id.*, Jour. Trop. Med. IV, 234, *nom. nud.* (1901).GENOTYPE, *Anopheles grabhami* Theob., by original designation.*Cyclolepidopteron*, Blanch. (*Cycloleppteron*, emend.), Moust., 185 (1905).*Grassia*, Theob., Jour. Trop. Med. V, 181 (1902).GENOTYPE, *Anopheles rossii* Giles, by original designation.*Myzomyia*, Blanch., Comp. rend. Soc. Biol. Paris, LIV, 795 (1902); *nom. nov.* for *Grassia*, Theob., preocc. Fisch., 1885.*Id.*, Theob., Monog. Culic. III, 12,² 24; IV, 41, 42, tab. spp.; V, 16, tab. spp.: *id.*, Gen. Ins., Fasc. 26, 7: Blanch., Moust., 177: James and List., Monog. Anoph. Mosq. Ind., 2nd ed., 40: James, Rec. Ind. Mus. IV, 98: Leices., Culic. Malaya, 23: Brun., Rec. Ind. Mus. I, 305; IV, 417; X, 66 (*mia lapsus*): Alcock, Entom. Medic. Officers, 79.*Howardia*, Theob., Jour. Trop. Med. V, 181 (1902).GENOTYPE (presumably), *Anopheles costalis* Lw.*Pyretophorus*, Blanch., Comp. rend. Soc. Biol. Paris LIV, 795 (1902); *nom. nov.* for *Howardia*, Theob., preocc. Dalla Torre, 1897 in *Insecta*.*Id.*, Theob., Monog. Culic. III, 66; IV, 63, 64, list and tab. spp.; V, 36, tab. spp.: *id.*, Gen. Ins., Fasc. 26, 8: Blanch. Moust., 186: James and List., Monog. Anoph. Mosq. Ind., 2nd ed., 41: James, Rec. Ind. Mus. IV, 99: Brun., *loc. cit.*, I, 312; IV, 423; X, 70: Leices., Culic. Malaya, 37.*Rossia*, Theob., Jour. Trop. Med. V, 181 (1902).GENOTYPE, *Anopheles sinensis* Wied., by original designation.*Myzorhynchus*, Blanch., Comp. rend. Soc. Biol. Paris LIV, 795 (1902): *nom. nov.* for *Rossia*, Theob., preocc. Owen, 1838 in *Mollusca*, also used by Bonaparte in same year.

¹ Full references to this well-known genus, *sensu lato*, are not given, as Schiner's description amply portrays the view of the older authors. These references can be gleaned from Kertész's Catalogue. For interesting descriptions of larvae of Malayan *Anopheles* see Stanton, "The Larvae of Malayan *Anopheles*," Bull. Ent. Res. VI, 159. For tables to determine Indian Anophelines see James, Paludism. No. 2, 52.

² Theobald here suggests *funesta*, Giles, as type of *Myzomyia*, which is impossible, having already selected *rossii*. Edwards has noted this.

- Id.*, Theob., Monog. Culic. III, 84, pl. v, wing scales; IV, 81, list and tab.; V, 49, tab. spp.: Theob., Gen. Ins., Fasc. 26, 9: Blanch., Moust., 190: James, Rec. Ind. Mus. IV, 102: Leices., Culic. Malaya, 29 (*Myzorhynchus*): Brun., Rec. Ind. Mus. I, 313; IV, 424; X, 66: Alcock, Entom. Medic. Officers Ind., 74.
- Laverania*, Theob., Jour. Trop. Med. V, 181 (1902).
GENOTYPE, *Anopheles argyrotarsis* R. Desv., by original designation.
- Nyssorhynchus*, Blanch., Comp. rend. Soc. Biol. Paris LIV, 795 (1902): *nom. nov.* for *Laverania*, Theob., preocc. Billett, 1895 in Protozoa; also again by Grassi and Filletti in 1900.¹
- Id.*, Theob., Monog. Culic. III, 92, pl. v, wing scales; IV, 95, 96, list and tab. spp.; V, 55, tab. spp.: Theob., Gen. Ins., Fasc. 26, 10: Blanch., Moust., 202: James and List. Anoph. Mosq. Ind., 2nd ed., 43: James, Rec. Ind. Mus. IV, 100: Leices., Culic. Malaya, 39: Brun., Rec. Ind. Mus. I, 317; IV, 428; X, 67: Alcock, Entom. Medic. Officers Ind., 85.
- Stethomyia*, Theob., Jour. Trop. Med. V, 181 (1902).
Theob., Monog. Culic. III, 62, pl. viii, wing scales; IV, 59, tab. spp.; V, 35, tab. spp.: Theob., Gen. Ins., Fasc. 26, 8: Blanch., Moust., 186: Brun., Rec. Ind. Mus. I, 312; IV, 423; X, 72.
GENOTYPE, *S. nimba* Theob., by original designation.
- Cellia*, Theob., Monog. Culic. III, 107 (1903).
Theob., *loc. cit.*, IV, 105, list and tab. spp.; V, 67, tab. spp.: Theob., Gen. Ins., Fasc. 26, 11: Blanch., Moust., 214: James, Rec. Ind. Mus. IV, 102: Leices., Culic. Malaya, 46: Brun., Rec. Ind. Mus. I, 321; IV, 431; X, 57: Alcock, Entom. Medic. Officers Ind., 90.
GENOTYPE, *Anopheles pharoensis* Theob.²
- Aldrichia*, Theob., Monog. Culic. III, App., 353 (1903).
Theob., Gen. Ins., Fasc. 26, 11: Blanch., Moust., 624: Brun., Rec. Ind. Mus. I, 322; X, 55.
GENOTYPE, *Aldrichia error* Theob., sp. nov., by original designation.
- Arribalzagia*, Theob., Monog. Culic. III, 81 (1903).
GENOTYPE, *A. maculipes* Theob., sp. nov., by original designation.
Alcock, Entom. Medic. Officers Ind., 77.
- Christya*, Theob., Rep. Sleep. Sick. Royal Soc. VII, p. 34 (1903).
Giles, Revis. Anophelinae, 40 (*Christya*) (1904): Alcock, Entom. Medic. Officers Ind., 77.

¹ Blanchard suggested *Anopheles albimanus* W., as type species, but as his name is only a *nomen novum* the original genotype must stand. Even Theobald himself (Monog. III, 14) suggested *maculatus* Theob., as type species, which is impossible after originally selecting *argyrotarsis*, Rob. Desv. Edwards has noted this.

² This species was apparently selected as genotype by Theobald, but his original paper is not accessible to me.

GENOTYPE, *C. implexa* Theob., sp. nov., by original designation.

Lophomyia, Giles, Jour. Trop. Med. VII, 366 (1904).

GENOTYPE, *Lophoscelomyia asiatica* Theob., sp. nov., by original designation.

Lophoscelomyia, Theob., Entom. XXXVII, 12 (1904).

Blanch., Moust., 635: Theob., Monog. Culic. IV, 91: Leices., Culic. Malaya, 21: Brun., Rec. Ind. Mus. IV, 427; X, 64: Theob., Gen. Ins., Fasc. 26, 10 (*Lophocelomyia*): Brun., Rec. Ind. Mus. I, 316 (*Lophocelomyia*): Giles, Jour. Trop. Med. VII, 366 (*Lophomyia*).

GENOTYPE, *L. asiatica* Theob., by original designation.

Kerteszia, Theob., Ann. Mus. Hung. III, 66 (1905).

Theob., Monog. Culic. IV, 117: Brun., Rec. Ind. Mus. IV, 431: X, 63.

GENOTYPE, *K. boliviensis* Theob., sp. nov., by original designation.

Nototricha, Coq., Class. Mosq. N. Amer., 12 (1906).

Theob., Monog. Culic. V, 33 (*Notonotricha*, emend.) (1910).

GENOTYPE, *Cyclolepteron mediopunctatus* Theob., by original designation.

Coelodiazesis, Dyer and Knab, Jour. N. Yk. Ent. Soc. XIV, 77 (1906).

GENOTYPE, *Anopheles barberi* Coq., by original designation.

Erected on larval characters only and therefore inadmissible.

Feltinella, Theob., Monog. Culic. IV, 56 (1907).

GENOTYPE, *F. pallidopalpi* Theob., sp. nov., by original designation.

Myzorhynchella, Theob., loc. cit., IV, 78 (1907).

GENOTYPE, *M. nigra* Theob., sp. nov., loc. cit.

Neocellia, Theob., loc. cit., IV, 111 (1907).

Theob., loc. cit., V, 73, tab. spp.: James, Rec. Ind. Mus. IV, 101: Brun., loc. cit., IV, 432; X, 66: Alcock, Entom. Medic. Officers Ind., 89.

GENOTYPE, *N. indica* Theob., by personal designation, 1914.

Manguinhosia, Cruz in Peryassu, Os. Culic. do Brasil, 112 (1908).

GENOTYPE, *M. lutzi* Cruz, sp. nov., the only species.

Patagiamyia, James, Rec. Ind. Mus. IV, 98 (1910).

James and List., Monog. Anoph. Mosq. Ind., 2nd ed., 41: Brun., Rec. Ind. Mus. IV, 417; X, 68.

GENOTYPE, *Anopheles gigas* Giles, by original designation.

Neostethopheles, James, Rec. Ind. Mus. IV, 98 (1910).

James and List., Monog. Anoph. Mosq. Ind., 2nd ed., 40: Brun., Rec. Ind. Mus. IV, 422; X, 67.

GENOTYPE, *Anopheles aitkenii* James, in Theob., by original designation.

Nyssomyzomyia, James, Rec. Ind. Mus. IV, 101 (1910).

James and List., Monog. Anoph. Mosq. Ind., 2nd ed., 43: Brun., Rec. Ind. Mus. IV, 422; X, 67.

GENOTYPE.¹

Christophersia, James, Rec. Ind. Mus. IV, 103 (descr.) (1910).
James, Paludism, I, 33 (*nom. nud.*) (July 1910): Brun., Rec. Ind. Mus. IV, 431; X, 57.

GENOTYPE, *C. hallii* James, by original designation.

Dactylomyia, Newstead and Carter, Ann. Trop. Med. IV, 377 (1910).

GENOTYPE, *Anopheles tessellatus* Theob., as *D. ceylonica*, sp. nov., by original designation.

Neomyzomyia, Theob., Monog. Culic. V, 29 (1910).

James, Rec. Ind. Mus. IV, 109: Brun., *loc. cit.*, IV, 417; X, 68.

GENOTYPE, *Anopheles elegans* James, in Theob., by original designation.

Aldrichinella, Theob., Monog. Culic. V, 77 (1910); *nom. nov.* for *Aldrichia*, Theob., preocc. Coq., 1894 in Diptera: Brun., Rec. Ind. Mus. IV, 433; X, 55.

Memnemyia, Strickland, Ind. Jour. Med. Res. III, 201 (1915) for *Anopheles brevipalpis*, Roper.

Of the synonyms of *Anopheles* as given by me in my recent "Critical Review" all are accepted as such by Messrs. Howard, Dyar and Knab except *Conchyliaestes* and *Coelodiaezesis*. The former is sunk by the American authors in *Psorophora*, a culicine genus, and the latter is erected on larval characters, the adult not being recognizable from *Anopheles*, with which genus Edwards and other of the most recent writers incorporate it.

sinensis Wied., Auss. Zweifl. I, 547, ♂ ♀ (1828).

Anopheles sinensis, Frnfl., Verh. zool. bot. Ges. Wien, XVII, 449: Theob., Monog. Culic. I, 137, ♀, fig. 30, wing scales; pl. xxxvii, 146, ♀, typical form, full ins. col.: pl. A, wing scales: Giles, Handbk., etc., 160: *id.*, 2nd ed., 305; pl. viii, 9, wing scales, ♀: Stanton, Ind. Jour. Med. Res. III, 255: Chris., *loc. cit.*, III, 480: Waterston, Bull. Ent. Res. IX, 5.

Myzorhynchus sinensis, Theob., Monog. Culic. III, 89, fig. 53, palpus, ♀: Giles, Jour. Trop. Med. VII, 365: Blanch., Moust., 190, ♀, fig. 170, wing scales: Leices., Culic. Malaya, 30, ♂ ♀ (*Myzorhynchus*): Brun., Rec. Ind. Mus. I, 315; IV, 426: James, Monog. Anoph. Mosq. Ind., 2nd ed., 120, col. pl. xv, full ins. ♀: Strickland, Bull. Ent. Res. IV, 139.

Anopheles sinensis, subspecies *indiensis*, Theob., Monog. Culic. I, 145.

Anopheles nigerrimus, James and List. (*nec* Giles), Monog. Anoph. Mosq. Ind., 79, ♀; col. pl. iii, full ins.

Anopheles plumiger, Don., Insektenborse XVIII, 37, ♂ ♀ (1901).

Anopheles jesoensis, Tsuzuki, Centr. fur Bakteriologie XXXI, 763 (1902).

¹ Major James selects *rossii*, which is impossible, as this species was definitely made the type of *Grassia*, Theob. (renamed *Myzomyia*, Blanchard). Major James places two other species in *Nyssomyzomyia*, *ludlowi*, Theob., and *punctulata*, James and List.

Don., Zeits. f. Hyg. XLIII, 231.

Anopheles pseudopictus, Grassi, Atti R. Acad. Lincei Rendic. VIII, 109 (1899).

Myzorrhynchus peditaeniatus, Leices., Culic. Malaya, 31, ♂ ♀ (*Myzorrhynchus*) (1908).

Id., *id.*, Brun., Rec. Ind. Mus. IV, 425: Strickland, Bull. Ent. Res. IV, 135.

Anopheles argyropus, Swellengrebel, Tydsch. Ned. Ind. Geneesk. LIV, 334 (1914).

Widely distributed throughout the East. India, Assam, Burma, Java, Sumatra, Siam, Philippines, Formosa, China, Japan. Type of *sinensis*, presumably at Vienna, that of *plumiger* in Berlin Museum, described from East India and Hongkong: of *peditaeniatus* in Leicester collection. *A. jesoensis*, Tsuzuki is synonymous with *barbirostris*, Wulp, according to Christophers. The latter writer would also sink *nigerrimus*, Giles; *nigerrimus*, James and Liston, and *minutus*, Theobald, as synonyms of *sinensis*, W., typical form, or of the variety *vanus* of Theobald, with which Christophers only queries *vanus* of Walker (which takes precedence in the name) as synonymous. The synonymy of *pseudopictus* is on the authority of Waterston.

plumbeus Hal., Zool. Jour. III, 501 (descr. sine nom.); 503 (named by Steph.) (1828).

Anopheles plumbeus, Chris., Ind. Jour. Med. Res. III, 489; pl. xxviii, var. chars.

A. (Coelodiaezis) plumbeus, Edw., Bull. Ent. Res. VII, 201.

nigripes, Staeg. Nat. Hist. Tij. II, 252 (1839).

barianensis, James and List., Monog. Anoph. Mosq. Ind., 2nd ed., 76 (1911): Brun., Rec. Ind. Mus. IV, 415.

barberi, Coq., Can. Ent. XXXV, 310 (1903).

Barien, Murree Hills, Punjab, 7000 ft. [*Wredde*].

A. plumbeus was described (without a name) by Haliday amongst notes on other insects from the north of Ireland, the species being christened *plumbeus* by Stephens two pages further on in the same journal. The Kertesz Catalogue attribute the species to Stephens and sinks it as synonymous with *bifurcatus*, L.

pictus Lw., Dipt. Beit. Posen, p. 4 (1845).

Theob., Monog. Culic. IV, 124: Brun., Rec. Ind. Mus. IV, 434.

Type in Berlin Museum. A species of uncertain validity according to some authors.

Recorded from Haut Tonkin and Harioi.

vanus Walk., Proc. Linn. Soc. Lond. IV, 91, ♀ (1860).

Giles, Handbk., etc., 158.

Myzorrhynchus vanus, Blanch., Moust., 196, ♂ ♀: Brun., Rec. Ind. Mus. I, 316; IV, 427.

Anopheles sinensis, W., "sub. sp., *annularis* Wulp," Theob., Monog. Culic. I, 142, ♂ ♀; fig. 32, head; pl. v, 18, ♀; pl. A, wing scales.

Myzorhynchus sinensis annularis, Theob., Monog. Culic., I, 142; III, 90.

nec annularis, Wulp, Notes Leyd. Mus. VI, 249 (1884).

Myzorhynchus minutus, Theob., Monog. Culic. III, 91, ♀ (1903).

Brun., Rec. Ind. Mus. I, 314; IV, 425.

Anopheles minutus, Strickland, Bull. Ent. Res. IV, 136.

Lahore [Christophers]; Sambalpur, Centr. Prov., India [D. O'C. Murphy]; Calcutta, x, xi, xii; Port Canning; Quilon, 27-vii-01 [James]; Malacca; Madras [Cornwall]; Dondra, Ceylon, 4-xii-07 [Green]; Galle, Ceylon, 8-xii-07 [Green]; Perak [Wright]; Taiping [Wray]; Penang [Freer]; Kuala Lumpur [Durham]; Dindings; Luzon, 7-ix-01, Pangasinan, Manila, Albay, Laguna, Bulacan, Tayabas, Cavite, Bataan, Rizul, Nueva Ecija [all Philippine Is. iv, and vi to i, Ludlow]; Celebes, Java, China. Types of *vanus* and *minutus* in British Museum, the former much damaged.

Leicester thinks *vanus* synonymous with *sinensis*, Wied., but Theobald and Strickland keep them distinct. Major Christophers considers the *vanus* of Theobald only a variety of *sinensis*, Wied., but seems uncertain whether it is identical with *vanus*, Walk. In this var. *vanus*, Theob., he also sinks *nigerrimus*, Giles and *minimus*, Theob.

barbirostris Wulp, Notes Leyd. Mus. VI, 248, ♀ (1884).

Wulp, Tijd. v. Ent. XXVIII, 79, ♀; pl. iv, 1: Theob., Monog. Culic. I, 146, ♀; fig. 33, head; fig. 34, wing (also see p. 151); pl. A, wing scales: Giles, Handbk., etc., 146; 2nd ed., 308, ♂ ♀; pl. viii, 13a, wing, ♂ ♀; James and List., Anoph. Mosq. India, 77, ♀; pl. x, 1, wing scales; pl. v, larval figs.; col. pl. ii, full insect ♀; Roper, Bull. Ent. Res. V, 146; Strickland, Ind. Jour. Med. Res. III, 253; Chris., Ind. Jour. Med. Res. III, 462.

Myzorhynchus barbirostris, Theob., Monog. Culic. III, 86, fig. 25, larval hairs; pl. iii, wing: Theob., Gen., Ins. Fasc. 26; pl. i, 3, full insect ♀; Blanch., Moust. 197; James, Monog. Anoph. Mosq. Ind., 2nd ed., col. pl. xiv, full ins. ♀; Brun., Rec. Ind. Mus. I, 314; IV, 424; Strickland, Bull. Ent. Res. IV, 139.

Myzorhynchus barbirostris, Leic., Culic. Malaya, 33, ♂ ♀.

India, common and widely distributed. Assam, Burma, Kuala Lumpur, Klang, Malay States, Siam, Sumatra, Papua, Philippines (var. locs. common), Andamans, also occurs in old Calabar, West Africa and Japan. Type in Leyden Museum.

Christophers gives *jesoensis*, Tsuzuki (with the reference, Arch. f. Schiff. vi [1902]), as synonymous with *barbirostris*, but other authors have regarded it as identical with *sinensis*.

annularis Wulp., Notes Leyd. Mus. VI, 249, ♂ ♀ (1884).

Wulp., Tijd. v. Ent. XXVIII, 80, pl. iv, 2, head, profile: Giles, Handbk., etc., 159.

Myzomyia annularis, Brun., Rec. Ind. Mus. IV, 418.

Myzorhynchus annularis, Brun., Rec. Ind. Mus. I, 314.
nec "annularis, Wulp." of Theob., etc.

Ardjoeno, Java [*Hekmeijer*]. Type in Leyden Museum.

By some authors (Theob., Monog. I, 142) once regarded as a variety of *sinensis*, W.; but the *annularis* of Theobald is *vanus*, Walk.

subpictus Grassi, Atti R. Accad. Lincei Rendic., VIII, 1 (1899).

Brun., Rec. Ind. Mus. I, 323; IV, 434.

Recorded from "India Orientalis" in the Kertesz Catalogue (I, 254), but it seems an uncertain species. Type in Rome University Museum.

rossii Giles, Jour. Trop. Med. II, 63, ♂ ♀ (1899).

Theob., Monog. Culic. I, 154, ♂ ♀, fig. 37, wing and cross-veins, fig. 38, palpus ♂, thorax ♀, costal border, ♂ ♀, unguis ♂; pl. A, wing scales; pl. iii, 10, ♂, 9, ♀ (both full ins. col.); Giles, Handbk., etc., 149; 2nd ed., 311, ♂ ♀; pl. ix 11, wing, ♂ ♀, claws: James and List., Anoph. Mosq. India, 109, pl. vi, 1, larval figs.; pl. x, 3, wing scales; col. pl. xii, full ins. ♀: Patton and Cragg, Textbk. Med. Entom., 196, larva; Chris., Ind. Jour. Med. Res. III, 476.

Myzomyia rossii, Theob., Monog. Culic. III, 45, fig. 23, wing, ♀; figs. (pp. 46, 47), larval hairs; pl. iii, wing; pl. vi, wing scales: Blanch., Moust., 178, ♂ ♀; fig. 162, wing; fig. 163, transverse veins: Leices., Culic. Malaya, 25, ♀ ♀: Brun., Rec. Ind. Mus. I, 309; IV, 421 (*rossii*, lapsus), (copious notes).

Nyssomyzomyia rossii, James, Monog. Anoph. Mosq. Ind., 2nd ed., col. pl. xi, full ins. ♀.

Anopheles vagus, Don., Beit. z. Kennt. Anoph., 80 (1902).

var. **indefinita** Ludl., Can. Ent. XXXVI, 299, ♀ (*Myzomyia rossii*, var. *indefinita*) (1904).

Ludl., Bull. Ent. Res. VI, 155 (synonymy cleared up).

Myzomyia indefinita, Theob., Monog. Culic. IV, 47: Brun., Rec. Ind. Mus. IV, 47.

Myzomyia rossii, subspecies *indefinita*, Brun., Rec. Ind. Mus. I, 310.

Myzomyia parangensis, Ludl., Psy. XXII, 137—140 (1915).

India, widely distributed in the plains throughout the year. Sylhet, i, ii, iv, v, vi [*Hall*]; Chittagong, vii, viii, ix [*Hall*]; Ceylon, abundant; Penang [*Free*]; Straits, var. locs. [*t. Leices.*]; Perak [*Wray, Wright*]; Kuala Lumpur [*Durham*]; Singapore, 22-vii-99 [*Hanitsch*]; Phrapatoon, Siam, i, iii, viii, ix [*Woolley*]; Philippines, numerous localities, common [*Ludlow*]; Deli, iii-14; Korinchi Valley, Sungei Penoh, 2700 ft., iii-14; Lampongs, vi-14 [all Sumatra and *t. Stanton*]. The typical form mainly confined to British India and Ceylon, the var. *indefinita* mainly to the Philippines.

fuliginosus Giles, Handbk. Gnats Mosq., 160 (1900).

- Giles, Handbk., etc., 2nd ed., 298, ♂ ♀; pl. viii, 7, wing, palpus ♂ ♀, scutellum scale: Theob., Monog. Culic. I, 132, ♀, figs. 27, 28, pl. i, 3 (full ins. col. ♀): James, Sci. Mem. Med. Off. India, No. 2, fig. 18 (p. 39, larval chars.): James and Liston, Anoph. Mosq. India, 91, pl. v, 2, larval figs.; pl. x, 4, wing scales; col. pl. v, full ins. ♀: Chris., Ind. Jour. Trop. Med. III, 464.
- Nyssorhynchus fuliginosus*, Theob., Monog. Culic. III, 93; IV, 100 (life-history): Blanch., Moust., 206, fig. 179, normal wing: Brun., Rec. Ind. Mus. I, 317; IV, 428: James, Anoph. Mosq. Ind., 2nd ed., col. pl. vii, full ins. ♀.
- Anopheles jamesii*, Liston, Ind. Med. Gaz. XXXVI, p. 441 (no sex stated) (1901).
- nec jamesii*, Theob., Monog. Culic. I, 134.
- Anopheles leucopus*, Don., Insektenborse XVIII, 37, ♂ ♀ (1901).
- var. *adiei* James and List.
- Nyssorhynchus fuliginosus*, Giles, var. *adiei*, James and Liston, Anoph. Mosq. Ind., 2nd ed., 89 (1911): Chris., Ind. Jour. Med. Res. III, 465.
- Anopheles nagpori*, James and Liston, Anoph. Mosq. Ind., 1st ed., 101 (1904).
- Anopheles fuliginosus*, var. *nagpori*, *id.*, *loc. cit.*, 2nd ed., 90 (1911).
- var. *nivipes* Theob.
- Nyssorhynchus nivipes*, Theob., Entom. XXXVI, 258, ♂ (1903).
- Anopheles fuliginosus*, var. *nivipes*, Chris., Ind. Jour. Med. Res. III, 466.
- Theob., Monog. Culic. IV, 101, ♀; Leices., Culic. Malaya, 44, ♂ ♀; (*Nyssorhynchus*): Brun., Rec. Ind. Mus. I, 319; IV, 429.
- var. *pallida* Theob.
- Nyssorhynchus fuliginosus*, Giles, var. *pallida*, Theob., Monog. Culic. I, 134, fig. 28b (p. 133), wing (1901).
- Blanch., Moust., 205, fig. 179b, wing, var. *pallida*.
- A. fowleri*, Chris., Paludism II, 64 (1911)
- The typical form of *fuliginosus*, Giles, is common throughout India and Burma up to 3000 ft. occurring in many localities during the greater part of the year; probably generally distributed further east. Ceylon, Assam, Java, Samatra, Phil. Is. (var. locs.). The *adiei* variety occurs from the N.W. Frontier to Assam during winter: var. *nivipes* comes from Burma, Kuala Lumpur, i [Durham], and Klang; also from the Andamans. Type of *fuliginosus* probably lost. According to Christophers the typical specimens in the British Museum are probably others supplied by Ross. Type of *nagpori* in Central Malaria Bureau, Kasauli, of *nivipes*, *pallida* and *jamesii* in British Museum. An apparent discrepancy occurs in *fowleri* figuring in this Catalogue further on as a separate species. From Major Christophers' recent paper I understand him to mean that the

forms *howleri* and var. *pallida* are probably identical, but that he grants the former name specific rank *pro. tem.*

nigerrimus Giles, Handb. Gnats Mosq., 161, ♀ (1900).

Theob., Monog. Culic. I, 150: Giles, Handb., etc., 2nd ed., 306.

Myzorrhynchus nigerrimus, Blanch., Moust., 197: Brun., Rec. Ind. Mus. I, 314; IV, 425.

nec "*nigerrimus*, Giles," James and Liston, Anoph. Mosq. Ind., 79 (1900).

Naini Tal [Giles]; Sylhet [Hall]; Calcutta [Alcock, Annandale, Daniels]; Port Canning [Annandale]; Jalpaiguri, Dacca, Lahore, Travancore [James]; Madras. Type in British Museum. Christophers, in a recent paper, sinks this species as synonymous with var. *vanus*, Theob., of *sinensis*, Wied., including also the *nigerrimus* of James and Liston.

lindsayi Giles, Handb. Gnats Mosq., 166, ♀ (*lindsaii*) (1900).

Giles, Handb., etc., 2nd ed., 323, ♀, pl. x, 8, wing ♀ (*lindsaii*): Theob., Monog. Culic. I, 203, pl. v, 19, ♀ (full ins. col.): James and List., Anoph. Mosq. Ind., 117, col. pl. xv (full ins. ♀): Blanch., Moust., 169: Brun., Rec. Ind. Mus. I, 304; IV, 416.

Chris., Ind. Jour. Med. Res. III, 470.

Patagiamyia lindsayi, James, Anoph. Mosq. Ind., 2nd ed., 62, redescr., figs., col. pl. ii, full ins. ♀.

Bakloh, Punjab, 6000 ft., vii [Lindsay]; Ferozepore [Nurse]; Raneghat, 4000 ft. [James]; Himalayas, var. locs., 2000 up to 7000 ft., Naini Tal [Giles]; Mussoorie; Bhim Tal, ix, 06 [Annandale]; Murree, 7000 ft.; Tret, 4600 ft. Dehra Dun [Thomson]; Kurseong, Palni Hills, Nilgiri Hills, Assam.

var. *maculata* Theob., Rec. Ind. Mus. IV, 1, ♀ (1910).

Brun., Rec. Ind. Mus. IV, 416.

Kurseong, 5-vii-08 [Annandale]. Type of *lindsayi*, in British Museum. of *maculata* in Indian Museum.

gigas Giles, Ent. Month. Mag. XXXVII, 196, ♂ ♀ (1901).

Theob., Monog. Culic. II, 308, ♂ ♀: James and List., Anoph. Mosq. Ind., 118: Giles, Handbk., etc., 2nd ed., 316, ♂ ♀; pl. x, 2, wing, ♂ ♀: Brun., Rec. Ind. Mus. I, 303; IV, 416: Chris., Ind. Jour. Med. Res. III, 467.

Myzomyia gigas, Blanch., Moust., 184.

Patagiamyia gigas, James, Rec. Ind. Mus. IV, 98.

Patagiamyia simlensis, James and List., Anoph. Mosq. Ind., 2nd ed., 66, ♂ ♀ (1911).

Anopheles simlensis, Brun., Rec. Ind. Mus. IV, 416.

Anoph. gigas, var. *simlensis*, Alcock, Jour. Lond. Sch. Trop. Med. II, No. 3, p. 161 (1913).

Anoph. gigas, var. *refutans*, *id.*, *loc. cit.*, p. 161.

Himalayas; Mahasu, near Simla, 8000 ft., viii; Kathlighat, Garhwal Distr., 6000 ft. [Imms]; Murree [Maj. Smith]; Deesa,

W. Centr. India [Nurse]; Coonoor, Nilgiri Hills, 5000—6000 ft. [Price]; Khasi Hills; Pachmarhi Hills; Ceylon (hills) [t. Alcock; Green]. Type of *gigas* in British Museum; of *simlensis* in Central Malaria Bureau, Kasauli.

culicifacies Giles, Ent. Month. Mag. XXXVII, 197, ♀ only, the ♂ renamed *turkhudi*, List. (1901).

Theob., Monog. Culic. II, 309, ♀ *nec* ♂ (t. Theob., *loc. cit.*, III, 48); Giles, Handbk., etc., 2nd ed., 317, pl. ix, 12, ♀ *nec* ♂; James and List., Anoph. Mosq. Ind., 106, pl. ix, 2, wing scales, pl. viii, 1, larval figs.; col. pl. xi, full ins. ♀; Chris., Ind. Jour. Med. Res. III, 463.

Myzomyia culicifacies, Theob., Proc. Royal Soc. LXIX, 379, ♀; *id.*, Monog. Culic. III, 39, fig. (p. 40), larval hairs; pl. iii, wing; pl. viii, wing scales; IV, 51; V, 25; Blanch., Moust., 182, ♀; James, Monog. Anoph. Mosq. Ind., 2nd ed., pl. iii, full ins. col. ♀; Brun., Rec. Ind. Mus. I, 305; IV, 404 (criticism on name of species), 418.

Myzomyia culicifacies, var. *punjabensis*, James, Anoph. Mosq. Ind. 72 (1911).

Anopheles listoni, Giles, Ent. Month. Mag. XXXVII, 197, ♂ ♀ (1901).

Id., Handbk., etc., 2nd ed., 319, ♂ ♀; pl. x, 4, wing, ♂ ♀; head, ♂ ♀; Theob., Monog. Culic. II, 311, App. ♀.

nec listoni, Liston,

Anopheles indicus, Theob., *loc. cit.*, I, 183, ♀ (1901).

Anopheles indicus, Giles, Handbk., etc., 2nd ed., 320, ♀.

Pyretophorus sergenti, Theob., Monog. Culic. IV, 68 ♀, fig. 16, wing (1907).

India, common, generally distributed up to 3000 ft.; Mandalay [Annandale]; Dr. Gough says it ranges from Algeria to India, and Edwards records it from as far west as Lake Tiberias. *Types* of *culicifacies* and *indicus* in British Museum; also of *listoni*, presumably the type ♂ attributed to *culicifacies*, Giles; of *punjabensis* in Central Malaria Bureau, Kasauli.

According to the strict rules of zoological nomenclature, since two species were confused at the time of description, the name *culicifacies* should have been retained for the ♂ (now known as *turkhudi*, List.), and the ♀ (still known as *culicifacies*) should have been renamed (*v. my* Cat. Or. Culic. Supp., p. 404).

theobaldi Giles, Ent. Month. Mag. XXXVII, 198, ♀ (1901).

Giles, Handbk., etc., 2nd ed., 299, ♀; Theob., Monog. Culic.

II, 311 (App.) ♀; James and List., Anoph. Mosq. Ind., 97, col. pl. viii, full ins. ♀; Chris., Ind. Jour. Med. Res. III, 482.

Nyssorhynchus theobaldi, Theob., Monog. Culic. III, 95; Blanch., Moust. 210; Brun., Rec. Ind. Mus. I, 320; IV, 430.

James, Monog. Anoph. Mosq. Ind., 2nd ed., col. pl. x, full ins. ♀.

Anoph. maculatus, var. *theobaldi*, Alcock, Jour. Lond. Sch. Trop. Med. II, No. 3, p. 164 (1913).

Ellichpur (Centr. Ind.) [Liston]; Shahjahanpur, 19-x-00 [Giles]; Dacca [Macrae]; Nagpur [Stephens]; Sambalpur [Murphy]; Lahore, x; Bombay; Benguet, Phil. Is., iii, iv, xi. Also occurs in the Aden hinterland. *Type* in British Museum.

kochi Don., Insektenborse XVIII, 36, ♂ ♀ (1901).

Theob., Monog. Culic. I, 174, ♀; pl. iv, 16 ♀, full ins. col. Edwards, Bull. Ent. Res. IV, 222; Gough, Bull. Ent. Res. V, 142; Roper, *loc. cit.*, V, 129, life-history; Chris., Ind. Jour. Med. Res. III, 469.

Nyssorhynchus kochi, Blanch., Moust., 208.

Cellia kochi, Theob., Monog. Culic. III, 110; Leices., Culic. Malaya, 46, ♂ ♀: Brun., Rec. Ind. Mus. I, 321; IV, 431.

Anopheles ocellatus, Theob. (MS. name only), Monog. Culic. II, 306, note as to name.

Giles, Handbk., etc., 2nd ed., 315, ♀, pl. ix, 5, wing ♀, dorsum of thorax.

Cellia flava, Ludl., Can. Ent. XL, 32, ♂ ♀ (1908): Brun., Rec. Ind. Mus. IV, 431.

Nyssorhynchus flava, Ludl., Psy. XVIII, 126 (1911).

Christophersia hallii, James, Paludism I, 32, ♂ ♀, figs. var. chars. (1910). James and List., Anoph. Mosq. Ind., 2nd ed., 123: Brun., Rec. Ind. Mus. IV, 431.

Sylhet, 4-ii-05; 7-iv-05; vi, 31-vii-05; 15-xii-05 [all Hall]; Singapore [Biro]; Taiping, Perak, Kuala Lumpur, xii-02 [Durham]; Malay Peninsula, var. locs.; Padarg, Sumatra [*t. Donitz*]; Deli, iii-14; Lampongs, vi-14 [both Sumatra, *t. Stanton*]; Serang and Tjimahi, Java [*t. Donitz*]; Tayabas, Phil. Is., ix-07. *Type* of *ocellatus* in British Museum; of *flava* in Army Medical Museum, Washington; of *hallii* in Indian Museum.

Theobald had described this species in the MS. of his Monograph as *ocellatus*, but as the description of Donitz's *kochi* was published just before Theobald's work, he allowed his own description to stand as that of *kochi*, and in Vol. II, 306, he again notes that *ocellatus*, Theob., is a MS. synonym of *kochi*, in case specimens may exist in any collection under the MS. name.

punctulatus Don., Insektenborse XVIII, 36, ♂ ♀ (1901).

Roper, Bull. Ent. Res. V, 144.

nec Anopheles, id., Theob., Monog. Culic. I, 175.

Cellia punctulata, Theob., Monog. Culic. IV, 109: Brun., Rec. Ind. Mus. IV, 432.

Stephansort, Papua; Herbertshohe, Bismarck Archipelago [both *t. Donitz*]. See note under *A. tessellatus*.

Mr. Roper (Bull. Ent. Res. V, 144) makes *tessellatus*, Theob., a synonym of this species, but I have retained Mr. Edwards' view in keeping them distinct.

leucosphyrus Don., Insektenborse XVIII, 37, ♀ (1901).

Theob., Monog. Culic. II, 307, App. ♀: Roper, Bull. Ent. Res. V, 193; Chris., Ind. Jour. Med. Res. III, 469.

Myzomyia leucophyrus, Giles, Handbk., etc., 2nd ed., 312, fig. 44, wing: James and List., Anoph. Mosq. Ind. 82; Leices., Culic. Malaya, 28, ♂ ♀: Brun., Rec. Ind. Mus. I, 307.

Nyssorhynchus leucophyrus, Blanch., Moust., 213, ♀.

Neomyzomyia leucophyrus, Brun., Rec. Ind. Mus. IV, 417.

Myzomia elegans, James in Theob., Monog. Culic. III, 51, ♀; fig. 28, wing scales, cross veins, 29, wing (1903).

Myzomyia elegans, Brun., Rec. Ind. Mus. I, 306.

Anopheles leucosphyrus, var. *elegans*, James and List., Monog. Anoph. Mosq. Ind., 82, ♀; pl. ix, 4, wing scales; pl. xii, wing, palpus, leg, larva.

Pyrethophorus, id., Theob., Monog. Culic. IV, 77.

Neomyzomyia elegans, James, Theob., Monog. Culic. V, 30, ♂ ♀: two figs. of ♀, wing and other chars. (This description by Theobald): Brun., Rec. Ind. Mus. IV, 417.

Karwar, iv [Cogill]; Meenglas, Jalpaiguri [Wallich]; Andaman Is. [Ray White]; Kajoe Janam, N. of Padang, Sumatra [t. Donitz]; Deli, iii-14; Lampongs, vi-14 [both Sumatra and t. Stanton]; Moearah Tewah, Borneo [t. Donitz]; Malay Peninsula. *Type* of *elegans* in British Museum.

turkhudi Liston, Ind. Med. Gaz. XXXVI, 441, no sex stated, apparently ♀ (1901).

Giles, Handbk., etc., 2nd ed., 320, ♀ (footnote): James, Sci. Mem. Ind. No. 2, 49; fig. 27, wing; fig. 28, larval chars.: James and List., Anoph. Mosq. Ind., 115; pl. viii, 2, larval figs.; col. pl. xiv, full ins. ♀: James, Anoph. Mosq. Ind., 2nd ed., col. pl. v, full ins. ♀: Gough, Bull. Ent. Res. V, 133, new syns. and notes: Chris., Ind. Jour. Med. Res. III, 483.

Myzomyia turkhudi, Theob., Monog. Culic. III, 48, ♀; pl. iii, wing (1903).

Blanch, Moust., 183: Brun., Rec. Ind. Mus. I, 311; IV, 422.

Anopheles culicifacies, Giles, Ent. Month. Mag. XXXVII, 197 (1901).

Theob., Monog. Culic. II, 309, ♂ only: Giles, Handbk., etc., 2nd ed., 317, ♂; pl. ix, 12, ♂ ♀: Theob., Proc. Roy. Soc. LXIX, 379, ♂; fig. 2 (p. 380), genitalia ♂.

Anopheles multicolor, Camboulin, C. R. Acad. Sci. Paris, CXXV, 704 (1902).

Anopheles impunctus, Don., Zeits. f. Hygiene, XLI, 67; pl. ii, 15 (1902).

Pyrethophorus chaudoyei, Theob., Monog. Culic. III, 68 (1903).

P. nigri-fasciatus, id., loc. cit., IV, 65 (1907).

P. myzomyfasciatus, id., loc. cit., IV, 69 (1907).

Pyrethophorus cleopatra, Willcocks, nom. nud., Ann. Trop. Med. Parasit. III, 586 (1910).

Myzomyia azriki, Patton, Jour. Bomb. Nat. Hist. Soc. XVI, 632 (1905).

M. hispaniola, Theob., Monog. Culic. III, 49 (1903).

India, generally distributed (Ellichpore, Cashmere, Nagpur, Hoshangabad, Lahore, Ferozepore, Aurungabad); Andaman Is. [*Maj. Anderson*]. Dr. Gough states the species ranges continuously from Teneriffe through the north coast of Africa to India. *Types* of *turkhudi* and *nigri-fasciatus* in British Museum.

Theobald's description of the ♂ was drawn up from Giles' type, which at that time was considered to be the ♂ of *culicifacies*, Giles, and was first described as such by both Giles and Theobald. *Vide* notes under *culicifacies*.

vincenti Laveran, Comp. Rend. LIII, 993 (1901).

Blanch., Moust., 175, ♀: Theob., Monog. Culic. V, 184:

Brun., Rec. Ind. Mus. I, 322; IV, 434.

Van-Linh (Haut Tonkin) [*t. Laveran*]. *Type*, location unknown to present author.

maculatus Theob., Monog. Culic. I, 171, ♂ ♀ (*maculata*), fig. 48, var chars. (1901).

James, Sci. Mem. Med. Off. Ind. No. 2, 47, fig. 25, wing and leg; Giles, Handbk., etc., 2nd ed., 301, ♂ ♀; pl. ix, 2, head, wing ♂, palpus ♀; James and List., Anoph. Mosq. Ind., 99; Roper, Bull. Ent. Res. V, 143; Chris., Ind. Jour. Med. Res. III, 471.

Nyssorhynchus maculatus, Theob., Monog. Culic. III, 96; Blanch.,

Moust., 207, ♂ ♀; fig. 180, wing; Leices., Culic. Malaya, 41, ♂ ♀; Brun., Rec. Ind. Mus. I, 318; IV, 429.

Nyssorhynchus willmori, Leicester, Culic. Malaya, 42, ♂ ♀. *nec willmori*, James.

Nyssorhynchus pseudowillmori, Theob., Monog. Culic. V, 65, ♀ (1910) Brun., Rec. Ind. Mus. IV, 430.

Anopheles maculatus, var *willmori*, Alcock, Jour. Lond. Sch. Trop. Med. III, No. 3 (1913).

Himalayas up to 6000 ft. from N.W. Frontier to Assam; Lahore, iii, iv [*Christophers*]; Jeypore; Jalpaiguri, 13-vii-09 [*Wallich*]; Kurseong, Nara Ghat, 25—26-ii-08 and Thaumaspur, 18—20-ii-08, ♂ ♀ (the two latter on the Bengal-Nepalese Frontier); Perak; Dindings, xi; Korinchi Lake (Sandarang Agong, Sumatra, 2450 ft.), v-14; Batak Highlands, Sumatra 4000 ft., iv-15; Hongkong, ix—x. *Types* originally stated to be in Dr. Ree's collection, but the *type* ♂ is now in the British Museum; the *type* ♀ is not in existence. *Type* of *pseudowillmori* in Ind. Mus.

Roper recently considers *willmori*, James in Theob. and *dudgeoni*, Theob., to be synonymous with *maculatus*, but Christophers states this view to be erroneous. This latter author (Ind. Jour. Med. Res. III, 471) clears up a difficulty set forth by Stanton (Jour. Lond. Sch. Trop. Med. II, No. 1, p. 7) making it obvious that *maculatus*, Theob., was actually described from both sexes of that species. He states that Theobald in his later descriptions mixed *maculata* with *karwari*.

tessellatus Theob., Monog. Culic. I, 175, ♀ (as *Anopheles punctulatus*, Don.), fig. 49, thorax, wing, hind leg; pl. xxxvii. 148, ♀, full ins. col. (1901).

Giles, Handb., etc., 2nd ed., 305, ♀; pl. ix, 7, wing ♀, dorsum of thorax, hind tarsus (*tesselatus*, lapsus): Edw., Bull. Ent. Res. IV, 221: Stanton, Ind. Jour. Med. Res. III, 255.

Myzomyia tessellata, Theob., Monog. Culic. III. 55 (*tessellatum*); IV, 42 (*tessellata*): Brun., Rec. Ind. Mus. I, 311; IV, 422.

Anopheles punctulatus, James and List., Anoph. Mosq. Ind., 84, pl. xi, wing, head, leg: Chris., Ind. Jour. Med. Res. III, 482.

Myzomyia punctulata, Leices., Culic. Malaya, 27, ♂ ♀: Brun., Rec. Ind. Mus. I, 309.

Nyssorhynchus punctulatus, Blanch., Moust., 208.

Nyssomyzomyia punctulata, James and List., Monog. Anoph. Mosq. Ind., 2nd ed., 104, pl. xiv, wing, head, leg.

Anopheles deceptor, Don., Zeits. for Hyg. u. Insek., XLI, 60.

Myzomyia deceptor, Theob., Monog. Culic. V, 29: Brun., Rec. Ind. Mus. IV, 418.

Myzomyia thornoni, Ludl., Can. Ent. XXXVI, 69, ♀ (1904): Blanch., Moust., 183: Brun., Rec. Ind. Mus. I, 311; IV, 422.

Dactylomyia ceylonica, Newstead and Carter, Ann. Trop. Med. Paras. IV, 377 (1910).

Trincomalee, Ceylon [Green]; Taiping, v; 22-xi-99; 21-xii-99 [Wray]; Sumatra [t. Donitz]; Deli, iii-14; Lampongs, vi-14; [both Sumatra and t. Stanton]. Type of *tessellatus* in British Museum; of *thornoni*, in Army Med. Mus. Washington; of *ceylonica* in Liverpool Sch. of Trop. Med.

Theobald had given the name *tesselatus* to this species in the MS. of his Monograph, but just before publication he considered it identical with *punctulatus*, Don., and changed the name to that of Donitz's species, confirming this later (*l.c.*, II, 306), as specimens under his own MS. name had been sent out to correspondents. In Vol. III. 55, he says Donitz informed him the two species were distinct.

minimus Theob., Monog. Culic. I, 186, ♀, fig. 55, wing, thorax, cross-veins (1901).

Giles, Handbk., etc., 2nd ed., 321, ♀; pl. x, 7, wing ♀, thorax scales: Chris., Ind. Jour. Med. Res. III, 473.

Pyretophorus minimus, Giles, Jour. Trop. Med. VII, 365: Blanch., Moust., 188, ♀, fig. 169, wing, transverse veins: Brun., Rec. Ind. Mus. I, 313; IV, 423.

Anopheles christopheri, Theob., Proc. Roy. So. LXIX, No. 456, p. 378, ♀, pl. v, 3, wing (Mar. 1902): Theob., Monog. Culic. V, 24: James and List., Anoph. Mosq. Ind., 103, pl. vii, 1, larval figs.: col. pl. x, full ins. ♀: Giles, Handbk., etc., 2nd ed., 512: Edw., Bull. Ent. Res. IV, 222: Ludl., *loc. cit.*, VI, 155 (synonymy).

Anopheles formosaensis I, Tsuchi, Arch. f. Schiff. IV, 9 (May, 1902).

A. aconitus, var. *cohaesa*, Don., Zeits. f. Hyg. XLIII, 233 (1903);
Myzomyia funesta, Ludl., Can. Ent. XXXVII, 135 (1905).
nec *funestus*, Giles.

Myzomyia mangyana, Banks, Phil. Jour. Sci. I, 991, ♀ (1906).
Brun., Rec. Ind. Mus. I, 309.

Anopheles febrifer, Banks, Phil. Jour. Sci. IX, Sect. D, 405 (1914);
(prelim. descr. by Drs. Walker and Barber, *loc. cit.*, IX, Sect.
B, 384 (1914).

var. *aconitus*, Don., Zeits. f. Hyg. XLI (1902).

Chris., Ind. Jour. Med. Res. III, 475.

Myzomyia albirostris, Theob., Monog. Culic. III, 24, ♂ ♀, fig.
11 (p. 25), palpus and proboscis (1903).

Leices., Culic. Malaya, 23, ♂ ♀: Brun., Rec. Ind. Mus. I,
305; IV, 418.

Myzomyia christophersi, var. *alboapicalis*, Theob., Monog. Culic.
V, 25 (1910).

A. brahmacharii, Chris., Paludism, No. 5, p. 11 (1912).

Id., Brahmachari, Ind. Med. Gaz. XLVI, 268 (1911).

Myzomyia flavirostris, Ludl., Bull. No. 4, Surg. Gen. Office (1913):
Psy. XXI, 30 (1914).

Sylhet, Central India, Meenglass, Jalpaiguri, 13-vii-07 [*Wallich*]; Calcutta [*Alcock*]; Burma, Assam, Kangra Valley, Kuala Lumpur, Pokfulam, Hongkong [*Rees*]; Pampanga and Phil. Is. (var. locs.), ii, and v to xi [*t. Ludlow*]. *Mangyana* comes from Rio Baco, Chicago, Mindoro (Phil. Is.), v [*McGregor*]. Type of *minimus* in Dr. Rees collection. Types of *christophersi* and *alboapicalis* in British Museum, of *brahmacharii* in Central Malaria Bureau, Kasauli, of *flavirostris* in Army Medical Museum, Washington, of *febrifer*, ♂ ♀, No. 18,015 in Bureau of Science, Entom. Coll., Manila; of *mangyana* in Ent. Coll. No. 3290, Bureau of Science, Manila.

Christophers in a recent paper sinks this species as synonymous with var. *vanus*, Theob. of *sinensis*, Wied.

stephensi Liston, Ind. Med. Gaz. XXXVI, 441, no sex stated, apparently ♀ (1901).

James, Sci. Mem. Ind., No. 2, 45, fig. 23, wing, 24, larval chars.: Giles, Handbk., etc., 2nd ed., 331, ♀ (footnote): James and List., Anoph. Mosq. Ind., 113, pl. vi, 2, larval figs.; pl. x, 6, wing scales; col. pl. xiii, full ins. ♀: Chris., Ind. Jour. Med. Res. III, 481.

Neocellia stephensi, James, Monog. Anoph. Mosq. Ind., 2nd ed., col. pl. xii, full ins. ♀.

Nyssorhynchus stephensi, Theob., Monog. Culic. III, 93, ♀; fig. 54, var. wing marks; fig. 55, wing; figs., pp. 40, 47, larval hairs: Blanch., Moust., 210: Brun., Rec. Ind. Mus. I, 319; IV, 430.

Anopheles metaboles, Theob., Proc. Royal So. LXIX, 374, ♀; pl. v, 1, wing, ♀ (1902): Giles, Handbk., etc., 2nd ed., 512.

Neocellia intermedia, Rothwell, Entom., ♀ (1907).

Theob., Monog. Culic. IV, 115, ♀; V, 73, ♂ ♀, fig. 29, wing
♂: Brun., Rec. Ind. Mus. IV, 432.

N.W. Frontier, Deesa, W. Centr. India, i, viii, Mian Mir, Ferozepore, v to xi [*t. Adie*]; Nagpur, Ellichpore, Madras City, Karachi, common, Calcutta, 1-viii-07 [*Annandale*]; Lushai Hills, Assam, 9-vii-04 [*MacLeod*]; Burma, Philippines [*t. Ludlow*]. *Types of metaboles and intermedia* in British Museum; location of that of *stephensi* unknown to present author.

Major Christophers notes that though Theobald (Monog. IV, 96) denies that *metaboles*, Theob., is synonymous with *stephensi*, List., yet his (Theobald's) *description of metaboles* is one of *stephensi*.

funestus Giles, Mem. Liv. Sch. Trop. Med., p. 50, pl. v (1900).

Giles, Handbk., etc, 162, pl. vii, 3, wing: Theob., Monog. Culic. I, 178, ♂ ♀; var. chars. figd.; pl. iv, 13, full ins. ♀, col; IV, fig. 10, ♂ genitalia; V, 17: Newstead, Ann. Trop. Med. Paras. I, 9.

Myzomyia funesta, Theob., Monog. Culic. III, 34, pl. ii, wing ♀: *id.*, Gen. Ins., Fasc. 26, pl. i, 2, full ins. col. ♀: Brun., Rec. Ind. Mus. I, 306; IV, 419: Blanch., Moust., 180, ♂ ♀.

var. *listoni* List.

Anopheles listoni, List., Ind. Med. Gaz. XXXVI, 441, ♀ (1901). nec *listoni*, Giles, ♀.

Myzomyia listoni, Theob., Monog. Culic. III, 27, ♀; fig. 12, palpus ♂ and wing scale; fig. 13, wing; fig. 17 (p. 40), larval hairs: James, Monog. Anoph. Mosq. Ind., 2nd ed., pl. iv, full ins. col. ♀ (on plate as *Anoph. fluviatilis*): Brun., Rec. Ind. Mus. I, 308; IV, 420: James, Sci. Mem. Med. Off. Ind., No. 2, 31, fig. 9.

Anopheles fluviatilis, James, Sci. Mem. Med. Off. India, No. 2, p. 31, fig. 9 (1902): Theob., Proc. Roy. So. LXIX, 378, ♀; pl. v, 3, wing, ♀ (1902): James and List., Anoph. Mosq. Ind., 103, pl. vii, 1, larval figs.: col. pl. x, full ins. ♀.

var. *leptomeres* Theob.

Myzomyia leptomeres, Theob., Monog. Culic. III, 38, ♀ (1903).

Id., loc. cit., IV, 124; V, 29: Brun., Rec. Ind. Mus. I, 307; IV, 419.

Type of funestus doubtful if in existence, of *leptomeres* in British Museum.

A. funestus, Giles, is an African species, of which the typical form has not been recorded from India though it is common in the Philippines, and references to it are given at the top of this list to aid identification. Theobald describes two African varieties, *umbrosa* and *subumbrosa*, of which the latter may be identical with the *leptomeres* variety. Giles is wrong in considering the var. *leptomeres* identical with *pictus*, Lw. The form *listoni* is generally distributed in India, N.W. Frontier, Assam, Ceylon [*Green*]; Central India, Burma, Western India, Sylhet, 21-i-05; 2-ii-05; 2-v-05 [*Hall*]; Perak. Some con-

fusion has arisen here over the name *listoni* by the form being at first considered identical with Giles' "*listoni*," as described by Capt. Liston, who, however, was unknowingly describing a new species; and as the real *listoni* of Giles proves synonymous with *culicifacies*, Giles, the name *listoni* must stand good for Capt. Liston's variety. See my notes and synonymy, Rec. Ind. Mus. I, 305, 306, 308 and IV, 420; also Major Christophers, Ind. Jour. Med. Res. III, 181.

- maculipalpis** Giles, Handbk., etc., 2nd ed., 279, ♂ (1902).
James and List., Anoph. Mosq. Ind., 95, pl. iv, larval figs.; pl. x, 5, wing scales; col. pl. vii, full ins. ♀: Chris., Ind. Jour. Med. Res. III, 472.
- Nyssorhynchus maculipalpis*, Theob., Monog. Culic. III, 96, ♀; fig. 56, var. chars.; pl. ii, two ♀ wings; fig. (p. 98), larval hairs: Brun., Rec. Ind. Mus. I, 319; IV, 429; James, Monog. Anoph. Mosq., Ind., 2nd ed.; col. pl. ix, full ins. ♀.
- Nyssorhynchus maculipalpis*, Giles, var. *indiensis*, Theob., Monog. Culic. III, 99 (1903): Brun., Rec. Ind. Mus. I, 319.
- Nyssorhynchus indiensis*, Theob., Monog. Culic. IV, 98: Brun., Rec. Ind. Mus. IV, 428.
- Anopheles indiensis*, James and List., Anoph. Mosq. Ind., 95, pl. ii, figs.
- Anopheles jamesii*, Theob., Monog. Culic. I, 134, ♀; pl. i, 2, ♀, full ins. col. (1901).
Giles, Handbk., etc., 2nd ed., 299, ♀: James, Sci. Mem. Med. Off. Ind., No. 2, 41: James and List., Anoph. Mosq. Ind., 93; col. pl. vi, full ins., ♀: Chris., Ind. Jour. Med. Res. III, 468: Stephens and Chris., Rep. Roy. So. Malaria, 1902.
- Nyssorhynchus jamesii*, Blanch., Moust., 206: Brun., Rec. Ind. Mus. I, 318; IV, 429; James, Monog. Anoph. Mosq. Ind., 2nd ed., col. pl. viii, full ins. ♀: Stephens and Chris., Rep. Roy. So. Malaria, 1902.
- India [*Christophers*, Stephens]; N.W. Frontier, Indo-Gangetic Plain, Central Provinces, Lower Ranges Himalayas, Shahjahanpore, Ferozepore, Nagpur, Ellichpore, Bombay, Karwar, Decan, Goa, Shamnagar, Calcutta, Port Canning, Quilon, Travancore, Ceylon, Pangasinam, Phil. Is.; also occurs in Mashonaland and Mauritius. *Type of maculipalpis* in British Museum.
- Theobald considered (Monog. IV, 98) his "*maculipalpis*, Giles, var. *indiensis*, Theob." (of Monog. III, 99) to be a distinct species from *maculipalpis*, Giles, typical form, but he ranked "*maculipalpis*, Giles of James and Liston" as synonymous with *indiensis*.
- aconitus** Don., Beit. Kennt., z. d. Anoph. 70, ♀ (1902).
Don., Zeits. f. Hygiene XLIII, 233: Theob. Monog. Culic. III, 30; fig. (p. 31), wing ♀.
- Myzomyia aconita*, Brun., Rec. Ind. Mus. I, 305.
- Anopheles aconitus*, *id.*, *loc. cit.*, IV, 415 (*aconita*): Stanton, Ind. Jour. Med. Res. III, 252.

? *A. formosaensis*, Tsuzuki.¹

Kajoe Tanam; Willen Is.; Soekaboemi, Java [*t. Donitz*]; Deli, Sumatra, iii-14; Lampongs, Sumatra, vi-14 [*t. Stanton*]. *Type*, location unknown to present author.

Vide Brun., *Rec. Ind. Mus.* IV, 415 for note on presumed variety *formosaensis*, and suggested change of name to *cohaesa*. Stanton considers (*loc. cit.*) that *albirostris*, Theob., is synonymous with this species.

jeyporiensis James, *Sci. Mem. Ind.*, No. 2, 32; fig. 11, wing; fig. 12, larval chars. (1902).

James and List., *Anoph. Mosq. Ind.*, 101, pl. vii, 2, larval figs.; col. pl. ix, full ins. ♀: Chris., *Ind. Jour. Med. Res.* III, 468.

Pyrelophorus jeyporiensis, Theob., *Monog. Culic.* III, 66; pl. viii, wing scales; James, *Anoph. Mosq. Ind.*, 2nd ed., col. pl. vi, full ins. ♀; fig. (p. 67), palpus ♂: Brun., *Rec. Ind. Mus.* I, 313 (*jeyporensis*).

Jeypur State [*Christophers, Stephens*]; Jakot, S. India [*Aitken*]; Nagpur [*James*]; Bombay, Central Provinces, India, Castle Rock, N. Kanara Distr. *Types* in British Museum.

formosaensis Tsuzuki, *Archiv. f. Schiffv. v. Tropen Hygiene* VI, 289 (1902).

Theob., *Monog. Culic.* V, 84; Brun., *Rec. Ind. Mus.* I, 323; IV, 415, notes.

Formosa. *Type*, location unknown to present author.

There are apparently two species of the same name by the same author from North and South Formosa respectively. *Vide* my notes (*loc. cit., supra*). Theobald (*Monog. V*) says that Donitz (*Zeits. f. Hygiene* XLIII, 233) says this species is only a variety of *aconita*, Don., and proposes, unnecessarily, to change the name to *cohaesa*.

pulcherrimus Theob., *Proc. Roy. So.* LXIX, 396, ♀; pl. v, 2, wings, ♀ (1902).

James, *Sci. Mem. Ind.*, No. 2, p. 48; fig. 26, wing; James and List., *Anoph. Mosq. Ind.*, 86, col. pl. iv, full ins. ♀; Giles, *Handbk., etc.*, 2nd ed., 510, ♀; Chris., *Ind. Jour. Med. Res.* III, 476.

Cellia pulcherrima, Theob., *Monog. Culic.* III, 107; Blanch., *Moust.*, 215, ♀; Brun., *Rec. Ind. Mus.* I, 321; IV, 432.

James, *Monog. Anoph. Mosq. Ind.*, 2nd ed., col. pl. xiii, full ins., ♀.

Mian Mir, near Lahore [*James*]; Ferozepore, vii to xii [*Adie*]; Goa [*James and List.*]; Purneah, 7-viii-07 [*Paiva*]; also occurs at Kokand, Turkestan [*Univ. Coll. Helsingfors*]. *Type* ♀ in British Museum, ♂ apparently still unknown.

pseudobarbirostris Ludl., *Jour. N.Yk. Ent. So.* X, 127, ♀ (*Myzorrhynchus*) (1902).

¹ *Vide A. formosaensis*, Tsuzuki, p. 105.

- Myzorhynchus pseudobarbistrois*, Theob., Monog. Culic. IV, 83, ♀: Brun., Rec. Ind. Mus. I, 315; IV, 426.
Philippines, generally distributed, May to October. Type in Army Medical Museum, Washington.
- philippinensis** Ludl., Jour. N.Yk. Ent. So. X, 128 (1902); also Jour. Amer. Med. Assn. XXXIX, 426 (1902).
Nyssorhynchus philippinensis, Theob., Monog. Culic. IV, 103, ♀: Brun., Rec. Ind. Mus. I, 319; IV, 430.
Philippines, common, June to January. Type in Army Medical Museum, Washington.
- aitkenii** James in Theob., Monog. Culic. III, 22, ♀ (1903).
James and List., Anoph. Mosq. Ind., 119; pl. ix, 3, wing scales; pl. xiii, larval figs. and wing: Blanch., Moust., 620: Brun., Rec. Ind. Mus. I, 303; IV, 415: Chris., Ind. Jour. Med. Res. III, 461.
Neostethopheles aitkenii, James, Anoph. Mosq. Ind., 2nd. ed., 59; figs.
- Stethomyia fragilis*, Theob., Entom. XXXVI, 257, ♂ (1903): Brun., Rec. Ind. Mus. I, 312; IV, 423.
- Stethomyia pallida*, Ludl., Can. Ent. XXXVII, 129, ♀ (1905): Brun., Rec. Ind. Mus. I, 312; IV, 423.
Goa Frontier [*Aitken*]; Karwar [*Dr. Cogill*]; Meenglas, Jalpaiguri, 9-viii-07 [*Wallich*]; Bengal Duars, Nilgiri Hills, Bengal, Assam, Kuala Lumpur, i, xii [*Durham*]; Burma, Pampanga, ix-05 [*t. Ludl.*]. Type of *aitkenii* in British Museum.
Major Christophers states that there is an individual labelled "type specimen" by James in the Central Malaria Bureau, Kasauli. He also considers *Stethomyia fragilis*, Theob., *S. pallida*, Ludl. and *Anopheles treacherii*, Leices., synonymous with *aitkeni*.
- immaculatus** James, Sci. Mem. Ind., No. 2, 35 (1902).
Theob., Monog. Culic. III, 23, ♀ (1903): James and List., Anoph. Mosq. Ind., 120: Brun., Rec. Ind. Mus. I, 304; IV, 416: Chris. Ind. Jour. Med. Res. III, 468.
Ennur, Madras Coast [*James and List.*]. Type in British Museum, unique.
On the ground of the brief description of fine words only, "wings entirely unspotted, legs unbanded" being insufficient I attributed the species in my Catalogue to Theobald instead of James, but on Major Christophers' plea that such diagnosis differentiated it from all other known species, the original author's name is here restored, though such "descriptions" are to be deprecated.
- ludlowi** Theob., Monog. Culic. III, 42 (*Myzomyia*) (1903).
Roper, Bull. Ent. Res. V, 145: Chris., Ind. Jour. Med. Res. III, 470.
Myzomyia ludlowi, Banks, Phil. Jour. Sci. II, 519, life-history; pls. i to xi; pl. v, full ins. ♂ ♀: Brun., Rec. Ind. Mus. I, 309; IV, 420.

Nyssomyzomyia ludlowi, James and List., Monog. Anoph. Mosq. Ind., 2nd ed., 101.

Madras, Ceylon, Singapore [*Biro*]; Burma; Deli, Sumatra, iii-14, Andaman Is., Phil. Is., common, var. locs. [*l. Ludlow*]. Type in British Museum.

palestinensis Theob., Monog. Culic. III, 71, ♂ ♀; fig. 46, head and thorax; 47, two wings, ♀; ♂ genitalia; pl. vi, wing scales, ♀ (*Pyretophorus*) (1903).

Edw., Jour. Asiat. Soc. Beng. IX, 48 (1913): Waterston, Bull. Ent. Res. IX, 4.

Anopheles nursei, Chris., Ind. Jour. Med. Res. III, 475.

Pyretophorus nursei, Theob., Monog. Culic. IV, 66, ♀ (1907).

Brun., Rec. Ind. Mus. IV, 424.

Pyretophorus cardamitis, Newstead and Carter, Ann. Trop. Med. Paras. IV, 379 (1910).

? *Anopheles superpictus*, Grassi, Reale Accad. Linc., 78 (1900).

Anopheles superpictus, var. *macedoniensis*, Cot. and Hovasse, Bull. Soc. Path. Exot. X, 890, fig. (1917).

Quetta, xi [*Nurse*]. Occurs in Palestine on Beyrouth River, vi [*Cropper*]; Cyprus, common up to 4000 ft. [*Miss D. M. A. Bate*]; Athens, Kashan, Persia, 8-xi-11, ♀ [*Cropper*]. Type of *nursei* in British Museum, location of that of *palestinensis* unknown to present author.

umbrosus Theob., Monog. Culic. III, 87, ♀; fig. 87, wing (*Myzorhynchus*) (1903).

Roper, Bull. Ent. Res. V, 146.

Myzorhynchus umbrosus, Leic., Culic. Malaya, 35, ♀ (*Myzorhynchus*): Brun., Rec. Ind. Mus. I, 316; IV, 427; Strickland, Bull. Ent. Res. IV, 140.

Malay States, Pahang [*Dr. Durham*]; Kuala Lumpur, Deli, Sumatra, iii-14. Type in British Museum.

albotaeniatus Theob., Monog. Culic. III, 88, ♀; pl. i, wing; pl. v, wing scales (*Myzorhynchus*) (1903).

Roper, Bull. Ent. Res. V, 147; Stanton, Ind. Jour. Med. Res. III, 252.

Myzorhynchus albotaeniatus, Leices., Culic. Malaya, 34, ♀ (*Myzorhynchus*): Brun., Rec. Ind. Mus. I, 313; IV, 424; Strickland, Bull. Ent. Res. IV, 140.

Myzorhynchus alboannulatus, James and List., Anoph. Mosq. Ind., 81 (1904).

Perak [*Wright*]; Kuala Lumpur; Port Swettingham; Deli, Sumatra, iii-14. Type in British Museum.

var. *montanus* Stanton and Hacker, Bull. Ent. Res. VII, 273, ♂ ♀ (1916-17).

Ulu Gombak, Selangor, 600 ft.

willmori James in Theob., Monog. Culic. III, 100, ♀; fig. 59, palpus ♀, fig. 60, var. scales; (*Nyssorhynchus*) (1903).

Chris., Ind. Jour. Med. Res. III, 484.

- Nyssorhynchus willmori*, Blanch., Moust., 624: Brun., Rec. Ind. Mus. I, 321; IV, 430
nec *willmori*, Leices.
- Neocellia willmori*, var. *maculosa*, James and List., Anoph. Mosq. Ind., 2nd ed., 112 (1911).
- Neocellia indica*, Theob., Monog. Culic. IV, iii, ♂ ♀ (1907): Brun., Rec. Ind. Mus. IV, 432
- Neocellia dudgeoni*, Theob., Monog. Culic. IV, 112, ♀ (1907).
Brun., Rec. Ind. Mus. IV, 432.
Kashmir, 4800 ft. [*Willmore*]; Dehra Dun, ii, iii; Lahore, Jalpaiguri, 13-vii-07 [*Wallich*]; Kangra Valley, 4500 ft., vi, vii [*Dudgeon*]; Pundaluoya, Ceylon, vii [*Green*]; Kuala Lumpur, Jugra, Malay Peninsula [*t. Leicester*]; Ula Gombak. Type of *maculosa* in Central Malaria Bureau, Kasauli; of *willmori*, *indica* and *dudgeoni* in British Museum: Roper thinks (Bull. Ent. Res. V, 143) that this species is synonymous with *maculatus*, Theob.
- karwari** James in Theob., Monog. Culic. III, 102, ♀; fig. 61 (p. 103), wing; (*Nyssorhynchus*) (1903).
- Anopheles karwari*, James and List., Anoph. Mosq. Ind., 89; pl. xiv, var. chars.; fig., p. 90, wing: Chris., Ind. Jour. Med. Res. III, 469.
- Nyssorhynchus id.*, Leices., Culic. Malaya, 39, ♂ ♀ (*Nyssorhynchus*): Brun., Rec. Ind. Mus. I, 318; IV, 429.
- Anopheles nigrans*, Stanton, J. Lond. Sch. Trop. Med. II, No. 1, 7 (1912).
Karwar, vi [*Cogill*]; Goa, 2000 ft., ii [*Aitken*]; Kuala Lumpur, 27-iv-02 [*Durham*]. Types of *karwari* and *nigrans* in British Museum: Christophers says that in Theobald's later descriptions of his *maculatus* he has confused *maculatus* and *karwari*.
- error** Theob., Monog. Culic. III, 353, ♀ (*Aldrichia*) (1903).
Theob., Gen. Ins., Fasc. 26, 13.
- Aldrichia error*, Brun., Rec. Ind. Mus. I, 322.
- Aldrichinella error*, Theob., Monog. Culic. V, 77: Brun., Rec. Ind. Mus. IV, 433.
"India, probably Calcutta." Type in British Museum, now broken: According to Col. Alcock this is not a good species, the unique type being a *Myzomyiarossii*, Giles, with the abdomen of some other species attached.
- culiciformis** Cogill., Jour. Bomb. Nat. Hist. So. XV, 333 (1903).
- Anopheles culiciformis*, James and List., Anoph. Mosq. Ind., 122, ♂, ? ♀; pl. xv, larval figs: Chris., Ind. Jour. Med. Res. III, 463: Chris. and Chand, Ind. Jour. Med. Res. III, 638, ♂ ♀ all stages descr.; pl. lix (var. chars.) (*Coelodiazesis*).
- Stethomyia culiciformis*, Theob., Monog. Culic. IV, 62: Brun., Rec. Ind. Mus. I, 312; IV, 423.

- Neostethopheles culiciformis*, James and List., *Anoph. Mosq. Ind.*, 2nd ed., 61, figs.
Karwar, Bombay Presidency [*Dr. Cogill*]. *Type*, location unknown to present author.
- pitchfordi** Giles, *Jour. Trop. Med.* VII, 365 (*Pyretophorus*) (1904).
Pyretophorus pitchfordi, Brun., *Rec. Ind. Mus.* I, 313.
Pampang [Whitmore]. *Type* in British Museum.
- asiatica** Leic., *Entom.* XXXVII, 13, ♂ ♀ (*Lophoscelomyia*) (1904).
Chris., *Ind. Jour. Med. Res.* III, 462.
Lophoscelomyia asiatica, Theob., *Monog. Culic.* IV, 92, ♂ ♀:
Brun., *Rec. Ind. Mus.* I, 317; IV, 427: Leices., *Culic. Malaya*, 21, ♂ ♀.
Lophomyia asiatica, Giles, *Jour. Trop. Med.* VII, 366.
Myzorhynchus asiaticus, Strickland, *Bull. Ent. Res.* IV, 141.
Kuala Lumpur. *Type* in British Museum.
- freerae** Banks, *Phil. Jour. Sci.* I, 993, ♀ (*Pyretophorus*) (1906).
Pyretophorus freerae, Theob., *Monog. Culic.* V, 43: Brun., *Rec. Ind. Mus.* I, 312; IV, 423.
Manila, x [*Banks*]. *Type*, apparently unique, No. 5975, in *Entom. coll.*, Bureau of Science, Manila.
- nigrifasciatus** Theob., *Monog. Culic.* IV, 65, ♀ (*Pyretophorus*) (1907).
Pyretophorus nigrifasciatus, Brun., *Rec. Ind. Mus.* IV, 424.
Peshin, India, iv [*Maj. Nurse*]; also occurs in Cyprus. *Type* in British Museum.
- treacheri** Leices., *Culic. Malaya*, 19, ♂ ♀ (1908).
Brun., *Rec. Ind. Mus.* IV, 417.
Malay Peninsula, widely distributed. *Type* in Leicester's collection. Christophers ranks this as synonymous with *aikhenii*, James.
- separatus** Leices., *Culic. Malaya*, 36 (sex not stated, apparently ♀) (1908).
Roper, *Bull. Ent. Res.* V, 145.
Myzorhynchus separatus, Brun., *Rec. Ind. Mus.* IV, 426.
Kuala Lumpur. *Type* in Leicester's collection.
Possibly a variety of *sinensis*, Wied.
- watsoni** Leices., *Culic. Malaya*, 38, ♀ (*Pyretophorus*) (1908).
Pyretophorus watsoni, Brun., *Rec. Ind. Mus.* IV, 424.
Klang [*Dr. Watson*]. *Type*, a unique ♀, in Leicester's collection.
- formosus** Ludl., *Can. Ent.* XLI, 22, ♀ (1909).
Theob., *Monog. Culic.* V, 8, ♀: Brun., *Rec. Ind. Mus.* IV, 416.
Benguet, *Phil. Is.*, iii-o8. *Type* in Army Medical Mus., Washington.
- mcgregori** Banks, *Phil. Jour. Sci.* IV, 548, ♂ (*Kerteszia*) (1910).
Kerteszia mcgregori, Brun., *Rec. Ind. Mus.* IV, 431.

- Basalin Is., Phil. Is., 28 xii-06 [McGregor]. Type in Entom. coll. (No. 6666), Bureau of Science, Manila.
- aurirostris** Watson, Ann. Trop. Med. Parasit. IV, 251, ♀; fig. wing (1910).
Bred, Klang, xii-09.
- fowleri** Chris., Paludism II, 64 (*Neocellia*) (1911).
Neocellia fowleri, id., loc. cit., III, pl. vi, var. chars.
Anopheles fuliginosus, var. *fowleri*, Alcock, Jour. Lond. Sch. Trop. Med. II, No. 3 (1913).
Amritsar; Cent. Prov. India, xi to ix; Punjab; South India.
Christophers thinks more recently (Ind. Jour. Med. Res. III, 466) that this may be the *pallida*, Theob., variety of *fuliginosus*, Giles.
- wellingtonianus** Alcock, Jour. Lond. Sch. Trop. Med. II, 1, 2 text-figs. (*Myzorhynchus*) (1912).
Larut Hills, 4000 ft., Malay Peninsula. Type in British Museum.
- brevipalpis** Roper, Bull. Ent. Res. V, 141, ♂ ♀, fig. 1, wing, proboscis ♀, front claw ♂ (1914).
Strickland, Ind. Jour. Med. Res. III, 201, larva.
Borneo. Types in British Museum. Strickland (*loc. cit.*) suggests *Memnemyia* as a new genus to receive this species on account of minor differences.
- schnuffneri**, Stanton, Bull. Ent. Res. V, 373 ♀; fig. 1, proboscis, palpus (1915).
Lampongs Distr., South Sumatra, v, vi, 1914 [*Dr. Schuffner*].
- hunteri** Strickland, Ind. Jour. Med. Res. IV, 263, ♂ ♀, figs. var. chars. (*Myzorhynchus*) (1917).
Tebong, Negri Sembilan, Malay Peninsula, vi-14; Bandar Penggaram, ix-15; Sungei Simpang, Kanan, Johore State [all *Strickland*]; Singapore [*Dr. Hunter*].
- novumbrosus** Strickland, Ind. Jour. Med. Res. VI, 271, ♂ ♀ (*Myzorhynchus*) (1917).
Malay Peninsula.
- annandalei** Prashad, Rec. Ind. Mus. XV, 123, ♂, pl. xvii, wing, larva, var. chars; life-history also described (1918).
Sureil, Darjiling District, 5000 ft., 28-x-17 [*Annandale* and *Gravely*].
- pampangensis** nom. nov. (*philippinensis* Ludl. preocc.).
Pyretophorus philippinensis, Ludl., Can. Ent. XXXVII, 135 (1905); Brun., Rec. Ind. Mus. I, 313.
Pampanga [*Whitmore*]. Type in Army Med. Mus., Washington.
This species of Miss Ludlow's has to be renamed as *Nysorhynchus philippinensis*, Ludl. is of course an *Anopheles* in the present author's view of the genus.

In an editorial note in the Ind. Jour. Med. Res. IV, 611, mention is made of a new species *A. (Myzorhynchus) similis*, Strickland, from the Malay States. The larva is "described" in three lines, the adult is "described" as being "indistinguishable from *umbrosus*."

Bironella, Theob.

Ann. Mus. Hung. III, 69 (1905).

Theob., Monog. Culic. IV, 120, ♂; fig. ♂, wing (p. 122).

GENOTYPE, *B. gracilis* Theob., by original designation.

gracilis Theob., Ann. Mus. Hung. III, 69, ♂; fig. 3, scales; pl. ii, wing ♂; pl. iii, wing scales ♂ ("♀" lapsus).

Theob., Monog. Culic. IV, 121, ♂: fig. of wing, p. 122:

Brun., Rec. Ind. Mus. I, 322; IV, 433.

Muina, Papua, 31-xii-1900 [*Biro*]. *Type*, unique, in the Hungarian Museum. Howard, Dyar and Knab think this genus may be a transitional form between *Anopheles*, *Megarhinus* and *Uranotaenia* (Mosq. N. and Cent. Am. and W. Ind. 899).

Calvertina Ludlow.

Can. Ent. XLI, 234, emend. from *Calvertia* Ludl., *loc. cit.*, 22 (1909).

Calvertia preocc. Warren in Lepidoptera.

GENOTYPE, *Chagasia lineata* Ludl., by original designation.

lineata Ludl., Can. Ent. XL, 50 (*Chagasia*) (1908).

Calvertia lineata, Ludl., Can. Ent. XLI, 22.

Calvertina lineata, *Id.*, *loc. cit.*, XLI, 234: Theob., Monog.

Culic. V, 77: Brun., Rec. Ind. Mus. IV, 433.

Pangasinan, viii. *Type* in Army Med. Mus., Washington.

Sect. CULICINI.

This "Section" comprises the whole of the subfamily Culicinae except the Anophelines, but is not herein assigned equivalent or any definite taxonomic value, being only adopted as a temporary convenience to separate them from the Anophelines.

Megarhinus, R. Desv.

Mém. Soc. Hist. Nat. Paris III, 412 (1827).

Macq., Hist. Nat. Dipt: Skuse, Proc. Linn. Soc.

N.S. Wales III, 1720: Theob., Monog. Culic. I,

215; III, 113; IV, 128, tab. spp.: Blanch., Moust., 200, 221, tab. spp.: Theob., Gen. Ins., Fasc. 26, 12: Brun., Rec. Ind. Mus. I, 323; IV, 435; X, 65: How., Dyar and Knab, Mosq. N. Amer., etc., IV, 927.

Megarhina, Giles, Handbk., etc., 119; 2nd ed., 265: *Auctl.*

GENOTYPE, *Culex haemorrhoidalis* Fab., by original designation.

The genus *Megarhinus* is not Oriental, but the species belonging to *Toxorhynchites* were originally assigned to it, therefore a few references are given here.

Toxorhynchites, Theob.

Monog. Culic. I, 244 (1901).

Theob., *loc. cit.*, III, 119, notes on life-history; V, 95, notes and tab. spp.: *Id.*, Gen. Ins., Fasc. 26, 13: Blanch., Moust., 229: Leices., Culic. Malaya, 59: Brun., Rec. Ind. Mus. I, 324; IV, 435; X, 72: Giles, Handbk., etc., 2nd ed., 278. GENOTYPE, *T. brevipalpis* Theob., by original designation.

Teromyia Leices., Culic. Malaya, 49 (1908).

GENOTYPE, *T. acaudata*, Leic., sp. nov., *loc. cit.* by my designation, 1914.

Brun., Rec. Ind. Mus. IV, 438, notes.

Worcesteria, Banks, Phil. Jour. Sci. I, 779 (1906).

GENOTYPE, *W. grata* Banks, sp. nov., *loc. cit.*

Theob., Monog. Culic. V, 110: Brun., Rec. Ind. Mus. I, 326; IV, 439; X, 73.

splendens Wied., Zool. Mag. III, 2, ♂ (*Culex*) (1819).

Culex splendens, Wied., Auss. Zweifl. I, 3, ♂: Macq., Hist. Nat. Dipt. I, 33.

Megarhinus splendens, Sch., Reise Novara, Dipt., 31: Wulp, Mid-Sumatra, Dipt. 8; pl. i, 2, wing: Theob., Monog. Culic. I, 235, pl. viii, 31, ♀, full ins. col: Giles, Handbk., etc., 124; 2nd ed., 271 (*Megarhina*): Blanch., Moust., 225: Brun., Rec. Ind. Mus. I, 324.

Toxorhynchites splendens, Brun., Rec. Ind. Mus. IV, 437.

Java, Sumatra, Singapore. *Types* in Wiedemann's and Westermann's collections. Not recorded in recent years, probably one or more of the more recently described species may be synonymous with it.

amboinensis Dol., Nat. Tijd. Ned. Ind. XIV, 381, ♂, pl. v, 5 (*Culex*) (1857).

Megarhina amboinensis, Giles, Handbk., etc., 133; 2nd ed., 276.

Megarhinus amboinensis, Theob., Monog. Culic. I, 243 ♂:

Blanch., Moust., 226, ♂ ♀ : Brun., Rec. Ind. Mus. I, 323, notes.

Toxorhynchites amboinensis, *id.*, *loc. cit.*, IV, 435.
Amboina, Ceram. Type in Vienna Museum.

subulifer Dol., Nat. Tijd. Ned. Ind. XIV, 382, ♀, pl. v, 2 (*Culex*) (1857).

Megarhinus subulifer, Theob., Monog. Culic. I, 242: Giles, Handbk., etc., 125; 2nd ed., 272 (*Megarhina*).

Toxorhynchites subulifer, Theob., Monog. Culic. V, 101: Brun., Rec. Ind. Mus. IV, 437: Edw., Bull. Ent. Res. VII, 202.
Amboina, Ceram. Type in Vienna Museum.

In my Catalogue this species was given as synonymous with *immisericors*, Walk., but, following other authors, the two were ranked as distinct in the Supplement. Edwards notes¹ that there is nothing in Doleschall's description of *subulifer* to prevent it being the same as *immisericors*, in which case the Dutch author's name takes precedence. He adds that Doleschall also described *amboinensis*, which he afterwards took to be the ♂ of his *subulifer*, and if that opinion should prove correct, the species is distinct from *immisericors*. This would be true but if *amboinensis* and *subulifer* are respectively ♂ and ♀ of a single species, the former name must be used because the description precedes that of *subulifer* by a page (pp. 381 and 382 respectively). Until the identity of all three alleged species is settled they may be regarded as provisionally distinct.

immisericors Walk., Proc. Linn. Soc. Lond. IV, 90, ♂ (*Megarhinus*) (1860).

Megarhinus immisericors, Theob., Monog. Culic. I, 225, ♂ ♀, pl. vii, 28, ♂, full ins. col.; pl. ix, 33, ♀ (as *M. gilesii*): Giles, Handbk., etc., 129; 2nd ed., 273 (*Megarhina*).

Megarhinus gilesii, Theob., Monog. Culic. I, 227, ♂ ♀; pl. ix, 33, ♀, full ins. col. (1901).

Toxorhynchites immisericors, Theob., Monog. Culic. III, 123, fig. 67, larva, pupa; IV, 141; V, 97, ♀: *id.*, Rec. Ind. Mus. IV, 4, ♀ redescr.: Blanch., Moust., 230, ♂: Brun., Rec. Ind. Mus. I, 324; IV, 436, notes: Green, Spol. Zeyl. II, 159, pl. life-history: Paiva, Rec. Ind. Mus. V, 187, larval habits: Edw., Bull. Ent. Res. IV, 222.

Toxorhynchites gilesii, Theob., Monog. Culic. V, 99.
Brun., Rec. Ind. Mus. IV, 435.

Toxorhynchites argenteotarsis, Ludl., Can. Ent. XXXVIII, 367, ♀ (1906).

Theob., Monog. Culic. V, 100, ♀: Brun., Rec. Ind. Mus. IV, 435.

¹ Bull. Ent. Res. IV, 223 under *immisericors*.

Megarhinus immisericors, Walk., or *amboinensis*, Dol., Ost. Sack., Berl. Ent. Zeits. XXVI, 96.

Worcesteria grata. Banks, Phil. Jour. Sci. I, 780, ♂ ♀, pl. palpus ♂, genitalia ♂, cross-veins ♂ ♀, scales, etc. (1906): Theob., Monog. Culic. V, 111: Brun., Rec. Ind. Mus. I, 326.

Culex regius, Thwaites, Proc. Linn. Soc. Lond. VIII, 102 (1864).¹

Toxorhynchites regius, Edw., Bull. Ent. Res. VII, 202.

? *Megarhinus subulifer*, Dol., Nat. Tijd. Ned. Ind. XIV, 382 (1857).

Common and widely distributed in the East. Bhim Tal, 19-22-x-06 [Annandale]; Sikkim, 1800 ft., vi [Dudgeon]; Sylhet, 5-v-05 and vi; Chittagong, 3-ix-08; Aijal, Lushai Hills, 3600 ft., 24-iv-04; Upper Burma [Watson]; Calcutta, i, vi, vii, viii [Annandale, etc.]; Peradeniya (Botan. gardens), 1500 ft., not uncommon; Pundaluoya, Ceylon, 4000 ft. [Green and T. B. Fletcher]; Trincomalee Hot Wells, Travancore, Mysore, Macassar; Andaman Is., Waigion; North Ceram; Mysol; Settleberg (Huon Gulf), Papua [Biro]; Bago, Negros Occidental, Cebu (Phil. Is.). Types of *immisericors* and *gilesii* in British Museum in bad condition; of *C. regius* in Hope coll.; Oxford University; of *W. grata* in Entom. Coll., Bureau of Science, Manila, Nos. 6071, ♂ ♀.

inornatus Walk., Proc. Linn. Soc. Lond. VIII, 102, ♂ (*Megarhinus*) (1865).

Megarhinus inornatus, Theob., Monog. Culic. I, 223, ♂ ♀, pl. vii, 26, ♂, 25, ♀ (both full ins. col.): Giles, Handbk, etc., 137, 2nd ed., 271, ♂ ♀ (*Megarhina innornata*, lapsus): Blanch., Moust., 223, ♂ ♀.

Toxorhynchites inornatus, Brun., Rec. Ind. Mus. I, 325; IV, 436. Papua. Type ♂ ♀ in British Museum.

leicesteri Theob., Entom. XXXVII, 36, ♂ ♀ (1904).

Theob., Monog. Culic. IV, 142, ♂ ♀: Leices, Culic. Malaya, 59, ♂ ♀: Brun., Rec. Ind. Mus. I, 325; IV, 437.

Kuala Lumpur; Papua. Types in British Museum.

¹ I decline to accept Edwards' suggestion to adopt the name *regius* by Ternent (1859) in place of *immisericors*. A "description" of a line or two (in this case six words only) is no scientific description at all properly so called, even in the case of a species recognizable by it. Moreover, there is, as Edwards himself admits, a second though rare species of *Toxorhynchites* occurring in Ceylon and presumably the "description" of *regius* ("one with a formidable hooked proboscis") would equally apply to the other species, *minimus*, Theob., the character being generic. Ternent is not known, so far as I am aware, as an entomologist, the species was not scientifically described nor is "Ceylon" the journal in which it was published a scientific publication.

In the same paper (Bull. Ent. Res. VII, 202) Edwards notes having found on re-examining Walker's types of *immisericors*, from Celebes, that they are different from the species generally known from the East under this name, and he believes them probably identical with *subulifer*, Dol. I do not care, without a close examination of Doleschall's type if such be still in existence, to alter the present view of *immisericors* (that is, Theobald's), but if Edwards' latest opinion should prove correct, it leaves Theobald's form without a name.

- metallicus** Leices. in Theob., Entom. XXXVIII, 37, ♂ ♀ (1904).
 Leices., Culic. Malaya, 61, ♂ ♀: Theob., Monog. Culic. IV,
 144, ♂ ♀: Brun., Rec. Ind. Mus. I, 325; IV, 437.
 Kuala Lumpur [*Leicester*]. *Types* in British Museum.
- lewaldii** Ludl., Can. Ent. XXXVI, 233, ♂ (*Megarhinus*) (1904).
 Theob., Monog. Culic. IV, 139, ♂; Brun., Rec. Ind. Mus. IV,
 437.
Megarhinus lewaldii, Brun., Rec. Ind. Mus. I, 324.
Megarhinus waldii, Blanch., Moust., 625.
 Salog (Guimaras Is.), Philippine Is., iv [*Lewald*]. *Type* in
 Army Medical Museum, Washington.
- minimus** Theob., Jour. Bomb. Nat. Hist. Soc. XVI, 237, ♂,
 pl. A, fig. 1, palpus, wing, tip of abdomen (1905).
 Brun., Rec. Ind. Mus. I, 324; IV, 437.
Megarhinus minimus, Theob., Monog. Culic. IV, 138, ♂: Brun.,
 Rec. Ind. Mus. I, 324.
 Yatiyantota, Ceylon. *Type* in British Museum.
- acaudata** Leices., Culic. Malaya, 49, ♂ ♀ (*Teromyia*) (1908).
Teromyia acaudata, Brun., Rec. Ind. Mus. IV, 438.
 Singapore (*Finlayson*). *Types* in *Leicester's* collection.
- quasiferox** Leices., Culic. Malaya, 51, ♂ ♀ (*Teromyia*) (1908).
 Edw., Bull. Ent. Res. IV, 223.
Teromyia quasiferox, Brun., Rec. Ind. Mus. IV, 439.
Toxorhynchites javaensis, Theob., Tijd. v. Ent. LIV, 233 (1911).
 Brun., Rec. Ind. Mus. IV, 436.
 Java, Singapore, Malay Peninsula. *Type* of *quasiferox* in
Leicester's collection; of *javaensis*, a unique ♀ from Java, in
 Amsterdam Museum.
- magnificus** Leices., Culic. Malaya, 54, ♂ ♀ (*Teromyia*) (1908).
 Edw., Bull. Ent. Res. IV, 223.
Teromyia magnificus, Brun., Rec. Ind. Mus. IV, 438.
Teromyia funesta, Leices., Culic. Malaya, 58, ♂ (1908).
 Malay Peninsula. *Type* of *magnificus* in *Leicester's* coll.,
 of *funestus*, a unique ♂ bred from bamboo at Kuala Lumpur, in
 same collection.
- raris** Leices., Culic. Malaya, 56, ♂ (*Teromyia*) (1908).
Teromyia raris, Brun., Rec. Ind. Mus. IV, 439.
 Ulu Klang, Fed. Malay States, a unique bad specimen, the
type apparently lost, not in *Leicester's* collection.
- ater** Daniels, Studies from Institute for Medical Research, Fed.
 Malay Sts., III, 265, ♂ ♀ (*Teromyia*) (1908).
Teromyia ater, Brun., Rec. Ind. Mus. IV, 438.
 Pahang, Malay Peninsula. *Type* apparently lost, not in
Leicester coll.

Mucidus, Theob.

Monog. Culic. I, 268 (1901).

Theob., Gen. Ins., Fasc. 26, 17: *id.*, Monog.
 Culic. III, 132, larva; V, 125 tab. spp.: Giles,

Handbk., etc., 2nd ed., 346; Blanch. Moust., 243; Leices., Culic. Malaya, 69; Brun., Rec. Ind. Mus. IV, 440; X, 39, 66; Alcock, Ent. Med. Off., 104.

GENOTYPE, *Culex alternans* Westw., by personal designation.

laniger Wied., Dipt. Exot. 9, ♀ (*Culex*) (1821).

Culex laniger, Id., Auss. Zweifl. I, 5, ♀; Macq., Dipt. Exot. I, 2, 176; Giles, Handbk., etc., 211.

Mucidus laniger, Blanch., Moust., 244, ♀ (1905); Edw., Bull. Ent. Res. IV, 224.

Mucidus mucidus, Leices. (*nec* Karsch), Culic. Malaya, 69, ♂ ♀ (1908).

Java, Coromandel, Deli, Sumatra, iii-14. Type in Copenhagen Museum (*t. Vienna Museum, in litt.*); that of *mucidus*, Leicester, in Leicester's coll.

alternans Westw., Ann. Soc. Ent. Fran. IV, 681 (*Culex*) (1835).

Giles, Handbk., etc., 2nd ed., 347, ♂ ♀, pl. xii, 2, wing ♀.

Culex commovens, Walk., Ins. Saund. Dipt. pt. 5, 432 (1896).

Culex hispidosus, Skuse, Trans. Linn. Soc. N.S. Wales, 1726 (1889).

Papua. Type in Copenhagen Museum.

mucidus Karsch, Ent. Nachr., 25 (*Culex*) (1887).

nec Mucidus mucidus, Leices.

Culex mucidus, Giles, Handbk., etc., 214.

Mucidus mucidus, Theob., Monog. Culic. I, 272, ♀, pl. xi, 42, full ins. col., pl. B, wing scales (1901); Giles, Handbk., 2nd ed., 549, ♀, pl. xii, 3, wing ♀; Brun., Rec. Ind. Mus. IV, 440; Edw., Bull. Ent., Res. II, 246.

Kuala Lumpur; Philippine Is. Type in Berlin Museum.

africanus, Theob., Monog. Culic. I, 274 (1901).

grahami, *id.*, *loc. cit.*, V, 127 (1910).

scatophagoides Theob., Monog. Culic. I, 277, ♀, pl. E, wing scales, fig. 81 (p. 278), wing, thorax, scales (1901).

Giles, Handbk., etc., 2nd ed., 348, ♀, pl. xii, 1a, full ins., 2a, venation; Blanch., Moust., 245, ♀; Brun., Rec. Ind. Mus. I, 327; Edw., Bull. Ent. Res. II, 246.

sudanensis, Theob., 3rd Rep. Welle. Labor., 252 (1908).

Bauria, Bengal, 17-viii-07 [*Tyrie*]; Damukdia Ghat, Ganges Riv., 22-viii-07 [*Paiva*]; Purnea, 5-viii-07 [*Paiva*]; Moradabad, N.W. Pro. Ind., ix [*Close*]; Myingan, Burma. Type in British Museum.

Pardomyia, Theob.

Monog. Culic. IV, 280 (1907).

Brun., Rec. Ind. Mus. IV, 460; X, 68.

GENOTYPE, *P. aurantia* Theob., sp. nov., by original designation.

Ekrinomyia, Leices., *Culic. Malaya*, 71 (1908).

GENOTYPE, *E. aureostriata* Leices, sp. nov., by original designation.

aurantia Theob., *Monog. Culic.* IV, 280, ♀ (1907).

Edw., *Bull. Ent. Res.* IV, 224.

Ekrinomyia aureostriata, Leices., *Culic. Malaya*, 71, ♂ ♀ (1908).

Brun., *Rec. Ind. Mus.* IV, 440.

Kuching, Sarawak, xi [Barker]; Kuala Lumpur, Klang.

Type of *aurantia* in Brit. Mus., of *aureostriata* in Leicester coll.

quadripunctis Ludl. in Theob., *Monog. Culic.* V, 608 (1910).

Parang, Mindanao, Philippine Is., 26-x. *Type* in Army Medical Mus., Washington.

Armigeres, Theob.

Monog. Culic. I, 322 (1901).

Armigeres, Giles, *Handbk.*, etc., 2nd ed., 384:

Edw., *Bull. Ent. Res.* IV, 224: VII, 202, diag. and tab. spp.: Brun., *Rec. Ind. Mus.* X, 55

Desvoidya, Blanch., *Comp. Rend.* LIII, 1035 (1901),

nom. nov., for *Armigeres*, Theob., under presumption of preoccupation by *Armiger*, Hartmann,

1840-1842, in *Mollusca*: Blanch., *Moust.*, 265:

Theob., *Gen. Ins.*, Fasc. 26, 17: *id.*, *Monog.*

Culic., IV, 163, tab. spp.; V, 141, tab. spp.:

Brun., *Rec. Ind. Mus.* I, 327: Leices., *Culic.*

Malaya, 74.

Desvoidea, Theob., *Monog. Culic.* III, 134, larva.

Blanchardiomyia, Brun., *Rec. Ind. Mus.* IV, 440,

nom. nov., notes; X, 156

GENOTYPE of *Armigeres*, *Culex obturbans*, Walk.,

by original designation.

On the structure of the larva and of the male genitalia Edwards provisionally recognises this group as a valid genus, whilst admitting that the female adults cannot be distinguished from *Ochlerotatus*, *Stegomyia* and *Aedes*

The name *Armigeres* is not really preoccupied (*t. mihi*, *Rec. Ind. Mus.* X, 55), *Armiger* not being a homonym. Edwards also notes this fact, but in his latest paper he sinks the genus under *Stegomyia*.

obturbans Walk., *Proc. Linn. Soc. Lond.*, IV, 91, ♀ (*Culex*), (1860).

Culex obturbans, Giles, *Handbk.*, 338.

Armigeres obturbans, Theob., *Monog. Culic.* I, 323, ♂ ♀; fig.

104, wing, 105, palpus ♀ (incorrect), unguis ♀, ♂ genitalia;

fig. 106, ♂, unguis: Edw., *Bull. Ent. Res.* VII, 205, fig. 2,

♂ genit.

Desvoidea obturbans, Theob., *Monog. Culic.* III, 138, fig. 75,

middle claws ♂, fig. 77, corrected ♂ palpus, fig. 78, ♂

clasper: Theob., *loc. cit.*, IV, fig. 47, head of larva; fig. 48,

siphon of larva.

Desvoidya obturbans, Blanch., Moust., 266, ♂ ♀; Leices, Culic. Malaya, 75, ♂ ♀; notes on varietal forms: Brun., Rec. Ind. Mus. I, 328.

Blanchardiomyia obturbans, Brun., *loc. cit.*, IV, 442 notes.

Culex ventralis, Walk., Proc. Linn. Soc. Lond., V, 144 (1861): Giles, Handbk., 339.

nec ventralis, Walk., Proc. Linn. Soc. Lond., VIII, 103.

Armigeres ventralis, Theob., Monog. Culic., pl. xv, 57, full ♀, ins. col.: Giles, Handbk., etc., 2nd ed., 385, ♂ ♀, pl. xiv, 11, venation, 12, claws.

Naini Tal (Giles); Kurseong, 5-viii-08 [Annandale]; Sukna, 1-viii-08 [Annandale]; Sylhet [Hall]; Lushai Hills; Behar [Green]; Mozufferpore; Rajmahal, 31-vii-07 [Hodgart]; Meerut, 25-iv-05 [Brunetti]; Madras [Cornwall]; Trivandrum, 14-xi-08 [Annandale]; Travancore [James]; Madulsima, Ceylon, 26-ix-07 [Green]; Selangor, 28-x-99 [Butler]; Perak [Wray]; Singapore, 25-viii-99; Batavia, 27-vi to 9-vii-06 [Brunetti]; Semarang, i, ii, iii-04 and viii-05 [Jacobson]; Waria River, British Papua [Fleming Jones]; Amboina, Celebes, Mysol, Waigiou, North Ceram, Tinghai, Formosa, West Lake, Hankow, 28-viii-07 [Cornford]; Foochow [Rennie]. Type in British Museum, a ♀.

Uranotaenia malayi, Theob., Monog. Culic. II, 258 (1901): Giles, Handbk., 2nd ed., 494, ♀: Blanch., Moust., 410.

Armigeres malayi, Edw., Bull. Ent. Res. VII, 207, fig. 3, ♂ genit.

Desvoidea fusca, Theob., Monog. Culic. III, 135 (1903), types only.

Selangor, 28-x-99; Kuala Lumpur. Type in British Museum.

fusca Theob., Monog. Culic. III, 135, ♂ ♀, fig. 75, middle claws ♂, palpus ♂, fig. 76, pupal chars., pl. xvii, larval figs. (*Desvoidea*), (1903).

Desvoidea fusca, Theob., Monog. Culic. IV, 165; V, 142: Brun., Rec. Ind. Mus. I, 327; Leices., Culic. Malaya, 78, ♂ ♀.

Blanchardiomyia fusca, Brun., Rec. Ind. Mus. IV, 441, notes.

Sylhet, iii to vi [Hall]; Lushai Hills, Assam, viii [Macleod]; Calcutta, v, vii to xii; Jugra and Kuala Lumpur [both Durham]; Angeles, Phil. Is. [Whitmore]. Types in British Museum.

This is a species of uncertain validity. Edwards says that only the type ♂ and ♀ in the British Museum are synonymous with *malayi*, but as other specimens have been recorded from various and widely distributed localities, these latter may represent a valid species, which of course will require a new name.

joloensis Ludl., Can. Ent. XXXVI, 236 (*Desvoidea fusca*, var. *joloensis*), (1904).

Desvoidea fusca, var. *joloensis*, Theob., Monog. Culic. IV, 165: Brun., Rec. Ind. Mus. I, 327.

Desvoidya joloensis, Theob., Monog. Culic. V, 143.

- Armigeres joloensis*, Edw., Bull. Ent. Res. IV, 225 (1908).
Blanchardiomyia joloensis, Brun., Rec. Ind., Mus. IV, 441.
Desvoidya jugraensis, Leices., Culic. Malaya. 77, ♂ ♀ (1908).¹
Blanchardiomyia jugraensis, Brun., Rec. Ind. Mus. IV, 441.
 Jolo, Philippine Is., Jugra, Ampang and E. Coast of Malay Peninsula; Borneo. Type of *joloensis* in Army Medical Museum, Washington, of *jugraensis*, in Leicester coll. Stanton records the *jugraensis* form from Delhi, Sumatra, iii, 14.
- aureolineatus** Leices., Culic. Malaya, 79, ♀ (*Desvoidya*), (1908).
Blanchardiomyia aureolineata, Brun., Rec. Ind. Mus. IV, 441, notes.
Armigeres aureolineatus, Edw., Bull. Ent. Res. VII, 205, fig. 1, ♂ genit.
 Ampang, near Kuala Lumpur. Type in Leicester coll.
- apicalis** Theob., Rec. Ind. Mus. IV, 5, ♀ (*Desvoidia*), (1910).
Armigeres apicalis, Edw., Bull. Ent. Res. IV, 225.
Blanchardiomyia apicalis, Brun., Rec. Ind. Mus. IV, 441.
Stegomyia crassipes, Theob. (nec Wulp), Monog. Culic., I, 320.
 ♀, pl. xxxiv, 134, ♀, full ins. col. (1901): Giles, Jour. Trop. Med. VII, 367: *id.*, Handbk., etc., 2nd ed., 381: Brun., Rec. Ind. Mus. I, 329; IV, 445.
 Balighai, near Puri, Orissa, 24-x-08 [*Annandale*].
 Type of *apicalis* in Indian Museum, of *crassipes* in British Museum. Erroneously identified as "*crassipes*, Wulp" it is recorded from Thayetmo, Upper Burma, viii [*Watson*] and from Pampanga, Philippine Is.
- moultoni** Edw., Bull. Ent. Res. V, 125, ♂ ♀ (1914-15).
 Kuching, Sarawak, 19-ii-6-iii-1914 [*Moulton*]; Jugra, 15-ix-1904 [*Leicester*]. Types in British Museum.
- brevitibia**, *id.*, *loc. cit.*, 125, ♀ (1914-15); VII, 204, ♂.
 Tabuan Swamp, Kuching, Sarawak, 25-26-ii-1914 [*Moulton*]. Type in British Museum.
- hybridus**, *id.*, Bull. Ent. Res. V, 126, ♂ ♀ (1904-15).
 Kuching, Sarawak, ♂ ♀ [*Moulton*]. Types in British Museum.
- conjungens**, *id.*, *loc. cit.*, 126, ♂ ♀ (1914-15).
 Kuching, 25-ii-5-iii-1914 [*Moulton*]. Types in British Museum.
- confusus**, *id.*, *loc. cit.*, 283, ♂ ♀ (1914-15).
 Kuching, Sarawak [*Moulton*, *Hewitt*, ? *Lewis*]. Types in British Museum.
- kuchingensis**, *id.*, *loc. cit.*, 283, ♂ ♀ (1914-15).
 Kuching environs, Sarawak, 24 and 27-vii-14 [*Moulton*]. Type in British Museum.

¹ Edwards says (Bull. Ent. Res. V, 283) that *jugraensis* is not a synonym of *joloensis* as he at first supposed, adding "specimens of the latter represented a colour variation of *obturans*."

breinli Taylor, Tr. Ent. So., 186 (*Neosquamomyia*), (1914).
Papua.

durhami Edw., Bull. Ent. Res. VII, 206, ♂ ♀ (1916-17).

Bukit Kutu, Fed. Malay States, 3300 ft., 11-v-93; type ♂ ♀
bred from larvae in tub near bungalow; 17-v-03 [*Leicester*],
20-ii-03 [*Durham*]. Types in British Museum.

maiae, *id.*, *loc. cit.*, 208, ♂ ♀ (1916-17).

Gombak Rd., Kuala Lumpur, 29-ii-04, 5 and 12-xii-03
[*Leicester*]; Pahang Rd., jungle, 6 miles from Kuala Lumpur,
15-v-04 (type ♂).

Banksinella, Theob.

Monog. Culic. IV, 468 (1907).

Brun., Rec. Ind. Mus. IV, 477 (*Banksiella*,
lapsus); X, 56: Theob., Monog. Culic. V, 402,
tab. spp.

GENOTYPE, *Culex luteolateralis* Theob., by origi-
nal designation.

luteolateralis Theob., Monog. Culic. II, 71, ♂ ♀, pl. xxvii, 108,
♀, full ins. col. (*Culex*), (1901).

Banksinella luteolateralis, Theob., Monog. Culic. IV, 469, ♀:
Brun., Rec. Ind. Mus. IV, 477 (*Banksiella*, *lapsus*): Edw., Bull.
Ent. Res. II, 245; III, 6.

Culex luteolateralis, Giles, Handbk., etc., 2nd ed., 448, ♂ ♀:
Blanch., Moust., 278, ♂ ♀: Leices.; Culic. Malaya, 160, ♂ ♀:
Brun., Rec. Ind. Mus. I, 349

Taeniorhynchus lineatopennis, Ludl., Can. Ent. XXXVII, 133
(1905). Theob., Monog. Culic. IV, 489, ♀: Brun., Rec. Ind.
Mus. IV, 479.

Sylhet, 23-xi-04, 5-x-14 [*Hall*]; Kuala Lumpur [*Leicester*];
Perak [*Wray*]; Pampanga, Philippine Is. [*Whitmore*]; Manila
[*Banks*]; Luzon: Bayambang (Pangasinan, Philippine Is.), ix
[*Chamberlain*]. Occurs also in Durban and Mashonaland.

Type of *luteolateralis* in British Museum; of *lineatopennis* in
Army Medical Museum, Washington.

B. chrysothorax, Theob., was at first thought to be the ♂ of
luteolateralis, but Edwards notes (Bull. Ent. Res. IV, 47) that
it is the ♂ of *taeniarostris*, an African species.

Stegomyia, Theob.

Howard's "Mosquitoes, etc." 233 (Jan. 1, 1901).
Theob., Monog. Culic. I, 285, tab. spp.; 284,
map of distribution; IV, 170, 171, list and tab.
spp.; V, 151, tab. spp.: *id.*, Gen. Ins., Fasc.
26, 18: Giles, Handbk., etc., 2nd ed., 368;
369, tab. spp.: Blanch., Moust., 247, fig. 194,
larva; 248, tab. spp.: Brun., Rec. Ind. Mus. I,

328; IV, 443: Leices., Culic. Malaya, 81, tab. Malay spp.: Alcock, Ent. Med. Off., 100: Edw., Bull. Ent. Res. VII, 203.

GENOTYPE, *Culex fasciatus* F., as *calopus*, Mg.

Scutomyia, Theob., Entom. XXXVII, 77 (1904).

Theob., Monog. Culic. IV, 196; V, 199, tab. spp.: *id.*, Gen. Ins., Fasc. 26, 19: Leices., Culic. Malaya, 105: Brun., Rec. Ind. Mus. I, 336; IV, 452; X, 71.

GENOTYPE, *Culex sugens* Wied., by my own designation.

Quasistegomyia, Theob., 2nd Rep. Gord. Coll. Wellcome Laborat. 69 (1916).

Theob., Monog. Culic. IV, 165; V, 132, tab. spp.: Brun., Rec. Ind. Mus. IV, 443; X, 70.

GENOTYPE, *Q. unilincata* Theob., sp. nov., by original designation.

Gymnometopa, Coq., Proc. Ent. Soc. Wash. VII, 183 (1906).

GENOTYPE, *Stegomyia mediiovittata* Coq., by original designation.

Kingia, Theob., Monog. Culic. V, 135 (1910).

Brun., Rec. Ind. Mus. X, 63.

GENOTYPE, *Stegomyia luteocephala* Newstead, by original designation.

fasciata F., Syst. Antl., 36 (*Culex*), (1805).

Culex fasciatus, Auctt. var.

Stegomyia fasciata, Theob., Monog. Culic. I, 289, ♂ ♀; figs. 86—89 var. chars.; map of distribution, p. 292; pl. xiii, 49, ♂, 50, ♀ both full ins. col.; pl. B, wing scales: Theob., Gen. Ins., Fasc. 26, pl. i, ii, ♀, full ins. col.: Giles, Handbk., etc., 2nd ed., 372; pl. xiv, 2, venation, 3, thorax: Leices., Culic. Malaya, 85, ♂ ♀: Brun., Rec. Ind. Mus. I, 330 and IV, 446, copious notes.

Culex calopus, Meig., Syst. Besch. I, 3 (1818): R. Desv., Ess. Culic. 407 (1827): Giles, Handbk., etc., 237.

Stegomyia calopus, Blanch., Moust., 249, ♂ ♀; fig. 196, ♂, 195, ♀, 197—201, var. chars.: Ludl., Mosq. Phil. Is. 33.

Aedes calopus, Howard, Dyar and Knab, Mosq. N. Amer. IV, 824.

Culex taeniatus, Wied., Auss. Zweifl. I, 10, ♂ ♀ (1828).

C. konoupi, Brullé, Exped. Morea. Zool. III, 289, pl. xlvi, 1 (1832).

C. mosquito, R. Desv., Ess. Culic. 407 (1827): Arrib., Revist. La Plata, 60 (1891).

C. viridifrons, Walk., List Dipt. Brit. Mus. I, 3, ♀ (1848).

C. formosus, *id.*, loc. cit., I, 4, ♀ (1848).

C. inexorabilis, *id.*, loc. cit., I, 4, ♀ (1848).

C. excitans, *id.*, loc. cit., I, 4, ♀ (1848).

C. exagitans, *id.*, Ins. Saund. Dipt. pt. 5., 430, ♀ (1856).

C. impatibilis, *id.*, Proc. Linn. Soc. Lond. IV, 91, ♂ (1860). Giles, Handbk., etc., 255.

C. toxorhynchus, Macq., Dipt. Exot. I, 25 (1838).

C. annulitarsis, *id.*, loc. cit., Supp. I, 136, ♀ 1838).

- C. bancroftii*, Skuse, Proc. Linn. Soc. N.S. Wales III, 1740 (1886).
C. elegans, Ficalbi, Bull. Soc. Ent. Ital., 251 (1896).
C. rossii, Giles, Jour. Trop. Med. 64 (1899).
C. luciensis, Theob., Monog. Culic. I, 297, ♂ ♀; pl. xiii, 50 (only the separate figure of a leg), (1901).
C. queenslandensis, *id.*, *loc. cit.*, I, 297, ♀ (1901).
 Subspecies *persistans* Banks, Phil. Jour. Sci. I, 996 (1906), Brun. Rec. Ind. Mus. I, 331.

The references to this species as *Stegomyia fasciata* are so numerous and so well known that it is unnecessary to list them all here.¹ The absolute identity of *fasciata* with *calopus*, Mg, is not proved, and as the description of Villiers' *Culex fasciatus* is quite unintelligible and the type non-existent no possible gain is obtained by endeavouring to alter this mosquito's well-known name. The Kertesz Catalogue does not mention Villiers' species, which, moreover, may not even have been a *Culex* at all. Messrs. Howard, Dyar and Knab in their monumental work on the North American Mosquitoes actually seriously propose to change the name to *argenteus*, Poiret, Jour. de Physique xxx, 245 (1787). *S. fasciata* is the sole carrier of yellow fever.

- variegata* Dol., Nat. Tijd. Ned. Ind. XVII, 77 (*Culex*), (1858).
Stegomyia variegata, Edw., Bull. Ent. Res. VII, 209.
Culex scutellaris, Walk., Proc. Linn. Soc. Lond. III, 77 (1859).
Culex zonatipes, Walk., Proc. Linn. Soc. Lond. V, 229 (1861).
 Giles, Handbk., etc., 287.
Stegomyia pseudoscutellaris, Theob., Entom. XLIII, 156 (1910).
 Amboina, Aru, Papua, New Caledonia, Christmas Is., the Solomon Is., Samarai Is. and the Fiji Is.

In Edwards' latest opinion the true *Culex scutellaris* of Walker is not the common oriental species generally known as such, but is identical with the *zonatipes* of Walker, and the *pseudoscutellaris* of Theob. It has not, however, been possible for him to examine Doleschall's type of *C. variegatus*, and the identity must be received with some caution. I have already accepted his previous synonymy of the generally accepted "*scutellaris*, Walk." with *albopicta*, Skuse. Type of *variegata*, location unknown; of *scutellaris*, Walk., *zonatipes*, Walk., and *pseudoscutellaris*, Theob., in British Museum.

- albopicta* Skuse, Ind. Mus. Notes III, 20 (*Culex*), (1895).
Culex albopictus, Giles, Handbk., etc., 228.
Stegomyia albopicta, Edw., Bull. Ent. Res. VII, 209.
Stegomyia albopicta, Theob., Monog. Culic. I, 298: Brun., Rec. Ind. Mus. I, 334.
 nec *Stegomyia scutellaris*, Walk., vera.
Stegomyia scutellaris, Theob. nec Walk., Monog. Culic. I, 298, ♂ ♀, fig. 91, ♂ claws; pl. xiv, 53, ♀ (full ins. col.), (1901).

¹ Howard, Dyar and Knab give nearly 150 references.

Id., id., Giles, Handbk., etc., 2nd ed., 374, ♀, pl. xiv, 4, venation, 5, head, thorax, ♂: Blanch., Moust. 257, ♂: Leices., Culic. Malaya, 86, ♂ ♀: Brun., Rec. Ind. Mus. I, 333; IV, 448, notes.

Culex scutellaris, Giles, Handbk., etc., 224.

Culex notoscriptus, Skuse, Proc. Linn. Soc. N.S. Wales, III, 1738 (1889).

Stegomyia notoscripta, Theob., Monog. Culic. I, 286, ♂ ♀; fig. 84, wing ♀; 85, wing scales; III, 145: Blanch., Moust., 257, ♂ ♀; Brun., Rec. Ind. Mus. I, 336; IV, 452.

Scutomyia notoscripta, Theob., Monog. Culic. IV, 198; V, 200: Giles, Handbk., etc., 225.

? *Culex variegatus*, Giles, Handbk., etc., 282.

Subspecies *samarensis* Ludl., Jour. N. Yk. Ent. Soc. XI, 138 (1903).

Ludl., Can. Ent. XXXVI, 71: Banks, Phil. Jour. Sci. I, 985: Theob., Monog. Culic. IV, 179: Brun., Rec. Ind. Mus. I, 333; IV, 449: Ludl., Psyche XVIII, 126 (specific rank claimed).

Scutomyia notoscripta, Skuse, subspecies *samarensis*, Theob., Monog. Culic. V, 200: Brun., Rec. Ind. Mus. I, 336; IV, 452.¹

Culex crassipes Wulp, Mid-Sum. Dipt., 9, pl. i, 4, head (*Culex*), (1892).

nec *S. crassipes*, Theob., *et auctt.*

A very common species throughout the East, almost universally known as *scutellaris*, Walk.², occurring in India throughout the year except perhaps December to February. Sylhet, Lushai Hills, Manipur, Colombo, Straits (commonly distributed), Celebes, Aru, Amboina, Ceram, Papua, Borneo, Sumatra, Java, Assam, Siam (abundant), China, Japan and the Philippines. It also occurs in Krakatua, Mauritius, Fiji, the Seychelles, Christmas Is., Pitcairn Is., Honolulu.

Crassipes, Wulp, was described from Soeroelangoen, Sumatra.

Type of *samarensis* in the Army Medical Museum, Washington.

microptera Giles, Jour. Bomb. Nat. Hist. Soc. XIII, 609 (*Culex micropterus*, provis. descr.): Giles in Theob. Monog. Culic. II, 281, ♂ ♀ (*Wyeomyia? micropterus*), (1901).

Stegomyia microptera, Giles, Handbk., etc., 2nd ed., 380, ♂ ♀, pl. xiv, 24, head, thorax, 25, head, 26, venation: Theob., Monog. Culic. III, 147, note: Blanch, Moust., 263: Brun., Rec. Ind. Mus. I, 332; IV, 447, notes.

Allahabad, Lucknow, Jhansi. *Type* "appears to be lost." (Theob., Monog. V, 607 App.).

albolineata (I), Giles, Jour. Bomb. Nat. Hist. Soc. XIII, 609 provis. descr. (*Culex*), (1901).

¹ For fuller date, v. Brun., Rec. Ind. Mus. I, 333; IV, 449.

² Entered by error, twice in my Cat. Supp. 449 and 452.

- Brun., Rec. Ind. Mus. I, 336.
Shahjahanpur, N.W. India, 20-x-1900. *Type*, a unique ♀.
- sexlineata** Theob., Monog. Culic. I, 308, ♀, fig. 94, var. chars. (1901):
Giles, Jour. Trop. Med. VII, 367: *id.*, Handbk., etc., 2nd ed., 377: Brun., Rec. Ind. Mus. I, 334; IV, 449.
Angeles, Pampanga. *Type* in British Museum.
Described originally from a unique perfect ♀ taken at Agua Santa in Trinidad. The identity of the species seems established. Theobald (Monog. V, 219) has doubted it.
- periskelata** Giles, Handbk., etc., 2nd ed., 371, ♂, pl. xiv, 22, head ♂ (1902).
Theob., Monog. Culic. III, 145, ♂: Blanch., Moust., 264: Brun., Rec. Ind. Mus. I, 322; IV, 448, notes.
Shahjahanpur (N.W. India). *Type* probably lost, not in the British Museum. Apparently an uncertain species.
- punctolateralis** Theob., Entom. XXXVI, 156, ♂ ♀, (1903).
Giles, Jour. Trop. Med. VII, 367.
Pampanga [*Whitmore*]. Also occurs in Queensland, i [*Dr. Bancroft*]. *Type* in British Museum.
- amesii** Ludl., Jour. N. Yk. Ent. Soc. XI, 139 (*S. nivea, amesii*), (1903).
Theob., Monog. Culic. IV, 191 (copies Ludl.'s descr.): Brun., Rec. Ind. Mus. I, 329.
Skusea amesii, Edw., Bull. Ent. Res. VII, 213.
Stegomyia fusca, Leices., Culic. Malaya, 92, ♂ ♀ (1908).
Id., *id.*, Brun., Rec. Ind. Mus. IV, 447.
Oras, Samar, Tacloban, Leyte, Twin Peaks, Banquet, Luzon (all Philippine Is.). Port Swettenham.
Type of *amesii* in Army Medical Museum, Washington; of *fusca* in Leicester collection. Edwards would remove this species to his subgenus *Skusea* of *Aedes*.
- albolineata** (II). Theob., Entom. XXXVII, 77, ♀. (1904).
Theob., Monog. Culic. IV, 197, ♀: Leices., Culic. Malaya, 105, ♂ ♀: Brun., Rec. Ind. Mus. I, 336; IV, 452.
Kuala Lumpur, vi [*Leicester*]. *Type* in British Museum.
- desmotes** Giles, Jour. Trop. Med. VII, 367, ♀ (1904).
Brun., Rec. Ind. Mus. IV, 445: Edwards, Bull. Ent. Res. IV, 225.
Stegomyia gracilis, Leices., Culic. Malaya, 81, ♂ ♀ (1908): Brun., Rec. Ind. Mus. IV, 447.
Stegomyia albipes, Theob., Rec. Ind. Mus. IV, 11, ♀ (1910): *id.*, Monog. Culic. V, 169 ♀.
Typical form from the Philippines, *gracilis* from the Malay Peninsula, *albipes* from Maddathorai, Travancore, 17-xi-08 [*Annandale*]. *Type* (*desmotes*) in British Museum, damaged; of *gracilis* in Leicester collection, of *albipes*, a unique specimen, in Indian Museum.

- leucomeres** Giles, Jour. Trop. Med. VII, 367, ♀ (1904).
Brun., Rec. Ind. Mus. IV, 447.
Philippines. *Type* in British Museum.
- striocrura** Giles, Jour. Trop. Med. VII, 367 (1904).
Brun., Rec. Ind. Mus. 449.
Angeles, Pampanga. *Type*, location uncertain, not in British Museum. A doubtful species.
- trilineata** Leices. in Theob., Entom. XXXVII, 163, ♂ ♀ (*Hulecoetomyia*), (1904).
Stegomyia trilineata, Edwards, Bull. Ent. Res. IV, 226
Hulecoetomyia trilineata, Theob., Monog. Culic. IV, 220, ♂ ♀ (1907). Leices., Culic. Malaya, 107, ♂ ♀: Brun., Rec. Ind. Mus. IV, 455.
Howardina chrysolineata, Theob., Monog. Culic. IV, 218, ♀; pl. i, wing scales, ♀ (1907).
Kurseong, 3-vii-08 [*Annandale*]; Kuala Lumpur, iv, v, vii, x, xii; Peradeniya [*Green*]; Bukit Kutu, v; Ulu Gombok [both Malaysia, and *t. Leicester*].
Types of *trilineata* and *chrysolineata* in British Museum.
- thomsoni** Theob., Monog. Culic. IV, 174 (1907).¹
Id., Gen. Ins., Fasc. 26, 18 (1905): Edw., Bull. Ent. Res. VII, 210, fig. 5d, ♂ genit.
N.W. India; bred at Pusa [*Howlett*]. *Type* in British Museum.
- w-alba** Theob., Ann. Mus. Hung. III, 74, ♀, fig. 4, head, thorax, femur (1905).
Id., Monog. Culic. IV, 180, ♀, fig. 51, head, thorax, femur: Brun., Rec. Ind. Mus. I, 335: Edwards, Bull. Ent. Res. IV, 225.
Matheran, Bombay Presidency, 800 metres [*Biro*].
S. imitator, Leices., Culic. Malaya, 89, ♀ (1908): Brun., Rec. Ind. Mus. IV, 447.
S. minutissima, Theob., Rec. Ind. Mus. IV, 9, ♀; V, 168, ♀; fig. 61, wing (1910): Brun., Rec. Ind. Mus. IV, 448.
Matheran, Bombay Presidency, 800 metres [*Biro*] (*w-alba*); Kuala Lumpur [*t. Leices., imitator*]; Sukna, 1-vii-08 [*t. Theob., minutissima*].
Type (*w-alba*) in Hungarian Museum, a unique specimen; of *imitator* in Leicester collection; of *minutissima* in Indian Museum.
- annulirostris** Theob., Jour. Bomb. Nat. Hist. Soc. XVI, 239, ♀ (1905).
Id., Monog. Culic. IV, 173.
Peradeniya, i, 1902.
Type in British Museum, a unique specimen.

¹ The Genera Insectorum "description" cannot be considered as such, being confined to five lines.

- mediopunctata** Theob., Jour. Bomb. Nat. Hist. Soc. XVI, 240, ♀ (1905).
Theob., Monog. Culic. IV, 187; ♀.
Peradeniya, xi [Green]. *Type*, a unique specimen, in British Museum.
- gardneri** Ludl., Can. Ent. XXXVII, 99, ♂ ♀ (1905).
Quasistegomyia gardneri, Brun., Rec. Ind. Mus. IV, 443.
Pseudostegomyia gardneri, Ludl., Mosq. Phil. Is., 10 (*lapsus* for *Quasistegomyia*).
Bulaco, viii [Gardner]; Pampanga [Whitmore].
Type in Army Medical Museum, Washington.
- aurostriata** Banks, Phil. Jour. Sci. I, 995, ♀ (1906).
Theob., Monog. Culic. V, 181; Brun., Rec. Ind. Mus. IV, 445.
Mt. Siya Siya, Canlaon Volcano, Negros Is., 760 metres, 24-vi-06.
Type in Entom. Coll., No. 6082, Bureau of Science, Manila.
- argenteomaculata** Theob., Monog. Culic. IV, 184, ♀ (1907).
Brun., Rec. Ind. Mus. IV, 444.
Narcondam Island (Bay of Bengal), 80 miles from Andamans and Nicobars [Rogers].
Type in British Museum.
- tripunctata** Theob., Rec. Ind. Mus. II, 288, ♀ (1901).
Id., *loc. cit.*, IV, pl. i, wing; pl. iii, wing scales; V, 182, fig. 68, wing: Brun., Rec. Ind. Mus. IV, 450.
Lushai Hills, Assam, 6-vi-04 [Macleod].
Type in Indian Museum.
- albolateralis** Theob., Rec. Ind. Mus. II, 289, ♀ (1908).
Id., *loc. cit.*, IV, pl. i, wing; pl. iii, wing scales: *id.*, Monog. Culic. V, 179, fig. 67, wing: Brun., Rec. Ind. Mus. IV, 444.
Sylhet, ix [Hall]; Lungleh, Lushai Hills, vii.
Type in Indian Museum.
- assamensis** Theob., Rec. Ind. Mus. II, 290, ♀ (1908).
Id., Monog. Culic. V, 174, ♀: Brun., Rec. Ind. Mus. IV, 445.
Sylhet, 13-iv-05 [Hall]; Pallode, Travancore, 15-xi-08 [Annandale]. *Type* in Indian Museum.
- perplexa** Leices., Culic. Malaya, 83, ♂ ♀ (*Culex*), (1908).
Brun., Rec. Ind. Mus. IV, 448.
Kuala Lumpur, v, x, xi. Leicester says possibly a hybrid.
Type in Leicester collection.
- dissimilis** Leices., Culic. Malaya, 91, ♂ ♀ (1908).
Brun., Rec. Ind. Mus. IV, 445.
Ampang (Straits). *Type* in Leicester collection.
- annandalei** Theob., Rec. Ind. Mus. IV, 10 (1910).
Brun., Rec. Ind. Mus. IV, 444.
Kingia annandalei, Theob., Monog. Culic. V, 139: Brun., Rec. Ind. Mus. IV, 443.
Sukna, vii-08 [Annandale]. *Type*, unique, in Ind. Mus.

nigritia Ludl., Can. Ent. XLII, 194 (1910).

Philippines. *Type* in Army Medical Museum, Washington.

gregoryi Ludl., Psyche XVIII, 128, ♀ (1911).

Parang, Mindanao, i.

Type, presumably in Army Medical Museum, Washington.

quasinigritia Ludl., Psyche XVIII, 129 (1911).

Turucan, Mindanao, Philippine Is, xi. *Type*, unique, presumably in Army Medical Museum, Washington.

Howardina, Theob.

Monog. Culic. III, 287, pl. xv, wing scales (1903).

Theob.; *loc. cit.*, IV, 214; V, 200, tab. spp.:

Gen. Ins., Fasc. 26, 21: Blanch., Moust., 415:

Brun., Rec. Ind. Mus. I, 338; IV, 456; X, 62.

GENOTYPE, *Culex walkeri*, Theob., by Dyar's designation (Proc. Ent. So. Wash. VII, 49).

greenii Theob., Monog. Culic. III, 289, ♀; fig. 160 (p. 289), head, fig. 161, wing (1903).

Blanch., Moust., 416

Peradeniya, ii [*Green*]. *Type*, unique, in British Museum.

Of the three recorded oriental species, Edwards sinks *H. chrysolineata* as synonymous with *Stegomyia 3-lineata*, and *H. himalayana* as synonymous with *Ochlerotatus pulchriverter*, leaving only *H. greenii* unaccounted for. This species is, therefore, retained here under *Howardina* until disposed of.

Grabhamia, Theob.

Monog., Culic. III, 243 (1903).

Theob., *loc. cit.*, IV, 284, list. spp.: *id.*, Gen.

Ins., Fasc. 26, 23: Blanch., Moust., 396: Brun.,

Rec. Ind. Mus. I, 341; IV, 461; X, 60.

GENOTYPE, *Culex jamaciensis* Theob., by original designation.

Ceratocystia, Dyar and Knab, Jour. N. Yk. Ent.

Soc. XIV, 183 (1906).

GENOTYPE, *Culex discolor* Coq., by original designation.

Feltidia, Dyar, Proc. Ent. Soc. Wash. VII, No. 1, 47 (1905).

GENOTYPE, *Culex jamaciensis* Theob., by original designation.

sollicitans Walk., Ins. Saund. Dipt., pt. 5, 427 (*Culex*), (1856).

Culex sollicitans, Theob., Monog. Culic. I, 368, ♀; pl. xvi, 64, ♀, full ins. col.

Grabhamia sollicitans, Theob., Monog. Culic. III, 248; fig. 130,

wing ♀; IV, 291, life-history notes, fig. 97, 98, larval chars.:

Blanch., Moust., 295, ♀; fig. 212: Brun., Rec. Ind. Mus. I,

341; IV, 461, notes.

Formosa.

- spencerii* Theob., Monog. Culic. II, 99, ♀; pl. xxvi, 104, full ins. col. (*Culex*), (1901).
Culex spenceri, Theob., *loc. cit.*, II, fig. 198, var. chars.: Giles, Handbk., etc., 2nd ed., 431, ♀.
Grabhamia spenceri, Theob., Monog. Culic. III, 250: Brun., Rec. Ind. Mus. I, 341.
 Recorded by Theobald from the Philippines, but Banks (Phil. Jour. Sci. I, 986) doubts the identification. *Type* in British Museum.
- bitaeniorhynchus** Giles, Jour. Bomb. Nat. Hist. Soc. XIII, 607 (provis. descr.) (*Culex*), (1901).
 Edwards, Bull. Ent. Res. IV, 231.
Culex bitaeniorhynchus, Giles, Entom. XXXIV, 196, ♂ (1901).
Culex infula, Theob., Monog. Culic. I, 370, ♀ (1901).
 Giles, Handbk., 2nd ed., 407, ♀ (*infulus*): Blanch., Moust., 298, ♀: Leices., Culic. Malaya, 146 (Theob.'s descr. copied): Brun., Rec. Ind. Mus. I, 348.
Taeniorhynchus ager, Giles, in Theob., Monog. Culic. II, 199, ♂; fig. 248, abdomen, palpus, proboscis, wing scales.
 Giles, Handbk., 2nd ed., 365, ♂ ♀: Blanch., Moust., 385, ♂ ♀: Leices., Culic. Malaya, 168, ♂ ♀: Brun., Rec. Ind. Mus. I, 357; IV, 478.
Taeniorhynchus tenax, Theob., Monog. Culic. II, 198, ♀; pl. xvii, 65, full ins. col. (1901).
 Theob., *loc. cit.*, III, 259, fig. 236, wing: Giles, Handbk., 2nd ed., 365, ♀: Blanch., Moust., 386, ♀: Leices., Culic. Malaya, 167, ♀: Brun., Rec. Ind. Mus. I, 358; IV, 480.
Taeniorhynchus tenax, var. *ocellatus*, Theob., Monog. Culic. IV, 488, ♀ (1907).
Grabhamia ambigua, *id.*, *loc. cit.*, III, 248, ♂ (1903).
Grabhamia taeniarostris, *id.*, *loc. cit.*, IV, 299, ♀ (1907).
Oculecmyia sarawaki, *id.*, *loc. cit.*, IV, 515, ♀; fig. 236, 237; pl. vi, wing scales, ♀ (1907).
 ? *Taeniorhynchus domesticus*, Leices., Culic. Malaya 169, ♂ ♀ (1908): Brun., Rec. Ind. Mus. IV, 479.
 ? *Culicelsa abdominalis*, Taylor, Report Austral. Inst. Trop. Med. for 1911, p. 53 (1913).
 Sylhet i to vi [Hall]; Calcutta, 2-ii-08 [Annandale]; Balighai, near Puri, 26-x-08, and 23-x-08 at light [both Annandale]; Shahjahanpur (Unit. Prov. India), [Giles]; Madras [Giles and Cornwall]; Pallode, 14-xi-08; Quilon vii [James]; Kerumadi, S. end of Vembanaad Lake, Travancore, 6-xi-08 [Annandale]; Peradeniya, i [Green]; Ceylon [Giles]; Mandalay, 12-ii-08 [Annandale]; Manipur, viii [Gourlay]; Perak [Wray, Wright]; Taiping; Sarawak [Barker]; Philippines [t. Ludlow].
Types of *tenax*, *infula*, *ambigua*, *taeniarostris* and *sarawaki* in British Museum; of *domesticus* in Leicester collection.
 Edwards includes *domesticus*, Leices., temporarily as a variety, but notes that should it prove to be of specific rank it must be renamed, owing to the existence of *C. domesticus*, Germ.

- deniedmanni* Ludl., Can. Ent. XXXVI, 234, ♀ (*Grabhamia*), (1904).
Philippines.
- epidesmus* Theob., Rec. Ind. Mus. IV, 22, ♀ (*Taeniorhynchus*), (1910).
Id., Monog. Culic. V, 429, ♀ (*Taeniorhynchus*).
Culex epidesmus, Edw., Bull. Ent., Res. IV, 231.
Taeniorhynchus luteoabdominalis, Theob., Rec. Ind. Mus. IV, 423, ♀ (1910): *id.*, Monog. Culic. V, 423; Brun., Rec. Ind. Mus. IV, 479.
Grabhamia ochracea, Theob., Jour. Econ. Biol. I, 25 ♀ (1905).
Theob., Monog. Culic. IV, 300, ♀.
Bhogaon, 2-x-08; Katihar, 5-x-08 [both Bengal and *Paiva*]; India [*Christophers*].
Types of epidesmus and luteoabdominalis in Indian Museum; of *ochracea* in British Museum.

Ochlerotatus, Arrib.

- Rev. Mus. La Plata II, 143 (1891).
Brun., Rec. Ind. Mus. X, 67, note *re* genotype.
Edwards, Bull. Ent. Res. II, 246; VII, 203.
GENOTYPE, *Culex albifasciatus* Macq.¹
- Finlaya*, Theob., Monog. Culic. III, 281 (1903).
Theob., *loc. cit.*, IV, 520; Gen. Ins., Fasc. 26, 32: Blanch., Moust., 415, 630.
Brun., Rec. Ind. Mus. I, 360; IV, 485; X, 60.
GENOTYPE, *F. poicilia*, Theob., by my own designation (1914).
- Aedimorphus*, Theob., Monog. Culic. III, 290 (1903).
GENOTYPE, *Uranotaenia domestica*, Theob., by original designation.
- Danielsia*, Theob., Entom. XXXVII, 78 (1904).
Theob., Gen. Ins., Fasc. 26, 21: *id.*, Monog. Culic. IV, 240; Leices., Culic. Malay., 117.
GENOTYPE, *D. albotaeniata*, *id.*, *loc. cit.*, sp. nov.
- Culicelsa*, Felt, N. Yk. State Mus. Bull. 79, Ent. 22. App., p. 391 (1904).
GENOTYPE, *Culex taeniorhynchus* W., by original designation.
- Culicada*, Felt., *loc. cit.*, p. 391b (1904).
Theob., Jour. Econ. Biol. I, 26 (1905): *id.*, Monog. Culic. IV, 318 and p. 10; V, 294, tab. spp.: Brun., Rec. Ind. Mus. IV, 461; X, 58. [tion.
GENOTYPE, *Culex canadensis* Theob., by original designation.
- Ecculex*, Felt, *loc. cit.*, p. 391c (1904). [tion.
GENOTYPE, *Culex sylvestris* Theob., by original designation.

¹ See Brun., Rec. Ind. Mus. X, 67. Edwards agrees to *albifasciatus*, Macq., as the type species, though Coquillett has suggested *confirmatus*, Arrib.

- Protoculex*, Felt, *loc. cit.*, p. 391d (1904).
 GENOTYPE, *Culex serratus* Theob., by original designation.
- Pseudoculex*, Dyar, Proc. Ent. Soc. Wash. VII, 45 (1905).
 GENOTYPE, *Culex aurifer* Coq.
- Chrysoconops*, Goeldi, Os. Mosq. do Para, 114 (1905).
 Theob., Monog. Culic. IV, 491; V, 433, tab. spp.: Brun.
 IV, 480; X, 157.
 GENOTYPE, unknown to present author.
- Reedomyia*, Ludl., Can. Ent. XXXVII, 94 (1905).
 Theob., Monog. Culic. IV, 257, ♂ ♀, list. and tab. spp.; V,
 252: Brun., Rec. Ind. Mus. I, 362; IV, 486, note; X, 70.
 GENOTYPE, *R. pampangensis* Ludl., by original designation.
- Lepidotomyia* I, Theob., Ann. Mus. Hung. III, 80 (1905).
 nec *Lepidotomyia* II, Theob., Gen. Ins., Fasc. 26, 22 (1905).
 GENOTYPE, *L. alboscuteclata*, *id.*, *loc. cit.*, sp. nov.
- Pecomyia*, Theob., Jour. Econ. Biol. I, No. 1, 24 (1905).
 Theob., Monog. Culic. IV, 265.
 GENOTYPE, *P. maculata* Theob., by original designation.
- Pseudograbhamia* Theob., Jour. Bomb. Nat. Hist. Soc. XVI,
 243 (1905).
 Theob., Monog. Culic. IV, 314: Brun., Rec. Ind. Mus.
 IV, 460; X, 69.
 GENOTYPE, *P. maculata* Theob., by original designation.
- Phagomyia*, Theob., Gen. Ins., Fasc. 26, 21 (1905).
 Theob., Monog. Culic. IV, 223; V, 224, tab. spp.: Brun.,
 Rec. Ind. Mus. I, 338; IV, 456; X, 68.
 GENOTYPE, *Stegomyia gubernatoris* Giles.¹
- Polyleptomyia*, Theob., Gen. Ins., Fasc. 26, 21 (1905).
 Theob. Monog. Culic. IV, 223.
 GENOTYPE, *P. albocephala* Theob., by original designation.
- Lepidotomyia* II,² Theob., Gen. Ins., Fasc. 26, 22 (1905).
 Theob., Monog. Culic. IV, 249; V, 249: Brun., Rec. Ind.
 Mus. IV, 458, note on names; X, refs. to names.
 nec *Lepidotomyia* I, Theob., Ann. Mus. Hung. III, 80; which
 = *Reedomyia*, Ludl.
 GENOTYPE, *L. magna* Theob., by original designation.
- Lepidoplatus*, Coq., Science XXIII, 314 (1906).
 GENOTYPE, *Culex squamiger* Coq., by original designation.
- Pseudoskusea*, Theob., Monog. Culic. IV, 192 (1907).
 GENOTYPE, *Skusea multiplex* Theob., by my own designa-
 tion, 1914.
- Pseudohowardina*, Theob., Monog. Culic. IV, 223 (1907).
 GENOTYPE, *Culex trivittatus* Coq., by original designation.
- Neomacleaya*, Theob., Monog. Culic. IV, 238 (1907).

¹ Theobald does not definitely select this species, but he places it first out of three species on three separate occasions, the third species being admittedly doubtfully congeneric.

² Theobald's first genus of the name of *Lepidotomyia* was admittedly synonymous with *Reedomyia*, but his second *Lepidotomyia* was (in his view) a valid genus.

- Brun., Rec. Ind. Mus. IV, 458.
 GENOTYPE, *N. indica*, *id.*, *loc. cit.*, sp. nov.
Protomacleaya, Theob., Monog. Culic. IV, 253 (1907).
 GENOTYPE, *Culex triseriatus* Say.
Duttonia, Newstead, Ann. Trop. Med. Parasit. I, 17 (1907).
 GENOTYPE, *D. tarsalis*, Newstead by original designation.
Mimetecculex, Theob., 3rd Rep. Wellcome Res. Labor. Gordon College, 258 (1908).
 Theob., Monog. Culic. V, 408.
 GENOTYPE, *M. kingii* Theob., by original designation.
Geitonomyia, Leices., Culic. Malaya, 134 (1908).
 GENOTYPE, *Culex caccus* Theob., by original designation.
Myxosquamus, Theob., Monog. Culic. V, 225 (1910).
 GENOTYPE, *M. confusus* Theob., by original designation.
Neopecomyia, Theob., Monog. Culic. V, 261 (1910). [tion.
 GENOTYPE, *N. uniannulata* Theob., by original designa-
Stenoscutus, Theob., Monog. Culic. V, 263 (1910)
 GENOTYPE, *S. africanus* Theob., by original designation.
Bathosomyia, Theob., Monog. Culic. V, 267 (1910).
 GENOTYPE, *B. abnormalis* Theob., by original designation.
Pseudocarrollia, Theob., Rec. Ind. Mus. IV, 12 (1910).
Id., Monog. Culic. V, 186: Brun., Rec. Ind. Mus. IV, 454;
 X, 69. [tion.
 GENOTYPE, *P. lophoventralis* Theob., by original designa-
Leslieomyia, Christophers, Paludism, No. 2, 68 (1911).
 GENOTYPE, *L. taeniorhynchoides* Christophers, by original
 designation.
Acartomyia, Theob., Monog. Culic. III, 251 (1903).
 GENOTYPE, *A. zammittii* Theob., by original designation.
 ? *Gilesia*, Theob., Monog. Culic. III, 233 (1903).
 ? *Gualteria*, Lutz in Bourroul's Mosq. Bras. 49 (? 54) (1904).
 Theob., Gen. Ins., Fasc. 26, 21; *id.*, Monog. Culic. IV,
 240; Leices., Culic. Malaya, 117. [(1906).
 ? *Cacomyia*, Coq., U.S. Dep. Agric. Bull. Tech. Ser. II, 16
 ? *Stegoconops*, Lutz, Imprensa Medica (1906). (? *nom. nud.*)
 Peryassu, Os. Culic. do Bras., 34 (1908).
 ? *Molpemyia*, Theob., Monog. Culic. V, 479 (1910).
 ? *Andersonia*, Strickland, Entom., 250 (1911).
- vexans** Mg., Syst. Besch. VI, 241 (*Culex*), (1830).
Ochlerotatus vagans, Edw., Bull. Ent. Res. VII, 218.
Culex articulatus, Rond., Bull. Soc. Ent. It. IV, 70 (1832).
Culex sylvestris, Theob., Monog. Culic. I, 406 (1901).
Culex vagans, *id.*, *loc. cit.*, I, 411 (1901).
Culicada nipponii, *id.*, *loc. cit.*, IV, 337 (1907).
Culicada minuta, *id.*, *loc. cit.*, IV, 338 (1907).
Culex stenoetrus, *id.*, *loc. cit.*, IV, 395 (1907).
Ochlerotatus stenoetrus, Edw., Bull. Ent. Res. IV, 228.
Culicada eruthrosops, Theob., Monog. Culic. V, 299, ♀, fig. 140,
 wing (1910).

Culex pseudostenocrus, *id.*, *loc. cit.*, V, 343, fig. 154, wing (1910).

Culex hirsutum, Ludl., Psy. XVIII, 126 (1911).

? *Culex nocturnus*, Theob., Monog. Culic. III, 159 (1903).

A very widely distributed species. India [*Christophers*]; Maskeliya, Ceylon, iv [*Green*]; Hakgala, Ceylon, v, viii-07 [*Green*]; Trincomalee, xi-06. Also found in China, Japan, the Philippines, Borneo, England and other parts of Europe, in Canada and possibly in Fiji.

Edwards finds the male genitalia of specimens from many of these localities agree exactly. He finds three definite varieties, *vexans* (*articulatus*, *sylvestris*); *stenocrus* (*minuta*, etc.), and *nipponii* (*vagans*, Theob., nec. Wied.). Types of *minuta*, *stenocrus*, *cruthrosops* and *pseudostenocrus* in British Museum.

imprimens Walk., Proc. Linn. Soc. V, 144, ♀ (*Culex*), (1861).

Edwards, Bull. Ent. Res. IV, 228.

Culex imprimens, Giles, Handb., 262; 2nd ed., 411, ♀ (*imprimiens*, *lapsus*): Blanch., Moust., 306, ♀: Theob., Gen. Ins., Fasc. 26 (*imprimiens*): Brun., Rec. Ind. Mus. I, 348.

Culicada suknaensis, Theob., Rec. Ind. Mus. VI, 21, ♀ (1910).

Id., Monog. Culic. V, 297, ♀, fig. 139, wing: Brun., Rec. Ind. Mus. IV, 462.

? *Culex auratus*, Leices., Culic. Malaya, 153, ♀: Brun., Rec. Ind. Mus. IV, 467.

Amboina, Sukna, 1-2-vii-08 [*Annandale*]; Malaysia [*t. Leicester*].

Type of *imprimens*, a remnant, in British Museum: of *suknaensis* in Indian Museum.

conopas Frnfl., Verh. zool.-bot. Ges. Wien. XVII, 451 (*Culex*) (1867).

Culex conopas, Giles Handbk., 286.

Taeniorhynchus conopas, Theob., Monog. Culic. II, 202, ♂, fig. 249, wing, wing scales, scutellum; pl. xxiii, 90, ♀, full ins. col.; pl. E, wing scales: Giles, Handbk., 2nd ed., 360, ♂: Blanch., Moust., 387, ♀ (*conopus*): Brun., Rec. Ind. Mus. I, 358: Leices., Culic. Malaya, 166, ♀.

Chrysoconops conopas, Brun., Rec. Ind. Mus. IV, 481.

Selangor, 28-x-99 [*Butler*]; Kuala Lumpur [*Durham*]; Klang [*Leicester*]; Perak [*Wray*]; Dinding, Straits, vi, xii; Formosa, 8-i-1900 and vi [*Ford*]; Philippines [*t. Ludlow*].

Type, location unknown to present author.

kochi Don., Insektenborse XVIII, 38, ♀ (*Culex*), (1901).

Culex kochi, Theob., Monog. Culic. II, 217: Blanch., Moust., 272, ♀.

Finlaya kochi, Theob., Monog. Culic. III, 284: Brun., Rec. Ind. Mus. I, 361.

Papua. Type, location unknown to present author.

gubernatoris Giles, Jour. Bomb. Nat. Hist. Soc. XIII, 607 (*Culex*), (provis. descr.), (1901).

Culex gubernatoris, id., Entom. XXXIV, 194: Theob., Monog. Culic. I, 314, ♀, fig. 97, thorax ♀.

Ochlerotatus gubernatoris, Edw., Bull. Ent. Res. IV, 226.

Stegomyia gubernatoris, Giles, Handbk., 2nd ed., 380, ♀: Blanch., Moust., 261, ♀.

Phagomyia gubernatoris, Theob., Gen. Ins., Fasc. 26, 21: Brun., Rec. Ind. Mus. I, 338.

Lepidotomyia (II) *magna*, Theob., Gen. Ins., Fasc. 26, 22 (1905).¹ Theob., Monog. Culic. IV, 250, ♂ ♀.

Allahabad, vii; North India; Bombay, 19-viii-02 [James]; Purnea, Bengal, 6-viii-07 [Paiva]; Pampanga, Philippine Is. [Whitmore].

Types of gubernatoris (damaged but recognizable), and *magna* in British Museum; of *lophoventralis*, a unique specimen, in Indian Museum.

As regards the *L. magna* of Theobald it must be remembered that he had two genera named *Lepidotomyia*, the first being synonymous with the previously described *Reedomyia*, Ludl., the second represented solely by *magna*.

pulchriventer Giles, Jour. Bomb. Nat. Hist. Soc. XIII, 608 (*Culex pulchriventer*, provis. descr.), (1901).²

Culex pulchriventer, Theob., Monog. Culic. II, 48, ♂ ♀; pl. xxiii; 92, ♀, full ins. col., fig. 170, ♂ ♀, var. chars.; fig. 172, wing, ♂; fig. 171, wing, ♀; fig. 173, larva: Giles, Handbk., 2nd ed., 449, ♂ ♀; pl. xvii, var. chars.: Blanch., Moust., 338, ♂ ♀: Brun., Rec. Ind. Mus. I, 351.

Ochlerotatus pulchriventer, Edw., Bull. Ent. Res. IV, 228.

Howardina himalayana, Giles, Jour. Trop. Med. VII, 384, ♀ (1904): Brun., Rec. Ind. Mus. IV, 456.

Naini Tal, viii.

Types of pulchriventer and himalayana in British Museum.

caecus Theob., Monog. Culic. I, 413, ♀; fig. 147; head, 148, scutellum and scales; pl. xx, 77, full ins. col. (*Culex*), (1901).

Culex caecus, Theob., loc. cit., IV, 268; fig. 86, wing ♀; pl. i, wing scales, ♀: Giles, Handbk., 2nd ed., 415, ♀: Blanch., Moust., 305, ♀.

Geitonomyia caecus, Leices., Culic. Malaya, 135, ♂ ♀.

Pecomomyia cacca, Theob., Monog. Culic. IV, 268; fig. 86, wing ♀; pl. i, wing scales, ♀: Brun., Rec. Ind. Mus. IV, 459.

Selangor, 28-x-99 [Butler]; Klang; Ipoh, Perak; Straits [Durham]; Philippines [t. Ludlow]. *Type* in British Museum.

mediolineatus Theob., Monog. Culic. II, 113 (*Culex*), (1901).

Culex mediolineatus, Giles, Handbk., 2nd ed., 341, ♀: Blanch., Moust., 369, ♀.

Ochlerotatus mediolineatus, Edw., Bull. Ent. Res. IV, 228.

Culex trilineatus, Theob., Monog. Culic. II, 105, ♀ (1901); loc.

¹ ² A "description" of five lines only in each case.

cit., V, 359: Giles, Handbk., 2nd ed., 464, ♀: Blanch., Moust., 330, ♀: Brun., Rec. Ind. Mus. IV, 476.

Types of both *mediolineatus* and *trilineatus* in British Museum.

aurites Theob., Monog. Culic. II, 209, ♀; fig. 253, 254, var. chars.; pl. xxii, 88, ♀, full. ins. col. (*Taeniorhynchus*), (1901).

Taeniorhynchus aurites, Giles, Handbk., 2nd ed., 362, ♀: Brun., Rec. Ind. Mus. I, 357.

Chrysoconops aurites, Brun., Rec. Ind. Mus. IV, 480.

Dindings, Perak [*Wright*]; Kuala Lumpur, 10-xi-02; 25-v-02 [*Durham*]; Philippines [*t. Ludlow*].

Type in British Museum, from Dr. Annett's collection.

pipersalatus Giles in Theob., Monog. Culic. II, 316, ♂ ♀ (*Stegomyia*), (1901).

Stegomyia pipersalata, Giles, Handbk., 2nd ed., 372, ♀ only¹; pl. xvi, 1 a, b, venation (given as ♂, Edwards says it is ♀) (1902); V, 607, App. (note as to generic position): Blanch., Moust., 264 (♀ only): Brun., Rec. Ind. Mus. I, 332; IV, 448.

Pseudograhamia maculata, Theob., Jour. Bomb. Nat. Hist. Soc. XVI, 243, ♂ ♀ (1905): Theob., Monog. Culic. IV, 314, ♂ ♀; fig. 109, wing ♀.

Ochlerotatus pipersalatus, Edw., Bull. Ent. Res. IV, 227.

Jhansi; Gonda, N.W. India; Galgamuwa, Ceylon, viii [*Green*]; Madras town, 30-x-08 [*Hodgart*].

Types of *pipersalatus* and *maculata* in British Museum. Theobald (Monog. I, 316) attributes this species to Giles, and this appears to be the original description, but he adds as a reference "Handbk., 2nd ed., 1901," giving no page, which makes it appear as though the latter work had not appeared at the time. Giles, 2nd ed., is dated 1902, a year later than Theobald's Monograph Vol. I, so the latter reference must stand.

ochraceus Theob., Monog. Culic. III, 263, ♀; fig. 140, scutellum (*Taeniorhynchus*), (1903).

Ochlerotatus ochraceus, Edw., Bull. Ent. Res. II, 250.

Taeniorhynchus ochraceus, Leices., Culic. Malaya, 164.

Taeniorhynchus ochraceus, Brun., Rec. Ind. Mus. I, 358.

Chrysoconops ochraceus, *id.*, *loc.*, *cit.* IV, 481.

Kuala Lumpur [*Durham*]. Type in British Museum.

poicilia Theob., Monog. Culic. III, 283, ♀; fig. 156, wing scales (slightly incorrect) (*Finlaya*), (1903).

Finlaya poicilia, Giles, Jour. Trop. Med. VII, 366 (*poialia, lapsus*): Theob., Monog. Culic. IV, 520; fig. 238 (corrected wing ♂): Blanch., Moust., 415: Brun., Rec. Ind. Mus. I, 361; IV, 486.

Penang, 24-x-07 [*Freear*]; Friedrich Wilhelmshafen, Seleo Berlinhafen and Mt. Hausemann, Astrolabe Bay [all Papua and

¹ Edwards says (Bull. Ent. Res. IV, 227) that the ♂ belongs to a different species at present undescribed. He should, however, have retained the specific name for the ♂ and renamed the ♀.

Biro]; Pasir Ganting, W. Coast Sumatra, vi-14; Lampongs, Sumatra, vi-14 [both *t. Stanton*]; Pampanga [*Whitmore*]; Negros I., Philippines [*Banks*]. Also occurs in N. Queensland. *Type* in British Museum.

multiplex Theob., Monog. Culic. III, 293, ♀; fig. 165, claws (*Skusea*), (1903).

Skusea multiplex, Brun., Rec. Ind. Mus. I, 336.

Pseudoskusea multiplex, Theob., Monog. Culic. IV, 192, ♂: Brun., Rec. Ind. Mus. IV, 450.

Friedrich Wilhelmshafen, Stephansort; Muina; Ins. Graget [all Papua and *Biro*]. *Type* in Hungarian Museum.

niveus Ludl., Jour. N. Yk. Ent. Soc. XI, 139 (*Stegomyia nivea*), (1903).

Stegomyia nivea, Leices., Culic. Malaya, 87, ♂ ♀.

Scutomyia nivea, Theob., Monog. Culic. V, 203: Brun., Rec. Ind. Mus. I, 336; IV, 452.

Ochlerotatus niveus, Edw., Bull. Ent. Res. IV, 227.

Stegomyia pseudonivea, Theob., Ann. Mus. Hung. III, 75 (1905).

Theob., Monog. Culic. IV, 188, ♀; V, 176, ♂, fig. 64, head: Brun., Rec. Ind. Mus. I, 332; IV, 448.

Fed. Malay States; Oras, Samor, Philippine Is., v, vi, 1902; Andaman Is. [*Lowis*].

Type of *niveus* in Army Medical Museum, Washington, of *pseudonivea* in Hungarian National Museum.

See note under *S. amesii*, Brun., Rec. Ind. Mus. I, 329.

albotaeniatus Leices. in Theob., Entom. XXXVII, 111, ♂ ♀ (*Danielsia*), (1904).

Danielsia albotaeniata, Theob., Monog. Culic. IV, 241, ♂ ♀; fig.

72 (cephalic and scutellar adornment): Brun., Rec. Ind. Mus. I, 338; IV, 458: Leices., Culic. Malaya, 117, ♂ ♀.

Malay Peninsula. *Type* in Leicester coll.

melanopterus Giles, Jour. Trop. Med. VII, 367 (*Finlaya*), (1904).

Finlaya melanoptera, Banks, Phil. Journ. Sci. I, 990 (1906).

Ochlerotatus melanopterus, Edw., Bull. Ent. Res. VII, 214.

Popea palawanensis, Ludl., Psy. XXI, 31 (1914).

Philippines. *Type* of *melanopterus* in British Museum.

flavipennis Giles, Jour. Trop. Med. VII, 367 (*Finlaya*), (1904).

Pampanga [*Whitmore*]. *Type* in British Museum.

niveoscutellatus Theob., Jour. Econ. Biol. I, No. 1, 22; pl. iii, 5, ♀ (*Reedomyia*), (1905).

Theob., Monog. Culic. IV, 259, ♂ ♀; fig. 79, ♂, fore claws: Brun., Rec. Ind. Mus. IV, 487.

India; Philippines [*t. Ludlow*]. *Type* in British Museum.

alboscutecllatus Theob., Ann. Mus. Hung. III, 80, ♀ (*Lepidotomyia*), (1905).

Reedomyia alboscutecllata, Theob., Monog. Culic. IV, 261, ♀; fig. 80, wing, ♀: Brun., Rec. Ind. Mus. IV, 487.

- Lepidotomyia alboscuteolata*, Leices., Culic. Malaya, 132, ♂ ♀:
Brun., Rec. Ind. Mus. I, 339.
Simbang (Huon Gulf), and Friedrich Wilhelmshafen [both
Papua and *Biro*]; Kuala Lumpur. Type in Hungarian Mus.
- pampangensis* Ludl., Can. Ent. XXXVII, 94, ♀ (*Reedomyia*),
(1905).
Reedomyia pampangensis, Theob., Monog. Culic. IV, 258, ♀.
Brun., Rec. Ind. Mus. I, 362.
Anzeles, Pampanga, ix [*Whitmore*].
Type in Army Medical Museum, Washington.
- aranetana* Banks, Phil. Jour. Sci. I, 1001, ♂ ♀ (*Finlaya*), (1906).
Finlaya aranetana, Brun., Rec. Ind. Mus. I, 361.
Bago, Negros I., on the Siya Siya Peak of the Canlaon Vol-
cano, 700 metres, vi.
Types in Entom. coll. No. 6066, ♂ ♀, Bureau of Science,
Manila.
- pallidostriatus* Theob., Monog. Culic. IV, 410, ♂ ♀, fig. 175,
wing ♂ (*Culex*), (1907).
Ochlerotatus pallidostriatus, Edw., Bull. Ent. Res. IV, 228.
Ochlerotatus ochraceus, Edw. (pt.), Bull. Ent. Res. II, 250.
Culex pallidostriatus, Edw., Bull. Ent. Res. IV, 228.
Culex parascelos, Theob., Rec. Ind. Mus. IV, 18, ♀ (1910): *id.*,
Monog. Culic. V, 379: Brun., Rec. Ind. Mus. IV, 473.
Mimteculex kingii, Theob., 3rd Rep. Wellc. Labor., 258 (1908).
India [*James, Christophers*]; Madras town, 30-x-08 [*Hodgart*];
Peradeniya, xii [*Green*].
Type of *pallidostriatus* in British Museum; of *parascelos* in
Indian Museum. Edwards now considers *ochraceus*, Theob., typi-
cal form, to be confined to the Ethiopian Region.
- taeniatus* Leices., Culic. Malaya, 133, ♂ ♀ (*Lepidotomyia*), (1908).
Lepidotomyia taeniata, Brun., Rec. Ind. Mus. IV, 459.
Malaysia. Type in Leicester collection.
- ostentatio* Leices., Culic. Malaya, 193, ♀ (*Aiorctomyia*), (1908).
Ochlerotatus ostentatio, Edw., Bull. Ent. Res. IV, 228.
Aiorctomyia ostentatio, Brun., Rec., Ind. Mus. IV, 490.
Pseudohowardina chrysoseuta, Theob., Monog. Culic. V, 228, ♀;
fig. 94, wing (1910).
Malaysia; Peradeniya, iv-07 [*Green*].
Type of *ostentatio* in Leicester collection; of *chrysoseuta* in
British Museum. Edwards notes that as the ♂ is unknown, the
location here of this species is provisional.
- lowisii* Theob., Monog. Culic. V, 257, ♂ ♀; fig. 121, head ♀, 122
wing, 123, head ♂, 124, wing, ♂ (*Reedomyia*), (1910).
Galle, Ceylon, 6 iv-07 [*Fletcher*]; Andaman Is. [*Lewis and Ray
White*]. Type in British Museum.
- lophoventralis* Theob., Rec. Ind. Mus. IV, 13 (*Pseudocarrollia*),
(1910).
Ochlerotatus lophoventralis, Edw., Bull. Ent. Res. VII, 214.

- Pseudocarrollia lophoventralis*, Theob., Monog. Culic. V, 186, ♀:
Brun., Rec. Ind. Mus. IV, 454.
Purnea, Bengal (*type*) [*Paiva*]; Pusa [*Howlett*].
Type in Indian Museum.
- alboannulis* Ludl., Psyche XVIII, 132, ♀ (*Duttonia*), (1911).
Mindanao, Philippine Is., i, xi, xii.
Type in Army Medical Museum, Washington.
- taeniorhynchoides* Chris., Paludism, No. 2, p. 68 (*Leslicomyia*),
(1911).
Ochlerotatus taeniorhynchoides, Edwards, Bull. Ent. Res. IV, 227.
Pecomomyia maculata, Theob. (*nec* Meigen), Jour. Econ. Biol. I, 24;
pl. iv, 7 (1905): Theob., Monog. Culic. IV, 266, ♂ ♀; figs. 82,
83, 84, wings ♀, India [*James*; *Christophers*].
Type, location of that of *taeniorhynchoides* unknown to present
author; that of *maculata* in British Museum. Mr. Edwards
says *maculata* may possibly be distinct, in which case it must
be renamed, as *Culex maculatus*, Mg. falls in this genus.
- annulifemur* Edw., Bull. Ent. Res. V, 77, ♂ (1914).
Jhansi, N.W. India, viii, 1900 [*Giles*].
Type in British Museum.
The specimen was the *type* ♂ of *Stegomyia pipersalata*, Giles,
but that author had confounded two species under one name.
Edwards fully describes it.
- jamesii* Edw., *loc. cit.*, 77 (1914).
Colombo [*James*]; Galle [*Fletcher*]. *Type* in Brit. Mus.
- oreophilus* Edw., Bull. Ent. Res. VI, 357, ♀ (1915—16).
Gharia, Murree Hills, W. Himalayas, 16-viii. Unique *type* in
British Museum, sent by Pusa Agricultural Research Institute.
- koreicus* Edw., Bull. Ent. Res. VII, 212, ♂ ♀ (1916—17).
Korea [*Dr. Mills*]. *Types* presented to British Museum by
Dr. Stanton.

Aedes, Mg.

- Syst. Besch. I, 13 (1818).
Sch., F. Austr. II, 630: Ficalbi, Bull. Soc. Ent.
It., 299 (1896): Theob., Monog. Culic. II, 224;
III, 285, restricted; V, 482, tab. spp.: *id.*, Gen.
Ins., Fasc. 26, 35: Giles, Handb., 343; 2nd ed.,
480; tab. spp., 346: Blanch., Moust., 399, ♂
♀: Leices., Culic. Malaya, 183: Brun., Rec. Ind.
Mus. IV, 489; X, 45, tab. *Aedes* group: Edw.,
Bull. Ent. Res. IV, 229; VII, 203: Alcock, Ent.
Med. Off., 102: Howard, Dyar and Kuab, Mosq.
N. Am., etc., IV, pt. 2, 612.
GENOTYPE, *A. cinereus* Mg., by original designa-
tion.
Heteronycha, Arrib., Rev. Mus. La Plata. I, 397
(1891).

- GENOTYPE, *Culex aestuans* Wied., as *dolosa*, sp. nov., by original designation.¹
- Skusea*, Theob., Monog. Culic. III, 291 (1903).
Id., loc. cit., IV, 542; V, 488, tab. spp.: Gen. Ins., Fasc. 26, 19; Blanch., Moust., 416; Brun., Rec. Ind. Mus. I, 335; IV, 450, notes; X, 71; Edw., Bull. Ent. Res. VII, 203.
 GENOTYPE, *S. pembraensis* Theob., by Blanchard's designation, 1905.
- Verrallina*, Theob., Monog. Culic. III, 295 (1903).
 Theob., loc. cit., V, 494, tab. spp.: Blanch., Moust., 417.
 Leices., Culic. Malaya, 196, ♂ ♀.
 GENOTYPE, *Aedes butleri* Theob., by Coquillett's designation² (1910).
- Macleaya*, Theob., Entom. XXXVI, 154 (1903).
 Theob., Monog. Culic. IV, 203.
 GENOTYPE, *M. tremula*, sp. nov., loc. cit.
- Neomacleaya*, Theob., Monog. Culic. IV, 238 (1907).
 GENOTYPE, *N. indica* Theob., by original designation.
- Aioretomyia*, Leices., Culic. Malaya, 185 (1908).
 Brun., Rec. Ind. Mus. IV, 490; X, 55.
 GENOTYPE, *A. varietas* Leices., by my own designation.
- Catageiomyia*, Theob., Monog. Culic. V, 115 (1907).
 This is a *nomen nudum*, of which no further information seems available.

Howard, Dyar and Knab give nearly fifty generic synonyms of *Aedes*. As Edwards recognises several of them as valid genera, viz. *Banksinella*, *Grabhamia*, *Howardina*, *Ochlerotatus*, *Stegomyia* and *Taeniorhynchus*, I have adopted all these latter with the respective synonymy of each as worked out by that author, the remainder of the American authors' synonyms being recognised here as such.

- pseudotaeniatus* Giles, J. Bomb. Nat. Hist. Soc. XIII, 606, prov. descr., 1901; Entom. XXXIV, 192 (*Stegomyia*), (1901).
- Hulecoetomyia pseudotaeniata*, Brun., Rec. Ind. Mus. I, 337; IV, 455; Theob., Monog. Culic. V, 222 (♂ descr.); fig. 89, wing, 90, head ♂, 91, ♂ genitalia.
- Stegomyia pseudotaeniata*, Giles, Handbk., etc., 2nd ed., 379, ♂ ♀; pl. xiv, 8, venation, 9, body, 10, larva; Theob., Monog. Culic. I, 312, ♀; fig. 96, thorax, head ♀, wing scales; 314, larva described: loc. cit., III, fig. 16 (p. 28), larva; Blanch., Moust., 261, ♀.
- Dehra Dun [Thomson]; Bakloh, Punjab; Naini Tal, 6000—8000 ft. [Giles]; Lushai Hills, 29-vii-14; Rizal, Manila, 1 [Banks]. Type in British Museum.

¹ *Culex aestuans*, Wied., has been regarded by some as synonymous with *C. fatigans*, Wied., in which case the generic synonymy here seems doubtful.

² Edwards in selecting *butleri* as genotype (Bull. Ent. Res. IV, 229) overlooks Coquillett's designation of the same species in 1910.

- uncus** Theob., Monog. Culic. II, 53 (*Culex*), (1901).
Aedes uncus, Edwards, Bull. Ent. Res. IV, 229; VII, 223, fig. 8d, ♂ genitalia.
Culex uncus, Giles, Handbk., 2nd ed., 452, ♀: Blanch., Moust., 350, ♀.
Neomacleaya indica, var. *simplex* only, Theob., Rec. Ind. Mus. II, 291. ♀ (1908).
 ? *Verrallina malayi*, Leic., Culic. Malaya, 198, ♀ (1908)
 Sylhet, vi [Hall]; Klang; Kuala Lumpur. Types of *uncus* and *simplex* in British Museum; of *malayi* in Leicester's collection.
- butleri** Theob., Monog. Culic. II, 230, ♀ (1901).
 Giles, Handbk., 2nd ed., 481, ♀: Brun., Rec. Ind. Mus. I, 367; IV, 489; Edwards, Bull. Ent. Res. IV, 229.
Verrallina butleri, Blanch., Moust., 417; Leices., Culic. Malaya, 196, ♂ ♀: Brun., Rec. Ind. Mus. IV, 491.
Skusea diurna, Theob., Entom. XXXVI, 259, ♀ (1903).
Id., Monog. Culic. IV, 547, ♀: Brun., Rec. Ind. Mus. I, 335; IV, 451.
Stegomyia hatiensis, Carter, Entom. XLIII, 275 (1910).
 Malaysia; Selangor; Jugra, ix [Durham]; Philippine Is. [t. Ludlow]. Types of *butleri*, *diurna* and *hatiensis* in Brit. Mus.
- funereus** Theob., Monog. Culic. III, 292, ♀, fig. 164, head, abdomen (*Skusea*), (1903).
Skusea funerea, var. *ornata*, Theob., Ann. Mus. Hung. III, 79, ♀, pl. i, wing (1905); Theob., Monog. Culic. IV, 542; fig. 248, wing ♀: Brun., Rec. Ind. Mus. I, 335; IV, 451.
 Sattelburg (Huon Gulf) and Friedrich Wilhelmshafen [both Papua and Biro]: Philippines [t. Ludlow]. Type of *funereus* in British Museum; of *ornata* in Hungarian Museum.
- culiciformis** Theob., Ann. Mus. Hung. III, 77, ♀, pl. i, wing; pl. iv, wing scales (*Skusea*), (1905).
Skusea culiciformis, *id.*, Monog. Culic. IV, 546, ♀, fig. 251, wing ♀: Brun., Rec. Ind. Mus. I, 335.
 Paumomu Riv., Papua [Loria]. Unique type in Hungarian Museum.
- indicus** Theob., Monog. Culic. IV, 238, ♀ (*Neomacleaya*), (1907).
Aedes indicus, Edw., Bull. Ent. Res. IV, 229.
Neomacleaya indica, Brun., Rec. Ind. Mus. IV, 458.
 nec *N. indica* var. *simplex*, Theob.
Skusea mediofasciata, Theob., Monog. Culic. IV, 544, ♂ ♀; fig. 249, wing ♀; 250, ♂ genitalia; pl. vii, wing scales ♀; pl. ix, ♂ genitalia (1907); Brun., Rec. Ind. Mus. IV, 451.
Pseudoskusea nigratarsis, Ludl., Can. Ent. XL, 52 (1908).
Pseudoskusea mediolineata, Ludl., *loc. cit.*, XL, 332 (1908).
 India [Christophers]; Woodlands, Ceylon, 9-x-07; Infanta, Tayabas, Philippine Is. Type of *indicus* and *mediofasciatus* in British Museum.

- nigrescens** Theob., Monog. Culic. IV, 540, fig. 246, wing ♀, 247, ♂ genitalia; pl. vi, wing scales, ♀ (1907).
Brun., Rec. Ind. Mus. IV, 490.
Castle Rock, Kanara Dist., S.W. India, i, ii, iii. Type in British Museum.
- jugraensis** Leices., Culic. Malaya, 109, ♂ ♀ (*Helecoctomyia, lapsus*), (1908).
Brun., Rec. Ind. Mus. IV, 455.
Jugra, near Kuala Lumpur. Type probably lost; not in Leicester's collection.
- fluviatilis** Leices., Culic. Malaya, III, ♂ ♀ (*Hulecoctomyia*), (1908).
Brun., Rec. Ind. Mus. IV, 455.
Ulu Gombok, 13 miles from Kuala Lumpur. Type probably lost; not in Leicester's collection.
- varietas** Leices., Culic. Malaya, 185, ♂ ♀ (*Aioretomyia*), (1908).
Brun., Rec. Ind. Mus. IV, 491.
Kuala Lumpur; Ulu Gombak; Pangkor Haut; Klang. Type in Leicester collection.
- singularis** Leices., Culic. Malaya 188, ♂ (*Aioretomyia*), (1908).
Aedes singularis, Edw., Bull. Ent. Res. VII, 223, fig. 8c, ♂ genitalia.
Kuala Lumpur; Kuching, Sarawak [*Moulton*]. Type in Leicester's collection.
- taeniatus** Leices., Culic. Malaya, 190, ♀ (*Aioretomyia*), (1908).
Kuala Lumpur. Type in Leicester's collection.
- perdita**, *id.*, *loc. cit.*, 192, ♂ (*Aioretomyia*), (1908).
Aioretomyia perdita, Brun., Rec. Ind. Mus. IV, 490.
Type lost.
- fragilis** Leices., Culic. Malaya, 199, ♂ (*Verrallina*), (1908).
Aedes fragilis, Edw., Bull. Ent. Res. IV, 230.
Verrallina fragilis, Brun., Rec. Ind. Mus. IV, 491.
Verrallina indecorabilis, Leices., Culic. Malaya, 200, ♂ ♀ (1908).
Brun., Rec. Ind. Mus. IV, 492.
Kuala Lumpur. Type of *fragilis* and *indecorabilis* in Leicester's coll.
- imitator** Leices., Culic. Malaya, 201, ♀ (*Verrallina*), (1908).
Kuala Lumpur. Type, damaged, in Leicester's collection.
- virilis**, *id.*, *loc. cit.*, 197, ♂ (*Verrallina*), (1908).
Kuala Lumpur. Type, a unique ♂, in Leicester's collection.
- pseudodiurna** Theob., Rec. Ind. Mus. IV, 32, ♂ (*Skusea*), (1910).
Skusea pseudodiurna, *id.*, Monog. Culic. V, 491, ♂: Brun., Rec. Ind. Mus. IV, 451.

- Sukna, 1-vii-08 [*Annandale*]. *Type* in Indian Museum, a unique specimen.
- uniformis** Theob., Rec. Ind. Mus. IV, 33, ♀ (*Skusea*), (1910).
Skusea uniformis, *id.*, Monog. Culic. V, 491, ♀.
 Pallode, Travancore, 15-xi-08 [*Annandale*]. *Type*, unique, in Indian Museum.
- pseudomediofasciatus** Theob., Monog. Culic. V, 489, ♂ (*Skusea*), (1910).
 Peradeniya and Hakgala, Ceylon, iii, iv, 07 [*Green*]. *Type* in British Museum.
- panangensis** Ludl., Psyche XXI, 159 (1914).
 Philippine Is.
- ceylonicus** Edw., Bull. Ent. Res. VII, 221, ♂, fig. 8a, ♂ genitalia (1916—17).
 Colombo, a unique ♂ [*K. McGahey*]. *Type* in British Museum.
- yerburyi** Edw., *loc. cit.*, 222, ♂, fig. 8g, ♂ genitalia (1916—17).
 Kitli Station, Ceylon, 29-xi-91 [*Yerbury*]. *Type* (unique) in British Museum.
- leicesteri** Edw., *loc. cit.*, 222, ♂, fig. 8e, ♂ genitalia (1916—17).
 Kuala Lumpur, one ♂, by stream in ampang jungle, 21-v-04 [*Leicester*].
- A. ? curtipes** Edw., Bull. Ent. Res. V, 283, ♀ (1914—1915).
 Membakut, N. Borneo [*Roper*]; Klang [*Leicester*]; also two ♀ ♀, including *type* of species from old collection of Sarawak Museum. *Type* in British Museum.

Leicesteria, Theob.

- Entom. XXXVII, 211 (1904).
 Theob., Gen. Ins., Fasc. 29, 20: *id.*, Monog. Culic. IV, 201; V, 212: Leices., Culic. Malaya, 94: Edwards, Bull. Ent. Res. IV, 255, 257, tab. orient. spp.: Brun., Rec. Ind. Mus. I, 337; IV, 454; X, 63.
 GENOTYPE, *L. longipalpis* Leices., by original designation.
- Chaetomyia*, Leices., Culic. Malaya, 100 (1908).
 GENOTYPE, *C. flava* Leices., by original designation.
- Brevirhynchus*, Theob., Rec. Ind. Mus. II, 293 (1908).
Id., Monog. Culic. V, 144, tab. spp.: Brun., Rec. Ind. Mus. IV, 442; X, 56.
 GENOTYPE, *B. magnus* Theob., by original designation.

Leicesteryomyia, Brun., Rec. Ind. Mus. IV, 452,
nom. nov. for *Chaetomyia* Leices, preocc. Brauer
and Berg., 1892 in Tachininae: Brun., Rec. Ind.
Mus. X, 63.

longipalpis Leices. in Theob., Entom. XXXVII, 211, ♂ ♀ (1904).
Theob., Monog. Culic. IV, 201, ♂ ♀: Leices., Culic. Malaya,
94, ♂ ♀. Edw., Bull. Ent. Res. IV, 261, fig. 4, ♂ geni-
talia.

Kuala Lumpur [*Leicester*]. Types in British Museum.

magna Theob., Rec. Ind. Mus. II, 293, ♂ ♀ (*Brevirhynchus*),
(1908).

Id., loc. cit., IV, pl. i, wing; pl. iii, wing scales: *id.*, Monog.
Culic. V, 145, ♂ ♀; fig. 51, wing; fig. 52, head, side view
abdom. segments: Brun., Rec. Ind. Mus. IV, 443: Edw.,
Bull. Ent. Res. IV, 259, fig. 2, ♂ genitalia.

Leicestertia annulitarsis, Leices, Culic. Malaya, 100, ♂ only
(1908): Brun., Rec. Ind. Mus. IV, 454.

Toxorhynchites rectirostris, Giles in Theob., Monog. Culic. V, 214,
nom. nud. (1910).

Sylhet, v [*Hall*]; Sukna, 1-viii-08 [*Annandale*]; Maddathorai,
Travancore, 17-xi-08 [*Annandale*]; Kuala Lumpur.

Types of *magna* ♂ ♀, in Indian Museum; of *annulitarsis* ♂,
in Leicester's collection.

dolichocephala Leices., Culic. Malaya, 95, ♂ ♀ (*dolichocephala*),
(1908).

Brun., Rec. Ind. Mus. IV, 455: Edw., Bull. Ent. Res. IV,
259.

Kuala Lumpur; Bukit Kutu (Malaysia). Types in Leices-
ter's collection.

cingulata Leices., Culic. Malaya, 97, ♀ (1908).

Brun., Rec. Ind. Mus. IV, 454: Edw., Bull. Ent. Res. IV,
260, fig. 3, ♂ genitalia.

Kuala Lumpur, iv, vi. Type in Leicester's collection.

annulitarsis Leices., Culic. Malaya, 99, ♀ only (1908).

Brun., Rec. Ind. Mus. IV, 454: Edw., Bull. Ent. Res. IV,
260.

Kuala Lumpur. Type in Leicester collection.

N.B.—*L. annulitarsis*, Leices. ♂ = *L. magna*, Theob.

flava Leices., Culic. Malaya, 101, ♂ ♀ (*Chaetomyia*), (1908).

Leicestertia flava, Edw., Bull. Ent. Res. IV, 258, fig. 1, ♂ geni-
talia.

L. apicalis, Theob., Rec. Ind. Mus. II, 291, ♂ ♀ (1908); IV,
pl. i, wing; pl. iii, wing scales: *id.*, Monog. Culic. V, 213,
♂ ♀; fig. 86, wing: Brun., Rec. Ind. Mus. IV, 454.

Brevirhynchus apicalis, Theob., Rec. Ind. Mus. IV, 7, ♀ (1910):
Theob., Monog. Culic. V, 148: Brun., Rec. Ind. Mus. IV,
443.

Leicesteriomyia flava, Brun., Rec. Ind. Mus. IV, 453.

Malay Peninsula. *Types (flava)* in Leicester collection; of *apicalis* in Indian Museum.

annulipalpis Theob., Rec. Ind. Mus. IV, 6, ♀ (*Brevirhynchus*), (1910).

Leicesteria annulipalpis, Edw., Bull. Ent. Res. IV, 257, fig. 1, ♂ genitalia.

Brevirhynchus annulipalpis, Theob., Monog. Culic. V, 148: Brun., Rec. Ind. Mus. IV, 442.

Maddathorai, Travancore, 16-xi-08 [*Annandale*]. *Type*, unique in Indian Museum. May possibly not belong to *Leicesteria*.

pendula Edw., Bull. Ent. Res. IV, 261, ♂, fig. 5, ♂ genitalia (1914).

Between Selangor and Pahang, Malay States, 16-vi-04 [*Leicester*]. *Type* in British Museum.

digitata, *id.*, *loc. cit.*, IV, 262, ♂ ♀, fig. 6, ♂ genitalia (1914).

Ulu Gombak, Malay States, 28-viii-03; 2-ix-03, *types* [*Daniels*]; between Selangor and Pahang, 16-iv-04 [*Leicester*]; Pahang, near Kuala Lumpur, 4-iv-03; 18-x-03 [*Leicester*]; Mailum Negros, Philippine Is., 9-vi-06 [*Banks*]. *Types* in British Museum.

pectinata, *id.*, *loc. cit.*, IV, 263, ♂ ♀, fig. 7, ♂ genitalia (1914).

Jugra, 3-7-x-04, *type* ♂ [*Leicester*]; Sarawak [*Hewitt*]. *Type* ♂ in British Museum.

omissa, *id.*, *loc. cit.*, V, 76, ♂ ♀ (1914).

Colombo [*James*]; Peradeniya [*Green*]. *Types* in British Museum. Possibly = *dolichocephala*, Leices., var.

Taeniorhynchus, Arrib.

Rev. Mus. La Plata, II, 147 (1891).

Theob., Monog. Culic. II, 190 (restricted); *loc. cit.*, IV, 483, generic notes, etc.; V, 419, tab. spp.: *id.*, Gen. Ins., Fasc. 26, 30, modified: Giles, Handb., 2nd ed., 358: Blanch., Moust., 381, fig. 244, ♂ ♀, generic chars.: Leices., Culic. Malaya, 163: Edw., Bull. Ent. Res. II, 251: Brun., Rec. Ind. Mus. I, 356; IV, 478; X, 72, note on genotype.

GENOTYPE, *Culex titillans* Walk., as *C. taeniorhynchus* W.

Mansonia, Blanch., Comp. rend. So. biol. Paris (1901), LIII, 1046, *nom. nov.* for *Panoplites*, Theob., preocc. Gould, 1853, in Aves.

Blanch., Moust., 375, 356, tab. spp.: Theob., Gen. Ins., Fasc. 26, 31: *id.*, Monog. Culic. IV, 494: V, 446, tab. spp.: Leices., Culic. Malaya, 171: Brun., Rec. Ind. Mus. I, 358; IV, 81; X, 64: Howard, Dyar and Knab, Mosq. N. Amer., etc., III, pt. 1, 501.

- Panoplites*, Theob., Rep. Coll. Mosq. Brit. Mus., 5 (1900).
Id., Monog. Culic. II, 173 (1901): Giles, Handb., 2nd ed., 350.
 GENOTYPE, *Culex titillans* Walk., as *Taeniorhynchus taeniorhynchus*, Wied., by Neveu Lemaire's designation.¹
- Coquillettidia*, Dyar, Proc. Ent. Soc. Wash. VII, 47 (1905).
 GENOTYPE, *Culex perturbans*, Walk. [(1910).
Rhynchotaenia, Brethes., Ann. Mus. Buen. Ayres. XX, 470
 GENOTYPE, *Taeniorhynchus fasciolatus* Arrib., by designation of Howard, Dyar and Knab (1915).
- acer** Walk., List. Dipt. Brit. Mus. I, 8, ♀ (*Culex*), (1848).
 nec *acer*, Theob., Monog. Culic. II, 211 (1901).
 Friedrich Wilhelmshafen; Mt. Hauseman, Astrolabe Bay; Yomba [all Papua and Boro]. Also occurs in Queensland and New Zealand. Type in British Museum.
- annulipes** Walk., Proc. Linn. Soc. Lond. I, 6, ♀ (*Culex*), (1857).
Mansonia annulipes, Leices., Culic. Malaya, 172, ♂ ♀: Brun., Rec. Ind. Mus. I, 359; IV, 482, notes.
Panoplites annulipes, Theob., Monog. Culic. II, 185, ♀; pl. xxx, 119, ♀, full ins. col.
Culex dives, Sch., Reise Novara, 31 (1868): Giles, Handbk., etc., 261.
Culex nero, Doles., Nat. Tijd. Ned. Ind. XIV, 383, pl. v, 3 (1857): Giles, Handbk., etc., 284.
Mansonia nero, Blanch., Moust., 380, ♀.
 Calcutta, 6-x-04 [*Brunetti*]; Selangor, 28-x-99 and ix [*Butler*]; Perak, "very abundant nocturnal species," [*Wright*]; Dindings, Straits, xi, xii [*Wright*]; Kuala Lumpur [*Durham* and *Leicester*]; Klang; Port Swettenham and Jugra [all *Leicester*]; Batavia [*t. Schiner*]; Rio Baco, Mindoro, Philippine Is. [*McGregor*]; Gombong, Mid-Java [*t. Doleschall*]. Types: I have no information about any of these.
- confinnis** Arrib., Revis. Mus. La Plata II, 149 (1891).
 Theob., Monog. Culic. III, 289.
Culex confinnis, Giles, Handbk., etc., 247.
 Chaca, Formosa.
 Much doubt attaches to the correct identity of this species, which is a South American one.
- brevicellulus** Theob., Monog. Culic. II, 212, ♂ ♀; fig. 255, 256, var. chars.; pl. xxiii, 89, full ins. col. ♂ (1901).
Taeniorhynchus brevicellulus, *id.*, *loc. cit.*, III, 268 (correction in wing): Giles, Handbk., 2nd ed., 363, ♀ (*brevicellula*): Blanch., Moust., 389, ♂ ♀ (*brevicellula*): Leices., Culic. Malaya, 163, ♂ ♀: Brun., Rec. Ind. Mus. I, 357: Edw., Bull. Ent. Res. IV, 230.
Chrysoconops brevicellulus, Brun., Rec. Ind. Mus. IV, 481.

¹ For selection of and identification of type species, see Howard, Dyar and Knab, Mosq. N. Amer., etc., III, pt. 1, 502.

- Taeniorhynchus acer*, Theob. nec Walk., Monog. Culic. II, 211 (1901).
- Chrysoconops pygmaeus*, Theob., Rec. Ind. Mus. II, 300, ♀ (1908): *id.*; Monog. Culic. IV, 25, ♂; pl. i, wing; iii, wing scales: *id.*, *loc. cit.*, V, 435, ♂ ♀; fig. 192, wing, 193, head, ♂, 194, ♂ genitalia, 195, wing, ♀: Brun., Rec. Ind. Mus. IV, 481.
- Chrysoconops fuscopteron*, Theob., Tijd. v. Ent. LIV, 239 (1911).
Sylhet [Hall]; Saugar; Manipur, viii [Gourlav]; Manipur Haut, ii, v, vi; Calcutta, vii, viii; Purnea and Rajmahal, Bengal; Selangor; Perak; Thayetmo, Upper Burma, viii; Ceylon [Green]; Kuala Lumpur [Leicester]; Sungei Penoh, Korinchi Valley, Sumatra, 2,700 ft.; Siolak-Daras, 3,100 ft., Sumatra, viii-14 [both *t. Stanton*]; Philippine Is. [*t. Ludlow*].
Types: *brevicellulus*, in British Museum; *pygmaeus*, ♂ ♀, in Indian Museum (*cotype* in British Museum); *acer*, Theob. (nec Walk.), in Brit. Mus.; *fuscopteron*, in Amsterdam Museum.
- annulatus* Leices., Culic. Malaya, 174, ♀ (*Mansonia*), (1908)
Brun., Rec. Ind. Mus. IV, 482.
Kuala Lumpur. *Type* in Leicester's collection.
- aureosquammatum* Ludl., Can. Ent. XLI, 234, ♀ (1909).
Theob., Monog. Culic. V, 425 (*aureosquamata*, *lapsus?*).
Taeniorhynchus pagei, Ludl., Psyche XVIII, 128: Theob., Monog. Culic. V, 618.
Parang, Mindanao, Philippine Is., 27-x-xii. *Types* of *aureosquammatum* and *pagei* in Army Medical Museum, Washington.
- chrysozona* Knab., Ent. News. Phil. XX, 386, ♂ ♀ (*Mansonia*), (1909).
Brun., Rec. Ind. Mus. IV, 482.
Chrysoconops aurites, Ludl. (nec Theob.).
Parang, Mindanao, Philippine I., 31-v-06. *Type* in U.S. Nat. Museum, cat. no. 12,626.
- nigrosignatus* Edw., Bull. Ent. Res. VII, 224, nom. nov. for *conopas*, Theob. nec. Fraunfld. (1916-17).
Taeniorhynchus conopas, Theob., Monog. Culic. I, 202.
nec *Culex conopas*, Frnfd.
? *Pseudotaeniorhynchus conopas*, var. *giblini*, Taylor, Tr. Ent. Soc. Lond., p. 198, ♀ (1914).
Lake Kamu Gold Field, Papua [Dr. Giblin]. *Type* in Australian Institute of Tropical Medicine. Edwards considers *giblini* sufficiently distinct to warrant ranking it as a good species.

Mansonioides, Theob.

Monog. Culic. IV, 498 (1907).

Theob., *loc. cit.*, V, 452, tab. spp.: Brun., Rec. Ind. Mus. IV, 483; X, 64: Edw., Bull. Ent. Res. II, 253.GENOTYPE, *M. guttata* Theob., by original designation.

uniformis Theob., Monog. Culic. II, 180, ♀; pl. xxx, 118, full ins. col., ♀ (*Panoplites*), (1901).

Mansonia uniformis, *id.*, *loc. cit.*, III, 270, fig. 144, pupa: Blanch., Moust., 379, ♀: Brun., Rec. Ind. Mus. I, 359; IV, 483; Leices., Culic. Malaya, 171, ♂ ♀: Edw., Bull. Ent. Res. II, 254.

Panoplites uniformis, Giles, Handbk., 2nd ed., 353; pl. xiii, 3 a-d, ♂, head and venation ♂ ♀ (1902).

Panoplites africanus, var. *reversus*, Theob., Monog. Culic. II, 189 (1901).

nec *P. africanus*, typical form, Theob., *loc. cit.*, 187.

Panoplites australiensis, Theob. in Giles, Handbk., 2nd ed., 355 (1902).

? *Mansonia seguini*, Laveran, Comp. Rend. Soc. Biol. Paris LIII, 992 (*Panoplites*), (1901): Blanch., Moust., 380, ♀: Theob., Monog. Culic. IV, 496; Brun., Rec. Ind. Mus. IV, 482.

Sylhet, i, ii, iii, v, vi, vii, xii [Hall]; Manipur, vi, ix; Shahjahanpur, N.W. India, x [Giles]; Bhogaon, 30-ix-08 and x; Kati-har, x [both Bengal and Paiva]; Balighai, near Puri, Orissa, 23-x-08; Gopkuda I., Chilka Lake, viii; Calcutta, 2-vii-07; Travancore and Cochin States, 19-xi-08, 4-xi-08 [all Annandale]; Quilon, 7-iv-00 [James]; Taiping [Wray]; Dilo, Friedrich Wilhelmshafen and Ins. Graget (all Papua) [Biro]; Bayembang Pangasinan, Philippine Is. [Chamberlain]; Manila; Rizal; Fort McKinley [Banks, Schultze, Craig, Araneta]; Rangoon, 25-ii-08 [Annandale]; Kuala Lumpur [Leicester]. Type of *uniformis* in British Museum.

According to Edwards' latest investigations, *africanus*, Theob., typical form, is a valid species, with *Mansonia major*, Theob. and *Mansonia nigerrima*, Theob., as synonyms, the species not being oriental; but *africanus*, var. *reversus* is synonymous with *uniformis*.

annulifera Theob., Monog. Culic. II, 183, ♀ (*Panoplites*), fig. 224, wing; pl. xxx, 120, ♀, full ins. coll. (1901).

Panoplites annulifera, Giles, Handbk., 2nd ed., 356, ♀; pl. xiii, 8, hind leg.

Mansonia annulifera, Theob., Monog. Culic. III, 274: Blanch., Moust., 380, ♀: Leices., Culic. Malaya, 174, ♀ (Theob.'s desc. copied): Brun., Rec. Ind. Mus. I, 359; IV, 482.

Mansonioides annuliferus, Edw., Bull. Ent. Res. IV, 230.

Mansonia 7-punctata, Theob., Ann. Mus. Hung. III, 187 (1905).

Id., Monog. Culic. IV, 494, ♀.

Mansonioides 7-guttata, Theob., Monog. Culic. IV, 499, ♀; figs. 226, 227, wings, ♀ (1907).

Calcutta, viii, xii; Port Canning, xii [Annandale]; Purneah, x; Bhogaon, 7-viii-09 [both Bengal and Paiva]; Bengal [Macrae]; Dacca; Sylhet, v, vi [Hall]; Manipur [Gourlay]; Madras, 12-xii-99 [Goodrich]; Quilon [James]; Ceylon [t. Banks];

Singapore [Durham]; Perak [Wright]; Sarawak [Barker]; Friedrich Wilhelmshafen, Papua, xi, 10 [Biro]; Bayembang (Pangasinan, Philippine Is.) [Chamberlain]; Manila [Banks].

Type of *annulifera* and *7-guttata* in British Museum; that of *7-punctata* in Hungarian Museum.

Culiciomyia, Theob.

Monog. Culic. IV, 227 (1907).

Theob., *loc. cit.*, V, 229, tab. spp.: Brun., Rec. Ind. Mus. IV, 456; X, 58; Edw., Bull. Ent. Res. IV, 234.

GENOTYPE, *C. inornata* Theob., by my own designation.

Neomelanoconion, Theob., Monog. Culic. IV, 514 (1907), ♂ only (*t.* Edw., Bull. Ent. Res. II, 254). GENOTYPE, *Culex rima* Theob., by original designation.

Pectinopalpus, Theob., Ann. Mag. Nat. Hist. (8) V, 375 (1910).

GENOTYPE, *P. fuscus* Theob., by original designation.

Trichorhynchus, Theob., Jour. Bomb. Nat. Hist. Soc. XVI, 241 (1905); Theob., Monog. Culic. IV, 270; Brun., Rec. Ind. Mus. I, 356; X, 73.

Trichorhynchomyia, Brun., Rec. Ind. Mus. IV, 477, *nom. nov.* for *Trichorhynchus*, Theob., preocc Balbiani 1887 in Protozoa: Brun., *loc. cit.*, X, 73.

viridiventer Giles, Jour. Bomb. Nat. Hist. Soc. XIII, 609 (*Culex*, provis. descr.), (1901).

Culex viridiventer, Theob., Monog. Culic. II, 128, ♂ ♀, fig. 219, claws, ♂ ♀, abdom. segments, wing scales; ♂ palpus and proboscis, fig. 220, larva; pl. xxix, 116, ♀, full ins. col.: Giles, Handbk., etc, 2nd ed., 445, ♂ ♀; pl. xvii, 12, ♂ claw, venation, abdomen, larva: Blanch., Moust., 346, ♂ ♀: Brun., Rec. Ind. Mus. I, 354; Edw., Bull. Ent. Res. IV, 235.

Culex angulatus, Theob., Monog. Culic. II, 324, ♀ (1901). Blanch., Moust., 362, ♀.

Culex longifurcatus, Theob. (nec Becker), Rec. Ind. Mus. IV, 19, ♂ ♀ (1910).

Culex pseudolongifurcatus, Theob. (*nom. nov.* for *C. longifurcatus*, preocc. by Becker), Monog. Culic. V, 366 (1910).

Naini Tal, 4000 ft., vi; 7000 ft., vi, vii [both Giles]; Katmandu, Nepal; Dahawanga Hill, Bengal, Nepalese Frontier, 16-ii-08. Types of *viridiventer*, *angulatus* and *longifurcatus* in British Museum; that of *pseudolongifurcatus* in Indian Museum.

pallidothorax Theob., Jour. Econ. Biol. I, 32 (*Culex*), (1905).

Theob., Monog. Culic. IV, 446, ♂ ♀ (*pallidothorax* emend.). Edw., Bull. Ent. Res. IV, 235.

Culex albopleura, Theob., Monog. Culic. IV, 456, ♀ (1907).

- Culiciomyia annuloabdominalis*, Theob., Monog. Culic. V, 236, ♂ ♀, fig. 102, occiput, 103, head (1910).
India [James, Christophers]; Peradeniya, ii-07 and Hakgala, Ceylon, v-07 [both Green]. Types of *pallidothorax*, *albopleura* and *annuloabdominalis* in British Museum.
- pulla* Theob., Ann. Mus. Hung. III, 87, ♀, fig. 6, head (*Culex pullus*), (1905).
Theob., Monog. Culic. IV, 232, fig. 66, head scales.
Muina, Papua, 1900 [Biro]. Type, unique, in Hungarian Museum.
- fusca* Theob., Jour. Bomb. Nat. Hist. Soc. XVI, 242, ♀; pl. A. 2, head, palpus, clypeus, antenna, scutellum (*Trichorhynchus fuscus*), (1905).
Culiciomyia fusca, Edw., Bull. Ent. Res. IV, 235.
Trichorhynchus fuscus, Theob., Monog. Culic. IV, 270, ♀; Edw., Bull. Ent. Res. III, 33; IV, 235.
Trichorhynchomyia tuscus, Brun., Rec. Ind. Mus. IV, 477.
Culiciomyia inornata, Theob., Monog. Culic. IV, 227, ♂ ♀, fig. 61, head scales; 62, wing, ♀; 63, ♂ genitals (1907).
Peradeniya, xii [Green]; Kuching, xi [Barker]; Philippines [t. Ludlow]. Type: Theobald wrote me that it was in the Hungarian Museum, subsequently (Monog. V), stating it to be in the British Museum; that of *inornata* is there.
- annulata* Theob., Monog. Culic. IV, 230, ♂ ♀, fig. 64, wing, ♀; 65, ♂ genitalia; pl. i, wing scales, ♀ (1907).
Brun., Rec. Ind. Mus. IV, 457.
Kuching, Sarawak [Barker]; Philippines [t. Ludlow].
Type in British Museum.
- ceylonica* Theob., Monog. Culic. V, 236, ♂ ♀, fig. 70, wing, ♀ (1907).
Peradeniya, ii-07; Maskeliya, Ceylon, iv-07 [both Green].
Types in British Museum, a unique pair.
- bahri* Edw., Bull. Ent. Res. V, 79, ♂ ♀ (1910).
Badulla, Ceylon; a series bred from larvae, 3 ♂ ♂, including type, and 4 ♀ ♀ in British Museum; *coltypes* in London School of Tropical Medicine; Hakgala, Ceylon; Peradeniya [Green].
Type in British Museum.
- spathifurca* Edw., Bull. Ent. Res. V, 284, ♂ (1914-15).
Kuching, Sarawak, 22-vii-14, in house [Moulton]; a ♀ from Kuching reservoir, 30-vii-14, may be this species. Type in British Museum.

Lophoceratomyia, Theob.¹

Ann. Mus. Hung., III, 93 (1905).

Theob., Jour. Bomb. Nat. Hist. Soc. XVI, 245:

id., Monog. Culic. IV, 471 (generic notes):

Leices., Culic. Malaya, 119.

¹ In Edwards' latest paper he ranks this as a subgenus only of *Culex*, and gives a table of oriental species.

GENOTYPE, *L. fraudatrix* Theob., by my own designation (Rec. Ind. Mus. X, 64)

Philodendromyia, Theob., Monog. Culic. IV, 623 (1907).

GENOTYPE, *P. barkeri* Theob., the original species.

Oculeomyia, pt. Theob., *loc. cit.*, IV, 515 (1907).

GENOTYPE, *O. sarawaki* Theob., the original species.

uniformis Theob., Ann. Mus. Hung. III, 93, ♂ ♀; pl. A, 3, antenna, ♂; pl. B, 4, palpus (1905).

Id., Jour. Bomb. Nat. Hist. Soc. XVI, 245: *id.*, Monog. Culic. IV, 473, ♂ ♀; Edw., Bull. Ent. Res. V, 80; VII, 227. Peradeniya, v [Green]. Type in British Museum.

fraudatrix Theob., Ann. Mus. Hung. III, 94, ♂ ♀, fig. 7, palpus ♂ ♀, 8, antennal organs (1905).

Id., Monog. Culic. IV, 474, ♂ ♀, fig. 208—211, var. chars.: Edw., Bull. Ent. Res. IV, 235.

var. **variata** Leices., Culic. Malaya, 121, ♂ ♀ (1908).

Friedrich Wilhelmshafen and Stephansort [both Papua and *Biro*]; Malay Peninsula. Type of *fraudatrix* in Hungarian Museum; of *variata* in Leicester coll.

brevipalpis Theob., Ann. Mus. Hung. III, 96, ♂, fig. 9, var. chars. (1905).

Id., Monog. Culic. IV, 477, ♂, fig. 12, var. chars.: Leices., Culic. Malaya, 129 (Theob. descr. copied): Edw., Bull. Ent. Res. IV, 236.

eminentia, Leices., Culic. Malaya, 131, ♂ (1908).

Singapore [*Biro*]; Kuala Lumpur. Type of *brevipalpis* in Hungarian Museum; of *eminentia* in Leicester collection.

Edwards suggests the use of the name *eminentia*, Leices. instead of the earlier *brevipalpis*, to avoid confusion with *Culex* (*Cyathomyia*) *brevipalpis*, Giles, but this does not appear to me to be legitimate.

minutissima Theob., Monog. Culic. IV, 135, ♀ (*Culiciomyia*), (1907).

Lophoceratomyia minutissima, Edw., Bull. Ent. Res. IV, 235.

Culiciomyia nigerrima, Theob., Monog. Culic. V, 233, ♀, fig. 100, wing (1910).

Melanoconion juxtapallidiceps, *id.*, *loc. cit.*, V, 456, ♀ (1910).

Peradeniya, ii [Green]; Trincomalee, x, 1907 [Green].

Types: all three in the British Museum.

barkeri Theob., Monog. Culic. IV, 623, ♂, fig. 289, head, scutellum, palpus, ♂; fig. 290, wing, ♂; pl. vii, wing scales, ♀ (*Philodendromyia*), (1907).

Brun., Rec. Ind. Mus. IV, 503: Edw., Bull. Ent. Res. IV, 236; VII, 227.

- sylvestris*, Leices., Culic. Malaya, 125, ♂ ♀ (1908): Brun., Rec. Ind. Mus. IV, 465.
- quadripalpis*, Edw., Bull. Ent. Res. V, 80, ♂ (1914).
Sarawak, vii [Barker]; Malaysia. Types of *barkeri* and *quadripalpis* in British Museum; paratypes in Cambridge Museum.
- rubithoracis** Leices., Culic. Malaya, 119, ♂ ♀ (1908).
Brun., Rec. Ind. Mus. IV, 464: Edw., Bull. Ent. Res. IV, 236.
Oculeomyia fulleri, Ludl., Can. Ent. XLI, 97, ♀ (*Oculiomyia*), (1909): Theob., Monog. Culic. V, 478, ♀ (Ludl. descr. copied):
Brun., Rec. Ind. Mus. IV, 485.
- ? *Lophoceratomyia niger*, Leices., Culic. Malaya, 123 (1908).
Kuala Lumpur; Parang; Mindanao, Philippines, x. Type of *rubithoracis* in Leicester's collection; of *fulleri* in Army Medical Museum, Washington.
- nigra** Leices., Culic. Malaya, 123, ♂ ♀ (*niger*), (1908).
Brun., Rec. Ind. Mus. IV, 464.
Kuala Lumpur. Types in Leicester collection.
Edwards notes that if *Lophoceratomyia* be considered only a subgenus of *Culex* the name *nigra* is preoccupied, but thinks Leicester's species is probably synonymous with *rubithoracis*.
- minor** Leices., Culic. Malaya, 126, ♂ ♀ (1908).
Brun., Rec. Ind. Mus. IV, 464: Edw., Bull. Ent. Res. VII, 227.
Malaysia. Type in Leicester collection.
It is possible, as Edwards notes, that this name may be antedated by *Culex minor*, Theobald, Oct. 1908.
- taeniata** Leices., Culic. Malaya, 127, ♂ ♀ (1908).
Brun., Rec. Ind. Mus. IV, 465.
Kuala Lumpur; Klang. Type in Leicester collection.
See note under *nigra*, the same argument applying here, *taeniata* possibly being synonymous with some other species (Edwards here repeating the name *taeniata*, apparently in error for some other).
- mammilifer** Leices., Culic. Malaya, 128, ♂ ♀ (1908).
Brun., Rec. Ind. Mus. IV, 464: Edw., Bull. Ent. Res. IV, 236.
bicornuta, Theob., Rec. Ind. Mus. IV, 25, ♂ (1910).
Id., Monog. Culic. V, 412, ♂, figs. 178-182, var. chars.:
Brun., Rec. Ind. Mus. IV, 463.
Kuala Lumpur; Raub (Malaysia); Kawkareik, base of Dawna Hills, 4-iii-08 [Annandale]. Type of *mammilifer* in Leicester coll.; of *bicornuta* a dissected ♂ in the Indian Museum.
- hewitti** Edw., Bull. Ent. Res. V, 80, ♂ ♀ (1914).
Matang, Sarawak [Hewitt]. Type in British Museum; paratypes in Cambridge Museum.
- curtipalpis** Edw., Bull. Ent. Res. V, 127, ♂ (1914-15).
Kuching reservoir, Sarawak, 6-iii-14 [Moulton]. Type in British Museum.

Culex, Linn.

Syst. Ent., Ed. X, 602 (1758).

Sch., Fauna Austr. II, 625: Giles, Handbk., etc., 186; 2nd ed., 386: Theob., Monog. Culic. I, 326; IV, 387; V, 323, tab. spp.: *id.*, Gen. Ins., Fasc. 26, 24; 25, list spp. world: Blanch., Moust., 267, 269, tab. spp. in 9 sections; 627, supp. tab.; 372, list of uncertain species: Leices., Culic. Malaya, 138, tab. of groups: Brun., Rec. Ind. Mus. I, 342; IV, 465; X, 39. *Culex* group of genera; 42, subgenera; 43, synonyms: Alcock, Ent. Med. Off., 98: How., Dyar and Knab, Mosq. N. Amer., etc., III, pt. 1, 215, list synonyms; 217, chars.

GENOTYPE, *Culex pipiens* L., by universal acceptance.

Lutzia, Theob., Monog. Culic. III, 155 (1903).¹

How., Dyar and Knab, Mosq. N. Amer., etc., III, 466.

GENOTYPE, *Culex bigotii* Bell, by original designation.

Melanoconops, Theob., Monog. Culic. III, 178, *nom. nud.* (1903).²

Lasioconops, Theob., *loc. cit.*, III, 235 (1903).

GENOTYPE, *L. poecilipes*, *id.*, sp. nov., *loc. cit.*

Melanoconion, Theob., Monog. Culic. III, 238 (1903).

Theob., *loc. cit.*, IV, 507; V, 455, tab. spp.: *id.*, Gen. Ins., Fasc. 26, 32: Blanch., Moust., 395 (*Melanoconium*, emend.): Glen Herrick, Ent. News., 382: Leices., Culic. Malaya, 136: Brun., Rec. Ind. Mus. I, 360; IV, 484; X, 65.

GENOTYPE, *Culex atratus* Theob., by Dyar's designation.

Heptaphlebomyia, Theob., Monog. Culic. III, 336 (1903).

Theob., *loc. cit.*, IV, 531; V, 481: Gen. Ins., Fasc. 26, 41: *id.*, Entom. XXX, 156.

GENOTYPE, *H. simplex* Theob., sp. nov., *loc. cit.*

Culicella, Felt., N. Yk. State Mus. Bull. 79: Ent. 22, App., p. 391c (1904).

How., Dyar and Knab, Mosq. N. Amer., etc., III, 457 (as subgenus).

GENOTYPE, *Culex dyari* Coq., by original designation.

Pseudoheptaphlebomyia, Ventrillon, Bull. Mus. Paris XI, 427, *nom. nud.* (1905).

GENOTYPE, *P. madagascarensis* Ventr.

Neoculex, *id.*, *loc. cit.*, VII, 47 (1905).

GENOTYPE, *Culex territans* Walk., by original designation.

¹ *Lutzia* is ranked as a valid genus by Howard, Dyar and Knab, with a suggestion (p. 218), that it might be a subgenus of *Culex*, the same opinion applying to *Carrollia* also.

² The name merely used in the text and not set up as a new genus even as a *nom. nud.*

- Gnophodeomyia*, Theob., Jour. Econ. Biol. I, No. 1, 21 (1905):
Id., Monog. Culic. IV, 251.
 GENOTYPE, *G. inornata*, sp. nov., *loc. cit.*
- Trichopronomyia*, Theob., Ann. Mus. Hung. III, 98 (1905).
 Theob., Monog. Culic. IV, 479: Brun., Rec. Ind. Mus.
 I, 356; IV, 477; X, 73.
 GENOTYPE, *T. annulata*, sp. nov., *loc. cit.*
- Melanoconium*, Blanch., Moust., 395 (*Melanoconion*, emend.),
 (1905).
- Popea*, Ludl., Can. Ent. XXXVII, 95 (1905).
 Theob., Monog. Culic. IV, 211: Brun., Rec. Ind. Mus.
 IV, 485; X, 69.
 GENOTYPE, *P. lutea*, sp. nov., *loc. cit.*
- Mochlostyrax*, Dyar and Knab, Jour. N. Yk. Ent. Soc. XIV,
 223 (1906).
 GENOTYPE, *M. caudelli* Dyar and Knab, by original designation.
- Jamesia*, Chris., Sci. Mem. Med. Off. India (n. s.) XXV, 12
 (1906).
 No definite type species quoted. *Culex concolor*, Rob.
 Desv. and *C. nigripes*, Zeit. are quoted as belonging to
 it.
- Tinolestes*, Coq., Proc. Ent. Soc. Wash. VII, 185 (1906).
 GENOTYPE, *T. latisquama*, *id.*, sp. nov., *loc. cit.*
- Maillotia*, Theob., Monog. Culic. IV, 274 (1907).
 GENOTYPE, *M. pilifera*, *id.*, sp. nov., *loc. cit.*
- Aporoculex*, *id.*, *loc. cit.*, 316 (1907).
 GENOTYPE, *A. punctipes*, *id.*, sp. nov., *loc. cit.*
- Leucomyia*, *id.*, *loc. cit.*, 372; pl. ix, ♂ genitalia (1907).
Id., *loc. cit.*, V, 311, tab. spp.
 GENOTYPE, *Culex gelidus* Theob., by original designation.
- Microculex*, *id.*, *loc. cit.*, IV, 461 (1907).
 GENOTYPE, *M. argenteoumbrosus*, *id.*, sp. nov., *loc. cit.*
- Neomelanoconion*, *id.*, *loc. cit.*, IV, 514, ♀ only (1907).¹
 GENOTYPE, *Culex rima* Theob., by original designation.
- Pneumaculex*, Theob., Monog. Culic. IV, 523 (1907).
 Dyar, Proc. Ent. Soc. Wash. VII, No. 1, *nom. nud.*
 GENOTYPE, *Culex signifer*, Coq.
 Dyar's reference was a *nom. nud.*, and the genus must
 stand to Theobald.
- Aedinus*, Lutz in Peryassu, Os Culic. do Bras., 36 (1908).
 GENOTYPE, *A. amazonensis*, sp. nov., *loc. cit.*
- Gnophodromyia*, Theob., Monog. Culic. V, 114 (1910).
 Apparently a *nomen nudum* figuring only in a table of
 genera, probably a misprint for *Gnophodeomyia*.

¹ Howard, Dyar and Knab say that Newstead, Dutton and Todd used this name previously (Ann. Trop. Med. Paras. 1, 31, 1907), with *N. palpus*, sp. nov., as genotype. Edwards ranks the ♂ of *Neomelanoconion*, as synonymous with *Culiciomyia*.

- Rachisoura*, Theob., *loc. cit.*, V, 207 (1910).
 GENOTYPE, *R. sylvestris*, sp. nov., *loc. cit.*
- Mimelcomyia*, *id.*, *loc. cit.*, V, 210 (1910).
 GENOTYPE, *E. apicotriangulata*, sp. nov., *loc. cit.*
- Eumelanomyia*, *id.*, *loc. cit.*, V, 240 (1910).
 GENOTYPE, *E. inconspicua*, sp. nov., *loc. cit.*
- Protomelanoconion*, *id.*, *loc. cit.*, V, 462 (1910).
 GENOTYPE, *P. fusca*, sp. nov., *loc. cit.*
- Thomasina*, Newstead and Carter, Ann. Trop. Med. Paras. IV, 553, fig. 1, head, ♂ (1910—11).
 GENOTYPE, *Mansonia longipalpis* Newst. and Carter, by original designation.
- Diceromyia*, Theob., 4th Rep. Wellc. Lab., Vol. B, 151 (1911).
 GENOTYPE, *D. africana*, *id.*, sp. nov., *loc. cit.*
- Theobaldiomyia*, Brun., Rec. Ind. Mus. IV, 462; nom. nov. for *Leucomyia*, Theob., preocc. Braur and Berg., 1892 in Sarcophaginae.
 Brun., Rec. Ind. Mus. X, 72 (1912).
- Climacura*, Howard, Dyar and Knab, Mòsq. N. Amer., etc., III, 452 (1915).
 GENOTYPE, *Culex melanurus* Coq., by original designation.
 Ranked as a subgenus by the authors.

Of the synonyms of *Culex* as given by me in my recent "Critical Review," 16 (not counting my *nomina novae*) are accepted as such by Messrs. Howard, Dyar and Knab, whilst 31 others are considered synonymous with *Aedes*, with which they include *Ochlerotatus*, as understood by Edwards, two or three of the remainder of my *Culex* synonyms being split up between *Culex* and *Aedes*.

- cantans* Mg., Syst. Besch. I, 6, ♀ (1818).
Culex stimulans, Walk., List Dipt. Brit. Mus. I, 4, ♀ (1848).
C. fumipennis, Steph., Zool. Jour. I, 453 (1825).
 Coonoor, 6,000 ft.; Madras; North India [Dr. Price]. Types in Paris Museum, two specimens.

- concolor* R. Desv., Mem. So. Hist. Nat. Paris IV, 405 (1825).
 Theob., Monog. Culic. II, 107, ♂ ♀, fig. 203, ♂ palpus; pl. xxviii, 109 ♂, 110 ♀ both full ins. col.: Giles, Jour. Trop. Med. VII, 368: *id.*, Handb., 2nd ed., 454, ♂ ♀; pl. xvii, 8, a, b, venation, ♂ ♀: Blanch., Moust., 365, ♂ ♀: Leices., Culic. Malaya, 154, ♂ ♀: Brun., Rec. Ind. Mus. I, 343; IV, 468, notes.

Practically common throughout India in the summer and many parts of the East, occurring in some places during the greater part of the year, Calcutta for instance. Outside of India it is found in Sylhet, ii, iv, v, xi, xii [Lt.-Col. Hall]; Upper Burma in August [Watson]; Selangor, 28-x-99 [Butler]; Kuala Lumpur [Durham]; Perak [Wray and Wright]; Samarang [Jacobson]; Phrapatoon, Siam, viii-06 [Woolley]; Pampanga, Phil. Is. [Whit-

more]; Hongkong; Foo Chow (China), Andaman Is. [*Ray White*].
Type location unknown to present author.

fuscatus W., Auss. Zweifl. I, 6 (1828).

Giles, Handbk., etc., 301: Theob., Monog. Culic. II, 167:
id., Gen., Ins. Fasc. 26, 30 (quotation incorrect): Giles,
Handbk., etc., 2nd ed., 455: Blanch., Moust., 275: Brun.,
Rec. Ind. Mus. I, 345.

East India; Singapore; Malacca; Sarawak. Type (no sex
stated), location unstated.

Much doubt attaches to this species (*vide* Brun., Rec. Ind.
Mus. I, 345).

fatigans Wied., Auss. Zweifl. I, 10, ♂ ♀ (1828).

Giles, Handbk., 298, 2nd ed., 438; Theob., Monog. Culic.
II, 151, ♂ ♀, fig. 234, ♂, wing; fig. 235, ♂ genitalia; fig.
236, wings, map of distribution, p. 155; pl. xxix, 114, ♂,
115, ♀ (both full ins. col.); pl. D, wing scales: list. vars.
and locs., p. 156, *et. seq.*; fig. 238, ten variations of wings:
Theob., Gen. Ins. Fasc. 26; pl. ii, 2, ♀ (full ins. col.): Giles,
Handbk., 2nd ed., 438, ♂ ♀, fig. 45 (var. chars.); list of
subspecies, p. 440: Theob., Monog. Culic. IV, pl. xvi, lar-
val characters: Blanch., Moust., 353, ♂ ♀, fig. 230, ♂ ♀
(as *pungens*); 231, ♀; 232, head, ♂ ♀ (as *anxifer*); 233,
adult larva, nymph (*pungens*), 234, head of larva (*pungens*):
Leices., Culic. Malaya, 157, ♂ ♀: Brun., Rec. Ind. Mus. I,
344; IV, 468.

Culex aestuans, Wied., Auss. Zweifl. I, 11 (1828).

Culex pallipes, Mg, Syst. Besch. VII, 1 (1838).

Culex anxifer, Coquerel, in Big., Ann. So. Ent. Fr. (1858).

Culex skusii, Giles, Handbk., 1st ed., 292 (1900).

Heteronychia dolosa, Arrib., Dipt. Argent., 56 (1891). [(1901).

Culex panalectoris, Giles, Jour. Bomb. N.H. Soc. XIII, 608

Armigeres panalectoros, *id.*, Handbk., etc., 2nd ed., 386.

Desvoidia panalectros, *id.*, t. Theob., Rec. Ind. Mus. IV, 5.

Desvoidia panalectros, Blanch., Moust., 266, ♂.

Blanchardiomyia panalectros, Brun., Rec. Ind. Mus. IV, 442.

Culex quasipipiens, Theob., Monog. Culic. II, 136, ♀, fig. 223,
head, wing, veins (1901): Giles, Handbk., 2nd ed., 438:
Blanch., Moust., 344, ♀: Brun., Rec. Ind. Mus. I, 351.

Culex fochowensis, Theob., Monog. Culic. II, 137, ♂ ♀, figs. 224,
225, var. chars. (1901): Blanch., Moust., 344, ♂ ♀ (*fouchow-*
ensis): Brun., Rec. Ind. Mus. I, 345.

Culex reesii, Theob., Monog. Culic. II, 145, ♂ ♀, fig. 232, palpus
♂, thorax ♀, claws ♂ (1901): Giles, Handbk., 2nd ed., 449,
♂ ♀: Blanch., Moust., 361, ♂ ♀: Brun., Rec. Ind. Mus. I,
351.

Culex sericeus, Theob., Monog. Culic. II, 147, ♀, fig. 233, var.
chars. (1901): Giles, Handbk., 2nd ed., 452, ♀: Blanch.,
Moust., 362, ♀: Brun., Rec. Ind. Mus. I, 352; IV, 475.

Culex christophersi, Theob., Monog. Culic. IV, 453, ♂ ♀ (1907).

Culex osakensis, Theob., Monog. Culic. IV, 439, ♂ only (1907).

Culex 5-fasciatus, Dyar and Knab (? nec Say), Proc. Ent. Soc. Wash. XI, 34 (1909).

Culex goughii, Theob., U.S. Afr. Dep. Agric., 1st Rep. Vet. Res. 268, ♂ only (1911).

subspecies *luteoannulatus* Theob., Monog. Culic. II, 159 (1901).

? *Culex pungens*, Wied., Auss. Zweifl. I, 9 (1828).

? *Culex macleayi*, Skuse, Pr. Linn. Soc. N.S. Wales, p. 1745 (1896).

The most widely distributed species in the Orient. Records of its occurrence during every month can be found amongst Indian localities. Ceylon (Kurmregalla, Badulla, Balangoda, Keleni Valley), i, iii, xi [Green]; Nepal; Moulmein; Manipur; at sea between Calcutta and Rangoon, 22-23-ii-08 [Annandale]; Rangoon, 24-xii-04 to 3-i-05, and 9-ii-06 [Brunetti]; Dindings [Wray]; Perak [Wright]; Penang, 3-8-viii-06 [Brunetti]; Singapore, 4-xi-99 and vii [Hanitsch, Biro]; 21-vi-06 [Brunetti]; Malay Peninsula, widely distributed and common near houses [Leicester]; Phrapatoon, Siam, 10-18-i; 19-iii; 30-xii-07; viii and xi-06 [Dr. Woolley]; Batavia, 27-vi; 9-vii-06 [Brunetti]; Soerabaya, 16-25-vii-06 [Brunetti]; Papua (Friedrich Wilhelmshafen and Stephansort), [Biro]; Sarawak; Hong Kong, 8-i-00; vii [Ford]; Foochow [Rennie]; Shaohyling [Cornford]; Shanghai, 16-25-vii-06: swarming in ditch; Hankow, 22-26-iv-06; Manila, 10-16-iii-06 [all three Brunetti]; Seistan; E. Persia [Annandale and Kemp]. It also occurs in Japan, East Africa, Natal, Zanzibar, Mombasa, the Soudan, Egypt, British Central Africa, Senegambia, Mauritius, the Seychelles, Australia, Fiji and the Southern part of N. America, Central and S. America, and the West Indies. Types of *fatigans*, *aestuans* and *pungens* in Wiedemann coll.; of *macleayi* in Sydney Museum; of *panalectros* in British Museum. No type specimen set up of *skusei*; a typical specimen in British Museum.

sitiens Wied. Auss. Zweifl. I, 542, ♀ (1828).

Edws., Bull. Ent. Res. IV, 232.

Culex impellens, Walk. (nec Theob.), Proc. Linn. So. Lond. IV, 91, ♀ (1860).

C. annulirostris, Skuse, Proc. Linn. So. N.S. Wales (2) III, 1737 (1889).

C. microannulatus, Theob., Monog. Culic. I, 353, ♂ ♀; pl. xviii, 69, ♀, full ins. col.; fig. 118 b, head, d, fore ungues ♂ (1901): Blanch., Moust., 292, ♂ ♀; Leices., Culic. Malaya, 140, ♂ ♀; Brun., Rec. Ind. Mus. I, 349; IV, 472.

C. gnophodes, Theob., Monog. Culic. III, 163, ♀ (1903).

Leices., Culic. Malaya, 145 (Theob., descr. copied).

C. somaliensis, Nev.-Lemaire, Arch. Parasit. X, 254 (1906).

C. ronaldi, Charm., Ann. Trop. Med. II, 259 (1908).

C. salus Theob., Third Rep. Wellcome Lab., 256 (1909).

Apparently a very widely distributed species throughout the East, common nearly everywhere.

Sylhet, i, ii, iv, v, xi, xii; Calcutta, July to November, common; N.W. Provinces, India, common [Giles]; Shahjahanpur, xii [Giles]; Mukerian; Hoshiarpur [Dr. Datta]; Purnea, Bengal, viii; between Bolpore and Rampore Haut, Bengal, viii; in railway carriage [Paiva]; Lake Chilka, Orissa, viii; Gopkuda I.; Trincomalee and Hakgala, Ceylon, 24-viii—, 9-ix-07, and Peradeniya [all three Green]; Singapore [Dr. Finlayson]; Dindings; Straits, xi; Makassar; Manila, Cavite and Pangasinan (all three Phil. Is.) [Banks, Stiff and Chamberlain respectively]. It also occurs in Mauritius. *Type* (*sitiens*), said to be in the Copenhagen Museum; that of *annulirostris* at Sydney Museum; of *microannulatus*, *impellens*, *salus* and *gnophodes* in the British Museum, of *ronaldi* at the Liverpool School of Tropical Medicine.

vagans Wied., Auss. Zweifl. I, 545, ♀ (1828).

Theob., Monog. Culic. I, 411, ♀; fig. 146, wing, scutellum: *id.*, loc. cit., IV, 14, ♂; V, 347, ♂; Giles, Handbk., etc., 279; 2nd ed., 414, ♀; pl. xvi, 14, venation ♀: *id.*, Jour. Trop. Med. VII, 368; Blanch., Moust., 304, ♀; Brun., Rec. Ind. Mus. I, 353; IV, 476.

Madras city, 31-x-08 [Hodgart]; Hongkong, x; Shanghai [Lindesay]; Pampanga [Whitmore]; Foochow [*t. Blanch.*].

Type in Dr. Trentepohl's collection.

nigripes Zett., Ins. Lapp., 807 (1838).

Ficalbi, Bull. So. Ent. Ital., 292 (1896): Theob., Monog. Culic. II, 93, ♂ ♀; fig. 194, wing, etc.; fig. 260 (p. 219), map of distribution: Brun., Rec. Ind. Mus. I, 350.

? *impiger*, Walk., List. Dipt. Brit. Mus. I, 6, ♀ (1848).

? *implacabilis*, *id.*, loc. cit., I, 7, ♀ (1848).

Deosai Plateau, between Kashmir and Shardo, 13,000 ft. [Dr. Neve]. A circumpolar species. *Type* (*nigripes*) in Paris Museum.

rubrithorax Macq., Dipt. Exot. Supp. IV, 9, ♀ (1850).

Theob., Monog. Culic. I, 416, ♀; fig. 150, head, thorax, apex of abdomen: Giles, Handbk., etc., 2nd ed., 412, ♀: *id.*, Jour. Trop. Med. VII, 368; Skuse, Proc. Linn. So. N.S. Wales, p. 1735 (1896): Brun., Rec. Ind. Mus. I, 352.

Pampanga, Ph. Is. [Whitmore]. *Type* in Paris Museum. Described originally from Australia.

aureostriatus Dol., Nat., Tijd. Ned. Ind. XIV, 385, ♀; pl. vi, 1 (1857).

Giles, Handbk., 280: Theob., Monog. Culic. I, 387, ♀; Blanch., Moust., 310, ♀; Brun., Rec. Ind. Mus. I, 355; IV, 467.

Amboina. Blanchard says it occurs in Tokio. *Type*, location unknown to present author.

longipalpis Wulp, Mid-Sum. Dipt., 9, pl. i, 3, head (1892).

Giles, Handbk., etc., 249; 2nd ed., 423, ♀: Blanch., Moust., 318, ♀: Brun., Rec. Ind. Mus. IV, 471.

Soceroelangoen, Sumatra. *Type*, location unknown to present author.

mimeticus Noe, Bull. Ent. So. Ital. XXXI, 240 (1899).

Giles, Handbk., etc., 202; 2nd ed., 389; pl. xv, 16, wing, ♀; 17, palpi and proboscis ♂; 18, claws ♂: Theob., Monog. Culic. I, 329, ♀; pl. xvi, 63, ♀, full ins. col.: Blanch., Moust. 271, ♂ ♀: Leices., Culic. Malaya, 139, ♂ ♀: Brun., Rec. Ind. Mus. I, 349; IV, 472.

? *hyrcanus*, Pallas, Reisen Russ. Reich. (1871), "near Caspian Sea."

Theog., 8000 ft., 2-v-07 [*Annandale*]; Punjab, 6000 ft., iii [*Lindesay*]; Canara District [*Aitken*]; Nilgiri Hills [*Giles*]; Lushai Hills, 1-vi-04 [*Macleod*]; Thaumaspur, Nepal, 18-20-ii-08; Malacca [*t. Blanch.*]; Peradeniya, 17-ix-07 [*Green*]; Kuala Lumpur [*Durham*]; Perak [*Wright*]. Occurs also in Italy, Cyprus, Macedonia and Palestine. See note under *C. mimulus*, Edw. *Type* in Rome University Museum.

tigripes Grandpré and Charmay, Planters Gazette Press (1900).¹

Giles, Handb., 2nd ed., 407, ♂ ♀; pl. xvi, 4, wing, ♂ ♀, head ♂, thorax var.: *id.*, Jour. Trop. Med. VII, 368: Theob., Monog. Culic. III, figs. 120-122, larva, pupa, diagrams: Brun., Rec. Ind. Mus. I, 352; IV, 475: Edw., Bull. Ent. Res. II, 261.

maculicrura, Theob., Monog. Culic. II, 34, ♂ ♀; pl. xxii, 85, ♀, full ins. col. (1901).

Sylhet, ii, iv, v to xii [*Lt.-Col. Hall*]; Kurseong, 4-viii-08; Damukdia Ghat (Riv. Ganges, E. Bengal), vii; Calcutta, viii-08; xii, 1905; Port Canning, xii; Mandalay, 11-iii-08 [*Annandale*]; Manipur, viii [*Gourlay*]; Kirindi, 20-xi-08; Weligama, 3-i-08; Dondra, 28-iv-08; Mandulsima, 14-xii-08 (all four, Ceylon); Dindings, Straits, xii; Pampanga, Ph. Is. [*Whitmore*]. Also occurs in West Africa, Natal, Mauritius, Queensland.

Type, location unknown to present author. Edwards has suggested that this species may be identical with *concolor*, R. Desv.

tritaeniorhynchus Giles, Jour. Bomb. Nat. Hist. So. XIII, 606 (provis. descr.) (May 1901).

Giles, Entom. XXXIV, 192 (July 1901): *id.*, Handbk., 2nd ed., 401, ♂ ♀: Theob., Monog. Culic. I, 364, ♂ ♀; fig. 124, wing, ♂ ♀: Blanch., Moust., 294, ♂ ♀: Brun., Rec. Ind. Mus. I, 355: Edw., Bull. Ent. Res. IV, 233; VII, 224.

Culex sitiens, Theob. (*nec* Wied.), Monog. Culic. I, 360, fig. 121, wing, proboscis (Nov. 1901): Giles, Handbk., 2nd ed., 400,

¹ Edwards (Bull. Ent. Res. II, 261) gives as reference "Grandpré, Les Moustiques, 1900." Presumably he refers to Blanchard's work.

- ♀: Blanch., Moust., 293, ♀: Leices., Culic. Malaya, 143, ♂ ♀: Brun., Rec. Ind. Mus. I, 352; IV, 475.
- Culex annulus*, Theob., Monog. Culic. I, 358, ♀ (1901).
Giles, Handbk., 2nd ed., 405, ♀: Blanch., Moust., 293, ♀: Leices., Culic. Malaya, 144 (copies Theob.'s descr. of ♀).
- Culex vishnui*, Theob., Monog. Culic. I, 355, ♂ only, (♀ = *vishnui*, Theob., *vera*), fig. 119, claws, wing tips; 120a, forms of abdomen (♂ only), claws, ♂ (1901): Giles, Handbk., 2nd ed., 399, ♂; pl. xvi, 5a, abdomen vars., 5b, fore tarsal claws, ♂: Theob., Monog. Culic. IV, 387, ♂ only; fig. 163, genitalia ♂: Blanch., Moust., 293, ♂ only: Brun., Rec. Ind. Mus. I, 354; IV, 476.
- Culex biroi*, Theob., Ann. Mus. Hung. III, 82, ♂ ♀; pl. i, wing, ♂ ♀: *Id.*, Monog. Culic. IV, 390, ♂ ♀; fig. 165, wing ♂; 164, wing ♀.
Sambalpur (Cent. Prov., India), 26-x-1900 [*Murphy*]; Calcutta, viii, ix; Bombay [*Biro*]; Madras, xi, xii [*Cornwall*]; Dacca; Travancore, type [*James*]; Quilon, 27-vii-1900 and ii [*James*]; Ceylon, xi; 27-xii-99 [*Bartholomew*]; Dindings (Straits), x, xi, xii; Malaysia; Malacca; Sumatra; Celebes [all three *t. Blanchard*]; Tai-Po; Pokfulam; Hongkong; Lamma, Philippines [*t. Ludlow*].
Types of tritaeniorhynchus and vishnui ♂ in British Museum; of *biroi* in Hungarian Museum; of *annulus* in Dr. Rees collection.
- vishnui* Theob., Monog. Culic. I, 355, ♀ only (the ♂ = *tritaeniorhynchus*, Giles); figs. 119, 120 var. chars. (♀ only); pl. xvii, 66, ♀, full ins. col. (1901).
Id., *loc. cit.*, IV, fig. 162, wing ♀; pl. v, wing scales, ♀: Blanch., Moust., 293 (♀ only): Leices., Culic. Malaya, 141 (♀ only): Brun., Rec. Ind. Mus. I, 354; IV, 576: Edw., Bull. Ent. Res. IV, 233; VII, 225.
- impellens*, Theob. (*nec* Walk.), Monog. Culic. I, 362, ♀; fig. 122, head; 123, wing; *loc. cit.*, IV, 15 (long list locs.): Giles, Handbk., 2nd ed., 399, ♀ only; pl. xvi, 3a head: Blanch., Moust., 294, ♀: Leices., Culic. Malaya, 142 (Theob.'s descr. copied): Brun., Rec. Ind. Mus. I, 347; IV, 470.
- Culex perplexus*, Leices., Culic. Malaya, 150, ♂ ♀ (1908): Brun., Rec. Ind. Mus. IV, 473.
- ? *C. microtaeniatus*, Theob., Tijds. v. Ent. LIV, 236 (1911).
? *C. pseudoinsula*, Theob., *loc. cit.*, p. 237 (1911).
? *C. parvus*, Taylor, Bull. N. Terr. Austral., 1a, p. 27 (1912).
- Very generally distributed through the East; widely throughout India, as far north as Nepal and Sylhet, i, ii [*Hall*]. Outside of India it is recorded as follows; Ceylon, xi and 27-xii-99 [*Bartholomew*]; Batticalora, Ceylon, iv, [*Green*]; Moulmein, ii; Mandalay, 11, 12-iii-08 [*Annandale*]; Chittagong, 26-vii-08 [*Lt.-Col. Hall*]; Rangoon, 24-xii-04 to 3-i-05 [*Brunetti*]; Kuala Lumpur, v [*Durham, Leicester*]; Perak [*Wray, Wright*]; Makassar [*t. Walker*]; Batavia, 27-vi-06 to 9-vii-06 [*Brunetti*]; Pampanga [*Whitmore*]. Also taken on board ship 5 miles off Aleppey, Mala-

- bar Coast, 4-v-08 [*Paiva*]. Type of *vishnui* ♀ and of *perplexus* ♂ ♀ in British Museum.
- annulioris** Theob., Liver. Sch. Trop. Med. Mem. V, App. (1901).
Id., Monog. Culic. I, 371, ♀ (1901): Brun., Rec. Ind. Mus. IV, 466: Edw., Bull. Ent. Res. II, 259.
 Phillipines. Described from a single ♀ from Mashonaland. Type in British Museum.
- hirsutum** Theob., Monog. Culic. I, 392, ♂ ♀; pl. xx, 80, ♀ full ins. col., fig. 137, palpus ♂, apex ♂ antenna (1901).
 Brun., Rec. Ind. Mus. I, 347.
 Phillipines. Types in British Museum.
- gelidus** Theob., Monog. Culic. II, 20, ♀; pl. xxiv, 93, ♀, full ins. col.; fig. 158, thorax and hind tarsus (*Culex*) (1901).
Culex gelidus, Giles, Handbk., 2nd ed., 421, ♀: Blanch., Moust., 316, ♀: Brun., Rec. Ind. Mus. I, 346.
Leucomyia gelida, Theob., Monog. Culic. IV, 372: Leices., Culic. Malaya, 147, ♂ ♀.
Theobaldiomyia gelidus (lapsus) Brun., Rec. Ind. Mus. IV, 462.
 Calcutta, August, 6-x-04 [*Brunetti*]; Purnea, Bengal, 6-viii-07, ix, x [*Paiva*]; Dacca; Rajshahi, Bengal, 1-6-ii-07; Madras town, 30-31-x-08 [*Hodgart*]; Peradeniya, vii, ix: Maskeliya, iv; Kelani Valley [both Ceylon and *Green*]; Travancore and Cochin States, xi-08 [both *Annandale*]; Selangor [*Butler*]; Rangoon, 25-ii-08; Dindings (Straits) in November; Perak; Sarawak, xi [*Butler*]; Bayembang, Phil. Is. [*Chamberlain*].
- var. **bipunctata** Theob., Monog. Culic. IV, 374, ♂ (*Leucomyia*) (1907).
 India [*Major Aldrich*]; Sarawak [*Barker*].
- var. **cuneata** Theob., Monog. Culic. II, 22, ♀, fig. 159, var. chars. (*Culex*) (1901).
Culex gelidus cuneatus, Giles, Jour. Trop. Med. VII, 368: Brun., Rec. Ind. Mus. I, 346.
 Sylhet, v [*Lt.-Col. Hall*]; Calcutta, July to December, common in houses, at light, in railway carriages and in the open and on plants, [*Annandale, Brunetti*]; Katihar; Bhogaon, ix, x; Purnea, viii, x [*Paiva*]; near Puri, 23-x-08, at light; Madras, 23 and 30-x-08; Travancore, 5 and 19-xi-08 [*Annandale*]; Malabar, 4-xi-08 [*Annandale*]; Quilon, vii [*James*]; Ceylon [*Green*]; Tai-pang, Perak [*Wray*]; Manila [*Banks*]; Pampang, Phil. Is. [*Whitmore*]; Pangasinan, Phil. Is.
 Type, location unknown to present author.
- univittatus** Theob., Monog. Culic. II, 29, ♂ ♀; fig. 161, head, leg, abdomen; pl. xxii, 86, ♀, full ins. col. (1901).
 Giles, Handbk., 2nd ed., 428, ♂ ♀ (*univittatus*, lapsus):
 Blanch., Moust., 321, ♀: Brun., Rec. Ind. Mus. I, 353.

Singapore, vii, ix [*Hainitsch*]. Occurs in Africa. Type in British Museum.

quasiunivittatus Theob., Monog. Culic. II, 32, ♀; fig. 164, head (1901).

Brun., Rec. Ind. Mus. I, 351.

Pampanga [*Whitmore*]. Described from single ♀ from Mashonaland. Type in British Museum.

hirsuteron Theob., Monog. Culic. II, 98; fig. 196, claws ♂; 197 wing, ♀ (1901).

Id., Gen. Ins., Fasc. 26, 27: Giles, Handbk., 2nd ed., 451 (*hirsuteros*): *id.*, Jour. Trop. Med. VII, 368 (*hirsuteros*): Brun., Rec. Ind. Mus. I, 347.

Pampanga [*t. Banks*]. Said to occur in Virginia, U.S.A., which is doubted by Banks. Type in British Museum.

tipuliformis Theob., Monog. Culic. II, 327, ♀; fig. 306, wing, leg, abdomen (1901).

Giles, Handbk., 2nd ed., 443, ♂: Blanch., Moust., 363, ♀: Brun., Rec. Ind. Mus. I, 353.

Bakloh (N.W. India), 5000 ft., iii [*Lindesay*]. Type in British Museum.

fragilis Ludlow, Jour. N.Yk. Ent. So. XI, 142 (1903).

Theob., Monog. Culic. IV, 424, ♂ ♀.

Oras (Samar, Phil. Is.), 6-viii [*t. Ludlow*]. Type in Army Medical Museum, Washington.

alis Theob., Monog. Culic. III, 167, ♂ ♀ (1903).

Philippines [*t. Ludlow*]. Also occurs on Christmas Island. Type in British Museum.

sinensis Theob., Monog. Culic. III, 180, ♀ (*C. gelidus*, var. *sinensis*) (1903).

Culex sinensis, Edw., Bull. Ent. Res. IV, 231.

Leucomyia sinensis, Theob., Monog. Culic. V, 313 (1910): *id.*, Rec. Ind. Mus. IV, 20.

Theobaldiomyia sinensis, Brun., Rec. Ind. Mus. IV, 463.

Culex sepositus, Leices., Culic. Malaya, 152, ♀ (1908).

Taeniorhynchus tenax, Leices. (*nec* Theob.), Culic. Malaya, 167, ♀ (1908).

Balighai, near Puri, 23-x-08, at light [*Annandale*]; Kuala Lumpur [*Leicester*]; Shaohyling, China [*Cornford*]. Types of *sinensis* and *tenax* in British Museum; that of *sepositus*, lost.

halifaxi Theob., Monog. Culic. III, 231, ♀ (1903).

Id., loc. cit., V, 350: Brun., Rec. Ind. Mus. IV, 470: Edw., Bull. Ent. Res. IV, 234.

Culex multimaculosus, Leices., Culic. Malaya, 155, ♂ ♀ (1908).

Brun., Rec. Ind. Mus. IV, 473.

Dindings, Strait Settlements, xii. Type of *halifaxi*, a unique ♀, in British Museum; of *multimaculosus*, in Leicester coll.

whitmorei Giles, Jour. Trop. Med. VII, 367 (*Taeniorhynchus*) (1904).

Culex whitmorei, Edw., Bull. Ent. Res. IV, 232; VII, 225.

Taeniorhynchus whitmorei, Theob., Monog. Culic. V, 431; Brun., Rec. Ind. Mus. IV, 480.

Taeniorhynchus argenteus, Ludlow, Can. Ent. XXXVII, 98, ♀ (1905): Theob., Monog. Culic. IV, 487, ♀ (copies Miss Ludlow's descr.): *loc. cit.*, V, 426, fig. 191, wing.

Theobaldiomyia argentea, Brun., Rec. Ind. Mus. IV, 462.

Leucomyia plegipennis, Theob., Monog. Culic. IV, 375 (1907).

Culex albus, Leices., Culic. Malaya, 148, ♀ (1908).

Culex loricatus, Leices., *loc. cit.*, 151, ♀ (1908).

Angeles, Pampanga, ix [*Whitmore*]; Kuala Lumpur. *Types* of *whitmorei*, *plegipennis* and *loricatus*, in British Museum; of *argenteus* in Army Medical Museum, Washington; of *albus* in Leicester collection.

trimaculatus Theob., Ann. Mus. Hung. III, 86, ♀; fig. 5, thoracic marks (1905).

Id., Monog. Culic. IV, 427, ♀; fig. 187, thorax, ♀.

Bombay [*Biro*]. *Type*, a unique ♀, in Hungarian Museum.

annulatus Theob., Ann. Mus. Hung. III, 98, ♂, fig. 10, proboscis apex, scales (*Trichopronomyia*) (1905).

Id., Monog. Culic. IV, 479, ♂; fig. 213, var. chars.; 214, wing, ♂.

Friedrich Wilhelmshafen, Papua [*Biro*]. *Type* in Hungariau Museum.

ornatus Theob., Ann. Mus. Hung. III, 100, ♀ (*Melanoconion*) (1905).

Id., Monog. Culic. IV, 508, ♀; fig. 231, wing, ♀.

Friedrich Wilhelmshafen, Papua, xii, 1900 [*Biro*]. *Type* in Hungarian Museum, a unique specimen.

pallidiceps Theob., Ann. Mus. Hung. III, 101, ♂ (*Melanoconion*) (1905).

Id., Monog. Culic. IV, 509, ♂.

Friedrich Wilhelmshafen, Papua, xii-1900 [*Biro*]. *Type* in Hungarian Museum.

lutea Ludl., Can. Ent. XXXVII, 96, ♂ (*Popea*) (1905).

Id., *loc. cit.*, XLII, 193, ♀ (1910).

Pampanga, Phil. Is. [*Whitmore*]. *Type* in Army Medical Museum, Washington.

macropus Blanch., Comp. Rend. LIII, 1045, nom. nov. for *C. longipes*, Theob., preocc. (1905).

Culex macropus, Blanch., Moust., 327.

Culex longipes, Theob., Monog. Culic. II, 68, ♀ (1901): Giles, Handbk., 2nd ed., 468, ♀.

Singapore, 4-ix-99, *type* [*Hanitsch*], and 27-vii. *Type* of *longipes* in British Museum.

- annuliferus** Ludl., Jour. N.Yk. Ent. Soc. XI, 141 (*annulifera*) (1903).
Theob., Monog. Culic. IV, 388, ♂ ♀: Brun., Rec. Ind. Mus. I, 343.
Culex ludlowi, Blanch., Moust., 630, nom. nov. for *annuliferus*, Ludl. preocc. Blanch., 1852: Brun., Rec. Ind. Mus. IV, 471.
Bayembang. Phil. Is. [*Chamberlain*]; Mangarin and Dagupan, ii to iv (both Phil. Is.). Type in Army Medical Museum, Washington.
- rizali** Banks, Phil. Jour. Sci. I, 999, ♀ (1906).
Theob., Monog. Culic. V, 391: Brun., Rec. Ind. Mus. IV, 474.
Siya Siya Mt. of Canlaon Volcano, Negros I. (Phil. Is.), vi [*Banks*]. Type in Entomological collection (No. 6083), Bureau of Science, Manila.
- fuscocephalus** Theob., Monog. Culic. IV, 420 ♀ (1907).
Edw., Bull. Ent. Res. IV, 234; VII, 225.
C. taytayensis, Banks, Phil. Jour. Sci. IV, 545, ♂ ♀ (1910).
Brun., Rec. Ind. Mus. IV, 475.
C. luteola, Theob., Monog. Culic. V, 378, ♀ (1910).
Brun., Rec. Ind. Mus. IV, 471.
C. uniformis, Leices., Culic. Malaya, 159, ♀ (1908): Brun., Rec. Ind. Mus. IV, 476.
C. minor, Theob., Rec. Ind. Mus. II, 298, ♂ ♀ (1901): Theob., loc. cit., IV, pl. i, wing: *id.*, Monog. Culic. V, 363, ♂ ♀; fig. 150, wing: Brun., Rec. Ind. Mus. IV, 473.
Sylhet [*Hall*]; Calcutta, xii [*Annandale*]; Pallode, Travancore, 15-xi-08 [*Annandale*]; Peradeniya, iii, ix [*Green*], x, 1900; Hakgala, Ceylon, 5-xi-07 [*Green*]; Lungleh, Lushai Hills, vi, vii [*Macleod*]; Rizal, Phil. Is. [*Banks*]; Batu Gajah, Malaysia. Type of *fuscocephalus* and of *luteola*, in British Museum; of *taytayensis* in Bureau of Science, Manila, Cat. No. II,459; of *minor* in Indian Museum; that of *uniformis*, probably lost, not in Leicester collection.
- nigricephalus** Leices., Culic. Malaya, 149, ♂ ♀ (*nigricephala*) (1908).
Brun., Rec. Ind. Mus. IV, 473.
Batu Gajah (Malaysia), [*Dr. Daniels*]. Type probably lost, not in Leicester collection.
- auratus** Leices., loc. cit., 153, ♀ (1908).
Brun., Rec. Ind. Mus. IV, 467.
Malayasia. Type probably lost, not in Leicester's collection.
- graminis** Leices., Culic. Malaya, 158, ♂ ♀ (1908).
Brun., Rec. Ind. Mus. IV, 470.
Malaysia. Type probably lost, not in Leicester's collection.
- minimus** Leices., Culic. Malaya, 160, ♂ ♀ (1908).
Brun., Rec. Ind. Mus. IV, 472.
Kuala Lumpur. Type in Leicester collection.

- pettigrewii** Theob., Rec. Ind. Mus. IV, 15, ♀ (1910).
Theob., Monog. Culic. V, 351, ♀.
Ukhrul, Manipur, viii-08 [*Pettigrew*].
Type, a unique ♀, in Indian Museum.
- japonicus**, var. **ceylonicus** Theob., Monog. Culic. V, 391 (*ceylonica*) (1910).
Brun., Rec. Ind. Mus. I, 348; IV, 471.
[*C. japonicus* typical form], Theob., Monog. Culic. I, 385, ♀ (1901), and III, 158 (1903).
Peradeniya, vi and x [*Green*]; a single ♀. *Types* of *japonicus* (typical form), and of *japonicus*, var. *ceylonicus* (presumably) in British Museum. The typical form is apparently not found in the East.
- aureopunctis** Ludlow, Can. Ent. XLII, 195, ♀ (1910).
Cottabato, Mindanao.
Type in Army Medical Museum, Washington.
- argentinotus** Banks, Phil. Jour. Sci. IV, 547, ♂ ♀ (1910).
Rizal, Phil. Is., 13-16-v-09 [*Banks*]. *Types* ♂ ♀ in Entomological collection, Cat. No. 11,460, Bureau of Science, Manila.
- palawanensis** Ludl., Psy. XXI, 30 (*Popea*) (1914).
Palawan, Phil. Is.
Type, location unknown to present author.
- mimulus** Edw., Bull. Ent. Res. V, 284 (1914-1915).
Malay States; India; Ceylon. Edwards thinks that probably all specimens hitherto recorded from the Orient as *mimeticus*, Noe, are this species, and those in the British Museum from Hong Kong, North India, Palestine and Cyprus are true *mimeticus*.
- nilgircicus** Edw., Bull. Ent. Res. VI, 358, ♂ ♀, fig. 1, ♂ genit. pt. (1915-16).
Utakamand (? Ootacamund), Nilgiri Hills, Madras Presid., 7500 ft., 24-31-xii-13 [*T. B. Fletcher*].
Types in British Museum, presented by Agric. Instit., Pusa.

Culex, spp. *unrecognisable*.

- molestus** W., Dipt. Exot. I, 39 (1821).
Id., Auss. Zweifl. I, 542: Giles, Handbk., 341, 2nd ed., 470.
? *C. filipes*, Walk.
Sumatra. *Type* (defective) in Dr. Trentepohl's collection (Vienna Museum).
- luridus** Dol., Nat. Tijds. Ned. Ind. XIV, 384; pl. v, 1 (1857).
Giles, Handbk., 341, 2nd ed., 469: Brun., Rec. Ind. Mus. I, 355.
? *inflictus*, Theob., Monog. Culic. II, 115 (1901).
Gombong, Mid-Java [*t. Doleschall*].
Type, location unknown to present author.

- setulosus** Dol., Nat. Tijd. Ned. Ind. XIV, 384: pl. v, 4 (1857).
Giles, Handbk., 335, 2nd ed., 470: Brun., Rec. Ind. Mus. I, 355.
Mid-Java [*t. Doleschall*]. Type in Vienna Museum.
- filipes** Walk., Proc. Linn. So. Lond. V, 229, ♀ (1861).
Giles, Handbk., 343.
? = *molestus*, W.
Dorey, Papua. Type in British Museum, unrecognisable.
- ventralis** Walk. II, Proc. Linn. So. Lond. VIII, 103, ♀ (1865).
Brun., Rec. Ind. Mus. I, 356.
Papua [*t. Walker*]. The second species of this name by Walker in *Culex*; the first is synonymous with *Armigeres obturbans*, Walk. Type in British Museum.
- doleschalli** Giles, Handbk., 1st ed., 338 (1900): *nom. nov.* for *cingulatus*, Dol.
Brun., Rec. Ind. Mus. I, 354.
Culex cingulatus, Dol., Nat. Tijd. Ned. Ind. X, 405; pl. vii, 2; prooc., Fab., 1805 for a Brazilian species.
Ambarawa, Java, common all the year round.
Type, location unknown to present author.

Micraedes, Coq.

- Proc. Ent. Soc. Wash. VII, 185 (1906).
GENOTYPE, *M. bisulcatus* Coq., by original designation.
- Acalleoymia*, Leices., Culic. Malaya, 194 (1908).
GENOTYPE, *A. obscura* Leices., by original designation.
- malayi** Leices., Culic. Malaya, 184, ♀ (*Aedes*) (1908).
Brun., Rec. Ind. Mus. IV, 490.
Micraedes malayi, Edw., Bull. Ent. Res. VII, 228.
Aioretomyia aedes, Leices., Culic. Malaya, 189 (1908).
Kuala Lumpur. Type in Leicester coll.
- obscurus** Leices., Culic. Malaya, 194, ♂ ♀ (*Acalleoymia obscurus*) (1908).
Micraedes obscurus, Edw., Bull. Ent. Res. IV, 237.
Malaysia. Type in Leicester coll.

Cyathomyia, de Meij.

- Ann. Jard. Bot. Buitenzorg, Ser. 2, Supp. III, 921 (1910).
GENOTYPE, *C. jenseni* Meij., by original designation.
- brevipalpis** Giles, Handb., 2nd ed., 384, ♂ ♀; pl. xiv, 17—19, wings; 20, head ♂ (*Stegomyia*) (1902).
Culex brevipalpis, Theob., Monog. Culic. III, 146, ♂ ♀; V, 607, App.

- Stegomyia brevipalpis*, Blanch., Moust., 264, ♂ ♀: Brun., Rec. Ind. Mus. I, 329; IV, 445.
Cyathomyia brevipalpis, Edw., Bull. Ent. Res. IV, 237.
Melanoconion uniformis, Leices., Culic. Malaya, 136, ♂ ♀: Brun., Rec. Ind. Mus. IV, 484.
 Shahjahanpore (N.W. India). *Types of brevipalpis* in British Museum; of *uniformis* in Leicester coll.

Theobaldia, Nev. Lemaire.

Comp. Rend. Soc. Biol. Paris LIV, 1331 (1902).

Theob., Monog. Culic. III, 148; pl. x, wing scales, var. spp.; IV, 275, note on name; V, 270, tab. spp.: *id.*, Gen., Ins. Fasc., 26, 23; Edw., Bull. Ent. Res. II, 251; Brun., Rec. Ind. Mus. I, 339; IV, 459, note; X, 72.

GENOTYPE, *Culex annulatus* Schrk.

Theobaldinella, Blanch., Moust., 390, nom. nov. (generic chars. given), on assumption of pre-occupation in Mollusca by *Theobaldius*, Nevike: Brun., Rec. Ind. Mus. X, 72.

Culiseta, Felt. N.Yk. State Mus. Bull., 79, App., 391e (1904).

GENOTYPE, *Culex absobrinus* Felt., by original designation.

Pseudotheobaldia, Theob., Monog. Culic. IV, 271 (1907): Brun., Rec. Ind. Mus. IV, 478; X, 70.

GENOTYPE, *P. niveitaeniata* Theob., Monog. Culic. IV, 271, sp. nov.

annulata Schrk., Beitr. z. Naturg., 97 (*Culex*) (1776).

Culex annulatus, F., Ent. Syst. IV, 400; Meig. Syst. Besch. I, 3; Macq., Hist. Nat. Dipt. I, 35; Sch., F. Austr. II, 626; Zett., Dipt. Scand. IX, 3640; Wulp., Dipt. Neerl., 325; Theob., Monog. Culic. I, 331, ♂ ♀, fig. 108, abd. segments, claws, ♂ ♀, palpus, ♂; pl. xv, 58, full ins. col.: Giles, Handbk., etc., 2nd ed., 391; pl. xv, abd. segments, claws, ♂ ♀, wing ♀, head ♂, genitalia ♂: Blanch., Moust., 280, ♂ ♀: fig. 206, claws and genitalia, 207 adult larva.

Theobaldia annulata, Theob., Gen. Ins. Fasc. 26; pl. i, 12, ♀ full ins. col.: Brun., Rec. Ind. Mus. I, 339; IV, 459.

Culex variegatus, Schrk., Enum. Ins. Austr., 482 (1781).

Culex affinis, Steph., Zool. Jour. No. IV (1825).

Bakloh, Punjab, 5000 ft., xi [*Lindesay*]; also recorded from Brianon, France, 4000 ft., and from Durango State, Mexico, 8000 ft. *Type*, Prof. Bezzi informs me the *type* of *annulata*, Schrk., is no longer in existence. That of *affinis* is in the Hope collection at Oxford University, that of *variegatus* probably lost or unrecognizable.

glaphyoptera Sch., F. Austr. II, 628 (*Culex*) (1864).

Culex glaphyopterus, Giles, Handbk., 216 (Sch.'s descr. copied).

Theobaldia glaphyoptera, Edw., Bull. Ent. Res. IV, 237.

Pseudotheobaldia niveitaeniata, Theob., Monog. Culic. IV, 272, fig. 87, ♂ genitalia, 88, wing, ♂ (1907): Brun., Rec. Ind. Mus. IV, 478.

Dehra Dun, ii, iii [Thompson]; Theog., Simla Distr., 2-v-07 [Annandale]. Type of *glaphyoptera*, location unknown to me; that of *niveitaeniata* in British Museum.

spathipalpis Rond., Bull. Soc. Ent. Ital. IV, 31 (*Culex*) (1872).

Culex spathipalpis, Theob., Monog. Culic. I, 339, ♂ ♀; III, 154, ♀; pl. x, wing scales; IV, 276 (larva descr.): Giles, Handbk., etc., 2nd ed., 392, ♂ ♀; pl. xv, 23, wing ♀, 24, head ♂, 25, genitalia, ♂: Ficalbi, Bull. Soc. Ent. Italy XXI, 86; XXVIII, 242; pl. ii, 8, 9; pl. iii, 7, 12; pl. iv, 18, 21, 22; pl. v, 31, 34; XXXI, 194, figs. 72-75, 146: Blanch., Moust., 283, ♂ ♀; fig. 209, claws and genitalia.

Theobaldia spathipalpis, Brun., Rec. Ind. Mus. I, 340; IV, 459, notes.

Culex serratipes, Beck., Mitt. Zool. Mus. Berl. IV, 78 (1908).

Naini Tal, 7000 ft. Occurs also in Italy, Palestine, Cyprus, Madeira, Teneriffe, Algeria, Gibraltar, Malta, the Transvaal, and Cape Colony. Type in Florence Museum.

In Edwards' latest paper he suggests sinking this species as synonymous with (*Culex*) *longiareolatus*, Macq., but a comparison with the type does not appear to have been made and until such is the case the present name may as well stand. Mr. James Waterston records the species under Macquart's name as common in Macedonia.

pagei Ludl., Psyche XVIII, 128, ♀ (1911).

Theob., Monog. Culic. V, 618.

Fort Pikit, Mindanao (Phil. Is.).

Type (presumably) in Army Medical Museum, Washington.

Ficalbia, Theob.

Monog. Culic. III, 296 (1903).

Theob., *loc. cit.*, IV, 577, tab. spp.; V, 541, tab. spp.: Gen. Ins., Fasc. 26, 36: Blanch., Moust., 418: Leices., Culic. Malaya, 228: Edwards, Bull. Ent. Res. IV, 237: Brun., Rec. Ind. Mus. I, 362; IV, 494; X, 60.

GENOTYPE, *Uranotaenia minima* Theob., by Edwards' designation.

Etorleptiomyia, Theob., 1st Rep. Wellc. Labor., 71 (1904).

Id., Gen. Ins., Fasc. 26, 44 (1905): *id.*, Monog. Culic. IV, 505: Leices., Culic. Malaya, 178: Brun., Rec. Ind. Mus. I, 368; IV, 484; X, 60: Blanch., Moust., 632.

- GENOTYPE, *E. mediolineata* Theob., by original designation.
- O'Reillia*, Ludl., Can. Ent. XXXVII, 101 (1905).
- GENOTYPE, *O. luzonensis* Ludl., ♀, by original designation.
- Ludlowia*, Theob., Monog. Culic. IV, 193 (1907).
- Id.*, loc. cit., V, 190, tab. spp.: Brun., Rec. Ind. Mus. IV, 493; X, 64.
- Edw., Bull. Ent. Res. II, 244.
- GENOTYPE, *Mimomyia chamberlaini* Ludl., by original designation.
- Megaculex*, Theob., Monog. Culic. IV, 282 (1907).
- GENOTYPE, *Culex albitarsis* Theob., by original designation.
- Boycia*, Newstead, Ann. Trop. Med. and Parasit. I, 33 (1907):
- GENOTYPE, *B. mimomyiaformis*, *id.*, sp. nov., loc. cit.
- Radioculex*, Theob., Rec. Ind. Mus. II, 295 (1908).
- Theob., Monog. Culic. V, 192: Brun., Rec. Ind. Mus. IV, 465; X, 70.
- GENOTYPE, *R. clavipalpus*, *id.*, sp. nov., loc. cit.
- Dasymyia*, Leices., Culic. Malaya, 102 (1908).
- Brun., Rec. Ind. Mus. IV, 453; X, 59.
- GENOTYPE, *D. fusca*: *id.*, sp. nov., loc. cit.
- Conopomyia*, *id.*, loc. cit., 113 (1908).
- Brun., Rec. Ind. Mus. IV, 453; X, 58.
- GENOTYPE, *C. metallica* Leices., by my own designation.
- Hispidimyia*, Theob., Monog. Culic. V, 245 (1910).
- GENOTYPE, *H. hispida*, *id.*, sp. nov., loc. cit.
- Ingramia*, Edw., Bull. Ent. Res. III, 43 (1912).
- GENOTYPE, *Mimomyia malfeyti* Newstead, by original designation.
- minima** Theob., Monog. Culic. II, 262, ♂, fig. 281, wing, costal border, wing scales (*Uranotaenia*) (1901).
- Uranotaenia minima*, Giles, Handb., 2nd ed., 488, ♂.
- Ficalbia minima*, Blanch., Moust., 418, ♂: Brun., Rec. Ind. Mus. IV, 494: Edw., Bull. Ent. Res. IV, 238.
- Mimomyia minuta*, Theob., Rec. Ind. Mus. II, 301, ♂ (1908).
- Id.*, loc. cit., IV, 30, ♀; pl. i, wing; pl. iii, wing scales: *id.*, loc. cit., V, 531, ♂ ♀, fig. 226, wing.
- Quilon, 7-iii-00 and ii [James]; Sylhet, 27-xi-04 [Hall]; Calcutta, 30-vii and 3 or 4-viii-07 [Annandale].
- Types of *minima* and *minuta* in British Museum.
- simplex** Theob., Monog. Culic. III, 297, ♂ (1903).
- Id.*, loc. cit., V, 541, ♀, fig. 235, wing: Blanch., Moust., 418, ♂.
- Aedes simplex*, Edw., Bull. Ent. Res. IV, 238.

- Skusea simplex*, *id.*, *loc. cit.*, VII, 224, fig. 9b, 9c, ♂ genitalia.
Kurumegalla, Ceylon, ix [Green]; Trincomalee, xi, 1906 [Green].
Type in British Museum.
- chamberlaini** Ludl., Can. Ent. XXXVI, 297, ♂ (*Mimomyia*) (1904).
Ludlowia chamberlaini, Theob., Monog. Culic. IV, 194, ♂ (Miss Ludlow's descr. copied): Brun., Rec. Ind. Mus. IV, 493.
Bayembang, Phil. Is., v [Chamberlain].
Type in British Museum.
- luzonensis** Ludl., Can. Ent. XXXVII, 101 (*O'Reillia*) (1905).
Etorleptomyia luzonensis, Ludl., Can. Ent. XXXVIII, 185.
Theob., Monog. Culic. IV, 506, ♀.
Ficalbia luzonensis, Edw., Bull. Ent. Res. IV, 238.
Etorleptomyia completiva, Leices., Culic. Malaya, 178, ♂ (1908).
Brun., Rec. Ind. Mus. IV, 484.
Bayembang (Pangasinan, Ph. Is.) [Chamberlain]; Singapore [Finlayson]. Type of *luzonensis* in Army Medical Museum, Washington; of *completiva* in Leicester's collection.
- clavipalpus** Theob., Rec. Ind. Mus. II, 295, ♂ ♀ (*Radioculex*) (1908).
Theob., *loc. cit.*, IV; pl. i, wing; pl. ii, head: *id.*, Monog. Culic. V, 193, ♂ ♀: Brun., Rec. Ind. Mus. IV, 465.
Calcutta, vii, xi, xii; Berhampur, Bengal, 1-i-08 [Lloyd];
Katihar, 4-5-x-08 [Paiva]; Vaikam coast of Travancore, 5-xi-08 [Annandale]; Rangoon, 25-ii-08 [Annandale]
Type in Indian Museum, *cotypes* in British Museum.
- fusca** Leices., *loc. cit.*, 102, ♂ ♀ (*Dasymyia*) (1908).
Brun., Rec. Ind. Mus. IV, 453.
Kuala Lumpur; Bukit Kutu (Malaysia).
Type in Leicester collection.
- metallica** Leices., Culic. Malaya, 113, ♂ ♀ (*Conopomyia*) (1908).
Kuala Lumpur and neighbourhood [Leicester].
Type in Leicester collection.
- hybrida** Leices., Culic. Malaya, 115, ♂ ♀ (1908).
Brun., Rec. Ind. Mus. IV, 453.
Kuala Lumpur [Leicester]; Singapore [Dr. Finlayson].
Type in Leicester collection.
- aurea** Leices., Culic. Malaya, 116, ♂ ♀ (1908).
Kuala Lumpur and Raub, Malaysia.
Type in Leicester collection.
- longirostris** Leices., Culic. Malaya, 228, ♂ ♀ (1908).
Skusea longirostris, Edw., Bull. Ent. Res. VII, 224.
Brun., Rec. Ind. Mus. IV, 494.
Kuala Klang, January. Type in Leicester collection.
- tenax** de Meij., Ann. Jard. Bot. Buitenzorg, Supp. 3, p. 928 (1910).
Java. Type in Amsterdam Museum.

dofleini Guenther, Zeits. wiss. Insek. IX, 204 (1914).

Ceylon. *Type*, location unknown to present author.

ludlowi *nom. nov.* for *minima*, Ludl., *preocc.*

Mimomyia minima, Ludl., Can. Ent. XXXIX, 413, ♂ ♀ (1907).

Ludlowia minima, Theob., Monog. Culic. V, 191, ♂ ♀ (Miss L.'s descr. copied): Brun., Rec. Ind. Mus. IV, 493.

Uranotaenia, Arrib.

Revista Mus. La Plata I, 405 (1891).

Theob., Monog. Culic. II, 241, fig. head, 242, map of distribution, 243; tab. spp.; pl. D, wing scales; V, 498, tab. spp.: *Id.*, Gen. Ins., Fasc. 26, 36: Blanch., Moust., 406, ♂ ♀: Giles, Handbk., 2nd ed., 485, 487, tab. spp.: Leices., Culic. Malaya, 203, tab. Malayan spp.: Brun., Rec. Ind. Mus. I, 363; IV, 495; X, 73: Alcock, Ent. Med. Off. 104: How., Dyar and Knab, Mosq. N. Amer., etc., IV, 898.

GENOTYPE, *U. pulcherrima* Arrib., by designation of Neveu-Lemaire, 1902.

Pseudouranotaenia, Theob., Jour. Econ. Biol. I, 33 (1905).

Theob., Monog. Culic. IV, 566, fig. 262 (p. 567), wing, ♂: Brun., Rec. Ind. Mus. IV, 494; X, 70. GENOTYPE, *P. rowlandii* Theob., by original designation.

Anisocheleomyia, Theob., Entom. XXXVIII, 52 (1905).

Theob., Monog. Culic. IV, 570: Brun., Rec. Ind. Mus. I, 363; IV, 493; X, 55.

GENOTYPE, *A. nivipes* Theob., by my own designation, 1914.

caeruleocephala Theob., Monog. Culic. II, 256; fig. 276, thorax, scutellum, head scales (1901).

Brun., Rec. Ind. Mus. I, 363.

Cottabatto (Mindanao, Phil. Is.) [*Vedder*]. Also occurs in Gambia, the Sudan and old Calabar.

Type in British Museum.

nivipes Theob., Entom. XXXVIII, 53 (*Anisocheleomyia*) (1905).

Uranotaenia nivipes, Edw., Bull. Ent. Res. IV, 238.

Uranotaenia nivea, Leices., Culic. Malaya, 211, ♂ (1908): Brun., Rec. Ind. Mus. IV, 497.

Pseudouranotaenia triangulata, Ludl., Can. Ent. XL, 331, ♂ (1908): Theob., Monog. Culic. V, 525 (Miss Ludlow's descr. copied).

Kuala Lumpur; Reine Regente (Mindanao, Phil. Is.), ii
Type, *nivipes* in British Museum; *triangulata* in Army Medical Museum, Washington.

alboannulata Theob., Entom. XXXVIII, 54 (*Anisocheleomyia*) (1905).

Uranotaenia alboannulata, Edw., Bull. Ent. Res. IV, 238.

Anisocheleomyia alboannulata, Theob., Monog. Culic. IV, 573, ♂; fig. 267, wing, ♂; pl. vii, wing scales, ♀: Brun., Rec. Ind. Mus. IV, 493.

trilineata, Leices., Culic. Malaya, 204, ♂ ♀ (1908): Brun., Rec. Ind. Mus. IV, 498.

India [*James*]; Malaysia [*t. Leices.*]. *Types of alboannulata and trilineata* in British Museum.

testacea Theob., Ann. Mus. Hung., III, 113, ♀; fig. 14, basal segments of antennae; pl. ii, wing; pl. iii, wing scales (1905).

Id., Monog. Culic. IV, 560, ♀; fig. 258, wing ♀: Edw., Bull. Ent. Res. IV, 239.

unilineata, Leices., Culic. Malaya, 220, ♂ ♀ (1908): Brun., Rec. Ind. Mus. IV, 498.

falcipes, Banks, Phil. Jour. Sci. I, 1004, ♂ ♀ (1906).

Singapore, 1902 [*Biro*]; Base of Dawna Hills, 4-iii-08 [*Annan-dale*]; Rizal; Manila, ii [*Banks, Schultze*]; Malaysia.

Type of testacea in British Museum, of *unilineata* in Leicester collection, of *falcipes* in Entom. coll. No. 5210, Bureau of Science, Manila.

atra Theob., Ann. Mus. Hung. III, 114, ♀ (1905).

Id., Monog. Culic. IV, 563, ♀: Edwards, Bull. Ent. Res. IV, 238.

ceylonica, Theob., Monog. Culic. V, 503, ♀; fig. 213, wing (1910).

Muina, Papua [*Biro*]; Galle, Ceylon, 10-vi-07 [*T. B. Fletcher*].

Type of atra in Hungarian Museum; of *ceylonica* in British Museum.

albitarsis Ludl., Can. Ent. XXXVII, 131, ♀ (*Anisocheleomyia*) (1905).

Theob., Monog. Culic. IV, 576 (Miss L.'s descr. copied): Brun., Rec. Ind. Mus. I, 363; IV, 494.

Camp Stotsenberg, Angeles, Phil. Is. [*Ludlow*].

Type in Army Medical Museum, Washington.

lateralis Ludl., Can. Ent. XXXVII, 385, ♀ (*caeruleocephala*, Theob., var.) (1905).

Id., Mosq. Phil. Is., 10 (*caeruleocephala*, var.).

Cottabatto, Mindanao, Philippines.

Type in Army Med. Mus., Washington.

Specific validity doubtful, as Miss Ludlow says, if Theobald's *type* was a rubbed specimen it became her var. *lateralis*; Theobald considers the species valid.

micans Leices., Culic. Malaya, 206, ♂ ♀ (1908).

Brun., Rec. Ind. Mus. IV, 497.

Malaysia. *Types* in Leicester's collection.

bimaculiala Leices., Culic. Malaya, 208, ♂ ♀ (1908).

Brun., Rec. Ind. Mus. IV, 495.

Kuala Lumpur; Raub. *Types* in Leicester's collection.

- unimaculiala** Leices., *Culic. Malaya*, 210, ♀ (1908).
 Brun., *Rec. Ind. Mus.* IV, 498.
 Malaysia. *Type* in Leicester's collection.
 Erroneously recorded by me as *unimaculata* (*Cat. Or. Culic. Supp.*).
- campestris** Leices., *Culic. Malaya*, 213, ♂ ♀ (1908).
 Brun., *Rec. Ind. Mus.* IV, 495.
 Malaysia. *Types* in Leicester's collection.
- argyrotarsis** Leices., *Culic. Malaya*, 214, ♂ ♀ (1908).
 Brun., *Rec. Ind. Mus.* IV, 495.
 Kuala Lumpur. *Types* in Leicester's collection.
- cancer** Leices., *Culic. Malaya*, 215, ♂ ♀ (1908).
 Brun., *Rec. Ind. Mus.* IV, 496.
 Port Swettenham; Klang. *Types* in Leicester's collection.
- longirostris** Leices., *Culic. Malaya*, 217, ♂ (1908).
 Brun., *Rec. Ind. Mus.* IV, 496.
 Malaysia. *Type*, apparently unique, in Leicester's collection.
- modesta** Leices., *Culic. Malaya*, 218, ♂ ♀ (1908).
 Brun., *Rec. Ind. Mus.* 497.
 Ampang Jungle, Malaysia. *Types* in Leicester's collection.
- nivipleura** Leices., *Culic. Malaya*, 219, ♂ ♀ (1908).
 Brun., *Rec. Ind. Mus.* IV, 497.
 Selangor. *Types* in Leicester's collection.
- maxima** Leices., *Culic. Malaya*, 221, ♀ (1908).
 Brun., *Rec. Ind. Mus.* IV, 497.
 Selangor. *Type* in Leicester's collection.
- lutescens** Leices., *Culic. Malaya*, 222, ♂ ♀ (1908).
 Brun., *Rec. Ind. Mus.* IV, 496.
 Kuala Lumpur.
- maculipleura** Leices., *Culic. Malaya*, 223, ♀ (1908).
 Kuala Lumpur. *Type*, unique, in Leicester's collection.
- bicolor**, *id., loc. cit.*, 225, ♂ ♀ (1908).
 Brun., *Rec. Ind. Mus.* IV, 495.
 Kuala Lumpur. *Types* in Leicester's collection.
- bimaculata** Leices., *Culic. Malaya*, 226, ♀ (1908).
 Brun., *Rec. Ind. Mus.* IV, 495.
 Kuala Lumpur. *Type* in Leicester's collection.
- parangensis** Ludl., *Can. Ent.* XLI, 24, ♀ (*Pseudouranotaenia*) (1909).
 Theob., *Monog. Culic.* V, 524 (Miss L.'s descr. copied).
 Parang, Mindanao, Phil. Is.
Type in Army Medical Museum, Washington.
- powelli** Ludl., *Can. Ent.* XLI, 235, ♀ (1909).
 Theob., *Monog. Culic.* V, 519, ♀: Brun., *Rec. Ind. Mus.* IV, 497.

- Tayabas, Phil. Is., January.
Type in Army Medical Museum, Washington.
- ascidiicola* de Meij., Ann. Jard. Bot. Buitenzorg, Supp. 3, 925 (1910).
 Java. *Type* in Amsterdam Museum.
- leicesteri* Edw., Bull. Ent. Res. IV, 239 (*nom. nov.* for *fusca*, Leices.) (1913).
 nec *U. fusca*, Theob., 1907.
fusca, Leices., Culic. Malaya, 227, ♂ ♀ (1908).
 Brun., Rec. Ind. Mus. IV, 496.
 Sungei Limbing (Malay Peninsula) [Dr. Daniels].
Type in Leicester's collection.
- metatarsata* Edw., Bull. Ent. Res. V, 81, ♂ (1914).
 Kuala Lumpur, 28-viii-03; 9-x-03 [Leicester].
Type in British Museum.
- moultoni*, *id.*, *loc. cit.*, V, 128, ♂ ♀ (1914-15).
 Kuching reservoir, Sarawak, 6-iii-14 [Moulton].
Type in British Museum.
- brevirostris*, *id.*, *loc. cit.*, V, 284, ♂ ♀ (1914-15).
 Kuching, Sarawak; bred from pitcher plant, 1907 [J. Hewitt].
Types in British Museum.
- obscura*, *id.*, *loc. cit.*, V, 285, ♂ ♀ (1914-15).
 Sarawak. From old collection, Sarawak Museum.
Type in British Museum.

Orthopodomyia, Theob.

Entom. XXXVII, 236 (1904).

Theob., Monog. Culic. IV, 527; V, 470, tab. spp.: Leices., Culic. Malaya, 175; Brun., Rec. Ind. Mus. I, 362; IV, 486; X, 67; Edw., Bull. Ent. Res. IV, 239; How., Dyar and Knab, Mosq. N. Amer. IV, 878.

GENOTYPE, *O. albipes* Theob., by original designation.

Bancroftia, Lutz, in Bourroul's Mosq. do Bras., 59 (1904).

GENOTYPE, *B. albicosta* Lutz, by original designation.

- anopheloides* Giles, Jour. Trop. Med. VI, 315 (*Mansonia*) (1903).
 Edw., Rec. Ind. Mus. IV, 239.
Finlaya anopheloides, Brun., Rec. Ind. Mus. I, 361.
O. maculata, Theob., Rec. Ind. Mus. IV, 29, ♂ (1910).
Id., Monog. Culic. V, 473, ♂.
O. albipes var. *nigritarsis*, Leices., Culic. Malaya, 177.
 Maddathorai, 17-xi-08 [Annandale], *type* of *maculata*; Pangkor Haut, *type* of *nigritarsis* [Dr. Daniels]. *Types* of *anophe-*

loides, ♂ ♀ in British Museum; of *maculata* in Indian Museum; of *nigritarsis* in Leicester collection.

albipes Leices. in Theob., Entom. XXXVII, 237, ♂ ♀ (1904).
Theob., Monog. Culic. IV, 527, ♂ ♀: Leices., Culic. Malaya, 176, ♂ ♀: Brun., Rec. Ind. Mus. I, 362; IV, 486.
Kuala Lumpur, iv [*Leicester*]. Type in British Museum.

maculipes Theob., Monog. Culic. V, 470; fig. 208, head ♀; 209, wing, ♀, 210, wing ♀ (? error for ♂) (1910).

Maddathorai, Travancore, 17-xi-08 [*Annandale*]; Peradeniya, 5-vii-09 [*Green*]; Andaman Is., 22-vii-08 [*Lewis and Ray White*].
Type ♂ in Indian Museum; ♀ in British Museum.

Edwards thinks both *albipes* and *maculipes* probably colour varieties only of *anopheloides*.

Aedeomyia, Theob.

Theob., Monog. Culic. II, 218; fig. 259, scales; 260, map of distribution; (sp. allotted) (1901).

Theob., Jour. Trop. Med. IV, 235 (*nom. nud.*) (1901): *id.*, Monog. Culic. V, 486: *Id.*, Gen. Ins., Fasc. 26, 35: Blanch., Moust., 403, ♂ ♀; fig. 255, var. chars.: Leices., Culic. Malaya, 181: Brun., Rec. Ind. Mus. I, 366; IV, 488; X, 54: How., Dyar and Knab, Mosq. N. Amer., etc., IV, 893.

GENOTYPE, *Aedes squammipennis* Arrib., by my own designation (1914).

Aedomyia. Giles, Handbk., etc., 2nd ed., 478.

Edw., Bull. Ent. Res. III, 24.

squamipenna Arrib., El. Nat. Argent. I, 151 (*Aedes squammipennis*) (1878).

Brun., Rec. Ind. Mus. I, 366; IV, 488.

Aedes squammipenna, Arrib., Dipt. Argent., 62 (1891).

Aedes squampennis, Giles, Handbk., etc., 347.

Aedeomyia squammipenna, Theob., Monog. Culic. II, 219, ♂ ♀; fig. 261, var. chars., ♂ ♀; pl. xxxi, 124, ♀, full ins. col.; pl. E, wing scales; *loc. cit.*, III, 307: Leices., Culic. Malaya, 182, ♂ ♀: Paiva, Rec. Ind. Mus. V, 202, notes.

Aedeomyia squammipennis, Theob., Gen. Ins., Fasc. 26; pl. ii, 9, ♀, full ins. col.: Giles, Handb., etc., 2nd ed., 479.

Aedeomyia squampennis, Blanch., Moust., 404, ♀.

Dawna Hills, 2-iii-08 [*Annandale*]; Calcutta, xi, "at light"; Madras [*Cornwall*]; "at light" at sea, 4 miles off Tuticorin, 25-v-08 [*Paiva*]; Ceylon; Kuala Lumpur; Perak [*Wray*]; Seleo Berlinhafen and Friedrich Wilhelmshafen [both Papua and *Biro*]; Manila [*Banks*]. Also occurs in South America, West Indies and the Sudan. Type of *squamipenna*, location unknown to present author.

catasticta Knab, Ent. News Phil., XX, 387, ♀ (1909).

Brun., Rec. Ind. Mus. IV, 488; Edw., Bull. Ent. Res. III, 25. Samal, Bataan Parang; Mindanao [both *Philippines*]; also Africa.

Type, Cat. No. 12,627, United States National Museum.

Edwards states that British Museum specimens from the Sudan, Gold Coast and both North and South Nigeria agree exactly with Knab's description.

Leptosomatomyia, Theob.

Ann., Mus. Hung. III, 110 (1905).

Theob., Monog. Culic. IV, 548; Brun., Rec. Ind. Mus. I, 362; IV, 488; X, 63.

GENOTYPE, *L. lateralis* Theob., by original designation.

lateralis Theob., Ann. Mus. Hung. III, 110, ♂; fig. 13, head, scutellum, claws (1905).

Id., Monog. Culic. IV, 548, ♂; fig. 252, head, scutellum, claws.

Muina, Papua. 31-xii-00 [*Biro*].

Type (partly dissected) in Hungarian Museum.

Harpagomyia, de Meij.

Tijd. v. Ent. LII, 165 (1909).

Id., loc. cit., LIV, 126 (metamorphoses); pl. xiv, var. chars.: Theob., Monog. Culic. V, 547; Brun., Rec. Ind. Mus. IV, 504; X, 61.

GENOTYPE, *Malaya genurostris* Leices., as *Harpagomyia splendens* de Meij., sp. nov., loc. cit.

Malaya, Leices., Culic. Malaya, 258 (1908).

Brun., Rec. Ind. Mus. IV, 503; X, 64.

GENOTYPE, *M. genurostris*, *id.*, sp. nov., loc. cit.

This name, as pointed out by Edwards, is preoccupied by *Malaia*, Heller, 1891, in Coleoptera.

Grahamia, Theob., Rep. on Dr. Graham's coll. (1909), and Monog. Culic. V, 497, footnote, and 548.

genurostris Leices., Culic. Malaya, 258 (*Malaya*) (1908).

Edw., Bull. Ent. Res. IV, 240.

Harpagomyia splendens, Meij., Tijd. v. Ent. LIII, 167, ♂ ♀, pl. x, 1-9, var. chars. (1909).

Brun., Rec. Ind. Mus. IV, 504.

H. caeruleovittata, Ludl., Psyche XVIII, 131 (1911).

Malay Peninsula; Mindanao, Phil. Is., vii.

Type of *genurostris* in British Museum; of *splendens* in Amsterdam Museum; of *caeruleovittata* in Army Med. Mus., Washington.

Hodgesia, Theob.

Jour. Trop. Med. VII, 17 (1904).

Id., Monog. Culic. IV, 579; V, 544: Gen. Ins., Fasc. 26, 40: Giles, Jour. Trop. Med. VII, 368: Leices., Culic. Malaya, 229: Brun., Rec. Ind. Mus. I, 367; IV, 500; X, 62: Blanch., Moust., 631.

GENOTYPE, *H. sanguinae* Theob., by original designation.

sanguinae Theob., Jour. Trop. Med. VII, 17 (1904).

Giles, *loc. cit.*, VII, 368: Brun., Rec. Ind. Mus. I, 367.

Angeles, Phil. Is. [*Whitmore*]; Luzon. Also occurs in Uganda.

Type in British Museum.

quasisanguineae Leices., Culic. Malaya, 230, ♀ (1908).

Edw., Bull. Ent. Res. IV, 240.

niveocaputis, Ludl., Psyche XVIII, 130, ♀ (1911).

Kuala Lumpur; Fort Pikit, Mindanao, Phil. Is., xi.

Type of *quasisanguinea* in Leicester's collection; of *niveocaputis* presumably in Army Medical Museum, Washington.

malayi Leices., Culic. Malaya, 231, ♂ ♀ (1908).

Brun., Rec. Ind. Mus. IV, 500.

Kuala Lumpur. *Type* in Leicester's collection.

trimaculata Taylor, Trans. Ent. Soc. Lond., p. 204 (1914).

Papua.

Topomyia, Leices.

Culic. Malaya, 238 (1908).

Brun., Rec. Ind. Mus. IV, 499; X, 72.

GENOTYPE, *T. minor*, sp. nov., *loc. cit.*, by my own designation.

Pseudograhamia, Theob., Rec. Ind. Mus. IV, 26 (1910).

Id., Monog. Culic. V, 551: Brun., Rec. Ind. Mus. I, 340; IV, 460; X, 69.

GENOTYPE, *P. aureoventer* Theob., by original designation.

minor Leices., Culic. Malaya, 238, ♂ ♀ (1908).

Brun., Rec. Ind. Mus. IV, 500.

Kuala Lumpur. *Type* in Leicester's collection.

decorabilis Leices., Culic. Malaya, 239, ♂ ♀ (1909).

Selangor. *Type* in Leicester's collection.

argenteoventralis Leices., *loc. cit.*, 240, ♂ ♀ (*argenteoventralis*) (1908).

Brun., Rec. Ind. Mus. IV, 499: Edw., Bull. Ent. Res. IV, 240.

Pseudograhamia aureoventer, Theob., Rec. Ind. Mus. IV, 27, ♀ (1910): Theob., Monog. Culic. V, 551, ♀: Brun., Rec. Ind. Mus. IV, 460, 489.

Selangor; Pallode, Travancore, 16-xi-08 [Annandale]. Type of *argenteoventralis* in Leicester's collection; of *aureoventer* in Indian Museum. Theobald says (Monog. V) that the type of *aureoventer* is in the British Museum, but it is in the Indian Museum.

argyropalpis Leices., Culic. Malaya, 242, ♂ ♀ (1908).

Selangor, iv; Kuala Lumpur, iii. Types (a unique pair) in Leicester's collection.

rubithoracis, *id.*, *loc. cit.*, 243, ♂ ♀ (1908).

Malaysia. Types in Leicester's collection.

gracilis, *id.*, *loc. cit.*, 244, ♂ ♀ (1908).

Selangor, v. Type in Leicester's collection.

nigra, *id.*, *loc. cit.*, 245, ♂ ♀ (1908).

Ampang, near Kuala Lumpur. Type in Leicester's collection.

dubitans, *id.*, *loc. cit.*, 246, ♀ (1908).

Selangor, v. Type, a unique specimen, probably lost, not in Leicester's collection.

tipuliformis, *id.*, *loc. cit.*, 247, ♂ (1908).

Brun., Rec. Ind. Mus. IV, 500.

Selangor [Raub.] Type in Leicester's collection.

T. argyroventralis may be the ♀ of this species, according to Leicester, instead of the ♀ described as such, although the way he mentions it is that *tipuliformis* may be the real ♂ of *argenteoventralis*. If this proves to be the case the name of the ♂ (*tipuliformis*) should stand, in spite of *argenteoventralis* being described on a previous page.

Zeugomyia, Leices.

Culic. Malaya, 231 (1908).

Brun., Rec. Ind. Mus. IV, 498; X, 73.

GENOTYPE, *Z. gracilis* Leices., by original designation.

gracilis Leices., Culic. Malaya, 232, ♂ ♀ (1908).

Brun., Rec. Ind. Mus. IV, 498.

Kuala Lumpur; Jugra, x, xi, xii. Type in Leicester's collection.

Rachionotomyia, Theob.

Jour. Bomb. Nat. Hist. Soc. XVI, 248 (1905).

Theob., Monog. Culic. IV, 518; Edw., Bull.

Ent. Res. IV, 240; Brun., Rec. Ind. Mus. I, 368; IV, 485; X, 70.

GENOTYPE, *R. aranoides*, Theob., as *R. ceylonensis*, sp. nov., by original designation.

Runchomyia, Theob., Monog. Culic. III, 319 (1903).

Theob., *loc. cit.*, IV, 585, note on name; V, 555, note

on name: *id.*, Gen. Ins., Fasc. 26, 38: Brun., Rec. Ind. Mus. I, 365; IV, 493; X, 71.

GENOTYPE, *R. frontosa*, *id.*, *loc. cit.*, sp. nov.

Binotia, Blanch., Arch. de Parasit. VIII, 478 (1904): *nom. nov.*, under suggested preoccupation by *Rhynchomyia*, Rob. Des. in Muscinae.

Polylepidomyia, Theob., Ann. Mus. Hung. III, 118 (1905).

Theob., Monog. Culic. IV, 625: Brun., Rec. Ind. Mus. I, 366; IV, 502; X, 68.

GENOTYPE, *P. argenteiventris* Theob., by original designation.

Colonemyia, Leices., Culic. Malaya, 233 (1908).

Brun., Rec. Ind. Mus. IV, 498; X, 58.

GENOTYPE, *C. caeruleocephala*, Leices., by my own designation.

Skeiromyia, *id.*, *loc. cit.*, 248 (1908).

Brun., Rec. Ind. Mus. IV, 500; X, 71.

GENOTYPE, *S. fusca*, Leices., by original designation.

Squamomyia, Theob., Rec. Ind. Mus. IV, 28 (1910).

Id., Monog. Culic. V, 529: Brun., Rec. Ind. Mus. IV, 489; X, 71.

GENOTYPE, *S. inornata*, Theob., by original designation.

aranoides Theob., Monog. Culic. II, 274. ♀ (*Wyeomyia*) (1901).

Rachionotomyia aranoides, Edw., Bull. Ent. Res. IV, 241.

Wyeomyia aranoides, Giles, Handbk., 2nd ed., 499, ♀: Blanch., Moust., 425.

Rachionotomyia ceylonensis, Theob., Jour. Bomb. Nat. Hist. Soc.

XVI, 248, ♀; pl. B. 6, scutellum (1905): Brun., Rec. Ind. Mus. I, 369.

Skeiromyia fusca, Leices., Culic. Malaya, 248, ♂ ♀ (1908).

Brun., Rec. Ind. Mus. IV, 500.

Squamomyia inornata, Theob., Rec. Ind. Mus. IV, 28, ♂ (1910).

Id., Monog. Culic. V, 529, ♂.

Straits; Malaysia; Peradeniya, x-01 [Green]; Dawna Hills, 2000—3000 ft., 2—3-iii-08 [Annanale]. Types of *aranoides* and *ceylonensis* in British Museum; of *fusca* in Leicester's collection; of *inornata* in Indian Museum.

nitidiventer Giles, Jour. Trop. Med. VII, 368 (*Uranotaenia*) (1904).

Rachionotomyia nitidiventer, Edw., Bull. Ent. Res. IV, 241.

Runchomyia philippinensis, Giles, Jour. Trop. Med. VII, 368 (1904): Theob., Monog. Culic. V, 555 (Gile's descr. copied):

Brun., Rec. Ind. Mus. IV, 493.

Pampanga, Phil. Is. [Whitmore].

Types of both forms in British Museum.

argenteiventris Theob., Ann. Mus. Hung. III, 118, ♀, fig. 15, head, scutellum, bristles (*Polylepidomyia*) (1905).

Id., Monog. Culic. IV, 625, fig. 291, head, etc.; fig. 292, wing, ♀.

Paumomu River (Papua), ix, xii, 1892 [*Loria*].

Type in Hungarian Museum.

caeruleocephala Leices., *Culic. Malaya*, 233, ♂ ♀ (*Colonemyia*) (1908).

Brun., *Rec. Ind. Mus.* IV, 498.

Bukit Kutu; Ulu Klang. Type in Leicester's collection.

similis Leices., *Culic. Malaya*, 235, ♂ ♀ (*Colonemyia*) (1908).

Bukit Kutu; Malaysia. Type in Leicester's collection.

hybrida, *id.*, *loc. cit.*, 237, ♀ (*Colonemyia*) (1908).

Bukit Kutu; Malaysia. Type in Leicester's collection.

mendacis Daniels, *Studies Inst. Med. Res. Fed. Malay States*, III, 266, ♂ ♀ (1908), (*Colonemyia*) (1908).

Brun., *Rec. Ind. Mus.* IV, 499.

East Coast Malay Peninsula.

Type probably lost, not in Leicester's collection. Dr. Daniels' paper follows that of Leicester's in the same publication.

affinis Edwards, *Bull. Ent. Res.* IV, 241, *nom. nov.* for *Phoniomyia caeruleocephala*, Theob. (*nec* Leices.).

Phoniomyia caeruleocephala, Theob., *Monog. Culic.* V, 577, ♂ ♀, fig. 252, wing (1910).

nec Colonemyia caeruleocephala, Leices., *Culic. Malaya*, 233 (1908).

Hakgala, Ceylon, iii-07 [*Green*].

Types, a unique pair, in British Museum.

vicina Edw., *Bull. Ent. Res.* V, 128, ♂ ♀ (1914).

Kuching Reservoir, 6-iii-14 [*Moulton*].

Types in British Museum.

nepenthis, *id.*, *loc. cit.* V, 285, ♂ ♀ (1914).

Bred from pitcher plant, xi-07 [*J. Hewitt*].

proxima, *id.*, *loc. cit.* V, 285, ♂ ♀ (1914).

Ulu Klang [*Leicester*] from old collection, Sarawak Museum, including type: Ulu Gombak [*Stanton*].

Types in British Museum.

Wyeomyia, Theob.

Monog. Culic. II, 267 (1901).

Theob., *Jour. Trop. Med.* II, 235 (*nom. nud.*);

id., *Monog. Culic.* III, 310, restricted (1903); V,

579, notes: *id.*, *Gen. Ins.*, Fasc. 26, 38; Blanch.,

Moust., 423; Giles, *Handbk.*, etc., 2nd ed. 495;

Leices., *Culic. Malaya*, 250; Brun., *Rec. Ind.*

Mus. I, 365; IV, 501; X, 73; How., *Dyar and*

Knab, Mosq. N. Amer., etc., IV, pt. 1, 49.

GENOTYPE, *W. grayii* Theob., by designation of Neveu-Lemaire (1902).

Phoniomyia, Theob., *Monog. Culic.* III, 311

(1903); Theob., *loc. cit.* IV, 598; *Gen. Ins.*, Fasc.

26, 38: Blanch., Moust., 425: Leices., Culic. Malaya, 253: Brun., Rec. Ind. Mus. I, 365; IV, 502; X, 68.

GENOTYPE, *Wyeomyia longirostris* Theob., by designation of Dyar.

Dendromyia, Theob., *loc. cit.* III, 313 (1903).

Id., *loc. cit.* IV, 603, list, spp.; V, 587, tab. spp.: *id.*, Gen. Ins., Fasc. 26, 39: Blanch., Moust. 426: Leices., Culic. Malaya, 25; Brun., Rec. Ind. Mus. IV, 502; X, 59.

GENOTYPE, *D. ulocoma* Theob., by personal designation (1914). Howard, Dyar and Knab give *Wyeomyia luteoventralis* Theob.

Bothodeomyia, Theob., Rec. Ind. Mus. IV, 31 (1910): *Id.*, Mong. Culic. V, 581: Brun., Rec. Ind. Mus. IV, 492; X, 56.

GENOTYPE, *B. complex*, Theob., sp. nov., by original designation.

Heizmannia, Ludl., Can. Ent. XXXVII, 130 (*Heizmannia*, lapsus) (1905): Banks, Phil. Jour. Sci. I, 99: Brun., Rec. Ind. Mus. I, 366; X, 61. GENOTYPE, *H. scintillans* Ludl., by original designation.

greenii Theob., Jour. Bomb. Nat. Hist. Soc., XVI, 247, ♂ ♀; pl. B. 5, antenna (1905).

Id., Monog. Culic. IV, 596: Brun., Rec. Ind. Mus. I, 366; IV, 501.

Peradeniya [Green], a perfect pair. *Types* in British Museum.

bimaculipes Theob., Ann. Mus. Hung. III, 114, ♀ (*Phoniomyia*) (1905): *id.*, Mong. Culic. IV, 600, ♀.

Moroka, Papua, 1300 metres alt., vii—ix; Fredrich Wilhelmshafen, Papua [Biro]. *Type* in Hungarian Museum.

indica Theob., Ann. Mus. Hung. III, 115, ♂ ♀; pl. ii, wing ♀; pl. iii, wing scales ♀ (*Phoniomyia*) (1905).

Phoniomyia indica, Theob., Mong. Culic. IV, 601, ♂, fig. 275, wing ♀: Leices., Culic. Malaya, 253, ♂ ♀: Brun., Rec. Ind. Mus. I, 365; IV, 502, notes.

Kuala Lumpur; Singapore. *Type* in British Museum.

scintillans Ludl., Can. Ent. XXXVII, 130, ♀ (*Heizmannia*, lapsus) (1905).

Angeles, Pampanga, Phil. Is., ix?

Type in Army Medical Museum, Washington.

metallica Leices., Culic. Malaya, 251, ♀ (1908).

Bukit Kutu, Malaysia.

Type, a unique ♀, in Leicester's collection.

funerea, *id.*, *loc. cit.*, 252, ♀ (1908).

Kuala Lumpur. *Type*, a unique ♀, in Leicester's collection

- communis**, *id.*, *loc. cit.*, 254, ♂ ♀ (*Dendromyia*) (1908).
Selangor. *Types* in Leicester's collection.
- aureochaetae**, *id.*, *loc. cit.*, 255, ♀ (*Dendromyia*) (1908).
Kuala Lumpur, iv, v. *Type* in Leicester's collection.
- achaetae**, *id.*, *loc. cit.*, 257, ♀ (*Dendromyia*) (1908).
Brun., *Rec. Ind. Mus.* IV, 503.
Malaysia, a unique ♀. *Type* in Leicester's collection.
- nepenthicola** Banks, *Phil. Jour. Sci.* IV, 550, ♂ ♀ (1910).
Benguet (Trinidad, *Phil. Is.*), 30-xii-07 [*Banks*].
Type in Entomological collection, Cat. No. 8159, Bureau of Science, Manila.
- complex** Theob., *Rec. Ind. Mus.* IV, 31, ♂ ♀ (*Bolbodeomyia*) (1910).
Id., *Mong. Culic.* V, 581, ♂ ♀, fig. 253, ♂ genitalia, 254, wing, ♂; 255, wing, ♀: Brun., *Rec. Ind. Mus.* IV, 492.
Dawna Hills, 2000—3000 ft., 1-3-iii-08 [*Annandale*].
Types, a unique pair, in Indian Museum.

Sub-family CORETHRINAE.

- Giles, *Handbk.*, 2nd. ed., 500.
Brun., *Rec. Ind. Mus.* IV, 317; 504; *loc. cit.*, VI, 227.

Chaoborus, Lichtenstein.¹

- Arch. Zool.* I, 174 (1800).
GENOTYPE, *Tipula plumicornis* F., as *antisepticus*, sp. nov.
Sayomyia, Coq., *Can. Ent.* XXXV, 189 (1903).
Theob., *Gen. Ins.*, Fasc. 26, 42: Brun., *Rec. Ind. Mus.* I, 368; X, 71.
GENOTYPE, *Corethra punctipennis* Say., sp. nov.
- manilensis** Sch., *Nov. Reise, Dipt.* 30 (*Corethra*) (1868).
Edw., *Bull. Ent. Res.* IV, 242.
Sayomyia manilensis, Brun., *Rec. Ind. Mus.* I, 368.
Corethra manillensis, Giles, *Handbk.*, etc., 360, 2nd ed., 504.
Corethra asiatica, Giles, *Jour. Bomb. Nat. Hist. Soc.* XIII, 610 (1901)²: Brun., *Rec. Ind. Mus.* I, 368.
Chaoborus asiatica, Brun., *Rec. Ind. Mus.* IV, 505.
Chaoborus manillensis (larva and pupa); Bains Prashad, *Rec. Ind. Mus.* XV, 154, pl. xix, 1-7.
Calcutta, v, vii, viii, ix, xi, xii, often at light [*Annandale*];
Sibpur, near Calcutta, viii; Katihar, Purnea Distr., at light;
Shajahanpur, Punjab [*Giles*]; Manila; Yenna R., Medha, Satara

¹ For discussion on generic synonymy in Corethrinae, see Brun., *Rec. Ind. Mus.* IV, 317 and VI, 227.

² A two-line "description."

Dist., Bombay Presidency. *Type*: location unknown to present author.

cornfordi Theob., Monog. Culic. III, 339, ♀ (*Corethra*) (1903).

Shaohyling, China, v, vi [*Cornford*]. *Type*, location unknown to present author.

Corethrella, Coq.

Jour. N. Yk. Ent. Soc. X, 191 (1902).

GENOTYPE, *C. brakeleyi*, Coq., by original designation.

Ramcia, Annand., Spol. Zeyl. VII, 187 (1911).

GENOTYPE, *R. inepta*, *id.*, sp. nov., *loc. cit.*

inepta Annand., Spol. Zeyl. VII, 189, fig. (p. 188) and pl. (larva, pupa, wing, etc.) (1911).

Diyatalawa, Ceylon, 4,300 ft. Larva, pupa and one imago, a ♂ taken by Maj. MacDougall, R.A.M.C.

Type in the British Museum; *co-type* of larva in the Indian Museum.

Family DIXIDAE.

Dixa, Mg.

Syst. Besch. I, 316 (1818).

Sch., F. Austr. II, 641: Brun., Fauna Brit.

Ind., 258, fig. 38, var. chars.: *id.*, Rec. Ind. Mus.

IV, 264, tab. Ind. spp.

GENOTYPE, *Dixa maculata* Mg., by Curtis's designation.

montana Brun., Rec. Ind. Mus. IV, 265, ♂ ♀ (1911).

Id., Fauna Brit. Ind., 262, ♂ ♀, pl. xii, 9, wing.

Simla, 10-v-09, *type* ♂; Phagu, 11-v-09, *type* ♀; Barogh, 5000 ft., 10-v-09 [all Simla Distr. and all Annandale].

Types in Indian Museum.

maculipennis Brun., Rec. Ind. Mus. IV, 266, ♂ ♀ (1911).

Id., Fauna Brit. Ind., 263, ♂ ♀, pl. xii, 10, wing.

Darjiling, 29-v-10, *type* ♂ [*Brunetti*]; Matiana, 8000 ft., Simla Distr., 28—30-iv-07, *type* ♀ [*Annandale*].

Types in the Indian Museum, a unique pair.

ochrolineata Brun., Rec. Ind. Mus. IV, 267, ♀ (*ochrolineata*) (1911).

Id., Fauna Brit. Ind. 261, ♀, pl. xii, 8, wing.

Kurseong, 9-ix-09 [*Annandale*].

Type, unique, in Indian Museum.

bistriata Brun., Rec. Ind. Mus. IV, 269, ♂ (1911).

Id., Fauna Brit. Ind. 264, ♂.

Darjiling, 29-v-10. [*Brunetti*].

Type, unique, in Indian Museum.

- bifasciata* Brun., Rec. Ind. Mus. IV, 269, ♀ (1911).
Id., Fauna Brit. Ind. 264, ♀.
 Phagu, Simla Distr., 12-v-09 [Annandale].
Type, unique, in Indian Museum.
- binotata* Edw., Jour. Fed. Malay Sts. Mus., VIII, 9, ♂ (1919).
 Korinchi Peak, Sumatra.

Family TIPULIDAE.

Sub-family PTYCHOPTERINAE.

Ptychoptera, Mg.

Illig. Mag. II, 262 (1803)

Sch., F. Austr. II, 495: Brun., Fauna Brit.

Ind., 279: *Auctt.*GENOTYPE, *Tipula contaminata* L., designated
 by Latreille (1810).? *Ctenoceria*, Rond., Dipt. Ital. Prod. I, 187 (1856).

- distincta* Brun., Rec. Ind. Mus. VI, 232, ♀ (1911).
Id., Fauna Brit. Ind., 281, ♀; pl. v, 1, wing (1912).
 Darjiling, 7000 ft., 3-9-vi-09 [Howlett].
Type in Pusa collection, a unique specimen.
- tibialis* Brun., Rec. Ind. Mus. VI, 233, ♂ ♀ (1911):
Id., Fauna Brit. Ind. 282, ♂ ♀ (1912): *id.*, Rec. Ind. Mus.
 XV, 256.
atritarsis, Brun., Rec. Ind. Mus. VI, 234 ♀ (1911): *id.*, Fauna
 Brit. Ind., 283, ♀ (1912): *id.*, Rec. Ind. Mus. XV, 256.
 Darjiling, 7000 ft., 7-16-x-05, 30-ix-08, 7-vi-17, 1st week
 in June 1917, very common [all *Brunetti*]; Siliguri, 18-20-
 vii-07; Lebong, 6000-6600 ft., near Ghoom, 13-vi-14 [Gravelly].
Type ♂ of *tibialis* in Indian Museum; *type* ♀ and *cotype* ♂ ♂
 in my collection; *cotype* ♀ in Vienna Museum; *type* of *atritarsis*
 in Indian Museum.
- annandalei* Brun., Rec. Ind. Mus. XV, 296, ♂ (1918).
 Kalaw, 4000-45000 ft., South Shan States, 10-iii-17 [Gravelly].
 Two ♂ ♂. *Type* in Indian Museum.

Tanyderus, Phil.

Verh. zool.-bot. Ges. Wien, XV, 780, pl. xxix, 57 (1865).

Ost. Sack., *loc. cit.*, XXIX, 517-522 (1879):

Brun., Rec. Ind. Mus. XV, 255.

GENOTYPE, *Cylindrotoma ornatissima* Dol., by
 original designation.

- ornatissimus* Dol., Nat. Tijd. Ned. Ind. XVII, 80, ♂ (*Cylindro-*
toma) (1858).
 Ost. Sack., Berl. Ent. Zeits., XXXI, 228, ♂ ♀: Brun., Rec.
 Ind. Mus. VI, 235: Ender., Zool. Jahr. XXII, 84.
 Amboina, Insel Obi. *Type* in Vienna Museum, damaged.

mirabilis de Meij., Tijds. v. Ent. LVIII, 104, fig. 1, wing (1915).

Id., Nov. Guin. Res. XIII, 51 (1915).

Papua.

Sub-family *TIPULINÆ*.

Section *CTENOPHORINI*.¹

Xiphura, Brullé.

Ann. Soc. Ent. Fran. I, 206 (1832).

GENOTYPE, *Tipula atrata* L., the original species.

indica Brun., Rec. Ind. Mus. XV, 257, pl. viii, 1, antenna (1918).

Darjiling 6900 ft., 1-vi-17. A perfect unique ♂ in my own collection, taken by me.

Ctenophora, Mg.

Illig. Mag. II, 263 (1803).

Sch., F. Austr. II, 498: Ost. Sack., Berl. Ent. Zeits. XXX, 164: Brun., Rec. Ind. Mus. VI, 235: Fauna Brit. Ind., 288: *Auctt.*

GENOTYPE, *Tipula flaveolata* F. designated by Rondani, 1856.²

Tanyptera, Latr., Nouv. Dict. Hist. Nat., 428 (1804).

¹ For characters, v. Ost. Sack., Berl. Ent. Zeits., xxx, 164, *et seq.*; and Brun., Fauna, 287.

² A little investigation will, I think, show that Coquillett's desire (1910) to restrict *Ctenophora* to *atrata* L. and its allies, and to set up *Phorocentia* for the remaining species of *Ctenophora* in a wide sense, is wrong. He accepts Latreille's designation (1810) of *atrata* L. as type species of *Ctenophora*, which would have therefore left the bulk of the remaining species of the genus without a name, but there may be a flaw in his reasoning which will save the name of Meigen's genus for the group of species so long associated with it. Meigen in 1803 (Illig. Mag.) set up *Ctenophora* for *pectinicornis* L., *bimaculata* L., *atrata* F., and *flaveolata* F. Two years later Latreille erected *Tanyptera* for *atrata* L. So far as can be judged at this length of time, the genus was not favourably received, all the species being still retained in *Ctenophora*. In 1810 Latreille apparently himself acquiesced in this view as he selected *atrata* as the type of *Ctenophora*. In 1832 Brullé proposed *Xiphura* with *atrata* as the sole species, and in 1833 *Dictenidia* with *A. bimaculata* L. as type and these genera were practically universally adopted. Rondani in his list of Italian genera (Prodrome 1856) accepted *atrata* as type species of *Xiphura* and selected *flaveolata* (one of Meigen's four original species) as type of *Ctenophora*, and this view, so far as I am aware, was not challenged. By general consent for the last three quarters of a century, *atrata* has been regarded as not congeneric with the other species of the old *Ctenophora*, and the only question to settle is in which genus it should be placed. Meigen selected no type species out of the four on which he founded *Ctenophora*; therefore Latreille in selecting *atrata* as type of his *Tanyptera* (1805) has undoubtedly prior claim to the species, as it was the sole exponent of his genus, and the fact that the dipterologists of his day apparently ignored the genus does not prevent it being resuscitated if originally legitimately founded. Therefore when Latreille himself, in 1810, acquiescing in the suppression of *Tanyptera*, selected *atrata* as the type species of *Ctenophora*, he went beyond his rights. *Xiphura* in 1832, also founded on *atrata* alone becomes an absolute synonym of *Tanyptera*. Rondani, recog-

xanthomelana Walk., List. Dipt. Brit. Mus. I, 77, ♀ (1848).

Ost. Sack., Berl. Ent. Zeits. XXX, 166; Wulp. Tijd. v. Ent. XXVIII, 81. Brun., Rec. Ind. Mus. VI, 235, 241; Fauna Brit. Ind., 289. East India. Type in British Museum.

This species is retained under *Ctenophora* but the ♂ is not known and it probably belongs to *Pselliophora*. *C. melanura*, Walk. (List Dipt. Brit. Mus. I, 78; 1848) is a *Sargus* (*Stratiomyidae*). Alexander now suggests that *xanthomelana* is probably identical with *Prionota nigriceps*, Wulp. If so, of course Walker's name takes precedence.

Pselliophora, Ost. Sack.

Berl. Ent. Zeits. XXX, 165 (1886).

de Meij., Tijd. v. Ent. LIV, 63, tab. Javan spp.: Brun., Rec. Ind. Mus. VI, 236, tab. orient. spp. and notes: *id.*, Fauna Brit. Ind., 290.

GENOTYPE, *Tipula laeta* F., by my own designation (Fauna, 290).

laeta Fab., Ent. Syst. IV, 239 (*Tipula*) (1792).

Ctenophora laeta, *id.*, Syst. Antl., 19; Wied., Dipt. Exot. I, 20 (*Ctenophora*): *id.*, Auss. Zweifl. I, 38¹; Macq., Dipt. Exot. I, 1, 42, pl. ii, 11 (full ins.): Brun., Rec. Ind. Mus. VI, 240, notes: XV, 257; *id.*, Fauna Brit. Ind., 291, ♂ ♀, pl. vi, 9, antenna (*Pselliophora*).

Widely distributed in India and the East; Dehra Dun, Purneah, Sind, Calcutta, Bangalore, Travancore, Bombay, Goa, Nilgiri Hills, Ceylon. A variable species.

Type presumably in Fabricius's collection.

var. *trilineata* Brun., Rec. Ind. Mus. VI, 240 (1911).

Id., Fauna Brit. Ind., 292, pl. v, 2, wing, var.

Unique type of this variety in Indian Museum.

ardens Wied., Dipt. Exot. I, 20, ♀ (*Ctenophora*) (1821).

Id., Auss. Zweifl. I, 39, ♀ (*Ctenophora*).

Pselliophora, *id.*, de Meij., Tijd. v. Ent. LIV, 61; Brun., Rec. Ind. Mus. VI, 238 (♂ noted): Alex., Proc. U.S. Nat. Mus. XLIX, 179, pl. xlv, 31, wing.

Batavia and environs, x [Jacobson]; Lombok; Sumatra; Buitenzorg, iii-09 [Bryant and Palmer].

Type (a ♀) in Wiedemann's collection: the ♂ was apparently first described by me from a specimen in the Vienna Museum.

nizing *atrata* as not congeneric with the other species of *Ctenophora*, retained *Xiphura* with *atrata* as type, and in selecting *flaveolata* F. as type of *Ctenophora* was entirely within his rights, and the latter species must be accepted as genotype of the genus in its generally accepted sense. As regards *Tanyptera* versus *Xiphura*, on the grounds of continuity before priority I retain the latter as it has been recognized for nearly ninety years. *Ctenophora* is almost certainly not oriental, but it is as well to keep the name in the lists until the discovery of the ♂ of Walker's *xanthomelana* decides this question.

¹ Some authors erroneously report a figure of this species by Wiedemann.

compedita Wied., Dipt. Exot. I, 21, ♀ (*Ctenophora*) (1821).

Id., Auss. Zweifl. I, 39, ♀ [*Ctenophora*].

Pselliophora, *id.*, Wulp, Tijds. v. Ent. XXIII, 156; pl. x, 1: *id.*, Notes Leyd. Mus. VI, 25; Brun., Rec. Ind. Mus. VI, 238; IX, 257; Alex., Proc. U.S. Nat. Mus. XLIX, 180, pl. xiv, 32, wing.

Darjiling, 1000—3000 ft. [Lord Carmichael's colln.]; Semarang, xi [Jacobson]; Depok, iii, 09 [Bryant and Palmer]; Celebes; Aru.

Type of species (a ♀) in Westermann's coll.; ♂ in Leyden Museum.

fuscipennis Macq., Dipt. Exot., Supp. I, 10, pl. i, 4 wing (*Ctenophora*) (1846).

Pselliophora, *id.*, Ost. Sack., Berl. Ent. Zeits. XXX, 172; Brun., Rec. Ind. Mus. VI, 238; IX, 254.

Ctenophora javanica, Dol., Nat. Tijds. Ned. Ind. X, 406, pl. ix, 1, full ins. ♂ (1856); Wulp, Tijds. v. Ent. XX, 156, ♂ noted, pl. x, 2, wing; Brun., Rec. Ind. Mus. VI, 238.

Djokjokarta, Java.

Type in Bigot coll. A ♂ and ♀ from the Darjiling District in the Indian Museum appear to be this species, also two specimens with a red thorax in Lord Carmichael's collection from Singla, 1500 ft. and Darjiling District, 1000—3000 ft. Originally recorded erroneously by Macquart from Brazil.

taprobanes Walk., List Dipt. Brit. Mus., I, 77, ♀ (*Ctenophora*) (1848).

Brun., Rec. Ind. Mus. VI, 240, ♂ noted: *id.*, Fauna Brit. Ind., 293, ♀, pl. v. 3, wing, pl. vi, 10, 11, ♂ genitalia.

Ceylon, not uncommon. May to September.

Type ♀ in British Museum.

fumiplena Walk., Ins. Saund Dipt., pt. v. 449, ♀ (*Ctenophora*) (1856).

Ost. Sack., Berl. Ent. Zeits. XXX, 170; Brun., Rec. Ind. Mus. VI, 240; XV, 258; Edw., Ann. Mag. Nat. Hist. (8) XVIII, 256, notes.

China.

Type ♀ in British Museum. This species is probably out of place in this Catalogue as apparently it comes from North China, from whence Mr. Edwards informs me the ♀ ♀ in the British Museum come.

chrysophila Walk., Proc. Linn. Soc. Lond. I, 6, ♀ (*Ctenophora*) (1857).

Brun., Rec. Ind. Mus. VI, 238; *id.*, Fauna Brit. Ind., 295, ♀.

Bhim Tal, 19—22-ix-06 [Annandale]; Singapore; Tura, Assam, viii-17 [Kemp].

Type ♀ in British Museum, the ♂ apparently still unknown.

incunctans Walk., Proc. Linn. Soc. Lond. IV, 93, ♂ ♀ (*Ctenophora*) (1860).

- Wulp., Notes Leyd. Mus. VI, 255 (*Ctenophora*): *id.*, Cat. Dipt. S. Asia, 37, note: Ost. Sack., Berl. Ent. Zeits. XXX, 171: Brun., Rec. Ind. Mus. VI, 239 (varieties noted).
Ctenophora velutina, Wulp, Notes Leyd. Mus. VI, 256, ♀.
 Celebes, Tondano, Saugir.
 Type in British Museum, in good condition.
- gaudens** Walk., Proc. Linn. Soc. Lond. IV, 93, ♂ ♀ (*Ctenophora*) (1860).
 Ost. Sack., Ann. Mus. Gen. XVI, 399 (*Ctenophora*): *id.*, Berl. Ent. Zeits. XXX, 171: Brun., Rec. Ind. Mus. VI, 238: IX, 257, pl. xiv, 8, ♂ genitalia.
 Celebes.
 Type ♂ ♀ in British Museum. A ♂ in the Indian Museum, from Singla, Darjiling District, v-13, may be this species.
- suspirans** Ost. Sack., Berl. Ent. Zeits. XXVI, 93, ♂ (*Ctenophora*) (1882).
 Brun., Rec. Ind. Mus. VI, 241.
 Philippines. Bezzi (Phil. J. Sci. XII, Sect. D, p. 109, ♂ ♀) describes a variety from three localities in Luzon.
 Type, location unknown.
- idalia** Ost. Sack., Berl. Ent. Zeits. XXVI, 94, ♂ (*Ctenophora*) (1882).
 Philippines.
 Type, a unique ♂; location unknown.
- dolens** Ost. Sack., Berl. Ent. Zeits. XXVI, 94, ♂ ♀ (*Ctenophora*) (1882).
 Brun., Rec. Ind. Mus. VI, 240.
 Philippines.
 Types, location unknown.
- curvipes** Wulp, Notes Leyd. Mus. VI, 254, ♀ (*Ctenophora*) (1884).
 Brun., Rec. Ind. Mus. VI, 238.
 Gorontalo.
 Type, a unique ♀, in Leyden Museum.
- annulosa** Wulp, Notes Leyd. Mus. VII, 1, ♀ (*Ctenophora*) (1885).
 Brun., Rec. Ind. Mus. VI, 241.
 Java.
 Type, a unique ♀, in Leyden Museum.
- rubra** Ost. Sack., Berl. Ent. Zeits. XXX, 171, ♂ (1886).
 Brun., Rec. Ind. Mus. VI, 238: Alex., Proc. U.S. Nat. Mus. XLIX, pl. xlv, 33, wing.
 Laos, Siam [*Mouhot*]; Buitenzorg, iii-09; 4-iv-09 [*Bryant and Palmer*].
 Type in Berlin Museum.
- nigrithorax** Wulp in de Meij., Bijd. tot de Dierk. XVII, 87, ♂ ♀ (1904).
 Brun., Rec. Ind. Mus. VI, 238.
 Java.
 Types in Amsterdam Museum.

- insignis** de Meij., *Bijd. tot de Dierk.*, XVII, 87, ♂ (1904).
Brun., *Rec. Ind. Mus.* VI, 241.
Java (Tosari).
Type in Amsterdam Museum, a unique ♂.
- stigmatica** de Meij., *Bijd. tot de Dierk.* XVII, 88, ♀, pl. viii,
1 full ins. (1904).
Brun., *Rec. Ind. Mus.* VI, 238.
Lahago, Central Nias Island.
Type, a unique ♀, in Amsterdam Museum.
- bifascipennis** Brun., *Rec. Ind. Mus.* VI, 241, ♀ (1911).
Shanghai.
Type, a unique ♀, labelled "Consul Haas" in Vienna Museum. Possibly out of place in this Catalogue.
- divisa**, *id.*, *loc. cit.*, 242, ♂ (1911).
Edw., *Ann. Mag. Nat. Hist.* (8) XVIII, 254, notes: Brun.,
Rec. Ind. Mus. XV, 258.
"East Indies, 1892" [*Felder*]; Pashok Spur, Darjiling Distr.
[*Lister*]; Sikkim, v, 96; Bhutan; Lat Bua, Kao, East Siam
[*C. Boden Klöss*].
Type (a ♀) in Vienna Museum.
- immaculipennis** Brun., *Rec. Ind. Mus.* VI, 243, ♀ (1911).
Id., *Fauna Brit. Ind.* 294, ♀, pl. vi, 7, full ins., 8, antenna.
Sylhet [*Hall*].
Unique *type* in Indian Museum.
- terminalis**, Brun., *Rec. Ind. Mus.* VI, 244, ♀ (1911).
Montes Mauson, 2000—3000 ft., Tonkin, iv or v [*Fruhstorfer*].
Unique *type* in Vienna Museum.
- ctenophorina** Riedel, *Ent. Mitt. Berl.* II, 273 (1913).
Kankau (Koshun), Formosa, v, vi, vii, ix, 1912.
Edwards records a ♂ in the British Museum from Koannania,
S. Formosa, 15 vi-06 [*Wileman*].
Types and *cotypes* in *Deuts. Ent. Mus. Berl.*, *cotypes* in Riedel
collection.
- strigipennis** de Meij., *Tijd. v. Ent.*, LVI, Supp. p. 5, ♀ (1913)
(Mar. 1914).
Java.
- speciosa** Edw., *Ann. Mag. Nat. Hist.* (8) XVIII, 254, ♂ (1916).
Khasi Hills, Assam.
Unique *type* in British Museum.
- semirufa** Edw., *loc. cit.*, 255, ♀ (1916).
Taihoku, Formosa, xi-09 [*Nitobe*].
Unique *type* in British Museum.
- luctuosa** de Meij., *Tijd. v. Ent.* LIX, 199, ♀ (1916).
Gunung Susuruh, Preanger, Java [*Corporaal*].
Unique *type* in Amsterdam Museum.
- praefica** Bezzi, *Phil. Jour. Sci.* XII, Sect. D, 110, ♂ (1917).
Mindanao, *Phil. Is.* [*Baker*].

- tripudians id.*, *loc. cit.*, III, ♀ (? *praefica* ♀) (1917).
Laguna, Phil. Is. [Baker].
- approximata* Brun., Rec. Ind. Mus. XV, 259, ♀ (1918).
Talewadi, N. Kanara Dist., 3--10-x-16 [Kemp].
Unique *type* in Indian Museum.
- flavofasciata id.*, *loc. cit.*, 259, ♂ ♀, pl. viii, 2, abdomen, 3, wing (1918).
Above Tura, Garo Hills, Assam, 3500--3900 ft, vii-17 [Kemp].
Types in Indian Museum.
- aurantia, id.*, *loc. cit.*, 260, ♀ (1918).
Above Tura, Garo Hills, Assam, 3500--3900 ft., viii-17 [Kemp].
Type in Indian Museum.
- latifascipennis, id.*, *loc. cit.*, 260, ♀ (1918).
Pollibetta, Coorg, S. India, 24-x--16-xi-15 [Fletcher].
Type, presented to British Museum.
- ? *P. elongata* Edw., Ann. Mag. Nat. Hist. (8) XII, 202, ♀ (1913).
Hakgala, Ceylon, 4500 ft., v-1911.
Unique *type* in British Museum.

Prionota, Wulp.

Notes Leyd. Mus. VII, 1 (1885).

Edw., Jour. Fed. Malay Sts. Mus. VIII, 15 (1919).

Prionocera, Ender., nec Lw., Zool. Jahr. XXXII, 28 (1912).

GENOTYPE, *P. nigriceps*, sp. nov., the original species.

nigriceps Wulp, Notes Leyd. Mus. VII, 2, ♂ ♀ (1885).

Id. Tijds. v. Ent. XXVIII, 82, pl. iv, 3, full ins.: Brun., Rec. Ind. Mus. VI, 245 [*Prionota*]: Alex., Proc. U.S. Nat. Mus. XLIX, 181, pl. xlv, 34, wing; pl. xlvii, 50, antenna, part, 51, 52, genitalia ♂ [*Prionota*].

Java; Mt. Salak, Java, 15-v-09 [Bryant and Palmer]; Tjibodas, Mt. Gedé, Java, 20-iv-09 [Bryant and Palmer].

Type in Leyden Museum. Alexander (*loc. cit.*, *antea*) says *Prionota* and *Prionocera* (*Stygeropsis*, Lw.) are distinct. Kieffer uses *Prionota* again in Cecidomyidae in 1894. Edwards recently states that *Ctenophora xanthomelana*, Walk., is probably identical with *nigriceps*.

serraticornis Brun., Rec. Ind. Mus. VI, 242, ? ♂ (*Pselliophora*) (1911).

Id., Fauna Brit. Ind., 296, ? ♂.

Ceylon, 1861 [Felder].

Type, a unique specimen with tip of abdomen missing, in Vienna Museum. Edwards recently refers my species here.

flaviceps Ender., Zool. Jahr. XXXII, 28, ♀ (*Prionocera*) (1912).
Sumatra.

Unique *type* in Stettin Zoological Museum.

elongata Edw., Ann. Mag. Nat. Hist. (8) XII, 202, ♀ (? *Pselliophora*) (1913).

Hakgala, Ceylon, 4500 ft., v-1911.

Unique *type* in British Museum.

Section TIPULINI.

Ctenacroscelis, Ender.

Zool. Jahr. XXXII, 1 (1912).

GENOTYPE, *C. dohrnianus* Ender., sp. nov.

Ctenacroscelis, Brun., Rec. Ind. Mus. XV, 261.

praepotens Wied., Auss. Zweifl. I, 40 (*Tipula*) (1828).

Tipula, *id.*, Macq., Dipt. Exot., Supp. 3, 7, pl. i, 1: Ost. Sack.,

Ann. Mus. Gen. XVI, 399: *id.*, Berl. Ent. Zeits. XXX, 183:

Brun., Rec. Ind. Mus. VI, 246.

Ctenacroscelis praepotens, Alex., Proc. U.S. Nat. Mus. XLIX, 182, pl. xlv, 35, wing.

East Indies; Java; Sumatra; Amboina; Mt. Salak, 15-v-09; Pelaboean Ratoe [both Java and *Bryant and Palmer*].

Type in Leyden Museum, Wiedemann quoting no sex, but both sexes are present among the old specimens in that collection.

monochrous Wied., Auss. Zweifl. I, 41, ♀ (*Tipula*) (1828).

Alex., Proc. U.S. Nat. Mus. XLIX, 182, pl. xlv, 36, wing; xlvii, 53, 54, genitalia ♂.

Tipula monochroa, Macq., Dipt. Exot., Supp. 3, 6, pl. i, 2, wing:

Brun., Rec. Ind. Mus. VI, 246.

Gunung Ungaran, Java, v [*Leeuwen*]; Mt. Salak, 15-v-09; Pelaboean Ratoe, x, 09 [both Java and *Bryant and Palmer*]; Sumatra; Celebes.

Type in Leyden Museum. Wiedemann mentions only the ♀, but both sexes are present in the old collection.

umbrinus Wied., Auss. Zweifl. I, 49, ♂ (*Tipula*) (1828).

Alex., Proc. U.S. Nat. Mus. XLIX, 182, pl. xlv, 37, wing; xlviii, 55, 56, genitalia ♂.

Tipula umbrina, Wulp, Tijd. v. Ent. XXIII, 158: *id.*, Notes

Leyd. Mus. VII, 4: Ost. Sack., Ann. Mus. Gen. XVI, 399:

Brun., Rec. Ind. Mus. VI, 246: Edw., Jour. Fed. Malay Sts.

Mus. VIII, 11.

Tipula castanea, Macq., Dipt. Exot. I, 1, 54, ♂ (1838).

Tipula congruens, Walk., Proc. Linn. Soc. Lond. V, 231, ♀ (1861).

East Indies; Sumatra; Sinabang (Simalur Is. near Sumatra), iii, v, vii; Celebes; Obi Island; Dorey; Papua; Waigou, i; Java, var. locs., i, iv, v, x [*t. de Meij.*]; Tjibodas, Mt. Gede, 20-iv-09; Mt. Salak, 15-v-09; Depok, 10-ii-09 [all Java and *Bryant and Palmer*]; Siolak Daras, Sumatra, 3000 ft.; Sungei Penoh, Korinchi Valley, Sumatra, 2600 ft., iii, 14.

Type in Leyden Museum. Edwards recently doubts the identity of *castanea*, Macq., with this species.

infidens Walk., Proc. Linn. Soc. IV, 92, ♀ (*Tipula*) (1860).

Brun., Rec. Ind. Mus. VI, 247.

Ceram, Macassar.

Type in British Museum in fair condition. Walker described it as a ♀ from Makassar, but the specimen labelled "type" is a ♂ from Ceram. Some error seems to have crept in here, which probably cannot now be rectified. I have to thank Mr. Edwards and Mr. Hill for these notes.

majesticus Brun., Rec. Ind. Mus. VI, 248, ♂ ♀ (*Tipula*) (1911).

Id., Fauna Brit. Ind., 303, ♂ ♀ (*Tipula*).

N.E. Frontier, India, 4000 ft., 15-iii-12 [*Kemp*]; Naini Tal; Darjiling, 3-9-vi-09, *type* ♂ [*Howlett*]; Singla, vi-12; Tindharia, 3000 ft., viii-12 [both Darjiling Distr. and Lord Carmichael collection]; Kalimpong, 600-4500 ft. [*Gravely*]; Sureil, iv, v-17 [*Kemp*]; Pashok, 26-v-14-vi-16 [*Gravely*]; Darjiling District, 1000-3000 ft. [Lord Carmichael coll.]; Kurseong, 24-vi-10, *type* ♀ [*Annandale*]; Nilgiri Hills, 3500 ft., ix-10 [*Andrewes*]; Talewadi, N. Kanara Dist. [*Kemp*]; between Kalek and Misshing, 4000 ft., Abor Exped. [*Kemp*]; Mazbat, Mangaldai, Assam [*Kemp*]; Ginting Bidai, 2000 ft., Selangor-Pahang Frontier, Fed. Malay States, iv-17 [*Boden Kloss*].

Type ♂ in Pusa coll., *type* ♀ in Indian Museum. A specimen of *majestica* in the Indian Museum collection measures almost exactly 100 mm. in full expanse of wings which easily surpasses both my *T. carmichaeli* and *Cten. rex*, Alex.

fulvolateralis Brun., Rec. Ind. Mus. VI, 249, ♂ (*Tipula*) (1911).

Id., Fauna Brit. Ind., 304, ♂ ♀, pl. vi, 15, genitalia (*Tipula*); *id.*, Rec. Ind. Mus. XV, 263 (*Tipula*).

Himalayas, common in most localities from 600 ft. upwards, March to October; Mussoorie, ix-06; Bhim Tal, 4500 ft., 19-22-ix-06 [*Annandale*]; Sikkim; Sureil; Kalimpong; Singla; Pashok; Soom; Kurseong; Dhikala, Gahrwal Distr., W. Himalayas, 10-iii-10; Shillong, 5500-6400, 29-viii-5-ix-15 [*Kemp*]; Mazbat, Mangaldai Distr., Assam, 11-15-x-10, *type* ♀ [*Kemp*].

Type ♂ and ♀ in Indian Museum. I have stated my *T. fulvolateralis* to be a true *Tipula*, but at the time of writing I had no specimens for comparison and wrote from memory, but a closer inspection of several specimens proves the species to belong here. Alexander thinks my *fulvolateralis* may be synonymous with *umbrinus*, Wied.

flavus, *id.*, Rec. Ind. Mus. VI, 252, ♂ (*Tipula*) (1911).

Tipula, *id.*, Fauna Brit. Ind., 308, ♂: Rec. Ind. Mus. IX, 257, ♀ noted.

Sikkim, iii or iv, *type* ♂ [*Fruhstorfer*].

Type in Vienna Museum. The specimens referred to this species by me recently (Rec. Ind. Mus. IX, 257) are *flavoides*, Brun., the *type* ♂ being the only example of *flavus* known to me

- ornatithorax**, *id.*, Rec. Ind. Mus. VI, 258 ♂ ♀ (*Tipula*) (1911).
Id., Fauna Brit. Ind., 331, ♂ ♀ (*Tipula*).
 Darjiling, 3-9-vi-09, *type* ♂ [*Howlett*]; 21-29-v-06 [*Annandale*]; Kurseong, 5000 ft. [*Annandale*]; Pashok, 2000-2500 ft. [*Gravely*]; Bhowali, Kumaon 5700 ft., vii-09, *type* ♀, and 10-vii-10 [*Imms*]. Recorded doubtfully from Sumatra.
Type ♂ in Pusa coll., *type* ♀ in Indian Museum.
- ochripes**, *id.*, Rec. Ind. Mus. VI, 260, ♂ ♀ (*Tipula*) (1911).
Id., Fauna Brit. Ind., 334, ♂ ♀ (*Tipula*).
 Pashok Spur, Darjiling District [*Lister*]; Castle Rock, N. Kanara Dist., 11--26-x-16 [*Kemp*]; Nilgiris, S. India, 6000 ft. [*Andrewes*]; Kandy, 20-v-10, *type* ♂ [*Gravely*]; 31-x-09, *type* ♀ [*Green*]; Man-San, N. Hsenwi, Shan States, Upper Burma, 2400 ft., ii to iv-15 [*Coggin-Brown*].
Type ♂ ♀ in Indian Museum; *cotype* in Vienna Museum from Ceylon.
- dives** Brun., Fauna Brit. Ind. 307, ♂ (*Tipula*) (1912).
 Kurseong, 23-vi-10 [*Annandale*]; Soom, Darjiling District, 5000-7000 ft., 14-vi-14 [*Gravely*].
Type in Indian Museum.
- carmichaeli** Brun., Rec. Ind. Mus. IX, 257, ♂; pl. xiv, 6, full ins., 9, genitalia (*Tipula*) (1913).
 Sureil, Darjiling District, vi-13; captured by Lord Carmichael in a bath-room.
Type in Indian Museum. A second specimen was recently seen by me, in the possession of Capt. Balfour.
- borneensis** Brun., Rec. Ind. Mus. XV, 263, *nom. nov.* for *pallida*, Walk., *procc. Lw.*, 1863 (*Tipula*) (1918).
Tipula pallida, Walk. nec Lw., Proc. Linn. Soc. Lond. VIII, 105 (1865).
 Brun., Rec. Ind. Mus. VI, 247; de Meij., Nova Guin. Res. IX, 311. Papua, Riverkamp, ii; Alkmaar, x; Regen Is., ix, x (all Papua).
Type in good condition in British Museum.
- flavoides**, *id.*, *loc. cit.*, 267, ♂ ♀; pl. viii, 5, antenna (as "sp. nov." only, *lapsus*) (*Tipula*) (1918).
 Singla, Darjiling District, 1500 ft., v-13; vi-12; Shoirakund, Assam-Bhutan Frontier, 18--22-x-12; Darjiling, 1000-3000 ft., vi-12; above Tura, Garo Hills, 3500-3900 ft., ix-17 [*Kemp*]; Sidapur, Coorg, S. Ind., 16-v-17.
Types in Indian Museum.
- dohrnianus** Enderl., Zool. Jahr. XXXI, 1, ♀, fig. A, wing (1912).
 Soekaranda, Sumatra.
Type, a unique ♀, in Stettin Zoolog. Museum.
- sikkimensis**, *id.*, *loc. cit.*, 4, ♂ (1912).
 Darjiling.
Type, a unique ♂, in Stettin Zoolog. Mus.

- sumatranus**, *id.*, *loc. cit.*, 5, ♀ (1912).
Soekaranda, Sumatra.
Type, a unique ♀, in Stettin Zoological Museum.
- rex** Alex., Insec. Insit. Menst. V, 21, ♂ (1917).
Brun., Rec. Ind. Mus. XV, 261.
Taungoo Distr., Burma.
Type in Entomological collection, Philadelphia, *cotype* in Alexander's coll. *Rex* and my *Tipula carmichaeli* are probably amongst the largest species of diptera known.
- penumbrinus** Edw., Jour. Fed. Malay Sts. Mus. VIII, 11, ♀ (1919).
Siolak Daras, Sumatra, 3000 ft.

Tipula, L.

Syst. Nat. Ed. X, 585 (1758).

Sch., F. Austr., II, 509: de Meij., Tijd. v. Ent. LIV, 63, tab. Java sp.: Brun., Rec. Ind. Mus. VI, 246; XV, 262, notes Orient. spp.: *id.*, Fauna Brit. Ind., 297.

GENOTYPE, *T. oleracea* L., by universal acceptance.

Tipulodina, Ender., Zool. Jahr. XXXII, 30 (1912): Brun., Rec. Ind. Mus. XV, 270, 322, criticism.
GENOTYPE, *Tipula magnicornis* Ender., sp. nov., by original designation.

- pedata** Wied., Dipt. Exot. I, 23, ♀ (1821).
Id., Auss. Zweifl. I, 45, ♀: Wulp., Dipt. Sum. Exped., 10: Ost. Sack., Berl. Ent. Zeits. XXVI, 92: de Meij., Tijd. v. Ent. LIV, 64: Brun., Rec. Ind. Mus. VI, 246: Alex., Proc. U.S. Nat. Mus. XLIX, 183, pl. xlv, 38, wing; XLVIII, 57—60, genitalia ♂.
Java, var. locs., i, xi, xii; Buitenzorg, iii, 09; 10-iv-09; Pelaboean Ratoe, 12-x-09; Mt. Salak, 3000 ft., 15-v-09 [all three Java and *Bryant* and *Palmer*]; Sumatra; Borneo; Philippine Is.; Assam; Ceylon.
Type, a ♀, in Westermann's collection.
- venusta** Walk., List Dipt. Brit. Mus. I, 64 (1848).
Brun., Rec. Ind. Mus. VI, 246: *id.*, Fauna Brit. Ind., 302, pl. v, 5, wing.
Sylhet; Siliguri; Calcutta [*Targett*]; Myitkyina, Upper Burma, 30-viii—1-ix-14 [*Fletcher*]; Tura, Garo Hills, Assam, viii-17 [*Kemp*].
Type recorded originally as in the British Museum, sex unstated, but now no longer to be found there.
- reposita** Walk., List Dipt. Brit. Mus. I, 67, ♂ ♀ (1848).
Brun., Rec. Ind. Mus. VI, 246: Fauna, 315.
Nepal.

Type ♂ ♀ in British Museum, ♂ in fair condition, ♀ reduced to a fragment.

melanomera Walk., List Dipt. Brit. Mus. I, 68, ♂ (1848).

Brun., Rec. Ind. Mus. VI, 247; XV, 263: Brun., Fauna Brit. Ind., 330.

Sureil, 5000 ft., iv—v-17 [*Kemp*]; Maymyo, Upper Burma [*Fletcher*]; Nepal.

Type in British Museum.

vicaria Walk., Ins. Saund. Dipt., pt. V, 444, ♂ (1856).

Brun., Rec. Ind. Mus. VI, 247: Fauna Brit. Ind., 332.

India, according to the *type*. Walker's description says East Indies.

Type in British Museum, too damaged for comparison. The name *vicaria* is preoccupied by Walker himself in 1848 for a South African species. The present species appears to be represented by the original specimen only, which is useless. If subsequently identified from the description a new name can be set up.

vilis Walk., Proc. Linn. Soc. I, 108, ♂ (1857).

Brun., Rec. Ind. Mus. VI, 247: de Meij., Tijd. v. Ent. LIV, 69 (♀ descr.).

Sarawak, *type* ♂; Buitenzorg, iii; Gunung Ungaran, xii [both Java and *Jacobson*].

Type in British Museum, beyond recognition.

inordinans Walk., Proc. Linn. Soc. IV, 93, ♂ (1860).

Brun., Rec. Ind. Mus. VI, 247.

Macassar.

Type, a unique ♂, in British Museum, in good condition.

fumifinis Walk., *loc. cit.*, V, 145, ♂ (1861).

Brun., Rec. Ind. Mus. VI, 247, notes.

Amboina.

Type, a unique ♂, in British Museum, in good condition. Walker thought his species identical with *longicornis*, Dol., but Van der Wulp considered them distinct.

punctifrons Rond., Ann. Mus. Gen. VII, 463 (1875).

Brun., Rec. Ind. Mus. VI, 247.

Sarawak. Location of *type* uncertain, said to be in Genoa Museum.

serrata Wulp, Notes Leyd. Mus. VII, 5, ♂ (1885).

Brun., Rec. Ind. Mus. VI, 247.

Seharan, Sumatra.

Type in Leyden Museum, also a ♀ from Sumatra in same collection.

pilosula Wulp, Notes Leyd. Mus. VII, 5, ♂ (1885).

de Meij., Tijd. v. Ent. LIV, 67: Brun., Rec. Ind. Mus. VI, 247.

Java; Gunung Kenepai; Borneo.

Type in Leyden Museum.

- leucopyga** Wulp. Notes Leyd. Mus. VII, 6, ♂ ♀ (1885).
Id., Tijd. v. Ent. XXVIII, 84, pl. iv, 6: Brun., Rec. Ind. Mus. VI, 247.
 Java; Depok, Java, ii [*Jacobson*]
Types ♂ ♀ in Leyden Museum.
- thibetana** de Meij., Bijd. tot de Dierk. XVII, 89, pl. viii, 2, 3, wing (1904).
 Brun., Rec. Ind. Mus. VI, 248.
 Tatsienlou, Thibet.
Type in Amsterdam Museum. Perhaps incorrectly included in this Catalogue but may occur in the Himalayas.
- omissinervis** de Meij., Nova Guin. Res. V, 71, fig. 4, wing, part (*Tanypremna*) (1906).
Id., loc. cit., IX, 312, note: Brun., Rec. Ind. Mus. VI, 268.
 Wendesi, Papua, vi; Regen Is., Papua, ix-x-09; Waigou, i.
Type in Leyden Museum, *cotype* in Amsterdam Museum.
- cinctipes** de Meij., Tijd. v. Ent. LIV, 64, ♂ (1911).
 Gunung Kenepai (Borneo, Pondok).
Type, a unique ♂, in Leyden Museum.
- gedehana**, *id.*, loc. cit., 66, ♀ pl. iv, 47, wing (1911).
 Alex., Proc. U.S. Nat. Mus. XLIX, 185, pl. xlix, 63, 64, genitalia ♂.
 Kraton des Gunung, Gedeh, Java [*Buttiker*]; Tjibodas, Mt. Gede, Java, 4000, 8000, 9000 ft., ix-09 [*Bryant and Palmer*].
Type in Leyden Museum.
- cinereifrons**, *id.*, loc. cit., 68, ♂, pl. iv, 48, wing (1911).
 Java [*t. de Meijere*]; Sumatra [*t. Enderlein*].
Type in Leyden Museum.
- inconspicua**, *id.*, loc. cit., 70, ♂, pl. iv, 49, wing (1911).
 Wonosobo, Java, iv [*Jacobson*].
Type in Amsterdam Museum, a unique ♂.
- walkeri** Brun., Rec. Ind. Mus. VI, 246, *nom. nov.* for (and notes on) *fulvipennis*, Walk., preocc. De Geer (1911).
Tipula fulvipennis, Walk., List. Dipt. Brit. Mus. I, 67 (1848).
 Nepal.
Type in British Museum, sex unstated, valueless for comparison. De Geer's *fulvipennis*, a European species, was set up in 1776 and is synonymous with *lutescens*, F.
- schummeli** Brun., Rec. Ind. Mus. VI, 247, *nom. nov.* for *longicornis*, Dol. (1911).
longicornis, Dol., Nat. Tijd. Ned. Ind. XVII, 79 (1858).
 Amboina.
Type in Vienna Museum; no sex stated. The name *longicornis*, preoccupied by Schummel in 1833 for a European species, and also used for some other species subsequently.
- fumifasciata** Brun., Rec. Ind. Mus. VI, 250, ♂ ♀ (1911).
Id., Fauna Brit. Ind., 308, ♂ ♀, pl. v, 7, wing.

Ukhral, Manipur, 6400 ft., viii 18, *types* [Pettigrew]; Central China.

Type ♂ and ♀ in Indian Museum, *cotypes* in Vienna Museum.

fumipennis, *id.*, Rec. Ind. Mus. VI, 250, ♀ (1911).

Id., Fauna, Brit. Ind., 305, ♀, pl. v, 6, wing.

Darjiling, 3-9-vi-09 [Howlett], *type*; 6000-7000 ft. [Gravelly]; 7000 ft. [Brunetti]; Mussoorie, vi-09.

Type in Pusa collection.

pluto, *id.*, Rec. Ind. Mus. VI, 251, ♀ (1911).

Tonkin (Montes Maussion, 2000-3000 ft., iv or v) [Fruhstorfer].

Type, a unique ♀, in Vienna Museum.

cinerea, *id.*, *loc. cit.*, 252, ♀ (1911).

Id., *loc. cit.*, XV, 274.

Lombok, 2000 ft., v or vi-1896 [Fruhstorfer].

Type in Vienna Mus., a unique ♀.

sciariformis, *id.*, Rec. Ind. Mus. VI, 259, ♀ (1911).

Tonkin [Fruhstorfer].

The unique ♀ *type* in Vienna Museum.

demarcata, *id.*, *loc. cit.*, 259, ♀ (1911).

Id., Fauna Brit. Ind., 335, ♀.

Kurseong, 4700-5000 ft [Annandale]; Almora, Kumaon, 5000 ft. [Paiva]; Kandy, v-1910, *type* [Green]; Peradeniya, 25-xii-01 [Dr. Uzel].

Type in Indian Museum, *cotype* from Peradeniya in Vienna Museum.

himalayensis, *id.*, Rec. Ind. Mus. VI, 252, ♂ ♀ (1911).

Id., Fauna Brit. Ind., 315, ♂ ♀, pl. v, 11, wing; pl. vi, 16, 17, genitalia: Rec. Ind. Mus. XV, 262, allied spp.

Himalayas, the commonest species, generally distributed at all hill stations from east to west during summer and autumn, at least from Shillong to Mussoorie.

Type ♂ and ♀ in Indian Museum, *cotype* in my collection.

robusta, *id.*, Rec. Ind. Mus. VI, 254, ♂ (1911).

Id., Fauna Brit. Ind., 319, ♂.

Mussoorie, 22-vi-05 [Brunetti].

Type, a unique ♂, in my collection.

elegans, *id.*, Rec. Ind. Mus. VI, 255, ♀ (1911).

Id., Fauna Brit. Ind., 323, ♀: *id.*, Rec. Ind. Mus. XV, 274.

Mussoorie, 18-vi-05 [Brunetti].

Type, a unique ♀, in my collection.

interrupta, *id.*, Rec. Ind. Mus. VI, 256, ♀ (1911).

Id., Fauna Brit. Ind., 327, ♀.

Darjiling, 3-9-vi-09 [Howlett].

Type, a unique ♀, in Pusa coll.

nigroapicalis, *id.*, Rec. Ind. Mus. VI, 257, ♂ ♀ (1911).

Id., Fauna Brit. Ind., 328, ♂ ♀.

- Darjiling, 16—20-x-05, *type* ♂ ♀, 10—16-x-05 [both *Brunetti*].
Type ♂ ♀ and other specimens in my collection.
- divisa*, *id.*, Rec. Ind. Mus. VI, 261, ♂ (1911).
Id., Fauna Brit. Ind., 337, ♂.
 Darjiling, 3--9-vi-09 [*Howlett*].
Type, a unique ♂, in Pusa collection.
- gracilis*, *id.*, Rec. Ind. Mus. VI, 262, ♀ (1911).
Id., Fauna Brit. Ind., 338, ♀.
 Darjiling, 7-x-05.
Type, a unique ♀, in my collection.
- gracillima* Brun., Fauna Brit. Ind., 302, ♀ (1912).
 Edw., Ann. Mag. Nat. Hist. (8) XII, 202 (1913): Brun., Rec. Ind. Mus. XV, 262.
 Peradeniya, 20-vii-10, *type* [*Gravelly*]: Cherambadi, Wynaad, S. India, x-13 [*Howlett*]; Parambikulam, Cochin, 1700—3200 ft., 16—24-ix-14 [*Gravelly*].
Type in Indian Museum.
- princeps* Brun., Fauna Brit. Ind., 306, ♂ ♀ (1912).
 Kurseong, 18—19-vi-10, at light [*Annandale*]; Darjiling, 7000 ft.: 1—10-vi-17 [*Brunetti*]; Soom, Darjiling District, 4000—5000 ft., 16-vi-14 [*Gravelly*].
Types in Indian Museum.
- serricornis*, *id.*, *loc. cit.*, 309, ♂ ♀ (1912).
 Western Himalayas, Naini Tal.
Types in Indian Museum.
- pulcherrima*, *id.*, *loc. cit.*, 310, ♂ ♀; pl. v, 8, wing; pl. vi, 12, thorax, 13, genitalia (1912).
 Darjiling, 3—9-vi-09 [*Howlett*]; v-13; Tura, Garo Hills, Assam, vii-17 [*Kemp*]; Bhowali, 5700 ft., 27-vi-10, *type* ♂, at light; vii-09, *type* ♀, 26-vi-10 [all *Imms*].
Types in Indian Museum, *cotype* ♀ in Pusa coll.
- fuscinervis*, *id.*, *loc. cit.*, 312, ♀ pl. v, 9, wing (1912).
 Kurseong, 21—29-v-06 [*Annandale*].
Type in Indian Museum, a unique ♀.
- patricia*, *id.*, *loc. cit.*, 313, ♀ (1912).
 Kurseong, 26-vi-10 [*Annandale*].
Type in Indian Museum, a unique ♀.
- splendens*, *id.*, *loc. cit.*, 314, ♂, pl. v, 10, wing (1912).
 Badrinath, Gahrwal Distr., 21-v-10 [*Imms*].
Type, a unique ♂, in Indian Museum.
- tessellatipennis*, *id.*, *loc. cit.*, 317, ♂ ♀, pl. v, 13, wing (1912).
 Naini Tal, 6400 ft., v or vi, 1893.
Types in Indian Museum, a unique pair.
- marmoratipennis*, *id.*, *loc. cit.*, 319, ♀, pl. v, 13, wing (1912).
 Darjiling, 24-iv-10, *type* [*Brunetti*]; 10-vi-17, 7-viii-09 [*Paiva*].
Type in Indian Museum.

- Darjiling, 16-20-x-05, *type* ♂ ♀, 10-16-x-05 [both *Brunetti*].
Type ♂ ♀ and other specimens in my collection.
- divisa*, *id.*, Rec. Ind. Mus. VI, 261, ♂ (1911).
Id., Fauna Brit. Ind., 337, ♂.
 Darjiling, 3-9-vi-09 [*Howlett*].
Type, a unique ♂, in Pusa collection.
- gracilis*, *id.*, Rec. Ind. Mus. VI, 262, ♀ (1911).
Id., Fauna Brit. Ind., 338, ♀.
 Darjiling, 7-x-05.
Type, a unique ♀, in my collection.
- gracillima* Brun., Fauna Brit. Ind., 302, ♀ (1912).
 Edw., Ann. Mag. Nat. Hist. (8) XII, 202 (1913): Brun., Rec. Ind. Mus. XV, 262.
 Peradeniya, 20-vii-10, *type* [*Gravelly*]: Cherambadi, Wynaad, S. India, x-13 [*Howlett*]; Parambikulam, Cochin, 1700-3200 ft., 16-24-ix-14 [*Gravelly*].
Type in Indian Museum.
- princeps* Brun., Fauna Brit. Ind., 306, ♂ ♀ (1912).
 Kurseong, 18-19-vi-10, at light [*Annandale*]; Darjiling, 7000 ft.: 1-10-vi-17 [*Brunetti*]; Soom, Darjiling District, 4000-5000 ft., 16-vi-14 [*Gravelly*].
Types in Indian Museum.
- serricornis*, *id.*, *loc. cit.*, 309, ♂ ♀ (1912).
 Western Himalayas, Naini Tal.
Types in Indian Museum.
- pulcherrima*, *id.*, *loc. cit.*, 310, ♂ ♀; pl. v, 8, wing; pl. vi, 12, thorax, 13, genitalia (1912).
 Darjiling, 3-9-vi-09 [*Howlett*]; v-13; Tura, Garo Hills, Assam, vii-17 [*Kemp*]; Bhowali, 5700 ft., 27-vi-10, *type* ♂, at light; vii-09, *type* ♀, 26-vi-10 [all *Imms*].
Types in Indian Museum, *cotype* ♀ in Pusa coll.
- fuscinervis*, *id.*, *loc. cit.*, 312, ♀ pl. v, 9, wing (1912).
 Kurseong, 21-29-v-06 [*Annandale*].
Type in Indian Museum, a unique ♀.
- patricia*, *id.*, *loc. cit.*, 313, ♀ (1912).
 Kurseong, 26-vi-10 [*Annandale*].
Type in Indian Museum, a unique ♀.
- splendens*, *id.*, *loc. cit.*, 314, ♂, pl. v, 10, wing (1912).
 Badrinath, Gahrwal Distr., 21-v-10 [*Imms*].
Type, a unique ♂, in Indian Museum.
- tessellatipennis*, *id.*, *loc. cit.*, 317, ♂ ♀, pl. v, 13, wing (1912).
 Naini Tal, 6400 ft., v or vi, 1893.
Types in Indian Museum, a unique pair.
- marmoratipennis*, *id.*, *loc. cit.*, 319, ♀, pl. v, 13, wing (1912).
 Darjiling, 24-iv-10, *type* [*Brunetti*]; 10-vi-17, 7-viii-09 [*Paiva*].
Type in Indian Museum.

- quasimarmoratipennis*, *id.*, *loc. cit.*, 320, ♀, pl. v, 14, wing (1912).
Darjiling, 23-v-10, *type* [Brunetti]; Kurseong, 9-ix-09 [Annandale]; Ukhrul, Manipur, 6400 ft. [Pettigrew]; Tura, Garo Hills, Assam, ix-17 [Kemp]. *Type* in Indian Museum.
- griseipennis*, *id.*, *loc. cit.*, 321, ♂, pl. v, 15, wing (1912).
Badrinath, Gahrwal Distr., 27-v-10 [Imms]; Matiana [Annandale]. *Type* in Indian Museum.
- nigrotibialis*, *id.*, *loc. cit.*, 324, ♂ (1912).
Darjiling, 28-v-10 [Brunetti].
Type in Indian Museum.
- striatipennis*, *id.*, *loc. cit.*, 325, ♀, pl. v, 16, wing (1912).
Kurseong, 6-vii-08 [Annandale]. *Unique type* in Indian Museum.
- subtineta*, *id.*, *loc. cit.*, 326, ♀, pl. vi, 18, genitalia (1912).
Darjiling, 29-v-17; 2 and 8-vi-12 (all Brunetti); Kurseong, 7-ix-09 [Annandale]; Bhowali, 5-700 ft. [Imms]; Maupun, Khasi Hills [Warren]. *Type* in Indian Museum.
- continuata*, *id.*, *loc. cit.*, 328, ♂ ♀ (1912).
Darjiling, 25—29-v-10, *type* ♂ ♀; 16—20-x-05 [both Brunetti]; Chitral, N.W. Kashmir, 9500 ft. [Wall].
Types in Indian Museum.
- quadrinotata*, *id.*, *loc. cit.*, 330, ♀ (1912).
Darjiling [Paiva]; Ukhrul, Manipur, 6400 ft. [Pettigrew].
Type in Indian Museum.
- brunnicauda*, *id.*, *loc. cit.*, 332, ♂ ♀ (1912).
Simla, 7000 ft.; 11-v-08 [Annandale]; Theog, 8000 ft. [Annandale] Kalighat, Gahrwal Distr., 6000 ft., 4-vi-10 [Imms].
Types in Indian Museum.
- tenuipes*, *id.*, *loc. cit.*, 333, ♀ (1912).
Sylhet [Hall]. *Unique type* in Indian Museum.
- flavescens*, *id.*, *loc. cit.*, 334, ♀ (1912).
Hakgalla, Ceylon, 29-ix-09 [Green].
Unique type in Indian Museum.
- munda*, *id.*, *loc. cit.*, 336, ♂ ♀ (1912).
Mussoorie, 18-vi-09; Darjiling, 23—29-v-10; 10-vi-17 [all Brunetti].
Type ♂ ♀ in Indian Museum, *cotype* ♂ ♂ in my collection.
- cinctoterminalis*, *id.*, *loc. cit.*, 338, ♀ (1912).
Kurseong, 7-ix-09, *type* [Annandale]; Bhowali, 15-vi-10 [Imms].
Type in Indian Museum.
- elegantula*, *id.*, *loc. cit.*, 339, ♂; pl. v, 17, wing (1912).
Almora, Kumaon, 5500 ft. [Paiva]; Castle Rock, N. Kanara Dist., ? ♀ [Kemp]; Shevaroy Hills, Vercaud, S. Ind., 4500 ft.

- 17—21-xii-13 [*Fletcher*]; Mazbat, Assam, 11—15-x-10 [*Kemp*]; Cherrapunji, Assam, 4400 ft., 2—8-x-14, incl. *type* [*Kemp*]; Shillong, x-16 [*Fletcher*]; 5500—6400 ft., 29-viii—5-ix-15 [*Kemp*]. *Type* in Indian Museum.
- coquilletti** Ender., Zool. Jahr., XXII, 7, *nem. nov.* for *nubifera* Coq.
nec. *nubifera*, Wulp (1912).
nubifera, Coq., Proc. U.S. Nat. Mus. XXI, 305 (1898).
Arisan, Formosa, 8000 ft., 10-x-12 [*Nitobe*]; also occurs in Japan.
- imperfecta**, *id.*, *loc. cit.*, 260, ♀ (1913).
Darjiling, 1000—3000 ft., iv-1912.
Type, a unique ♀ in fair condition, in Indian Museum.
- tropica** de Meij., Nova Guin. Res. IX, 311, ♀ (1913).
Regen Is., Papua, ix, x.
Type unique, in Amsterdam Museum.
- novae guineae**, *id.*, *loc. cit.*, 312, ♂ ♀ (1913).
Lorentz-Fluss, S.W. Papua, ix-09.
Types, a unique pair, in Amsterdam Museum.
- divergens**, *id.*, *loc. cit.*, 312, ♂ ♀ pl. x, 6, wing; 7, ♂ genitalia (1913).
Bivak Is., Papua, xii-09, i-10. *Types* in Amsterdam Museum.
- dentata**, *id.*, *loc. cit.*, 313, ♂ ♀ (1913).
Regen Is., Papua, 16-ii-10.
Types, a unique pair, in Amsterdam Museum.
- de meijerei** Edw., Trans. Zool. Soc. Lond. XX, 393, ♀, pl. xxxviii, 1, full ins. col. (1915).
Upper Utakwa Valley, 5000—10,000 ft., Dutch Papua, ii, iii-13, *type*.
- aetherea** de Meij., Tijd. v. Ent. LVIII, 7, ♀ (1915).
Laulo, viii; Sinabang vii, both Simalur Island, near Sumatra.
Type in Amsterdam Museum.
- sinabangensis**, *id.*, *loc. cit.*, 9, ♂ ♀ (1915).
Sinabang, Simalur Island, ii, near Sumatra.
Type in Amsterdam Museum.
- pumila**, *id.*, *loc. cit.*, 9, ♂ (1915).
Sinabang, Simalur Island, ii, near Sumatra.
Type in Amsterdam Museum.
- klossi** Edw., Ann. Mag. Nat. Hist. (8) XVII, 351, ♀ (1916).
Kedah Peak, 3200 ft., Malay Peninsula [*Stanton*].
Type in British Museum, *paratype* in Kuala Lumpur Museum.
- shirakii** Edw., *loc. cit.*, XVIII, 258, ♀ (1916).
Arisan, Formosa, 8000 ft., 10-x-12 [*Nitobe*]; Khasi Hills, Assam. *Type* in British Museum.

- rufomedia*, *id.*, *loc. cit.*, p. 259, ♂ ♀, pl. xii, 5-7, genitalia (1916).
Horisha, Formosa [*Maki*]; "Formosa" [Brit. Mus. coll.].
Type in British Museum.
- umbrinoides* Alex., Proc. U.S. Nat. Mus. XLIX, 185, ♂, pl. xlvi, 39, wing; xlvii, 61, 62, ♂ genitalia (1916).
Mt. Salak, Java, 3000 ft., 4-vii-09 [*Bryant and Palmer*].
Type, No. 19,027, in U.S. Nat. Mus.
- sunda*, *id.*, *loc. cit.*, 186, ♂, pl. xlvi, 40, wing; xlix, 65, 66, ♂ genitalia (1916).
Tjibodas, Mt. Gedé, Java, 9000 ft. [*Bryant and Palmer*].
Type, No. 19,028, in U.S. Nat. Mus.
- flavicosta*, *id.*, *loc. cit.*, 187, pl. xlvi, 41, wing (1916).
Pelaboean-Ratoe, Java, 16-x-09 [*Bryant and Palmer*].
Type, No. 19,029, in U.S. Nat. Mus.
- tjibodensis*, *id.*, *loc. cit.*, 188, ♂ ♀, pl. xlvi, 42, wing; pl. l, 68, 69, genitalia ♂ (1916).
Tjibodas, Mt. Gedé, Java, 9000 ft. [*Bryant and Palmer*].
Type, No. 19,030, in U.S. Nat. Mus.
- salakensis*, *id.*, *loc. cit.*, 189, ♂, pl. xlvi, 43, wing; xlix, 67, genitalia ♂ (1916).
Mt. Salak, Java, 1909 [*Bryant and Palmer*].
Type, No. 19,031, in U.S. Nat. Mus.
- gedericola*, *id.*, *loc. cit.*, 189, ♂, pl. xlvi, 44, wing; l, 70, 71, 72, genitalia ♂ (1916).
Tjibodas, Mt. Gedé, Java, 5000 ft., 20-iv-09.
Type, No. 19,032, in U.S. Nat. Mus.
- gravelyi* Brun., Rec. Ind. Mus. XV, 264, ♂ (1916).
Darjiling, 7000 ft., 13-vi-14 [*Gravely*]; vii-12; E. Himalayas, [*Dracott*]. *Type* in Indian Museum.
- contigua*, *id.*, *loc. cit.*, 265, ♂ (1918).
Above Tura, Garo Hills, Assam, 3500-3900 ft., viii-17 [*Kemp*]. *Unique type* in Indian Museum.
- simillima*, *id.*, *loc. cit.*, 265, ♂ (1918).
Castle Rock, N. Kanara Distr., 11-26-x-16 [1918].
Type in Indian Museum.
- fumifascipennis*, *id.*, *loc. cit.*, 266, ♂ ♀ (1918).
Darjiling, 6000-7000 ft., 11-vi-14, *type* ♂, 12-vi-14, *type* ♀ [both *Gravely*]; 28-v-10 [*Brunetti*].
Types in Indian Museum.
- fumicosta*, *id.*, *loc. cit.*, 266, ♂ (1918).
Shillong, 5500-6400 ft., 28-viii-5-ix-15, *type* [*Kemp*], 4900 ft., 10-12-x-14 [*Kemp*]; Tura, Garo Hills, x-17 [*Kemp*].
Type in Indian Museum.

filicornis, *id.*, *loc. cit.*, 267, ♂ (1918).

Pashok, Darjiling District, 3000 ft., 26-v-14-vi-16 [*Gravelly*].
The unique *type* in the Indian Museum.

rufiventris, *id.*, *loc. cit.*, 268, ♀ (1918).

Pashok, Darjiling Distr., vi-16 [*Hartless*] *type* ♂; Soom,
Darjiling Dist., 3000—3500 ft., 15-vi-14, *type* ♀ [*Gravelly*]; above
Tura, 3500—3900 ft., ix-17 [*Kemp*].
Types in Indian Museum.

flavithorax, *id.*, *loc. cit.*, 268, ♂ (1918).

Parambikulam, Cochin State, S. India, 1700—3200 ft., 16—
24-ix-14 [*Gravelly*]. The unique *type* in the Indian Museum.

fasciculata, *id.*, *loc. cit.*, 269, ♂ ♀ (1918).

Darjiling, 7000 ft., 25-v-10, *type* ♂ [*Brunetti*]; Tonglu, Dar-
jiling District, 10,000 ft., 21-iv-10 [*Beebe*]; Senchal, Darjiling
District, 8000 ft., v-13 [Lord Carmichael coll.], also *type* ♀ from
same locality. *Types* in Indian Museum.

brevis, *id.*, *loc. cit.*, 270, ♂ ♀ (1918).

Shillong, 4900 ft., 10-12-x-14 [*Kemp*].
Type in Indian Museum.

jacobsoni Edw., Jour. Fed. Malay Sts. Mus., VIII, 12, ♀ pl. iv, 9,
wing (1919).

? *Tipula* sp. de Meij., Tijd. v. Ent. LIV, 71 (1911).
Korinchi Peak, Sumatra, 7300 ft. v-1914, a unique ♀.

korinchiensis Edw., Jour. Fed. Malay Sts. Mus. VIII, 13, ♀
(1919).

Sungei Kumbang, Korinchi, 4500 ft., Sumatra, iv-1914.

robinsoni Edw., *loc. cit.* 14, ♂ ♀ (1919) pl. iii, 3 a, b, genitalia.
Sandaran Agong, Korinchi, 2450 ft., iv-v-1914, a single pair
in cop.

Longurio, Lw.

Berl. Ent. Zeits. XIII, 3 (1869).

GENOTYPE, *L. testaceus*, *id.*, sp. nov., *loc. cit.*

rubriceps Edw., Ann. Mag. Nat. Hist. (8) XVIII, 261, ♂ pl. xii, 9,
genitalia (1916).

Shinten, Formosa, 400 ft. [*Shiraki*].
Unique *type* in British Museum.

Brithura, Edw.

Ann. Mag. Nat. Hist. (8) XVIII, 262 (1916).

GENOTYPE, *B. conifrons* Edw., sp. nov.

conifrons Edw., *loc. cit.*, 263, ♂ pl. xii, 10-11, genitalia (1916).

Arisan, Formosa, 8000 ft., 10-x-12 [*Nitobe*].
Unique *type* in British Museum.

crassa, *id.*, *loc. cit.*, 264, ♂ (1916).
 "East Indies" (? India). Unique *type* in British Museum.

Pachyrhina Macq.

Hist. Nat. Dipt. I, 88 (1834).

Sch., F. Austr. II, 503: de Meij., Tijd. v. Ent. LIV, 72, notes: Brun., Rec. Ind. Mus. VI, 262, notes Orient, spp.: *id.*, Fauna Brit. Ind., 339.
 GENOTYPE, *Tipula crocata* L., by Westwood's designation.

javana Wied., Dipt. Exot. I, 27, ♂ (*Tipula*) (1821).

Id., Auss. Zweifl. I, 57, ♂ (*Tipula*): Brun., Rec. Ind. Mus. VI, 262 (*Pachyrhina*).

Java. *Type* in Westermann's collection. Specimens taken at Coonor by Mr. Howlett, 4 and 5, ix-ii, ♂ ♂, and 1 and 4-ix-ii, ♀ ♀, may be this species.

bombayensis Macq., Dipt. Exot. Supp. V, 35, ♂, pl. i, 1, wing (1854).

Wulp, Tijd. v. Ent. XXIII, 158: *id.*, Mid. Sumat. Exped. Dipt., 11: de Meij., Tijd. v. Ent. LIV, 79: Brun., Rec. Ind. Mus. VI, 263: *id.*, Fauna Brit. Ind., 351, ♂ ♀.

Bombay; Pusa (nearly all the year round), Chapra, Bengal; Katihar, 7—31-viii-10 [*Paiva*]; Damukdia Ghat, N. Bengal; Calcutta [*Anmandale*]; Bandung, Java, vi; Tandjong Priok, Java [both *Jacobson*]. *Type* in Bigot coll.

delta Walk., Ins. Saund., Dipt. pt. v, 445, ♀ (1856).

Brun., Rec. Ind. Mus. VI, 263.

East Indies. *Type* in British Museum.

Van der Wulp quotes East India in his catalogue, but my information from an examination of the *type* is that it is from the East Indies.

javensis Dol. (nec *javana*, W.), Nat. Tijd. Ned. Ind. X, 406 (*Tipula*), pl. iii, 2, full insect, head, profile (1856).

Brun., Fauna Brit. Ind., 348, ♀ redescr.: Edw., Ann. Mag. Nat. Hist. (8) XVIII, 266, notes: Brun., Rec. Ind. Mus. XV, 274.

doleschalli, Ost. Sack., Ann. Mus. Gen. XVI, 399, ♀ (1881): de Meij., Tijd. v. Ent. LIV, 77: Brun., Rec. Ind. Mus. VI, 263, 264, notes: Alex, Proc. U.S. Nat. Mus. 190, pl. xlvi, 45, wing.

fallax, de Meij., Bijd. tot de Dierk. XVII, 90, ♂ ♀ (1904). (t. Meij., Tijd. v. Ent. LIV, 77: 1911).

? *fasciata* Macq., Hist. Nat. Dipt. I, 90, ♂ (1834).

Naini Tal Distr., 3—4-iv-10; Coonor, S. Ind., 1 and 4-ix-11 [*Howlett*]; Ceylon; Tura, Garo Hills, ix, x-17 [*Kemp*]; Djokjakarta; Pangerando, x; Wonosobo, iv [both Java and

- Jacobson*]; Buitenzorg, vi-09; Tjibodas, Mt. Gedé, iv-09
[*Bryant and Palmer*]; Sumatra. Type ♂ in Vienna Museum.
- tripartita* Walk., Proc. Linn. Soc. Lond. V, 231, ♂ (1861).
Papua. Type, location unknown, probably lost.
- tenuis*, *id.*, *loc. cit.*, VIII, 106, ♂ (1865).
Papua. Type, location unknown, probably lost.
- colorata*, *id.*, *loc. cit.*, VIII, 131, ♀ (1865).
de Meij., Tijd. v. Ent. LIV, 77; Brun., Rec. Ind. Mus. VI, 263.
Salawatti; Dutch South Papua (Digul Fluss).
Type, location unknown, probably lost.
De Meijere records a ♀ from Dutch South Papua which is probably of this species.
- familiaris* Ost. Sack, Ann. Mus. Gen. XVI, 401, ♂ ♀ (1880).
de Meij., Tijd. v. Ent. LIV, 74 (augmented description):
Brun., Rec. Ind. Mus. VI, 263.
Sumatra; Pulu Babi Is. off W. Coast Sumatra, iv; Java; Wonosobo, Java, v [*Jacobson*]. Types in Genoa Museum.
- melanura* Ost. Sack., Ann. Mus. Gen. XVI, 401, ♀ (1880).
Papua. Type in Genoa Museum.
- laconica*, *id.*, Berl. Ent. Zeits. XXVI, 92, ♀ (1882).
Philippines. Type, location unknown, a unique specimen.
- ortiva*, *id.*, *loc. cit.*, 93, ♀ (1882).
Brun., Rec. Ind. Mus. VI, 263.
Philippines. Type, location unknown.
- nigro-annulata* Wulp, Notes Leyd. Mus. VII, 8, ♂ ♀ (1885).
Brun., Rec. Ind. Mus. VI, 264.
Morotai. Type ♂ ♀ in Leyden Museum.
- quadrivittata*, *id.*, *loc. cit.*, 9, ♂ (1885).
Java. Type in Leyden Museum.
- triplasia*, *id.*, *loc. cit.*, 10 ♂ ♀ (1885).
Java. Type in Leyden Museum.
- immaculata* Wulp, Tijd. v. Ent. XXXIV, 196, ♂ (1892).
Java [*t. Wulp*]; Tankoeban Prahoe, Java, vi [*Jacobson*].
var. *pangerangensis* Alex., Proc. U.S. Nat. Mus. XLIX, 190, pl. xlvi, 47, wing (1916).
Pangrango, Java, 9000 ft; Tjibodas, Mt. Gedé, Java, 4500 and 9000 ft. [*Bryant and Palmer*].
Type of *immaculata* in Leyden Museum; of *pangerangensis*, No. 19,033, in U.S. Nat. Mus.
- virgata* Coq., Proc. U.S. Nat. Mus. XXI, 306 (1898).
Edw., Ann. Mag. Nat. Hist. (8) XVIII, 266, notes.

Arisan, Formosa, 8000 ft., 10-x-12 [*Nitobe*]. Described originally from Japan.

dorsalis de Meij., *Bijd. tot de Dierk.* XVII, 89, ♀ (1904).

Java. *Type* in Amsterdam Museum.

scurroides, *id.*, *loc. cit.*, 90, ♀ (1904).

Id., *Tijd. v. Ent.* LIV, 75, ♂ (1911).

Java; South Formosa; Wonosobo, Java, iv [*Jacobson*]; Tosari, Java, iii [*Beaufort*]. *Type* in Amsterdam Museum.

dimidiata de Meij., *Bijd. tot de Dierk.* XVII, 90, ♀ (1904).

Id., *Nova Guin. Res.* IX, 315.

Java; Alkmaar, Papua, ii. *Type* in Amsterdam Museum.

dorsopunctata Brun., *Rec. Ind. Mus.* VI, 265, ♂ ♀ (1911).

Id., *Fauna Brit. Ind.*, 350, ♂ ♀ : *Rec. Ind. Mus.* XV, 275.

Naini Tal District, 3-v-10; Pashok, Darjiling District, 2000—3500 ft. [*Gravelly*]; Siliguri [*Annandale*]; Dinapore, 16-xii-14 to 23-i-15 [*Caunter*]; Katihar, 30-xi-09, *type* ♂; Bhogaon, 20-xii-09; Madhupur, 13-x-09 [all Bengal and *Paiva*]; Pusa, 31-iii; 4-iv-11 [*Gravelly*]; 27-xi-14 [*Fletcher*]; Calcutta [*Annandale*]; Maddathorai, Travancore, 17-xi-08, *type* ♀ [*Annandale*]; Ceylon; Assam-Bhutan Frontier, 30-xii-10 [*Kemp*].

Type ♂ ♀ in Indian Museum.

Two ♂♂ in the Pusa collection 16-ii and 26-ii-10 bred from larvae in earth from rice-fields at Pusa may be this species.

consimilis Brun., *Rec. Ind. Mus.* VI, 266, ♂ ♀ (1911).

Id., *Fauna Brit. Ind.*, 346, ♂ ♀ ; *Rec. Ind. Mus.* XV, 275.

Ghoom, above Darjiling, 6000—7000 ft., 11-vi-14 [*Gravelly*]; Darjiling, 7000 ft., 23—28-v-10, *type* ♂ ♀ ; 1-x-08; 17-v—10-vi-17; very common [all *Brunetti*]; Ghumti, Darj. Distr., 4000 ft. [*Gravelly*]; 5—10-viii-09 [*Paiva*]; Pashok, Darj. Distr., 2000—3500 ft., 23-iv—11-v-15 [*Gravelly*]; Kalimpong, Darj. Distr., 600—4500 ft., 24-iv—10-v-15 [*Gravelly*]; Kurseong, 4700—5000 ft., 19-vi-10; 4-ix-09; 19—24-vi-10; 5-vii-08; 5—8-ix-09 [all *Annandale*]; Gangtok, Sikkim, 22-v-05; Bhim Tal., 17—19-ix-06 [*Annandale*]; Mussoorie, 22-v-05 [*Brunetti*]; Tarnab, Peshawar Distr., v-16 [*Fletcher*]; Cherrapunji, Assam, 4400 ft., 2—3-x-14 [*Kemp*]; Shillong, 5500—6400 ft., 29-viii—5-ix-15 [*Kemp*]; x-16 [*Fletcher*].

The species appears generally distributed throughout India. *Type* ♂ ♀ in Indian Museum; *cotypes* in Vienna Museum, Pusa collection and my own.

serricornis Brun., *Fauna Brit. Ind.*, 341 ♂ (1912).

Mussoorie, 7000 ft. [*Brunetti*]; Almora, Kumaon, 5500 ft. [*Paiva*]; Darjiling, 6-viii-09 [*Paiva*]; Kurseong, 4-ix-09, *type* ♂; 5-vii-08; 22—25-vi-10 [*Annandale*]; Sureil, iv—v-17 [*Kemp*]; Gangtok, Sikkim, 6150 ft., 10-ix-09; Siliguri, 18—20-vii-07; Pusa, ii, iii, iv, vii, ix, xii rather common; Lahore, 8-iv-14 [*Lord*]; Shillong, 5000 ft., x-16 [*Fletcher*]; 19-x-11; Tengueh, Yunnan [*Coggin Brown*]. *Type* in Indian Museum.

Some ♂♂ taken by me at Yokohama, 20--26-vi-06, are probably the same species.

puncticornis, *id.*, *loc. cit.*, 343, ♂ 1912.

Siliguri, 30-vi-08. Type in the Indian Museum.

pleurinotata, *id.*, *loc. cit.*, 343, ♂ ♀ (1912).

Id., Rec. Ind. Mus. XV, 274.

Pusa, 14-iii-13 on guava; Namoya, Ceylon, ix-1909 [Green]. Types in Indian Museum.

demarcata Brun., Fauna Brit. Ind., 344, ♂ ♀ (1912).

Darjiling, 5-9-viii-09 [Paiva]; Kurseong, 5000 ft. [Annan-dale]. Types in Indian Museum.

concolorithorax, *id.*, *loc. cit.*, 346, ♂ ♀ (1912).

Sylhet, 18-i-04, type ♀ [Hall]; Pusa, 21-vi-11 [Fletcher]; Shillong [Kemp]; Khasi Hills, 1000-3000 ft., iii-07 type ♂. Type ♂ in Pusa coll.: type ♀ in Indian Museum.

gamma, *id.*, *loc. cit.*, 347, ♀ (1912).

Id., Rec. Ind. Mus. XV, 275.

Darjiling to Soom, 6000-7000 ft. [Lord Carmichael colln.]; 11 and 12-vi-14 [Graveley]; Mangaldai Distr., Assam-Bhutan Frontier, 30-31-xii-10, type ♀ [Kemp]. Type in Indian Museum.

speculata de Meij., Nova Guin. Res. IX, 314, ♀ (1913).

Alkmaar, Papua, xii, ii. Unique type in Amsterdam Museum.

ochripleuris de Meij., Tijd. v. Ent. LVI, Supp. p. 6, ♂ ♀ (1913) (Mar. 1914).

Alex., Proc. U.S. Nat. Mus. XLIX, pl. xlvi, 46, wing.

Java; Tjibodas, Mt. Gedé, Java, 26-viii-09, 7000 ft. [Bryant and Palmer]. Type in Amsterdam Museum.

citrina Edw., Ann. Mag. Nat. Hist. (8) XVIII, 265, ♀ (1916).

Taihoku, Formosa [Shiraki].

Unique type in British Museum.

parva Edw., *loc. cit.*, 266, ♂ (1916)

Arisan, Formosa, 10-x-12 [Nitobe].

Unique type in British Museum.

formosensis Edw., *loc. cit.*, 267, ♀ (1916).

Kammotu, Formosa [Nitobe].

Unique type in British Museum.

fuscoflava Brun., Rec. Ind. Mus. XV, 275, ♀ (1918).

Dungagali, 8000 ft. 21-24-v-15.

Type sent to British Museum, cotype in Pusa collection.

parvinotata, *id.*, *loc. cit.*, 276, ♂ ♀ (1918).

Taru, Peshawar Distr., 16-29-v-15, type ♂ ♀; Haripur Hazara, N.W. Frontier, 27-v-15 [Fletcher].

Types sent to British Museum, cotypes in Pusa collection.

hypocrites, *id.*, *loc. cit.*, 276, ♂ ♀ (1918).

Sureil, Darjiling District, 5000 ft., iv—v-17 [*Kemp*].

Types in Indian Museum.

Section DOLICHOPEZINI.¹

Scamboneura, Ost. Sack.

Berl. Ent. Zeits. XXVI, 95 (1882).

GENOTYPE, *S. dotata*, the original species.

vittifrons Walk., Proc. Linn. Soc. Lond. V, 144, ♀ (*Limnobia*) (1861).

Brun., Rec. Ind. Mus. VI, 267.

Amboina.

Type, reduced to a fragment, in British Museum.

dotata Ost. Sack., Berl. Ent. Zeits. XXVI, 95, fig. 1, ♂ (1882).

Philippines. Location of *type* unknown.

quadrata de Meij., Tijd. v. Ent. LVI, Supp. p. 8, 1913 (Mar. 1914).

Java. *Type* in Amsterdam Museum.

Megistocera, Wied.

Dipt. Exot. I, 41 (*Maekistocera*) (1821).

Wied., Auss. Zweifl. I, 55: Ost. Sack., Berl. Ent. Zeits. XXX, 158 (recharacterisation of genus and notes).

GENOTYPE, *Tipula filipes* F., the second species included by Wiedemann, by designation of Macquart, Dipt. Exot. I, 59.

fuscana Wied., Dipt. Exot. I, 29, ♀ (*Nematocera*) (1821).

Id., *loc. cit.*, 41, ♀ (*Maekistocera*): *id.*, Auss. Zweifl. I, 55, ♀: Macq., Hist. Nat. Dipt. I, 114, ♂, pl. ii, 15, full insect: *id.*, Dipt. Exot., Supp. I, 18, pl. ii, 4, wing: Ost. Sack., Ann. Mus. Gen. XVI, 399: Wulp, Notes Leyd. Mus. VII, 3: *id.*, Tijd. v. Ent. XXVIII, 83, pl. iv, 4: Ender., Zool. Jahr. XXXII, 30, figs., wing (1912): Brun., Rec. Ind. Mus. VI, 267.

Batavia, viii—xi; Bandong, vi [both *Jacobson*]; Celebes; Sumatra; Aru Is.

Type in Westermann's collection (Vienna Museum).

¹ For characters and comparison see Ost. Sack., Monog. Dipt. N. Amer. IV, 289, and Brun., Fauna Brit. Ind., 353.

Mitopeza, Edw.

Ann. Mag. Nat. Hist. (8) XVII, 349 (1916).

GENOTYPE, *M. nitidirostris* Edw., sp. nov.

nitidirostris Edw., *loc. cit.*, p. 350, ♀, fig. 1, p. 356, genitalia (1916).

Kedah Peak, 3200 ft., Malay Peninsula [Stanton].

The unique *type* in the British Museum.

Nesopeza, Alex.

Can. Ent. XLVI, 157 (1914).

GENOTYPE, *Dolichozeza gracilis* de Meij.

gracilis de Meij., Tijds. v. Ent. LIV, 60, ♂, pl. iv, 46, wing (*Dolichozeza*) (1911).

Wonosobo, Java, v; Tjibodas, Mt. Gedé, Java, 4500 ft. [Bryant and Palmer]. *Type* in Amsterdam Museum.

albitarsis Brun., Rec. Ind. Mus. XV, 278, ♂ ♀, pl. vii, 1, wing (1918).

Lebong, 6000—6600 ft., 13-vi-14, *type* ♂ [Gravelly]; between Darjiling and Soom, 5000—7000 ft., 14-vi-14 [Gravelly].

The *types*, a unique pair, in Indian Museum.

longicornis, *id.*, *loc. cit.*, 278, ♂ (1918).

Above Tura, Garo Hills, 3500—3900 ft., vii-1917 [Kemp]. *Type* in Indian Museum.

picticornis, *id.*, *loc. cit.*, 279, ♂ (1918).

Inle, Southern Shan States, 3000 ft., 18-ii-17 [Gravelly]. *Type* in Indian Museum.

Dolichozeza, Curt.

Brit. Ent., 62 (1825).

Sch., F. Austr. II, 525: Ost. Sack., Berl. Ent. Zeits. XXX, 157: Brun., Fauna Brit. Ind., 353.

GENOTYPE, *D. sylvicola* Curt.,¹ by original designation.

Leptina, Mg., Sys. Besch. VI, 283, pl. lxx, 10, in text as *Dolichozeza*, on plate as *Leptina* (1830).

Apeilesis, Macq., Dipt. Exot. Supp. I, 8, pl. i, 3 (1846): *id.*, Ost. Sack., Berl. Ent. Zeits. XXX, 158.

GENOTYPE, *A. cinerea*, sp. nov., by original designation.

¹ Some desire to change this name to *albipes*, Strom, but *sylvicola* has stood so long unchallenged that I decline to accept any alteration.

orientalis Brun., Fauna Brit. Ind., 354, ♂ ♀, pl. v, 19, wing (1912).

Darjiling, 7000 ft., 4-vi-17 [Brunetti]; 11-vi-14 [Gravely]; Kurseong, 8-vii-08, type ♂; 4-ix-09; 22-vi-10, type ♀ [Annandale]; Tura, Garo Hills, viii-17 [Kemp]; Ginting Bidai, Federated Malay States, iv-17 [Boden Kloss].

Types in Indian Museum.

obscura, *id.*, *loc. cit.*, 355, ♂ ♀ (1912).

Kurseong, 4700 ft., 14-iv-11, type ♂, 29-vi-10, type ♀ [both Annandale]; Lebong, 6000—6600 ft. [Lord Carmichael colln.].

Types in Indian Museum.

postica *id.*, *loc. cit.*, App. 564, ♂ ♀ (1912).

Ghumti, Darjiling Distr., 6000 ft., vii-1911, including ♂ ♀ in cop. [Gravely]; Tura, Garo Hills, viii-17 [Kemp].

Types in the Indian Museum.

infuscata, *id.*, *loc. cit.*, App. 565, ♂ (1912).

Nilgiri Hills, 6000 ft., v-1911 [Andrewes].

Type in Indian Museum, a unique specimen.

costalis Brun., Rec. Ind. Mus. XV, 277, ♂, pl. viii, 6, wing (1918).

Kavalai, Cochin State, 1300—3000 ft., 24—27-ix-14 [Gravely].

Unique type in Indian Museum.

pallidithorax de Meij., Tijd. v. Ent. LVI, Supp., p. 4, ♂; pl. i, 3. 1913 (Mar. 1914).

Java. Type in Amsterdam Museum.

Sub-family LIMNOBIINAE.

Section CYLINDROTOMINI.¹

Stibadocera, Ender.

Zool. Jahr. XXXII, 83 (1912).

GENOTYPE, *S. bullans* Ender., by original designation.

quadricellula Brun., Rec. Ind. Mus. VI, 268, ♂ (*Cylindrotoma*) (1911).

Id., Fauna Brit. Ind., 361, ♂, pl. vii, 1, antenna, 2, wing.

Kurseong, 18-v-10, type, 23-vi-10; 6-vii-08 [Annandale].

Type in Indian Museum.

bullans Ender., Zool. Jahr. XXXII, 84, ♂, fig. A 2, wing (1912).

Sumatra (Liangagos).

Type, a unique ♂, in Stettin Museum.

metallica Alex., Proc. U.S. Nat. Mus. XLIX, 178, pl. xlv, 30, wing (1916).

¹ For characters and comparisons v. Ost. Sac., Monog. Dipt. N. Amer. IV, 289 and Brun., Fauna, 353: *id.*, Rec. Ind. Mus. XV, 280 on generic characters.

Tjibodas, Mt. Gedé, Java, 9000 ft. [Bryant and Palmer];
Pangrango, Java. Type, No. 19,026, in U.S. Nat. Mus.

Stibadocerella, Brun.

Rec. Ind. Mus. XV, 283 (1918).

GENOTYPE, *S. pristina*, *id.*, sp. nov., *loc. cit.*

pristina Brun., *loc. cit.*, 283, ♂ ♀ (1918).

Tura, Garo Hills, 3500—3900, vii-17 [Kemp].

Type in Indian Museum.

N.B.—In my second paper on Oriental Tipulidae a “new species *latefurcata*” (of *Cylindrotoma*) is mentioned. This was a MS. name for *pristina*, and referred to the late bifurcation of the second and third veins. On establishing *Stibadocerella* the name *bifurcata* was not so applicable, but in one instance the MS. name by oversight was not changed.

Section LIMNOBIINI.¹

Dicranomyia, Steph.

Cat. Brit. Ins. II, 243 (1829).

Sch., F. Austr. II, 565, as *Limnobia*, pt.: Ost.

Sack., Monog. Dipt. N. Amer. IV, 53: *id.*, Berl.

Ent. Zeits. XXXI, 172: Brun., Fauna Brit.

Ind., 363.

GENOTYPE, *Limnobia modesta* Mg., designated by
Coquillett, 1910.

Siagona, Mg., Syst. Besch. VI, pl. lxxv, 7 (1830).

GENOTYPE, *Glochina sericata* Mg., by original
designation.

Numantia, Big., pt., Ann. Soc. Ent. Fran. (3) II,
470 (1854).

GENOTYPE, *Limnobia fusca* Mg., by Coquillett's
designation.

Thrypticomyia, Skuse, Proc. Linn. Soc. N.S. Wales
(2) IV, 775, pl. xxi, 6, xxiv, 44, 45 (1890).

GENOTYPE, *T. auripennis*, *id.*, sp. nov., *loc. cit.*

? *Glochina*, Mg., Syst. Besch. VI, 280, pl. lxxv, 7 (as
Siagona on plate) (1830).

*saltens*² Dol., Nat. Tijd. Ned. Ind. XIV, 390, pl. ii, 3 (*Limnobia*)
(1857).

Ost. Sack., Berl. Ent. Zeits. XXVI, 88: de Meij., Tijd. v.

Ent. LIV, 22, pl. i, 1, wing: Brun., Rec. Ind. Mus. VI, 269:

id., Fauna Brit. Ind., 373, ♂ ♀.

¹ For table of oriental genera, v. Brun., Fauna, 363.

² Sometimes (erroneously) spelt *saltans*. If the genus *Thrypticomyia*, Skuse, be valid, *saltens*, *kobusi*, *cuneiformis*, *longivena*, *absens* and *bicinctipes* come in it.

Travancore, 14-xi-09 [*Annandale*]; Cherambadi, Wynaad, S. India, x-13 [*Fletcher*]; Dibrugarh, Assam, 17-19-xi-11 [*Kemp*]; Java (var. locs.). *Type*: location unknown to me.

This species may possibly be synonymous with *Limnobia apicalis*, Wied.

kobusi de Meij., *Bijd. tot de Dierk.* XVII, 91, pl. viii, 5, wing, 6, tip of leg (1904).

Brun., *Rec. Ind. Mus.* VI, 270: *id.*, *Fauna Brit. Ind.*, 371, ♂ ♀: de Meij., *Tijd. v. Ent.* LVI, 343, pl. xvi, 8, ♂ genitalia.

Kurseong, 20-25-vi-10; 5-vii-08, including ♂ ♀ in cop. [*Annandale*]; Lebong, 6000-6600 ft., 13-vi-14; Nedumangad, Travancore [*Annandale*]; Dibrugarh, Assam (Abor. Exped.) [*Kemp*]; Nusa, Kambangan, Java, iii [*Jacobson*].

Types ♂ ♀ in Amsterdam Museum.

pulchra de Meij., *Bijd. tot de Dierk.* XVII, 92, ♀, pl. viii, 7, full ins. (*Rhipidia, id.*) (1904).

Id., *Tijd. v. Ent.* LIV, 27, pl. ii, 14-16 antennae, ♂ ♀ genitalia (*Dicranomyia, id.*): Brun., *Rec. Ind. Mus.* VI, 240.

Batavia, ii, x; Semarang, ii [both *Jacobson*].

Type in Amsterdam Museum.

cuneiformis de Meij., *Tijd. v. Ent.* LIV, 23, ♂, pl. i, 2, wing (1911).

Brun., *Rec. Ind. Mus.* VI, 270; XV, 284: *id.*, *Fauna Brit. Ind.*, 372, ♂ ♀.

Talewaddi, Castle Rock, N. Kanara Dist., 10-x-16 [*Kemp*]; Dawna Hills, 2000-3000 ft., 2-3-iii-08 [*Annandale*]; Wonosobo, Java, iv [*Jacobson*].

Type ♂ in Amsterdam Museum; the ♀ from which I described this sex in Indian Museum.

convergens de Meij., *Tijd. v. Ent.* LIV, 24, ♂ ♀, pl. i, 3-6, wing, and ♂ ♀, genitalia (1911).

Tosari [*Kobus*]; Semarang, ii; Tankuban Prahau [both *Jacobson*, all Java]. *Types* in Amsterdam Museum.

umbrata, id., loc. cit., 25, ♂ ♀, pl. i, 7, wing (1911).

Id., *Tijd. v. Ent.* LVI, 343, pl. xlvi, 9, 10, ♂ genitalia.

Semarang, xii; Batavia and environs, x, xi [*Jacobson* and *Dr. Rogge*]. *Types* in Amsterdam Museum.

punctulata, id., Tijd. v. Ent. LIV, 26, ♂; pl. i, 8, wing, 9, ♂ genitalia (1911).¹

Semarang, i; Wonosobo, v [*Jacobson*].

Type in Amsterdam Museum.

nervosa, id., loc. cit., 26, ♀, pl. i, 10, wing (1911).

Wonosobo, Java, v [*Jacobson*].

Type, a unique ♀, in Amsterdam Museum.

¹ De Meijere employs both sex signs at the beginning of his description though he stated his specimen to be unique. Probably only the ♂ sex was before him.

tenella, *id.*, *loc. cit.*, 27, ♂ ♀, pl. i, 11—13, wing, ♀ genitalia (1911).

Pangerango, Java, x; Semarang, xi [both *Jacobson*].

Types in Amsterdam Museum, a unique pair.

longivena, Edwards, *Ann. Mag. Nat. Hist.* (8) VIII, 59, ♀ (1911).

Brun., *Fauna Brit. Ind.*, 374, ♀.

Dondra, Ceylon, 3-xii-07 [*Fletcher*].

Type in British Museum.

marmoripennis Brun., *Fauna Brit. Ind.*, 369, ♂ ♀, pl. vii, 3, wing (1912).

Darjiling, 16-vi-14 [*Gravelly*]; 8-viii-09 [*Paiva*]; 3-9-vi-06 [*Howlett*]; Kurseong, 5-ix-09, *type* ♂ ♀ [*Annandale*]; Katihar, 7-31-viii-10 [*Paiva*]; Pusa, 29-vi-11; 1-viii-11 sweeping grass [both *Howlett*]; Bangalore, 3000 ft. [*Annandale*]; Shillong, x-16 [*Fletcher*]; 5500—6400 ft., 29-viii—5-ix-15 [*Kemp*]; Lashio, Upper Burma, 3000 ft., 23, 24-viii-14 [*Fletcher*]; Kobo, Assam (Abor Exped.), 400 ft., 2-3-xii-11 [*Kemp*]. *Types* in Indian Museum.

demarcata, *id.*, *loc. cit.*, 370, ♀ (1912).

Kurseong, 15-iv-11 [*Annandale*]. *Type* in Indian Museum.

absens, *id.*, *loc. cit.*, 372, ♀, pl. vii, 4, wing (1912).

Darjiling, 5-viii-09 [*Paiva*]; Kurseong, 25-vi-10, *type* [*Annandale*]; Hakgala, Ceylon. *Type* in Indian Museum.

Two ♂♂ and two ♀♀ taken by me at Darjiling are probably this species, two have the discal cell open (specific character) and two have it closed.

pulchripennis, *id.*, *loc. cit.*, 376, ♂ ♀, pl. vii, 8, wing; xi, 2, ♂ genitalia (1912).

Mussoorie, 18-vi-05; 17-viii-05 [both *Brunetti*]; Simla, x-08; Bhowali, 5700 ft., x-09 [*Imms*]; Kangra Valley, W. Himalayas [*Dudgeon*]; Kurseong [*Annandale*]; Shillong, 5500—6400 ft., 29-viii—5-ix-15 [*Kemp*]; x-16 [*Fletcher*].

Types in Indian Museum; *cotypes* in my collection.

puncticosta, *id.*, *loc. cit.*, 377, ♂ ♀, pl. vii, 7, wing (1912).

Kurseong, 4-ix-09, *type* ♂ [*Annandale*]; Darjiling, 8-viii-09, *type* ♀ [*Paiva*]. *Types* in Indian Museum.

fraterna, *id.*, *loc. cit.*, 378, ♀, pl. vii-5, wing (1912).

Darjiling, 5--8-viii-09 [*Paiva*]. *Type* in Indian Museum.

fascipennis, *id.*, *loc. cit.*, 379, ♀, pl. vii, 9, wing (1912).

Almora, Kumaon, 5500 ft., 24-vi-11, *type* [*Paiva*], 17—19-ix-11 [*Paiva*]; Pusa, in spiders' webs, 1-i-14 [*Howlett*]; Kurseong, 5-vii-08, *type* ♀ [*Annandale*]; Shillong, 29-viii—5-ix-15 [*Kemp*]; Coonoor, at light, 4-ix-11; Peradeniya [*Gravelly*].

Types in Indian Museum.

- subfascipennis**, *id.*, *loc. cit.*, 380, ♂ ♀, pl. vii, 6, wing (1912).
Kurseong, 4-vii-08 [*Annandale*]; Ghumti, 4000 ft. [*Gravely*].
Types in Indian Museum.
- ornatipes**, *id.*, *loc. cit.*, 380, ♂, pl. vii, 10, wing (1912).
Vaikam, Travancore, 5-xi-08, *type* [*Annandale*]; Balighai, near
Puri, 24-x-08; Khurda Rd., Orissa; Dawna Hills, 2000—3000
ft., 2—3-iii-08 [all *Annandale*]. *Type* in Indian Museum.
- cinerascens**, *id.*, *loc. cit.*, 381, ♂ ♀, pl. vii, 11 (1912).
Darjiling, 8-iv-10, *type* ♂; 10--20-x-05 [*Brunetti*]; 7000 ft.
[*Beebe*]; Kurseong, 24-iii-10, *type* ♀. *Type* in Indian Museum.
- sordida**, *id.*, *loc. cit.*, 382 (1912).
Kurseong, 4-ix-09, *type* ♂; 9-ix-09 [*Annandale*]; Darjiling, 7-
viii-09, *type* ♀ [*Paiva*]. *Type* in Indian Museum.
- delicata**, *id.*, *loc. cit.*, 383, ♂ ♀ (1912).
Darjiling, 6-viii-09, *type* ♂, 9-viii-09, *type* ♀ [both *Paiva*].
Types in Indian Museum.
- flavobrunnea**, *id.*, *loc. cit.*, 384, ♂ ♀ (1912).
Calcutta, 18-xi-07, *type* ♂ ♀; "at light," 9-vi-07; 6-xi-07; 20-
xii-07; Purulia, Manbhum Distr., Chota Nagpur, 10-ii-12 [*Annandale*
and *Gravely*]; Tenmalai, Travancore [*Annandale*].
Types in Indian Museum.
- simplex**, *id.*, *loc. cit.*, 384, ♀ (1912).
Calcutta, 6-xi-10 [*Gravely*]; Shillong, Khasi Hills, 5500—6400
ft. [*Kemp*]. *Type* in Indian Museum.
- fortis**, *id.*, *loc. cit.*, 385, ♂ (1912).
Darjiling, 6-viii-09, *type* [*Paiva*]; 9-viii-09 [*Brunetti*].
Type in Indian Museum.
- nigrithorax**, *id.*, *loc. cit.*, 385, ♂ (1912).
Darjiling, 28-vi-10, *type*; 26-v-10 [all *Brunetti*].
Type in Indian Museum.
- subtessellata**, *id.*, *loc. cit.*, App., 565, ♂ ♀ (1912).
Pattipola, Ceylon, 6000 ft., 14-x-11 [*Annandale*]; Castle Rock,
Goa, 11—26-x-16 [*Kemp*]. *Types* in Indian Museum.
- bicinctipes**, *id.*, *loc. cit.*, App., 566, ♂ (1912): Rec. Ind. Mus. VII,
447.¹
"Thingannyinaung to Sukli," Dawna Hills, 900—2100 ft., 23
—27-xi-11 [*Gravely*]. *Type* in Indian Museum.
- columbina**, *id.*, *loc. cit.*, App., 567, ♀ (1912).
Peradeniya, 22-vii-10, *type*; 7-ii-10; 3-x-10 [*Green*]; Shillong
[*Kemp*]. *Type* in Indian Museum.

¹ The first reference antedates the second by about a month. The species comes in the *Thrypticomyia* group.

- approximata**, *id.*, *loc. cit.*, App., 567, ♀ (1912).
Ghumti, Darjiling Distr., 4000 ft., vii-1911 [*Gravelly*].
Type in Indian Museum, a unique ♀.
- innocens**, *id.*, *loc. cit.*, App. 568, ♂ ♀ (1912).
Almora, Kumaon Div., 18—25-vi-11 [*Paiva*].
Types in Indian Museum.
- debeauforti** de Meij., *Bijd. tot de Dierk.* XIX, 47, ♂ ♀ (1913).
Saonek, xii, i. *Types* in Amsterdam Museum.
- alta** de Meij., *Tijd. v. Ent.* LVI, 341, ♂, pl. xvi, 6, wing (1913).
Gunung Ungaren, ix [*Jacobson*].
Type, a unique ♂, in Amsterdam Museum.
- nongkodjadjarensis**, *id.*, *loc. cit.*, 343, ♂, pl. xvi, 7, wing (1913).
Nongkodjadjar, Java, i [*Jacobson*].
Type, a unique ♂, in Amsterdam Museum.
- novae-guineae**, *id.*, *loc. cit.*, LVIII, 101 (*Thrypticomyia*) (1915).
Hollandia, North Papua *Type* in Amsterdam Museum.
- tinctipennis**, *id.*, *loc. cit.*, LVIII, Supp. 66 (1916).
Fort de Kock, Sumatra. *Type* in Amsterdam Museum.
- fullowayi** Alex., *Can. Ent.* LVII, 79 (1915).
Arisan, Formosa, 8000 ft., 10-x-12 [*Nitobe*]. Described originally from the Ladrone Islands.
- alticola** Edw., *Ann. Mag. Nat. Hist.* (8) XVIII, 246, ♀, pl. xii, 1, genitalia (1916).
Arisan, Formosa, 8000 ft., 10-x-12 [*Nitobe*]; Horisha, Formosa. *Type* in British Museum.
- albitarsis** Alex., *Proc. U.S. Nat. Mus.* XLIX, 159, ♂; pl. xlii, 1, wing (1916).
Tjibodas, Mt. Gedé, Java, 4500 ft. [*Bryant and Palmer*].
Type, Cat. No. 19,007, in U.S. Nat. Mus.
- atrescens**, *id.*, *loc. cit.*, 160, ♂ (1916).
Tjibodas, Mt. Gedé, Java, 8000 ft. [*Bryant and Palmer*].
Type, Cat. No. 19,008, in U.S. Nat. Mus.
- erythrina**, *id.*, *loc. cit.*, 161, ♂, pl. xlii, 2, wing (1916).
Tjibodas, Mt. Gedé, 4500 ft., Java [*Bryant and Palmer*].
Type, Cat. No. 19,009, in U.S. Nat. Mus.
- excelsa**, *id.*, *loc. cit.*, 161, ♂ ♀, pl. xlii, 3, wing (1916).
Pangrango, Java, 9000 ft., *type* ♂ [*Bryant and Palmer*];
Tjibodas, Mt. Gedé, 8000 ft., *type* ♀.
Type, Cat. No. 19,010, in U.S. Nat. Mus.
- simplissima**, *id.*, *loc. cit.*, 162, ♀ pl. xlii, 4, wing (1916).
Tjibodas, Mt. Gedé, Java, 9000 ft. [*Bryant and Palmer*].
Type, Cat. No. 19,011, in U.S. Nat. Mus.

- carneotincta*, *id.*, *loc. cit.*, 162, ♂, pl. xlii, 5, wing (1916).
Singapore [Bryant and Palmer].
Type, Cat. No 19,012, U.S. Nat. Mus.
- bicolor* Brun., Rec. Ind. Mus. XV, 285, ♂ (1918).
Cherrapunji, Assam, 4,400 ft., 2-8-x-14 [Kemp].
Unique *type* in Indian Museum.
- prominens*, *id.*, *loc. cit.*, 285, ♂ ♀ (1918).
Mormugao, Goa, ix-16 [Kemp]. *Types* in Indian Museum.
- niveiapicalis*, *id.*, *loc. cit.*, 285, ♂ (1918).
N. Kanara Distr., W. India, 11-26-x-16 [Kemp].
The unique *type* in the Indian Museum.
- pictipes*, *id.*, *loc. cit.*, 286, ♂ ♀ (1918).
Mormugao, Goa, ix-16 [Kemp]. *Types* in Indian Museum.

Geranomyia, Hal.

- Ent. Month. Mag. I, 154 (1833).
Sch., F. Austr. II, 564: Ost Sack., Monog.
Dipt. N. Amer. IV, 78; *id.*, Berl. Ent Zeits.
XXXI, 173: Brun., Fauna Brit. Ind., 387:
Auct.
GENOTYPE, *G. unicolor*. Hal., by original designation.
- Limnobiaorhynchus*, ♀ only, Westw., Ann. Soc
Ent. Fran. (1) IV, 683 (1835).
Plettusa, Phil., Verh. Zool. bot. ges. Wien, XV, 597,
pl. xxiii, 1 (1865).
GENOTYPE, *P. virescens*, sp. nov., *loc. cit.*
- sorbillans* W., Auss. Zweifl. I, 551, ♀ (*Limnobia*) (1828).
Brun., Rec. Ind. Mus. VI, 274.
Sumatra. *Types* in Wiedemann's and Trentepohl's collections at Vienna Museum.
- nitida* de Meij., Tijd. v. Ent. LIV, 28, ♀ (1911).
Alex., Proc. U.S. Nat. Mus. XLIX, pl. xlii, 6, wing.
Wonosobo, Java, iv [Jacobson]; Batavia, 1-iv-09 [Bryant
and Palmer]. *Type* in Amsterdam Museum.
- argentifera* de Meij., Tijd. v. Ent. LVI, 29, ♂ ♀, pl. ii, 18, wing,
19, ♂ genitalia (1911).
Semarang, 1-vii [Jacobson]. *Types* in Amsterdam Museum.
- montana*, *id.*, *loc. cit.*, 29, ♂ ♀ (1911).
Pangerango, Java, x [Jacobson]. *Type* in Amsterdam Museum.

notata, *id.*, *loc. cit.*, 30, ♀ (1911).

Batavia environs [*Jacobson*]. Type in Amsterdam Museum.

fletcheri Edwards, Ann. Mag. Nat. Hist. (8) VIII, No. 43, 60, ♀ (1911).

Brun., Fauna Brit. Ind. VI, 393, ♀.

Madulsima, Ceylon, 19-v-08, *type*; 21-xii-07.

Type in British Museum.

vinaceobrunnea Brun., Rec. Ind. Mus. VI, 274, ♀ (1911).

Id., Fauna Brit. Ind., 392, ♀.

Simla, x-08 [*Howlett*]. Type in Pusa collection.

genitalis, *id.*, Rec. Ind. Mus. VI, 275, ♂ ♀ (1911).

Id., Fauna Brit. Ind., 388, ♂ ♀, pl. xi, 7, ♂ genitalia: *id.*,

Rec. Ind. Mus. XV, 287.

Pashok Spur, Darjiling, 2000—3500 ft., 25-iv—ii-v-15 [*Gravelly*]; Tenmalai, Travancore, 21-xi-08, *type* ♂ [*Annandale*]; Nongpoh, Assam, ix-06, *type* ♀.

Type ♂ in Indian Museum, *type* ♀ in Pusa coll.

semifasciata Brun., Rec. Ind. Mus. VI, 276, ♀ (1911).

Id., Fauna Brit. Ind., 395, ♀, pl. vii, 16, wing.

Darjiling, 3—9-vi-09 [*Howlett*]; Kalimpong, 600—4500 ft. [*Gravelly*]; Sureil, iv, v-17 [*Kemp*]; Almora, Kumaon, 5500 ft. [*Paiva*]; Heho, S. Shan States, 7—9-iii-17 [*Gravelly*].

Type in Pusa collection.

semistriata Brun., Rec. Ind. Mus. VI, 277, ♀ (1911).

Id., Fauna Brit. Ind., 394, ♀, pl. vii, 17, wing.

Pashok, Darjiling Distr., 2000—3500 ft. [*Gravelly*]; Paresnath, W. Bengal, 4300—4500 ft., 15-iv-09 [*Annandale*]; Talewadi, 3—10-x-16 and Castle Rock [both N. Kanara Dist. and *Kemp*]; Shillong, 4900 ft., 10—12-x-14; Cherrapunji, Assam [both *Kemp*].

flavicosta Brun., Fauna Brit. Ind., 389, ♀, pl. viii, 2, wing, pl. xi, 6, dorsum of thorax (1912).

On board a launch, "at light" off Chandpal, Ganges Delta, 26-viii-09 [*Jenkins*]. Type, an incomplete unique specimen in Indian Museum.

circipunctata, *id.*, *loc. cit.*, 390, ♂ ♀, pl. viii, 3, wing, pl. xi, 3, thorax dorsum.

Port Canning [*Annandale*]; Lower Bengal, and Ganges Delta, var. locs., ii, iv, viii, ix, xi, xii (including Sorabkatti, 7-xii-09, *type* ♀, Calcutta; Sonarpur, near Calcutta, 5-ii-10, "at light" in railway carriage, *type* ♂ [*Annandale*]; "at light" on board steam launch off coast of Ganges Delta, var. locs. [*Jenkins*]; Sunderbunds [*Kemp*]; on board ship 10 miles off Masulipatam, Madras Coast, 4—5-iv-08 [*Paiva*]; Puri, 12-xi-10 [*Annandale*]. Types in Indian Museum.

- tridens*, *id.*, *loc. cit.*, 391, ♂ ♀, pl. viii, 4, wing; pl. xi, 45, thorax (1912).
Ganges Delta, var locs. in the Sunderbunds, including *type* ♂ from Sonadigee, 6-xii-09, and *type* ♀ from Chennia, 11-xi-09, all "at light" on steam launch. *Types* in Indian Museum.
- pulchripennis*, *id.*, *loc. cit.*, 393, ♀, pl. viii, 1, wing (1912).
Kurseong, 7-ix-09 [*Annandale*]; Shillong, 6400, 12-x-14 [*Kemp*]. *Type*, a unique ♀, in Indian Museum.
- notatipennis*, *id.*, *Rec. Ind. Mus.* VIII, 152, ♂, pl. vi, 1, wing (1913).
Dibrugarh, N.E. Assam, 17—19-xi-11 [*Kemp*].
Type, a unique specimen, in Indian Museum.
- decemguttata* de Meij., *Tijd. v. Ent.* LVI, 345, ♂ ♀, pl. xvii, 11, wing (1913).
Semarang, viii [*Jacobson*].
Types in Amsterdam Museum.
- brunnescens* de Meij., *loc. cit.*, LVIII, Supp. 10, ♂ (1916).
Sinabang, Simalur Is., Sumatra, ii [*Jacobson*].
Unique *type* in Amsterdam Museum.
- septemnotata* Edw., *Ann. Mag. Nat. Hist.* (8) XVIII, 246, ♀ (1916).
Arisan, Formosa, 8000 ft., 10-x-12 [*Nitobe*].
- linearis* Alex., *Proc. U.S. Nat. Mus.* XLIX, 163, ♂; pl. xlii, 7, wing (1916).
Tjibodas, Mt. Gedé, Java, 4500 ft. [*Bryant and Palmer*].
Unique *type*, Cat. No. 19,013, in U.S. Nat. Mus.
- javanica* Alex., *loc. cit.*, 164, ♀; pl. xlii, 8, wing (1916).
Batavia, 1-iv-09 [*Bryant and Palmer*].
Unique *type*, Cat. No. 19,014, in U.S. Nat. Mus.
- cornigera* Alex., *Ins. Menstr.* I, 137 (1914).
Pettit Barracks, Luzon [*Ludlow*]. *Type*: location unknown to present author.
- nigronotata* Brun., *Rec. Ind. Mus.* XV, 287, ♀ (1918).
Cherambadi, Wynaad, Malabar Distr., x-13 [*Howlett*].
Type, a unique ♀, presented to British Museum.
- flaviventris*, *id.*, *loc. cit.*, 288, ♂, pl. vii, 2, wing (1918).
Pashok, Darjiling, 3000 ft., 14-vi-16 [*Gravelly*].
Type in Indian Museum.

Aporosa,¹ Macq.

In Webb and Berth, *Hist. Nat. d'iles Canaries*:
Entom. Dipt. 100 (1838).
Macq., *Dipt. Exot.*, I, 1, 62 (1838).

¹ *Geranomyia* and *Aporosa* differences: v. Brun., *Rec. Ind. Mus.* XV, 288. Kertész ranks *Aporosa* as synonymous with *Geranomyia*, but Bergroth more recently keeps them apart. See note on *Elephantomyia*, p. 230.

GENOTYPE, *Geranomyia canariensis* Berg., as *maculipennis*, sp. nov. by Coquillett's designation.

aurantia Brun., Rec. Ind. Mus. XV, 289, ♀ (1918).

Above Tura, Garo Hills, 3,500—3,900 ft., vii-17 [Kemp].
Type in Indian Museum.

Limnobia, Mg.

Syst. Besch, I, 116 (1818).

Sch., F. Austr. II, 565: Ost. Sack., Monog. Dipt. N. Amer. IV, 84: *id.*, Berl. Ent. Zeits. XXXI, 177: Brun., Fauna Brit. Ind., 396: *Auctt.*

GENOTYPE, *Tipula 3-punctata* F., according to (but not designated by) Coquillett, 1910.

Numantia, Big. pt., Ann. Soc. Ent. Fran. (3) II, 470 (1854).¹

Limnomyza, Rond., Dipt. Ital. Prod. I, 185 (1856).

GENOTYPE, *Tipula 3-punctata* F., by original designation.

costalis Wied., Anal. ent. 10 (1824).

Id., Auss. Zweifl. I, 35: Brun., Rec. Ind. Mus. VI, 271.

East India. Type in Copenhagen Museum, in bad condition. Possibly not a true *Limnobia*. No sex of type stated, and now indeterminable.

apicalis W., Auss. Zweifl. I, 551, ♂ (1828).

Brun., Rec. Ind. Mus. VI, 271.

Sumatra. Possibly *Dicranomyia saltens*, Dol., is identical with this species.

Type, a fragment, in Winthem's collection, Vienna Museum.

bibula W., Auss. Zweifl., 552, ♀ (1828).

Brun., Rec. Ind. Mus. VI, 271.

China. Type in Trentepohl's collection, Vienna Museum.

Probably a true *Limnobia*, according to Osten Sacken, but possibly it should have no place in oriental lists.

sanguinea Dol., Nat. Tijd. Ned. Ind. XIV, 391, pl. ii, 5, full insect (1857).

Brun., Rec. Ind. Mus. VI, 271, notes.

Java. Type, location unknown to present author.

infixa Walk., Proc. Linn. Soc. Lond. VIII, 104 (1865).

Ost. Sack., Berl. Ent. Zeits. XXXI, 178: Brun., Rec. Ind. Mus. VI, 271.

Papua. Type in British Museum, in bad condition.

¹ See under *Dicranomyia*, p. 212.

- cinctiventris* Brun., Fauna Brit. Ind., 382, ♀ (*Dicranomyia*) (1912).
vitripennis, *id.*, *loc. cit.*, 405 ♀ (1912).
 Kurseong, 15-iv-11, *type*, *cinctiventris* [*Annandale*]; Darjiling, 30-ix-08 (*type*, *vitripennis*) 6900 ft. [*Brunetti*]; Mahableswar, Satara Distr., 4200 ft. [*Gravely*]. *Types* in Indian Museum.
- festiva* Brun., Fauna Brit. Ind., 400, ♂, pl. vii, 13, wing (1912).
Id., Rec. Ind. Mus. XV, 289.
 Darjiling, 7000 ft., 11-vi-14 [*Lord Carmichael's colln.*]; Kurseong, vi-1910. *Type* in Indian Museum.
- tinctinervis*, *id.*, Faun. Brit. Ind., 401, ♀, pl. vii, 12, wing (1912).
 Darjiling, 10-viii-09 [*Paiva*].
Type, a unique ♀, in Indian Museum.
- indica*, *id.*, *loc. cit.*, 401, ♂ ♀, pl. vii, 15, wing, pl. xi, 1, ♂ genitalia (1912).
 Mussoorie, ix-06 [*Howlett*]; Calcutta, not uncommon, Sept. to Feb.; Katihar, 7-31-viii-10; Madhupur, Bengal, 13-x-09 [*Paiva*]; Monghyr, Bihar [*Jenkins*]; Pusa, 18-20-ii-08, on moss; 18-iii-13; 20-ix-13; ix-07; 17-xii-12; Puri [*Paiva*]; Kalyan, Bombay, 7-x-13; Maddathorai, and Nedumangad [both Travancore and *Annandale*]; Ernakulam, Cochin, 11-14-ix-14 [*Gravely*]. *Types* in Indian Museum.
- trimaculata*, *id.*, *loc. cit.*, 402, ♂ ♀, pl. vii, 16, wing (1912).
 Ghumti, 4000 ft. [*Gravely*]; Kurseong, 3-5-vii-08; 18-29-vi-09 [*Annandale*]; on low herbage and "at light"; Lucknow [*Caunter*]; Shillong, 5500-6400 ft. [*Kemp*].
Types in Indian Museum.
- longinervis*, *id.*, *loc. cit.*, 403, ♀, pl. vii, 14, wing (1912).
 Kurseong, 5-ix-09 [*Annandale*].
Type, a unique ♀, in Indian Museum.
- centralis*, *id.*, *loc. cit.*, 403, ♂ ♀ (1912).
 Kurseong, 18-23-vi-10 [*Annandale*].
Types in Indian Museum.
- niveipes*, *id.*, *loc. cit.*, 404, ♀ (1912).
 Darjiling, 28-v-10 [*Brunetti*].
Type, a unique ♀, in Indian Museum.
- nigra*, *id.*, *loc. cit.*, 404, ♂ ♀ (1912).
 Kurseong, 4700-5000 ft. [*Annandale*]; Pallode, Travancore, 15-xi-08 [*Annandale*]; Talewadi, N. Kanara Dist., 3-10-x-16 [*Kemp*]. *Types* in Indian Museum.
- triangularis*, *id.*, *loc. cit.*, 406, ♀ (1912).
 Barogh, Simla Hills, 5000 ft., 10-v-10 [*Annandale*].
Type, a unique specimen, in Indian Museum.
- annulifemur* de Meij., Tijds. v. Ent. LVI, 344, ♂, pl. xvii, 12, wing (1913).
 Nongkodjadar, Java, i [*Jacobson*]; Tjibodas, Mt. Gedé, Java, 8000 ft. [*Bryant and Palmer*]; Korinchi, Sumatra, 7300 ft., v-14. *Type*, unique, in Amsterdam Museum.

- crocea** Edw., Ann. Mag. Nat. Hist. (8) XVII, 353, ♂ (1916).
Sungkie, Siam, 9-ii-02 [*Robinson and Annandale*].
Unique type in British Museum.
- nitobei**, *id.*, *loc. cit.*, XVIII, 247, ♂, pl. xii, 2, genitalia (1916).
Arisan, Formosa, 8000 ft. 10-x-12 [*Nitobe*].
Unique type in British Museum.
- flavocincta** Brun., Rec. Ind. Mus. XV, 289, ♀ (1918).
Mahabaleshwar, Satara Distr., W. India, 4200 ft., 13—16-iv-12 [*Gravelly*]. Type in Indian Museum.
- marginata**, *id.*, *loc. cit.*, XV, 290, ♀, pl. viii, 7, wing (1918).
Mahabaleshwar, Satara Distr., 4,200 ft., 13—16-iv-12 [*Gravelly*].
Type in Indian Museum.
- confinis**, *id.*, *loc. cit.*, 290, ♂ ♀ (1918).
Above Tura, Garo Hills, Assam, 3500—3900 ft., vii, viii, 17 [*Kemp*]. Types in Indian Museum.
- bipunctata**, *id.*, *loc. cit.*, 291, ♂ (1918).
Taken in company with the above.
- tritincta**, *id.*, *loc. cit.*, 291, ♂ ♀, pl. viii, 9, wing (1918).
Taken in company with *L. confinis*.
Type, ♂ ♀ in cop., in Indian Museum.
- 5-notata**, *id.*, *loc. cit.*, 292, ♂, pl. viii, 8, wing (1918).
Taken in company with *L. confinis*.
Unique type in Indian Museum.
- longipennis**, *id.*, *loc. cit.*, 292, ♂ (1918).
Taken in company with *L. confinis*.
Unique type in Indian Museum.
- nigrescens**, *id.*, *loc. cit.*, 293, ♂ (1918).
Taken in company with *L. confinis*.
Type in Indian Museum.
- punctithorax**, *id.*, *loc. cit.*, 293, ♂, fig. 1, thorax, side view (1918).
Talewadi, N. Kanara Distr., 9—10-x-16 [*Kemp*].
Unique type in Indian Museum.
- trigonia** Edw., Jour. Fed. Malay Sts. Mus. VIII, 15, ♂, pl. iii, fig. 4 a b, genitalia; pl. iv, fig. 10, wing (1919).
Siolak Daras, Sumatra, 3000 ft., iii-14, a unique ♂.
- striopleura**, *id.*, *loc. cit.*, 16, ♀ (1919).
Korinchi Peak, Sumatra, 7300 ft., v-14, a unique ♂.
- sp. incert.: **aterrima** Walk., Ins. Saund. Dipt., 434, ♀ (1856).
Brun., Rec. Ind. Mus. VI, 271.
East India.
Type, location unknown, not in the British Museum. Osten Sacken has suggested this may be an *Eriocera*.

Rhipidia, Mg.

Syst. Besch. I, 153 (1818).

Sch., F. Austr. II, 564: Ost. Sack., Monog. Dipt. N. Amer. IV, 81: *id.*, Berl. Ent. Zeits. XXXI, 174: *Auctt.*GENOTYPE, *R. maculata* Mg., by original designation.*Ceratostephanus*, Brun., Rec. Ind. Mus. VI, 271 (1911).GENOTYPE, *C. antennatus*, *id.*, sp. nov., *loc. cit.***javanensis** de Meij., Tijd. v. Ent. LIV, 31, ♂ (1911).

Alex., Proc. U.S. Nat. Mus. XLIX, 164, pl. xlii, 9, wing.

Tosari, Java, iii-1910 [*Beaufort*].

Tjibodas, Mt. Gedé, Jaya, 4500, 8000, 9000 ft.

Type in Amsterdam Museum.**antennatus** Brun., Rec. Ind. Mus. VI, 272, ♂ (*Ceratostephanus*) (1911).*Id.*, Fauna Brit. Ind., 407, ♂, pl. xi, 17, antenna.Simla, 24-iv-07 [*Annandale*].*Type*, a unique ♂, in Indian Museum.**bioculata** de Meij., Tijd. v. Ent. LVIII, Supp., 11, ♂ (1913).

Lasikin, Simalur Is., Sumatra, iv.

Unique *type* in Amsterdam Museum.**rostrifera** Edw., Ann. Mag. Nat. Hist. (8) XVII, 352, ♂ (1916).Kedah Peak, 3200 ft., Malay Peninsula [*Stanton*].*Type* in British Museum. Originally described from a single specimen but both sexes subsequently received from the same locality: one ♀ from Korinchi Peak, Sumatra.**Atypophthalmus, Brun.**

Rec. Ind. Mus. VI, 273 (1911).

Brun., Fauna Brit. India, 408.

GENOTYPE, *A. holopticus* Brun., by original designation.**holopticus** Brun., Rec. Ind. Mus. VI, 273 (1911).*Id.*, Fauna Brit. Ind., 408, ♂ ♀, pl. xi, 9, ♂ genitalia.Calcutta, 8-ix-10, *type* ♂ in spider's web, 20-viii 09, *type* ♀ in house at night [both *Annandale*].*Types*, unique, in Indian Museum.Alexander queries the validity of this genus and would merge it in *Rhipidia* Mg., but he overlooks the fact that the antennae are of normal cylindrical form and not pectinate.**Goniodineura, Wulp.**

Tijd. v. Ent. XXXVIII, 37 (1895).

GENOTYPE, *G. nigriceps*, *id.*, sp. nov., *loc. cit.*

nigriceps Wulp, *loc. cit.*, 38, pl. II, 3, ♀ (1895).

Alex., Proc. U.S. Nat. Mus. XLIX, 165, pl. xlii, 10, wing.

Java [*t. Wulp*]; Buitenzorg, iii-09 [*Bryant and Palmer*]; Pulu Pandjang Is. off W. Coast, Sumatra, ii.

Type lost; originally in Amsterdam Museum.

Dapanoptera, Ost. Sack. in Westw.

Trans. Ent. Soc. Lond., 365 (1881).

Ost. Sack., Berl. Ent. Zeits. XXXI, 178.

GENOTYPE, *Limnobia perdecora* Walk., by my designation.

perdecora Walk., Proc. Linn. Soc. V, 230, ♂ (*Limnobia*) (1861).

Papua. *Type* in British Museum in good condition.

auroatra Walk., *loc. cit.*, VII, 202, ♂ (*Limnobia*) (1864).

Mysol. *Type* in British Museum, in good condition.

plenipennis Walk., *loc. cit.*, VIII, 103, ♂ (*Limnobia*) (1865).

Westw., Trans. Ent. Soc. Lond., 366, pl. xvii, 2 (1881).

Papua. *Type* in British Museum, in fair condition.

latifascia Walk., Proc. Linn. Soc. Lond., VIII, 104, ♀ (*Limnobia*) (1865).

Papua. *Type* in British Museum, in good condition.

lorentzi de Meij., Nov. Guin. Res. IX, 307, ♂ ♀, pl. x, 1, wing (1913).

S.W. Papua; Bivak Is., Regen Is.

Type in Amsterdam Museum.

fascipennis, *id.*, *loc. cit.*, 307, ♂ (1913).

Alkmaar, S.W. Papua. Unique *type* in Amsterdam Museum.

pallida, *id.*, *loc. cit.*, 307, ♂ (1913).

Alkmaar, S.W. Papua, x, 1909. Utakwa River, 4000—6000 ft., Dutch Papua, i, ii-13. *Type* in Amsterdam Museum.

pulchra de Meij., Tijd. v. Ent. LVIII, 103, ♂ (1915).

North Papua.

versteegi de Meij., Nov. Guin. Res. XIII, 52 (1915).

Papua.

Libnotes, Westw.

Trans. Ent. Soc. Lond., 505 (1876).

Ost. Sack., Berl. Ent. Zeits. XXXI, 179, notes and tab. spp.

Brun., Rec. Ind. Mus. VI, 278, notes on oriental spp.: *id.*, Fauna Brit. Ind., 409.

GENOTYPE, *L. thwaitesiana* Westw., by original designation.

- impressa* Walk., Proc. Linn. Soc. Lond. I, 106, ♀ (*Limnobia*) (1857).
Sarawak. Type in British Museum, useless for comparison.
- aurantiaca* Dol., Nat. Tijd. Ned. Ind. XVII, 78, ♂ (*Limnobia*) (1858).
Ost. Sack., Berl. Ent. Zeits. XXXI, 181: Brun., Rec. Ind. Mus. VI, 278.
Amboina. Type, damaged, in Vienna Museum.
- imponens* Walk., Proc. Linn. Soc. Lond. IV, 92, ♀ (*Limnobia*) (1860).
Makassar. Type in British Museum, useless for comparison.
- strigivena* Walk., *loc. cit.*, V, 229, ♂ (*Limnobia*) (1861).
de Meij., Tijd. v. Ent. LIV, 35, notes: *loc. cit.*, LVI, 346: Brun., Rec. Ind. Mus. VI, 278.
Papua (Dorey). Ungaren, ix; Nongkodjadjar [both Java and Jacobson]; Mt. Silak, Java, 15-v-09 [Bryant and Palmer]. Type in British Museum, in bad condition, but the wing markings are sufficiently conspicuous to identify the species. De Meijere (*loc. cit.*) notes the apparent ♀ from Bandung, Java, vi [Preanger].
- quadrifurca* Walk., Proc. Linn. Soc. Lond. V, 230, ♀ (*Limnobia*) (1861).
Papua (Dorey). Type in British Museum, in bad condition: the wing markings conspicuous.
- innotabilis* Walk., *loc. cit.*, VII, 231, ♀ (*Limnobia*) (1864).
Ceram. Type in British Museum, in bad condition.
- ¹*thwaitesiana* Westw., Trans. Ent. Soc. Lond. 505, ♂, pl. iii, 6 (1876).
L. rufa de Meij., Tijd. v. Ent. LIV, 39, ♀, pl. ii, 24, wing (1911).
Id., Brun., Rec. Ind. Mus. VI, 279 (♂ descr.): *id.*, Fauna Brit. Ind., 414, ♂ ♀.
Ceylon; Semarang, iii [Jacobson]; Calcutta, 5-viii-08 [Annandale]; Pusa, 17-xi, 2-xii-13, both on bark of jackfruit tree; Peradeniya, 13, 17-vii-10 [Gravely]; x, xi, 11 [t. Edwards]; Talewadi, N. Kanara dist. [Kemp]. Type (*thwaitesiana*) in Hope coll., Oxford University Museum, in inferior condition, but recognition possible; type of *rufa* in Amsterdam Museum.
- notata* Wulp, Tijd. v. Ent. XXI, 194, pl. xii, 5, ♂ (1878).
Id., Dipt. Sum. Exp. 12: de Meij., Tijd. v. Ent. LIV, 34 (♀ descr.) (1911); Enderl., Zool. Jahr. XXXII, 82: Brun., Rec. Ind. Mus. VI, 279.

¹ Omitted by error from my Fauna volume, but *thwaitesiana* was included as *rufa*, Meij.

Sumatra; Laulo, Simalur Is., nr. Sumatra, viii; Semarang, iii, xi [Jacobson]; Myithyina, Up. Burma, 30-viii—1-ix-14 [Fletcher]; Peradeniya [Gravely]; Talewadi [Kemp].

Type originally in Amsterdam Museum, now lost; other specimens present.

simplex Ost. Sack., Ann. Mus. Gen. XVI, 402, ♂ (1880).

Ternate. Type in Genoa Museum in poor condition, apparently unique.

¹ **poeciloptera**, *id.*, *loc. cit.*, 403, ♂ ♀ (1880).

Brun., Rec. Ind. Mus. VI, 279; Enderl., Zool. Jahr. XXXII, 82.

Pangerando, Java, x [Jacobson]; Tenggergebirge, Java [Piepers]; Sumatra; Pundaluoya, Ceylon [Green]; Sinabang, ii; Sibigo, viii, both Simalur Is., off Sumatra. Types originally in Genoa Museum, but only one specimen now remains.

semperi Ost. Sack., Berl. Ent. Zeits. XXVI, 88, ♀ (1882).

Philippines. Type, unique, originally in Osten Sacken's collection, now apparently lost.

termitina Ost. Sack., *loc. cit.*, 88, ♂ ♀ (1882).

Philippines. Types, a unique pair, originally in Osten Sacken's collection, now apparently lost.

familiaris Ost. Sack., *loc. cit.*, 89, ♂ (1882).

de Meij., Tijd. v. Ent. LIV, 37, ♂ ♀ descr., pl. ii, 22, wing; *id.*, Nova Guin. Res. IX, 309; Brun., Rec. Ind. Mus. VI, 279, notes.

Philippines; Wonosobo, Java, v; Batavia, ii [both Jacobson]; Riverkamp, S. W. Papua, 12-ii-10 [Lorentz]; Sinabang, Simalur Is., off Sumatra, iii.

Type originally in Osten Sacken's collection, now apparently lost.

punctipennis de Meij., Tijd. v. Ent. LIV, 35, ♂ ♀, pl. ii, 20, wing (1911).

Brun., Rec. Ind. Mus. VI, 279; *id.*, Fauna Brit. Ind., 413, ♂ ♀, pl. viii, 6, wing.

Darjiling, 6-viii-09 [Paiva]; Kavalai, Cochin, 1300—3000 ft., 24—27-ix-14 [Gravely]; Parambikulam, Cochin, 1700—3200 ft., 16—24-ix-14 [Gravely]; Talewadi and Castle Rock, N. Kanara Dist. [Kemp]; Tura, Garo Hills, vii-17 [Kemp]; Mazbat, Assam, 11—15-x-10 [Kemp]; Dibrugarh, 17—19-xi-11 [Kemp]; Peradeniya, 5-viii-10, 15-x-09 [Green; Gravely]; Batavia, ii; Wonosobo, iv, Semarang, i [all Java and Jacobson].

Type in Amsterdam Museum.

nervosa de Meij., Tijd. v. Ent. LIV, 36, ♂, pl. ii, 21, wing (1911).

Wonosobo, v; Semarang, ii [both Java and Jacobson]; Sinabang, ii, vi; Lasikin, iv; Pulu Babi, iv, all off W. Coast Sumatra; Buitenzorg, 9-iv-09 [Bryant and Palmer].

Type in Amsterdam Museum.

¹ Omitted by error from my Fauna volume.

- forcipata**, *id.*, *loc. cit.*, 38, ♂ ♀, pl. ii, 23, wing (1911).
Wonosobo, Java, iv [Jacobson]. Types in Amsterdam Museum, a unique pair.
- fuscinervis** Brun., Fauna Brit. Ind., 411, ♂, pl. viii, 5, wing, pl. xi, 8, ♂ genitalia (1912).
Id., Rec. Ind. Mus. XV, 295.
Darjiling, 9-viii-09 [Paiva]; 5-viii-09; 22-ix-08 [Brunetti]; Kalimpong, 600—4500 ft. [Gravelly]; N.E. Ind. Frontier, 4000 ft., 18-iii-12 [Kemp]. Type in Indian Museum.
- notatinervis** Brun., Fauna Brit. Ind., 412, ♀, pl. viii, 7, wing (1912).
Kurseong, 6-ix-09 [Annandale]; Ghumti, 4000 ft. [Gravelly]; Pattipola, Ceylon, 6000 ft. [Annandale]; Shillong [Kemp]. Type in Indian Museum.
- affinis de Meij.**, Nova Guin. Res. IX, 308, ♀, pl. x, 2, wing (1913).
Id., Tijd. v. Ent. LVI, 346.
Bivak Is., S.-W. Papua, xii; Gunung, Ungaren, ix [Jacobson]. Type in Amsterdam Museum, unique.
- punctatissima de Meij.**, Tijd. v. Ent. LVIII, 58, ♂ (1915).
North Papua. Unique type in Amsterdam Museum.
- scutellata** Edw., Ann. Mag. Nat. Hist. (8) XVII, 353, ♂ ♀, fig. 2, p. 356, genitalia (1916).
Talum, Perak-Pahang boundary, 18-i-02 [Robinson; Annandale].
Type in British Museum.
- stantoni**, *id.*, *loc. cit.*, 354, ♀ (1916).
Kedah Peak, Malay Peninsula, 3200 ft. [Stanton].
Type in British Museum, a unique ♀.
- limpida**, *id.*, *loc. cit.*, 355, ♀ (1916).
Taken in company with above.
Type, a unique ♀, in British Museum.
- lutea**, *id.*, *loc. cit.*, 357, ♀ (1916).
Taken in company with *stantoni*.
Type in British Museum, a unique specimen.
- transversalis de Meij.**, Tijd. v. Ent. LIX, 198 (1916).
Gedeh, Java, 1625—2400 metres, v; 1 ♂ [Konigsberger] (1916). Unique type in Amsterdam Museum.
- sumatrana** Edw., Jour. Fed. Malay Sts. Mus. VIII, 17, ♂, pl. iii, fig. 5, genitalia; pl. iv, fig. 11, wing (1919).
Korinchi Peak, Sumatra, 7300 ft. v-14; Sungei Kumbang, Sumatra, 4500 ft., iv-14.
- regalis** Edw., *loc. cit.*, XVIII, 248 (1916).
Taihoku, Formosa [Shiraki]. The much damaged unique type in British Museum.
- montivagans** Alex., Proc. U.S. Nat. Mus. XLIX, 166, ♂ ♀, pl. xliii, 12, wing (1916).
Tjibodas, Mt. Gedé, Java, 8000 ft. [Bryant and Palmer].
Type, Cat. No. 19,015, in U.S. Nat. Mus.

- nigricornis*, *id.*, *loc. cit.*, 166, ♂ ♀, pl. xliii, 13, wing (1916).
Tjibodas, Mt. Gedé, Java, 4,500 ft. [Bryant and Palmer].
Type, Cat. No. 19,016, in U.S. Nat. Mus.
- opaca* Bezzi, Phil. Jour. Sci. XII, Sect. D, 116, ♂ (1917).
Luzon [Baker].
- marginalis*, *id.*, *loc. cit.*, 116, ♂ (1917).
Luzon [Baker].

Section RHAMPHIDINI.

Rhamphidia, Mg.

Syst. Besch. VI, 281 (1830).

Sch., F. Austr. II, 558: Ost. Sack., Monog. Dipt. N. Amer. IV, 103: Berl. Ent. Zeits. XXXI, 183: Brun., Fauna Brit. Ind., 416: *Auctt. var.*GENOTYPE, *Limnobia longirostris* Mg., by Westwood's designation (1840).*Megarhina*, St. Farg. et Serv., Encyc. Meth. Ins. X, 585 (1825).GENOTYPE, *Limnobia longirostris* Mg., by original designation.*Helius*, *id.*, *loc. cit.*, 831 (1835).Nom. nov. for *Megarhina* under supposed pre-occupation by *Megarhinus* Rob. Desv.*Leptorhina*, Steph., Cat. Brit. Ins. II, 243 (1829).

GENOTYPE, unknown to present author.

ferruginosa Brun., Fauna Brit. Ind., 418, ♂, fig. 41, head, profile; pl. viii, 8, wing, pl. xi, 11, ♂ genitalia (1912).Pashok, Darjiling Distr., 2000—3500 ft., 23-iv—11-v-15 [Gravelly]; base of Dawna Hills, Lower Burma, 2-iii-08 [Annandale]. *Type* in Indian Museum.*unicolor*, *id.*, *loc. cit.*, 419, ♂ (1912).*Id.*, Rec. Ind. Mus. XV, 296.

Darjiling, 7000 ft., 8-viii-09 [Paiva].

Type in Indian Museum.*inconspicua* Brun., Fauna Brit. Ind., 419, ♂ (1912).

Kurseong, 16-iv-11 [Annandale]; Kalimpong, 600—4500 ft., 24-iv—10-v-15 [Gravelly]; Pachmarhi, 20-ix-13, near waterfall.

Type in Indian Museum.*kambangani* de Meij., Tijds. v. Ent. LVI, 346, pl. xvii, 13, wing (1913).

Nusa, Kambangan, an island off S. coast of Java [Jacobson].

Type in Amsterdam Museum.*apicalis* Alex., Proc. U.S. Nat. Mus., XLIX, 167, ♂ ♀ pl. xliii, 15, wing (1916).Tjibodas, Mt. Gedé, Java, 4500 ft., 8000 ft. [Bryant and Palmer]. *Type*, No. 19017, in U.S. Nat. Mus.

- nigriceps** Edw., Ann. Mag. Nat. Hist. (8) XVII, 358, ♂ ♀ ; fig. 3, p. 356, genitalia (1916).
Bukit Besar, Siam, type ♂ ; Talum, Perak-Pahang boundary, 18-i-01 [Robinson and Annandale]. Types in British Museum.
- rufescens** Edw., loc. cit., 358, ♀ (1916).
Bukit Kutu, Selangor, ii-03 [Durham].
Type in British Museum.
- fratella** Brun., Rec. Ind. Mus., XV, 296, ♂ ♀ (1918).
Castle Rock, N. Kanara Dist., 11-26-x-16 [Kemp]: Ginting Bidai, 2000 ft.; Selangor-Pahang Frontier, Federated Malay States, iv-17 [C. Boden Kloss]. Types in Indian Museum.
- abnormalis** Brun., Rec. Ind. Mus., XV, 296, ♀ , pl. vii, 3, wing (1918).
Cherrapunji, Assam, 4400 ft., 2-3-x-14 [Kemp].
The unique type in the Indian Museum.

Subgenus *Eurhamphidia*, Alex.

- Proc. U. S. Nat. Mus. XLIX, 168 (1916).
GENOTYPE, *Rhamphidia niveitarsis* Skuse, by original designation.
- monticola** Edw., Jour. Fed. Malay Sts. Mus., VIII, 18, ♀ , pl. iv, fig. 12, wing (1919).
Korinchi Peak, Sumatra, 7300 ft., iv-14.
Type in British Museum.
- niveitarsis** Skuse, Proc. Linn. Soc. N.S. Wales (2) IV, 791 (*Rhamphidia*) (1889).
Alex., Proc. U.S. Nat. Mus. XLIX, 168, ♂ ♀ , pl. xliii, 16, wing.
Tjibodas, Mt. Gedé, 8000, 9,000 ft.; Pangranggo, 9000 ft. [both Java and Bryant and Palmer].
Type: location unknown to present author.

Sub-genus, *Rhampholimnobia*, Alex.

- Proc. U.S. Nat. Mus. XLIX, 169 (1916).
GENOTYPE, *R. reticularis*, id., sp. nov., loc. cit.
- reticularis** Alex., loc. cit., 169, ♀ , pl. xliii, 17, wing (1916).
Tjibodas, Mt. Gedé, Java, 4,500 ft. [Bryant and Palmer].
Type, Cat. No. 19018, in U.S. Nat. Mus.

Limnobiorrhynchus, Westw.¹

- Ann. Soc. Ent. Fran. IV, 683, ♀ only (1835).
Brun., Rec. Ind. Mus. XV, 298.
GENOTYPE, *Limnobiorrhynchus westwoodi*, nom. nov. for *L. brasiliensis*, Westw., sp. nov., ♀ only; by my own designation, 1918.

¹ See Rec. Ind. Mus. XV, 297 on the *Elephantomyia* group.

incertus Brun., Fauna Brit. Ind., 422 ♀; pl. viii, 9 wing; xi, 12, ♀ genitalia (*Toxorhina*) (1912).¹

Id. Rec. Ind. Mus. XV, 298.

Kurseong, 27-vi-10 [Annandale]. Unique type in Indian Museum. The genus *Toxorhina* does not now appear to be oriental.²

Ceratocheilus, Wesché.

Jour. Linn. Soc. XXX, 358 (1910).

GENOTYPE, *C. winn-sampsoni*, sp. nov., *loc. cit.*, the 1st species, by present designation.

Neostyringomyia, Alex., Canad. Ent. XLIV, 85 (as subgenus of *Styringomyia*) (1912).

GENOTYPE, *Styringomyia cornigera* Speis., by original designation.

Conithorax, Brun., Rec. Ind. Mus. XV, 298 (1918).

GENOTYPE, *C. latifrons*, sp. nov., *loc. cit.*

I have to thank Mr. Chas. Alexander for the above synonymy (*in litt.*). *Ceratocheilus* at its inception was considered by Wesché to represent a new subfamily having affinities with the Ptychopterinae. The exact position of *Styringomyia* is also uncertain. Some authors refer it to the Rhamphidiini, others to the Eriopterini.

latifrons Brun., *loc. cit.* 299, ♀, fig. 2, antenna, 3, wing (*Conithorax*) (1918).

Bidai, Selangor-Pahang Frontier, Federated Malay States, iv-17 [*C. Boden Kloss*]. Unique type in Indian Museum.

brevifrons, *id.*, *loc. cit.*, 300, ♀ (*Conithorax*) (1918).

Above Tura, Garo Hills, 3500—3900, viii-17 [*Kemp*]. Unique type in Indian Museum.

Elephantomyia, Ost. Sack.

Proc. Acad. Nat. Sci. Philad. 220 (1859).

Ost. Sack., Monog. Dipt. N. Amer. IV, 106, pl. i, 5, wing, iii, 8, forceps: *id.*, Berl. Ent. Zeits. XXXI, 184; Brun., Rec. Ind. Mus. XV, 297, *Elephantomyia* group.

GENOTYPE, *E. westwoodi* Ost. Sack. (as *Limnobia rhynchus canadensis*, Westw.), by original designation.

¹ In the note after the description of this species the first sentence is misleading. In the description itself it is stated that the discal cell is open in one wing and closed in the other in the unique type. In penning the phrase commencing "though the discal cell is open, etc. (Fauna, p. 423), I must have overlooked this anomaly and observed only the left wing, in which the cell is open; and, although this explanation makes the text intelligible, it admittedly does not excuse a want of care in examination. My figure shews the cell closed as it normally is in this genus and as is stated both in my generic description (*l.c.* p. 421), and in Loew's.

² See Rec. Ind. Mus. XV, 300, for discussion.

- argento cincta** Walk., Proc. Linn. Soc. I, 107, ♀ (*Limnobia*) (1857).
Sarawak. Type in British Museum, useless for comparison and apparently unique.
- delectata**, *id.*, *loc. cit.*, VII, 231, ♂ ♀ (*Limnobia*) (1864).
Ceram. Types in British Museum, useless except for conspicuous wing markings.
- filiformis**, *id.*, *loc. cit.*, VIII, 131, ♂ (*Limnobia*) (1865).
Salawatti.
Type in British Museum, valueless, apparently unique.
- fuscomarginata** Ender., Zool. Jahr. XXXII, 64, ♂ fig. N, 1, wing (1912).
Sumatra (Sinabong).
Type, a unique ♂, in Stettin Zoological Museum.
- egregia** de Meij., Tijd. v. Ent. LVI, 347, ♂, pl. xvii, 14, wing (1913).
Nongkodjadjar, Java, i [*Jacobson*].
Type, a unique ♂, in Amsterdam Museum.

Dicranoptycha, Ost. Sack.

- Proc. Acad. Nat. Sci. Philad., 203 and 217 (1859).
Ost. Sack., Monog. Dipt. N. Amer., IV, 116, pl. i, 8, wings.
GENOTYPE, *D. germana* Ost. Sack., by Coquillett's designation.
- signaticollis** Wulp, Tijd. v. Ent. XXXVIII, 36, pl. ii, 1, ♂ (1895).
Brun., Rec. Ind. Mus. VI, 280.
Java. Type lost, originally in Amsterdam Museum.

Orimarga, Ost. Sack.

- Monog. Dipt. N. Amer. IV, 120, pl. i, 9, wing (1869).
Ost. Sack., Berl. Ent. Zeits. XXXI, 186:
Brun., Fauna Brit. Ind., 423.
GENOTYPE, *Limnobia alpina* Zett., by original designation.
- Ninguis**, Wallengren, Entom. Tidskr. II, 183 (1881).
GENOTYPE, *Limnobia virgo* Zett., by Coquillett's designation.
- ? **Spyloptera**, Rond., Dipt. Ital. Prod. I, 181 (1856).
- borneensis** Brun., Rec. Ind. Mus. VI, 280, ♀ (1911).
Sarawak, 10 miles s. of Kuching, 27-vi-10 [*Beebe*]. Type in Indian Museum.

peregrina, *id.*, Fauna Brit. Ind., 424, ♂, pl. viii, 11, wing (1912).

Brun., Rec. Ind. Mus. XV, 309.

Kurseong, 29-vi-10, [Annandale]; Ghumti, Darjiling Distr., vii-11 [Gravely]; Pashok, 3500 ft. [Hartless].

Type, a unique specimen, in Indian Museum.

javana de Meij., Tijds. v. Ent. LVI, 348, ♀, pl. xvii, 15, wing (1913).

Nongkodjadar, i [Jacobson].

Type, a unique ♀, in Amsterdam Museum.

Antocha, Ost. Sack.

Proc. Acad. Nat. Sci. Philad., 219 (1859).

Sch., F. Austr. II, 558.

Ost. Sack., Berl. Ent. Zeits. XXXI, 187; Brun., Fauna Brit. Ind., 425.

GENOTYPE, *A. opalizans* Ost. Sack., as *saxicola* Ost. Sack.

Taphrophila, Rond., Dipt. Ital. Prod. I, 185 (1856).

GENOTYPE, *Limnobia inusta* Mg., by original designation.

Orimargula, Mik., Wien. Ent. Zeitg. II, 198, fig. (1883).

GENOTYPE, *O. alpigena*, sp. nov., *loc. cit.*

indica Brun., Fauna Brit. Ind., 426, ♂ ♀, pl. viii, 12, wing (1912).

Phagu, 12-v-09; Theog., 2-v-07 [all Annandale]; Kurseong, 9-ix-09, *type* ♀; 16-iv-11, *type* ♀; Kalimpong, 600—4500 ft., 1—24-iv—10-v-15 [Gravely]; Pashok 4500 ft. [Gravely]; Almora, Kumaon, 5500 ft., 11—12-vi-11 [Paiva]; Darrang Distr., Assam-Bhutan Frontier, 26-xii-10 [Kemp].

Type in Indian Museum.

unilineata, *id.*, *loc. cit.*, 427, ♀ (1912).

Mundali, Dehra Dun, 9000 ft., 10-v-10.

Type, unique, in Indian Museum.

javanensis, Alex., Proc. U.S. Nat. Mus. XLIX, 171, ♀, pl. xliii, 18, wing (1916).

Pelaboean Ratoe, Java, 19-x-09; [Bryant and Palmer].

Type, Cat. No. 19,019, in U.S. Nat. Mus.

Teucholabis, Ost. Sack.

Proc. Acad. Nat. Sci. Philad., 222 (1859).

Ost. Sack., Monog. Dipt. N. Amer. IV, 129,

pl. i, 12, wing; pl. iii, 9, forceps: *id.*, Berl.

Ent. Zeits. XXVI, 188: *id.*, *loc. cit.*, XXXI,

188: Brun., Fauna Brit. Ind., 427.

GENOTYPE, *T. complexa* Ost. Sack., by original designation.

- exclusa** Walk., Proc. Linn. Soc. VIII, 105, ♀ (*Limnobia*) (1865).
Papua. Type in British Museum, much damaged.
- bicolor** Ost. Sack., Ann. Mus. Gen. XVI, 404, ♂ (1880).
de Meij., Tijd. v. Ent. LVI, 348.
Sumatra; Sinabang, Simalur Is., off Sumatra, i, ii; Gunung
Ungaren, Java, ix [*Jacobson*].
Type in Genoa Museum in poor condition.
- fenestrata** Ost. Sack., Berl. Ent. Zeits. XXXI, 188 ♂ ♀ (1887).
Brun., Rec. Ind. Mus. VI, 283; XV, 304: *id.*, Fauna Brit.
Ind., 429, ♂ ♀.
Common and widely distributed in India, from April to
October. Naini Tal, Darjiling, Kurseong, Lake Chilka, Mad-
ras, Peradeniya; Mazbat, Assam; Tura, Garo Hills.
Types in the Leyden Museum and Bigot collection.
- determinata** Ost. Sack., Berl. Ent. Zeits. XXXI, 188, ♂ (1887).
Brun., Rec. Ind. Mus. VI, 283.
Sula (Indo-Malay Archipelago).
Type, a unique ♂, in British Museum.
- insignis** Brun., Fauna Brit. Ind., 430, ♂, pl. viii, 13, wing (1912).
Tenmalai, Travancore, 22-xi-08 [*Annandale*].
Type, a unique ♂, in Indian Museum.
- bi-annulata**, *id.*, *loc. cit.*, 430, ♂ (1912).
Kurseong, 26-vi-10. [*Annandale*]; Rotung, 27-xii-11 and
Damda, 3000 ft., both N.-E. Ind. Frontier [*Kemp*].
Type in Indian Museum.
- plecioides** de Meij., Tijd. v. Ent. LVI, 348, ♀ (1913).
Nongkodjadjar, Java, i; Sinabang, Simalur Is. off Sumatra,
ii, vii [*Jacobson*]. Type in Amsterdam Museum.
- glabripes**, *id.*, *loc. cit.*, 349, ♂ (1913).
Nongkodjadjar, Java, i [*Jacobson*].
Type, a unique ♂, in Amsterdam Museum.
- femorata**, *id.*, *loc. cit.*, LVIII, 67, ♂ (1916).
Fort de Kock, Simalur Island, Sumatra, x.
The unique type in Amsterdam Museum.
- nigerrima** Edw., Ann. Mag. Nat. Hist. (8) XVIII, 248 (1916).
Horisha, Formosa, 10-v-13 [*Maki*]; Taihoku, Formosa
[*Shiraki*]. Type in British Museum.
- angusticapitis** Brun., Rec. Ind. Mus. XV, 305, ♂, pl. viii, 11, head
and thorax, 12, head in profile, 13, tip of proboscis (1918).
Above Tura, Garo Hills, 3500—3900 ft., viii-17 [*Kemp*].
Unique type in Indian Museum.
- ornata**, *id.*, *loc. cit.*, 305, ♂ (1918).
Peradeniya, 11-viii-10. Unique type in Indian Museum.
var. *assamensis*, *id.*, *loc. cit.*, 306, ♂ (1918).
Shillong, 5500—6400 ft., 29-viii—5-ix-15 [*Kemp*].

nocticolor Edw., Jour. Fed. Malay Sts. Mus., viii, 18, ♂ ♀, pl. iii, fig. 6a, b, genitalia; pl. iv, 13, wing (1919).
Siolak Daras, Sumatra, 3000 ft.

Gymnastes, Brun.¹

Rec. Ind. Mus. VI, 281 (1911).

Id., *id.*, Fauna Brit. Ind., 432: Rec. Ind. Mus. XV, 307: Edw., Ann. Mag. Nat. Hist. (8) XVII, 359.

GENOTYPE, *Teucholabis cyanea* Edw., as *G. violaceus*, sp. nov.

cyanea Edw., Ann. Mag. Nat. Hist. (8) VIII, 61, ♂ ♀ (*Teucholabis*) (1911).

Gymnastes violaceus, Brun., Rec. Ind. Mus. VI, 282 (1911).

id., Fauna Brit. Ind., 433, ♂, fig. 42 full ins. (1912).

Madulsima, Ceylon, 19-v-08, type ♂ [*Fletcher*]; Bentota, Ceylon, 13-vi-90, type ♀ [*Yerbury*]; Kandy, 22-v-10 (type of *violaceus*); Peradeniya, 15-vii-10 [*Gravelly*]; Cherambadi, Wynaad, S. India, x-13 [*Fletcher*]. Type ♂ ♀ *cyanea* in British Museum, of *violaceus* in Indian Museum.

ornatipennis de Meij., Tijds. v. Ent. LIV, 47, pl. iii, 34, wing (*Gnophomyia*) (1911).

Gymnastes, *id.*, Edw., Ann. Mag. Nat. Hist. (8) XVIII, 249.

Paratropesa, *id.*, Alex., Proc. U.S. Nat. Mus. XLIX, 167, pl. xliii, 14, wing.

Wonosobo, Java, iv; Telaga Mendjer, Java, v [both *Jacobson*]; Tjibodas, Mt. Gedé, Java, 4500 ft. [*Bryant and Palmer*]; Arisan, Formosa, 8000 ft., 10-x-12 [*Nitobe*]; Reidel also records it from Formosa.

Type in Amsterdam Museum, no sex stated.

Alexander refers this species to *Paratropesa*, Sch., I think wrongly.

pictipennis Edw., Ann. Mag. Nat. Hist. (8) xvii, 358, ♂, fig. 4, p. 356, genitalia (1916).

Bukit Besar, Malay Penin. [*Robinson and Annandale*].

Type in British Museum.

bistriatipennis Brun., Rec. Ind. Mus. XV, 307, ♂ (1918).

Talewadi, N. Kanara Dist., 3-10-x-16 [*Kemp*].

Type in Indian Museum; cotypes in my collection.

pennipes, *id.*, loc. cit., 308, pl. viii, 10, hind leg (1918).

Above Tura, Garo Hills, 3500-3900, vii-17 [*Kemp*].

Unique type in Indian Museum.

¹ Alexander regards this genus as synonymous with *Teucholabis*, but I still think it distinct (Rec. Ind. Mus. XV, 307) and Edwards supports its retention (Ann. Mag. Nat. Hist. (8) xvii, 358).

Atarba, Ost. Sack.

Monog. Dipt. N. Amer. IV, 127, pl. i, 13, wing (1869).

Brun., Fauna Brit. Ind., 434.

GENOTYPE, *A. picticornis*, *id.*, sp. nov., *loc. cit.*

Leiponeura, Skuse, Proc. Linn. Soc. N. S. Wales (2)
IV, 795 (1890).

GENOTYPE, *L. gracilis*, sp. nov., *loc. cit.*, the 1st
species by present designation.

nebulosa de Meij., Tijds. v. Ent. LIV, 42, ♂, pl. ii, 25, wing
(1911).¹

Batavia, viii [Jacobson]. Type in Amsterdam Museum.

pilifera, *id.*, *loc. cit.*, 43, ♂ ♀, pl. ii, 26, wing (1911).

Semarang, vi, vii [Jacobson]. Types in Amsterdam Museum.

diffusa, *id.*, *loc. cit.*, 43, ♀ (1911).

Semarang, i, xi [Jacobson]. Type in Amsterdam Museum.

flava Brun., Fauna Brit. Ind., 435, ♂ ♀, pl. viii, 15, wing (1912).

Darjiling, apparently common in autumn, 28-v-10, type ♂; 26-
v-20, type ♀; 22-ix-08, 1-x-08 [all Brunetti]; 9-viii-09 [Paiva].
Types in Indian Museum.

pallidicornis Edw., Ann. Mag. Nat. Hist. (8) XVIII, 249, ♀
(1916).

Arisan, Formosa, 8000 ft., 10-x-12 [Nitobe].

Unique type in British Museum.

fuscicornis Edw., *loc. cit.*, 250, ♀ (1916).

Arisan, Formosa, 8000 ft., 10-x-12 [Nitobe].

Unique type in British Museum.

javanica Alex., Proc. U.S. Nat. Mus. XLIX, 171, ♀, pl. xliii, 19,
wing (1916).

Tjibodas, Mt. Gedé, Java, 8000 ft. [Bryant and Palmer].

Type, Cat. No. 19,020, U.S. Nat. Mus.

Section ERIOPTERINI.²

Eriopterina, Ost. Sack., Eriopterinae, Kert.

Rhypholophus, Kol.

Wien. Ent. Monats. IV, 393 (1860).

Ost. Sack., Monog. Dipt. N. Amer. IV, 139, pl. i,

14, 15, wings: *id.*, Berl. Ent. Zeits. XXXI, 192:

Sch., F. Austr. II, 536: Brun., Fauna Brit. Ind.,

440, pl. xi, 20, antenna.

¹ Alexander regards *Leiponeura* as valid and places therein *nebulosa*, *pilifera* and *diffusa*. He thinks my *flava* the only oriental species of the genus *Atarba* except his own *javanica*.

² For characters and table of genera *v.* Brun., Fauna Brit. Ind. 436 *et seq.*, and Ost. Sack., Monog. Dipt. N. Amer. IV, 135.

GENOTYPE, *Rhypholophus phryganopterus* Kol.,
the original species.

Ilisomyia, Rond., Dipt. Ital. Prod. I, 180 (*Ilisomyia*) (1896).

GENOTYPE, *I. nubipennis*, sp. nov., *loc. cit.*

Ormosia, *id.*, *loc. cit.*, 180 (1856).

GENOTYPE, *Erioptera nodulosa* Macq., by original designation.

Dasyptera, Sch., Wien. Ent. Monats. VII, 221 (1863).

GENOTYPE, *Erioptera varia* Mg., by original designation.

Erioptera, Sch., *nec* Mg.

geniculatus Brun., Fauna Brit. Ind., 441, ♂ (1912).

Kurseong, 16-iv-11 [*Annandale*].

Type in Indian Museum.

pulcher, *id.*, *loc. cit.*, 442, ♀, pl. viii, 16, wing (1912).

Phagu, Simla Distr., 9000 ft., 11-v-09 [*Annandale*]; Naini Tal, 6400 ft., Kumaon Distr., x-1906 [*Annandale*].

Type in Indian Museum.

Molophilus, Curt.

Brit. Entom. 444 (1833).

Ost. Sack., Monog. Dipt. N. Amer. IV, 243, pl.

i, 16—19, wings, pl. iv, 14—16, forceps: *id.*, Berl.

Ent. Zeits. XXXI, 192: Brun., Fauna Brit.

Ind., 443

Erioptera Sch. (*nec* Meig.), Wien. Ent. Monats.

VII, 221 (1863): *id.*, *id.*, F. Austr. II, 541.

GENOTYPE, *Erioptera atra* Mg., as *brevipennis*

Curt., sp. nov.

bicolor de Meij., Tijds. v. Ent. LIV, 45, pl. iii, 27, wing (1911).

Wonosobo, Java, v [*Jacobson*].

Unique type in Amsterdam Museum.

inconspicuus Brun., Fauna Brit. Ind., 444, ♂ ♀ (1912).

Simla, 12-v-08; Ghumti, 4000 ft. [*Gravely*]; Kurseong, 4—

6-vii-08; 18—20-vi-10, type ♂ [all *Annandale*]; Maddathorai,

16-xi-08, type ♀; Tenmalai, 22-xi-08 [both Travancore and

Annandale]. Types in Indian Museum.

assamensis, *id.*, *loc. cit.*, 445, ♂ (1912).

Sylhet, 8-i-06 [*Hall*]. Unique type in Indian Museum.

costalis Edw., Ann. Mag. Nat. Hist. (8) XVIII, 251 (1916).

Arisan, Formosa, 8000 ft., 10-x-12 [*Nitobe*].

Type in British Museum.

sirius Alex., Can. Ent. XLVII, 82, ♂ ♀, fig. 1, wing (1915).

Philippine Islands, July.

Erioptera, Mg.

Illig. Mag. II, 262 (1803).

Sch., F. Austr. II, 541: Ost. Sack., Monog. Dipt. N. Amer. IV, 146, pl. i, 16—19, wings; iv, 14—16, forceps: *id.*, Berl. Ent. Zeits. XXXI, 193. *Auctt. var.*

Brun., Fauna Brit. Ind., 445: *id.*, Rec. Ind. Mus. XV, 309, notes and synonyms.

GENOTYPE, *Erioptera lutea* Mg., by Coquillett's designation (1910).

Chemalida Rond., Dipt. Ital. Prod. I, 180 (1856).

GENOTYPE, *Erioptera taenionota* Mg., by original designation.

Limnaea, *id.*, *loc. cit.*, 181.

GENOTYPE, *Tipula flavescens* L., by original designation.

Limnoica, *id.*, *loc. cit.* IV, p. 11, Corrigenda (1861), nom. nov. for *Limnaea*, preocc. Poli, 1791.

? *Octavia* Big., Ann. Soc. Ent. Fran. (3) II, 475 (1854).

javanensis de Meij., Tijd. v. Ent. LIV, 45, ♂; pl. iii, 28, wing (1911).

Wonosobo, Java, iv [Jacobson]; Pelaboean Ratoe, Java, 16-x-09; Buitenzorg, iii, 09 [Bryant and Palmer].

Type in Amsterdam Museum.

notata, *id.*, *loc. cit.*, 46, ♂ ♀, pl. iii, 29, wing, 30, ♀ genitalia, 31, ♂ genitalia (1911).

Semarang ii [Jacobson]; Sinabang, Simalur Is., off W. Coast Sumatra, v, vi. Types in Amsterdam Museum in alcohol.

punctipennis Brun., Fauna Brit. Ind., 449, ♀, pl. viii, 17, wing (1912).

Kurseong, 26—27-vi-10 [Annandale]; Tura, Garo Hills, vii, viii, 17 [Kemp]. Type in Indian Museum.

ferruginea, *id.*, *loc. cit.*, 450, ♂ (1912).

Pallode, Travancore, 15-xi-08 [Annandale].

Type in Indian Museum.

distans, *id.*, *loc. cit.*, 451, ♂ (1912).

Kurseong, 26-vi-10 [Annandale].

Type in Indian Museum.

incerta, *id.*, *loc. cit.*, 452, ♂ (1912).

Darjiling, 5-viii-09 [Paiva].

Type, a unique ♂, in Indian Museum.

parallela, *id.*, *loc. cit.*, 453, ♀ (1912).

Kurseong, vi-1910. Type in Indian Museum, a unique ♀, the abdomen now missing.

orientalis, *id.*, *loc. cit.*, 453, ♂ ♀ (1912).

Darjiling, 5-viii-09, type ♂ [Paiva]; type ♀ [Annandale];

- Kurseong, 4-9-ix-09: [*Annandale*]; Calcutta [*Annandale*]; Shanghai, 8-v-06 [*Brunetti*]. *Types* in Indian Museum.
- subtinctoria*, *id.*, *loc. cit.*, 455, ♂ ♀ (1912).
Darjiling, 5-viii-09 [*Paiva*].
Types, a unique pair, in Indian Museum.
- flava*, *id.*, *loc. cit.*, 455, ♂ ♀ (1912).
halterata, *id.*, *loc. cit.*, 457 ♂ ♀ (1912).
Darjiling [*Brunetti* and *Paiva*]; 6000—7000 ft., 11 and 13-vi-14 [*Gravely*]; 20-ix-08 at light, *type* ♂, *halterata* [*Brunetti*]; 6-11-viii-09, *type* ♀, *halterata* [*Paiva*]; Madhupur, Bengal, 17-x-09 at light, *type* ♂, *flava* [*Paiva*]; Calcutta, 12-ix-07, *type* ♀, *flava* [*Annandale*]; near Khulna, Ganges Delta, on board launch, at light, 28-viii-09 [*Jenkins*]; Medha, Yenna Valley, Satara Distr., 2200 ft., 17-23-iv-12 [*Gravely*]; Shillong, 5500—6400 ft., 29-viii-5-ix-15 [*Kemp*].
Types of *flava* ♂ ♀ and *halterata* ♂ ♀ in Indian Museum.
- grandior*, *id.*, *loc. cit.*, 456, ♀, pl. viii, 18, wing (1912).
Simla, 10-v-09 [*Annandale*]. *Type* in Indian Museum, unique.
- genitalis*, *id.*, *loc. cit.*, 456, ♂ ♀ (1912),
Bhowali, Kumaon Div., 27-vi-10, *type* ♂, at light; 12-vi-10, *type* ♀. *Types* in Indian Museum.
- fusca* de Meij., Tijds. v. Ent. LVI, 351, pl. xvii, 17, wing (1913).
Nongkojadjar, Java [*Jacobson*]. *Type*, unique, in Amsterdam Museum.
- nigripalpis*, *id.*, *loc. cit.*, 351, ♂ (♀ lapsus), pl. xvii, 18, ♂ genitalia (1913).
Nongkojadjar, Java, i [*Jacobson*].
Type, unique, in Amsterdam Museum.
- insignis* Edw., Ann. Mag. Nat. Hist. (8) XVIII, 251, ♂ ♀, (*Erioptera*, sub-gen.) (1916).
Arisan, Formosa, 8000 ft., 10-x-12 [*Nitobe*]; Tokyo.
Types in British Museum.
- albogutatta* Edw., *loc. cit.*, p. 252, ♂, pl. xii, 4, genitalia (*Erioptera*, sub-gen.) (1916).
Arisan, Formosa, 8000 ft., 10-x-12 [*Nitobe*].
Unique *type* in British Museum.
- subfusca* Edw., Jour. Fed. Malay Sts. Mus., VIII, 19, ♂, pl. iii, 7, genitalia (1919).
Korinchi Peak, Sumatra. *Type* in British Museum.

Acyphona, Ost. Sack.

Monog. Dipt. N. Amer. IV, 151 (1869).
GENOTYPE, *Erioptera venusta*, Ost. Sack., by present designation, no species having apparently been nominated.

Trichosticha, Sch., Wein. Ent. Monats. VII, 221
(1863).¹

Ilisia, Rond., Dipt. Ital. Prod. I, 182 (1856).²

fenestrata de Meij., Tijds. v. Ent. LVI, 352, ♀, pl. xvii, 19, wing
(1913).

Nongkodjadjar, Java, i [Jacobson]; Tjibodas, Mt. Gedé,
Java, 8000 ft. [Bryant and Palmer].

Type in Amsterdam Museum.

Apparently no genotype of *Acyphona* has hitherto been selected.

Mesocyphona, Ost. Sack.

Monog. Dipt. N. Amer. IV, 152 (1869).

Ost. Sack., Berl. Ent. Zeits. XXXI, 194:

Brun., Fauna Brit. Ind., 458.

GENOTYPE, *Erioptera caloptera* Say., by my own
designation, Fauna, 458.

nigripes Brun., Fauna Brit. Ind., 458, ♂, pl. ix, 1, wing (1912).

Kurseong, 7—9-ix-09 [Annandale]; Ghoom, 7500 ft., 19-ix-08;

Pashok, 2000 ft. [Gravely]; Kalimpong, 600—4500 ft., 24-iv—
10-v-15 [Gravely]. Type in Indian Museum.

gracilis Brun., Rec. Ind. Mus. XV, 310, ♀ (1918).

Bhoirakund, Assam-Bhutan Frontier, Darrang Distr., 18—
22-x-12 [Kemp]. Unique type in Indian Museum.

Empeda, Ost. Sack.

Monog. Dipt. N. Amer. IV, 183 (1869).

Ost. Sack., Berl. Ent. Zeits. XXXI, 203:

Brun., Fauna Brit. Ind., 475.

GENOTYPE, *E. stigmatica* Ost. Sack., by original
designation.

Ilisophila, Rond., Dipt. Ital. Prod. I, 180 (1856).

GENOTYPE, *Erioptera lutea* Mg., by original
designation.

gracilis de Meij., Tijds. v. Ent. LIV, 49 ♀; pl. iv, 37, wing
(1911).

Id., loc. cit., LVI, 353, pl. xvii, 20, ♂ genitalia.

Tosari, Java [Kobus]; Nungkodjadjar, Java, i [Jacobson].

Type in Amsterdam Museum.

brevior Brun., Fauna Brit. Ind., 452, ♂ ♀, pl. ix, 2, wing (*Eriop-*
ptera) (1912).

Empeda inconspicua, *id.*, loc. cit., 475, ♂ ♀; pl. ix, 10, wing.

^{1, 2} No genotype appears to have been erected for *Trichosticha*, and of the three species placed in it by Schiner at its creation only one, *maculata* Mg., is now referred to it, and this one was selected by Rondani as type of *Ilisia*.

Darjiling, 5 and 11-viii-09, type ♂ and ♀ [Paiva]; 25-29-v-10 [Brunetti]; 3-9-vi-09 [Howlett]; Kurseong, 22-26-vi-10; 7-ix-09 [Annandale]. Types of both sexes of *breviar* and *inconspicua* in Indian Museum.

antica Brun., Fauna Brit. Ind., 568, ♂ (*Gonomyia*) (1912).
Ghumti, 4000 ft. Type in Indian Museum.

femoralis Edw., Jour. Fed. Malay Sts. Mus., viii, 20, ♀ (1919).
Korinchi Peak, Sumatra. Type in British Museum.

Gonomyia, Mg.

Syst. Besch I, 146 (1818).

Sch., F. Austr. II, 542: Ost. Sack., Mong. Dipt. N. Amer. IV, 177-179: *id.*, Berl. Ent. Zeits. XXXI, 200 (incl. correction of Monog. ref.): Brun., Fauna Brit. Ind., 468: *id.*, Rec. Ind. Mus. XV, 311, notes.

GENOTYPE, *Limnobia tenella* Mg. (t. Coquillett, 1910).

Taphrosia, Rond., Dipt. Ital. Prod. I, 182 (1856).
GENOTYPE, *Limnobia tenella* Mg., by original designation.

Gonomyia, Ost. Sack., Monog. Dipt. N. Amer. IV, 177 (1869).¹

metatarsata de Meij., Tijd. v. Ent. LIV, 48, ♀, pl. iii, 35, wing (1911).

Semarang, i, vii [Jacobson]; Pasaruan, Java [Kobus].
Type in Amsterdam Museum.

nubeculosa, *id.*, *loc. cit.*, 48, ♂, pl. iii, 36, wing (1911).

Wonosobo, Java, iv [Jacobson].

Type, a unique ♂, in Amsterdam Museum.

incompleta Brun., Fauna Brit. Ind., 471, ♂ ♀, pl. ix-5, wing; pl. xi-19, ♂ genitalia (1912).

Kurseong, 3-vii-08 [Annandale]; Sukna, 1-vii-08 [Annandale]; Ghumti, Darjiling, 4000 ft., vii-11 [Gravely]; Calcutta, 26-vii-08, type ♂, 21-viii-09 type ♀ [both Annandale]; 16-viii-12 in house [Gravely]; Katihar, viii-ix-10 [Paiva]; Bhogaon, 9-x-08 [Paiva], Kushtea, Bengal, 7-8-x-09 [Jenkins]; Talewadi, N. Kanara Dist. [Kemp]. Types in Indian Museum, *cotypes* in my collection.

flavomarginata, *id.*, *loc. cit.*, 472, ♂ ♀, pl. ix, 6 (1912).

Simla, 7000 ft. [Annandale]; Kurseong, 20-vi-10, type ♂; 18 and 22-vi-10; 6-ix-09, type ♀; 20-vi-10 [all Annandale]; Darjiling, 7-viii-09 [Jenkins]; 7000 ft. [Paiva]; Almora, Kumaon Div., 5500 ft., 13-16-ix-11 [Paiva].
Types in Indian Museum.

¹ *Gonomyia* is merely a different spelling of the name *Gonomyia*.

affinis, *id.*, *loc. cit.*, 472, ♂ ♀, pl. ix, 7, wing (1912).

Kumaon, 4700 ft. and 5000 ft. [*Annandale*]; Kurseong, 13—16-vii-07, *type* ♂; 25—28-iii-10; 5—7-vii-08; 22-vi-10 [all *Annandale*]; Darjiling, 5-viii-09 *type* ♀; 9-viii-09 [*Jenkins*]; Castle Rock and Talewadi, N. Kanara Dist. [*Kemp*].

Types in Indian Museum.

aperta, *id.*, *loc. cit.*, 473, ♂, pl. ix, 8, wing (1912).

Katihar, Bengal, 29—31-xii-09 [*Paiva*].

Type in Indian Museum, a unique specimen.

proxima, *id.*, *loc. cit.*, 474, ♂ ♀, pl. ix, 9, wing (1912).

Tirvani, 27-xii-09; Noalpur, 15-ii-08 (both base of Nepalese Himalayas); Rajmahal, Bengal, 14-ii-10, *type* ♂ ♀ *in cop.* [*Chaudhuri*]; Rajshahi, E. Bengal, 1—6-ii-07 [*Annandale*]; Pusa river bank, 6-i-11 [*Fletcher*]; 19-iv-09.

Types in Indian Museum, *cotype* ♀ in Pusa coll.

bryanti Alex., Proc. U.S. Nat. Mus. XLIX, 173, ♂, pl. xlv, 20, wing; pl. xlvii, 48, 49, genitalia (1916).

Tjibodas, Mt. Gedé, Java, 8000 ft. [*Bryant and Palmer*].

Type, Cat. No. 19,021, U.S. Nat. Mus.

nebulosa Edw., Ann. Mag. Nat. Hist. (8) XVIII, 252 (*Lipophleps*) (1916).

Arisan, Formosa, 8000 ft., 10-x-12 [*Nitobe*].

Type in British Museum.

Mongoma, Westw.

Trans. Ent. Soc. Lond. 364, pl. xvii, 1 (1881).

Ost. Sack., Berl. Ent. Zeits. XXVI, 90; XXXI, 203; Brun., Rec. Ind. Mus. VI, 290, 292 (*Mongoma* group): *id.*, Fauna Brit. Ind., 476: *id.*, Rec. Ind. Mus. XV, 311, notes, tab. allied gen. and Ind. spp.

GENOTYPE, *M. fragillima* Westw., by original designation.

tenera Ost. Sack., Berl. Ent. Zeits. XXVI, 89, ♂ (1882).

Brun., Rec. Ind. Mus. VI, 295: *id.*, Fauna Brit. Ind., 480, ♂ ♀.

pallidiventris, Brun., Fauna Brit. Ind., 481, ♀ (1912).

Kurseong, 4700—5000 ft. [*Annandale*]; Ghumti, 4000 ft. [*Gravely*]; Sukna, 1-vii-08; 1-viii-08; Nedumangad, Travancore, 14-xi-09; Pallode, 15-xi-08, *type* of *pallidiventris* [both *Annandale*]; Tura, Garo Hills, vii, viii-17 [*Kemp*]; Ginting Bidai, Selangor-Pahang Frontier, Federated Malay States, 2000 ft., iv-17 [*Boden Kloss*]; Philippines.

pennipes Ost. Sack., Berl. Ent. Zeits. XXXI, 204, ♂ (1887).

de Meij., Tijd. v. Ent. LIV, 50, pl. iv, 38—41, var. chars.:

Brun., Rec. Ind. Mus. VI, 295: *id.*, Fauna Brit. Ind., 479, ♂ ♀, pl. ix, 14, wing; pl. xi, 13, ♂ genitalia.

Sukna, N. Bengal, 1-vii-08; Calcutta, not uncommon, Feb. to Sept.; Pusa, at light, 6-vii-13 [Howlett]; in plantain bark, 27-xi-12; Kavalai, Cochin, 1300—3000 ft., 24—27-ix-14 [Gravelly]; Peradeniya, 12—16-vii-10 [Green]; 16-ix-09 at light; Insein, Lower Burma, 31-vii—4-viii-14 [Fletcher]; Semarang, i, x [Jacobson]; Tumbong Hiong, Borneo.

cariniceps Ender., Zool. Jahr. XXXII, 60, ♂, fig. 1¹, wing (1912).

Soekaranda, Sumatra, Mt. Salak, Java, 15-v-09, ♂ ♀ [Bryant and Palmer]; Siolak Daras, Korinchi Peak, Sumatra, 3000 ft., iii-14. Type in Stettin Zoological Museum.

obscura de Meij., Bijd. tot de Dierk. XIX, 48, ♂, pl. i, 5, wing (1913).

Waigon, 1-i-10. Unique type in Amsterdam Museum.

albipennis de Meij., Tijd. v. Ent. LVI, 353, ♀ (1913).

Nungkodjadjar, Java, i [Jacobson].

Unique type in Amsterdam Museum.

splendida Brun., Rec. Ind. Mus. XV, 313, ♂ ♀ (1918).

Above Tura, Garo Hills, 3500—3900 ft., vii, viii-17 [Kemp].

Types in Indian Museum.

kempi, *id.*, *loc. cit.*, 313, ♀ (1918).

Above Tura, Garo Hills, 3500—3900 ft., viii-17 [Kemp].

Type in Indian Museum.

flava, *id.*, *loc. cit.*, 314, ♀ (1918).

A unique ♀ taken in company with the above species.

Type in Indian Museum.

Plesiomongoma, Brun.

Rec. Ind. Mus. XV, 314 (1918).

GENOTYPE, *P. venosa*, *id.*, sp. nov., *loc. cit.*

venosa Brun., *loc. cit.*, 314, ♀ (1918).

Shillong, 5500—6400 ft., 29-viii—5-ix-15 [Kemp].

Unique type in Indian Museum.

Trentepohlia, Big.

Ann. Soc. Ent. Fran. (3) II, 473 (1854).

Brun., Rec. Ind. Mus. XV, 315: Edw., Ann.

Mag. Nat. Hist. (8) VIII, 63 (1911).

Mongomioides, Brun., Rec. Ind. Mus. VI, 296 (1911).

Id., Fauna Brit. Ind., 481.

GENOTYPE, *Limnobia trentepohlii* Wied., by original designation; the type species of *Mongomioides* is the same.¹

¹ Though Bigot's description is quite valueless, Edwards recently noted (*vide ante*) one of the true generic characters (three posterior cells only), but I had overlooked his paper. Still, the nomination of a type species does not constitute the diagnosis of a genus.

trentepohlii Wied., Auss. Zweifl. I, 551, ♂, pl. vi, b, fig. 12 (*Limnobia*) (1828).

Macq., Hist. Nat. Dipt. I, 100 (*Limnophila*): Ost. Sack., Berl. Ent. Zeits. XXVI, 90; de Meij., Tijd. v. Ent. LIV, 51, notes, pl. iv, 42, wing; Brun., Rec. Ind. Mus. VI, 297: *id.*, Fauna Brit. Ind., 482, ♂ ♀, pl. ix, 13, wing.

Widely distributed throughout India, especially in the plains, Ceylon, Burma, Assam, Siam, Java and Sumatra. It extends northward at least to Sylhet [*Hall*]. The recorded months of appearance are February, and from August to November.

Type in Dr. Trentepohl's collection.

marmorata Brun., Fauna Brit. Ind. 483, ♀, pl. ix, 11, wing (*Mongomioides*) (1912).

Calcutta, 28-xi-07. Unique *type* in Indian Museum.

nigroapicalis, *id.*, *loc. cit.*, 483, ♂, pl. ix, 12, wing (*Mongomioides*) (1912).

Lucknow, 5-iii-11, *type*; Pusa, 20-iv-13, on pipal [*Howlett*]; 1-iv-16 [*Fletcher*]; Rajmahal, Bihar [*Chaudhuri*]; Sorabkalti, Ganges Delta [*Jenkins*]; Peradeniya, v-09; Tura, Garo Hills, Assam, vii-17 [*Kemp*]. *Type* in Indian Museum.

albogeniculata, *id.*, *loc. cit.*, App. 569, ♂ (*Mongomioides*) (1912).

Brun., Rec. Ind. Mus. VII, 448 (1912).¹

Dawna Hills ("3rd Camp to Misty Hollow"), 400—2400 ft., 22—30-xi-11 [*Gravelly*]. *Types* in Indian Museum.

speiseri Edw., Ann. Mag. Nat. Hist. (8) XII, 204, ♂ ♀, *nom. nov.*, fig. (1913).

Mongoma exornata Speiser (*nec* Bergr.), Berl. Ent. Zeits. LII, 135 (1907).

Peradeniya, II, 1912. Described from W. Coast of Africa.

Type in British Museum.

ornatipennis Brun., Rec. Ind. Mus. XV, 315, ♂, fig. 4, wing (*ornatepennis*, *lapsus*) (1918).

Castle Rock. N. Kanara Dist., 11—26-x-16 [*Kemp*].

Unique *type* in Indian Museum.

saucia Alex., Proc. U.S. Nat. Mus., XLIX, 174, ♀, pl. xlv, 21, wing (1916).

Tjibodas, Mt. Gedé, Java, 7800 ft. [*Bryant and Palmer*].

Type, Cat. No. 19,022, U.S. Nat. Museum.

pictipennis Bezzi, Phil. Jour. Sci. XII, Sect. D. 115, ♂ (1917).

Luzon [*Baker*].

Paramongoma, Brun.

Rec. Ind. Mus. VI, 295 (1911).

Brun., Fauna Brit. Ind., 484: *id.*, Rec. Ind.

Mus. XV, 314.

¹ The Fauna reference antedates the other by about a month.

GENOTYPE, *Cylindrotoma albitarsis* Dol., by original designation.

Mongomella, Ender., Zool. Jahr. XXXII, 61 (1912).

GENOTYPE, *Mongoma pallida* Will., by original designation.

albitarsis Dol., Nat. Tijds. Ned. Ind. XIV, 391, pl. iv, 1, full ins. (*Cylindrotoma*) (1857).

Ost. Sack., Berl. Ent. Zeits. XXVI, 90: Brun., Rec. Ind. Mus. VI, 295, notes: *id.*, Fauna Brit. Ind., 484.

Java, Amboina. Location of *type* unknown, probably lost.

Anchimongoma, Brun.

Rec. Ind. Mus. XV, 316 (1918).

GENOTYPE, *A. simplex*, sp. nov., *loc. cit.*

simplex Brun., *loc. cit.*, 316, ♂ ♀, fig. 5, wing (1918).

Castle Rock, N. Kanara Dist., 11-26-x-16 [Kemp].

The unique *types* in the Indian Museum.

Lechria, Skuse.

Proc. Linn. Soc. N.S. Wales (2) IV, 830 (1890).

Brun., Fauna Brit. Ind., 465.

GENOTYPE, *L. singularis* Skuse, by original designation.

lucida de Meij., Tijds. v. Ent. LIV, 53, ♀, pl. iv, 45, wing (1911).
Batavia, vi [Jacobson].

Type, unique, in Amsterdam Museum.

bengalensis Brun., Rec. Ind. Mus. VI, 301, ♂ ♀ (1911).

Id., Fauna Brit. Ind., 467, ♂ ♀, pl. ix, 4, wing: *id.*, Rec. Ind. Mus. XV, 317.

Pusa, 15-viii-08, *type* ♂; 16-vii-10; 24-viii-11 [Howlett]; Calcutta, 19-viii-07, *type* ♀; 18-vi-09; Bangalore, 12-x-10 [Annandale]; Talewadi, N. Kanara Dist. [Kemp]; 30-v-12 [Gravely].
Type ♂ in Pusa collection, *type* ♀ in Indian Museum.

nepalensis Brun., Rec. Ind. Mus. XV, 317, ♀ (1918).

Katmandu, Nepal. The unique *type* in the Indian Museum.

leucopeza de Meij., Tijds. v. Ent. LVI, Supp., p. 3, ♀ (1914).

Semarang, viii [Jacobson].

Unique *type* in Amsterdam Museum.

Styryngomyia, Lw.¹

Dipt. Beitr., I in "Zu der öffentlichen Prüfung der Schuler d. Königl. Fried. Wilh. Gymn. zu Posen," p. 6 (1845).

¹ Placed in error in Rhabdini in my second paper on Tipulidae. It belongs near *Lechria*.

Edw., Trans. Ent. Soc. Lond., 208 (1914)¹:
Brun., Rec. Ind. Mus. VI, 297; XV, 304:
Fauna Brit. Ind., 459.

GENOTYPE, *S. venusta* Lw., a fossil species, by original designation.

Pycnocrepis, Ender., Zool. Jahr. XXXII, 65 (1912).

GENOTYPE, *Styringomyia crassica* Speis., as *P. annulipes*, *id.*, sp. nov., *loc. cit.*

crassica Speis., Berl. Ent. Zeits. LII, 132 (*Idiophloeba*) (1907).
ceylonica Edwards, Ann. Mag. Nat. Hist. (8) VIII, 62, ♀ (July 1911): Brun., Rec. Ind. Mus. VI, 298, ♂ ♀ (erroneous redescription from series containing more than one species): Brun., Fauna Brit. Ind., 461 (same redescription copied).

didyma Grims., de Meij., Tijd. v. Ent. LIV, 40 (pt).

Pycnocrepis annulipes Ender., Zool. Jahr. XXXII, 65, ♂, fig. O¹, wing (1912).

Tarnab, Peshawar Distr. [*Fletcher*]; Bhogaon, N. Bengal [*Paiva*]; Puri, Orissa, 10-xi-10 [*Annandale*]; Kankondigee, Sunderbunds, Ganges Delta, 14-xi-09, at light on board launch [*Jenkins*]; Calcutta; Weligama, Ceylon, 9-ii-08 [*Fletcher*]; Taiping [*Wray Junr.*]; Pasuran, Java [*Kobus*]; Tainan, Formosa [*Sauter*]; Formosa ♂ ♀ (t. Ender.); Singgora, Siam [*Annandale*].
Type in British Museum.

The above localities are mainly from Mr. Edwards' corrected list in the Trans. Ent. Soc.

obscura Brun., Rec. Ind. Mus. VI, 300 ♀ (erroneously given as ♂ (1911)).

Id., Fauna Brit. Ind., 463, ♂ (as ♀).

Thaumaspur, Nepal, 18—20-ii-08.

Type in Indian Museum.

flava Brun., Rec. Ind. Mus. VI, 301, ♂ (1911).

Id., Fauna Brit. Ind., 464, ♂.

Tenmalai, Travancore, 22-xi-08, a unique ♂, "at light" [*Annandale*]. *Type* in Indian Museum.

nigrofemorata Edw., Trans. Ent. Soc., p. 215, ♀, pl. xxiii, 51 (1914).

Taiping, Malay States [*Wray Junr.*].

Unique *type* in British Museum.

formosana, *id.*, *loc. cit.*, 219, ♂ ♀, pl. xix, 9; xxi, 30, 31; xxv, 70, 71, 72 (1914).

Tainan, Formosa; Koshun, Formosa [*Sauter*].

Type in the Deuts. Ent. Museum, Berlin; *paratypes* in British Museum.

javana, *id.*, *loc. cit.*, 220, ♂, pl. xix, 8; xxii, 32, 33 (1914).

Nongkodjadjar, i, Java [*Jacobson*].

Type in Amsterdam Museum, a unique specimen.

¹ This paper by Edwards reviews all the known species, and includes a revised generic description and table. The figures, of which there are many, are mainly of genitalia. He corrects one or two errors of mine in the genus.

- jacobsoni**, *id.*, *loc. cit.*, 220, ♂ ♀; pl. xix, 6, thorax, 7; pl. xxii, 34, 35; xxv, 73 (1914).
didyma Grims., de Meij., Tijds. v. Ent. LIV, 40 (pt.).
 Semarang, i; Batavia, ii [*Jacobson*]; Sukabumi (Pare), xii [*Dammermann*]. Type in Amsterdam Museum.
- fryeri**, *id.*, *loc. cit.*, 221, ♂ ♀, pl. xxii, 36, 37; xxv, 74, 75 (1914).
 Peradeniya [*Fryer*]. Type in British Museum.
- himalayana**, *id.*, *loc. cit.*, 221, ♂ ♀, pl. xxii, 40; xxiii, 41; xxv, 77 (1914).
 Sukna, ♂ ♀ *in cop.* [*Annandale*]. Type in Indian Museum.
- nepalensis**, *id.*, *loc. cit.*, 222, ♂ ♀, pl. xxiii, 42, 43; xxv, 78, 79 (1914).
 Sukhwani, Nepal. Type in Indian Museum.

Gnophomyia, Ost. Sack

- Proc. Acad. Nat. Sci. Philad. 223 (1859).
 Ost. Sack., Monog. Dipt. N. Amer. IV, 172;
 pl. ii, 5, wing; iv, 19, 19a, genitals, ♂ ♀; Sch.,
 F. Austr. II, 535; Ost. Sack., Berl. Ent. Zeits.
 XXXI, 198; Brun., Fauna Brit. Ind., 487.
Id., Rec. Ind. Mus. XV, 318, notes on synonymy.
 GENOTYPE, *G. tristissima* Ost. Sack., by Coquillett's designation.
- Furina*, Jaen., Abh. Senck. Ges. VI, 318 (1867).
 GENOTYPE, *Limnobia rufithorax* W., by original designation.
- orientalis** de Meij., Tijds. v. Ent. LIV, 46, ♂ ♀; pl. iii, 32, wing, 33, ♂ genitalia (1911).
 Batavia, vi; Bekasssi, Java, viii [*Jacobson*]; Arisan, Formosa, 8000 ft., 10-x-12 [*Nitobe*]. Types in Amsterdam Museum.
- strenua** Brun., Fauna Brit. Ind., 492, ♀, pl. ix, 19, wing (1912).
 Kurseong, 5-ix-09 [*Annandale*].
 Type in Indian Museum, a unique specimen.
- nigra**, *id.*, *loc. cit.*, 493, ♂, pl. x, 2, wing (1912).
 Kumaon [Forest Zoology colln.]; Bhim Tal, 4500 ft. [*Annandale*]; Kurseong, 7-ix-09 [*Annandale*]; Nilgiri Hills, 3500 ft. [*Andrewes*]. Type in Indian Museum.

Dasymallomyia, Brun.¹

- Rec. Ind. Mus. VI, 304 (1911).
 Brun., Rec. Ind. Mus. XV, 318.
 GENOTYPE, *D. signata* Brun., sp. nov.

¹ The synonymy of *Dasymallomyia* with *Gnophomyia* was pointed out by Alexander. I did not know of the group of tropical American species with short hairy legs to which he refers. The genus *Gnophomyia* was imperfectly understood by me in the preparation of my work. However, Mr. Edwards is inclined to retain *Dasymallomyia* for one of the groups of species comprised in *Gnophomyia*, and it is here regarded provisionally as a valid genus.

- signata** Brun., *loc. cit.*, 304, ♀ (1911).
Id., Fauna Brit. Ind., 495, ♀, pl. x, 4.
 Kurseong, 7-ix-09 [Lynch]; Horisha, Formosa, 10-v-13 [Maki]. *Type* in Indian Museum.
- maculipleura** Edw., Ann. Mag. Nat. Hist. (8) XVII, 360, ♂ ♀, fig. 5, p. 356, genitalia (1916).
 Bukit Besar, Siam [Robinson and Annandale].
- fraterna**, *id.*, *loc. cit.*, 361, ♂, fig. 6, p. 356, genitalia (1916).
 Selangor [Butler].
- nigrescens**, *id.*, *loc. cit.*, 361, ♀ (1916).
 Talam, Perak, 3500 ft., 17-i-02 [Robinson and Annandale].
Types of last three species in British Museum.
- similis** Edw., Ann. Mag. Nat. Hist. (8) XVIII, 251, ♀, pl. xii, 3 genitalia (1916).
 Arisan, Formosa, 8000 ft., 10-x-12 [Nitobe].
Type in British Museum.

Oxydiscus, de Meij.

- Tijd. v. Ent. LVI, 350 (1913).
 GENOTYPE, *O. nebulosus* de Meij., the original species.
- nebulosus** de Meij., *loc. cit.*, 351, ♀ (1913).
 Tjinjiruan, West Java [v.d. Weele].
 Unique *type* in Amsterdam Museum.
- umbrosus** Edw., Ann. Mag. Nat. Hist. (8) XVII, 361, ♀ (1919).
 Kedah Peak, 3200 ft., Malay Peninsula [Stanton].
 Unique *type* in British Museum.

Clydonodozos, Ender.¹

- Zool. Jahr. XXXII, 57 (1912).
 Brun., Rec. Ind. Mus. XV, 318.
 GENOTYPE, *C. multistriatus*, *id.*, sp. nov., *loc. cit.*
- multistriatus** Ender., *l. c.*, 57, ♂ ♀, fig. J¹, wing (1912).
 Bekantiang, Sumatra.
Types in Stettin Zoological Museum.
- punctulatus**, *id.*, *loc. cit.*, 59 (sex?) fig. K¹, wing.
 Soekaranda, Sumatra.
 The unique *type* in Stettin Zoological Museum.
- griseiceps** de Meij., Tijd. v. Ent. LVIII, Supp. II, ♀ (1910).
 Sinabang, Simalur Is. off W. coast Sumatra, ii.
Type in Amsterdam Museum, a unique ♀.

¹ In Mr. Edwards' opinion this genus is doubtfully distinct from *Conosia*.

Conosia, Wulp.

Tijd. v. Ent. XXIII, 159, pl. x, 5—7 (1880).

Ost. Sack., Berl. Ent. Zeits. XXXI, 206:

Brun., Fauna Brit. Ind., 496.

GENOTYPE, *Limnobia irrorata* W., by original designation.

irrorata Wied., Auss. Zweifl. I, 574, ♀ (*Limnobia*) (1828).

Auctt. var.; Brun., Rec. Ind. Mus. VI, 283: Fauna Brit. Ind., 497, ♂ ♀, fig. 43, lateral view, pl. x, 5, wing.

Limnobia substituta Walk., List Dipt. Brit. Mus. I, 39 (1848).

Limnophila crux Dol., Nat. Tijd. Ned. Ind. XIV, 388, pl. iv, 3, full insect (1857).

Apparently distributed generally throughout the Orient, common in many places, and extending from Peshawar District [Fletcher] to Assam, Cochin, Java, Formosa, Sumatra, China, Japan and Australia. Dr. Annandale took it at Lake Tiberias in Palestine.

Type presumably in Westermanns' coll. at Vienna Museum.

Cladura, Ost. Sack.

Proc. Acad. Nat. Sci. Philad. p. 229 (1859).

Ost. Sack., Monog. Dipt. N. Amer. IV, 187:

Berl. Ent. Zeits. XXXI, 205: Brun., Rec. Ind.

Mus. VI, 283 (the *Cladura* group); XV, 319:

Fauna Brit. Ind., 499.

GENOTYPE, *C. flavoferruginea* Ost. Sack., by original designation.

flavescens Brun., Rec. Ind. Mus. VI, 284, ♂ ♀ (1911).¹

Id., Fauna Brit. Ind., 501, ♂ ♀, pl. x, 6, wing: Rec. Ind.

Mus. XV, 319 (retention of species in *Cladura*).

Darjiling, 7-viii-09, *type* ♂; 6-viii-09, *type* ♀ [both *Paiva*].

Types, a unique pair, in Indian Museum.

interrupta Brun., Rec. Ind. Mus., XV, 319, ♀ (1918).

Pashok, Darjiling Distr., 3500 ft., 26 v—14-vi-16 [*Gravelly*].

Type in Indian Museum.

Paracladura, Brun.²

Rec. Ind. Mus. VI, 286 (1911).

Id., Fauna Brit. Ind., 502: *id.*, Rec. Ind. Mus.

XV, 320 (position criticised).

GENOTYPE, *P. gracilis* Brun., by original designation.

¹ *C. flavescens* is almost certainly a *Cladura* in Osten Sacken's sense, agreeing with his description and with Needham's figure. I think Alexander is in error in claiming that my species is wrongly placed generically.

² For discussion on Alexander's objection to this genus, v. Brun., Rec. Ind. Mus. XV, 320.

gracilis Brun., Rec. Ind. Mus. VI, 287 ♂ ♀ (1911).

Id., Fauna Brit. Ind., 503, ♂ ♀, pl. x, 9, wing, 10 antenna.

Darjiling, 28—29-v-10 [Brunetti].

Types, a unique pair, in Indian Museum.

elegans Brun., Rec. Ind. Mus. VI, 288, ♀ (1911).

Id., Fauna Brit. Ind., 504, ♀.

Darjiling, 26—29-v-10 [Brunetti]. *Type* in Indian Museum.

Symplecta, Mg.

Syst. Besch. VI, 282 (1830).

Sch., F. Austr. II, 545: Ost. Sack., Monog.

Dipt. N. Amer. IV, 170, pl. i, 20, wing; iv, 21,

forceps: *id.*, Berl. Ent. Zeits. XXXI, 197:

Brun., Fauna Brit. Ind., 485.

GENOTYPE, *Limnobia punctipennis* Mg., as *hybrida*, by Westwood's designation.

Helobia St. Farg. et Serv., Encyc. Meth. Ins. X, 585 (1825).

GENOTYPE, *Limnobia hybrida* Mg. as *L. punctipennis* Mg., by original designation.

Idioneura Phil., Verh. zool. bot. Gesch. Wien. XV, 615, pl. xxiii, 4 (1865).

GENOTYPE, *I. macroptera*, *id.*, *loc. cit.*, sp. nov.

Symplectromorpha Mik., Wien. Ent. Zeits. V, 318 (1886).

GENOTYPE, *Limnobia stictica* Mg., by original designation.

punctipennis Mg., Syst. Besch. I, 147, pl. v, figs. 2, 3, 7 (*Limnobia*) (1818).

Aucttr var.: Brun., Fauna Brit. Ind., 486, ♂ ♀, pl. ix, 15, wing.

Peshawar, 19-iii-13 [Howlett]; Darjiling, 22—30-ix-08 [Brunetti]; 5—6-viii-09 [Paiva]; Matiana, Simla Distr., 8000 ft., 28—30-iv-07 [Annandale]. Also Europe.

Type in the Meigen collection at Paris Museum, but practically nothing remains of it but the wings.

Section AMALOPINI.

Ost. Sack., Monog. Dipt. N. Amer. IV, 259:

Bruñ., Rec. Ind. Mus. VI, 305: *id.*, Fauna Brit. Ind., 507.

Amalopina, Ost. Sack., Monog. Dipt. N. Amer. IV, 259.

Pediciinae, Kert., Kat. Dipt. II.

Trichocera, Mg.¹

Illig. Mag. II, 262 (1803).

Sch., F. Austr. II, 546; Ost. Sack., Mong. Dipt. N. Amer. IV, 233, pl. ii, 13, wing: *id.*, Berl. Ent. Zeits. XXXI, 217; Brun., Rec. Ind. Mus. VI, 305 (genus removed to Section Amalopini: *id.*, Fauna Brit. Ind., 508.

GENOTYPE, *Tipula hiemalis* Degeer, by original designation.

ocellata Walk., Ins. Saund. Dipt., pt. v, 433, ♀ (1856).

Brun., Fauna Brit. Ind., 510, ♀.

East India; Theog, Simla Hills, 2-v-07 [*Annandale*]. *Type* in British Museum, in inferior condition. My redescription is from a single ♀ from Theog, on the assumption that the species is correctly identified.

punctipennis Brun., Fauna Brit. Ind., 511, ♂ ♀, pl. x, 13, wing (1912).

Simla, 23—25-iv-07 [*Annandale*].

Types in Indian Museum.

flava, *id.*, *loc. cit.*, 512, ♀ (1912).

Darjiling, 28-v-10 [*Brunelli*].

Type, a unique ♀, in Indian Museum.

montana, *id.*, *loc. cit.*, 513, ♂ (1912).

Badrinath, Garhwal Distr., 10,200 ft., W. Himalayas, 27-v-10, ♂ [*Imms*]. *Type*, a unique ♂, in Indian Museum

Amalopsis, Hal. in Walk.

Ins. Brit. Dipt. III, p. xv (1856).

Sch., F. Austr., II, 527; Ost. Sack., Monog. Dipt. N. Amer. IV, 260, pl. ii, 15, wing; iv, 30 genitals: *id.*, Berl. Ent. Zeits. XXXI, 224; Brun., Fauna Brit. Ind. 513: *id.*, Rec. Ind. Mus. XV, 321; Bergr., Ann. Mag. Nat. Hist. (8) XI, 583 (*Amalopsis* v. *Tricyphona*).

GENOTYPE, *Linnobia occulta* Mg. (*t.* Coquillett, 1910).

Tricyphona, Zett., Ins. Lapp. Dipt., 851 (1838).

GENOTYPE, *Linnobia immaculata* Mg., by original designation.

Bophrosia, Rond., Dipt. Ital. Prod. I, 183 and Errata (1856).

GENOTYPE, the same species as of *Tricyphona* by original designation.

¹ For reasons for removing the genus here, v. Brun., Rec. Ind. Mus. VI, 305.

Crunobia, Kol., Wien. Ent. Monats. IV, 391 (1860).

GENOTYPE, *C. schincri*, sp. nov., *loc. cit.*

? *Nasiterna*, Walleng., Entom. Tidskr. II, 179 (1881).

glabripennis Brun., Fauna Brit. Ind., 515, ♂, pl. x, 12, wing (1912).

Darjiling, 1-x-08, 2 and 10-vi-17 [*Brunetti*]; Yembung, N.E. Ind. Frontier, 1100 ft., 15-i-12 [*Kemp*].

Type in Indian Museum.

elegans Brun., *loc. cit.*, 516, ♂ ♀, pl. x, 11, wing (1912).

Kurseong, 28-vi-10, type ♂; 20-vi-10, type ♀; 18-vi-10. Types in Indian Museum.

spectralis Brun., Rec. Ind. Mus. XV, 321, ♂, pl. vii, 4, wing (1918).

Pashok, Darjiling District, 3000 ft., 26-v—14-vi-16 [*Gravelly*]. Type in Indian Museum.

Amalopina, Brun.

Fauna Brit. Ind. 517 (1912).

GENOTYPE, *A. elegantula*, *id.*, sp. nov., *loc. cit.*

elegantula Brun., *loc. cit.*, 517, ♂ (1912).

Darjiling, 26-v-10 [*Brunetti*].

Type, a unique ♂ in indifferent condition, in Indian Museum.

Section LIMNOPHILINI.¹

Trichocerinae, Kert., Kat. Dipt. II, 218 (1902).

Rhaphidolabis, Ost. Sack.

Proc. Ent. Soc. Philad. p. 225 (1865).

Ost. Sack., Monog. Dipt. N. Amer. IV, 284, pl. ii, 17, wing: Brun., Fauna Brit. Ind., 517: *id.*, Rec. Ind. Mus. XV, 322, placed unintentionally in Eriopterini; notes *re* antennae.

GENOTYPE, *R. tenuipes* Ost. Sack., by original designation.

Plectromyia, Ost. Sack., Monog. Dipt. N. Amer. IV, 282 (1869).

GENOTYPE, *P. modesta*, *id.*, sp. nov., *loc. cit.*

? *Claduroides*, Brun., Rec. Ind. Mus. VI, 288 (1911): Fauna Brit. Ind., 505.²

GENOTYPE, *C. fascipennis*, *id.*, sp. nov., *loc. cit.*

sordida Brun., Rec. Ind. Mus. VI, 290, ♂ ♀ (*Claduroides*) (1911).

Id., Fauna Brit. Ind., 506, ♂ ♀ (*Claduroides*).

¹ For characters, etc. v. Ost. Sack., Monog. Dipt. N. Amer. IV, 190 and Brun., Fauna Brit. Ind., 521.

² See Brunetti, Rec. Ind. Mus. XV, 320, 322, for validity of *Claduroides*.

- Simla, 10-v-09, type ♂, 12-v-09; Kurseong, 4-ix-09, type ♀; Phagu [all *Annandale*]. Types in Indian Museum.
- fascipennis** Brun., Rec. Ind. Mus. VI, 289, ♂ ♀ (*Claduroides*) (1911).
Id., Fauna Brit. Ind., 505, pl. x, 7, wing, 8, antenna (*Claduroides*) (1912).
Gnophomyia longipennis, Brun., Fauna Brit. Ind., 489, ♀, pl. ix, 17, wing (1912).
Rhaphidolabis fascipennis, *id.*, loc. cit., 519, ♀, pl. x, 14, wing (1912).
 Darjiling, 20-ix-08, type [Brunetti]; 9-viii-09 [Paiva]; Kurseong, 7-ix-09; Phagu, 12-v-09 [both *Annandale*]; Dhakuri, Kumaon Div., 8900 ft. [Imms]. Types of *R. fascipennis*, *G. longipennis* and *C. fascipennis* all in Indian Museum.
- incompleta** Brun., Fauna Brit. Ind., 493, ♂, pl. x, 2, wing (*Gnophomyia*) (1912).
 Kurseong, 7-ix-09 [*Annandale*].
 Unique type in Indian Museum.
 Alexander has noted that this species would fall in *Plectromyia*, Ost. Sack., but this is now generally sunk in *Rhaphidolabis*.
- indica** Brun., Fauna Brit. Ind., 519, ♂ ♀, pl. x, 15, wing (1912).
 Theog, 27-iv-07, type ♂; Matiana, 28—30-iv-07, type ♀; Phagu [all *Annandale*]. Types in Indian Museum.
- brunettii** Edw., Ann. Mag. Nat. Hist. (8) XVIII, 254, *nom. nov.* for *aperta* Brun., preocc. Coq. (1916).
aperta Brun., Fauna Brit. Ind., 492, ♂, pl. x, 1, wing (*Gnophomyia*) (1912).
 Darjiling, 8-viii-09 [Paiva]; Arisan, Formosa, 8000 ft., 10-x-12 [Nitobe]. Type in Indian Museum.

Limnophila, Macq.

- Hist. Nat. Dipt. I, 95 (1834).
 Sch., F. Aust. II, 554: Ost. Sack., Monog. Dipt. N. Amer. IV, 196—202, pl. ii, 6, 7, 9, 10, wings; iv, 23—27, ♂ genitalia: Brun., Fauna Brit. Ind., 521: Rec. Ind. Mus. VI, 306, notes; XV, 323, tab. orient spp.
 GENOTYPE, *Limnobia ferruginea* Mg., according to Coquillett (1910), who replaces the generic name by *Phylidorea*, Big.
Limnomya, Rond., Dipt. Hal. Prod. IV, Corrigenda, II (1861): *nom. nov.* for *Limnophila*.
Pilaria, Sintensis, Sitzs. Naturf. Ges. Dorpat VIII, 398 (1888).
 GENOTYPE, *Limnobia pilicornis* Zett., by Coquillett's designation.

- terminalis** Walk., Proc. Linn. Soc. Lond. V, 230, ♀ (*Limnobia*) (1861).
de Meij., Nova Guin. Res. IX, 309, pl. x, 4, wing: Brun., Rec. Ind. Mus. VI, 306.
Dorey, Papua; Bivak Is., Papua, xii; Regen Is, Papua ix, x [Lorentz]. *Type* in British Museum, useless for comparison.
- selectissima** Walk., Proc. Linn. Soc. Lond. VII, 202, ♂ (*Limnobia*) 1864).
Mysol. *Type* in British Museum.
- contingens**, *id.*, *loc. cit.*, VIII, 104, ♂ (*Limnobia*) (1865).
Papua. *Type* in British Museum, abdomen and legs missing.
- trisignata**, *id.*, *loc. cit.*, VIII, 105, ♀ (*Limnobia*) (1865).
Brun., Rec. Ind. Mus. VI, 306.
Papua. *Type* in British Museum in fair condition.
- opaca** de Meij., Tijds. v. Ent. LIV, 52, ♀, pl. iv, 44, wing (1911).
Wonosobo, Java, v [Jacobson].
Type, a unique ♀, in Amsterdam Museum.
- genitalis** Brun., Fauna Brit. Ind., 490, ♂ ♀, pl. ix, 16, wing (*Gnophomyia*) (1912).
Bhowali, Kumaon Div., 27-vi-10 [Imms].
Types in Indian Museum, unique.
- furcata**, *id.*, *loc. cit.*, 491, ♂ ♀, pl. ix, 18, wing (*Gnophomyia*) (1912).
Darjiling, 5—10-viii-09 [Paiva].
Types in Indian Museum.
- pallidicoxa** Brun., *loc. cit.*, 523, ♀ (1912).
Kurseong, 4-vii-08 [Annandale]; 26-iii-10 [Gravely]; Bhogaon, [Paiva]; Bhowali, Kumaon, 14-vi-10, at light [Imms]. Cherrapunji, Assam, 4400 ft., 2—8-x-14 [Kemp].
Type in Indian Museum.
- simplex**, *id.*, *loc. cit.*, 523, ♀ (1912).
Pattipola (hills), Ceylon [Annandale]; Bhowali, 29-vi-10 [Imms]. *Type* in Indian Museum.
- multipunctata**, *id.*, *loc. cit.*, App. 569 (1912).
Nilgiri Hills. The unique *type* in the Indian Museum.
- honesta**, *id.*, *loc. cit.*, App., 570, ♀ (1912).
Kumaon Division. The unique *type* in the Indian Museum.
- claripennis**, *id.*, Rec. Ind. Mus. VIII, 153, ♀ (1913).
Yembung, N.E. Indian Frontier, 1100 ft., 13-i-12 [Kemp].
Types in Indian Museum.
- quartarius**, *id.*, *loc. cit.*, 154, ♂ ♀ (1913).
Rötung, N. E. Indian Frontier, 1400 ft., 26—27-xii-11 [Kemp].
Types in Indian Museum, the ♀ in bad condition.

- apicalis* de Meij., Nova Guin. Res. IX, 309, ♀, pl. x, 3, wing (1913).
Alkmaar, S.W. Papua [Lorentz].
Type in Amsterdam Museum, unique.
- javana* de Meij., Tijd. v. Ent. LIX, 198, ♂ ♀, pl. vii, 11, wing (1916).
Gedeh, Java, 1500—2000 metres, June [Konigsberger].
Unique ♂ in Amsterdam Museum.
- amica* Alex., Proc. U.S. Nat. Mus. XLIX, 175, ♂, pl. xlv, 22, wing (1916).
Java, Tjibodas, Mt. Gedé, Java, 4500 [Bryant and Palmer].
Type, Cat. No. 19023, U.S. Nat. Mus.
- palmeri*, *id.*, *loc. cit.*, 175, ♂ (1916).
Java, taken in company with above species.
Type, Cat. No. 19024, U.S. Nat. Mus.
- demeijeri* Edw., Tr. Zool. Soc. XX, 393, ♀, pl. xxxviii, full ins. col. (1915).
Upper Utakwa Valley, Papua, 5000—10,000 ft., ii—iii-13. The unique type (presumably) in British Museum.
- annulipes* Brun., Rec. Ind. Mus. XV, 324, ♂ (1918).
Talewadi, N. Kanara Distr., 3—10-x-16 [Kemp].
Unique type in Indian Museum.
- parvicellula*, *id.*, *loc. cit.*, 325, ♂ (1918).
Taken in company with *annulipes* [Kemp].
Unique type in Indian Museum.
- glabra*, *id.*, *loc. cit.*, 325, ♂ (1918).
Taken in company with *annulipes* (11—26-x-16) [Kemp].
Unique type in Indian Museum.
- fusca*, *id.*, *loc. cit.*, 326, ♂ ♀ (1918).
Sureil, Darjiling District, 5000 ft., iv—v-17 [Kemp].
Types in Indian Museum.
- incompleta*, *id.*, *loc. cit.*, 326, ♂ (1918).
Taken in company with *fusca*. Type in Indian Museum.
- inconsequens*, *id.*, *loc. cit.*, 326, ♂ (1918).
Castle Rock, N. Kanara Distr., 11—26-x-16 [Kemp]; Cooroor, 3-ix-11, banks of a rivulet [Howlett]. Type in Indian Museum.
- flavipennis*, *id.*, *loc. cit.*, 327, ♀, pl. viii, 14, wing (1918).
Darjiling, 6900 ft. 17-v-17 [Brunetti].
Unique type in my collection.
- ornatipennis*, *id.*, *loc. cit.*, 328, ♂ ♀, pl. vii, 5, wing (1918).
Soom, Darjiling District, 4000—5000 ft., 14-vi-14, type ♂;
Darjiling, 7000 ft., 12-vi-14, type ♀ [both Gravelly].
Types in Indian Museum.

sp. incert., *enchroma* Walk., Proc. Linn. Soc. Lond. VI, 4 (*Limnobia*) (1862).

Brun., Rec. Ind. Mus. VI, 306.

This species is doubtful, described originally from Gilolo, and apparently not seen since. Osten Sacken refrained from confirming the single specimen as a *Limnophila*, as the antennae were missing. The *type* is now legless also but the conspicuously marked wings would render identification of the species possible.

Dicranophragma, Ost. Sack.

Proc. Acad. Nat. Sci. Philad. 240 (1859).

Brun., Fauna Brit. Ind., 524.

GENOTYPE, *D. fuscovaria* Ost. Sack., by original designation.

pulchripennis Brun., Fauna Brit. Ind., 524, ♂ (1912).

Kurseong, 4—9-ix-09, *type* ♂; 27-vi-10 [both *Annandale*]; Darjiling, 8—11-viii-09, *type* ♀ [*Paiva*]; Kalimpong, 600—4500 ft., 24-iv—10-v-15 [*Gravely*]; Bhoirakund, Darrang Distr., Assam-Bhutan Frontier, 8—22-x-12 [*Kemp*] (with a doubt).

Types in Indian Museum.

gracilis, *id.*, Rec. Ind. Mus., VIII, 156, ♀, pl. vi, 3, wing (1913).

Yembung, N.E. Frontier, India, 1100 ft., 13-i-12 [*Kemp*].

Type in Indian Museum, a unique specimen.

remota de Meij., Tijds. v. Ent. LVI, Supp., p. 1 (1914).

Buitenzorg, iv-09 [*Bryant and Palmer*]; Nongkodjadar, Java, ♂ ♀ [*Jacobson*]; Tjibodas, Mt. Gedé, Java, 4500 and 9000 ft. [*Bryant and Palmer*]. *Types* in Amsterdam Museum.

multipunctipennis Brun., Rec. Ind. Mus. XV, 319, ♂; pl. viii, 17 (1918).

Darjiling, 15—19-v-17. Unique *type* in my collection.

Epiphragma, Ost. Sack.

Proc. Acad. Nat. Sci. Philad. 238 (1859).

Sch., F. Austr. II, 550: Ost. Sack., Monog.

Dipt. N. Amer. IV, 193, pl. i, 8: *id.*, Berl. Ent.

Zeits. XXXI, 208.

GENOTYPE, *Limnobia fascipennis* Say (as *pavonina*, sp. nov.), by Coquillett's designation.

insignis Wulp, Tijds. v. Ent. XXI, 196, pl. xii, 6 (1878).

Sumatra. *Type* in Amsterdam Museum, in bad condition.

signata de Meij., *loc. cit.*, LIV, 52, ♂, pl. iv, 43, wing (1911).

Buitenzorg environs, Java, xi [*Jacobson*]; Tjibodas, Mt. Gedé, Java, 4500 ft. [*Bryant and Palmer*].

Type, a unique ♂, in Amsterdam Museum.

- kempi** Brun., Rec. Ind. Mus. VIII, 155, ♀, pl. vi, 2, wing (1913).
Edw., Ann. Mag. Nat. Hist. (8) XII, 204 (1913): Brun.,
Rec. Ind. Mus. XV, 330.
N.E. Frontier, India, 4000 ft., 16-iii-12 [*Kemp*]; Hakgala,
Ceylon, 4500 ft., v-1911; Castle Rock and Talewadi, N. Kanara
Distr. [*Kemp*]; Tura, Garo Hills, vii, viii-17 [*Kemp*].
Type in Indian Museum.
- klossi** Brun., Rec. Ind. Mus. XV, 330, ♂, pl. viii, 16, wing (1918).
Ginting Bidai, 2,000 ft., Selangor-Pahang Frontier, Fed. Malay
Sts., iv-17 [*Boden Kloss*]. Unique type in Indian Museum.
- vicina**, *id.*, *loc. cit.*, 331, ♀; pl. viii, 15, wing (1918).
Sureil, Darjiling Distr., 5,000 ft., 11-31-x-17 type [*Annandale*
and *Gravelly*]; above Tura, Garo Hills, 3,500-3,900 ft., vii-17,
viii-17 [*Kemp*]. Type in Indian Museum.

Ephelia, Sch.

- Wien. Ent. Monats. VII, 222 (1863).
Sch., F. Austr. II, 549: Brun., Fauna Brit.
Ind., 525.
GENOTYPE, *Limnobia marmorata* Hf. in Mg., by personal
designation (Fauna, p. 525).
Elocophila, Rond., Dipt. Hal. Prod. I, 182 (1856).
GENOTYPE, *Limnobia marmorata* Hf. in Mg.
? *Nasiterna*, Wallengren, Ent. Tidskr. II, 179 (1881).
- fascipennis** Brun., Fauna Brit. Ind., 526, ♂, pl. x, 16, wing
(1912).
Kurseong, 13-viii-09 [*Paiva*]; Pashok, 3500 ft. [*Hartless*];
Arisan, Formosa, 8000 ft., 10-x-12 [*Nitobe*].
Type in Indian Museum.
- ornata**, *id.*, *loc. cit.*, 527, ♀ (1912).
Bhowali, Kumaon, 27-vi-10 [*Imms*].
Type in Indian Museum, a unique specimen.

Poecilostola, Sch.

- Wien. Ent. Monats. VII, 222 (1863).
Sch., F. Austr. II, 551.
GENOTYPE, *Limnobia pictipennis* Mg., by original
designation.
- pallens** Wulp, Notes Leyd. Mus. VII, 13 (1885).
Id., Tijd. v. Ent. XXVIII, 86, pl. iv, 8.
Java [*Kuhl*]. Type in Leyden Museum.

Ula, Hal.

- Entom. Mag. I, 153 (1833).
GENOTYPE, *Limnobia macroptera* Macq. as *U.*
mollissima, sp. nov.

javanica Alex., Proc., U.S. Nat. Mus., 176, pl. xlv, 24, wing (1916).

Tjibodas, Mt. Gedé, Java, 8000 ft. [Bryant and Palmer].

Type, Cat. No. 19,025, U.S. Nat. Mus.

Alexander removes this genus to the Linnophilini having bred a North American species and found the larva quite linnophiline.

Gynoplistia, Westw.

Lond. and Edinb. Philos. Mag. VI, 280 (1835).

Westw., Trans. Ent. Soc. Lond., 369 (1881), pl. xviii, 5, 6; Ost. Sack., Berl. Ent. Zeits. XXXI, 210; Brun., Rec. Ind. Mus. VI, 306 (notes on Orient. spp.).

GENOTYPE, *Ctenophora vilis* Walk., by personal designation, 1918.

Anoplistes, Westw., Zool. Jour., No. 20, ined.

GENOTYPE unknown to present author.

Caenarthria, Thoms., Eug. Resa, 445, pl. ix, 1.

GENOTYPE, *Gynoplistia viridis* Westw., by original designation.

Cloniophora, Sch., Verh. zool. bot. ges. Wien XVI, 932 (1866).

GENOTYPE, *Gynoplistia subfasciata* Walk., by original designation.

jurgiosa Walk., Proc. Linn. Soc. Lond. III, 78, ♂ ♀ (1859).

Aru. *Types*, in good condition, in British Museum.

fulviceps, *id.*, *loc. cit.*, V, 231, ♀ (1861).

Dorey, Papua. *Type* in British Museum in good condition, except that the abdomen is missing.

melancholica, *id.*, *loc. cit.*, VII, 203, ♂ ♀ (1864).

de Meij., Bijd. tot de Dierk. XIX, 41, notes.

Mysol, Waigon, 7-i-10. *Type* in British Museum in good condition, except that the abdomen is missing.

insolita Walk., Proc. Linn. Soc. Lond. VIII, 131, ♀ (1865).

Salawatti. *Type* in good condition in British Museum.

jucunda Ost. Sack., Ann. Mus. Gen. XVI, 405, ♂ ♀ (1880).

Kandari, Celebes. *Types* originally in Genoa Museum, only one now remaining.

8-fasciata Brun., Rec. Ind. Mus., VI, 307, ♀ (1911).

Patuhuang, South Celebes. Unique *type* in Vienna Museum.

occipitalis de Mej., Nova Guin. Res. IX, 310, ♂ (1913).

River Kamp, Papua, 12-ii-10.

Unique *type* in Amsterdam Museum.

Section ANISOMERINI.¹

Eriocera, Macq.

Dipt. Exot. I, 1, 74, pl. x, 2 (1838).

Ost. Sack., Mong. Dipt. N. Amer. IV, 244:
id., Berl. Ent. Zeits. XXXI, 220: Brun., Rec.
 Ind. Mus. VI, 307, notes, orient. spp.; XV, 334,
 tab. orient. spp.: *id.*, Fauna Brit. Ind., 530:
Auctt.

GENOTYPE, *Limnobia nigra* W., by Macquart's designation.

Caloptera, Guer., Voy. de la Coq., Zool., Ins., pl. xx, (1830) *vide Evanioptera*, *post.*

GENOTYPE, *C. fasciata*, *id.*, sp. nov., *loc. cit.*

Evanioptera, *id.*, *loc. cit.*, Zool. II. 287, pl. xx, 2, full ins., 2a, head in profile; 2b, wing. (in plate as *Caloptera*, in text renamed *Evanioptera*)² (1830), *nom. nov.* for *Caloptera* under alleged pre-occupation.

Pterocosmus, Walk., List Dipt. Brit. Mus. I. 78 (1848).

GENOTYPE, *P. hilpa* Walk., by Coquillett's designation.

Oligomera, Dol., Nat. Tijd. Ned. Ind. XIV, 387 (1857).

GENOTYPE, *Limnobia acrostacta* W., as *javensis*, sp. nov., by original designation.

Physecrania, Big., Ann. Soc. Ent. Fran. (3) VII, 123 (1859).

GENOTYPE, *P. obscura* Big., sp. nov., *loc. cit.*

Arrhenica, Ost. Sack., Proc. Acad. Nat. Sci. Phil., 243 (1859).

GENOTYPE, *A. spinosa* Ost. Sack., by Coquillett's designation.

Androclosma, Ender., Zool. Jahr. XXXII, 34 (1912).

GENOTYPE, *Megistocera verticalis* Wied., by original designation.

acrostacta Wied., Dipt. Exot. I, 14 (*Limnobia*) (1821).

Id., Auss. Zweifl. I, 26, ♂ ♀ (*Limnobia*): Macq., Dipt. Exot. I, 1, 68, pl. ix, 1, full ins., 1a, head profile (*Cylindrotoma*): *id.*, *loc. cit.*, Supp. I, 18, pl. ii, 6, wing, 6a, antenna (*Limnobia*): *id.*, *loc. cit.*, Supp. III, 7 (*Cylindrotoma*): Wulp, Notes

¹ For characters, *v.* Brun., Fauna Brit. Ind., 529.

² In the text, Guerin changes the name of the genus on account of having been previously employed by the Count of Casteinau in Coleoptera, but I cannot trace any such genus. The name *Caloptera*, however, has been suggested in four different cases subsequent to its use by Guerin.

Leyd. Mus. VII, 11: Alex., Proc. U.S. Nat. Mus. XLIX, 177, pl. xlv, 26, wing: Brun., Rec. Ind. Mus. XV, pl. viii, 19, wing.

Oligomera javensis, Dol., Nat. Tijd. Ned. Ind. XIV, 387, pl. vii, 3, full ins. (1857).

? *Limnobia diana*, Macq., Hist. Nat. Dipt. I, 107 (1834): Walk., List Dipt. Brit. Mus. I, 42 (1848).

Java [Muller]; Tjibodas, Mt. Gedé, Java, iv-09 [Bryant and Palmer]; ? Bengal.

Types in Wiedemann's and Westermann's collections.

basilaris Wied., Dipt. Exot. I, 15, ♂ ♀ (*Limnobia*) (1821).

Id., Auss. Zweifl. I, 27 (*Limnobia*): Alex., Proc. U.S. Nat. Mus. XLIX, 177, pl. xlv, 27, wing.

Java, var. locs., iii, iv, v, xii [*t. de Meij.*]; Depok, 10-ii-09; Tjibodas, Mt. Gedé, Java, iv-09 [Bryant and Palmer].

Type in Westermann's collection.

mesopyrrha Wied., Auss. Zweifl. I, 26, ♂ (*Limnobia*) (1828).

de Meij., Tijd. v. Ent. LIV, 55: Brun., Rec. Ind. Mus. VI, 307: Ender., Zool. Jahr. XXXII, 33 (*Physecrania*); Alex., Proc. U.S. Nat. Mus. XLIX, 177, pl. xlv, 28, wing.

Sumatra, Gunung Ungaran, Java, xii [Jacobson]; Buitenzorg, iii-09 [Bryant and Palmer]. Types in Leyden Museum.

verticalis Wied., Auss. Zweifl. I, 56, ♂ (*Megistocera*) (1828).

Wulp., Tijd. v. Ent. XXXVIII, 39, pl. ii, 6: de Meij., *loc. cit.*, LIV, 55: Brun., Rec. Ind. Mus., VI, 307; XV, 333: Alex., Proc. U.S. Nat. Mus. XLIX, 177, pl. xlv, 25, wing.

? *Megistocera atra* Dol., Nat. Tijd. Ned. Ind. XVII, 80 (1858).

Bantar Gebang, Java, 21-x-09 [Bryant and Palmer]; Semarang, xi; Wonosobo, iv, v [both Jacobson]; Taihoku, Formosa [Shiraki]; Laut Tawar, viii; Sinabang, ii; both Simalur Is., off Sumatra; Sandaran Agong, Sumatra, 2450 ft., v, vi-14.

Type from Java in Leyden Museum in bad condition.

nepalensis Westw., Ann. Soc. Ent. Fran. IV, 681 (*Caloptera*) (1835).

Id., Trans. Ent. Soc. Lond. 379 (*Caloptera*) (1881): Brun., Rec. Ind. Mus. VI, 308 also as *velutina*: *id.*, *loc. cit.*, XV, pl. vii, 13, wing: *id.*, Fauna Brit. Ind., 543.

velutina Walk., List Dipt. Brit. Mus. I, 79 (*Pterocosmus*) (1848).

Nepal, and generally distributed along the Himalayas, the Bhutan-Frontier and Assam, Shillong and Peshawar.

Type in the Hope collection at Oxford University.

bicolor Macq., Dipt. Exot. I, 1, 66, ♀, pl. vii, 2, full ins. (*Limnophila*) (1838).

Wulp., Mid Sum. Exped., Dipt. II, pl. 1, 5, 6¹: de Meij., Tijd. v. Ent. LIV, 57 (♂ descr.): Ender., Zool. Jahr. XXXII,

¹ Wulp erroneously gives figs. 4 and 5.

- 33 (*Physecrania*): Brun., Fauna Brit. Ind., 545: *id.*, Rec. Ind. Mus. XV, pl. vii, 6, wing.
Bengal; Koetoer, Sumatra; Sandaran Agong, Korinchi Lake, Sumatra, 2450 ft., iv, v-1914; Java.
Type in Paris Museum.
- sumatrensis* Macq., Dipt. Exot., Supp. IV, 16, ♂, pl. 1, 7, full insect (*Limnobia*) (1851).
Brun., Rec. Ind. Mus. XV, pl. vii, 14, wing.
Sumatra. *Type* in Paris Museum.
- leucotelus* Walk., Proc. Linn. Soc. Lond. I, 6, ♂ (*Limnobia*) (1857).
Singapore.
Type in British Museum, in fair condition, but legless.
- plecioides*, *id.*, *loc. cit.*, I, 6, ♂ (*Limnobia*) (1857).
Singapore.
Type in British Museum, only one leg remaining.
- dichroa*, *id.*, *loc. cit.*, I, 6, ♀ (*Limnobia*) (1857).
Brun., Rec. Ind. Mus. VI, 308.
Mount Ophir. *Type* in British Museum, in bad condition.
- rubescens*, *id.*, *loc. cit.*, I, 106, ♂ (*Limnobia*) (1857).
Borneo. *Type* in British Museum, in fair condition.
- pyrrhochroma*, *id.*, *loc. cit.*, I, 106, ♂ (*Limnobia*) (1857).
Borneo. *Type* in British Museum, in bad condition.
- lunigera*, *id.*, *loc. cit.*, I, 107, ♂ (*Pterocosmus*) (1857).
de Meij., Tijd. v. Ent. LIV, 56, ♂ ♀: Ender., Zool. Jahr. XXXII, 39, ♂ ♀ redescr.: Brun., Rec. Ind. Mus. VI, 308.
Borneo; Padang, etc., Sumatra; Buitenzorg, ix [Jacobson].
Type in British Museum in bad condition.
- infixa*, Walk., Proc. Linn. Soc. Lond., I, 107, ♂ ♀ (*Pterocosmus*) (1857).
Borneo.
- optabilis*, *id.*, *loc. cit.*, I, 107, ♀ (*Pterocosmus*) (1857).
Brun., Rec. Ind. Mus. VI, 309.
Borneo. *Type* in British Museum, valueless, apparently unique.
- combinata*, *id.*, *loc. cit.*, I, 107, ♀ (*Pterocosmus*) (1857).
Brun., Rec. Ind. Mus. VI, 309.
Borneo. *Type* in British Museum, damaged.
- diluta*, *id.*, *loc. cit.*, I, 108 (*Pterocosmus*) (1858).
Brun., Rec. Ind. Mus. VI, 309.
Borneo. *Type* in British Museum, valueless, sex unstated by Walker, now undiscoverable.
- albonotata* Loew, Peter's Natur. Reise Mozambique, Zool. V (Insek.) pl. i, 1 (*Limnobia*) (1862).
Ost. Sack., Berl. Ent. Zeits. XXXI, 223: v. Roder, Entom. Nachr. XIX, 234: Brun., Rec. Ind. Mus. VI, 308: *id.*, Fauna Brit. Ind., 547 ♂ ♀ redesc.

Peradeniya, 10-x [Humbert]; Caunia, near Trincomalee, 30-vii; Ceylon (var. locs. *t.* Edwards); Maskeliya [Green].

Described originally from Mozambique.

Type in Berlin Museum.

albipunctata Wulp, Tijds. v. Ent. XXIII, 158, pl. x, 3 antenna, 4, wing (1880).

Id., Notes Leyd. Mus. VII, 11: Brun., Rec. Ind. Mus. VI, 309.

Java, Semarang, i. *Type* originally in Amsterdam Museum, now lost, but other specimens (presumably *co-types*) present.

morosa Ost. Sack., Ann. Mus. Gen. XVI, 406, ♀ (1880).

Makassar. *Type* in Genoa Museum in fair condition.

selene, *id.*, *loc. cit.*, 406, ♀ (1880).

Enderl., Zool. Jahr. XXXII, 39: Brun., Rec. Ind. Mus. VI, 309.

Sumatra (Mt. Singalang).

Type in Genoa Museum in fair condition.

lunata Westw., Trans. Ent. Soc. Lond. 1881, 367, pl. xviii, 4.

Enderl., Zool. Jahr. XXXII, 35.

Sarawak. *Type*, unless in the Hope collection at Oxford, probably lost, no sex stated.

perennis Ost. Sack., Berl. Ent. Zeits. XXVI, 91, ♂ ♀ (1882).

Philippine Is. *Types*, a unique pair; location unknown.

mansueta, *id.*, *loc. cit.*, 92, ♂ ♀ (1882).

Philippines. *Types*, location unknown; probably originally in Osten Sacken's collection.

ferruginosa Wulp, Notes Leyd. Mus. VII, 13 (1885).

Enderl., Zool. Jahr. XXXII, 32, fig. S (wing) (*Physecrania*):

Brun., Rec. Ind. Mus. VI, 309.

Java [Kuhl]; Semarang [Jacobson]; Sumatra.

Type in Leyden Museum.

humberti Ost. Sack., Berl. Ent. Zeits. XXXI, 221, ♀ (1887).

Brun., Fauna Brit. Ind., 536, ♀.

Pundeloya Valley, Ceylon, 3,800—3,900 ft., 26-xi—27-xii-1859 [Humbert]. *Type* in Geneva Museum. Edwards thinks

a ♂ from Uva Prov., Madulsima, Ceylon, may be this species.

meleagris Ost. Sack., Berl. Ent. Zeits. XXXI, 221, ♀ (1887).

Brun., Fauna Brit. Ind., 537, ♀.

Between Nuwara Eliya and Gampola, Ceylon, 13—14-xii-1859 [Humbert]. Unique ♀ in Geneva Museum.

pachyrrhina Ost. Sack., Berl. Ent. Zeits. XXXI, 222, ♂ ♀ (1887).

Brun., Rec. Ind. Mus. VI, 309: *id.*, Fauna Brit. Ind., 537.

Kaduganawa, Ceylon, 8-x [Humbert].

Types, *in cop.*, in Geneva Museum.

crystalloptera Ost. Sack., Berl. Ent. Zeits., XXXI, 222, ♂ (1887).

Brun., Rec. Ind. Mus. VI, 309: *id.*, Fauna Brit. Ind., 541.

Ceylon (Rambodde) [*Nietner*]; Madulsima, Pundaluoya, Haputule (all Ceylon ♂ ♀ and *t.* Edwards).

Type (an imperfect ♂) in Berlin Museum.

nigripennis de Meij., *Bijd. tot de Dierk.* XVII, 92, ♀ (1904).

Java. Riedel notes a ♀ from Formosa. It would fall in *Physocrania*. *Type* in Amsterdam Museum.

cingulata de Meij., *Tijd. v. Ent.* LIV, 58, ♂ ♀ (1911).

Alex., *Proc. U.S. Nat. Mus.* XLIX, 178, pl. xlv, 29, wing.

Pangeranga, Java; Berg Ungarar, x, xii [*Jacobson*]; Depok, 10-ii-09 [*Bryant and Palmer*].

ctenophoroides Edw., *Ann. Mag. Nat. Hist.* (8) VIII, 64, ♂ ♀ (1911).

Brun., *Fauna Brit. Ind.*, 533.

Kandy, 19-v-92, *type* ♂; Kottawa, 29-iv-92, *type* ♀; Pallamadulla, 17-vi-92 [all Ceylon and *Yerbury*].

Types in British Museum. Edwards (*l.c.*) notes a variety.

scutellata Edw., *Ann. Mag. Nat. Hist.* (8) VIII, 65, ♂ ♀ (1911).

Brun., *Fauna Brit. Ind.*, 534.

Pundaluoya, Ceylon, ix-92 *type* ♂; x-1897, *type* ♀ [*E. E. Green*]. *Types* in British Museum.

tuberculifera Edw., *Ann. Mag. Nat. Hist.* (8) VIII, 66, ♀ (1911).

Brun., *Fauna Brit. Ind.*, 547.

Pundaluoya, Ceylon, xi-88 (*type*) [*E. E. Green*]; vii-89.

Type in British Museum.

fusca Edw., *Ann. Mag. Nat. Hist.* (8) VIII, 66, ♂ ♀ (1911).

Brun., *Fauna Brit. Ind.*, 541.

Pundaluoya, Ceylon, iv-89 [*E. E. Green*].

Types in British Museum.

badia Brun., *Rec. Ind. Mus.* VI, 310, ♀ (1911).

Id., *Fauna Brit. Ind.*, 538, ♀.

Peradeniya, 25-xii-01 [*Uzel*].

Type in Vienna Museum, a unique specimen.

rufibasis Brun., *Rec. Ind. Mus.* VI, 310, ♀ (1911).

Id., *Fauna Brit. Ind.* 538, ♀.

Tandong, Tenasserim, 4000 ft., May [*Fruhstorfer*].

Type in Vienna Museum, unique.

semilimpida Brun., *Rec. Ind. Mus.* VI, 311, ♂ (1911).

Id., *Fauna Brit. Ind.*, 546 ♂ pl. x, 17, wing.

Khasi Hills, ix-1906, *type*; Nongpoh, Assam, vi-1906.

Type in the Pusa Collection.

plumbicincta Brun., *Rec. Ind. Mus.* VI, 311, ♂ (1911).

Id., *Fauna Brit. Ind.*, 541, ♂, pl. xi, 16, abdomen: *id.*,

Rec. Ind. Mus. XV, pl. vii, 11, wing.

Ukhrul, Manipur, viii-08, *type* [*Pettigrew*]; Darjiling, 3-9-vi-09, *cotype* [*Howlett*]; Phooboering, Lebong, 5000 ft., ix-10 [*Lefroy*]. *Type* in Indian Museum, *cotype* in Pusa coll.

- fenestrata** Brun., Rec. Ind. Mus. VI, 312, ♂ ♀ (1911).
Id., Fauna Brit. Ind., 535, ♂ ♀, pl. xi, 14, wing (part) 15, abdomen.
 Central Tonkin, *type* ♂; Khasi Hills, 1000—3000 ft., iv-1905, *type* ♀. *Type* ♂ in Vienna Museum; *type* ♀ in Pusa collection.
- greeni** Brun., Rec. Ind. Mus. VI, 313, ♀ (1911).
Id., Fauna Brit. Ind., 539, ♀: Rec. Ind. Mus. XV, pl. vii, 8, wing.
 Kandy, 24-xi-09 [Green].
Type in Indian Museum, a unique specimen.
- angustipennis** Ender., Zool. Jahr. XXXII, 33, ♂, fig. T, wing (*Physocrania*) (1912).
 Soerakanda, Sumatra.
Type, a unique ♂, in Stettin Museum.
- pannosa**, *id.*, *loc. cit.*, 40, ♂ ♀, fig. W, wing (1912).
 Soerakanda. *Types*, a unique pair, in Stettin Museum.
- gamma**, *id.*, *loc. cit.*, 42, ♂ ♀, fig. X, wing (1912).
 Soerakanda and Liangagas, Sumatra.
Types in Stettin Museum.
- sauteriana**, *id.*, *loc. cit.*, 42 ♀, fig. Y, wing.
 Takao, South Formosa, Kotosho Is., Formosa, Arisan, Formosa, 8000 ft., 10-x-12 [Nitobe]. *Type* in Stettin Museum.
 Reidel notes two ♂♂ from Formosa.
- paenulata**, *id.*, *loc. cit.*, 43, ♂, fig. Z, wing.
 Soerakanda; Sungei Penoh, Korinchi Valley, Sumatra, 2600 ft., iii-14. *Type* in Stettin Museum.
- rufithorax** Brun., Fauna Brit. Ind., 534, ♂ ♀ (1912).
Id., Rec. Ind. Mus. XV, pl. vii, 10, wing.
 Kandy, viii-06, *type* ♂ ♀, ix-02; x-07, *cotype* ♂; Hadumalla, Ceylon [Green].
Type ♂ ♀ in Indian Museum, *cotype* ♂ in my collection.
- tenuis**, *id.*, *loc. cit.*, 539, ♂ (1912).
 Nilgiri Hills, 3500 ft., iv-1910 [Andrewes]; Gudalur, Nilgiri Hills, 3000, 13-ix-17 [Ras].
Type in Indian Museum, a unique specimen.
- aterrima**, *id.*, *loc. cit.*, 540, ♀ (1912).
 Maddathorai (Travancore), 18-xi-08 [Annandale].
Type, a unique ♀, in Indian Museum.
- elongatissima**, *id.*, *loc. cit.*, 542, ♂ (1912).
 Trivandrum, vi-93, *type*; Aspamboo, Travancore, South India, vi-93. *Type* in Indian Museum.
- flavipes**, *id.*, *loc. cit.*, 544, ♂ (1912).
Id., Rec. Ind. Mus. XV, 333, pl. vii, 15, wing.
 Kurseong, 9-ix-09 [Annandale].
Type in Indian Museum, a unique specimen.

- testacea*, *id.*, Fauna Brit. Ind., 548, ♀ (1912).
Nilgiri Hills, iv-1910 [*Andrewes*].
Type, a unique ♀, in Indian Museum.
- nigerrima*, *id.*, Fauna Brit. Ind., App. 571, ♀ (1912).
Id., Rec. Ind. Mus. XV, pl. vii, 16, wing.
Ghunti, vii-1911 [*Gravely*].
Unique type in Indian Museum.
- triangularis*, *id.*, Fauna Brit. Ind., App. 572, ♀ (1912).
Nilgiri Hills, 6000 ft., v-1911 [*Andrewes*].
Type in Indian Museum.
- nigrina* Riedel, Ent. Mitt. Berl., II, 273 (1913).
Kankao, Formosa, ♂ ♀, vii, ix-1912.
Type in Deut. Ent. Mus., *cotype* in Riedel coll.
- xanthopyga* de Meij., Tijd. v. Ent. LVI, Supp. p. 3, ♂ ♀, 1913
(Mar. 1914).
Nusa Kambangan, Java, iii [*Jacobson*].
- unicolor* de Meij., Tijd. v. Ent. LVIII, Supp. 12, ♂ ♀ (1916).
Simalur Island, Sumatra; Horisha, Formosa, 10-x-12 [*Nitobe*];
1000 ft. [*Maki*]. Types in Amsterdam Museum.
- simalurensis*, *id.*, *loc. cit.*, 13, ♂ ♀ (1916).
Sinabang, ii, Laulo, viii, both, Simalur Island, off W. Coast,
Sumatra. Types in Amsterdam Museum, a unique pair.
- rubriceps* Edw., Ann. Mag. Nat. Hist. (8) XVIII, 253, ♀ (1916).
Taipin, Formosa, 12-x-10 [*Shiraki*].
Unique type in British Museum.
- lativentris* Bezzi, Phil. Jour. Sci. XII, Sect. D, 113, ♂ (1917).
Luzon [*Baker*].
- crassipes*, *id.*, *loc. cit.*, 114, ♂ ♀ (1907).
Luzon [*Baker*].
- cincta* nom. nov., Brun., Rec. Ind. Mus. XV, 333, pl. vii, 7, wing
(1918).
E. cingulata Brun., Fauna Brit. Ind., App. 570, ♂ (1912),
preocc. de Meij., 1911.
- aurantia* Brun., Rec. Ind. Mus. XV, 335, ♀, pl. vii (1918).
Pashok, 2500 ft. [*Gravely*]; Kalimpong, Darjiling, 600—4500
ft., 24-iv—10-v-14 [*Gravely*]. A ♂ from Darjiling, 7000 ft.,
12-vi-14 [*Gravely*], is probably the ♂ of this species.
Type in Indian Museum.
- nigroapicalis* Brun., *loc. cit.*, 335, ♀ (1918).
Kalimpong, Darjiling, 24-iv—10-v-14 [*Gravely*].
Unique type in Indian Museum.
- pulchrithorax*, *id.*, *loc. cit.*, 336, ♀ (1918).
On forest tramway, miles 10 to 14, 0—300 ft., Cochin State,
28—29-ix-14 [*Gravely*]. Unique type in Indian Museum.

- rufiventris*, *id.*, *loc. cit.*, 336, ♂ ♀ (1918).
Parambikulam, Cochin State, 16-24-ix-14 [*Gravelly*]; Talewadi, 3-10-x-16; Castle Rock, 11-26-x-16; Sidapur, Coorg, S. Ind., 7-vi, 2-vii, 20-ix-17. *Types* in Indian Museum.
- gravelyi*, *id.*, *loc. cit.*, 337, ♂ ♀, pl. vii, 9, wing (1918).
Pashok, Darjiling District, 1000-2500 ft., 26-v-14-vi-16 [*Gravelly*]; Sureil, iv-v-17 [*Kemp*]; Tura, Garo Hills, vii-17; x-17 [*Kemp*]. *Types* in Indian Museum.
- decorata*, *id.*, *loc. cit.*, 337, ♂, pl. vii, 12, wing (1918).
Pashok, Darjiling District, 2000 ft., 26-v-14-vi-16 [*Gravelly*].
Type in Indian Museum.
- tripunctipennis*, *id.*, *loc. cit.*, 338, ♂ ♀, fig. 7, wing (1918).
Castle Rock, N. Kanara Dist., 11-26-x-16 *type* ♂; Talewadi, N. Kanara Dist., 3-10-x-16, *type* ♀ [*Kemp*].
Types in Indian Museum.
- caliginosa*, *id.*, *loc. cit.*, 339, ♀ (1918).
Talewadi, N. Kanara Dist., 3-10-x-16 [*Kemp*].
Type in Indian Museum.
- kempi*, *id.*, *loc. cit.*, 339, ♂ ♀ (1918).
Castle Rock, N. Kanara Dist., 11-26-x-16 [*Kemp*].
Types in Indian Museum.
- maculiventris*, *id.*, *loc. cit.*, 340, ♂ ♀, pl. viii, 18, wing (1918).
Tura, Garo Hills, 1200-1500 ft., vii-17; above Tura, 3900 ft., vii, viii, x-17 [all *Kemp*].
- pyrrhomesa* Edw., Jour. Fed. Malay Sts. Mus. VIII, 21, ♂ ♀ (s.g. *Physcerania*) (1919).
Sandaran Agong, Korinchi Peak, Sumatra, 2450 ft., v, vi-14.
Types in British Museum.
- klossi*, *id.*, *loc. cit.*, 22, ♀, pl. iv, 14, wing (s.g. *Ericcera*) (1919).
Sandaran Agong, Sumatra, 2450 ft., v, vi-14.
Type in British Museum.

Family RHYPHIDAE.

Rhyphus, Latr.

Hist. Nat. Crust. et Ins. XIV, 291 (1804).

Sch., F. Austr. II, 494: Brun., Rec. Ind. Mus. IV, 259, tab. Ind. spp.: *id.*, Fauna Brit. Ind., 500.GENOTYPE, *Tipula fenestralis* Scop., the original species, as *fenestratum* Scop.*Sylvicola*, Harris, Expos. Engl. Ins., 104 (1776).GENOTYPE, *Tipula fenestralis* Scop., as *brevis*, sp. nov.; by Coquillett's designation.*Anisopus* Mg., Illig. Mag. II, 264 (1803).No species given in 1803. In "Klass." p. 103, *Tipula fuscata* F. as *fuscus*, sp. nov., is given as genotype, according to Coquillett.

fenestralis Scop.,¹ var. **indicus** Brun., Rec. Ind. Mus. IV, 261, ♂ ♀ (1911).

Brun., Fauna Brit. Ind., 554, ♂ ♀, pl. xii, 3 (typical), 4 (var.).
Simla, 24-iv-07, 10-v-09; Matiana, 28—30-iv-07; [both *Annandale*]; Darjiling 5—9-viii-09 [*Paiva*]; Kurseong, xi-10 [*D'Abreu*]; Ukhurul, Manipur, Assam, 6400 ft. [*Pettigrew*].

Type of fenestralis Scop., location uncertain; of var. *indicus* in Indian Museum. This form seems present at all the Hill Stations in Northern India.

punctatus F., Mant. Ins. II, 333 (*Rhagio*) (1787).

Musca nigricans, L., Fauna Suec., Ed. II, 553 (1761).

Rhagio punctatus, F., Mant. Ins. II, 333 (1787).

Musca bilineata, Gmel., Syst. Nat. V, 2866 (1792).

Anisopus nebulosus, Mg., Klass. I, 103, pl. vi, 5, ♂ (1804).

Sciara punctata, F., Syst. Antl., 59 (1805).

Rhyphus marginatus, Say, Jour. Acad. Sci. Phil. III, 27 (1822), and Compl. Writ. II, 50 (1859); Wied. Auss. zweifl. I, 82 (1828).

Rhyphus punctatus Sch., Fauna Austr. II, 495 (1864); Brun., Rec. Ind. Mus. IV, 262: *id.*, Fauna Brit. Ind. 555, pl. xii, 5, wing.

Kurseong, 9—26-ix-09 [*Lynch*, *Annandale*].

Type: location uncertain.

maculipennis Wulp, Notes Leyd. Mus. VII, 14, ♂ (1885).

Id., Tijid. v. Ent. XXVIII, 87, pl. iv, 9: *id.*, loc. cit., XXXVIII, 41; LIV, 258; Brun., Rec. Ind. Mus. IV, 260: *id.*, Fauna Brit. Ind., 552, ♂ ♀, pl. xii, 1, wing.

Ardjoeno, Java [*Hekmeyer*]; Batavia, ii; Pulu Babi Is., off W. Coast, Sumatra, iv; Peradeniya, x, xii-07; also probably Assam (exemp. in Ind. Mus. coll. *sine loc.*, 30-vii-10).

Type in Leyden Museum, a ♂.

pulchricornis Brun., Rec. Ind. Mus. IV, 260, ♀ (1911).

Id., Fauna Brit. Ind. 553, ♀, pl. xii, 2, wing.

Siliguri, 18—20-vii-07; Ghumti, 4000 ft. [*Gravelly*].

Type in Indian Museum.

distinctus Brun., Rec. Ind. Mus. IV, 262, ♂ ♀ (1911).

Id., Fauna Brit. Ind. 556, ♂ ♀, pl. xii, 6, wing.

Darjiling, 5—8-viii-09 [*Paiva*]; Kurseong, 4-ix-9 [*Annandale*]; Kowpati, Mangaldai, Assam, 12-xii-10 [*Kemp*].

Types in Indian Museum.

divisus Brun., Rec. Ind. Mus. IV, 263, ♂ ♀ (1911).

Id., Fauna Brit. Ind. 557, ♂ ♀, pl. xii, 7, wing.

Darjiling, 5—12-viii-09 [*Jenkins*, *Paiva*, common]; Kurseong, 10—26-ix-09 [*Lynch*]; Gangtok, Sikkim, 9-ix-09; Phagu, 12-v-09 [*Annandale*]. Two pairs taken *in cop*; Darjiling, 5-viii-09 and Gangtok, 6150 ft., 8-ix-09.

Types in Indian Museum.

¹ For references to typical form see Kertész' Kat. Dip. I, 305.

flavipes de Meij., Tijds. v. Ent. LIV, 259, ♂ ♀, pl. xviii, 1, wing (1911).

Wonosobo, Java [*Jacobson*].

Types, a unique pair, in Amsterdam Museum.

hellwigi de Meij., Nova Guin. Res. IX, 315, ♀, pl. x, 8, wing (1913).

Hellwig-Gabirge, Papua, 2000 metres, xi.

Type in Amsterdam Museum.

Olbogaster, Ost. Sack.

Biol. Cent. Am. Dipt. I, 20 (1886).

GENOTYPE *O. cognata* Ost. Sack., sp. nov.

orientalis Edw., Ann. Mag. Nat. Hist. (8) XVI, 504, ♀ (1915).

Pundaluoya, Ceylon, vii-97 [*Green*].

The unique *type* in the British Museum.

II. APPENDIX TO CATALOGUE.

(A) Additions to the Family CECIDOMYIDAE.

Subfamily CECIDOMYINAE.

Sect. LASIOPTERARIAE.

Lasioptera, Mg.

manilensis Felt, Phil. Jour. Sci., XIII, D. 288, ♀ (1918).

Laguna, Luzon, reared from leaf galls on *Leea manillensis*, Walp.

Type, Cecid. a 2851, New Yk. State colln.; *paratype* ♂ ♀ in College of Agric., Los Banos, Phil. Is.

falcata Banks, Phil. Jour. Sci. XIV, 288, ♀ (1919).

Los Banos, Laguna Prov., Luzon, 3-i-17.

Type, Cecid. a 2887, New York State collection, a unique specimen.

Sect. OLIGOTROPHARIAE.

Luzonomyia, Felt.

Phil. Jour. Sci. XIII, D. 282 (1918).

GENOTYPE *L. symphoremæ*, sp. nov.

This genus is near *Oligotrophus*, Latr.

symphoremæ Felt, Phil. Jour. Sci. XIII, D. 282, ♂ ♀ pl. i, 1, ♂ genitalia; 2, ♀ genitalia (1918).

Luzon (var. locs.); reared from leaf galls on *Symphorema luzonicum*, F. Vill.

Type, Cecid. a 2850, New Yk. State colln.; *paratype* No. 18,315 in College of Agric., Los Banos.

Sect. ASPHONDYLARIAE.

Schizomyia, Kieff.

- acalyphae** Felt, Phil. Jour. Sci. XIII, D, 286, ♀ (1918).
Laguna, reared from leaf galls on *Acalypha stipulacea*, Klotz.
Type, Cecid. a 2848, New Yk. State colln.; part of *type* material in College of Agric., Los Banos.
- diplodisci** Felt, *loc. cit.*, 287, ♂ ♀ (1918)
Mt. Maquiling, Laguna, 29-viii and 3 and 6-ix-17.
Type, Cecid. a 2849, New Yk. State colln.; *paratypes* ♂ ♀, No. 18,314, College of Agric., Los. Banos.

Asphondylia, Loew.

- vitea** Felt, Phil. Jour. Sci. XIII, D, 284, ♂ ♀ (1918).
Manila, 1905; reared from stem galls on *Cissus trifolia*, L.
Type, Cecid. a 2839, New Yk. State colln.; *cotype* ♂ ♀ in Bureau of Science, Manila; Entom. Colln. No. 3252.
- callicarpae** Felt, Phil. Jour. Sci. XIII, D, 285, ♂ ♀ (1918).
Mt. Maquiling, Laguna, Luzon, 1917. Reared from leaf galls on *Callicarpa erioclona*, Schauer.
Type, Cecid. a 2843, New Yk. State colln.; *paratype* ♂ ♀ No. 18,147, College of Agric., Los Banos.
- grewiae** Banks, Phil. Jour. Sci. XIV, 289, ♂ (1919).
Mt. Maquiling, Laguna Prov., Luzon, 23-ix-17.
Type, Cecid. a 2880, New York State collection, a unique ♂. Bred from leaf galls on *Grewia stylocarpa*, Warb.

Diceromyia, Felt.

- Phil. Jour. Sci. XIII, D, 283 (1918).
GENOTYPE *D. vernoniae*, sp. nov.
This genus is near *Zalepidota*, Rubs.
- vernoniae** Felt, *loc. cit.*, 284, ♂, pl. i, 3, ♂ genitalia (1918).
Laguna, Luzon; reared from leaf galls on *Vernonia lancifolia*, Merr.
Type, Cecid. a 2842, New Yk. State colln.

Sect. CECIDOMYIARIAE.

Arthrocnodax, Rubs.

- coprae** Banks, Phil. Jour. Sci. XIV, 291, ♂ ♀ (1919).
Los Banos, Phil. Is, 6-x-17.
Type, Cecid. a 2883, New York State collection.

Kronodiplosis, Felt.

- Phil. Jour. Sci. XIII, D, 289 (1918).
GENOTYPE, *K. uichancoi*, sp. nov.

uichancoi Felt, *loc. cit.*, 289, ♂; pl. i, 4, 3rd antennal segment ♂.
Laguna, from leaf galls on *Barringtonia luzonensis*, Rolfe.
Type, Cecid. a 2847, New Yk. State colln.; *paratype* ♂
No. 18,307, College Agric., Los Banos.

Kamptodiplosis, Felt.

Phil. Jour. Sci. XIII, D, 290 (1918).

GENOTYPE, *K. reducta*, sp. nov.

reducta Felt, *loc. cit.*, 290, ♂ ♀; pl. i, 5, 5th segment of ♂ antenna, 6, ♂ genitalia (1918).
Laguna (Balong Bulo Hill, near Los Banos); reared from leaf galls on *Siphonodon celastrineus*, Griff.
Type, Cecid. a 2852, New Yk. State colln.

Heliodiplosis, Felt.

Phil. Jour. Sci. XIII, D, 291 (1918).

GENOTYPE, *H. spatholobi*, sp. nov.

spatholobi Felt, *loc. cit.*, 291, ♀, pl. i, 7, ovipositor (1918).
Laguna; reared from leaf galls on *Spatholobus gyrocarpus* (Wall) Benth.
Type, Cecid. a 2853, New Yk. State colln.

Profeltiella, Kieff.

Neue Gallmucken Gatt. Bitsch, p. 2 (1912).

Lestodiplosis Kieff., pt., Bull. Soc. Hist. Nat. Metz.
XXVI, 25 (1909).

orientalis Felt, Phil. Jour. Sci. XIII, D, 292, ♂ ♀ (1918).
Laguna; reared from galls on *Siphonodon celastrineus*, Griff.
Type, Cecid. a 2852 a, New Yk. State colln.

Hyperdiplosis, Felt.

Bull. N. Yk. State Mus. CXXIV, 405 (1908).

Cecidomyia Felt, pt., Rep. 22, N. Yk. State Mus.
CX, 136 (1907).

banksii Felt, Phil. Jour. Sci. XIII, D, 294, ♀ (1918).
Laguna; reared from leaf galls on *Cissus adnata*, Wall.
Type, Cecid. a 2846, New Yk. State colln.

Coccodiplosis, de Meij.

Tijd. v. Ent. LX, 238 (1917).

GENOTYPE, *C. pseudococi*, *id.*, sp. nov.

pseudococci de Meij., *loc. cit.*, 239, ♂ ♀, fig. 1 a—g.
Salatiga, Java [Dr. Roepke]. Parasitic on *Pseudococcus crotonis*.
Type in Amsterdam Museum.

relicta Felt, Phil. Jour. Sci. XIII, D. 295, ♀ (1918).

Manila, 1911.

Type, Cecid. a 2841, New Yk. State colln.; *paratype* No. 16,015, College of Agric., Los Banos.

Tricontarinia, Kieff.

luzonensis Felt, Phil. Jour. Sci. XIII, D. 293, ♂ ♀ (1918).

Laguna; reared from leaf galls on *Parashorea melaanonan*, (Blanco) Merrill.

Type, Cecid. a 2844, New Yk. State colln.; *paratype* No. 18,151, College of Agric., Los Banos.

Contarinia, Rond.

Atti. Soc. Sc. Nat. Milano, II, 287 (1860).

Diplosis, Lw., pt. Dipt. Beitr. IV, 34 (1850).

Eudiplosis, Kieff., Bull. Soc. Ent. Fran. 28 (1894).

GENOTYPE, no species set up; two undescribed new species mentioned, *isorbi* and *brizae*.

saltata Banks, Phil. Jour. Sci. XIV, 289, ♂ (1919).

Los Banos, Luzon, 6-ii-18. Taken in spider's webs.

Type, Cecid. a 2884, New York State collection.

Bremia, Rond.

Atti. Soc. Sc. Nat. Milano, II, 287 (1860).

macrofilum Banks, Phil. Jour. Sci. XIV, 290, ♂ ♀ (1919).

Kolambugan, Lanao Distr., Mindanao, Phil. Is., 10-vi-14. Taken in spiders' webs.

Type, Cecid. a 2888, New York State collection.

Cecidomyia, Mg.

paederiae Banks, Phil. Jour. Sci. XIV, 293, ♂ ♀ (*Itonida*) (1919).

Los Banos, Phil. Is., 13-x-17. Bred from leaf galls on *Paederia tomentosa*, Blume. *Type*, Cecid. a 2882, New York State collection.

philippinensis Banks, Phil. Jour. Sci. XIV, 294, ♀ (1919).

Mt. Maquiling, Laguna Prov., Luzon, 3-ix-17.

Type, Cecid., a 2889, New York State collection, a unique specimen.

Ctenodactylomyia.

antidesmae Banks, Phil. Jour. Sci. XIV, 287, ♀ (1919).

Mt. Maquiling, Laguna Prov., Luzon, 2-x-17.

Type, Cecid. a 2881, New York State collection. Reared from leaf galls on *Antidesma leptocladum*, Tul., a unique specimen.

N.B.—I have been unable to place all these genera in systematic order

(B) *Errata et Addenda.*

Page 2, after line 7, add: From a short correspondence quoted by Aldrich (Can. Ent. XLIII, 34) between him and the International Commission on Zoological Nomenclature, it would appear that that Institution regards Meigen's names of 1900 as *available*, but that the question of their validity rests with dipterologists on the basis of their correct identification or otherwise. This is, of course, a flagrant violation of the principle of adopting the customs of the times in which the earlier authors wrote.

Page 9, *Microperissia*, Kieff. Add the reference: Kieff., Gen. Ins. Fasc. 152, 41.

The *type* is *brachypsectra*, Kieff., by original designation.

Page 9, line 2 from bottom: The species *ischaemi*, Kieff., was placed provisionally in *Oligotrophus* (Kieff., Roy. Bot. Gard., Kew, Bull. 3, 71, 1910), but it is now referred to *Hormomyia*: (Fletcher, Agric. Res. Inst. Pusa, Bull. 89 "2nd hundred notes on Indian Insects," p. 46, 1912). The adult is from Coimbatore, South India, reared from galls on *Ischaenum pilosum*. The identity of the species is not quite certain.

Type Cecid. a 2781.

Page 11, line 11 from bottom: replace the term *DIPLSARIAE* by *CECIDOMYIARIAE*, with which it is synonymous.

Page 12, *Chrysodiplosis*, Prof. Kieffer informs me (*in litt.*) that he designated *pulchricornis*, Kieff., as *type*.

Page 15, add the following species after 2nd line. *Dyodiplosis andropogonis*, Felt, Ent. News, XXVIII, 369—372 (Oct. 1917). *Type* Cecid. a 2782.

Page 15, for *Pseudhormomyia fluviatilis* read *fluviialis*.

Page 15, *Pseudhormomyia fluviialis* and *cornea* are now referred to *Dyodiplosis*, according to Mr. Fletcher ("2nd hundred notes on Indian Insects," p. 49) (1919).

Page 17, *Epidosis*, Loew, add the reference: Kieff., Gen. Ins. Fasc. 152, 271.

Page 17, *Epidosis indicus*. *Type* belonging to Indian Museum.

N.B.—As the species *indicus* is now referred to *Dicroncurus*, the whole matter relating to *Epidosis* must be deleted.

Page 18, *Lestremia indica*. The *type* (a ♀) in Indian Museum.

Page 32, *Leia nigricoxa*. *Types* of *nigricoxa* and *spathulata* in Indian Museum.

Page 43, *Simulium striatum*. *Type* in Indian Museum.

Page 47, *Atrichopogon flaviceps*. *Type* in Deuts. Ent. Mus., Berlin-Dahlem.

Page 48, as regards the *type* species of *Prohelia*, Prof. Kieffer writes me that no species was definitely selected but that *ornaticrus* was the first one described and he evidently regards this species as genotype. It seems to me, however, my argument (Cat. p. 48, footnote) stands good.

Page 49, *Prohelia thcobromae*. Nine *paratypes* in Indian Museum.

Page 50, *Forcipomyia sauteri*. *Type* in Deuts. Ent. Mus., Berlin-Dahlem.

Page 50, line 28: for *conifera* read *conigera*.

Page 52, *Culicoides formosae* and *C. indecora*. *Types* of both species in Deuts. Ent. Mus., Berlin-Dahlem.

Page 53, *Palpomyia tailana*. *Type* in Deuts. Ent. Mus., Berlin-Dahlem.

Page 57, *Bezzia nigricans*. *Type* in Indian Museum.

Page 58, *Bezzia formosana* and *chrysolopha*. *Types* in Deuts. Ent. Mus., Berlin-Dahlem.

Page 62, *Trichotanypus* Kieff. Prof. Kieffer (*in litt.*) apparently regards *posticalis* Lund. as genotype.

Page 62, *Protenthes bilobatus*. *Type* should be in Indian Museum, but it is apparently not there.

Page 63, *Procladius formosanus*. *Type* in Deuts. Ent. Mus., Berlin-Dahlem.

Page 66, *Cricotopus carnosus*. *Type* in Deuts. Ent. Mus., Berlin-Dahlem.

Page 66, the only species included in *Thalassomyia* Johann. (*nec* Sch.) at its erection was *obscura* Johann.

Page 68, *Tanytarsus formosanus*. *Type* in Deuts. Ent. Mus., Berlin-Dahlem.

Page 68, *Microspectra* Kieff., Bull. Soc. Hist. Nat. Metz. XXVI, 50 (1909). GENOTYPE, the only species mentioned, *inermipes* Kieff.

Page 68, *Polypedilum* Kieff., Supp. Ent. Berl. I, 41 (1912). GENOTYPE, the only species included at its erection, *pelostolum* Kieff., sp. nov., *loc. cit.* Prof. Kieffer informs me that the generic diagnosis did not appear until 1913, in the Bull. Soc. Hist. Nat. Metz. XXVIII, 15, where only a single species, *emarginatum*, was mentioned, but he adds that this paper was written in 1911 or 1912. He therefore considers that *pelostolum* must be the genotype.

Page 69, *Polypedilum fasciatipenne*, in the reference to Dr. Bains Prasad's paper on the anatomy of a larva of *Polypedilum*: for Vol. XV, read vol. XIV; also add the date (1918).

Page 69, *Polypedilum pelostolum*. *Type* in Deuts. Ent. Mus., Berlin-Dahlem.

Page 69, for *Polyptedilum brumalei*, read *brumale*.

Page 69, *Paratendipes* Kieff., Bull. Soc. Hist. Nat. Metz, XXVII, 41 (1911). GENOTYPE by original designation *Chironomus albimanus*, Mg.

Page 70, *Glyptotendipes*, Kieff. The generic characters are given in Rec. Ind. Mus. IX, 197 (1913).

Page 70, *Prochironomus*, Kieff., Bull. Soc. Hist. Nat. Metz, XXVI, 53 (1909). GENOTYPE, by original designation *P. pallidus*, Kieff.

Page 71, line 5: for *stripes* read *atripes*.

Page 79, add following details to species: *formosae*, ♂ ♀, pl. ii, 4, ♂ genitalia; *sordidus* ♀; *tainanus* ♂ ♀, pl. ii, 5, ♂ genitalia; *formosicola*¹ ♂ ♀, pl. ii, 6, ♂ genitalia; *prasinellus* ♂ ♀, pl. ii, 7, antennae (part), 8, ♂ genitalia; *chrocephorus* ♂, pl. ii, 9, ♂ genitalia; *chlorostolus* ♂ ♀, fig. 10, antennae; 11, ♂ genitalia; *hlochloris* ♂ ♀, fig. 12, ♂ genitalia, part; *glaucoventris*, ♂ ♀, fig. 13, ♂ genitalia. Figures 10 to 13 are text-figures. The types of all the above species are in the Deuts. Ent. Mus., Berlin-Dahlem.

Page 80, for *Tripelina* read *Tripelma*.

Page 80, line 12: for *conifer* read *coniger*.

Page 87, *Phlebotomus himalayensis*. Type in Indian Museum.

Page 87, add the following species to *Phlebotomus*, Rond.:
nicnic Banks, Phil. Jour. Sci., XIV, 163, ♂ ♀ (1919).

Laguna and Los Banos, Luzon [Banks].

Type ♂ ♀, No. 18,492 in Entom. colln., College of Agric., Los Banos, Phil. Is. Some paratypes exist.

Page 92, Sect. Anophelini: For diagnostic table of genera in the "Anophelina subfamily," see Giles, "A Revision of the Anophelinae" (London 1904). *Anopheles* (restr.) p. 23; *Stethomyia*, 25; *Cyclolepteron*, 26; *Myzomyia*, 27; *Pyretophorus*, 33; *Myzorhynchus*, 37; *Arribalzagia*, 40; *Chrystya* (MS. name), 40; *Nyssorhynchus*, 41; *Cellia*, 45; *Aldrichia*, 46.

Page 92, Major James, in Paludism, No. 2, pp. 52-63 (Jan. 1911) contributes a paper entitled "Notes on Mosquitoes" in which is given a key to the Indian Anophelines.

Page 92, line 12: for *grabhami* read *grabhamii*.

Page 92, to footnote 1 add: For structure of ♂ genitalia in *Anopheles*, v. Christoph., Ind. Jour. Med. Res., III, 371, with excellent figures.

Page 102, line 20: for p. 32 read 33.

Page 105, line 6 from bottom: after V, 24 add (*Myzomyia*).

¹ Spelt incorrectly *formosicola* in Catalogue.

Page 111, line 19 should read, Occurs at Lake Tiberias in Palestine and on the Beyruth River in Syria.

Page 115, after 5th line from bottom, add: Christophers, in Paludism, No. 3, pp. 40—54, offers an interesting paper, "Aids to the identification of Culicidae other than *Anopheles*, with special reference to Indian species," July 1911.

Page 119, *Toxorhynchites minimus*, Theob., add the reference: White, Spol. Zeyl. XI, 189, ♀, described (1919).

Page 122, line 22: *malayi* is a good species, described as a *Uranotaenia*. *Armigeres malayi*, Edw.; and *Desvoidia fusca*, Theob., are identical. Delete the Kuala Lumpur locality as Edwards says the only specimens from this locality referred by Theobald to this species are *obturbans*.

Page 124, line 6 from bottom: for p. 285 read 283.

Page 125, add as a further generic synonym after *Scutomyia*: *Hulecoctomyia*, Theob., Entom. XXXVII, 163 (1904): Monog. Culic., IV, 219 (1907). GENOTYPE, *H. trilineata* Leices. in Theob., sp. nov., loc. cit., by my own designation. (Rec. Ind. Mus. X, 62, 1914).

Page 128, add to *albolineata*, Giles, the following references: Giles, Handbk., 2nd Ed., 430, ♀, pl. xvii, 10a, venation ♀ (1902): Theob., Monog. Culic. III, 192, ♀: Brun., Rec. Ind. Mus. IV, 452. Type in British Museum.

Page 131, add: *Oculeomyia*, Theob. pt. as a synonym of *Grabhamia*.

Page 134, delete last line beginning *Neomacleaya*.

Page 135, delete first two lines.

Page 141, the date of the species *Ochlerotatus orcophilus* is 1916, and of *koreicus*, 1917.

Page 145, the date of the species *ceylonicus*, *yerburyi* and *leicesteri* is 1917.

Page 152, the date of the species *Culiciomyia spathifurca* is 1915.

Page 154, the date of the species *Lophoceratomyia curtipalpis* is 1914.

Page 180, line 4 from bottom should read: GENOTYPE, *Wycomyia aranoides*, Theob., etc.

Page 180, *Topomyia tipuliformis*. In text, for *argyroventralis* and *argentoventralis*, read *argenteoventralis*.

Page 185, *Corethrella inepta* Annand. For notes on habitat see A. J. MacDougall, Spol. Zeyl. VIII, 71 (*Ramcia inepta*) (1918).

Page 185, *Dixa montana* Brun. For full description of larva and pupa see B. Prashad, Rec. Ind. Mus. XV, 156, pl. xix, var. chars. (1918).

Page 187, *Tanyderus mirabilis*, add the closer locality: Kloof-Biwak, 29-X-12 [Versteg]. Unique ♀ type belongs to the Buitenzorg Museum.

Page 187, in the footnote respecting the type species of *Ctenophora* I overlooked having myself suggested *pectinicornis* L. as genotype (Fauna Brit. India, Errata). At that time I had not studied the point closely, and in any case the suggested species would be invalid as Rondani's designation of *flaveolata* F. stands good.

Page 193, delete the species *clongata*, Edw., with full data. It is already recorded under *Pselliophora*, p. 192.

Page 204, after the species *robinsoni*, add the following:—
pseudofulvipennis de Meij., Bijd. Dierk., Feest-nummer, 18, ♂ ♀ (1919).

Korinchi Peak, Sumatra, viii.

Types in Amsterdam Museum. Dr. Meijere has also seen the species from Java, and it is very near the European *fulvipennis*.

Page 204, before the genus *Longurio*, insert the following new genus and species.

Sphaerionotus, de Meij. gen. nov.

Bijd. Dierk., Feest-nummer, 17 (1919).

GENOTYPE, *S. curtipennis*, id., sp. nov., loc. cit.

curtipennis de Meij., loc. cit. 18, pl. ii, 8 wing (1919).

Murao Sako, Sumatra, x. Unique ♂ type in Amsterdam Museum. The genus is allied to *Tipula*.

Page 209, add the following species of *Pachyrhina*:—

nigrithorax de Meij., Bijd. Dierk., Feest-nummer, 18, ♀ (1919).

Air Njuruk, Dempu, Sumatra, 1400 metres, viii.

The unique ♀ type in Amsterdam Museum.

Page 212, add the following new genus and species to *Cylindrotomini*:—

Agastomyia, de Meij., gen. nov.

Bijd. Dierk., Feest-nummer, 17 (1919).

GENOTYPE, *A. albitarsis*, id., loc. cit., sp. nov.

albitarsis de Meij., loc. cit. 17, pl. ii, 7, wing (1919).

Sungai Kumbang, Sumatra, ix.

Unique type ♂ in Amsterdam Museum.

Page 217, add following species to the genus *Dicranomyia*:—

signata de Meij., Bijd. Dierk., Feest-nummer, 13, ♂, pl. ii, 1, wing, 1919.

Korinchi Peak, Sumatra, viii. Unique ♂.

Type in Amsterdam Museum.

Page 222, add following species of *Limnobia* :—

stigmosa de Meij., Bijd. Dierk., Feest-nummer, 14, pl. ii, 2, wing (1919).

Suban Ajam, Sumatra, vii. Type in Amsterdam Museum, sex not stated.

Page 224, *Dapanoptera versteegi* de Meij. Add following data: fig. 2, wing. Dutch South Papua, 28-ix-12, one ♀.

Type belonging to Buitenzorg Museum.

Page 225, *Libnotes strigivena* Walk. Add new locality, Biwak Island, Papua, ix, one ♂.

Page 225, *Libnotes quadriurca* Walk. Add new locality, Beaufort-Fluss, Papua, xi, one ♂.

Page 228, add following species of *Libnotes* :—

tripunctata de Meij., Bijd. Dierk., Feest-nummer, 14, pl. ii, 3, wing (1919).

Suban Ajam, Sumatra, vii.

Unique ♀ type in Amsterdam Museum.

Page 239, add following species of *Acyphona* :—

punctulata de Meij., Bijd. Dierk., Feest-nummer, 14, pl. ii, 4, wing (1919).

Sungei Kumbang, Sumatra, ix.

Unique ♀ type in Amsterdam Museum.

Page 246, add following species of *Gnophomyia* :—

fascipennis de Meij., Bijd. Dierk., Feest-nummer, 15, pl. ii, 5, wing (1919).

Muara Sako, Sumatra, ix.

Unique ♂ type in Amsterdam Museum.

Page 242, add following species of *Mongoma* :—

nigriceps de Meij., Bijd. Dierk., Feest-nummer, 15 (1919).

Suban Ajam, Sumatra, vii.

Unique type ♀ and a damaged specimen in Amsterdam Museum.

Page 256, add following species of *Epiphragma*.

kerberti de Meij., Bijd. Dierk., Feest-nummer, 16, pl. ii, 6, wing (1919).

Air Njuruk, Dempu, Sumatra, 1400 metres, viii; Suban Ajam, Sumatra, vii. Type in Amsterdam Museum, sex not stated.

Page 268, *Diceromyia* Felt. This name is preoccupied in Culicidae by Theobald, 1911.

	Page		Page
alticola Edw. ..	216	Anomalomyia Hutton ..	30
" Kieff. ..	50	anomalous Kieff. ..	65
Amalopina Brun. ..	251	Anopheles Mg. ..	92, 273, 273
AMALOPINI (Amalopina) ..	249	ANOPHELINA ..	80
Amalopsis Hal. in Walk. ..	250	ANOPHELINAE ..	80
Amaurosiphon Rubs. ..	15	ANOPHELINES ..	89
amazonensis Lutz. ..	156	ANOPHELINI ..	80, 92, 273
ambigua Theob. ..	132	anopheloides Giles ..	176
ambiguus Mg. ..	46, 49	Anoplistes Westw. ..	257
amboinensis Dol. ..	116, 117, 118	antennatus Brun. ..	223
amesii Ludl. ..	128	anthracina de Meij. ..	28
amica Alex. ..	254	antica Brun. ..	240
analis Kieff. ..	56	antidesmae Banks ..	270
Anchimongoma Brun. ..	244	antisepticus Lichtenstein ..	184
Andersonia Strickland ..	135	Antocha Ost. Sack. ..	232
Androclosma Ender. ..	258	anxifer Coq. ..	158
Androdiplosis Felt ..	16	Apeilesis Macq. ..	210
Andropogonis Felt ..	271	aperta Brun. (<i>Gonomyia</i>) ..	241
angulatus Theob. ..	151	" " (<i>Rhaphidolabis</i>). ..	252
angusticapitis Brun. ..	233	apicalis Alex. ..	228
angustiforceps Kieff. ..	69	" Brun. ..	82
angustipennis de Meij. ..	86	" de Meij. (<i>Empheria</i>) ..	27
" Ender. ..	263	" " (<i>Linnophila</i>) ..	254
" Winn. ..	17	" Theob. (<i>Armigeres</i>) ..	123
Anisocheleomyia Theob. ..	173	" " (<i>Brevirhynchus</i>) ..	146
Anisomerini ..	258	" " (<i>Leicesteria</i>) ..	146, 147
Anisopus Mg. ..	265	" W. ..	212, 220
annandalei Brun. (<i>Isoneuromyia</i> , <i>Platyura</i>) ..	25, 26	apicipennis Brun. ..	25
" " (<i>Pericoma</i>) ..	83	apicotriangulata Theob. ..	157
" " (<i>Ptychoptera</i>) ..	186	Apistomyia Big. ..	36
" Kieff. (<i>Isoplastus</i>) ..	63	aplochirus Kieff. ..	76
" " (<i>Prochironomus</i>) ..	71	aploneurus ..	76
" Prashad ..	114	Apogon Rond. ..	46
" Theob. ..	130	Aporoculex Theob. ..	156
annularis Schrk. ..	169	Aporosa Macq. ..	219
" Theob. (<i>Culicomyia</i>) ..	152	appendiculata Brun. ..	82
" " (<i>Trichopronomyia</i>) ..	156	approximata Brun. (<i>Dicranomyia</i>) ..	216
" Theob. nec Wulp (<i>Anopheles</i>) ..	96	" " (<i>Pselliophora</i>) ..	192
" Wulp. nec Theob (<i>Anopheles</i>) ..	96, 97, 98	approximatus ..	40
annulatus Leices. ..	149	apricus Kieff. ..	15
" Schrk. ..	169	aranetana Banks ..	140
" Theob. ..	165	aranoides Theob. (<i>Wyeomyia</i>) ..	181, 274
annulicornis Brun. ..	33	arcuata Brun. ..	32
annulifemur Edw. ..	141	arcuatipes Kieff. ..	57
" de Meij. ..	221	ardens W. ..	188
annulifera Theob. ..	150, 151	arenariae Rubs. ..	14
annuliferus Edw. ..	150	argentea Ludl. Brun. (<i>Taeniorhynchus</i> , <i>Theobaldiomyia</i>) ..	165
" Ludl. ..	166	argenteiventris Theob. ..	181
annulioris Theob. ..	163	argenteomaculata Theob. ..	130
annulipalpis Theob. ..	147	argenteopunctata Brun. ..	84
annulipes Brun. ..	254	argenteosquamosa Ender. ..	30
" Ender. ..	245	argenteotarsis Ludl. ..	117
" Walk. ..	148	argenteoumbrosus Theob. ..	156
annulirostris Skuse ..	159	argenteoventralis Leices. ..	179, 180, 274
" Theob. ..	129	argenteus Ludl. ..	165
annulitarsis Leices. ..	146	argentifera de Meij. ..	217
" Macq. ..	125	argentinotus Banks ..	167
annuloabdominalis Theob. ..	152	argentipes Annand. and Brun. ..	86
annulosa Wulp ..	190	argentocincta Walk. ..	231
annulus Theob. ..	162	argentoventralis Leices. (<i>lapsus</i> , <i>mihi for argenteoventralis</i>) ..	180
Anomala Marsh. nec Steph. ..	31	argyrocinctum de Meij. ..	43
anomalipes Kieff. ..	79	argyropalpis Leices. ..	180
		argyropus Swellen. ..	96
		argyrotarsis Leices. ..	175
		" R. Desv. ..	93

	Page		Page
argyroventralis Leices. (lapsus, mihli for argenteoventralis) ..	180	auronitens Kieff. ..	47
armatifrons Kieff. ..	80	aurostriata Banks ..	130
armatipes Kieff. ..	57	australiensis Theob. in Giles ..	150
Armigeres Theob. ..	121	azriki Patton ..	103
Arrhenica Ost. Sack. ..	258	babu Annaud. ..	86
Arribalzagia Theob. ..	93, 273	badia Brun. ..	262
artemisiarum Kieff. ..	16	bahri Edw. ..	152
Arthrocnodax Rubs. ..	12, 268	bancrofti Skuse ..	126
articulatus Rond. ..	135, 136	Bancroftia Lutz. ..	176
ascidiicola de Meij ..	176	banksii ..	269
asiatica Giles ..	184	Banksinella Theob. ..	124, 142
.. Leices. ..	113	barbatitarsis Kieff. ..	75
.. Theob. ..	94	barberi Coq. ..	94, 96
ASPHONDYLARIAE ..	10	barbicornis F. ..	46
Asphondylia Lw. ..	10, 268	barbirostris Wulp ..	97
assamensis Brun. (<i>Molophilus</i>) ..	236	barianensis James and List. ..	96
.. .. (<i>Teucholabis ornata</i> , var.) ..	233	barkeri Theob. ..	153, 154
.. Theob. ..	130	basalis Brun. ..	27
Asthenia Westw. ..	36	basilaris ..	259
Atarba Os. Sac. ..	235	basilinea Brun. ..	34
ater Daniels ..	119	Bathosomyia Theob. ..	135
aterrima Brun. ..	263	bella Brun. ..	83
.. Walk. ..	222	.. Lw. ..	36
aterrimus Kieff. (<i>Atrichopogon</i>) ..	48	bengalensis Brun. (<i>Lechvia</i>) ..	244
.. .. (<i>Clinotanypus</i>) ..	61 (<i>Psychoda</i>) ..	81
atra Brun. ..	39	.. Kieff. (<i>Bezzia</i>) ..	56
.. Dol. ..	259 (<i>Peromyia</i>) ..	17
.. Macq. ..	38	Bezzia Kieff. ..	56
.. Mg. ..	235	biannulata Brun. ..	233
.. Theob. ..	174	Biblio Geoff. ..	39
Atractocera Mg. ..	42	BIBIONIDAE ..	37
atrata L. ..	187	BIBIONINAE ..	37
atratum de Meij. ..	43	bibula W. ..	220
atratus Kieff. (<i>Clinotanypus, Procladius</i>) ..	60, 61	bicinetipes Brun. ..	212, 215
atratus Theob. ..	155	bicolor Brun. (<i>Leia</i>) ..	32
atrescens Alex. ..	216 (<i>Dicranomyia</i>) ..	217
atricauda Skuse ..	31 (<i>Sciophila</i>) ..	26
Atrichopogon Kieff. ..	47	.. de Meij. ..	236
atripes Kieff. (stripes, lapsus) ..	71	.. Leices. ..	175
atrisquamis Brun. ..	84	.. Macq. ..	259
atritarsis Brun. ..	186	.. Ost. Sack. ..	233
atrophus Kieff. ..	80	.. Walk. ..	40
atrosignatus Kieff. ..	77	bicornuta Theob. ..	154
Atypophthalmus Brun. ..	223	bicornutus Kieff. ..	68
aurantia Brun. (<i>Aporosa</i>) ..	220	bifasciata Brun. ..	185
.. .. (<i>Eriocera</i>) ..	264	bifascipennis Brun. (<i>Mycomyia</i>) ..	27
.. .. (<i>Pselliophora</i>) ..	192 (<i>Pselliophora</i>) ..	191
.. Theob. ..	120, 121	bifurcatus L. ..	96
aurantiaca Dol. ..	225	bigotii Bell. ..	155
auratus Leice. ..	136, 166	bilineata Gmel. ..	269
aurea ..	172	bilobata Lw. ..	36
aureochaetae Leices. ..	184	bilobatus Kieff. (<i>Culicoides</i>) ..	52
aureohirtum Brun. ..	42 (<i>Procladius</i>) ..	63
aureolineatus Leices. ..	123 (<i>Tanypus, Trichotanypus, Protenthes</i>) ..	62, 272
aureopunctis Ludl. ..	167	bimacula Kieff. ..	54
aureosquamatus Ludl. ..	149	bimaculata Leices. ..	175
aureostriata Leices. ..	121	.. Walk. ..	32
aureostriatus Dol. ..	160	bimaculiala Leices. ..	174
aureoventer Theob. ..	179	bimaculipes Theob. ..	183
aurifer Coq. ..	134	binotata Brun. ..	35
auripennis Skuse ..	212	.. Edw. ..	186
aurirostris Watson ..	114	Binotia Blanch. ..	181
aurites Ludl. nec Theob. ..	149	bioculata de Meij ..	223
.. Theob. nec Ludl. ..	138	bionis Agharkar ..	36
auroatra Walk. ..	224	bipunctata Brun. ..	222
		bipunctata L. ..	46

	Page		Page
bipunctata Theob. ..	163	bullans Ender. ..	211
bipunctatus I. Kieff. (<i>Chironomus</i>) ..	72	burmae Kieff. ..	53
.. II. .. (<i>Chironomus</i>) ..	80	butleri Theob. ..	142, 143
birmanensis Kieff. ..	60	byssina Schrk. ..	64
biroi Theob. ..	162		
Bironella Theob. ..	115	Cacomyia Coq. ..	135
bisignata Kieff. ..	54	caeca Theob. (<i>Pecomyia</i>) ..	137
bispinosa Petzey. ..	22	caecus Theob. (<i>Culex</i>) ..	135, 137
bistriata Brun. ..	185	Caenarthria Thoms. ..	257
bistriatipennis Brun. ..	234	caeruleocephala Leices. nec	
bisulcatus Coq. ..	168	Theob. ..	181-182
bitaeniorhynchus, Giles ..	132	caeruleocephala Theob. nec Leices.	
Blanchardiomyia Brun. ..	121	(<i>Phoniomyia</i>) ..	182
Blepharocera Macq. (<i>Blepharocera</i>) ..	36	caeruleocephala Theob. (<i>Uranolaenia</i>) ..	173
BLEPHAROCERIDAE ..	36	caeruleocephala Theob. var. <i>lateralis</i> Ludl. ..	174
Bolbodeomyia Theob. ..	183	caeruleovittata Ludl. ..	178
Boletina Staeg. ..	29	calcarata Kieff. ..	49
bolwiensis Theob. ..	94	calcuttensis Kieff. (<i>Bezzia</i>) ..	58
bombayensis Macq. ..	205	.. (<i>Dasyhelia</i>) ..	46
Bophrosia Rond. ..	250	caliginosa Brun. ..	75
borneensis Brun. (<i>Ovinarga</i>) ..	231	callicarpae Felt ..	268
.. (<i>Tipula</i>) ..	195	callichirus Kieff. ..	75
Boycia Newstead ..	171	callicomus ..	74
Brachycampa Winn. ..	33	calligaster ..	76
Brachydierania Skuse ..	34	callimorphus Kieff. ..	77
BRACHYNEURARIAE ..	11	callinotus Felt (<i>Chironomus</i>) ..	75
brachypsectra Kieff. ..	271	.. (<i>Metricnemus</i>) ..	64
brahmacharii Chris. ..	106	callisphyrus Felt. ..	76
brakeleyi Coq. ..	185	callithorax ..	73
brasiliensis Westw. ..	220	Caloptera Guer. ..	258
breinli Taylor ..	124	caloptera Say ..	239
Bremia Rond. ..	270	calopus Mg ..	125
brevicellulus Theob. ..	148, 149	calotricha Kieff. ..	49
bevicornis Kieff. ..	66	Calvertia Ludl. ..	115
breviforceps Kieff. ..	75	Calvertina ..	115
brevifrons Brun. ..	230	Calyptopogon Kieff. ..	44
brevimannus Kieff. ..	51	campestris Leices. ..	175
brevior Brun. ..	230, 240	Camptocladus Wulp. ..	64
brevipalpis Giles ..	168, 169	camptogaster Kieff. ..	72
.. Roper ..	95, 114	canadensis Theob. ..	133
.. Theob. (<i>Lophoceraomyia</i>) ..	153	.. Westw. ..	230
.. (<i>Toxorhynchites</i>) ..	116	canariensis Berg. ..	220
brevipennis Curt. ..	236	cancer Leices ..	175
brevirostris Brun. ..	29	cantans Mg ..	157
.. Edw. ..	176	carbonaria Brun. ..	39
Brevirhynchus Theob. ..	145	cardamitis Newst. and Carter ..	111
brevis Brun. ..	204	Cardiocladius Kieff. ..	66
.. Harris ..	265	caricis Rubs. ..	15
brevispina Kieff. ..	53	cariniceps Ender. ..	242
brevistila ..	58	carmichaeli Brun. ..	195
brevistilus ..	48	carnea Lw. ..	17
brevitibia Edw. ..	123	carneotincta Alex. ..	217
breviventris Kieff. ..	67	carnosus Kieff. ..	66, 272
Brithura Edw. ..	204	Carrollia Lutz in Theob. ..	155
brumale Kieff. ..	273	castanea Macq. ..	193, 194
brumalei Kieff. (=brumale) ..	69-273	Catageomyia Theob. ..	142
Brunettia Annand. ..	84	catastieta Knab. ..	177
brunettii Edw. (<i>Delopsis</i>) ..	35	Catocha Hal. ..	18
.. (<i>Rhaphidolabis</i>) ..	252	caudelli Dyar and Kirab. ..	156
.. Kieff. (<i>Chironomus</i>) ..	79	caunteri Kieff. ..	70
.. (<i>Cricotopus</i>) ..	66	Cecidogona Lw. ..	17
.. (<i>Tanypus</i>) ..	60	Cecidomyia Felt ..	269
brunnea Brun. ..	24	Cecidomyia Mg. ..	15, 270
brunnescens Brun. ..	37	CECIDOMYIARIAE ..	268, 271
.. de Meij ..	219	CECIDOMYIDAE ..	8, 267
brunnicosta Brun. ..	201	CECIDOMYINAE ..	8, 267
bryanti Alex. ..	241		
bryoniae Sch. ..	6		

	Page		Page
Cellia Theob. ..	93, 273	cinctipes Fab. (<i>Tipula</i>) ..	198
cellularis Kieff. ..	16	cinctiventris Brun. (<i>Limnobia</i>) ..	221
centauriae Kieff. ..	15 (<i>Mycetophila</i>) ..	35
centralis Brun. ..	221	cinctoterminalis Brun. ..	204
Ceratocheilus Wesche ..	230	cinerascens Brun. ..	215
Ceratoecystia Dyar and Knab. ..	131	cinerea Brun. ..	199
Ceratolophus Kieff. ..	55	.. de Meij. ..	198
Ceratopogon Mg. ..	43, 46	.. Kieff. (<i>Palpomyia</i>) ..	53
CERATOPOGONINAE ..	44	.. Macq. (<i>Apeilexis</i>) ..	210
Cerastephanus Brun. ..	223 (<i>Lestramia</i>) ..	17
CEROPLATINAE ..	24	cinereifrons de Meij. ..	198
Ceroplatus Bosc. ..	24	cinereus Mg. ..	141
Cerotelion Rond. ..	25	cingulata Brun. ..	264
ceylanica Felt ..	13	.. de Meij. ..	262
.. Kieff. (<i>Clinodiplosis</i>) ..	16	.. Leices. ..	146
.. .. (<i>Dibezzia</i>) ..	58	cingulatus Dol. ..	168
.. .. (<i>Lestremia</i>) ..	18	circipunctata Brun. ..	218
.. .. (<i>Pachydiplosis</i>) ..	16	citrina Edw. ..	208
ceylanicus .. (<i>Cardiocladius</i>) ..	66	Cladura Ost. Sack. ..	248
.. .. (<i>Chironomus</i>) ..	73	Claduroides Brun. ..	251
.. .. (<i>Culicoides</i>) ..	52	claripennis ..	253
.. .. (<i>Dactylocladius</i>) ..	66	Clastobasis Skuse ..	31
.. .. (<i>Epidosis</i>) ..	17	clavata Kieff. ..	58
.. .. (<i>Lestodiplosis</i>) ..	14	clavatipes Kieff. ..	71
.. .. (<i>Misocosmus</i>) ..	17	clavatitarsis Kieff. ..	67
.. .. (<i>Tanytarsus</i>) ..	68	clavipalpus Theob. ..	171, 172
ceylonensis Theob. ..	181	cleopatra Willcocks ..	103
ceylonica Edw. ..	245	Climacura How. Dyar and Knab. ..	157
.. Newst. and Carter ..	95-105	Clinodiplosis Kieff. ..	13, 16
.. Theob. (<i>Culicomyia</i>) ..	152	Clinotanytus ..	60
.. .. (<i>Uranotaenia</i>) ..	174	Cloniophora Sch. ..	257
ceylonicus Edw. ..	145	Clydonodozos Ender. ..	247
Chaetomyia Leices. ..	145, 146	CLUNIONARIAE ..	63
Chagasia Ludl. ..	115	Cnephoeophila Phil. ..	27
chamberlaini Ludl. ..	171, 172	coccidivora Felt (<i>Androdiplosis</i>) ..	14
Chaoborus Licht. ..	184 (<i>Diadiplosis</i>) ..	16
chaudoyei Theob. ..	193	Coccodiplosis de Meij. ..	260
Chemalida Rond. ..	237	Coelodiazesis Dyar and Knab. ..	94, 95
Chiasmoneura de Meij. ..	28	Coelodiplosis Kieff. ..	12
CHIRONOMARIAE ..	66	Coelosia Winn. ..	34
CHIRONOMIDAE ..	43	cognata Ost. Sack. ..	267
CHIRONOMINAE ..	63	cohaesa Don. ..	106, 109
Chironomus Mg. ..	71	collaris Brun. ..	35
chlorogaster Kieff. I (<i>Chironomus</i>) ..	77	.. Ender. ..	35
.. .. II ..	78	Colonemyia Leices. ..	181
chlorophorus ..	79, 273	colorata Walk. ..	206
chlorostolus ..	79, 273	Colpodia Winn. ..	17
choricus ..	77	columbaeensis Schoub. ..	42
Christophersia James ..	95	columbina Brun. ..	215
.. Theob. (<i>Anopheles</i> , ..		combinata Walk. ..	260
.. .. <i>Myzomyia</i>) ..	105	commovens ..	120
.. Theob. (<i>Culex</i>) ..	159	communis Leices. ..	184
Christya Theob. (<i>Chrystya</i>) ..	93-273	.. Mg. ..	46
Chrysoconops Goeldi ..	134	compacta Brun. ..	21
Chrysodiplosis Kieff. ..	11, 271	compedita W. ..	189
chrysozona Knab. ..	149	completiva Leices. ..	172
chrysolineata Theob. ..	129, 131	complex Theob. ..	183, 184
chrysolopha Kieff. ..	58, 272	complexa Ost. Sack. ..	232
chrysophila Walk. ..	189	Conchyliastes Theob. ..	92-95
chrysosecula Theob. ..	140	concolor Bech. ..	34
chrysothorax ..	124	.. R. Desv. ..	156, 157, 161
Chrystya v. Christya ..	93	concolorithorax Brun. ..	208
cilicrus Kieff. ..	16	confinis Brun. ..	222
cincta Brun. ..	264	confinis Arrib. ..	148
.. de Meij. ..	30	confirmatus Arrib. ..	133
.. Fab. ..	59, 62	confluens Kieff. ..	59
cinctipes .. (<i>Mesochria</i>) ..	29	confundendus Kieff. ..	67
		confusus Edw. ..	123

	Page		Page
confusus Theob. ..	135	Culicelsa Felt. ..	133
congruens Walk. ..	193	CULICIDAE ..	88
conifrons Edw. ..	204, 205	culicifacies Giles ♂ ..	103
coniger (conifer) Kieff. ..	80, 273	" " ♀ ..	101
conigera Kieff. ..	50	culiciformis Cogill ..	112
coniophaga Winn. ..	16	" James and Liston ..	113
Conithorax Brun. ..	230	" Theob. ..	143
conjungens Edw. ..	123	CULICINA ..	90
" Kieff. ..	74	CULICINAE ..	88, 89
Conocladus Kieff. ..	64	CULICINES ..	90
conopas Frnld. nec Theob. ..	136	CULICINI ..	89, 115
conopas Theob. nec Frnld. ..	149	Culiciomyia Theob. ..	151
Conopomyia Leices. ..	171	Culicoides Latr. ..	43, 48, 51
Conosia Wulp ..	248	Culiseta Felt ..	169
consimilis Brun. ..	207	cuneatus Theob. ..	163
contaminata L. ..	186	cuneiformis de Meij ..	212, 213
Contarinia Rond. ..	270	curtimanus Kieff. ..	71
contigua Brun. ..	203	curtipalpis Edw. ..	154
contingens Walk. ..	253	curtipennis de Meij. ..	275
continuata Brun. ..	201	curtipes Edw. ..	145
convergens de Meij. ..	213	curtitarsis Kieff. ..	79
coprae Banks ..	268	curvilinea Brun. (<i>Mycetophila</i>) ..	35
Coprodiplosis Kieff. ..	14	" " (<i>Mycomyia</i>) ..	27
Coquillettia Dyar ..	148	curvipes Wulp. ..	190
coquillettii Ender. ..	202	cyanea Edw. ..	234
Corethrella Coq. ..	185	cyaneomaculatus Dol. ..	50
CORETHRINAE ..	184	Cyathomyia de Meij. ..	168
cornea Felt ..	15, 171	Cyclolepidopteron. Blanch. ..	92
cornfordi Theob. ..	185	Cyclolepteron Theob. ..	92-273
cornigera Alex. ..	219	Cylindrocera Lioy ..	11
" Speis. ..	230	Cylindrotoma Macq. ..	211, 212
costalis Brun. ..	211	CYLINDROTOMINI ..	211
" Edw. ..	236	Cyniphes G. Costa. ..	85
" Lw. ..	92		
" W. ..	220	Dactylocladius. Kieff. ..	66
costata Felt ..	15	Dactylomyia. Newst. and Cart. ..	95
Courteia Kieff. ..	12	Danielsia, Theob. ..	133
Crapitula Gimm. ..	37	Dapanoptera, Ost. Sack in Westw. ..	224
crassa Edw. ..	205	Daphnephila, Kieff. ..	11
crassicornis de Meij. ..	37	Dasyhelia, Kieff. ..	45
crassicosta Speis. ..	245	Dasymallomyia, Brun. ..	246
crassinervis de Meij. ..	52	Dasymyia Leices. ..	171
crassipes Bezzi ..	264	Dasyneura Rond. ..	19
" Theob. nec Wulp ..	123	Dasyptera Sch. ..	236
" Wulp nec Theob. ..	127	de beauforti, de Meij. ..	216
crassiuscula de Meij. ..	37	decemguttata de Meij ..	219
Cricotopus Wulp ..	66	deceptor Don ..	105
cristatum ..	69	decepiens Kieff. ..	48
crocata L. ..	205	decora Brun. ..	82
crocea Edw. ..	222	decorabilis Leices. ..	179
Crunobia Kolen. ..	251	decorata Brun. ..	265
erux Dol. ..	248	defectus Brun ..	41
" W. ..	62	deferenda Walk. ..	15
Cryptodiplosis Kieff. ..	15	DEINOCERATINAE ..	91
crystalloptera Ost. Sack. ..	261	DEINOCERATINES ..	91
Ctenacroscelis Ender. ..	193	delectata Walk. ..	231
Ctenoceria Rond. ..	186	delicata Brun ..	215
Ctenodaetylomyia ..	270	Delopsis Skuse ..	35
Ctenophora Mg. ..	187, 275	delta Walk. ..	205
ctenophorina, Riedel ..	191	demarcata Brun. (<i>Dicranomyia</i>) ..	214
CTENOPHORINI ..	187	" " (<i>Pachyrhina</i>) ..	208
ctenophoroides Edwards ..	262	" " (<i>Tipula</i>) ..	199
cubicularis Kieff. ..	48	de meijeri ..	32
cubiculorum Dol. ..	72	" " (<i>Tipula</i>) ..	202
Culex L. ..	155	" Edw. ..	254
Culicada Felt ..	133	Dendromyia Theob. ..	183
CULICALES ..	89	DENDROMYINAE ..	91
Culicella Felt ..	155	deniedmanni Ludl. ..	133

	Page		Page
dentata de Meij. ..	202	dolichotomus Kieff. ..	75
Dentifibula Felt ..	13	dolicocephala Leices. ..	146
desmotes Giles ..	128	dolosa Arreb. ..	142, 158
Desvoidya Blanch. (<i>Desvoidia</i> , <i>Desvoidia</i>) ..	121	domestica Theob. ..	133
determinata Ost. Sack. ..	233	domesticus Germ. ..	132
Diadiplosis Felt ..	14	.. Leices. ..	132
DIADOCIDIINAE ..	28	dorsalis de Meij. ..	207
diagraphis Kieff. ..	69	.. Walk. ..	38
diana Macq. ..	259	dorsopunctata Brun. ..	207
Dibezzia Kieff. ..	58	dotata Ost. Sack. ..	269
Diceromyia Felt ..	268, 276	dubiosa Brun. ..	28
Diceromyia Theob. ..	157	dubitans Leices. ..	180
dichroa Walk. ..	260	dudgeoni Theob. ..	112
Dicranomyia Steph. ..	212	durhami Edw. ..	124
Dicranophragma Ost. Sack. ..	255	Duttonia Newstead ..	135
Dicranoptycha Ost. Sack. ..	231	dyari Coq. ..	155
Dicroneurus Kieff. ..	16	Dyodiplosis Rubs. ..	14
Dictenidia Brullé ..	187	Ecculex Felt ..	133
didyma Grimshaw ..	245	egregia de Meij. ..	231
diffusa de Meij. ..	235	Ekrimomyia Leices. ..	121
digitata Edw. ..	147	Elacophila Rond. ..	256
Dilophus Mg. ..	41	elatus Kieff. ..	78
diluta Walk. ..	260	elegans Big. ..	36
dimidiata de Meij. ..	207	.. Brun. (<i>Amalopsis</i>) ..	251
Diomyza Meg. in Mg. ..	8 (<i>Macrocera</i>) ..	24
Diplecus, Kieff. ..	14 (<i>Paracladura</i>) ..	249
diplosisci Felt ..	268 (<i>Tipula</i>) ..	199
Diplonema Annand nec Lw. ..	84	.. Ficalbi ..	126
DIPLOSARIAE ..	11, 271	.. James in Theob. ..	95
Diplosis, Lw. ..	15, 270	elegantula Brun. (<i>Amalopina</i>) ..	251
discalis Brun. ..	40 (<i>Tipula</i>) ..	202
discolor Coq. ..	131	Elephantomyia Ost. Sack. ..	220, 230
discolor de Meij. ..	52	Eloeophila Rond. ..	256
disjuncta Kieff. ..	54	elongata Edw. ..	193-275
dispar Kieff. ..	11	elongatissima Brun. ..	263
dissimilis Leices. ..	130	eminentia Leices. ..	153
distans Brun. (<i>Erioptera</i>) ..	237	Empeda Ost. Sack. ..	239
.. .. (<i>Psychoda</i>) ..	81	Empheria Ost. Sack. ..	28
disticta Kieff. ..	53	Empheria Winn. ..	28
distincta Brun. (<i>Psychoda</i>) ..	81	Endaphis Felt ..	12
.. .. (<i>Ptychoptera</i>) ..	186	Ephelia Sch. ..	256
.. Meun. ..	29	EPIALURGI Alcock ..	89
distinctus Brun. ..	266	epidesmus Theob. ..	133
distinguenda Brun. ..	20	Epidosis Lw. ..	17, 271
ditaemius Kieff. ..	70	Epiphragma Ost. Sack. ..	255
diurna Theob. ..	143	Eriocera Macq. ..	258
divergens de Meij. ..	202	Erioptera Sch. = <i>Molophilus</i> Curt. ..	236
diversipes Brun. ..	20	.. Mg. nec Sch. ..	237
dives Brun. ..	195	ERIOPTERINI (<i>Erioptera</i> , <i>Erioptera</i> inae) ..	235
dives Sch. ..	148	error Theob. ..	93, 112
divisa Brun. (<i>Pselliophora</i>) ..	191	eruthrosops Theob. ..	135
.. .. (<i>Tipula</i>) ..	200	erythrina Alex. ..	216
divisus ..	266	erythropterus Kieff. ..	80
Dixa ..	185	Etorleptiomyia Theob. ..	170
DIXIDAE ..	185	eucera Kieff. ..	56, 57
dofleini Guenther ..	173	euchroma Walk. ..	255
dohrnianus Ender. ..	193, 195	Eudiplosis Kieff. ..	270
dolens Kieff. ..	70	Eumelanomyia Theob. ..	157
.. Ost. Sack. ..	190	Eumerozema Kieff. ..	11
doleschalli Giles ..	168	Eupeitenus Macq. ..	38
.. Ost. Sack. ..	205	Eurhamphidia Alex. ..	229
dolichocephala Leices. ..	146	Euryschalis Brun. ..	34
Dolichocladus Kieff. ..	65	evanescens Brun. ..	22
dolichogaster ..	74	Evanioptera Guer. ..	258
dolichomerus ..	76	exacta Brun. ..	20
Dolichozepea Curt. ..	210	exagitans Walk. ..	125
DOLICHOPEZINI ..	269		

	Page		Page
excelsa Alex.	216	filiformis Walk.	231
excilians Walk.	125	filipes F.	209
exclusa "	233	.. Kieff.	63, 64
Exechia Wiun.	34	.. Walk. (<i>Culex</i>)	167, 168
exilis Coq.	47	.. " (<i>Sciara</i>)	18, 19
eximium de Meij.	43	filitarsis Kieff. (<i>Chironomus</i>)	76
exornata Speis.	243	.. " (<i>Sphaeromyias</i>)	54
extremus Walk.	18	simbriatus Kieff.	71
		Finlaya Theob.	133
falcata Banks	267	hissicornis Kieff.	65
falcipes Banks	174	flammiventris Brun.	26
fallax de Meij.	205	flava Brun. (<i>Atarba</i>)	235
familiaris Ost. Sack. (<i>Libnotes</i>)	226	.. " (<i>Erioptera</i>)	238
.. " (<i>Pachyrhina</i>)	206	.. " (<i>Mongoma</i>)	242
fascialis Kieff.	59	.. " (<i>Styringomyia</i>)	245
fasciata F.	125	.. " (<i>Trichocera</i>)	250
.. Guer.	258	.. Leices. (<i>Chaetomyia</i>)	145, 146, 147
.. Macq.	206	.. Ludl.	102
.. Mg. (<i>Macrocera</i>)	24	flavellus Kieff.	48
.. " (<i>Platyura</i>)	25	flaveolata F.	187
.. " (<i>Rhynostia</i>)	33	flavescens Brun. (<i>Cladura</i>)	248
.. " (<i>Trichoptera</i>)	82	.. " (<i>Tipula</i>)	201
.. Westw.	36	.. Kieff.	56
fasciatipenne Kieff.	69, 272	.. L.	237
fasciatipennis Kieff.	70	flaviceps Kieff. (<i>Atrichopogon</i>)	47, 271
.. Say	255	.. " (<i>Forcipomyia</i>)	50
fasciatus F. (<i>Culex</i>)	125	.. Ender.	102
.. Mg.	54	flavicollis Brun. (<i>Parabrunettia</i>)	83
.. Say.	55	.. " (<i>Sciara</i>)	20
.. Walk.	55	.. Kieff.	65
fasciculata Brun.	204	flavicosta Brun. (<i>Geranomyia</i>)	218
fasciolatus Arrib.	148	.. " (<i>Macrocera</i>)	24
fascipennis Brun. (<i>Claduroides</i>)	251, 252	.. " (<i>Tipula</i>)	203
.. " (<i>Dicranomyia</i>)	214	flavidus Kieff.	61
.. " (<i>Ephelia</i>)	256	flavipalpis Kieff.	47
.. " (<i>Rhaphidolabis</i>)	252	flavipennis Brun.	254
.. " (<i>Sciara</i>)	21	.. Giles	139
.. de Meij. (<i>Dapanoptera</i>)	224	.. Skuse	35
.. " (<i>Gnophomyia</i>)	276	flavipes Brun.	263
.. Mg.	31	.. de Meij.	267
.. Say.	255	.. Macq.	35
fascipes Brun. (<i>Rhynostia</i>)	32	.. Mg.	46, 52
fatigans W.	158	flavipleura Brun.	22
febrifer Banks	106	flaviventrif Ludl.	106
febrilis L.	41	flaviventrif Brun.	22
Feltinella Theob.	94	flavithorax Brun. (<i>Mycomyia</i>)	27
Feltidia Dyar	131	.. " (<i>Tipula</i>)	204
Feltodiplosis Kieff.	12	flaviventrif Brun. (<i>Geranomyia</i>)	219
femorialis Dol.	19	.. " (<i>Mycomyia</i>)	27
.. Edw.	240	.. " (<i>Platyura</i>)	25
femorata Mg.	55	.. Kieff. (<i>Bezzia</i>)	56
.. de Meij.	233	.. " (<i>Chironomus</i>)	73
fenestrata Brun.	263	flavobrunnea Brun. (<i>Dicranomyia</i>)	215
.. de Meij.	239	.. " (<i>Leia</i>)	32
.. Ost. Sack.	233	flavocincta	222
fenestralis Scop.	265	flavofasciata	192
ferruginea Brun. (<i>Acrodicrania</i>)	31	flavofemorata	20
.. " (<i>Erioptera</i>)	237	flavoferruginea Ost. Sack.	248
.. " (<i>Macrocera</i>)	24	flavohirta Brun.	30
.. " (<i>Mycomyia</i>)	27	flavoides Brun.	194, 195
.. Mg.	252	flavolimbata	33
ferruginosa Brun. (<i>Rhamphidia</i>)	228	flavomarginata Brun. (<i>Gonomyia</i>)	240
.. Wulp.	261	.. " (<i>Platyura</i>)	26
festiva Brun.	221	flavus Brun.	194
.. Kieff.	54	.. Kieff.	64
Ficalbia Theob.	170	Flebotomus Rond.	85
filicornis Brun.	204	fletcheri Edw.	218
.. Kieff.	53	fletcheri Felt	17

	Page		Page
fluitans Felt	9	funesta Ludl. nec Giles	106
fluvialis	271	funestus Giles	107
fluviatilis	15	" Harris	39
" James	107	fungorum De Geer	34
" Leices.	144	furcata Brun.	253
foochowensis Theob.	158	Furcinerva Rond.	18
forcipata de Meij.	227	furens Poey	48
" Ost. Sack.	39	Furina Jaen.	246
Forcipomyia Mg	46, 49	fusca Brun.	254
forficularius Kieff.	75	" de Meij.	238
formosae Kieff. (<i>Chironomus</i>)	79, 273	" Edw.	262
" " (<i>Culicoides</i>)	52, 272	" Leices. (<i>Dasomyia</i>)	171, 172
" Tsuzuki I	105, 109	" " (<i>Skieromyia</i>)	181
" " II	109	" " (<i>Stegomyia</i>)	128
formosana Edw.	245	" " nec Theob. (<i>Urano-</i>	
" Ender.; de Meij.	30	" " " <i>taenia</i>)	176
" Kieff.	58, 272	" Mg. (<i>Cecidomyia</i>)	10
formosanus (<i>Procladius</i>)	63, 272	" " (<i>Linnobia</i>)	212
" " (<i>Tanytarsus</i>)	68, 272	" Theob. (<i>Armigeres</i> , <i>Des-</i>	
formosensis Edw.	208	" " " <i>voidea</i>)	122, 274
formosicola (formosicola) Kieff.	79, 273	" " (<i>Protomelanoconion</i>)	157
formosus Ludl.	113	" " (<i>Trichorhynchus</i>)	152
" Walk.	125	fuscana W.	209
fortis Brun.	215	fuscans W.	158
fowleri Chris	99, 100, 114	fuscicornis Edw.	235
fragilis Leices.	144	fuscinervis Brun. (<i>Libnotes</i>)	227
" Lw.	17	" " " (<i>Tipula</i>)	200
" Ludl.	164	fuscipennis Mcq.	189
" Theob.	110	fuscitarsis Kieff.	74
fragillima Westw.	241	fuscitibia Brun.	41
fratella Brun.	229	fuscocephalus Theob.	166
fratercula Brun.	20	fuscoflava, Brun.	208
fraterna	214	fuscolimbata Ender	22
" Edw.	247	fuscomarginata Ender.	231
fraudatrix Theob.	153	fuscopteron Theob.	149
freerae Banks	113	fuscofuscus Kieff.	61
frontosa Theob.	181	fuscofuscus Ost. Sack.	255
fruhstorferi Rubs.	23	fuscus Theob. (<i>Pectinopalpus</i>)	151
fryeri Edw. (<i>Macrocera</i>)	24	fusiger Kieff.	64
" " (<i>Styringomyia</i>)	245	galii Kieff.	10
fulginosus Giles	98	gamma Brun	208
" var. fowleri Alcock	114	" Ender.	263
fulleri Ludl.	154	gardneri Ludl.	130
fullowayi Alex.	210	gandens Walk.	190
fulva Felt	16	gedehana de Meij.	198
fulvescens Brun.	21	gedericola	203
" Kieff.	77	Geitomyia Leices.	135
fulviceps Walk.	257	gelidus Theob.	156, 163
fulvicollis F.	38	Geneja Lioy	24
fulvipennis Walk.	198	generosi Felt	15
fulvohirta Brun.	82	geniculata Brun.	81
fulvolateralis Brun.	194	geniculatus	236
fumicosta Brun.	204	genitalis Brun. (<i>Erioptera</i>)	238
fumifasciata Brun.	198	" " (<i>Geranomyia</i>)	218
fumifascipennis Brun.	203	" " (<i>Linnophila</i>)	253
fumifinis Walk.	197	" " (<i>Rhymosia</i>)	33
fumipennis Brun.	199	genurostris Leices.	178
" Kieff. (<i>Clinotanytus</i> ,		Geranomyia, Hal.	217
" " <i>Procladius</i>)	61	germana Os. Sac.	231
" Steph.	157	gibbosa Edw.	45
fumipes Brun	26	" W.	44
fumiplena Walk.	189	giblini Taylor	149
funerea Brun.	26	gigas Giles	94, 100
" Leices.	183	Glesia Theob.	135
funereus Theob.	143	gilesii	117
funesta Ender.	23	gilvipes Brun.	83
" Giles	92, 107	glabra	254
funesta Leices.	119		

	Page		Page
glabrimanus Kieff. ..	80	guttipennis de Meij. ..	56
glabripennis Brun. ..	251	Gymnastes Brun. ..	234
glabripes de Meij. ..	233	Gynnometopa Coq. ..	125
.. Kieff. ..	79	Gynoplistia Westw. ..	257
glandifex Kieff. ..	11	haasi Kieff. (<i>Daphnephila</i>) ..	11
Glaphyoptera Sch. ..	169, 170 (<i>Panteliola</i>) ..	9
glaphyoptera Winn. ..	31	Haasiella Kieff. ..	55
glauciventris Kieff. ..	79, 273	Haemasson Lw. ..	85
Glochina Mg. ..	212	haemorrhoidalis F. ..	116
gloriosus Kieff. ..	78	halifaxii Theob. ..	164
Glyptotendipes Kieff. ..	79, 273	hallii James ..	95, 102
Gnophodeomyia Theob. ..	156, 157	.. Kieff. ..	71
gnophodes Theob. ..	159	Hallomyia Kieff. ..	11
Gnophodromyia Theob. ..	156	halophila ..	45
Gnophomyia Ost. Sack. ..	246	halterata ..	238
Gnoriste Mg. ..	29	hamifera Kieff. ..	57
Goniodineura Wulp ..	223	Hammatorrhina Lw. ..	36
Goniomyia Ost. Sack. = Gonomyia	240	Harpagomyia de Meij. ..	178
Gonomyia Mg. ..	240	Harpomyia Felt ..	10
gossypii Felt. ..	10	hatiensis Carter ..	143
goughii Theob. ..	159	Hebotomus Rond. ..	85, 86
Grabhamia Theob. ..	131, 142	Heinzmannia Ludl. (<i>Heizmannia</i>)	183
grabhamii (grabhami) Theob. ..	92	Helioidiplosis Felt ..	269
graciliforceps Kieff. ..	71	Helius St. Farg. and Serv. ..	228
gracilipes Kieff. ..	56	heliwigi de Meij. ..	267
gracilis Brun. (<i>Dicranophragma</i>)	255	Helobia St. Farg. and Serv. ..	249
.. .. (<i>Mesocyphona</i>) ..	239	Heptaphlebomyia Theob. ..	155
.. .. (<i>Paracladura</i>) ..	248, 249	HEPTAPHLOEBOMYINA ..	91
.. .. (<i>Tipula</i>) ..	200	HEPTAPHLOEBOMYINAE ..	91
.. de Meij. (<i>Dolichopeza</i>) ..	210	heptatomus Kieff. ..	78
.. .. (<i>Eupeda</i>) ..	239	heterocera ..	50
.. Leices. (<i>Stegomyia</i>) ..	128	heterocerus ..	79
.. .. (<i>Topomyia</i>) ..	180	Heteromyia Say ..	43, 55
.. .. (<i>Zeugomyia</i>) ..	180	Heteronycha Arrib. ..	141
.. Theob. ..	115	heteroptera de Meij. ..	22
gracillima Brun. ..	200	heteropteris Kieff. ..	65
grahami Theob. ..	120	hewitti Edw. ..	154
Grahamia ..	178	hexatomus Kieff. ..	78
graminicola Kieff. ..	16	hiemalis De Geer ..	250
graminis Kieff. ..	12	hilpa Walk. ..	258
.. Leices. ..	166	himalayae Kieff. (<i>Culicoides</i>) ..	52
grandior Brun. ..	238 (<i>Dibezzia</i>) ..	58
grandis ..	25 (<i>Prohelia</i>) ..	48, 49
granifex Kieff. ..	15 (<i>Rhopalocladius</i>) ..	61
Grassia Theob. ..	92 (<i>Tanypus</i>) ..	60
grata Banks ..	116, 118	himalayana Edw. ..	246
gratiosus Big. ..	41	himalayana Giles ..	131, 137
gravelyi Brun. (<i>Eriocera</i>) ..	265	himalayanus Kieff. ..	77
.. .. (<i>Tipula</i>) ..	203	himalayensis Annand. ..	87-273
grayii Theob. ..	182	.. Brun. (<i>Mycetophila</i>) ..	35
greeni Brun. ..	263 (<i>Tipula</i>) ..	199
.. Theob. (<i>Howardina</i>) ..	131	hirsuteron Theob. (<i>hirsuteros</i>) ..	164
.. .. (<i>Wyeomyia</i>) ..	133	hirsutum Ludl. ..	136
Greenomyia Brun. ..	31	.. Theob. ..	163
gregoryi Ludl. ..	131	hirta Felt ..	12
grewiae Banks ..	268	.. Mg. ..	26
griseiceps de Meij. ..	247	Hirtea Mg. ..	39
griseifrons Brun. ..	42	hirtilineata Brun. ..	21
griseipennis ..	201	hirtipennis ..	82
griseolateralis Brun. ..	35	hirtipes de Meij. ..	46
griscens ..	42	.. Kieff. ..	67
griscus Annand. ..	87	hispaniola Theob. ..	103
grossipes Kieff. ..	73	hispida Theob. ..	171
Gualteria Lutz ..	135	Hispidimyia Theob. ..	171
gubernatoris Giles ..	136, 137	hispidosus Skuse ..	120
guttata Hutton ..	31	Hodgesia Theob. ..	179
.. Theob. ..	149	holopticus Brun ..	223
guttifer de Meij ..	51		

	Page		Page
honesta Brun. ..	253	inconspicua Theob. ..	157
Hormomyia ..	271	inconspicuous Brun. ..	236
hortulanoides Brun. ..	40	.. Kieff. (<i>Caprodiplosis</i>) ..	14
hortulanus L. ..	39 (<i>Diplocus</i>) ..	14
.. Mg. ..	46 (<i>Procladius</i>) ..	63
Howardia Theob. ..	92	incunctaus Walk. ..	189
Howardina ..	131, 142	indecora Kieff. (<i>Bezzia</i>) ..	57
Hulecoetomyia Theob. ..	142, 274 (<i>Culicoides</i>) ..	52, 272
humberti Ost. Sack. ..	261	indecorabilis Leices. ..	144
humeralis Brun. ..	33	indefinita Brun. ..	27
hunteri Strickland ..	114	.. Ludl. ..	98
hyalina Mg. ..	40	indianus Kieff. (<i>Atrichopogon</i>) ..	47
hybrida Leices. (<i>Colonymyia</i>) ..	182 (<i>Oligotrophus</i>) ..	9
.. .. (<i>Ficalbia, Conopomyia</i>) ..	172	indica Brun. (<i>Anomalomyia</i>) ..	31
.. Mg. ..	249 (<i>Antocha</i>) ..	232
hybridus Edw. ..	123 (<i>Blepharocera</i>) ..	30
Hyperdiplosis Felt ..	269 (<i>Limnobia</i>) ..	221
hybridus Edw. ..	123 (<i>Mycomyia</i>) ..	27
hypocrites Brun. ..	269 (<i>Odontopoda</i>) ..	30
Hypodiplosis Kieff. ..	13 (<i>Plecia</i>) ..	39
hycranus Pallas ..	161 (<i>Rhaphidolabis</i>) ..	252
idalia Ost. Sack. ..	190 (<i>Xiphura</i>) ..	187
Idioneura Phil. ..	249	.. Eaton (<i>Brunettia</i>) ..	84
Idiophloebe ..	245 (<i>Pericoma</i>) ..	84
ignicolis Walk. ..	38	.. Felt (<i>Harponomyia</i>) ..	10
Ilesia Rond. ..	239 (<i>Streptodiplosis</i>) ..	13
Ilisomyia Rond. ..	236	.. Kieff. (<i>Heteromyia</i>) ..	55
Iisophila ..	239 (<i>Lestremia</i>) ..	18, 271
imitator Leices. (<i>Slegomyia</i>) ..	129 (<i>Micropsectra</i>) ..	68
.. .. (<i>Verrallina</i>) ..	144 (<i>Schizomyia</i>) ..	10
immaculata Mg. ..	259	.. Theob. (<i>Anopheles, Neocellia</i>) ..	94, 112
.. Wulp ..	266 (<i>Neomacleaya</i>) ..	135
immaculatus James ..	110 (<i>Phonimyia</i>) ..	183
immaculipennis Brun. ..	191	.. Walk. ..	18
immisericos Walk. ..	117, 118	indica Theob. var. simplex ..	143
impatibilis ..	125	indicum Becher ..	42
impellens Walk. nec Theob. ..	159	indicus Theob. (<i>Neomacleaya</i>) ..	134, 143
.. Theob. nec Walk. ..	161	.. Kieff. (<i>Dicroneurus, Epidosis</i>) ..	17, 271
imperfecta Brun. ..	202 (<i>Isoplastus</i>) ..	63
impiger Walk. ..	160 (<i>Oligotrophus</i>) ..	9
implacabilis Walk. ..	160 (<i>Pachyleptus</i>) ..	55
implexa Theob. ..	94	indiensis Theob. (<i>Anopheles</i>) ..	95
imponens Walk. ..	225 (<i>Nyssorhynchus</i>) ..	108
impostor Brun. (<i>Plecia</i>) ..	39	indistincta Brun. ..	26
.. .. (<i>Sciara</i>) ..	21	indivisa Walk. ..	28
impressa Walk. ..	225	Indodiplosis Felt ..	13
imprimens ..	136	inepta Annand. ..	185, 274
impunctata Brun. ..	83	inermicus Kieff. ..	54
impunctus Don. ..	103	inermipes .. (<i>Micropsectra</i>) ..	272
inaequalis Kieff. ..	53 (<i>Stilobezzia</i>) ..	55
incerta Brun. (<i>Acrodicrania</i>) ..	31	inexorabilis Walk. ..	125
.. .. (<i>Erioptera</i>) ..	237	infindens Walk. ..	194
.. .. (<i>Polylepta</i>) ..	28	infixa Walk. (<i>Eriocera</i>) ..	260
.. .. (<i>Toxorhina</i>) ..	230 (<i>Limnobia</i>) ..	220
.. Kieff. ..	10	infictus Theob. ..	167
incertus Brun. (<i>Toxorhina, Limnobia</i>) ..	230	infula Theob. ..	132
incompleta Brun. (<i>Gonomyia</i>) ..	240	infuscata Brun. ..	211
.. .. (<i>Limnophila</i>) ..	254	Ingramia Edw. ..	171
.. .. (<i>Rhaphidolabis</i>) ..	252	innocens Brun. ..	216
inconsequens Brun. ..	254	innotabilis Walk. ..	225
inconspicua Brun. (<i>Empeda</i>) ..	239, 240	inordinans ..	197
.. .. (<i>Macrocera</i>) ..	24	inornata Theob. (<i>Culicomyia</i>) ..	151, 152
.. .. (<i>Rhamphidia</i>) ..	228 (<i>Gnophodeomyia</i>) ..	156
.. .. (<i>Sciara</i>) ..	21 (<i>Squanomyia</i>) ..	181
.. .. (<i>Tipula</i>) ..	198	inornatus Walk. ..	118
		insignis Brun. (<i>Leia</i>) ..	32
	 (<i>Teucholabis</i>) ..	233

	Page		Page
insignis Edw. ..	238	Kempi Kieff. (<i>Bezzia</i>) ..	58
.. de Meij. ..	191 (<i>Polypedilum</i>) ..	69
.. Wulp ..	255	Kempia Kieff. ..	45
insolita Walk. ..	257	kerberti de Meij. ..	276
instabilis ..	72	Kerteszia Theob. ..	94
intermedia Rothw. ..	106	kiefferi Patton ..	52
interrupta Brun. (<i>Cladura</i>) ..	248	Kingia Theob. ..	125
.. .. (<i>Tipula</i>) ..	109	kingii ..	135, 140
.. Kieff. ..	53	klossi Brun ..	256
inusta Mg ..	232	.. Edw. ..	202
iridescens de Meij. ..	43	kobusi de Meij. ..	212
iris Kieff. ..	11	kochi Don. (<i>Anopheles</i>) ..	102
irrorata W. ..	248 (<i>Ochlerotatus</i>) ..	136
ischaemi Kieff. ..	9, 271	konoupi Brullé ..	125
isoneuromyia Brun. ..	25	koreicus Edw. ..	141
Isoplastus Skuse ..	63	korinchiensis Edw. ..	204
		kowarzii Dzied. ..	34
jacobsoni de Meij. ..	47	Kronodiplosis Felt ..	268
.. Edw. (<i>Styringomyia</i>) ..	246	kuchingensis Edw. ..	123
.. .. (<i>Tipula</i>) ..	204		
jamaciensis Theob. ..	131	Labidomyia Steph. ..	46
Jamesia Chris. ..	156	laconica Ost Sack. ..	206
jamesii Blanch ..	108	lacteipennis Kieff. ..	56
.. Edw. ..	141	lacteitarsis Brun. ..	83
.. List. nec Theob. ..	99	laeta F. ..	188
.. Theob. nec Leices. ..	108	lampronotus Kieff. ..	74
japonica W. ..	37	lamprothorax ..	74
japonicus, var. ceylonicus Theob. ..	167	laniger W. ..	120
javaensis Theob. ..	119	lasiochirus Kieff. ..	76
javana de Meij. (<i>Linnophila</i>) ..	251	Lasioconops Theob. ..	155
.. .. (<i>Orimarga</i>) ..	232	Lasioptera Mg. ..	8, 267
.. Edw. ..	245	LASIOPTERARIAE ..	8, 267
.. W. ..	205	lasiopterus Kieff. ..	67
javanensis Alex. ..	232	lasioptus Kieff. ..	67
.. de Meij. (<i>Erioptera</i>) ..	237	Lasiosoma Winn. ..	26
.. .. (<i>Rhyptidia</i>) ..	223	latefurcata Brun. ..	212
.. Kieff., Edw. ..	44	.. Ender. ..	23
javanica Alex. (<i>Atarba</i>) ..	235	latelineata Brun. ..	21
.. .. (<i>Geranomyia</i>) ..	219	lateralis Ludl. ..	174
.. .. (<i>Ula</i>) ..	257	.. Theob. ..	178
.. Dol. ..	187	laticornis Mg ..	25
.. Kieff. ..	15	laticornis Walk. ..	18
.. Kieff. and Leeuwen ..	9	latifascia ..	224
javensis Dol. (<i>Eriocera</i>) ..	258, 259	latifasciipennis Brun. ..	192
.. .. (<i>Pachyrhina</i>) ..	205	lat frous Brun. ..	230
jenkinsi Kieff. ..	61	lat squama Coq. ..	156
Jenkinsia Kieff. ..	45	lativentris Bezzi ..	264
jenseni de Meij. ..	168	Laveraina Theob. ..	93
jesoensis Tsuzuki ..	95, 97	Lechria Skuse ..	244
jeyporiensis James ..	109	Leia Mg. ..	31
JOBLOTINA ..	90	leicesteri Edw. (<i>Aedes</i>) ..	145
JOBLOTINAE ..	90 (<i>Uranotaenia</i>) ..	176
johannis L. ..	40	.. Theob. ..	118
Johannseniella, Will. ..	55	Leicesteria Theob. ..	145
joloensis Ludl. ..	122, 123	Leicesteriomvia Brun. ..	146
josoensis Tsuzuki ..	95	Leiponenra Skuse ..	235
jucunda Ost. Sack. ..	257	Leiomya Rond. = Leia ..	31
jugraensis Leices (<i>Desoidya</i>) ..	123	Lejosoma	31
.. .. (<i>Hulecoetomyia</i>) ..	144	Lepidoplatus Coq. ..	134
juniperinus L. ..	9	Lepidotomyia Theob. I ..	134
jurgiosa Walk. ..	257 II ..	134
juxapallidiceps Theob. ..	153	Leptina Mg. ..	210
kambangani de Meij. ..	228	leptochirus Kieff. ..	76
Kamptodiplosis Felt ..	269	Leptoconops Skuse ..	58
karwari James in Theob. ..	112	leptogastrus Kieff. ..	67
Kempi Brun. (<i>Ebiphragma</i>) ..	256	leptomeres Th ob. ..	107
.. .. (<i>Eriocera</i>) ..	265	Leptomorphus Curt. ..	29
.. .. (<i>Mongoma</i>) ..	242	Leptorhina Steph. ..	228

	Page		Page
Leptosomatomyia Theob.	178	longipes Theob.	165
Leslieomyia Chris.	135	" Walk.	19
Lestodiplosis, Kieff.	14, 269	longirostris de Meij.	25
Lestremia Macq.	17, 18	" Leices. (<i>Ficalbia</i>)	172
LESTREMINAE	17	" " (<i>Uranotaenia</i>)	175
leucomeres Giles	129	" Mg	228
Leucomyia Theob.	156	" Theob.	183
leucopeza de Meij.	244	longispatha Kieff.	9
leucopogon Kieff.	53	longistila Kieff.	58
leucopus Don.	99	longitudinalis Brun.	22
leucopyga Wulp.	198	longivalvis Kieff.	72
leucosphyrus Don.	102	longivena Edw.	212, 213
leucotarsus Kieff.	73	Longurio Lw.	204
leucotelus Walk.	260	Lophocelomyia	94
leveillei, Kieff.	17	Lophoceratomyia Theob.	152
lewaldii Ludl.	119	Lophomyia Giles	94
Libnotes Westw.	224	Lophoscelomyia Theob.	94
lignicola Brun.	46	lophoventralis	135, 137, 140, 141
LIMATINAE	91	lorentzi de Meij.	224
limbata Brun.	26	loricatus Leices.	165
limbipennis Macq.	36	Lowiola Kieff.	15
Limnaea Rond.	237	lowisii Theob.	140
Limnobia Mg.	220	lucernarum Kieff.	76
LIMNOBIINAE	211	lucida de Meij.	244
LIMNOBIINI	212	luciensis Theob.	126
Limnobiorynchus, Westw.	217, 229	luctuosa de Meij.	191
Limnoica Rond.	237	ludlowi Blanch.	166
Limnomyia	252	" Brun. nom. nov.	173
Limnomyza	220	" Theob.	95, 110
Limnophila Macq.	252	Ludlowia	171
LIMNOPHILINI	251	lugens W.	33
limpida Edw.	227	lunata Westw.	261
linderae Kieff.	11	lunigera Walk.	260
lindesayi Giles	100	luridus Dol.	167
linearis Alex.	219	lutea Edw.	227
lineata F.	25	" Ludl.	156, 165
" Kieff.	55	" Mg. (<i>Erioptera</i>)	237, 239
" Ludl.	115	" " (<i>Macrocera</i>)	24
lineatopennis Ludl.	124	luteinervis de Meij.	52
lineatus Kieff.	17	luteiventris Brun.	20
lineola Kieff.	56	luteoabdominalis Theob.	133
Liponeura Skuse= <i>Leiponeura</i>	235	luteoannulatus Theob.	159
listoni Giles nec List.	101	luteocephala Newstead	125
" List. nec Giles	107	luteola Theob.	166
lobaticeps Kieff.	73	luteolateralis Theob.	124
lobaticollis	73	luteoventralis	183
lobatus	68	lutescens Leices.	175
longiareolatus Macq.	170	lutzi Cruz	94
longicalcar Macq.	50	Lutzia Theob.	155
longichaeta Brun.	85	luzonensis Felt	270
longicornis Brun.	210	" Ludl.	171, 172
" Dol.	198	Luzonomyia Felt	267
" Kieff.	46		
" Mg.	24	macedoniensis Cot. and Hovasse	111
longicosta Brun.	34	Macleaya Theob.	142
longicrus Kieff.	74	macleayi Skuse	159
longifurcata Brun.	26	Macrobrachius Dzied.	33
longifurcatus Theob. nec Becker	151	Macrocera Mg.	24
longinervis Brun. (<i>Limnobia</i>)	221	MACROCERINAE	24
longinervis Brun. (<i>Sciara</i>)	20	macrocerus Kieff.	59
longipalpis Kieff.	79	macrochaeta	60
" Leices.	145	macrochirus	67
" Leices. in Theob.	146	macrofilum Banks	270
" Newst. and Carter	157	Macropeza	44
" Wulp.	161	macroptera Macq.	256
longipennis Brun. (<i>Rhaphidolabis</i>)	252	" Phil.	249
" " (<i>Limnobia</i>)	222	macropus Blanch.	165
" " (<i>Sciara</i>)	22	Macrorhynchus	183

	Page		Page
macrorhynchus Kieff. ..	47	marginata Mg. ..	27
macroscelus Kieff. ..	78	marginatus Annand. ..	86
macrostoma	51	" Say	266
macrothrix	50	marginotata Brun. ..	83
Macroura, Berendt ..	24	marmorata Brun. ..	243
maculata Mg. (<i>Dixa</i>) ..	185	" Hffg. in Mg. ..	256
" " (<i>Rhyppidia</i>) ..	223	marmoratipennis Brun. ..	200
" " (<i>Trichosticha</i>) ..	239	marmoripennis Brun. ..	214
" Theob. (<i>Nyssorhynchus</i>) ..	104	matteiana Kieff. and Cecconi ..	13
" " (<i>Orthopodomyia</i>) ..	176	maxima Leices. ..	175
" " nec Mg. (<i>Pecomomyia</i>) 134, 141		mcgregori Banks	113
" " (<i>Pseudograhamia</i>) 134, 138		mediofasciata Theob. ..	143
" " var. of <i>Anoph.</i>		mediolineata Ludl. ..	143
<i>lindsayi</i> Giles. 100		" Theob	171
maculatus Mg. nec Theob. (<i>Culex</i> ,		mediolineatus	137, 138
<i>Oehlerotatus</i>)	141	mediopunctata	130
" Theob. 93, 104, 112		mediopunctatus Theob. ..	94
maculicoxa Ender.	32	mediovittata Coq.	125
maculicrura Theob.	161	Megaculex Theob.	171
maculipalpis Giles	108	MEGARLORRHINI	90
maculipennis Brun. (<i>Dixa</i>) ..	185	Megarhina Giles, = Megarhinus Rob.	
" " (<i>Psychoda</i>)	82	Desv.	115
" " Mcq. (<i>Aporosa</i>)	220	Megarhina St. Farg. and Serv. nec	
" " Mg.	92	Megarhinus Rob. Desv. ..	228
" " Say	32	MEGARHINES	90
" " Wulp.	266	MEGARHINI	90
maculipes Theob. (<i>Arribalzagia</i>)	93	MEGARHININA	90
" " (<i>Orthopodomyia</i>)	177	MEGARHININAE Theob	90
maculipleura Edw.	247	Megarhinus Rob. Desv. nec Me-	
" " Leices.	175	garhina St. Farg. and Serv. ..	115
maculiventris Brun.	265	Megistocera W.	209
maculosa James and List. ..	112	melanaspis W.	37
madagascarensis Theob. ..	155	melancholica Walk.	257
Maekistocera v. Megistocera ..	209	melanochirus Kieff.	75
magna Theob. (<i>Brevirhynchus</i>) ..	146	Melanoconion Theob.	155
" " (<i>Lepidotomyia</i>) 134-137		Melanoconium Blanch	155, 156
magnicornis Ender.	106	Melanoconops Theob.	155
magnifica Kieff.	12	melanomera Walk.	197
magnificus Leices.	119	melanophorus Kieff.	75
magnipennis Kieff.	12	melanopterus Giles	139
magnus Theob.	146	melanostictus Kieff.	76
maiae Edw.	124	melanostolus	74
Maillotia Theob.	156	melanothorax	69
majesticus Brun.	104	melanura Ost. Sack.	206
major Annand.	87	" Walk.	188
" Theob.	150	melanurus Coq.	157
malabaricus Theob.	87	" Dol.	59
Malala Heller	178	Meleagris Ost. Sack.	261
Malaya Leices.	178	Memnemyia Strick.	95
malayi Leices. (<i>Aedes</i>)	168	mendacis Daniels	182
" " (<i>Hodgesia</i>)	170	Mesochria Ender.	29
" " (<i>Verrallina</i>)	143	Mesocyphona Ost. Sack. ..	239
" Theob. (<i>Uranotaenia</i>) 122-274		mesopyrrha W.	259
malfeysi Newstead	171	metaboles Theob.	106, 107
mammifer Leices.	154	metallica Alex.	211
mangiferae Felt	13	" Leices. (<i>Ficalbia, Cono-</i>	
" Kieff.	9	" <i>pomyia</i>) 171, 172	
Manguinhosia Cruz	94	metallica Leices. (<i>Wyeomyia</i>) ..	183
mangyana Banks	106	metallicus Leices. in Theob. ..	119
manillensis Felt	267	METANOTOTRICA	91
manillensis Sch. (<i>manillensis</i> ,		metatarsale Brun.	42
<i>manilliensis</i>)	184	metatarsalis .. (<i>Pericoma</i>) ..	83
manillensis Sch.	59	metatarsata Edw.	176
Mansoniana Blanch.	147	" de Meij.	240
Mansonioides Theob.	149	" Metriocnemus Wulp.	64
mansueta Ost. Sack.	261	micans Leices.	174
marginalis Bezzi	228	Micraedes Coq.	168
marginata Brun.	222	microannulatus Theob. ..	150

	Page		Page
microcerus Kieff. ..	162	mucidus, Leices. nec Karsch ..	120
Microculex Theob. ..	155	multicolor Camboulin ..	103
Microperrisia Kieff. ..	9, 271	multifida Kieff. ..	14
Micropsectra ..	68, 272	multimaculosus Leices. ..	164
microptera Giles in Theob. ..	127	multiplex Theob. ..	134, 139
microtaeniatus Theob. ..	162	multipunctata Brun. ..	253
milnei Kieff. ..	69	multipunctipennis Brun. ..	255
Mimeteomyia Theob. ..	157	multistriatus E nder. ..	247
Mimetculex Theob. ..	135	munda Brun. ..	201
mimeticus Noe ..	161	musicus Say. ..	92
mimomyiaformis Newstead ..	171	Mycetina Rond. ..	35
Mimosciara Rond. ..	18	MYCETOBIIINAE ..	29
Mimulus Edw. ..	167	Mycetophila Mg. ..	35
minima Kieff. ..	46	MYCETOPHILIDAE ..	18
.. Ludl. ..	173	MYCETOPHILINAE ..	29
.. Theob. ..	170, 171	Mycodiplosis Rubs. ..	16
minimus Leices. ..	166	Mycomyia Rond. (<i>Mycomya</i>) ..	27
.. Theob. (<i>Anopheles</i>) ..	105	.. Johann. ..	28
.. .. (<i>Toxorhynchites</i>) ..	118, 119, 274	Mycozetoca Rond. ..	35
minor Leices. (<i>Lophoceratomyia</i>) ..	154	Myxosquamus Theob. ..	135
.. .. (<i>Topomyia</i>) ..	179	myzomyfacies ..	103
.. Theob. ..	166	Myzomyia Blanch ..	92, 273
minuta .. (<i>Culicada</i>) ..	135, 136	Myzorhynchella Theob. ..	94
.. .. (<i>Minomyia</i>) ..	171	Myzorhynchus Blanch. ..	92, 273
minutissima, Theob. (<i>Culiciomyia</i>) ..	153	nagpori James and List. ..	99
.. .. (<i>Stegomyia</i>) ..	129	Nasiteria Walleogr. ..	251, 256
minutus, Lw. nec Rond. ..	86	nebulosa Edw. ..	241
.. Rond. nec Lw. ..	86	.. de Meij. ..	235
.. Theob. ..	97	nebulosus de Meij. ..	247
mirabilis de Meij. ..	187, 275	.. Mg. ..	266
Misocosmus Kieff. ..	17	Nematocera W. ..	209
Mitopeza Edw. ..	210	Neocellia Theob. ..	94
mixta Brun. ..	83	Neoculex Dyar. ..	155
Mochlostyrax Dyar and Knab ..	156	Neoempheria Ost. Sack. ..	28
mo lesta Leices ..	175	Neoglaphyoptera Ost. Sack. ..	31
.. Mg. ..	212	Neomacleaya Theob. ..	134
.. Ost. Sack. ..	251	Neomelanoconion Theob. ..	151, 156
molestus Costa ..	86	Neomyzomyia ..	95
.. Kieff. ..	52	Neopecomyia Theob. ..	135
.. W. ..	167	Neostethopheles James ..	94
mollissima Hal. ..	256	Neostyringomyia Alex. ..	230
Molobraea Rond. ..	18	nepalensis Brun. ..	244
Molobrina ..	18	.. Edw. ..	246
Molobrus Latr. ..	18	.. Kieff. ..	78
Molophilus Curt. ..	236	.. Westw. ..	259
Molpemyia Theob. ..	135	Nepalia Kieff. ..	63
Mongomia Westw. ..	241	nepenthicola Banks ..	184
Mongomella Ender. ..	244	nepenthis Edw. ..	182
Mongomioides Brun. ..	242	nero Dol. ..	148
monochrous W. ..	193	nervosa de Meij. (<i>Dicranomyia</i>) ..	213
montana Brun. (<i>Dixa</i>) ..	185, 274 (<i>Libnotes</i>) ..	226
.. .. (<i>Trichocera</i>) ..	250	Nesopeza Alex. ..	210
.. de Meij. ..	217	nicnic Banks ..	273
montanus Stanton and Hacker ..	111	niger Annand. ..	86
monticola Edw. ..	229	.. Leices. ..	154
monticola Kieff. (<i>Camptocladius</i>) ..	64	nigerrima Brun. ..	264
monticola Kieff. (<i>Forcipomyia</i>) ..	50	.. Edw. ..	233
montivaga ..	50	.. Theob. (<i>Culiciomyia</i>) ..	153
montivagans Alex. ..	227 (<i>Mansonia</i>) ..	150
.. Kieff. ..	51	.. Giles, nec James and List. ..	100
morosa Ost. Sack. ..	261	.. James and List. nec Giles ..	95, 97, 100
mosquito R. Desv., Arrib. ..	125	nigra Brun. (<i>Gnophomyia</i>) ..	246
motschulskyi Gimm. ..	37 (<i>Leia</i>) ..	32
moultoni Edw. (<i>Armigeres</i>) ..	123 (<i>Limnobia</i>) ..	221
.. .. (<i>Uranotaenia</i>) ..	176	.. de Meij. ..	33
Mucidus Theob. ..	119	.. Leices. (<i>Lophoceratomyia</i>) ..	154
mucidus, Karsch. nec Leices. ..	120		

	Page		Page
nigra Leices. (<i>Topomyia</i>)	180	niveocaputis Ludl.	179
" Theob.	94	niveocincta Kieff.	9
" W.	258	niveoscutellatus Theob.	139
nigrans Stanton	112	niveus Ludl.	139
nigrescens Brun.	222	nivipes Theob. (<i>Anisocheteomyia</i>)	173
" Edw.	247	" " (<i>Anopheles</i>)	99
" Theob.	144	nivipleura Leices.	175
nigricans Kieff.	56, 57, 272	nobile de Meij.	42
" L.	266	nocticolor Edw.	234
nigricephalus Leices	166	nocti vaga Kieff.	49
nigriceps de Meij.	276	nocturnus Theob.	136
" Edw.	229	nodifex Kieff.	16
" Wulp (<i>Goniodineura</i>)	223, 224	nodulosa Mcq.	236
" " (<i>Prionota</i>)	188, 192	nongkodjarensis de Meij.	216
nigricornis Alex.	228	notabilis Skuse	63
nigricoxa Brun.	31, 32, 271	notata de Meij. (<i>Erioptera</i>)	237
nigrifasciatus Theob.	103, 113	" " (<i>Geranomyia</i>)	218
nigriforceps Kieff.	74	" L.	37
nigrina Riedel	264	" Wulp	225
nigripennis Brun. (<i>Bibio</i>)	41	notatinervis Brun.	227
" " (<i>Psychoda</i>)	81	notatipennis " (<i>Geranomyia</i>)	219
" " (<i>Sciara</i>)	20	" " (<i>Psychoda</i>)	82
" Wulp in de Meij.	262	notatus de Meij.	55
nigripalpis	238	Notonotricha Theob.	94
nigripes Brun.	239	notoscriptus Skuse	127
" Staeg.	96	Nototricha Coq.	94
" Zett.	156, 160	novaeguineae de Meij.	202
nigritarsis Ludl.	144	" " (<i>Dicranomyia</i>)	216
" Leices.	176	novaeguineensis de Meij.	72
nigrithorax Brun.	215	9-notata Brun.	85
" de Meij.	275	9-punctatus Kieff.	61
" Wulp in de Meij.	190	novumbrosus Strickland	114
nigritia Ludl.	131	nubeculosa de Meij.	240
nigriventris de Meij.	41	nubilipennis Kieff.	13
nigroannulata Wulp	206	nubipennis Rond.	236
nigroapicalis Brun. (<i>Eriocera</i>)	264	nudipes Kieff.	77
" " (<i>Mongoma</i> ,		Numantia Big.	212
" " (<i>Trentepohlia</i>)	243	nursei Theob.	111
" " (<i>Tiula</i>)	199	Nyssomyzomyia James	94
nigrocinctus Dol.	59	Nyssorhynchus Blanch.	93, 273
nigrofasciata Brun.	33	obediens Ost. Sack.	40
nigrofemorata Edw.	245	obscura Big.	258
" de Meij.	30	" Brun. (<i>Dolichopeza</i>)	211
nigromarginatus Kieff.	73	" " (<i>Plectia</i>)	39
nigronitida Brun.	37	" " (<i>Styringomyia</i>)	245
" (<i>Geranomyia</i>)	219	" de Meij.	242
nigrosignatus Edw.	149	" Edw.	176
nigrosparus Kieff.	73	" Johann.	272
nigrotibialis Brun.	201	" Leices.	168
nigriticus Edw.	167	" Rond.	10
nimba Theob.	93	obscurata Walk.	30
Ninguis Wallengr.	231	obscuripennis de Meij.	40
nipponi Theob.	135, 136	obscuripes	63
nitida de Meij.	217	obscurus Leices.	168
nitidirostris Edw.	210	obturbans Walk.	121, 122
nitidithorax Brun.	22	obtusilobae Felt	13
nitidiventer Giles	181	occipitalis de Meij.	257
nitidulus Kieff.	51	oculta Mg.	250
nitidus	79	ocellata Walk.	250
nitobei Edw.	222	ocellatus Theob. (<i>Anopheles</i>)	102
nivea Leices.	130	" " (<i>Taeniorhynchus</i>)	132
" Ludl.	139	Ochlerotatus Arrib.	133, 142
nivea-amesii Ludl.	128	ochracea Theob.	133
niveiapicalis Brun. (<i>Sciara</i>)	22	ochraceus Edw. part.	138, 140
" " (<i>Dicranomyia</i>)	217	" Theob.	138
niveipes	221	ochripleuris de Meij.	208
niveitarsis Skuse	229	ochrolineata Brun.	185
niveitaeniata Theob.	169, 170		

	Page		Page
nigra Leices. (<i>Topomyia</i>)	180	niveocaputis Ludl.	179
" Theob.	94	niveocincta Kieff.	9
" W.	258	niveoscutellatus Theob.	139
nigrans Stanton	112	niveus Ludl.	139
nigrescens Brun.	222	nivipes Theob. (<i>Anisocheleomyia</i>)	173
" Edw.	247	" " (<i>Anopheles</i>)	99
" Theob.	144	nivipleura Leices.	175
nigricans Kieff.	56, 57, 272	nobile de Meij.	42
" L.	266	nocticolor Edw.	234
nigricephalus Leices	166	nocti vaga Kieff.	49
nigriceps de Meij.	276	nocturnus Theob.	136
" Edw.	229	nodifex Kieff.	16
" Wulp (<i>Goniodineura</i>)	223, 224	nodulosa Mcq.	236
" " (<i>Prionota</i>)	188, 192	nongkodjadjarensis de Meij.	216
nigricornis Alex.	228	notabilis Skuse	63
nigricoxa Brun.	31, 32, 271	notata de Meij. (<i>Erioptera</i>)	237
nigrifasciatus Theob.	103, 113	" " (<i>Geranomyia</i>)	218
nigriforceps Kieff.	74	" L.	37
nigrina Riedel	264	" Wulp	225
nigripennis Brun. (<i>Bibio</i>)	41	notatinervis Brun.	227
" " (<i>Psychoda</i>)	81	notatipennis " (<i>Geranomyia</i>)	219
" " (<i>Sciara</i>)	20	" " (<i>Psychoda</i>)	82
" Wulp in de Meij.	262	notatus de Meij.	55
nigripalpis	238	Notonotricha Theob.	94
nigripes Brun.	239	notoscriptus Skuse	127
" Staeg.	96	Nototricha Coq.	94
" Zett.	156, 160	novaeguineae de Meij.	202
nigritarsis Ludl.	144	" " (<i>Dicranomyia</i>)	216
" Leices.	176	novaeguincensis de Meij.	72
nigrithorax Brun.	215	9-notata Brun.	85
" de Meij.	275	9-punctatus Kieff.	61
" Wulp in de Meij.	190	novumbrosus Strickland	114
nigritia Ludl.	131	nubeculosa de Meij.	240
nigriventris de Meij.	41	nubilipennis Kieff.	13
nigroannulata Wulp	206	nubipennis Rond.	236
nigroapicalis Brun. (<i>Eriocera</i>)	264	nudipes Kieff.	77
" " (<i>Mongoma</i> ,		Numantia Big.	212
" " (<i>Trentepohlia</i>)	243	nursei Theob.	111
" " (<i>Tiula</i>)	199	Nyssomyzomyia James	94
nigrocinctus Dol.	59	Nyssorhynchus Blanch.	93, 273
nigrofasciata Brun.	33	obediens Ost. Sack.	40
nigrofemorata Edw.	245	obscura Big.	258
" de Meij.	30	" Brun. (<i>Dolichopæa</i>)	211
nigromarginatus Kieff.	73	" " (<i>Plectia</i>)	30
nigronitida Brun.	37	" " (<i>Stringomyia</i>)	245
" (<i>Geranomyia</i>)	219	" de Meij.	242
nigrosignatus Edw.	149	" Edw.	176
nigrosparus Kieff.	73	" Johann.	272
nigrotibialis Brun.	201	" Leices.	168
nigriticus Edw.	167	" Rond.	10
nimba Theob.	93	obscurata Walk.	30
Ninguis Wallengr.	231	obscuripennis de Meij.	40
nipponi Theob.	135, 136	obscuripes	63
nitida de Meij.	217	obscurus Leices.	168
nitidirostris Edw.	210	obturbans Walk.	121, 122
nitidithorax Brun.	22	obtusilobae Felt	13
nitidiventer Giles	181	occipitalis de Meij.	257
nitidulus Kieff.	51	oculta Mg.	250
nitidus	79	ocellata Walk.	250
nitobei Edw.	222	ocellata Theob. (<i>Anopheles</i>)	102
nivea Leices.	130	" " (<i>Taeniorhynchus</i>)	132
" Ludl.	139	Ochlerotatus Arrib.	133, 142
nivea-amesii Ludl.	128	ochracea Theob.	133
niveiapicalis Brun. (<i>Sciara</i>)	22	ochraceus Edw. part.	138, 140
" " (<i>Dicranomyia</i>)	217	" Theob.	138
niveipes	221	ochripleuris de Meij.	208
niveitarsis Skuse	229	ochrolineata Brun.	185
niveitaeniata Theob.	169, 170		

	Page		Page
ochripes Brun. ..	195	osakensis Theob. ..	159
Octavia Big ..	237	ostentatio Leices. ..	140
8-fasciata Brun. ..	257	Oxydiscus de Meij. ..	247
8-segmentata Brun. ..	26	oxystoma Kieff. ..	52
Oculeomyia Theob. ..	131, 153, 274	Pachydiplosis Kieff. ..	15
Odontonyx Rubs. ..	23	Pachyleptus Walk. ..	55
Odontopoda Aldr. ..	30	Pachyrrhina Macq. ..	205
oecacta Poey. ..	48	pachyrrhina Ost. Sack. ..	261
Olbiogaster Ost. Sack. ..	267	paederiae Banks ..	270
oleracea L. ..	196	" Kieff. and Leeuwen ..	13, 14
Oligomera Dol. ..	258	paenulata Endr. ..	263
oligotrophariae ..	9, 267	pagei Ludl. (<i>Taeniorhynchus</i>) ..	149
Oligotrophus Latr. ..	9, 267, 271	" " (<i>Theobaldia</i>) ..	170
omissa Edw. ..	147	paivai Kieff. (<i>Clinotanypus</i>) ..	61
omissinervis de Meij. ..	198	" " (<i>Culicoides</i>) ..	52
opaca Bezzi. ..	228	" " (<i>Tanypus</i>) ..	59
" de Meij. ..	253	Palaeoanaclinia Meun ..	29
opacus Kieff. ..	51	palawanensis Ludl. ..	139
opalizans Ost. Sack. ..	232	palestinensis Theob. ..	111
opposita Brun. ..	19	pallens Wulp ..	256
optabilis Walk. ..	260	pallescens Brun. ..	21
orbicularis Brun. ..	82	pallida de Meij. ..	224
Oreillia Ludl. ..	171	" Ludl. ..	110
oreophilus Edw. ..	141	" Theob. ..	99
Orfelia A. Costa ..	25	pallidiceps, Theob. ..	165
Oribremia, Kieff. ..	14	pallidicornis de Meij. ..	37
orientalis Brun. (<i>Dolichopeza</i>) ..	210	" Edw. ..	235
" " (<i>Erioptera</i>) ..	237	pallidicoxa Brun. ..	253
" " (<i>Sciara</i>) ..	20	pallidipes Kieff. ..	60
" de Meij. (<i>Gnophomyia</i>) ..	246	palliditarsis de Meij. ..	37
" Edw. ..	267	pallidithorax de Meij. ..	211
" Felt ..	269	" Theob. ..	151
origenus Kieff. ..	47	pallidiventris Brun. ..	241
Orimarga Ost. Sack. ..	231	pallidopalpis Theob. ..	94
Orimargula Mik. ..	232	pallidostriatus Theob. ..	140
oriphila Kieff. ..	50	pallidothorax (pallidithorax) Theob. ..	151, 152
oriplanus Kieff. (<i>Chironomus</i>) ..	75	pallidum Kieff. ..	80
" " (<i>Orthocladius</i> , <i>Tri-</i> " " " <i>chocladius</i>) ..	65	pallidus .. (<i>Procladius</i>) ..	63
" " (<i>Tanypus</i>) ..	60	" " (<i>Prochironomus</i>) ..	273
orissae ..	80	" Walk. ..	195
Ormosia Rond. ..	236	pallipes Mg. ..	158
ornata Brun. (<i>Ephelia</i>) ..	256	palmeri Alex ..	254
" " (<i>Macrocera</i>) ..	24	Palpomyia Meg. in Mg. ..	52
" " (<i>Teucholabis</i>) ..	233	palpus Newstead, Dutton and Todd ..	156
" Theob. ..	143	paludicola Skuse ..	62
ornaticornis Kieff. ..	46	pampangensis Brun. ..	114
ornaticrus ..	49, 271	" Ludl. ..	134, 140
ornatipennis Brun. (<i>Limnophila</i>) ..	254	panalectoros Giles (toris, tros) ..	158
" " (<i>Trentepohlia</i>) ..	243	panangensis Ludl. ..	145
" de Meij. ..	234	pangerangensis Alex. ..	206
ornatipes Brun. ..	214	pannosa Ender. ..	263
" Kieff. ..	59	Panoplites Theob. ..	147, 148
ornatissimus Dol. ..	186	Pantehola Kieff. ..	9
" Kieff. (<i>Chironomus</i>) ..	79	papatasi Scop. ..	85
" " (<i>Clinotanypus</i> , " " " <i>Procladius</i>) ..	60	Parabrunettia Brun. ..	84
ornatithorax Brun. ..	195	Paracladura ..	248
ornatus Brun. ..	29	parallela Brun. (<i>Erioptera</i>) ..	237
" Dol. ..	62	" " (<i>Sciara</i>) ..	22
" Mg. ..	56	Paramongoma ..	243
" Theob. ..	165	parangensis Ludl. (<i>Myzomyia</i>) ..	98
Orseoliella Kieff. ..	15	" " (<i>Uranotaenia</i>) ..	175
ORTHOCLADIARIAE ..	64	Parapleciomyia Brun. ..	39
Orthocladus Wulp ..	65	parascelos Theob. ..	140
Orthopodomys Theob. ..	176	Paratendipes Kieff. ..	69, 273
ortiva Ost. Sack. ..	206	pardalis Dol. ..	62
oryzae Wood-Mason ..	16	Pardomyia Theob. ..	120

	Page		Page
Parexechia Bech. ..	34	picta Mg. ..	8
parva Edw. ..	208	picticornis Brun. ..	210
parvicellula Brun. ..	254	" Ost. Sack. ..	235
parvinotata Brun. ..	209	pictipennis Bezzi ..	243
parvus Taylor ..	162	" Edw. ..	234
Patagiamyia James ..	94	" Kieff. (<i>Dibezzia</i>) ..	58
patricia Brun. ..	200	" Mg. ..	256
pavonina Say. ..	255	pictipes Brun. ..	217
Pecomyia Theob. ..	134	" Kieff. ..	53
pectinata Edw. ..	147	pictiventris Kieff. (<i>Culicoides</i>) ..	52
Pectinopalpus Theob. ..	151	" " (<i>Prochironomus</i>) ..	70
pedata W. ..	196	pictus Dol. ..	72
PEDICINAE Kert. ..	249	" Lw. ..	96, 107
peditaeniatus Leices. ..	96	Pilaria Sintensis ..	252
pelargus Kieff. ..	70	pilicornis Zett. ..	252
pelochloris Kieff. ..	79-273	pilifera de Meij. ..	235
PELOPIINAE ..	59	" Theob. ..	156
pelostolum Kieff. ..	69, 272	pilosa Mg. ..	26
pembaensis Theob. ..	142	pilosipennis Kieff. ..	67
pendula Edw. ..	147	pilosula Wulp ..	197
pennipes Brun. ..	234	pini DeGeer. ..	15
" Ost. Sack. ..	241	pipericola Kieff. ..	10
pentatomus Kieff. ..	68	pipersalatus Giles in Theob. ..	138, 141
pentazonus ..	65	pipiens L. ..	155
Penthetria W. ..	37	pitchfordi Giles ..	113
penumbrinus Edw. ..	196	Planatella Westw. ..	18
perdecora Walk. ..	224	Planetes Walk. ..	18
perdita Leices. ..	144	planicollis Kieff. ..	75
peregrina Brun. ..	232	Platyura Mg. ..	25
peregrinus Kieff. ..	51	Plecia W. ..	38
perennis Ost. Sack. ..	261	plecioides de Meij. ..	233
perichlorus Kieff. ..	78	" Ost. Sack. ..	40
Pericoma Walk. ..	82	" Walk. ..	262
perileucus Kieff. ..	80	Pleciomyia Brun. ..	37
periskelata Giles ..	128	Plectromyia Ost. Sack. ..	251
perniciosus Newstead ..	87	plegipennis Theob. ..	165
Peromyia Kieff. ..	17	plenipennis Walk. ..	224
perplexa ..	46	Plesiomongoma Brun. ..	242
" Leices. ..	130	Plettusa Phil. ..	217
perplexus ..	162	pleurinotata Brun. ..	208
persistans Banks ..	126	plumbeus Hal. ..	96
perturbans de Meij. ..	87	plumbicincta Hal. ..	262
" Walk. ..	148	plumicornis F. ..	184
pettigrewii Theob. ..	167	plumiger Don. ..	95
Phagomyia ..	134	plumosa L. ..	71
phalaenoides L. ..	81	pluto Brun. ..	199
pharoensis Theob. ..	93	Plutodiplosis Kieff. ..	12
philippinensis Banks ..	270	Pneumaculex Theob. ..	156
" Giles ..	181	poeciloptera Ost. Sack. ..	226
" Ludl. (<i>Anopheles</i>) ..	110	Poecilostola Sch. ..	256
" " (<i>Pyretophorus</i>) ..	114	poecilus Kieff. ..	68
Philodendromyia Theob. ..	153	poicilia Theob. ..	133, 138
Philorus Kellogg. ..	36	poicilipes Theob. ..	155
PHLEBOTOMINAE ..	85	polius Kieff. ..	77
Phlebotomus Rond ..	85	Polylepidomyia Theob. ..	181
" sp. Giles ..	87	Polylepta Winn. ..	28
" " Howlett ..	86	Polyleptiomyia Theob. ..	134
Phoniomyia Theob. ..	182	Polypediulum Kieff. ..	68, 272
Phoroctenia Coq. ..	187	polypori Lw. ..	14
Phorodonta Coq. ..	23	polysticta Lw. ..	53
photophila Kieff. ..	54	polystictus ..	73
photophilus .. (<i>Isoplastus</i>) ..	63	pomonae F. Schrk. ..	39
" " (<i>Protenthes</i>) ..	62	ponderosa Walk. ..	19
Phronia Winn. ..	34	Popea Ludl. ..	156
phryganopterus Kolen. ..	236	PORRICONDYLIARIAE ..	16
Phylidorea Big. ..	252	postica Brun. ..	211
Phyllophaga Rond. ..	10	posticalis Lundb. ..	62, 272
Physecrania Big. ..	258	posticatus W. ..	92

	Page		Page
powelli Ludl.	175	Pterocosmus Walk.	258
praecox F. Mg.	40	Ptychoptera Mg.	186
praefica Bezzi	191	PTYCHOPTERINAE	186
praepotens W.	163	pulcher Brun.	236
prasinelus Kieff.	79-273	pulcherrima Arrib.	173
prasiogaster Kieff. (<i>Chironomus</i>)	77	" Brun.	200
" " (<i>Tanytarsus</i>)	68	" Kieff.	13
pretiosus Kieff.	78	pulcherrimus Theob. (<i>Anopheles</i> , <i>Cellia</i>)	109
princeps Brun.	200	pulchra de Meij. (<i>Dapanoptera</i>)	224
Prionocera Ender. nec Lw.	192	" " (<i>Dicranomyia</i>)	213
Prionomyia Steph.	55	pulchricornis Brun.	12, 271
Prionota Wulp	192	" Kieff.	12, 271
pristina Brun.	211, 212	pulchripennis Brun. (<i>Dicranomyia</i>)	214
Proanaclina Meun.	30	" " (<i>Dicranophragma</i>)	255
Probezzia Kieff.	56	" " (<i>Geranomyia</i>)	219
Prochironomus Kieff.	70, 273	pulchripes Kieff. (<i>Palpomyia</i>)	53
Procladius Skuse	62	" " (<i>Tanypus</i>)	59
Procontarinia Kieff. and Cecconi	13	pulchrithorax Brun.	264
Profeltiella Kieff.	269	pulchriventer Giles	131, 137
Prohelia	48, 271	pulicarius L.	50
Prokempia	45	pulla Theob.	152
Prolassioptera	9	pullata Harris	39
prominens Brun.	217	pullicauda Skuse	34
propinqua de Meij.	28	pulvinariae Felt	9
Protenthes Johann.	62	pumila de Meij.	202
Protoculex Felt	134	punctata F.	266
Protomacleaya, Theob.	135	" Latr.	51
Protomelanoconion Theob.	157	punctatipennis Kieff.	70, 72
proxima Brun. (<i>Gonomyia</i>)	241	punctatissima de Meij.	227
" " (<i>Pericoma</i>)	83	punctatus F. Sch.	266
" Edw.	182	puncticornis Brun.	208
proximus Brun.	40	puncticosta	214
pruinosa Rubs.	19	punctifrons Rond.	197
pruinosis Kieff.	73	punctipennis Brun. (<i>Erioptera</i>)	237
prussicus	70	" " (<i>Trichocera</i>)	250
Psectrocladius Kieff.	65	" de Meij.	226
Pseliophora Ost. Sack.	188	" Mg. (<i>Tanypus</i>)	62
Pseudhormomyia Kieff.	15, 271	" " (<i>Symplecta</i>)	249
pseudobarbistris Ludl.	109	" Say	184
Pseudocarrollia Theob.	135	punctipes Theob.	156
pseudococci de Meij.	269	" W.	67
Pseudoculex Dyar	134	punctithorax Brun.	222
Pseudodiurna Theob.	144	punctolateralis Theob.	128
pseudofulvipennis de Meij.	275	punctulata de Meij. (<i>Acyphona</i>)	276
Pseudograhamia Theob.	134	" " (<i>Dicranomyia</i>)	213
Pseudograhamia	179	" James and List.	95
Pseudoheptaphlebomyia Theob.	155	punctulatus Don.	102
Pseudohowardina Theob.	134	" Ender.	247
pseudoinfula Theob.	162	pungens de Meij.	51
pseudolongifurcatus Theob.	151	pungens W.	159
pseudomediofasciatus	145	punjabensis James	101
pseudonivea Theob.	139	purimanus Kieff.	69
pseudopictus Grassi	96	pusilla Dol.	18
pseudoscutellaris Theob.	126	Pycnocrepis Ender.	245
Pseudoskusea	134	pygmaeus Theob.	149
pseudostenotrus	136	Pyrethorus Blanch.	92, 273
pseudotaeniatus Giles	142	pyri F.	40
Pseudotheobaldia Theob.	169	pyrrhochroma Walk.	260
Pseudouranotaenia Theob.	173	quadrata de Meij.	209
pseudowillmori Theob.	104	quadri-cellula Brun.	211
psilochirus Kieff.	76	quadri-fasciata	35
Psilomegalosphys Ender.	23	quadri-furca Walk.	225, 276
psilopterus Kieff.	65	quadri-lobatus Kieff. (<i>Culicoides</i>)	51
Psorophora Rob. Desv.	95	" " (<i>Oligotrophus</i>)	9
psychasta Kieff.	50	quadri-notata Brun.	201
Psychoda Latr.	80	quadri-palpis Edw.	154
PSYCHODIDAE	80		
PSYCHODINAE	80		

	Page		Page
SCIARINAE	18	signifer Coq.	156
Sciobia Lw.	27	" de Meij.	59
scintillans Ludl.	183	sikkimensis Ender	195
Sciophila Mg. Winn.	26, 27	simalurensis de Meij.	264
SCIOPHILINAE	26	similis Edw.	247
Scottella Ender.	30	" Leices.	182
scottiana	29	" Strickland	115
scurroides de Meij.	207	simillima Brun.	203
scutellaris Giles	127	simlensis James and List.	100
" Kieff.	55	simplex Brun. (<i>Aechimongoma</i>)	244
" Theob. nec Walk.	126, 127	" " (<i>Dicranomyia</i>)	215
" Walk. nec Theob.	126	" " (<i>Limnophila</i>)	253
scutellata Edw. (<i>Eriocera</i>)	262	" " (<i>Phronia</i>)	34
" " (<i>Libnotes</i>)	227	" Ost. Sack.	226
Scutomyia Theob.	125	" Theob. (<i>Ficalbia</i>)	171
Scythroprochroa Ender.	23	" " (<i>Heptaphloebomyia</i>)	155
segmenticornis Brun.	21	" " var. of <i>N. indica</i>	143
Seguini Laveran	150	simplissima Alex.	216
selecta Walk.	19	simulacri Felt	16
selectissima Walk.	253	Simulia=Simulium	41
selene Ost. Sack.	261	SIMULIIDAE	41
semifasciata Brun.	218	Simulium Latr.	41
semiflava Kieff.	55, 56	sinabangensis de Meij.	202
semiflavus	75	sinensis Theob.	164
semifumata Brun.	34	sinensis annularis Theob.	97
semiglaber Kieff.	67	" W. (<i>Anopheles</i>)	92, 95
semilimpida Brun.	262	singhalensis Rubs.	19
seminiger Kieff.	74	singularis Leices.	144
semipilosus	50	" Skuse	244
semirufa Edw.	191	sirius Alex.	236
semistriata Brun.	218	sitiens Theob. nec W.	161
semiviridis Kieff.	77	" W nec Theob.	159
semperi Ost. Sack.	226	Skeiromyia Leices.	181
senile Brun.	43	Skusea Theob.	142
separatus Leices.	113	skusii Giles	158
sepositus	164	smithii Felt	14
7-guttata Theob.	150, 151	socius Walk.	71
7-notata Edw.	219	solita	19
7-punctata Edw.	150, 157	sollicitans Walk.	131
sergenti Theob.	101	somaliensis Nev. Lem.	159
serrata Wulp	197	sorbillans W.	217
serraticornis Brun.	192	sordida Brun. (<i>Claduroides</i>)	251
serratipes Beck.	170	" " (<i>Dicranomyia</i>)	215
serratus Theob.	134	sordidus Kieff.	79, 273
sericata Mg.	212	spathifurca Edw.	152
sericeus Theob.	158	spathipalpis Rond.	170
serricornis Brun. (<i>Pachyrhina</i>)	207	spatholobi Felt	269
" " (<i>Tipula</i>)	200	spathulata Brun. (<i>Gnoriste, Palao-</i> <i>anaclinia</i>)	29
Serromyia Meg. in Mg.	43, 55	" " " (<i>Leia</i>)	32, 271
sesami Felt	11	spatulicornis Kieff.	65
sessilis Kieff.	78	speciosa Edw.	191
setiger	51	speciosus Kieff.	72
setilineata Brun.	21	spectralis Brun. (<i>Amalopsis</i>)	251
setosicornis Kieff.	63	" " " (<i>Euryschalis</i>)	34
setosipennis	48	speculata de Meij.	208
" " (<i>Atrichopogon</i>)	48	speiseri Edw.	243
" " (<i>Jenkinsia</i>)	45	spenceri Theob.	132
" " (<i>Tanytarsus</i>)	67	Sphaerionotus de Meij.	275
setulosus Dol.	168	Sphaeromyias Curt.	43, 54
6-lineata Theob.	128	spincornis Brun.	82
6-punctatus Kieff.	77	spinigera Kieff.	58
6-setosa Brun.	21	spinosa Ost. Sack.	258
shirakii Edw.	203	splendens Brun.	200
Siagona Mg.	212	" de Meij.	178
sigillatus Kieff.	70	" W.	116
signata Brun.	246, 247	splendida Brun.	242
" de Meij. (<i>Agastomyia</i>)	275	spurius Kieff.	48
" " (<i>Epiphragma</i>)	255		
signaticollis Wulp.	231		

	Page		Page
Spyloptera Rond. ..	231	sunda Alex. ..	203
squamatipes Kieff. ..	12	superbus Kieff. ..	78
squamiger Coq. ..	134	superpictus Grassi ..	111
squamineris Brun. ..	83	superstes Annand. ..	84
squamipennis Blanch. ..	488	suspirans Ost. Sack. ..	190
" Brun. ..	84	sylvestris Leices. ..	154
squamimipennis Arrib. (<i>squami-</i>		Theob. ..	133, 135, 136
<i>penna, squamipennis</i> Auct. ..	177	" " (<i>Rachisoura</i>) ..	157
Squamomyia Theob. ..	181	sylvicola Curt. ..	210
stantoni Edw. ..	227	Sylvicola Harris ..	265
" Newstead ..	87	symphoremæ Felt ..	267
Stegoconops Lutz. ..	135	Symplecta Mg. ..	249
Stegomyia Theob. ..	124, 142	Symplectomorpha Mik. ..	249
stenoctrus ..	135, 136	taeniarostris Theob. ..	124, 132
Stenoscutus ..	135	taeniata Leices. (<i>Lepidotomyia</i>) ..	140
stephensi List. ..	106, 107	" " (<i>Lophoceratomyia</i>) ..	154
stercoraria de Geer ..	65	taeniatus ..	144
Stethomyia Theob. ..	93, 273	" " (<i>Aedes</i>) ..	144
Stibadocera Ender. ..	211	" " (<i>Aioretomyia</i>) ..	144
Stibadocerella Brun. ..	211	" " W. ..	125
stictica Mg. ..	249	taenionota Mg. ..	237
Stictodiplosis Kieff. ..	13	taeniorhynchoides Chris. ..	135, 141
stictogaster ..	78	Taeniorhynchus Arrib. ..	142, 147
stictonota ..	54	taeniorhynchus W. ..	133, 147, 148
stigmatica ..	191	taitana Kieff. ..	53, 272
" Ost. Sack. ..	239	tainanus ..	79, 273
stigmosa de Meij. ..	276	Tanyderus Phil. ..	186
Stilobezzia Kieff. ..	54	TANYPINAE ..	59
stimulans de Meij. ..	47	Tanypremna Ost. Sack. ..	198
" Walk. ..	157	Tanyptera Latr. ..	137
strenua Brun. ..	246	Tanypus Mg. ..	59
Streptodiplosis Kieff. ..	13	Tanytarsus Wulp ..	66
striata Mg. ..	28	Taphrophila Rond. ..	232
striatipennis Brun. ..	201	Taphrosia ..	240
" Kieff. ..	70, 74	taprobanes Walk. ..	189
striatum Brun. ..	43, 271	tarsalis Newstead ..	135
strigipennis de Meij. ..	191	tatayensis Banks ..	166
strigivena Walk. ..	225, 276	tenax de Meij. ..	172
striocera Giles ..	129	" Leices. <i>nec</i> Theob. ..	164
striopleura Edw. ..	222	" Theob. ..	132
Stygeropis Lw. ..	192	tenella de Meij. ..	213
stygius Skuse ..	58	" Mg. ..	240
Styringomyia Lw. ..	244	tenera Ost. Sack. ..	241
subfasciata Walk. ..	257	tenerrimus Kieff. ..	74
subfascipennis Brun. ..	214	tenuicrus ..	66
subfusca Edw. ..	238	tenuiforceps Kieff. ..	80
sublimatus Kieff. ..	48	tenuipes Brun. ..	201
subpictus Grassi ..	98	" Ost. Sack. ..	251
substituta Walk. ..	248	tenuis Brun. ..	263
subte sellata Brun. ..	215	" Walk. ..	206
subtineta Brun. (<i>Erioptera</i>) ..	238	tenuispatha Kieff. ..	9
" " (<i>Tipula</i>). ..	201	tenuitarsis Kieff. ..	71
subulifer Dol. ..	117, 118	tergorata Rond. ..	38
" Giles ..	117	terminalis Brun. ..	191
subumbrosa Theob. ..	107	" Walk. ..	253
subvarians Walk. ..	38	termitina Ost. Sack. ..	226
sudanensis, Theob. ..	120	Teromyia Leices. ..	116
suffusa Brun. ..	35	terrigans Walk. ..	155
suffusinervis Brun. ..	25	tessellatipennis Brun. ..	200
sugens W. ..	125	tessellatum Theob. (<i>lapsus</i>) ..	105
suknaensis Theob. ..	136	tessellatus ..	95
sulcata Wulp ..	19	" " " (<i>Anopheles</i>) ..	105
sumatrana Edw. ..	227	testacea Brun. ..	264
" Ender. ..	22	" Theob. ..	174
sumatranus ..	196	testaceus Lw. ..	204
sumatrensis Macq. ..	260	tetraclada Kieff. ..	48, 49
sumptuosus Kieff. ..	78	tetrasticta ..	53

	Page		Page
Spyloptera Rond. ..	231	sunda Alex. ..	203
squamatipes Kieff. ..	12	superbus Kieff. ..	78
squamiger Coq. ..	134	superpictus Grassi ..	111
squaminnervis Brun. ..	83	superstes Annand. ..	84
squamipennis Blanch. ..	488	suspirans Ost. Sack. ..	190
" Brun. ..	84	sylvestris Leices. ..	154
squamimipennis Arrib. (<i>squami-</i>		" Theob. ..	133, 135, 136
<i>penna, squamipennis</i> Anct. ..	177	" " (<i>Rachisoura</i>) ..	157
Squamomyia Theob. ..	181	sylvicola Curt. ..	210
stantoni Edw. ..	227	Sylvicola Harris ..	265
" Newstead ..	87	symphoremæ Felt ..	267
Stegoconops Lutz. ..	135	Symplecta Mg. ..	249
Stegomyia Theob. ..	124, 142	Symplectomorpha Mik. ..	249
stenoetrus ..	135, 136		
Stenoscutus ..	135	taeniarostris Theob. ..	124, 132
stephensi List. ..	106, 107	taeniata Leices. (<i>Lepidotomyia</i>) ..	140
stercoraria de Geer ..	65	" " (<i>Lophoceratomyia</i>) ..	154
Stethomyia Theob. ..	93, 273	taeniatus ..	144
Stibadocera Ender. ..	211	" " (<i>Aioretomyia</i>) ..	144
Stibadocerella Brun. ..	211	" " W. ..	125
stictica Mg. ..	249	taenionota Mg. ..	237
Stictodiplosis Kieff. ..	13	taeniorhynchoides Chris. ..	135, 141
stictogaster ..	78	Taeniorhynchus Arrib. ..	142, 147
stictionota ..	54	taeniorhynchus W. ..	133, 147, 148
stigmatica ..	191	taitana Kieff. ..	53, 272
" Ost. Sack. ..	239	tainanus ..	79, 273
stigmata de Meij. ..	276	Tanyderus Phil. ..	186
Stilobezzia Kieff. ..	54	TANYPINAÆ ..	59
stimulans de Meij. ..	47	Tanypremna Ost. Sack. ..	198
" Walk. ..	157	Tanyptera Latr. ..	187
strenua Brun. ..	246	Tanyptus Mg. ..	59
Streptodiplosis Kieff. ..	13	Tanytarsus Wulp ..	66
striata Mg. ..	28	Taphrophila Rond. ..	232
striatipennis Brun. ..	201	Taphrosia ..	240
" Kieff. ..	70, 74	taprobanes Walk. ..	189
striatum Brun. ..	43, 271	tarsalis Newstead ..	135
strigipennis de Meij. ..	191	tatayensis Banks ..	166
strigivena Walk. ..	225, 276	tenax de Meij. ..	172
striocrura Giles ..	129	" Leices. <i>nee</i> Theob. ..	164
striopleura Edw. ..	222	" Theob. ..	132
Stygeropis Lw. ..	192	tenella de Meij. ..	213
stygius Skuse ..	58	" Mg. ..	240
Styrgomyia Lw. ..	244	tenera Ost. Sack. ..	241
subfasciata Walk. ..	257	tenerrimus Kieff. ..	74
subfascipennis Brun. ..	214	tenuicrus ..	66
subfusca Edw. ..	238	tenuiforceps Kieff. ..	80
sublimatus Kieff. ..	48	tenuipes Brun. ..	201
subpictus Grassi ..	98	" Ost. Sack. ..	251
substituta Walk. ..	248	tenuis Brun. ..	263
subte sellata Brun. ..	215	" Walk. ..	266
subtineta Brun. (<i>Erioptera</i>) ..	238	tenuispatha Kieff. ..	9
" " (<i>Tipula</i>). ..	201	tenuitarsis Kieff. ..	71
subulifer Dol. ..	117, 118	tergorata Rond. ..	38
" Giles ..	117	terminalis Brun. ..	191
subumbrosa Theob. ..	107	" Walk. ..	253
subvarians Walk. ..	38	termitina Ost. Sack. ..	226
sudanensis, Theob. ..	120	Teromyia Leices. ..	116
suffusa Brun. ..	35	terrigans Walk. ..	155
suffusinervis Brun. ..	25	tessellatipennis Brun. ..	200
sugens W. ..	125	tessellatum Theob. (<i>lapsus</i>) ..	105
suknaensis Theob. ..	136	tessellatus ..	95
sulcata Wulp ..	19	" " " (<i>Anopheles</i>) ..	105
sumatrana Edw. ..	227	testacea Brun. ..	264
" Ender. ..	22	" Theob. ..	174
sumatranus ..	196	testaceus Lw. ..	204
sumatrensis Macq. ..	266	tetraclada Kieff. ..	48, 49
sumptuosus Kieff. ..	78	tetrasticta ..	53

	Page		Page
Teucholabis Ost. Sack.	232, 234	trigonia Edw.	222
textor Kieff.	9	trilineata Brun. (<i>Antocha</i>)	232
Thalassomyia Sch. Johann	66, 272	" " (<i>Apistomyia</i>)	36
theobaldi Giles	101	" " (<i>Mycomyia</i>)	27
Theobaldia Nev. Lem.	169	" " (<i>Pselliophora</i>)	188
Theobaldinella Blanch.	169	" " (<i>Sciara</i>)	19
Theobaldiomyia Brun.	157	" Leices (<i>Uranotaenia</i>)	174
theobromae Kieff.	49, 272	" " in Theob.	129, 274
thibetana de Meij.	198	trilineatus Theob.	137, 138
thomae L.	18	triobata Kieff.	9
Thomasina Newst. and Carter	157	trimaculata Brun.	221
thomsoni Theob.	129	" Taylor	179
thoracica Guer.	38	trimaculatus Theob.	165
thorntoni Ludl.	105	trinotata Kieff.	50
Thrypticomomyia Skuse	212	tripartita Walk.	206
thwatesiana Westw.	224, 225	tripartitum Kieff.	68
tibialis Brun.	186	Tripelina (lapsus = Tripelma)	80
" Mg.	66	Tripelma Kieff.	80, 273
tigripes Grandpré and Charmay	161	triplasia Wulp	206
tinctinervis Brun.	221	tripudians Bezzi	192
Tinearia Schlubrg.	81	tripunctata de Meij.	276
tinctipennis Brun.	27	" F.	220
" de Meij.	216	" Theob.	130
Tinoestés (Tinoletes) Coq.	156	tripunctipennis Brun.	265
Tipula L.	196	triseriatus Say	135
TIPULIDAE	186	tresignata Walk.	253
tipuliformis Theob.	164	trispinosa Kieff.	57
" Leices.	180	tristicta	50
TIPULINAE	187	tristis Wulp	39
TIPULINI	193	tristissima Ost. Sack.	246
Tipulodina Ender.	196	tritaeniorhynchus Giles	161
titillans Walk.	147, 148	tritincta Brun.	222
tjibodensis Alex.	203	trivittata Mg.	29
Topomyia Leices.	179	trivittata Coq.	134
Toxorhina Lw.	230	trochanteratus Thoms.	72
Toxorhynchites Theob.	116	tropica de Meij.	202
TOXORHYNCHITINAE	90	" Dol.	28
toxorhynchus Macq.	125	tryonii Skuse	31
transversa Brun.	82	tuberculifera Edw.	262
transversalis Kieff.	68	turkhudi List.	103
transversalis de Meij.	227	turrita de Meij.	22
travancorica Annand.	84		
treacherii Leices	110, 113	uichancoi Felt	268, 269
tremula Theob.	142	" Pettey	23
Trentepohlia Big.	242	Ula Hal.	256
trentepohlil W.	242, 243	ulocoma Theob.	183
triangularis Brun. (<i>Eriocera</i>)	264	umbrata de Meij.	213
" (<i>Limnobia</i>)	221	umbrinoides Alex.	203
triangulata Ludl.	173	umbrinus W.	193
Trichocera Mg.	250	umbrosa Theob.	107
TRICHO CERINAE Kert.	251	umbrosus Edw.	247
Trichocladus Kieff.	65	umbrosus Theob.	111
Trichocantarina Kieff.	270	uncus Theob.	143
Trichomegalosphys Ender.	23	undulata Winn.	28
Trichoperrisia Kieff.	10	uniannulata Theob.	135
Trichopronomyia Theob.	156	unicolor Brun. (<i>Pericoma</i>)	84
TRICHO PROSOPONINA	91	" " (<i>Rhamphidia</i>)	228
Trichoptera Mg.	81	" de Meij.	264
trichopus Thoms.	46	" Hal.	217
Trichorhynchomyia Brun.	151	uniformis Leices. (<i>Culex</i>)	166
Trichorhynchus Theob.	151	" " (<i>Melanocconion</i>)	169
Trichosticha Sch.	239	" Theob. (<i>Lophoceratomyia</i>)	153
Trichotanypus Kieff.	62, 272	" " (<i>Panoplites</i>)	150
Tricontanina Kieff.	270	" " (<i>Skusea</i>)	145
Tricyphona Zett.	250	unilineata Brun.	232
tridens Brun.	219	" Leices.	174
trifasciata Brun.	19	" Theob.	125
" Walk.	32	unimaculata Leices.	175

	Page		Page
univittatus Theob. ..	163	viridis Kieff. ..	67
Uranotaenia Arrib. ..	173	" Westw. ..	257
URANOTAENIAE ..	91	viridiventer Giles ..	151
urbicola Kieff. ..	50	viridiventris Kieff. (<i>Chironomus</i>)	77
		" " (<i>Sphaeromyias</i>)	54
vaga Kieff. ..	57	virilis Leices. ..	144
vagans Edw. ..	135	vishnui Theob. ..	162
" Theob. nec Wied. ..	135	vitea Felt ..	268
" Wied. nec Theob. ..	160	viticola Kieff. and Leeuwen ..	11
vagus Don. ..	98	vitis Rubs. ..	12
vanus Walk. ..	96	vitripennis Brun. ..	221
varia Mg. ..	236	vittata ..	81
variata Leices. var. of fraudatrix		vittifrons Walk. ..	200
Theob. ..	153	vittigera Brun. ..	31
variegata Dol. ..	126	vulgaris Kieff. ..	61
variegatus Giles ..	127		
" Schrk. ..	169	w-alba Theob. ..	129
varietas Leices. ..	142, 144	waldii Blanch. ..	119
varicornis Kieff. ..	79	walkeri Brun. ..	198
varicrus ..	49	" Curt. ..	29
variipes ..	63	" Theob. ..	131
varipes Walk. ..	19	walkeriana Felt ..	12
velata Ender. ..	23	watsoni Leices. ..	113
velutina Walk. ..	259	wellingtonianus Alcock. ..	114
" Wulp ..	190	westwoodi Brun. ..	230
venosa Brun. ..	242	" Ost. Sack... ..	229
ventralis Walk. I (<i>Culex</i>)	122	whitmorei Giles ..	165
" " II (<i>Culex</i>)	168	willmori James in Theob. nec	
venusta Lw. ..	245	Leices. ..	111, 112
" Mg. ..	56	" Leices. nec James in	
" Ost. Sack. ..	238	Theob. ..	104
" Walk. (<i>Platyura</i>)	25	winthemi Lehm. ..	31
" " (<i>Tipula</i>)	196	Worcesteria Banks ..	116
venustus W. ..	71	Wulpiella Kieff. ..	64
vernoniae Felt ..	268	Wyeomyia Theob. ..	182
Verrallina Theob. ..	142	WYEOMYINAE ..	91
verrucosus Kieff. ..	70		
versteegi de Meij. ..	224, 276	Xanthocoma Kieff. ..	52
verticalis Wied. ..	258, 259	xanthogaster Ender. ..	23
vexans de Meij. ..	49	xanthomelana Walk. ..	188, 192
" Mg. ..	135, 136	xanthophila Kieff. ..	50
viburni Felt ..	13	xanthopus ..	47
vicaria Walk. ..	197	xanthopyga de Meij. ..	264
vicarius ..	72	Xiphodiplosis Felt. ..	16
vicina Brun. (<i>Epiphragma</i>)	256	Xiphura Brulle ..	187
" " (<i>Platyura</i>)	25	Xylocrypta Kieff. ..	54
" Edw. ..	182		
vilis Walk. (<i>Glenophora</i>)	257	yerburyi Edw. ..	145
" " (<i>Tipula</i>) ..	197	Yposataea Rond. ..	18
vinaceobrunnea Brun. ..	218		
vincenti Laveran. ..	104	Zalepidota Rubs ..	268
violaceopennis Kieff. ..	60	zammittii Theob. ..	135
violaceus Brun. ..	234	Zeugnomyia Leices. ..	180
virescens Phil. ..	217	zeylanicus Annand. ..	87
virgata Coq. ..	207	zonatipes Walk. ..	126
virgo Zett. ..	231	Zygoneura Mg. ..	18, 23
viridifrons Walk. ..	125		

591-5405
ANN



RECORDS
OF THE
INDIAN
MUSEUM



VOL. XVII
1920

BOOK NO.

5015

ZOOLOGICAL
GARDEN
CALCUTTA