



Smithsonian Institution
United States National Museum



Bulletin 220

Type Specimens of
Reptiles and Amphibians in
The United States
National Museum

By DORIS M. COCHRAN
*Curator of Reptiles and Amphibians
United States National Museum*



WASHINGTON, D.C. • 1961

SMITHSONIAN INSTITUTION

UNITED STATES NATIONAL MUSEUM



BULLETIN 220

WASHINGTON, D.C.

1961

Type Specimens
of Reptiles and Amphibians
in the U.S. National Museum

By DORIS M. COCHRAN
Curator of Reptiles and Amphibians
United States National Museum



Publications of the United States National Museum

The scientific publications of the United States National Museum include two series, *Proceedings of the United States National Museum* and *United States National Museum Bulletin*.

In these series are published original articles and monographs dealing with the collections and work of the Museum and setting forth newly acquired facts in the fields of Anthropology, Biology, Geology, History, and Technology. Copies of each publication are distributed to libraries and scientific organizations and to specialists and others interested in the different subjects.

The *Proceedings*, begun in 1873, are intended for the publication, in separate form, of shorter papers. These are gathered in volumes, octavo in size, with the publication date of each paper recorded in the table of contents of the volume.

In the *Bulletin* series, the first of which was issued in 1875, appear longer, separate publications consisting of monographs (occasionally in several parts) and volumes in which are collected works on related subjects. *Bulletins* are either octavo or quarto in size, depending on the needs of the presentation. Since 1902 papers relating to the botanical collections of the Museum have been published in the *Bulletin* series under the heading *Contributions from the United States National Herbarium*.

This work forms number 220 of the *Bulletin* series.

REMINGTON KELLOGG,
Director, United States National Museum.

UNITED STATES
GOVERNMENT PRINTING OFFICE
WASHINGTON : 1961

Contents

	Page
Introduction	vii
Class Amphibia	3
Order Gymnopiona	3
Order Caudata	4
Order Salientia	29
Class Reptilia	81
Order Squamata	81
Suborder Sauria	81
Suborder Serpentes	157
Order Testudines	227
Order Loricata	236
Index	237

Introduction

As biologists know, an important function of a natural history museum is the preservation of those specimens from which new forms of life are described. Many museums have collections of such type material, the National Collection, in the United States National Museum, being among the largest. The value of such specimens is equally obvious since they serve as a standard with which later field collections can be compared, in order to validate the original claim to novelty and to establish the range and variation of the form.

A published list of the type material in the national collection of reptiles and amphibians is especially useful because of the intensive study being given these forms today. The present list gages the extent of the collection and makes known what is available to researchers; it provides a reference to the sources containing the names and original descriptions of the forms represented; and it complements similar lists issued by other museums. Some of these lists are:

- BARBOUR, T., and LOVERIDGE, A. Typical reptiles and amphibians. *Bull. Mus. Comp. Zool.*, vol. 69, pp. 205-360, 1929; supplement, vol. 96, pp. 59-214, 1946.
- GUIBÉ, J. Catalogue des types d'amphibiens de Muséum d'Histoire Naturelle, 71 pp., 1949.
- LEVITON, A. E. Catalogue of the amphibian and reptile types in the Natural History Museum of Stanford University. *Herpetologica*, vol. 8, pp. 121-132, 1953; supplement by A. E. Leviton and B. H. Banta, vol. 12, pp. 213-219, 1956.
- MARX, H. Catalogue of type specimens of reptiles and amphibians in the Chicago Natural History Museum. *Fieldiana: Zoology*, vol. 36, pp. 409-496, 1958.
- PETERS, J. A. Catalogue of type specimens in the herpetological collections of the University of Michigan Museum of Zoology. *Occ. Pap. Mus. Zool. Univ. Michigan*, No. 539, 55 pp., 1952.
- SLEVIN, J. R., and LEVITON, A. E. Holotype specimens of reptiles and amphibians in the collection of the California Academy of Sciences. *Proc. California Acad. Sci.*, series 4, vol. 28, pp. 529-560, 1956.

The national collection had its beginnings over a century ago. A brief review of its history, prior to indicating the content and arrangement of this list, seems appropriate.

James Smithson's legacy for the establishment of an institution "for the increase and diffusion of knowledge among men" resulted in Congress' passing a bill in 1846 for the incorporation of the Smithsonian Institution. With the appointment of Dr. Spencer Fullerton Baird as Assistant Secretary in 1850, announcement was made of the gathering of a collection of natural history objects. Dr. Baird's personal collection, deposited in the Smithsonian Institution in that year, was the forerunner of numerous other collections that began to arrive soon afterward. The Smithsonian encouraged such donations by distributing instructions on how to collect and preserve natural history materials and by sponsoring or assisting various exploring expeditions. Reptiles and amphibians were represented in these early collections. As the materials were received and cataloged, they were assigned to specialists for identification and description, and duplicate material received was given to or served as the basis of exchange with other institutions at home and abroad.

Much of this early material collected and deposited in the Smithsonian Institution issued from the 19th-century exploring expeditions abroad, and from the land surveys of the West. Perhaps earliest of these was the U.S. Exploring Expedition, commanded by Charles Wilkes, which sailed on its round-the-world voyage in 1838, touching at African, South American, Antarctic, Asian, and western North American coasts. Natural history objects were collected near the coast at every stop. In 1842 the expedition returned with most of its collections intact. The materials, at first deposited in the U.S. Patent Office, were in 1857, along with specimens collected by other expeditions, turned over to the Smithsonian Institution. Many of the animals secured on this journey represented forms as yet unknown to science and hence were in need of a scientific name and a description. Dr. Baird and his friend and colleague, Dr. Charles Girard, wrote several papers on the herpetological results of this expedition. Dr. Girard's later book and atlas became classics.

The La Plata Expedition, under the command of Thomas Jefferson Page, from 1853 through 1856 explored and mapped the tributaries of the La Plata River and the adjacent countries of Argentina, Paraguay, Brazil, and Bolivia. Zoological collections were made, and subsequent study yielded additional new forms of animal life.

In 1853 the U.S. Army Engineers, directed to find a "practicable and economical" route for the first railroad to be laid from the Mississippi to the Pacific, surveyed five possible routes between the 32d and 49th parallels. Collections of animals, plants, and minerals were made at every opportunity, and these excellent representations of the natural history of a vast part of the West eventually came to be deposited in the Smithsonian Institution.

In 1854 began the work of the commission headed by Major William H. Emory to survey lands to establish a permanent U.S.-Mexican boundary. The reptiles and amphibians of the boundary were listed by Dr. Baird in 1859, following earlier preliminary descriptions of new species by him and others.

Many new forms were described from "Indianola," a town no longer existing but formerly located in what is now Calhoun County, Texas. Collecting was equally abundant in some other regions visited by boundary surveyors. Among the members of the various groups, the names of J. H. Clark and D. N. Couch are perhaps more often found than others, a fact indicating their ability first to track down and catch the smaller, less conspicuous creatures, then to preserve, label, and dispatch them in boxes for the long journey east.

The geographical and geological explorations and surveys of lands west of the 100th meridian, begun in 1869 and continued through 1879, each year followed a different route. In order to scour the areas of greatest promise and to estimate rainfall, soil conditions, temperatures, and other natural conditions that might influence the success or failure of future agricultural ventures, the parties were subdivided. Some who went with these expeditions later acquired considerable fame in science—among others, Coues, Yarrow, and Cope. Many of the new species secured by these men were later also named and described by them.

Institutions and persons in foreign lands donated or exchanged many specimens. The Jardin des Plantes of Paris in 1858 sent specimens that had been collected in the Old World and that had been studied by Duméril and Bibron, two of the foremost herpetologists in Europe at the time. Part of the Bonaparte collection, received in 1869, contains European material still valuable for comparison. Felix Poey, a Cuban scholar and naturalist of renown, in 1856 began sending reptiles and amphibians of Cuba, and continued to do so throughout his life.

Much collecting was the work of enthusiastic amateurs from the United States wishing to fill in the blank spaces on the faunal maps of the world. Such collecting continues today. John Xantus de Vesey (J. Xantus), U.S. Consul stationed in Baja California in 1859, and in Colima in 1863, was a noteworthy collector. One of the defects of many 19th-century collections was the frequent lack of precise locality data, and Xantus' material from Mexico is worthy of mention for the care he took in numbering every specimen and entering the locality for each number in one of his numerous letters to officials at the Smithsonian.

W. L. Abbott of Philadelphia, a world traveler and sportsman, early in life took a keen interest in securing all kinds of exotic animals for the national collection. From 1893 until his death in 1937, he worked the lesser-known parts of Asia, the East Indies, Africa, and the West Indies. Many new forms were discovered and named from his material.

The National Geographic Society during the 20th century has financed many rewarding collecting trips to all parts of the globe to enrich still further our knowledge of the fauna of little-traveled regions. The medical research teams of the Armed Services continue to send fine collections from the areas in which they are stationed. Much of this recently deposited material has

not yet been studied but it, too, will probably yield its share of new forms. Some of my colleagues have also deposited for permanent safekeeping in the national collection specimens that they have named and described.

The Smithsonian Institution was fortunate in its early years in being aided by many brilliant scientists who were also herpetologists. First among these was Dr. Baird, who in 1850 was appointed Assistant Secretary of the Smithsonian Institution and in 1878 its Secretary, serving in that capacity until his death in 1887. In 1878 he appointed Dr. Henry C. Yarrow as the first curator of the Division of Reptiles and Amphibians. In addition to papers on reptiles and amphibians collected during the surveys west of the 100th meridian, in which he had been a participant, Yarrow wrote the "Check List of North American Reptilia and Batrachia, With a Catalogue of Specimens in the U.S. National Museum."¹

The second curator of the division, Dr. Leonhard Stejneger, was appointed in 1889 and worked until his death in 1943 at the age of 93. He had acquired world fame as an ornithologist and mammologist before entering the field of herpetology, and it was not long before his influence was felt here also. His "Herpetology of Japan and Adjacent Territory"² and his "Poisonous Snakes of North America"³ are models of careful scientific writing. The several editions of the "Check List of North American Amphibians and Reptiles" that he published in collaboration with Dr. Thomas Barbour have been essential in every herpetological library since the first one appeared in 1917.

Special mention should also be made of Edward Drinker Cope, who worked intermittently with the national collection from 1860 until his death in 1897. He described hundreds of new forms while doing brilliant work on anatomical relationships and on faunal distribution. He became one of the foremost American scientists of his day in herpetology, ichthyology, and paleontology and was noted for his perception of the finer points of specific variation, which had been largely disregarded before his time. Many of his 1,400 scientific papers were published by the Smithsonian Institution. One of his last papers, "The Crocodylians, Lizards, and Snakes of North America,"⁴ is indispensable to every herpetologist. His "Batrachia of North America"⁵ is now a collector's item.

Curators at the National Museum, realizing early that type specimens must be preserved for comparison with material still to come, have sought to maintain the types in the best possible condition, and to safeguard the

¹ U.S. National Museum Bulletin 24, pp. v+249, 1882.

² U.S. National Museum Bulletin 58, pp. xx+577, figs. 427, pls. 35, July 22, 1907.

³ Report of the United States National Museum for the year ending June 30, 1893, pt. 2, No. 1, pp. 339-487, figs. 70, pls. 19, 1895.

⁴ Report of the United States National Museum for the year ending June 30, 1898, pt. 2, pp. 153-1270, figs. 347, pls. 36, 1900.

⁵ U.S. National Museum Bulletin 34, pp. 525, figs. 119, pls. 86, 1889.

specimens have placed them in airtight jars in light-proof rooms. Although the patterns of some of the older type specimens have faded, they are still preserved very distinctly on many. With the passage of time, a few of the types have softened and disintegrated despite constant care, but instances of such loss are becoming fewer as the techniques for preserving material become better understood.

Some of the early collections, especially those from the western United States, were labeled poorly or not at all, because there were few towns or permanent settlements by which to fix the locations of the collecting sites. Often an expedition on the move had to carry its biological material in crowded boxes, which were then shipped east as opportunity offered, sometimes without any field journals or notes to indicate the route along which the specimens had been collected. Thus the specimens collected during a whole expedition might be credited in subsequent museum cataloging to the point from which they had been shipped, while their actual place or places of origin might have been several hundred miles distant. Such errors in recording type localities have usually been cleared up by later collecting and study by specialists, and in the following list the corrections are given.

The first specimens received by the Smithsonian Institution were not always cataloged individually. Often many examples presumed to be of a single species and received from the same locality and collector were all given the same catalog number. Later, when it was found necessary to refer to specific individuals from the lot, they were renumbered. This practice led to much confusion because the old label was frequently almost illegible, and thus correct data were not always inscribed after the new entry. When Dr. Stejneger took charge of the collection in 1889, and beginning with catalog number 14856, a single number was assigned to each specimen (except eggs and tadpoles), a change that has simplified our bookkeeping considerably and has allowed a particular specimen always to be indicated without possibility of confusion. An O in front of our catalog number indicates that the specimen was once a part of the collection of the Division of Comparative Anatomy, which functioned from 1885 to 1914. Most specimens so marked are skeletons, skulls, or dry skins.

Most of the type material is preserved in alcohol. No special mention is made in this list of the condition of each type specimen. It may be assumed to be good or fair unless a statement to the contrary is made.

Scientific material that has been studied by many scientists becomes valuable because it exemplifies the different interpretations that have been placed upon it. In this respect the national collection of reptile and amphibian types is one of the foremost because many of the herpetologists of the past century have done extensive research here. The considerable number of type names listed in the following pages is a tribute to their industry as well as to the extent and value of the collection itself.

All known paratypes are given in the list, because a paratype should be available to take the place of a lost or damaged holotype. If no formal designation of paratypes was made in the original description, I considered the "specimens examined" cited in the description as paratypes. In some instances I received the sanction of the describers of such species for this action. L. M. Klauber, Hobart M. Smith, and other herpetologists agree with me that any specimens used by an author and recognized by him as belonging to the species when he is describing it are actually paratypes, although we recognize his right to name only certain ones as paratypes in his original description if he so desires. Another reason for listing paratypes is to encourage their exchange with other museums. Common sense indicates the need for as wide a dispersal as possible of critical material where it may be most useful. Many museums, including the U.S. National Museum, are willing to trade when they have several paratypes of any given form.

For the spelling of geographic names, I referred to the "U.S. Directory of Post Offices," the "Columbia Lippincott Gazetteer of the World," the U.S. Board on Geographic Names, and the American Geographical Society's "Map of Hispanic American." If the locality data as given in the original description are too general or obviously in error, emendations and corrections are added in parentheses.

The list contains 1,742 names of types. The original generic, specific, and subspecific names followed by the name of the describer appear on the first line. I then cite the publication where the name and description first appeared. If the original name of the genus or species is no longer accepted, I give on the third line the current synonym in italics. Finally I give a complete listing of museum number, locality, collector, and date collected for the holotype, allotype, neotype, cotypes, paratypes, and lectotypes.

The accepted definitions of these terms are:

- | | |
|-----------|--|
| Holotype: | The designated single specimen on which the original description is based. |
| Allotype: | The designated single specimen of the sex opposite to that of the holotype. |
| Neotype: | The specimen chosen to take the place of the lost or destroyed holotype but not belonging to the original series of specimens used by the author. |
| Cotype: | One of several specimens used by the describer in preparing his original description, in which he did not designate a specific individual as a holotype. |
| Paratype: | The other specimen (or specimens) used by the author and recognized as belonging to the same species when he clearly designates one particular specimen as the holotype or "type." |

Lectotype: One of the cotypes that may be selected by a later writer as having been the specimen on which the original description was based.

If there is more than one cotype or paratype, I give the total number of each in parentheses.

The genera and species are listed alphabetically under the following classes, orders, and suborders:

<i>Classification</i>	<i>Number of names</i>
Class Amphibia	
Order Gymnopiona (caecilians)	12
Order Caudata (salamanders)	176
Order Salientia (frogs)	409
Class Reptilia	
Order Squamata	
Suborder Sauria (lizards)	558
Suborder Serpentes (snakes)	524
Order Testudines (turtles)	62
Order Loricata (crocodilians)	1

Students may wish to know their division into families, as follows:

<i>Families</i>	<i>Species</i>	<i>Families</i>	<i>Species</i>
Agamidae	10	Hynobiidae	6
Alligatoridae	1	Iguanidae	276
Ambystomidae	39	Kinosternidae	16
Amphisbaenidae	9	Lacertidae	7
Anguidae	29	Leiopelmidae	3
Atelopodidae	5	Leptodactylidae	122
Boidae	24	Leptotyphlopidae	13
Bufonidae	55	Microhylidae	23
Caeciliidae	12	Pelobatidae	8
Centrolenidae	9	Pipidae	2
Chamaeleontidae	2	Plethodontidae	111
Chelonidae	5	Proteidae	7
Chelydridae	1	Ranidae	51
Colubridae	412	Rhacophoridae	13
Crotalidae	46	Salamandridae	8
Cryptobranchidae	1	Scincidae	80
Dendrobatidae	7	Sirenidae	4
Elapidae	17	Teiidae	82
Emydidae	31	Testudinidae	5
Gekkonidae	53	Trionychidae	4
Helodermatidae	1	Typhlopidae	8
Hydrophidae	3	Viperidae	1
Hylidae	111	Xantusiidae	9

When referring to institutions, I use the following abbreviations:

<i>Abbreviation</i>	<i>Institution</i>
AM	Australian Museum, Sydney
AMNH	American Museum of Natural History, New York City
ANSP	Academy of Natural Sciences of Philadelphia, Pennsylvania
BM	Bergen Museum, Norway
BMNH	British Museum (Natural History), London
BYU	Brigham Young University, Provo, Utah
CAS	California Academy of Sciences, San Francisco
CM	Carnegie Museum, Pittsburgh, Pennsylvania
CNHM	Chicago Natural History Museum, Illinois
CPS	College of Puget Sound, Takoma, Washington
DZ	Departamento de Zoologia, Secretaria da Agricultura, São Paulo
FMNH	Field Museum of Natural History (later renamed Chicago Museum of Natural History)
IB	Instituto Butantan, Brazil
IML	Instituto Miguel Lillo, Argentina
JPP	Jardin des Plantes, Paris (affiliate of Muséum Nationale d'Histoire Naturelle, Paris)
LC	Dr. Adolpho Lutz' collection, Instituto Oswaldo Cruz, Rio de Janeiro
MCZ	Museum of Comparative Zoology, Cambridge, Massachusetts
MHNB	Muséum d'Histoire Naturelle Bâle, Switzerland
MHNP	Muséum Nationale d'Histoire Naturelle, Paris
MP	Museum Paulista, São Paulo
MZUM	Museum of Zoology of the University of Michigan, Ann Arbor, Michigan
NOAS	New Orleans Academy of Science, Louisiana
R	Rijksmuseum, Amsterdam
RPSF	Royal Palm State Park, Florida
SCMT	Science College Museum, Imperial University of Tokyo
SDSNH	San Diego Society of Natural History, California
SNCBSH	State of North Carolina Biological Station, Highlands
SNG	Senckenbergische Naturforschende Gesellschaft, Frankfurt-am-Main
SUM	Stanford University Museum, California
SYUM	Sun Yat-sen University Museum, China
TCM	Trowbridge College Museum (Museum of Zoology of the University of Michigan)
TUM	Tulane University Museum, New Orleans, Louisiana

<i>Abbreviation</i>	<i>Institution</i>
UCM	University of Colorado Museum, Boulder, Colorado
UIMNH	University of Illinois Museum of Natural History, Urbana, Illinois
USNM	U.S. National Museum, Washington, D.C.
WCUUM	West China Union University Museum, Chengtu
WIM	Wistar Institute Museum, Philadelphia, Pennsylvania
ZML	Zoological Museum of Leningrad
ZS	Zoologische Staatssammlung, Munich
ZSP	Zoological Society of Philadelphia, Pennsylvania

The names of some collectors have the same initials. In order to distinguish Hobart M. Smith from Hugh M. Smith and for the sake of economy of space, I identify them as H. M. and Hugh, respectively.

Expressions of gratitude are owed to colleagues for their help in checking data—especially Hobart M. Smith for the entire list, Norman Hartweg and John W. Crenshaw for the turtles, Richard Highton for the salamanders, and John M. Condit and Coleman J. Goin for the Hylidae.

**Type Specimens
of Reptiles and Amphibians
in the U.S. National Museum**



Class Amphibia

Order Gymnophiona

***Caecilia isthmica* Cope**

Proc. Amer. Philos. Soc., vol. 17 (1878), p. 91, 1877.

=*Caecilia tentaculata* Linnaeus

HOLOTYPE: 25188, Atlantic side, Isthmus of Darién (Panama-Colombia),
T. O. Selfridge.

***Cæcilia ochrocephala* Cope**

Proc. Acad. Nat. Sci. Philadelphia, vol. 18, p. 132, 1866.

=*Caecilia ochrocephala* Cope

HOLOTYPE: 29764, Panama, J. L. LeConte.

***Hypogeophis alternans* Stejneger**

Proc. U.S. Nat. Mus., vol. 16, p. 739, 1894.

HOLOTYPE: 20418, Mahé, Seychelles Islands (Indian Ocean), W. L.
Abbott, 1892.

PARATYPE: 20404, Seychelles Islands, MHNP.

***Ichthyophis kotoensis* Taylor**

Univ. Kansas Sci. Bull., vol. 40, No. 4, p. 110, 1960.

HOLOTYPE: 72293, Koh Tao, Gulf of Siam (Thailand), Hugh M. Smith,
Jan. 1, 1927.

PARATYPE: 76131, Koh Tao, Gulf of Siam, Hugh M. Smith, Sept. 18, 1928.

***Ichthyophis nigroflavus* Taylor**

Univ. Kansas Sci. Bull., vol. 40, No. 4, p. 101, 1960.

HOLOTYPE: 129462, within 20 miles of Kuala Lumpur, Selangor (Malayan
Peninsula), R. Traub and V. Tipton, April 1950.

***Ichthyophis paucidentulus* Taylor**

Univ. Kansas Sci. Bull., vol. 40, No. 4, p. 49, 1960.

HOLOTYPE: 70671, Kepahiang, Sumatra, H. C. Kellers, Jan. 12, 1926.

***Ichthyophis paucisuleus* Taylor**

Univ. Kansas Sci. Bull., vol. 40, No. 4, p. 103, 1960.

HOLOTYPE: 103565, Siantar, Sumatra, National Geographic Society-
Smithsonian Expedition to Sumatra, 1937.

Ichthyophis sumatranus Taylor

Univ. Kansas Sci. Bull., vol. 40, No. 4, p. 95, 1960.

HOLOTYPE: 70672, Kepahiang, Sumatra, H. C. Kellers, Dec. 10, 1925.

PARATYPES (3): 70667, 70669-70, Kepahiang, Sumatra, H. C. Kellers, Dec. 10, 1925.

Scolecormorphus uluguruensis Barbour and Loveridge

Mem. Mus. Comp. Zool., vol. 50, p. 180, 1928.

PARATYPES (2): 73236-7, Nyingwa, Uluguru Mountains, Tanganyika, 7,000-8,000 ft., A. Loveridge, Oct. 18, 1926.

Siphonops oligozonus Cope

Proc. Amer. Philos. Soc., vol. 17 (1878), p. 91, 1877.

=*Gymnopsis oligozona* (Cope)

HOLOTYPE: 25187, Guatemala?.

Siphonops proximus Cope

Proc. Amer. Philos. Soc., vol. 17 (1878), p. 90, 1877.

=*Gymnopsis multiplicata proxima* (Cope)

HOLOTYPE: 29762, coast of eastern Costa Rica, W. M. Gabb.

PARATYPE: 29763, same data.

Siphonops simus Cope

Proc. Amer. Philos. Soc., vol. 17 (1878), p. 91, 1877.

=*Gymnopsis multiplicata multiplicata* Peters

HOLOTYPE: 29765, Costa Rica, A. von Frantzius.

Order Caudata

Amblystoma annulatum Cope

Proc. Amer. Philos. Soc., vol. 23, p. 525, 1886.

=*Amblystoma annulatum* Cope.

HOLOTYPE: 11564, locality unknown (restricted to vicinity of Hot Springs, Garland County, Arkansas).

Amblystoma aterrimum Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 19 (1867), p. 201, 1868.

=*Dicamptodon ensatus* (Eschscholtz)

HOLOTYPE: 5242, crossing of Bitter Root River, North Rocky Mountains (Montana), S. P. Hildreth and J. Mullan, 1860.

Amblystoma cingulatum Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 19 (1867), p. 205, 1868.

=*Amblystoma cingulatum cingulatum* Cope

NEOTYPE: 129396, Robertsville, Jasper County, South Carolina, W. T. Neill, Nov. 17, 1947.

Amblystoma copeianum Hay

Proc. U.S. Nat. Mus., vol. 8, p. 209, pl. 14, 1885.

=*Ambystoma tigrinum tigrinum* (Green)

HOLOTYPE: 14112, Irvington, Indianapolis, Marion County, Indiana, G. H. Clarke, Apr. 7, 1885.

Amblystoma decorticaum Cope

Proc. Amer. Philos. Soc., vol. 23, p. 522, 1886.

=*Ambystoma gracile decorticaum* Cope

HOLOTYPE: 14493, Port Simpson, British Columbia, Canada, T. H. Streets, Dec. 9, 1885.

Amblystoma lepturum Cope

Proc. Amer. Philos. Soc., vol. 23, p. 524, 1886.

=*Ambystoma cingulatum cingulatum* Cope

HOLOTYPE: 14583, locality unknown (probably United States).

**Amblystoma macrodactyla** Baird

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 1, p. 292, 1850.

=*Ambystoma macrodactylum macrodactylum* Baird

COTYPE: 4042, Astoria (Clatsop County), Oregon, J. K. Townsend.

Amblystoma obscurum Baird*in* Cope, Proc. Acad. Nat. Sci. Philadelphia, vol. 19 (1867), p. 192, 1868.=*Ambystoma tigrinum tigrinum* (Green)

HOLOTYPE: 3994, Fort Des Moines, Polk County, Iowa, W. E. Moore.

Amblystoma paroticum Baird*in* Cope, Proc. Acad. Nat. Sci. Philadelphia, vol. 19 (1867), p. 200, 1868.=*Ambystoma gracile gracile* (Baird)

COTYPES (2): 4708a, Chiloweynek Lake, British Columbia, Canada, A. Campbell, 1859. 7021, Puget Sound (Washington), U.S. Exploring Expedition, 1841.

Amblystoma platineum Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 19 (1867), p. 198, 1868.

=*Ambystoma jeffersonianum* (Green)

COTYPES (2): 3998, 39444 (formerly 3998), Cleveland, Cuyahoga County, Ohio, J. P. Kirtland.

Amblystoma proserpine Baird and Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 173, 1852.

=*Ambystoma tigrinum mavortium* Baird

COTYPES (3): 4082 (3), Salado River, 4 miles east of San Antonio, Bexar County, Texas, L. A. Edwards.

Amblystoma tenebrosum Baird and Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 174, 1852.

=*Dicamptodon ensatus* (Eschscholtz)

HOLOTYPE: 4710, Oregon, U.S. Exploring Expedition, 1841.

Amblystoma trisruptum Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 19 (1867), p. 194, 1868.

=*Amblystoma tigrinum mavortium* Baird

HOLOTYPE: 4068, ♀, Ocate Creek, New Mexico, J. Potts.

Amblystoma xiphias Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 19 (1867), p. 192, 1868.

=*Amblystoma tigrinum tigrinum* (Green)

HOLOTYPE: 14470 (formerly 4135), Columbus, Franklin County, Ohio,
L. Lesquereux.

Amblystoma cingulatum bishopi Goin

Ann. Carnegie Mus., vol. 31, p. 300, pl. 1, figs. 3-5, 1950.

PARATYPES (3): 42861, 57389-90, Mobile, Mobile County, Alabama, J.
Hurter, Mar. 15, 1910.

Amblystoma fluvinatum Taylor

Copeia, No. 3, p. 144, figs. 2a-b, 1941.

PARATYPE: 137205 (formerly UIMNH 27234), Mojárichic, Chihuahua,
Mexico, I. Knobloch.

Amblystoma granulosum Taylor

Univ. Kansas Sci. Bull., vol. 30, p. 57, 1944.

PARATYPES (22): 116633-53, 11-15 km. west of Toluca, Mexico, Mexico,
E. H. Taylor and H. M. Smith, August 1939. 137206 (formerly UIMNH
27230), Km. 74, west of Toluca, Mexico, Mexico, E. H. Taylor and H. M.
Smith, August 1939.

Amblystoma lacustris Taylor and Smith

Proc. U.S. Nat. Mus., vol. 95, p. 532, pl. 18, figs. 1-2, 1945.

HOLOTYPE: 117410, Texcoco, Mexico, Mexico, H. M. Smith, Mar. 2,
1939.

PARATYPES (19): 116736-44, 116746-55, Lake Zumpango, Mexico,
Mexico, H. M. Smith, March-September 1939.

Amblystoma mabeei Bishop

Journ. Elisha Mitchell Sci. Soc., vol. 43, p. 157, pl. 23, figs. 1-2, 1928.

HOLOTYPE: 75058, low grounds of the Black River near Dunn, Harnett
County, North Carolina, W. B. Mabee, May 12, 1923.

Ambystoma mavortia Baird

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 1, p. 292, 1850.

= *Ambystoma tigrinum mavortium* Baird

HOLOTYPE?: 3990, New Mexico, J. L. LeConte (collector said to be Dr. Wislizenus in original description).

Ambystoma nebulosum Hallowell

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 209, 1852.

= *Ambystoma tigrinum nebulosum* Hallowell

HOLOTYPE: 4702a, San Francisco Mountains, Arizona (New Mexico in original description), S. W. Woodhouse.

Ambystoma ordinaria Taylor

Univ. Kansas Sci. Bull., vol. 26 (1939), p. 422, pl. 46, figs. 1-3, 1940.

= *Ambystoma ordinarium* Taylor

PARATYPE: 134276, near Puerto Hondo, Michoacán, Mexico, E. H. Taylor, Sept. 3, 1938.

Ambystoma rosaceum nigrum Shannon

Proc. U.S. Nat. Mus., vol. 101, p. 466, fig. 91, 1951.

HOLOTYPE: 123581, El Salto, Durango, Mexico, R. G. Miller, May 8, 1946.

Ambystoma rosaceum sonoraensis Shannon

Proc. U.S. Nat. Mus., vol. 101, p. 468, 1951.

HOLOTYPE: 17253, larva, Pinetos Camp, Sonora, Mexico, P. L. Jouy, June 14, 1891.

PARATYPE: 17255, larva, same data. 17254 was exchanged to MCZ in 1958.

Ambystoma stejnegeri Ruthven

Proc. U.S. Nat. Mus., vol. 41, p. 517, 1912.

= *Ambystoma macrodactylum macrodactylum* Baird

HOLOTYPE: 48598, Bloomfield, Davis County, Iowa, G. H. Berry, July 27, 1906.

PARATYPE: 48599, same data.

Ambystoma subsalsum Taylor

Copeia, No. 3, p. 152, figs. 1-3, 1943.

PARATYPE: 134277 (formerly UIMNH 27326), Lake Alchichica, Puebla, Mexico, E. H. Taylor and D. M. Forbes, Aug. 14-15, 1942.

Ambystoma tigrinum diaboli Dunn

Copeia, No. 3, p. 160, 1940.

PARATYPE: 13394, Ottawa?, Ontario, Canada, R. Bell, Apr. 21, 1883. This specimen was used as a paratype, but was not so listed.

Ambystoma tigrinum slateri Dunn

Copeia, No. 3, p. 159, 1940.

= *Ambystoma tigrinum melanostictum* (Baird)

HOLOTYPE: 108982, 5 miles southeast of Coulee Dam, Grant County, Washington, J. R. Slater.

Ambystoma tigrinum stebbinsi Lowe

Proc. Biol. Soc. Washington, vol. 67, p. 243, 1954.

PARATYPE: 134131 (formerly Univ. Arizona 3219), J.A. Jones Ranch in Parker Canyon, southwest side of Huachuca Mountains, Santa Cruz County, Arizona, around 5,000 ft., C. H. Lowe, Jr., Nov. 4, 1950.

Anaides ferreus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 21, p. 109, 1869.

= *Aneides ferreus* Cope

HOLOTYPE: 14451 (formerly probably 6794), Fort Umpqua, Oregon, E. P. Vollum.

Batrachoseps catalinae Dunn

Copeia, No. 109, p. 62, 1922.

= *Batrachoseps pacificus* (Cope)

HOLOTYPE: 57335, ♀, Catalina Island, California, J. Hurter, Jan. 23, 1908.

PARATYPES (3): 38362, 57334, 57336, same data. 38361 was exchanged to MCZ.

Batrachoseps caudatus Cope

U.S. Nat. Mus. Bull. 34, p. 126, fig. 27, 1889.

= *Batrachoseps attenuatus attenuatus* (Eschscholtz)

HOLOTYPE: 13561, Hassler Harbor, Alaska, H. E. Nichols, December 1881.

Batrachoseps leucopus Dunn

Copeia, No. 109, p. 60, 1922.

= *Batrachoseps attenuatus attenuatus* (Eschscholtz)

HOLOTYPE: 64319, ♀, North Island, Coronados Islands, Baja California, Mexico, A. W. Armstrong, Aug. 4, 1921.

Batrachuperus karlschmidti Liu

Chicago Nat. Hist. Mus., Fieldiana: Zool. Mem., vol. 2, p. 87, fig. 12, 1950.

PARATYPES (2): 124581-2, Lu-ho-hsien, Sikang, China, C. C. Liu.

Batrachuperus yenyuanensis Liu

Chicago Nat. Hist. Mus., Fieldiana: Zool. Mem., vol. 2, p. 99, figs. 14a-d, 1950.

PARATYPES (2): 124583-4, from a large pool on Peilinshan, between Yenyuanhsien and Yenpiensien, Sikang, China, H. W. Chang and Y. W. Kao.

Bolitoglossa arborea Taylor

Herpetologica, vol. 2, p. 62, figs. 4a-c, 5a-c, 1941.

=*Chiropterotriton arborea* (Taylor)

PARATYPE: 134278 (formerly UIMNH 27078), near Tianguistengo, Hidalgo, Mexico, E. H. Taylor and R. Roberts, Aug. 13, 1938.

Bolitoglossa borburata Trapido

Bol. Soc. Venezolana Cienc. Nat., vol. 8, p. 297, figs. 1-4, 1942.

HOLOTYPE: 115509, Valle del Río Borburata, Venezuela, P. Anduze, March 1940.

Bolitoglossa cephalica rubrimembris Taylor and Smith

Proc. U.S. Nat. Mus., vol. 95, p. 539, 1945.

=*Pseudoeurycea cephalica rubrimembris* (Taylor and Smith)

HOLOTYPE: 110661, ♀, 6 km. south of Santa Anita, Hidalgo, Mexico, about 4,500 ft., H. M. Smith, Oct. 10, 1939.

PARATYPES (2): 110659-60, same data.

Bolitoglossa chondrostega Taylor

Univ. Kansas Sci. Bull., vol. 27, p. 113, 1941.

=*Chiropterotriton chondrostega* (Taylor)

PARATYPE: 134279 (formerly UIMNH 27045), Durango, Hidalgo, Mexico, 5,000-6,000 ft., E. H. Taylor, Sept. 12, 1938.

Bolitoglossa cochranæ Taylor

Univ. Kansas Sci. Bull., vol. 29, p. 343, 1943.

=*Pseudoeurycea cochranæ* (Taylor)

PARATYPES (2): 116396, Cerro San Felipe, Oaxaca, Mexico, H. M. Smith, July 10-20, 1940. 134289 (formerly UIMNH 27183), Cerro San Felipe, Oaxaca, Mexico, E. H. Taylor, 1940.

Bolitoglossa dimidiata Taylor

Univ. Kansas Sci. Bull., vol. 26 (1939), p. 408, figs. 1-2, 1940.

=*Chiropterotriton dimidiata* (Taylor)

PARATYPE: 134280 (formerly UIMNH 27071), El Chico National Park, Hidalgo, Mexico, E. H. Taylor, 1938.

Bolitoglossa galaenæ (sic) Taylor

Proc. Biol. Soc. Washington, vol. 54, p. 83, 1941.

=*Pseudoeurycea galeanae* (Taylor)

PARATYPES (15): 110642-46, 116048-9, 116051-8, 15 miles west of Galeana, Nuevo León, Mexico, 5,200 ft., H. M. Smith, Oct. 13-14, 1939. 110647 was exchanged to MCZ, 1944; 116050 to MZUM, 1944.

Bolitoglossa lavae Taylor

Univ. Kansas Sci. Bull., vol. 28, p. 295, pl. 27, figs. 5-6, 1942.

=*Chiropterotriton lavae* (Taylor)

PARATYPES (3): 134281-3 (formerly UIMNH 27103-5), 2 miles west of La Joya, Veracruz, Mexico, E. H. Taylor et al., Aug. 16-17, 1941.

Bolitoglossa melanomolga Taylor

Proc. Biol. Soc. Washington, vol. 54, p. 81, 1941.

=*Pseudoeurycea melanomolga* (Taylor)

PARATYPES (2): 110640, 17 km. northeast of Limón, Veracruz, Mexico, H. M. Smith, Mar. 23, 1940. 110641, near Teziutlán, Veracruz, Mexico, H. M. Smith, June 25, 1940.

Bolitoglossa moreleti Smith

Herpetologica, vol. 3, p. 14, 1945.

HOLOTYPE: 116079, Palenque, Chiapas, Mexico, Dr. and Mrs. H. M. Smith, July 12, 1939.

PARATYPES (28): 116070, Piedras Negras, Petén, Guatemala, H. M. Smith, May 24, 1939. 116071-8, 116080-98, Palenque, Chiapas, Mexico, H. M. Smith, July 7-23, 1939.

Bolitoglossa nigroflavescens Taylor

Univ. Kansas Sci. Bull., vol. 27, p. 150, pl. 8, pl. 9, figs. 9-10, 1941.

=*Magnadigita nigroflavescens* (Taylor)

HOLOTYPE: 111169, Cerro Ovando, Chiapas, Mexico, 5,000-6,000 ft., H. M. Smith, Apr. 16, 1940.

PARATYPES (33): 111158-9, 111161-8, 111170-85, 111187-92, 134288, same data. 111160, 111186 were exchanged to MZUM in 1944.

Bolitoglossa nigromaculata Taylor

Univ. Kansas Sci. Bull., vol. 27, p. 141, text-figs. 1a-b, pl. 9, figs. 5-6, 1941.

=*Pseudoeurycea nigromaculata* (Taylor)

HOLOTYPE: 110635, Cuautlapan, Veracruz, Mexico, H. M. Smith, January-February 1940.

PARATYPES (6): 110631, 110633, 110636-8, same data. 110639, Cuautlapan, Veracruz, Mexico, H. M. Smith, Aug. 7-14, 1940. 110632 was exchanged to MZUM; 110634 to MCZ.

Bolitoglossa occidentalis Taylor

Univ. Kansas Sci. Bull., vol. 27, p. 145, text-fig. 1c, pl. 9, figs. 1-4, 1941.

HOLOTYPE: 111085, La Esperanza, Chiapas, Mexico, H. M. Smith, Apr. 28, 1940.

PARATYPES (24): 111069-84, 111086-93, La Esperanza, Chiapas, Mexico, H. M. Smith, Apr. 6-28, 1940. 111068 was exchanged to MZUM in 1944.

Bolitoglossa terrestris Taylor

Univ. Kansas Sci. Bull., vol. 27, p. 115, 1941.

=*Chiropterotriton terrestris* (Taylor)

PARATYPES (6): 116319-21, Zacualtipán, Hidalgo, Mexico, R. C. and E. H. Taylor, Aug. 10-11, 1938. 116322-3, 139717 (formerly UIMNH 26927), 4-10 miles south of Tianguistengo, Hidalgo, Mexico, E. H. and R. C. Taylor, July 1-2, 1940.

Bolitoglossa unguidentis Taylor

Herpetologica, vol. 2, p. 57, figs. 1a-c, 2c-d, 3a-b, 1941.

=*Pseudoeurycea unguidentis* (Taylor)

PARATYPE: 134292 (formerly UIMNH 27175), Cerro San Felipe, Oaxaca, Mexico, E. H. Taylor, summer 1940.

Bolitoglossa xolocalcae Taylor

Univ. Kansas Sci. Bull., vol. 27, p. 148, pl. 7, pl. 9, figs. 7-8, 1941.

=*Chiropterotriton xolocalcae* (Taylor)

HOLOTYPE: 111371, Cerro Ovando, Chiapas, Mexico, 6,800-8,000 ft., H. M. Smith, Apr. 16, 1940.

PARATYPES (71): 111400-70, same data.

Cryptobranchus bishopi Grobman

Occ. Pap. Mus. Zool. Univ. Michigan, No. 470, p. 6, 1943.

=*Cryptobranchus alleganiensis bishopi* Grobman

PARATYPES (3): 57042, Oregon County, Missouri, J. Hurter, 1900. 94356, Current River at Big Spring Park, Carter County, Missouri, A. A. Heinze, May 19, 1931. 99751, Montauk State Park, Dent County, Missouri, A. A. Heinze, Sept. 5, 1932.

Desmognathus aeneus Brown and Bishop

Copeia, No. 3, p. 163, fig. 1, pl. 1, 1947.

=*Desmognathus aeneus aeneus* Brown and Bishop

HOLOTYPE: 123977, ♂, 0.5 mile south-southeast of Peachtree, Cherokee County, North Carolina, J. C. Nichols, Jr., Oct. 22, 1946.

Desmognathus aureatagulus Weller

Proc. Jun. Soc. Nat. Hist. Cincinnati, vol. 1, No. 7, p. 3, 1930.

=*Desmognathus ochrophaeus carolinensis* Dunn

COTYPE: 93686, Mount Le Conte, Sevier County, Tennessee, R. Dury, H. Wyss, A. S. Loring and W. H. Weller, June 23, 1930.

Desmognathus brimleyorum Stejneger

Proc. U.S. Nat. Mus., vol. 17, p. 597, 1895.

=*Desmognathus fuscus brimleyorum* Stejneger

HOLOTYPE: 22157, Hot Springs, Garland County, Arkansas, H. H. and C. S. Brimley, October 1894.

PARATYPES (12): 22158-69, same data. 22170-1 were exchanged to MCZ, 1909.

Desmognathus chermocki Bishop and Valentine

Copeia, No. 1, p. 39, 1950.

=*Desmognathus aeneus chermocki* Bishop and Valentine

PARATYPES (2): 129267-8, Hurricane Creek, Tuscaloosa County, Alabama, B. O. Valentine, Mar. 30, 1948.

Desmognathus fuscus imitator Dunn

Copeia, No. 164, p. 84, 1927.

=*Desmognathus ochrophaeus carolinensis* Dunn

HOLOTYPE: 72762, near Indian Gap, Great Smoky Mountains, North Carolina, J. E. Benedict, Jr., June 9-10, 1927.

PARATYPES (3): 72761, 72763-4, same data. 72765 was exchanged to MZUM, 1944.

Desmognathus fuscus welteri Barbour

Copeia, No. 4, p. 277, 1950.

HOLOTYPE: 129312, ♂, north fork of Looney Creek, near Lynch, Harlan County, Kentucky, 2,300 ft., R. W. Barbour, June 14, 1948.

PARATYPES (42): 129313-54, same data.

Desmognathus monticola Dunn

Proc. Biol. Soc. Washington, vol. 29, p. 73, 1916.

=*Desmognathus monticola monticola* Dunn

HOLOTYPE: 38313, Elk Lodge Lake, near Brevard, Transylvania County, North Carolina, about 3,000 ft., R. and E. Tipping, July 13, 1908.

PARATYPES (9): 38314-6, 38320-1, 38323-6, same data.

Desmognathus monticola jeffersoni Hoffman

Journ. Elisha Mitchell Sci. Soc., vol. 67, p. 250, pl. 17, fig. 1, 1951.

HOLOTYPE: 126891, ♀, Saddle Hollow, 2 miles west of Crozet, Albemarle County, Virginia, 1,600 ft., R. L. Hoffman, Mar. 11, 1945.

PARATYPES (8): 123759-61, Saddle Hollow, 2 miles west of Crozet, Albemarle County, Virginia, 1,800 ft., R. L. Hoffman, Mar. 27, 1946. 126892-5, Saddle Hollow, 2 miles west of Crozet, Albemarle County, Virginia, 1,600 ft., R. L. Hoffman, Mar. 11, 1945. 130192, Charlottesville, Albemarle County, Virginia, R. L. Hoffman, Mar. 1, 1946. 126896 was exchanged to MCZ, 1958.

Desmognathus ochrophaea carolinensis Dunn

Proc. Biol. Soc. Washington, vol. 29, p. 74, 1916.

=*Desmognathus ochrophaeus carolinensis* Dunn

HOLOTYPE: 31135, Mount Mitchell, Mitchell County, North Carolina, over 6,000 ft., H. H. Brimley and F. Sherman, Jr., Oct. 5, 1902.

PARATYPES (5): 31133-4, 31136-8, same data.

Desmognathus ocoee Nichols

Journ. Tennessee Acad. Sci., vol. 24, p. 127, 1949.

HOLOTYPE: 128007, ♂, Ship's Prow Rock, Ocoee Gorge, 9 miles northwest of Ducktown, Polk County, Tennessee, J. C. Nichols, Jr., Nov. 14, 1948.

PARATYPE: 128008, ♀, same data.

Desmognathus wrighti King

Herpetologica, vol. 1, p. 57, pl. 5, 1936.

HOLOTYPE: 101794, ♂, Mount Le Conte, Great Smoky Mountains National Park, Sevier County, Tennessee, F. W. King, July 19, 1934.

PARATYPES (22, all from Great Smoky Mountains National Park, collected by F. W. King): 101790-3, 101795-6, Mount Le Conte, July 19, 1934. 101797-9, 101802, Mount Le Conte, Sept. 23, 1934. 101803-4, Pecks Corner, Aug. 15, 1934. 101805-6, Mount Hardison, Aug. 22, 1934. 101807-12, Brushy Lead, Aug. 8, 1934. 101813-4, White Rock, Oct. 14, 1934. 101800-1 were exchanged to MCZ in 1937.

Diemictylus viridescens evergladensis Peterson

Herpetologica, vol. 8, p. 103, fig. 1, 1952.

=*Diemictylus viridescens piaropicola* Schwartz and Duellman

PARATYPE: 134287 (formerly UIMNH 30492), 26 miles west of Miami, Dade County, Florida, H. W. Peterson, May 1, 1952.

Diemictylus viridescens vittatus Garman

Journ. Cincinnati Soc. Nat. Hist., vol. 19, p. 49, figs. b-c, 1897.

=*Diemictylus viridescens dorsalis* (Harlan)

COTYPES (3): 23797-9, Wilmington, New Hanover County, North Carolina, H. Garman, July 22, 1892.

Ensatina eschscholtzii xanthoptica Stebbins

Univ. California Publ. Zool., vol. 48, p. 407, pl. 1, figs. 2, 12, 1949

PARATYPE: 93606, Berkeley, Alameda County, California, B. Campbell, Feb. 2, 1931. This specimen was used as a paratype, but was not so listed.

Ensatina klauberi Dunn

Proc. U.S. Nat. Mus., vol. 74, No. 2767, p. 2, 1928.

=*Ensatina eschscholtzii klauberi* Dunn

HOLOTYPE: 75337, ♀, Descanso, San Diego County, California, J. Carter, Apr. 1, 1928.

PARATYPES (2): 75229-30, Banning, Riverside County, California, A. Gilman, Oct. 10, 1926. These specimens are considered by R. C. Stebbins to be intergrades between *Ensatina croceator* and *E. klauberi*.

Eurycea bislineata major Trapido and Clausen

Copeia, No. 3, p. 118, 1938.

=*Eurycea bislineata bislineata* (Green)



HOLOTYPE: 104239, ♂, Ouiatchouan River at Val Jalbert, Lake St. John County, Quebec, Canada, R. T. Clausen and H. Trapido, Sept. 17, 1937.

PARATYPES (3): 48056, 48081-2, Godbout, Quebec, Canada, U.S. Bureau of Biological Survey.

***Eurycea bislineata rivicola* Mittleman**

Proc. Biol. Soc. Washington, vol. 62, p. 93, 1949.

HOLOTYPE: 129397, ♂, Echo Canyon, Owen County, Indiana, M. E. and M. B. Mittleman, August 1942. Allotype, 8832, Cincinnati, Hamilton County, Ohio, J. N. B. Scarborough.

PARATYPES (20): 17465-8, Indianapolis, Marion County, Indiana, O. P. Hay. 17972-4, Lafayette, Tippecanoe County, Indiana, F. C. Test. 76825-6, 5 miles southeast of Camden, Preble County, Ohio, R. Greenfield, Apr. 3, 1929. 85686, 2 miles east of Shacklett, Cheatham County, Tennessee, C. E. Burt, June 5, 1932. 85689, 5 miles west of Belleview, Davidson County, Tennessee, C. E. Burt, June 5, 1932. 87615, 3 miles east of Pleasant Hill, Cumberland County, Tennessee, C. E. Burt, July 23, 1932. 88754, Bean Station, Grainger County, Tennessee, C. E. Burt, July 1932. 88755, Dowelltown, De Kalb County Tennessee, C. E. Burt, July 23, 1932. 118302-6, Marietta, Washington County, Ohio, F. W. King, May 21, 1930. 129398, ♀, Echo Canyon, Owen County, Indiana, M. E. and M. B. Mittleman, August 1942.

***Eurycea griseogaster* Moore and Hughes**

Copeia, No. 3, p. 139, fig. 1, 1941.

=*Eurycea multiplicata griseogaster* Moore and Hughes

PARATYPES (14): 113231-4, 113235 (10 larvae), Swimmer's Creek, 10 miles northeast of Gore, Sequoyah County, Oklahoma, A Seamster and G. A. Moore, Apr. 11, 1941.

***Eurycea latitans* Smith and Potter**

Herpetologica, vol. 3, p. 106, 1946.

HOLOTYPE: 123594, ♂, Cascade Cavern, 4.6 miles southeast of Boerne, Kendall County, Texas, F. E. Potter, Jr., May 15, 1946.

***Eurycea longicauda pernix* Mittleman**

Proc. New England Zool. Club, vol. 21, p. 101, pl. 20, 1942.

PARATYPE: 120708 (formerly MCZ 25574), Brown County State Park, 2.5 miles southeast of Nashville, Brown County, Indiana, M. E. and M. B. Mittleman, Aug. 22, 1942.

***Eurycea neotenes* Bishop and Wright**

Proc. Biol. Soc. Washington, vol. 50, p. 142, fig., 1937.

HOLOTYPE: 103161, ♂, Culebra Creek, 5 miles north of Helotes, Bexar County, Texas, M. R. Wright and S. C. Bishop, Apr. 1, 1936. Allotype, 103162, same data.

Eurycea tynereensis Moore and Hughes

Amer. Midl. Nat., vol. 22, p. 697, figs. 1-2, 1939.

PARATYPES (12): 108548 (12 larvae), Tyner Creek, 0.5 mile east of Proctor, Adair County, Oklahoma, J. T. Mizelle and G. A. Moore, May 29, 1939.

Gyrinophilus dunnii Mittleman and Jopson

Smithsonian Misc. Coll., vol. 101, No. 3638, p. 2, pl. 1, 1941.

=*Gyrinophilus danielsi dunnii* Mittleman and Jopson

HOLOTYPE: 113230, ♀, Clemson, Pickens County, South Carolina, A. Grobman, Apr. 8, 1941.

PARATYPES (4): 68168, Indian Cave, near Jefferson City, Jefferson County, Tennessee, 1,250 ft., J. D. Ives. 68820, Indian Cave, near Jefferson City, Jefferson County, Tennessee, 1,200 ft., J. D. Ives, Feb. 12, 1945. 102440, Caesar's Head, Greenville County, South Carolina, 3,000 ft., H. G. M. Jopson, Apr. 13, 1934. 102441, larva, east of Rabun Gap, Rabun County, Georgia, 2,300 ft., H. P. and H. G. M. Jopson, Apr. 6, 1934.

Gyrinophilus porphyriticus inagnoseus Mittleman

Proc. New England Zool. Club, vol. 20, p. 27, pl. 6, fig. F, text-fig. 1 (incorrectly given as "2" in original description), 1942.

HOLOTYPE: 115520, ♀, Salt Creek, 4 miles southwest of South Bloomingville, Hocking County, Ohio, M. B. Mittleman, Sept. 25, 1940.

Hemidactylium pacificum Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 17, p. 195, 1865.

=*Batrachoseps pacificus* (Cope)

HOLOTYPE: 6733, Santa Barbara, Santa Barbara County, California, W. W. Hayes.

Heredia oregonensis Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 8, p. 140, 1857.

=*Ensatina eschscholtzii oregonensis* (Girard)

COTYPES (2): 15479-80, Puget Sound (Washington), U.S. Exploring Expedition, 1841.

Hynobius retardatus Dunn

Proc. California Acad. Sci., ser. 4, vol. 12, p. 27, 1923.

PARATYPES (2): 64902, Kutara Lake, Iburi, Hokkaido, Japan, M. Oshima. 64903, estuary of Shirisatsumi River near Lake Shikotsu, Iburi, Hokkaido, Japan, M. Oshima.

Hynobius shihi Liu

Chicago Nat. Hist. Mus., Fieldiana: Zool. Mem., vol. 2, p. 77, 1950.

PARATYPE: 124580, Ta-pa-shan, Szechwan, China, C. C. Liu.

Hynobius stejnegeri Dunn

Proc. California Acad. Sci., ser. 4, vol. 12, p. 28, 1923.

HOLOTYPE: 23901, ♀, Kyushu, Higo Province, Kumamoto, Japan, Nakagawa, 1884.

Manculus quadridigitatus paludicolus Mittleman

Herpetologica, vol. 3, p. 220, 1947.

=*Manculus quadridigitatus* (Holbrook)

HOLOTYPE: 123979, Pollock, Grant Parish, Louisiana, P. Viosca, Jr., Sept. 6-9, 1937.

PARATYPES (9): 17700, Palestine, Anderson County, Texas, B. W. Evermann. 99176-7, 4 miles southwest of Many, Sabine Parish, Louisiana, C. E. Burt, June 16, 1934. 99762, Hillister, Tyler County, Texas, C. E. Burt, July 6, 1934. 99773-7, 5 miles southwest of Pendleton Ferry, Sabine County, Texas, C. E. Burt, June 17, 1934.

Manculus quadridigitatus uvidus Mittleman

Herpetologica, vol. 3, p. 221, 1947.

=*Manculus quadridigitatus* (Holbrook)

HOLOTYPE: 123980, Gayle, Caddo Parish, Louisiana, L. S. Frierson and J. K. Strecker.

Menobranchus lateralis latastei Garnier

Proc. Canadian Inst., ser. 3, vol. 5, p. 218, 1888.

=*Necturus maculosus maculosus* (Rafinesque)

COTYPES (2): 22331-2, Maitland River, Ontario, Canada, J. H. Garnier.

Menobranchus punctatus Gibbes

Proc. Amer. Assoc. Adv. Sci., p. 159, 1850.

=*Necturus punctatus* (Gibbes)

COTYPE: 11813, South Santee River, South Carolina, L. R. Gibbes, 1850.

Necturus alabamensis Viosca

Copeia, No. 2, p. 121, 1937.

=*Necturus maculosus beyeri* Viosca

HOLOTYPE: 102676, ♀, Black Warrior River, near Tuscaloosa, Tuscaloosa County, Alabama, Univ. Alabama, Jan. 22, 1914.

Necturus beyeri Viosca

Copeia, No. 2, p. 123, fig. 2, 1937.

=*Necturus maculosus beyeri* Viosca

HOLOTYPE: 102674, ♀, Oakdale, upper Calcasieu River, Allen Parish, Louisiana, P. Viosca, Jr., winter of 1928.

PARATYPE: 102675, same data.

Necturus lödingi Viosca

Copeia, No. 2, p. 126, fig. 4, 1937.

HOLOTYPE: 61752, ♀, Mertz Station, Eslava Creek, near Mobile, Mobile County, Alabama, C. Löding, February 1918.

Necturus louisianensis Viosca

Proc. Biol. Soc. Wash., vol. 51, p. 143, pls. 1-2, 1938.

=*Necturus maculosus louisianensis* Viosca

HOLOTYPE: 104238, ♀, Fishville, in Big Creek, east of Pollock, Grant Parish, Louisiana, P. Viosca and O. F.-R. Bruce, Feb. 22, 1938.

Necturus maculosus lewisi Brimley

Journ. Elisha Mitchell Sci. Soc., vol. 40, p. 167, 1924.

HOLOTYPE: 73848, Neuse River, near Raleigh, Wake County, North Carolina, F. B. Lewis, Feb. 25, 1921.

Oedipina ignea Stuart

Proc. Biol. Soc. Washington, vol. 65, p. 1, 1952.

HOLOTYPE: 127959, ♀, Río Las Brisas, south of Yepocapa, Department of Chimaltenango, Guatemala, 1,450 m., H. T. Dalmat, Aug. 2, 1947.

Oedipus bromeliacia Schmidt

Field Mus. Nat. Hist., Zool. Ser., vol. 20, p. 161, 1936.

=*Chiropterotriton bromeliacia* (Schmidt)

PARATYPES (2): 103339-40 (formerly FMNH 20294), Volcán Tajumulco, Guatemala, K. P. Schmidt, Feb. 19, 1934.

Oedipus flavimembris Schmidt

Field Mus. Nat. Hist., Zool. Ser., vol. 20, p. 158, 1936.

=*Magnadigita flavimembris* (Schmidt)

PARATYPE: 103331 (formerly FMNH 20323), Volcán Tajumulco, Guatemala, 7,200 ft., K. P. Schmidt, Feb. 22, 1934.

Oedipus flaviventris Schmidt

Field Mus. Nat. Hist., Zool. Ser., vol. 20, p. 148, 1936.

=*Bolitoglossa flaviventris* (Schmidt)

HOLOTYPE: 46922, ♀, Chicharras, Chiapas, Mexico, E. W. Nelson and E. A. Goldman, Feb. 6, 1896.

PARATYPES (2): 30305, Tehuantepec, Oaxaca, Mexico, F. Sumichrast. 46923, Chicharras, Chiapas, Mexico, E. W. Nelson and E. A. Goldman, Feb. 6, 1896.

Oedipus franklini Schmidt

Field Mus. Nat. Hist., Zool. Ser., vol. 20, p. 159, fig. 19, 1936.

=*Magnadigita franklini* (Schmidt)

PARATYPES (2): 103332-3 (formerly FMNH 20655-6), Volcán Tajumulco, Guatemala, 7,500 ft., K. P. Schmidt, Feb. 27, 1934.

Oedipus gadovii Dunn

Salamanders of the family Plethodontidae, Ann. Publ. Smith Coll., vol. 50, p. 437, 1926.

=*Pseudoeurycea gadovii* (Dunn)

PARATYPES (3): 19263-5, Mount Orizaba, Veracruz, Mexico, 12,000 ft., W. S. Blatchley, Aug. 2-4, 1891.

Oedipus goebeli Schmidt

Field Mus. Nat. Hist., Zool. Ser., vol. 20, p. 163, 1936.

=*Pseudoeurycea goebeli* (Schmidt)

PARATYPES (2): 103334-5 (formerly FMNH 20299, 20301), Volcán Tajumulco, Guatemala, 8,000-14,000 ft., K. P. Schmidt, Feb. 13-14, 1934.

Oedipus helmrichi Schmidt

Field Mus. Nat. Hist., Zool. Ser., vol. 20, p. 152, fig. 18c, 1936.

=*Bolitoglossa helmrichi* (Schmidt)

PARATYPE: 103338 (formerly FMNH 20727), Samar, Alta Verapaz, Guatemala, 5,000 ft., D. Clark, Mar. 27, 1934.

Oedipus multidentata Taylor

Univ. Kansas Sci. Bull., vol. 25 (1938), p. 289, pl. 29, fig. 1, 1939.

=*Chiropterotriton multidentata* (Taylor)

PARATYPES (3): 134284-6 (formerly UIMNH 26953-5), El Chico National Park, Hidalgo, Mexico, E. H. Taylor, Aug. 2, 1938.

Oedipus nasalis Dunn

Field Mus. Nat. Hist., Zool. Ser., vol. 12, p. 97, 1924.

=*Chiropterotriton nasalis* (Dunn)

PARATYPES (2): 103336-7 (formerly FMNH 4569-70), San Pedro, Honduras, 4,500 ft., K. P. Schmidt, Apr. 1, 1923.

Oedipus odonnelli Stuart

Misc. Publ. Mus. Zool. Univ. Michigan, No. 56, p. 10, 1943.

=*Bolitoglossa odonnelli* (Stuart)

PARATYPE: 118781 (formerly MZUM 89105), Finca Volcán, Alta Verapaz, Guatemala, L. C. Stuart, Mar. 20, 1938.

Oedipus robertsi Taylor

Univ. Kansas Sci. Bull., vol. 25 (1938), p. 287, pl. 26, fig. 2, 1939.

=*Pseudoeurycea robertsi* (Taylor)

PARATYPE: 134290 (formerly UIMNH 27210), Nevada de Toluca, Mexico, Mexico, 10,000-11,000 ft., E. H. Taylor, Aug. 31, 1938.

Oedipus smithi Taylor

Univ. Kansas Sci. Bull., vol. 25 (1938), p. 269, pl. 25, figs. 5-6, 1939.

=*Pseudoeurycea smithi* (Taylor)

PARATYPE: 134291 (formerly UIMNH 27219), Cerro San Felipe, Oaxaca, Mexico, E. H. Taylor, Aug. 20-22, 1938.

Oedipus striatulus Noble

Bull. Amer. Mus. Nat. Hist., vol. 38, p. 344, pl. 19, 1918.

=*Bolitoglossa striatula* (Noble)

PARATYPE: 37771, Turrialba, Costa Rica, J. F. Tristan, Feb. 6, 1906.

Oedipus townsendi Dunn

Proc. Biol. Soc. Washington, vol. 35, p. 5, 1922.

=*Parvimolge townsendi* (Dunn)

PARATYPE: 30352, Tehuantepec, Oaxaca, Mexico? (probably from Cerro de la Defensa, near Córdoba, Veracruz, fide Dunn), C. Sartorius?.

Plethodon cinereus angusticlavius Grobman

Ann. New York Acad. Sci., vol. 45, p. 302, fig. 6c, 1944.

=*Plethodon dorsalis angusticlavius* Grobman

PARATYPE: 57089, Stone County, Missouri, J. Hurter, June 20, 1910.

Plethodon cinereus dorsalis Baird

in Cope, U.S. Nat. Mus. Bull. 34, p. 138, 1889.

=*Plethodon dorsalis dorsalis* Baird

COTYPES (4): 3776 (4), Louisville, Jefferson County, Kentucky.

Plethodon clemsonae Brimley

Copeia, No. 164, p. 73, 1927.

=*Plethodon jordani clemsonae* Brimley

HOLOTYPE: 73849, ♂, Jocassee, Oconee County, South Carolina, 1,200-1,500 ft., J. A. Berly, Apr. 8, 1927.

Plethodon crassulus Cope

Proc. Amer. Philos. Soc., vol. 23, p. 521, 1886.

=*Aneides lugubris lugubris* (Hallowell)

HOLOTYPE: 9447, California, J. G. Cooper.

Plethodon dixi Pope and Fowler

Chicago Acad. Sci. Nat. Hist. Misc., No. 47, p. 2, 1949.

=*Plethodon wehrlei dixi* Pope and Fowler

PARATYPES (11): 127239-49, Dixie Caverns, 5 miles south of Salem, Roanoke County, Virginia, J. A. Fowler, June 21, 1948.

Plethodon dumni Bishop

Proc. Biol. Soc. Washington, vol. 47, p. 169, pl. 1, 1934.

HOLOTYPE: 95196, ♀, near Portland, Clackamas County, Oregon, S. G. Jewett, Jr., Jan. 13, 1934.

Plethodon glutinosus chlorobryonis Mittleman

Herpetologica, vol. 7, p. 108, 1951.

=*Plethodon glutinosus glutinosus* (Green)

Oedipus gadovii Dunn

Salamanders of the family Plethodontidae, Ann. Publ. Smith Coll., vol. 50, p. 437, 1926.

=*Pseudoeurycea gadovii* (Dunn)

PARATYPES (3): 19263-5, Mount Orizaba, Veracruz, Mexico, 12,000 ft., W. S. Blatchley, Aug. 2-4, 1891.

Oedipus goebeli Schmidt

Field Mus. Nat. Hist., Zool. Ser., vol. 20, p. 163, 1936.

=*Pseudoeurycea goebeli* (Schmidt)

PARATYPES (2): 103334-5 (formerly FMNH 20299, 20301), Volcán Tajumulco, Guatemala, 8,000-14,000 ft., K. P. Schmidt, Feb. 13-14, 1934.

Oedipus helmrichi Schmidt

Field Mus. Nat. Hist., Zool. Ser., vol. 20, p. 152, fig. 18c, 1936.

=*Bolitoglossa helmrichi* (Schmidt)

PARATYPE: 103338 (formerly FMNH 20727), Samar, Alta Verapaz, Guatemala, 5,000 ft., D. Clark, Mar. 27, 1934.

Oedipus multidentata Taylor

Univ. Kansas Sci. Bull., vol. 25 (1938), p. 289, pl. 29, fig. 1, 1939.

=*Chiropterotriton multidentata* (Taylor)

PARATYPES (3): 134284-6 (formerly UIMNH 26953-5), El Chico National Park, Hidalgo, Mexico, E. H. Taylor, Aug. 2, 1938.

Oedipus nasalis Dunn

Field Mus. Nat. Hist., Zool. Ser., vol. 12, p. 97, 1924.

=*Chiropterotriton nasalis* (Dunn)

PARATYPES (2): 103336-7 (formerly FMNH 4569-70), San Pedro, Honduras, 4,500 ft., K. P. Schmidt, Apr. 1, 1923.

Oedipus odonnelli Stuart

Misc. Publ. Mus. Zool. Univ. Michigan, No. 56, p. 10, 1943.

=*Bolitoglossa odonnelli* (Stuart)

PARATYPE: 118781 (formerly MZUM 89105), Finca Volcán, Alta Verapaz, Guatemala, L. C. Stuart, Mar. 20, 1938.

Oedipus robertsi Taylor

Univ. Kansas Sci. Bull., vol. 25 (1938), p. 287, pl. 26, fig. 2, 1939.

=*Pseudoeurycea robertsi* (Taylor)

PARATYPE: 134290 (formerly UIMNH 27210), Nevada de Toluca, Mexico, Mexico, 10,000-11,000 ft., E. H. Taylor, Aug. 31, 1938.

Oedipus smithi Taylor

Univ. Kansas Sci. Bull., vol. 25 (1938), p. 269, pl. 25, figs. 5-6, 1939.

=*Pseudoeurycea smithi* (Taylor)

PARATYPE: 134291 (formerly UIMNH 27219), Cerro San Felipe, Oaxaca, Mexico, E. H. Taylor, Aug. 20-22, 1938.

Oedipus striatulus Noble

Bull. Amer. Mus. Nat. Hist., vol. 38, p. 344, pl. 19, 1918.

=*Bolitoglossa striatula* (Noble)

PARATYPE: 37771, Turrialba, Costa Rica, J. F. Tristan, Feb. 6, 1906.

Oedipus townsendi Dunn

Proc. Biol. Soc. Washington, vol. 35, p. 5, 1922.

=*Parvimolge townsendi* (Dunn)

PARATYPE: 30352, Tehuantepec, Oaxaca, Mexico? (probably from Cerro de la Defensa, near Córdoba, Veracruz, fide Dunn), C. Sartorius?.

Plethodon cinereus angusticlavius Grobman

Ann. New York Acad. Sci., vol. 45, p. 302, fig. 6c, 1944.

=*Plethodon dorsalis angusticlavius* Grobman

PARATYPE: 57089, Stone County, Missouri, J. Hurter, June 20, 1910.

Plethodon cinereus dorsalis Baird

in Cope, U.S. Nat. Mus. Bull. 34, p. 138, 1889.

=*Plethodon dorsalis dorsalis* Baird

COTYPES (4): 3776 (4), Louisville, Jefferson County, Kentucky.

Plethodon clemsonae Brimley

Copeia, No. 164, p. 73, 1927.

=*Plethodon jordani clemsonae* Brimley

HOLOTYPE: 73849, ♂, Jocassee, Oconee County, South Carolina, 1,200-1,500 ft., J. A. Berly, Apr. 8, 1927.

Plethodon crassulus Cope

Proc. Amer. Philos. Soc., vol. 23, p. 521, 1886.

=*Aneides lugubris lugubris* (Hallowell)

HOLOTYPE: 9447, California, J. G. Cooper.

Plethodon dixi Pope and Fowler

Chicago Acad. Sci. Nat. Hist. Misc., No. 47, p. 2, 1949.

=*Plethodon wehrlei dixi* Pope and Fowler

PARATYPES (11): 127239-49, Dixie Caverns, 5 miles south of Salem, Roanoke County, Virginia, J. A. Fowler, June 21, 1948.

Plethodon dumni Bishop

Proc. Biol. Soc. Washington, vol. 47, p. 169, pl. 1, 1934.

HOLOTYPE: 95196, ♀, near Portland, Clackamas County, Oregon, S. G. Jewett, Jr., Jan. 13, 1934.

Plethodon glutinosus chlorobryonis Mittleman

Herpetologica, vol. 7, p. 108, 1951.

=*Plethodon glutinosus glutinosus* (Green)

HOLOTYPE: 129933, ♂, 13 miles north of New Bern, Craven County, North Carolina, C. B. Goodstein and M. B. Mittleman, Nov. 10, 1950. 129934, allotype, same data.

PARATYPES (6): 83469, Lake Waccamaw, Columbus County, North Carolina, J. E. Benedict, Jr. 84206-7, Williams Mill Pond, Gates County, North Carolina, M. K. Brady, Aug. 11, 1931. 91725-6, Lake Waccamaw, Columbus County, North Carolina, I. E. Gray, July 3, 1933. 129936, 4 miles north of Windsor, Bertie County, North Carolina, C. B. Goodstein and M. B. Mittleman, Nov. 10, 1950. 124354 from Perquimans County was exchanged to MCZ in 1958.

Plethodon hubrichti Thurow

Herpetologica, vol. 13, p. 59, 1957.

HOLOTYPE: 139087, ♂, Blue Ridge Parkway, 0.9 mile south of Milepost 80, Bedford County, Virginia, G. R. Thurow, Dec. 9, 1956.

PARATYPES (2): 139088-9, ♂, ♀, same data.

Plethodon huldae Grobman

Proc. Biol. Soc. Washington, vol. 62, p. 136, 1949.

=*Plethodon cinereus cinereus* (Green)

HOLOTYPE: 127955, foottrail to Hawksbill Mountain, near Skyline Drive, Madison County, Virginia, H. and A. Grobman, Sept. 5, 1947.

PARATYPES (3): 127956-8, foottrail to Hawksbill Mountain, Madison County, Virginia, H. I. Kleinpeter and S. A. Peabody, August 1947.

Plethodon idahoensis Slater and Slipp

Occ. Pap. Dept. Biol. Coll. Puget Sound, No. 8, p. 38, figs. 1-2, 1940.

=*Plethodon vandykei idahoensis* Slater and Slipp

HOLOTYPE: 110504, northeast corner of Coeur d'Alene Lake, Kootenai County, Idaho, 2,160 ft., J. R. Slater, 1939.

PARATYPE: 110505, northeast corner of Coeur d'Alene Lake, Kootenai County, Idaho, J. W. Slipp, 1939.

Plethodon intermedius Baird

Proc. Acad. Nat. Sci. Philadelphia, vol. 8, p. 207, 1857.

COTYPE: 4732a, Fort Tejon (Kern County), California, J. Xantus.

Plethodon jacksoni Newman

Herpetologica, vol. 10, p. 9, fig. 1, 1954.

=*Plethodon wehrlei jacksoni* Newman

HOLOTYPE: 134498, ♂, Trillium Vale near Blacksburg, Montgomery County, Virginia, 2,100 ft., W. B. Newman, Feb. 11, 1950.

Plethodon kentucki Mittleman

Herpetologica, vol. 7, p. 105, fig., 1951.

=*Plethodon glutinosus glutinosus* (Green)

HOLOTYPE: 129937, ♂, Pine Mountain, Harlan County, Kentucky, W. Cornett, August 1933. 129938, allotype, same data.

***Plethodon metcalfi* Brimley**

Proc. Biol. Soc. Washington, vol. 25, p. 138, pl. 6, figs. 1-2, 4, 7-11, 1912.

=*Plethodon jordani metcalfi* Brimley

HOLOTYPE: 49682, Sunburst, Haywood County, North Carolina, 3,500 ft., C. S. Brimley, May 1912.

PARATYPE: 49683, same data.

***Plethodon neomexicanus* Stebbins and Riemer**

Copeia, No. 2, p. 73, pl. 1, 1950.

PARATYPE: 129378, 12 miles west and 4 miles south of Los Alamos, Sandoval County, New Mexico, 8,750 ft., R. C. Stebbins, Aug. 14, 1949.

***Plethodon ouachitae* Dunn and Heinze**

Copeia, No. 3, p. 121, 1933.

HOLOTYPE: 92484, ♂, Rich Mountain in Ouachita National Forest, Polk County, Arkansas, A. A. Heinze and D. A. Boyer, May 30, 1933.

***Plethodon richmondi* Netting and Mittleman**

Ann. Carnegie Mus., vol. 27, p. 288, pl. 30, fig. 1, 1938.

=*Plethodon richmondi richmondi* Netting and Mittleman

PARATYPES (3): 107283-4, Athens County, Ohio, H. T. Gier, Mar. 29, 1938. 108549 (formerly CM 14188), Ritter Park, Huntington, Cabell County, West Virginia, N. D. Richmond and J. Green, Oct. 15, 1938.

***Plethodon richmondi popei* Highton and Grobman**

Herpetologica, vol. 12, p. 187, 1956.

PARATYPES (34): 135178, Nigger Mountain, southeast of Jefferson, Ashe County, North Carolina, L. Hubricht, Oct. 17, 1953. 135179, Milepost 236.2, Blue Ridge Parkway, Alleghany County, North Carolina, L. Hubricht, Oct. 10, 1953. 135180, 2 miles southeast of Sparta, Alleghany County, North Carolina, L. Hubricht, Oct. 17, 1953. 135181-98, Burkes Garden, Tazewell County, Virginia, R. L. Hoffman, July 12, 1953-Mar. 20, 1954. 135199-202, 6 miles west of Wytheville, Wythe County, Virginia, R. L. Hoffman, Mar. 30, 1954. 135207-13, Brushy Mountain, 5 miles east of Marion, Smyth County, Virginia, R. L. Hoffman, May 2, 1954. 137291-2, Comers Rock, Grayson-Wythe County line, Virginia, W. Auffenberg and A. B. Grobman, Aug. 30, 1951.

***Plethodon shermani* Stejneger**

Proc. U.S. Nat. Mus., vol. 30, p. 559, fig. 1, 1906.

=*Plethodon jordani shermani* Stejneger

HOLOTYPE: 36214, Nantahala Mountain, between Andrews and Aquone, North Carolina, F. Sherman, Jr., Aug. 24, 1904.

Plethodon vandykei larselli Burns

Herpetologica, vol. 10, p. 83, 1954.

HOLOTYPE: 134129, ♂, north slope of Larch Mountain, Multnomah County, Oregon, D. M. Burns, May 24, 1953.

PARATYPE: 134130, north slope of Larch Mountain, Multnomah County, Oregon, D. M. Burns, early June 1951.

Plethodon welleri Walker

Proc. Jun. Soc. Nat. Sci. Cincinnati, vol. 2, p. 48, 1931.

=*Plethodon welleri welleri* Walker

HOLOTYPE: 84135, ♂, Grandfather Mountain, near Linville, Avery County, North Carolina, above 5,000 ft., W. H. Weller and R. Dury, Aug. 27, 1930.

PARATYPES (2): 84136-7, same data.

Plethodon welleri ventromaculatum Throuow

Amer. Midl. Nat., vol. 55, p. 344, pl. 1, 1956.

PARATYPES (5): 124632-3, Mount Rogers, Grayson County, Virginia, R. L. Hoffman and H. I. Kleinpeter, July 1, 1947. 132323-4, 5.3 miles north of Carderview, Johnson County, Tennessee, R. L. Hoffman, May 31, 1952. 133073, White Top Mountain, Grayson County, Virginia, L. G. Carr, Aug. 15, 1946. 133074 was exchanged to MCZ in 1958.

Plethodon yonahlossee Dunn

Bull. Amer. Mus. Nat. Hist., vol. 37, p. 598, 1917.

PARATYPES (3): 36397-8, between Linville and Blowing Rock, North Carolina, F. Sherman, Jr., September 1902. 62017, Linville, Avery County, North Carolina, E. R. Dunn, Aug. 16, 1916.

Plethopsis wrighti Bishop

Herpetologica, vol. 1, p. 93, pl. 9, 1937.

=*Batrachoseps wrighti* (Bishop)

HOLOTYPE: 102445, ♂, 8.7 miles southeast of Sandy, Clackamas County, Oregon, S. C. Bishop, June 15, 1936. 102446, allotype, same data.

Pseudobranchius striatus axanthus Netting and Goin

Ann. Carnegie Mus., vol. 29, p. 183, pl. 1, figs. 1-2, 1942.

PARATYPES (10): 67352-3, Gainesville, Alachua County, Florida, T. Van Hyning, 1924. 92566, Gainesville, Alachua County, Florida, M. K. Brady, 1931. 107288, 107290-4, Newnans Lake, near Gainesville, Alachua County, Florida, C. R. Aschemeier, Feb. 24, 1938. 118790 (formerly MZUM 77150), Paines Prairie, near Gainesville, Alachua County, Florida, W. Clanton, Oct. 4, 1932. 107289 was exchanged to ZS in 1952.

Pseudobranchius striatus spheniscus Goin and Crenshaw

Ann. Carnegie Mus., vol. 31, p. 277, 1949.

PARATYPES (5): 128140-4, larvae, Putney Pond, Baker County, Georgia, J. W. Crenshaw and G. B. Rabb, July 1, 1948.

Pseudoeurycea brunnata Bumzahem and Smith

Herpetologica, vol. 11, p. 73, 1955.

PARATYPE: 137207 (formerly UIMNH 33720), near Soconusco, Chiapas, Mexico, E. Matuda, 1944-49.

Pseudoeurycea firscheini Shannon and Werler

Herpetologica, vol. 11, p. 82, fig. 2, 1955.

PARATYPE: 139718 (formerly UIMNH 25103), mountain crest above Aculzingo, Veracruz, Mexico, J. Werler, Jan. 7, 1951.

Pseudoeurycea werleri Darling and Smith

Trans. Kansas Acad. Sci., vol. 57, p. 180, 1954.

PARATYPE: 137208 (formerly UIMNH 33896), Volcán San Martín, Veracruz, Mexico, D. Darling, Feb. 22-28, 1953.

Pseudotriton duryi Weller

Proc. Jun. Soc. Nat. Hist. Cincinnati, vol. 1, p. 9, 1930.

=*Gyrinophilus porphyriticus duryi* (Weller)

HOLOTYPE: 84300, Cascade Caverns, Carter County, Kentucky, R. Dury and W. H. Weller, Apr. 6, 1930.

Pseudotriton montanus Baird

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 1, p. 293, 1849.

=*Pseudotriton montanus montanus* Baird

COTYPES (3): 3839 (3), Carlisle, Cumberland County, Pennsylvania, S. F. Baird.

Pseudotriton montanus floridanus Netting and Goin

Ann. Carnegie Mus., vol. 29, p. 175, pl. 1, fig. 5, 1942.

PARATYPES (2): 22819, Chuluota, Seminole County, Florida, R. A. Mills. 118789 (formerly MZUM 86419), Gainesville, Alachua County, Florida, Apr. 10, 1936.

Pseudotriton ruber vioscai Bishop

Occ. Pap. Boston Soc. Nat. Hist., vol. 5, p. 247, 1928.

HOLOTYPE: 75057, ♀, 10 miles west of Bogalusa, Washington Parish, Louisiana, P. Viosca, Jr., Apr. 10, 1926.

PARATYPE: 75080, Bogalusa, Washington Parish, Louisiana, P. Viosca, Jr., Feb. 13, 1928.

Ranodon olympicus Gaige

Occ. Pap. Mus. Zool. Univ. Michigan, No. 40, p. 2, pl. 1, 1917.

=*Rhyacotriton olympicus* (Gaige)

PARATYPES (2): 55277, 55331, Olympic Mountains, Lake Cushman, Washington, P. Putnam, Apr. 19, 1916 (55331), May 7, 1916 (55277).

Rhyacosiredon rivularis Taylor

Herpetologica, vol. 1, p. 171, pl. 17, fig. 1, 1940.

PARATYPE: 137209 (formerly UIMNH 27241), 8 miles west of Villa Victoria, Mexico, Mexico, E. H. Taylor, Sept. 11, 1938.

Rhyacosiredon zempoalaensis Taylor and Smith

Proc. U.S. Nat. Mus., vol. 95, p. 527, 1945.

HOLOTYPE: 116617, Zempoala, Morelos, Mexico, 10,000 ft., H. M. Smith, Feb. 23, 1939.

PARATYPES (14): 116614-6, 116118-28, Zempoala, Morelos, Mexico, 10,000 ft., H. M. Smith, Feb. 21-24, 1939.

Salamandra agilis Sager

Amer. Journ. Sci., ser. 1, vol. 36, p. 322, 1839.

=*Plethodon cinereus cinereus* (Green)

COTYPES? (15): 3770 (15), Detroit, Wayne County, Michigan, A. Sager.

Salamandra granulata De Kay

Zoology of New York, vol. 1, pt. 3 (Reptiles and Amphibia), p. 73, pl. 23, fig. 66, 1842 (in [New York State], Natural History of New York).

=*Ambystoma jeffersonianum* (Green)

COTYPES? (2): 3981 (2), New York.

Salamandra jeffersoniana Green

Contr. Maclurian Lyc., vol. 1, No. 1, p. 4, pl. 1, fig. 1, 1827.

=*Ambystoma jeffersonianum* (Green)

HOLOTYPE?: 3968, near Chartier's Creek, at Canonsburg, Washington County, Pennsylvania, J. Green.

Salamandra lurida Sager

Amer. Journ. Sci., ser. 1, vol. 36, p. 323, 1839.

=*Ambystoma tigrinum tigrinum* (Green)

HOLOTYPE: 39442 (formerly 3899), Detroit, Wayne County, Michigan, A. Sager.

PARATYPE: 3970, same data.

Salamandrella sinensis Sauvage

Bull. Soc. Philom. Paris, ser. 7, vol. 1, p. 117, 1877.

=*Batrachuperus pinchonii* David

PARATYPE: 10995, Kiang-si, China, MHNP.

Siredon dumerilii Dugès

Naturaleza, vol. 1, p. 241, pl. 5a, figs. 1-13, 1870.

=*Bathysiredon dumerilii* (Dugès)

COTYPES (2): 16201-2, Lake Pátzcuaro, Michoacán, Mexico, 2,055 ft., A. Dugès.

Siredon gracilis Baird

Report on reptiles collected on the survey, in Report U.S. Pacific R.R. Explor. and Surv. [California and Oregon], vol. 10, pt. 4, No. 4, p. 13, pl. 44, fig. 2, 1859.

= *Ambystoma gracile gracile* (Baird)

COTYPES (2): 4080 (2), larvae, Cascade Mountains, Oregon, latitude 44°, J. S. Newberry.

Siren intermedia nettingi Goin

Ann. Carnegie Mus., vol. 29, p. 211, 1942.

PARATYPE: 118793 (formerly MZUM 68382), Imboden, Lawrence County, Arkansas, B. C. Marshall, May 3, 1929.

Siren intermedia texana Goin

Herpetologica, vol. 13, p. 37, 1957.

PARATYPES (10): 4048 (7), 4075 (head only), Matamoras, Tamaulipas, Mexico, D. N. Couch. 10853, 10855, Upson, Maverick County, Texas, A. Türpe, 1880. 10861 also from Maverick County was exchanged to IML in 1946; 10857 to MCZ, 1958.

Siredon lermaensis Taylor

Univ. Kansas Sci. Bull., vol. 26 (1939), p. 427, pl. 48, 1940.

PARATYPE: 134297 (formerly UIMNH 27236), Lerma, Mexico, Mexico, E. H. Taylor and H. M. Smith, 1939.

Siredon lichenoides Baird

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 68, 1852.

= *Ambystoma tigrinum mavortium* Baird

COTYPES (2): 4061 (2), larvae, Spring Lake, at the head of Santa Fe Creek, New Mexico, R. H. Kern.

Spelerpes bilineatus borealis Baird

in Cope, U.S. Nat. Mus. Bull. 34, p. 165, 1889.

= *Eurycea bislineata bislineata* (Green)

COTYPES (12): 4735a (12), Lake Oquassoc (Franklin County), Maine, C. Girard.

Spelerpes collaris Stejneger

Proc. U.S. Nat. Mus., vol. 32, p. 465, 1907.

= *Oedipina collaris* (Stejneger)

HOLOTYPE: 37350, Topaz Mines, 90 miles northwest of Bluefields, Nicaragua, 400 ft., W. F. Thornton.

Spelerpes gibbicaudus Blatchley

Proc. U.S. Nat. Mus., vol. 16, p. 38, 1893.

= *Pseudoeurycea leprosa* (Cope)

HOLOTYPE: 19255 (formerly 6340), Orizaba, Veracruz, Mexico, F. Sumichrast. This specimen is also one of the cotypes of *Spelerpes leprosus* Cope.

Spelerpes leprosus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 21, p. 105, 1869.

=*Pseudoeurycea leprosa* (part) (Cope)

COTYPES (5): 6340 (2), 19255, 103591-2, Orizaba, Veracruz, Mexico, F. Sumichrast. 19255 is the lectotype of *leprosus*, and is also the holotype of *Spelerpes gibbicaudus* Blatchley. 103591 (formerly 6340) is *Pseudoeurycea cephalica cephalica* (Cope), and 103592 (formerly 6340) is an undetermined form according to Taylor, Univ. Kansas Sci. Bull., vol. 25, p. 274, 1939.

Spelerpes multiplicatus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 21, p. 106, 1869.

=*Eurycea multiplicata* (Cope)

COTYPES (5): 4038 (5), Red River (restricted to vicinity of Fort Towson, Choctaw County, Oklahoma), L. A. Edwards.

Spelerpes orizabensis Blatchley

Proc. U.S. Nat. Mus., vol. 16, p. 38, 1893.

=*Pseudoeurycea leprosa* (Cope)

COTYPES (2): 19266-7, Orizaba, Veracruz, Mexico, 11,000 ft., W. S. Blatchley, Aug. 2, 1891.

Spelerpes picadoi Stejneger

Proc. U.S. Nat. Mus., vol. 41, p. 285, 1911.

=*Chiropterotriton picadoi* (Stejneger)

HOLOTYPE: 48280, La Estrella, southeast of Cartago, Costa Rica, C. Picado, September 1911.

Spelerpes ruber schenki Brimley

Proc. Biol. Soc. Washington, vol. 25, p. 139, pl. 7, figs. 12-13, 15, 17-18, 20-22, 1912.

=*Pseudotriton ruber schenki* (Brimley)

HOLOTYPE: 49679, Sunburst, Haywood County, North Carolina, 3,200 ft., C. S. Brimley, May 1912.

PARATYPE: 49680, same data.

Spelerpes ruber sticticeps Baird

in Cope, U.S. Nat. Mus. Bull. 34, p. 178, 1889.

=*Pseudotriton ruber sticticeps* (Baird)

COTYPES (2): 11475 (2), Georgia? (restricted to Rabun County, Georgia), W. L. Jones.

Taricha laevis Baird and Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 302, 1853.

=*Taricha torosa torosa* (Rathke)

COTYPES (5): 4014 (5), San Francisco, San Francisco County, California, J. L. LeConte.

Thorius dubitus Taylor

Univ. Kansas Sci. Bull., vol. 27, p. 103, pl. 3, fig. 3, 1941.

PARATYPE: 134293 (formerly UIMNH 26829), summit of hill above Aculzingo, Veracruz, Mexico, E. H. Taylor, 1939.

Thorius maxillabrochus Gehlbach

Copeia, No. 3, p. 205, fig. 1c-d, 1959.

HOLOTYPE: 140293, ♂, 4 miles west of Zoquitlán, Puebla, Mexico, around 8,400 ft., R. H. Long, Jr. and F. C. Sibley, Aug. 12, 1954.

PARATYPE: 140294, same data.

Thorius narisovalis Taylor

Univ. Kansas Sci. Bull., vol. 26 (1939), p. 416, pl. 47, fig. 3, 1940.

PARATYPES (2): 134294-5 (formerly UIMNH 26853-4), Cerro San Felipe, Oaxaca, Mexico, E. H. Taylor, 1938.

Thorius pennatulus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 21, p. 111, 1869.

NEOTYPE: 111017, Cuautlapan, near Orizaba, Veracruz, Mexico, H. M. Smith, July 1940. See also Taylor, Univ. Kansas Sci. Bull., vol. 27, p. 107, 1941.

Thorius pulmonaris Taylor

Univ. Kansas Sci. Bull., vol. 26 (1939), p. 411, figs. 3-4, 1940.

PARATYPE: 139719 (formerly UIMNH 30983), Cerro San Felipe, Oaxaca, Mexico, E. H. Taylor, Aug. 22, 1938.

Thorius schmidti Gehlbach

Copeia, No. 3, p. 203, fig. 1a-b, 1959.

HOLOTYPE: 140295, ♂, 4 miles west of Zoquitlán, Puebla, Mexico, around 8,400 ft., F. C. Sibley, Aug. 14, 1954.

PARATYPES (5): 140296-300, same data.

Triton blasii de l'Isle

Ann. Sci. Nat. Zool., Ser. 4, vol. 17, p. 364, 1862.

=*Triturus marmoratus marmoratus* (Latreille)

COTYPES (2): 11011 (2), ♂, ♀, Nantes, France, MHNP.

Triton ensicauda Hallowell

Proc. Acad. Nat. Sci. Philadelphia, vol. 12 (1860), p. 494, 1861.

=*Triturus ensicaudus ensicaudus* (Hallowell)

COTYPE: (3): 7410 (3), ♂, ♂, ♀, Ousima (Amami, Ryukyu Islands), W. Stimpson. These specimens are also the cotypes of *Triton subcristatus* Hallowell.

Triton suberistatus Hallowell

Proc. Acad. Nat. Sci. Philadelphia, vol. 12 (1860), p. 494, 1861.

=*Triturus ensicaudus ensicaudus* (Hallowell)

COTYPES (3): 7410a-c, Ousima (Amami, Ryukyu Islands), W. Stimpson. These specimens are also the cotypes of *Triton ensicauda* Hallowell.

Triturus ensicaudus popei Inger

Chicago Nat. Hist. Mus., Fieldiana: Zool., vol. 32, p. 319, 1947.

PARATYPES (118): 7451 (2), ♂, ♀, Loo Choo (Ryukyu) Islands, W. Heine, Perry Expedition, May 1853. 36555, ♂, Okinawa, SCMT. 122118-33, Okinawa, W. B. S. Thomas, Oct. 5-20, 1945. 123032, Chizuka Canyon, Okinawa, R. Bohart, Sept. 14, 1945. 123033-130, Hentona, Okinawa, D. H. Johnson and O. A. Muennink, August-September 1945.

Triturus perstriatus Bishop

Occ. Pap. Mus. Zool. Univ. Michigan, No. 451, p. 3, 1941.

=*Diemictylus viridescens perstriatus* (Bishop)

PARATYPE: 118791 (formerly MZUM 74437), 3 miles southwest of Gainesville, Alachua County, Florida, A. Carr, Jr., Feb. 14, 1933.

Typhlomolge rathbuni Stejneger

Proc. U.S. Nat. Mus., vol. 18, p. 620, 1896.

HOLOTYPE: 22686, San Marcos, Hays County, Texas, U.S. Fish Commission, February 1896.

PARATYPES (6): 22687-9, 22691-3, San Marcos, Hays County, Texas, U.S. Fish Commission, Jan. 25-Mar. 1, 1896. 22690 was exchanged to IML in 1946.

Typhlotriton nereus Bishop

Copeia, No. 1, p. 1, figs. 1, 3b-b', 1944.

PARATYPES (7): 19778-9, 19781, 19784-5, 19787, Rock House Cave, Barry County, Missouri, G. E. Harris. 134296 (formerly UIMNH 27304), near Riverton, Cherokee County, Kansas, H. M. Smith et al., Mar. 26, 1931. 19778-9, 19781, 19784-5, and 19787 are also paratypes of *Typhlotriton spelaeus* Stejneger.

Typhlotriton spelaeus Stejneger

Proc. U.S. Nat. Mus., vol. 15, p. 116, pl. 9, 1892.

HOLOTYPE: 17903, Rock House Cave, Barry County, Missouri, F. A. Sampson, July 24, 1891.

PARATYPES (17): 17904, larva, same data. 17905-12, 19780, 19782-3, 19786, 19788-91, Rock House Cave, Barry County, Missouri, G. E. Harris, Sept. 21, 1891. 19776-7 was given to SNCBSH. 19778-9, 19781, 19784-5, and 19787 are also paratypes of *Typhlotriton nereus* Bishop.

Order Salientia

Acris gryllus paludicola Burger, Smith, and Smith

Journ. Tennessee Acad. Sci., vol. 24, p. 131, 1949.

PARATYPE: 129412 (formerly UIMNH 854), 5 miles west of Sabine Pass, Jefferson County, Texas, W. Robertson, May 2, 1948.

Acrodytes modesta Taylor and Smith

Proc. U.S. Nat. Mus., vol. 95, p. 594, pl. 27, fig. 2, pl. 28, figs. 2-3, 1945.

=*Phrynohyas modesta* (Taylor and Smith)

HOLOTYPE: 115013, ♂, Cruz de Piedra, near Acacoyagua, Chiapas, Mexico, Dr. and Mrs. H. M. Smith, Apr. 21, 1940.

PARATYPES (14): 115010-2, La Esperanza, Chiapas, Mexico, H. M. Smith, Apr. 7-June 3, 1940. 115014-9, 115021-3, Cruz de Piedra, near Acacoyagua, Chiapas, Mexico, H. M. Smith, Apr. 21-May 10, 1940. 115024-5, Colonia Soconusco, Chiapas, Mexico, H. M. Smith, May 4-9, 1940.

Agalychnis helenae Cope

Proc. Amer. Philos. Soc., vol. 22, p. 182, 1885.

HOLOTYPE: 13737, Nicaragua, J. F. Moser, Dec. 4, 1883.

Altirana parkeri Stejneger

Journ. Washington Acad. Sci., vol. 17, p. 318, 1927.

HOLOTYPE: 72328, ♂, Tingri, north of Mount Everest, Tibet, 15,000 ft., received from BMNH. Paratype 72327 was exchanged to WCUUM in 1947.

Ascaphus truei Stejneger

Proc. U.S. Nat. Mus., vol. 21, p. 900, text-fig. 1 and pl. 89, 1899.

=*Ascaphus truei truei* Stejneger.

HOLOTYPE: 25979, Humptulips, Gray's Harbor County, Washington, C. Rutter, Aug. 19, 1897.

Ascaphus truei californicus Mittleman and Myers

Proc. Biol. Soc. Washington, vol. 62, p. 63, 1949.

PARATYPES (5): 45362, northwest slope Craigy Peak, Siskiyou Mountains, Siskiyou County, California, N. Hollister, Sept. 26, 1909. 93779 (4 tadpoles), near Scotia, Humboldt County, California, G. S. Myers.

Ascaphus truei montanus Mittleman and Myers

Proc. Biol. Soc. Washington, vol. 62, p. 64, 1949.

HOLOTYPE: 102505, ♂, Lake McDonald, tributary to Lincoln Creek, Glacier National Park, Flathead County, Montana, L. P. Schultz and R. T. Smith, July 29, 1934.

PARATYPES (4, plus larvae, all from Glacier National Park, Flathead County, Montana): 102506-7, ♂, ♀, tributary to Lincoln Creek, L. P.

Schultz and R. T. Smith, July 29, 1934. 102508-9, juv., Lake Evangeline, L. P. Schultz, Aug. 16, 1934. 102510, larvae, Coal Creek, L. P. Schultz and L. Woodbury, July 24, 1934. 102511, larvae, Coal Creek, L. P. Schultz, July 23, 1934. 102512, larvae, 8 miles above mouth of Ole Creek, L. P. Schultz and L. Woodbury, June 28, 1934. 102513, larvae, Dutch Creek, L. P. Schultz and R. T. Smith, Aug. 19, 1934. 102514, larvae, 6 miles above Park Creek, L. P. Schultz and L. Woodbury, July 6, 1934. 102503 was exchanged to MP in 1949; 102504 to BM.

***Atelopus cruciger vogli* Müller**

Zool. Anz., vol. 108, p. 151, 1934.

PARATYPES (15): 107328-33, near Maracay, Venezuela, P. C. Vogely, 1933. 113265-66, 113268-74, Los Penas at Hacienda de la Trinidad, near Maracay, Venezuela, P. C. Vogely, 1933. 113267 was exchanged to MZUM in 1944.

***Atelopus gracilis* Barbour**

Bull. Mus. Comp. Zool., vol. 46, p. 102, 1905.

=*Atelopus elegans* (Boulenger)

COTYPES (2): 52401 (formerly MCZ 2423), Gorgona Island, Colombia, W. W. Brown, Jr., 1904. 118219, Gorgona Island, Colombia, W. W. Brown, Jr., 1904.

***Atelopus moreirae massarti* Cochran**

Bull. Mus. Hist. Nat. Belgique, vol. 24, No. 24, p. 1, 2 figs., 1948.

PARATYPES (2): 124585-6, Castanhal Grande near Obidos, Pará, Brazil, J. Massart, February 1922.

***Atelopus spurrelli certus* Barbour**

Occ. Pap. Mus. Zool. Univ. Michigan, No. 129, p. 12, 1923.

=*Atelopus varius certus* Barbour

PARATYPE: 118221 (formerly MCZ 9243), Mount Sapo, eastern Panama, T. Barbour and W. S. Brooks, 1922.

***Atelopus varius glyphus* Dunn**

Occ. Pap. Boston Soc. Nat. Hist., vol. 5, p. 396, 1931.

HOLOTYPE: 50230, Pirri Range, near head of Río Limon, Panama, Apr. 15, 1912, E. A. Goldman.

PARATYPES (36): 48594-5, Porto Obaldia, Panama, H. Pittier, Aug. 30, 1911. 50168-75, 50226, Cana, Panama, 1,800-2,000 ft., E. A. Goldman, March 1912. 50231-48, Pirri Range near head of Río Limon, Panama, 5,000 ft., E. A. Goldman, Apr. 15, 1912. 53964, Río Jappe near Cana, Panama, S. E. Meek and S. F. Hildebrand, Mar. 2, 1912. 54191-5, Cerro Azul, Panama, 1,500 ft., E. A. Goldman, Mar. 25, 1911. 54230, Cana, Panama, S. E. Meek and S. F. Hildebrand, Mar. 1, 1912. 53965, 54183-4, 53186-90 were exchanged to CNHM; 54185 was exchanged to LC, 1927, and 66316 to MP, 1949.

Batrachylodes trossulus Brown and Myers

Journ. Washington Acad. Sci., vol. 39, No. 11, p. 379, 1949.

HOLOTYPE: 119577, ♂, Torakina, Bougainville Island, Solomon Islands, W. L. Necker, July 12, 1944.

PARATYPES (6): 119586-8, Torakina, Bougainville Island, Solomon Islands, W. L. Necker, Aug. 24, 1944. 119787-9, Torakina, Bougainville Island, Solomon Islands, W. L. Necker, Sept. 3, 1944.

Borborocoetes stejneri Noble

Proc. Biol. Soc. Washington, vol. 37, p. 68, 1924.

=*Craspedoglossa stejneri* (Noble)

HOLOTYPE: 52608, ♂, Organ Mountains, Brazil, 1,500 m., J. N. Rose, Aug. 12, 1915.

Bufo aduncus Cope

Proc. U.S. Nat. Mus., vol. 11, p. 317, 1889.

=*Bufo woodhousei woodhousei* Girard

HOLOTYPE: 14100, probably from Gainesville, Cook County, Texas, G. H. Ragsdale, Apr. 9, 1885.

Bufo alvarius Girard

in Baird, Reptiles of the boundary, in Report U.S. and Mexican Bound. Surv., vol. 2, pt. 2, p. 26, pl. 41, figs. 1-6, 1859.

COTYPE: 2572, Fort Yuma (Imperial County), California, G. H. Thomas, 1855.

Bufo angustipes Taylor and Smith

Proc. U.S. Nat. Mus., vol. 95, p. 553, fig. 59, 1945.

HOLOTYPE: 116513, ♀, La Esperanza, Chiapas, Mexico, Dr. and Mrs. H. M. Smith, Apr. 7, 1940.

Bufo argillaceus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 20, p. 138, 1868.

=*Bufo marmoreus* Wiegmann

COTYPES (2): 27763-4, Colima, Mexico, J. Xantus.

Bufo auritus Cope

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8 (1875), p. 99, pl. 23, fig. 4, 1876. (Not of Raddi, 1821.)

=*Bufo gabbi* Taylor

COTYPE: 30676, east coast region, Costa Rica (probably Talamanca), W. M. Gabb.

Bufo beldingi Yarrow

Proc. U.S. Nat. Mus., vol. 5, p. 441, 1882.

=*Bufo punctatus* Baird and Girard

COTYPES (10): 12660 (5), 12670 (5), La Paz, Baja California, Mexico, L. Belding, 1882.

Bufo boreas Baird and Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 174, 1852.

=*Bufo boreas boreas* Baird and Girard

COTYPES (4): 15467-70, all ♂, Columbia River, Puget Sound (Washington), C. J. Pickering, May 1841.

Bufo boreas nelsoni Stejneger

North Amer. Fauna, No. 7, p. 220, pl. 3, figs. 4a-b, 1893.

HOLOTYPE: 18742, Oasis Valley, Nevada, F. Stephens, Mar. 16, 1891.

PARATYPES (8): 18738-41, same data. 18744-5, Resting Springs, Inyo County, California, A. K. Fisher, Feb. 7, 1891. 18746, Morans, Owens Valley, Inyo County, California, F. Stephens, July 4, 1891. 18747, Lone Pine, Owens Valley, Inyo County, California, E. W. Nelson, June 18, 1891. 18743 was exchanged to MCZ in 1928.

Bufo canaliferus Cope

Proc. Amer. Philos. Soc., vol. 17 (1878), p. 85, 1877.

COTYPES (10): 30315-24, western Tehuantepec, Oaxaca, Mexico, F. Sumichrast.

Bufo coccifer Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 18, p. 130, 1866.

HOLOTYPE: 6490, Arriba, Costa Rica, C. N. Riotta.

Bufo columbiensis Baird and Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 378, 1853.

=*Bufo boreas boreas* Baird and Girard

COTYPES (7): 4182 (7), Columbia River, Oregon, C. J. Pickering, 1841.

Bufo coniferus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 158, 1862.

HOLOTYPE: 4335, Turbo, New Granada (Colombia), A. Schott.

Bufo copei Yarrow and Henshaw

Report upon the collections of reptiles and batrachians collected during the years of 1875, 1876, and 1877 in California, Arizona, and Nevada, in Annual Report [of the Corps of Engineers] upon the U.S. Geogr. Surv. . . . West of the 100th Mer. . . ., Appendix L, p. 207, 1878.

=*Bufo terrestris copei* Yarrow and Henshaw

COTYPES (64): 5388 (17), 35925-71 (formerly 5937), James Bay, Hudson Bay (Canada), C. Drexler, June 1860.

Bufo debilis Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 7, p. 87, 1854.

=*Bufo debilis debilis* Girard

COTYPES (7): 2621 (7), juv., Matamoros, Tamaulipas, Mexico, D. N. Couch.

Bufo dorsalis Hallowell

Proc. Acad. Nat. Sci. Philadelphia, vol. 6 (1852), p. 181, 1853. (Not of Spix, 1824.)

=*Bufo woodhousei woodhousei* Girard, new name for *Bufo dorsalis* Hallowell

HOLOTYPE: 2531, San Francisco Mountains, Arizona (New Mexico in original description), S. W. Woodhouse.

Bufo dunni Barbour

Occ. Pap. Boston Soc. Nat. Hist., vol. 5, p. 192, pl. 13, figs. 1-2, 1926.

PARATYPES (3): 71160, 71552-3 (formerly MCZ 11083, 11085), Mina Carlota near Cumanayagua, Cuba, E. R. Dunn, July 1925.

Bufo exsul Myers

Occ. Pap. Mus. Zool. Univ. Michigan, No. 460, p. 3, pls. 1-3, 1942.

=*Bufo boreas exsul* Myers

PARATYPES (2): 104314-5, Deep Springs, Inyo County, California, G. F. Ferris, G. S. Myers, and I. L. Wiggins, May 1, 1937.

Bufo gabbi Taylor

Univ. Kansas Sci. Bull., vol. 35, p. 619, fig. 6, 1952. (New name for *Bufo auritus* Cope, preoccupied.)

COTYPE: 30676, probably Talamanca, Costa Rica, W. M. Gabb.

Bufo gemmifer Taylor

Univ. Kansas Sci. Bull., vol. 26 (1939), p. 490, pl. 53, figs. 3, 3a-b, 1940.

PARATYPE: 135257 (formerly UIMNH 25133), El Limoncito near La Venta, Guerrero, Mexico, E. H. Taylor, June 29, 1939.

Bufo granulosus Baird and Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 173, 1852. (Not of Spix, 1824.)

=*Bufo valliceps* Wiegmann

HOLOTYPE: 2595, Texas (original description: "One specimen was collected on the route between Indianola and San Antonio"), J. H. Clark. This specimen was renamed *Bufo nebulifer* by Girard, Proc. Acad. Nat. Sci. Philadelphia, vol. 7, p. 87, 1854.

Bufo güntneri Cochran

U.S. Nat. Mus. Bull. 177, p. 8, fig. 1, 1941.

HOLOTYPE: 59081, Port-au-Prince, Haiti, C. Bencomo, 1916.

PARATYPES (2): 59082, same data. 75993, Cape Haitien, Haiti, W. M. Mann, 1913.

Bufo haematiticus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 157, 1862.

COTYPES (2): 48448-9, Truandó, New Granada (Colombia), A. Schott. These very young specimens were reentered from 4344, of which the remaining one is the type of *Bufo haematiticus lachrymans* Cope.

Bufo haematiticus lachrymans Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 158, 1862.

=*Bufo haematiticus* Cope

COTYPE: 4344, Truandó, New Granada (Colombia), A. Schott.

Bufo halophila Baird and Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 301, 1853.

=*Bufo boreas halophilus* Baird and Girard

COTYPES (2): 2589 (2), Benicia, Solano County, California, J. L. LeConte.

Bufo hemiophrys Cope

Proc. Amer. Philos. Soc., vol. 23, p. 515, 1886.

=*Bufo woodhousei hemiophrys* Cope

COTYPES (7): 11927 (7), Pembina, Pembina County, North Dakota (northern boundary of Montana in original description), E. Coues, June-July 1874. One cotype was exchanged to MCZ in 1915.

Bufo houstonensis Sanders

Herpetologica, vol. 9, p. 27, figs. 1-3, 5, 7, 1953.

PARATYPES (4): 134433-6, Fairbanks, Harris County, Texas, J. C. Wottring and W. J. Greer, May 18, 1952.

Bufo ictericus mertensi Cochran

Journ. Washington Acad. Sci., vol. 40, p. 238, 1950.

HOLOTYPE: 105244, Itá, Nova Teutonia, Santa Catarina, Brazil, F. Plaumann, 1938.

PARATYPES (72): 98777-8, São Bento, Santa Catarina, Brazil, B. Behr, 1935. 103842, 103844-50, 103852-906, 105245-51, Itá, Nova Teutonia, Santa Catarina, Brazil, F. Plaumann, 1937-38. 103843 was exchanged to SNG in 1951; 103851 to CNHM, 1953.

Bufo inca Stejneger

Proc. U.S. Nat. Mus., vol. 45, p. 541, 1913.

HOLOTYPE: 49557, Huadquiña, Peru, 5,000 ft., Yale-Peruvian Expedition, August 1911.

Bufo insidios Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 7, p. 88, 1854.

=*Bufo debilis insidios* Girard

COTYPES (2): 2622 (2), Chihuahua, Mexico, T. H. Webb.

Bufo kelloggi Taylor

Univ. Kansas Sci. Bull., vol. 24 (1936), p. 510, pl. 45, figs. 1-3, 1938.

=*Bufo debilis kelloggi* Taylor

PARATYPE: 134258 (formerly UIMNH 23461), 2 miles east of Mazatlán, Sinaloa, Mexico, E. H. Taylor, July 24, 1934.

Bufo lamentor Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 11 (1859), p. 169, 1860.

=*Bufo boreas boreas* Baird and Girard

COTYPES (4): 4194 (4), Fort Bridger (Uinta County), Wyoming (Utah Territory in original description), C. Drexler, 1858.

Bufo lentiginosus pachycephalus Cope

U.S. Nat. Mus. Bull. 34, p. 288, 1889.

=*Bufo terrestris terrestris* (Bonnaterre)

COTYPES (2): 14681 (2), Micanopy, Alachua County, Florida, T. H. Bean.

Bufo longinasus Stejneger

Proc. U.S. Nat. Mus., vol. 28, p. 765, figs. 1-6, 1905.

HOLOTYPE: 27419, ♂, El Guamá, Pinar del Rio, Cuba, W. Palmer and J. H. Riley, Mar. 9, 1900.

Bufo lugubrosus Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 424, 1853.

=*Bufo spinulosus* Wiegmann

HOLOTYPE: 16359, Valparaiso, Chile, J. Drayton, January 1839.

Bufo melanochloris Cope

Proc. Amer. Philos. Soc., vol. 17 (1878), p. 85, 1877.

HOLOTYPE: 30592, eastern Costa Rica, W. M. Gabb.

Bufo microscaphus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 18 (1866), p. 301, 1867.

COTYPES (2): 4184, 132901, upper Colorado region, Arizona (restricted to Fort Mohave, Mohave County), H. B. Mollhausen. 4184 is the lectotype.

Bufo minshanicus Stejneger

Journ. Washington Acad. Sci., vol. 16, p. 446, 1926.

HOLOTYPE: 68567, Choni, on Tao River, Kansu, China, F. R. Wulsin, Aug. 24-31, 1923.

PARATYPES (27): 67779, 67781, 67783-6, 67810-2, 68808-10, Songpan, Szechwan, China, D. C. Graham, July 24-29, 1924. 68549-50, 68552-3, 68555, 68557-66, Choni, on Tao River, Kansu, China, F. R. Wulsin,

Aug. 24-31, 1923. 67780 and 67782 were exchanged to WCUUM in 1947; 68551 was exchanged to MCZ in 1944; 68554 to ZML, 1927; 68556 to MZUM, 1944.

Bufo monksiae Cope

Proc. Amer. Philos. Soc., vol. 18, p. 263, 1879.

=*Bufo occidentalis* Camerano

HOLOTYPE: 9896, Guanajuato, Mexico, A. Dugès, 1877.

Bufo nayaritensis Taylor

Univ. Kansas Sci. Bull., vol. 29, p. 349, pl. 26, figs. 1-1a, 1943.

=*Bufo mazatlanensis nayaritensis* Taylor

PARATYPE: 134259 (formerly UIMNH 24948), Tepic, Nayarit, Mexico, E. H. Taylor, July 31, 1934.

Bufo nebulifer Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 7, p. 87, 1854. (New name for *Bufo granulosus* Baird and Girard preoccupied.)

=*Bufo valliceps* Wiegmann

HOLOTYPE: 2595, Texas (original description: "One specimen was collected on the route between Indianola and San Antonio"), J. H. Clark. This specimen is the type of *Bufo granulosus* Baird and Girard (not of Spix, 1824).

Bufo paracnemis Lutz

Compt. Rend. Soc. Biol. Paris, vol. 93, p. 213, 1925.

COTYPES (2): 97238-9, ♂, ♀, Belo Horizonte, Minas Gerais, Brazil, A. Lutz, 1923.

Bufo perplexus Taylor

Univ. Kansas Sci. Bull., vol. 29, p. 347, pl. 27, figs. 1-2, 1943.

PARATYPE: 139720 (formerly UIMNH 21705), Tonolapam, Guerrero, Mexico, E. H. Taylor, July 18, 1936.

Bufo pictus Cope

in Yarrow, Report upon the collections of batrachians and reptiles . . . 1871, 1872, 1873, and 1874, in Report U.S. Geogr. and Geol. Expl. and Surv. West of One Hundr. Mer., vol. 5, p. 522, pl. 25, figs. 4-5, 1875.

=*Bufo boreas boreas* Baird and Girard

COTYPES (4): 8655, 25332-4, Utah, H. C. Yarrow, 1872.

Bufo politus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 158, 1862.

HOLOTYPE: 5600, Greytown, Nicaragua, H. C. Caldwell.

Bufo punctatus Baird and Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 173, 1852.

COTYPES (3): 2618 (3), Rio San Pedro of the Rio Grande (Devils River, Val Verde County), Texas, J. H. Clark under Col. J. D. Graham.

Bufo pygmaeus Myers and Carvalho

Zoologica, New York, vol. 37, p. 1, 1952.

PARATYPES (2): 132356-7, São João da Barra, Rio de Janeiro, Brazil, A. L. de Carvalho and L. de Castro Faria, December 1948.

Bufo rubescens Lutz

Compt. Rend. Soc. Biol. Paris, vol. 93, p. 214, 1925.

=*Bufo rufus* Garman

COTYPES (2): 97236-7, ♀, ♀, Belo Horizonte, Minas Gerais, Brazil, A. Lutz, May 1925.

Bufo smithi Stejneger

U.S. Nat. Mus. Bull. 58, p. 64, 1907.

HOLOTYPE: 31951, ♂, Kochi, Province of Toza, Shikoku, Japan, Hugh M. Smith, May 11, 1903.

PARATYPES (8): 31851, ♂, Suzaki, Toza, Shikoku, Japan, Hugh M. Smith, May 8, 1903. 31929, 31943, 31945-49, Kochi, Toza, Shikoku, Japan, Hugh M. Smith, May 11, 1903. 31950 was exchanged to MZUM in 1944; 31942 and 31944 were exchanged to MCZ in 1919 and 1944.

Bufo speciosus Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 7, p. 86, 1854.

=*Bufo compactilis speciosus* Girard

COTYPES (4): 2608, Ringgold Barracks (Rio Grande City, Star County), Texas, A. Schott. 2610, Brownsville, Cameron County, Texas, S. Van Vliet. 2611, 131559, Pesquiería Grande, Nuevo León, Mexico, D. N. Couch, April 1853.

Bufo tacanensis Smith

Copeia, No. 3, p. 176, pl. 1, 1952.

PARATYPE: 139721 (formerly UIMNH 24874), Volcán de Tacaná, above Cacahuatán, Chiapas, Mexico, C. J. and M. Goodnight, August 1950.

Bufo terrestris charlesmithi Bragg

Wasmann Journ. Biol., vol. 12, p. 245, figs. 1-4, 1954.

PARATYPE: 134260, 1.8 miles south and 7 miles east of Norman, Cleveland County, Oklahoma, A. N. Bragg, May 9, 1939.

Bufo woodhousii Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 7, p. 86, 1854. (New name for *Bufo dorsalis* Hallowell, preoccupied.)

—*Bufo woodhousii woodhousii* Girard

HOLOTYPE: 2531, San Francisco Mountains, Arizona, S. W. Woodhouse.

PARATYPES? (2): 4184 (2), Upper Colorado River region, Arizona, H. B. Möllhausen.

Bufo woodhousii velatus Bragg and Sanders

Wassman Journ. Biol., vol. 9, p. 366, pl. 1, figs. 1–4, 1951.

HOLOTYPE: 131869, ♂, Elkhart, Anderson County, Texas, O. Sanders and R. M. Sanders, Mar. 8, 1951.

PARATYPES (8): 131870–7, same data.

Centrolenella viridissima Taylor

Univ. Kansas Sci. Bull., vol. 28, p. 75, pl. 9, figs. 2–2b, 1942.

PARATYPE: 137198, Agua del Obispo, Guerrero, Mexico, E. H. Taylor, Aug. 27, 1941.

Cerathyla panamensis Stejneger

Proc. Biol. Soc. Washington, vol. 30, p. 31, 1917.

HOLOTYPE: 55320, ♀, Signal Loma, north coast 3 miles south of San Isabel, Panama, C. G. Holland, May 20, 1915.

PARATYPE: 55321, juv., same data.

Ceratophrys fusceiventris Lutz

Compt. Rend. Soc. Biol. Paris, vol. 95, p. 1012, 1926.

—*Zachaenus parvulus* Girard

COTYPE: 96313, Tijuca, Rio de Janeiro, Brazil, J. Venancio, Jan. 10, 1926.

Cochranella albotunica Taylor and Cochran

Univ. Kansas Sci. Bull., vol. 35, p. 1648, 1953.

HOLOTYPE: 96559, Serra da Bocaina, boundary of São Paulo and Rio de Janeiro, Brazil, A. Lutz, Jan. 2–19, 1930.

PARATYPES (2): 96557, same data. 96723, Bonito, Serra da Bocaina, boundary of São Paulo and Rio de Janeiro, Brazil, A. Lutz, March 1934.

Cochranella delicatissima Taylor and Cochran

Univ. Kansas Sci. Bull., vol. 35, p. 1640, 1953.

HOLOTYPE: 96481, Angra dos Reis, Rio de Janeiro, Brazil, A. Lutz.

Cochranella divaricans Taylor and Cochran

Univ. Kansas Sci. Bull., vol. 35, p. 1643, 1953.

HOLOTYPE: 96562, ♂, Serra da Bocaina, boundary of São Paulo and Rio de Janeiro, Brazil, A. Lutz, Jan. 2–19, 1930.

Cochranella dubia Taylor and Cochran

Univ. Kansas Sci. Bull., vol. 35, p. 1644, 1953.

HOLOTYPE: 96722, Bonito, Serra de Bocaina, boundary of São Paulo and Rio de Janeiro, Brazil, A. Lutz, December 1930–February 1931.

Cochranella lutzorum Taylor and Cochran

Univ. Kansas Sci. Bull., vol. 35, p. 1638, 1953.

HOLOTYPE: 101134, Independencia, Petrópolis, Rio de Janeiro, Brazil, D. M. Cochran, B. Lutz, and J. Venancio, May 1935.

Cochranella petropolitana Taylor and Cochran

Univ. Kansas Sci. Bull., vol. 35, p. 1637, 1953.

HOLOTYPE: 101135, Petrópolis, Rio de Janeiro, Brazil, A. and B. Lutz and D. M. Cochran, May 1935.

Cochranella surda Taylor and Cochran

Univ. Kansas Sci. Bull., vol. 35, p. 1630, fig. 1, 1953.

HOLOTYPE: 96916, Passa Quatro, Minas Gerais, Brazil, Zikan, November 1920.

PARATYPE: 96917, same data.

Cornufer worcesteri Stejneger

Proc. U.S. Nat. Mus., vol. 28, p. 345, 1905.

HOLOTYPE: 34784, Mount Apo, Mindanao, Philippine Islands, E. A. Mearns.

Corythomantis adpersa Lutz

Compt. Rend. Soc. Biol. Paris, vol. 93, p. 213, 1925.

=*Aparasphenodon bruno* Miranda-Ribeiro

COTYPE: 96413, Lagoa de Paquarema, Rio de Janeiro, Brazil, A. Lutz, May 29, 1924.

Cranopsis fastidiosus Cope

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8, p. 96, pl. 23, fig. 1, 1875.

=*Bufo fastidiosus* (Cope)

LECTOTYPE: 32585, Pico Blanco, District of Uren, Costa Rica, 2,500 ft., W. M. Gabb.

PARATYPES (3): 32584, 32586-7, same data.

Craspedoglossa sanctae-catharinae Müller

Blatt. Aquar. Terr. Kunde, vol. 33, p. 168, figs. 1-3, 1922.

PARATYPE: 103186, Rio Novo, Santa Catarina, Brazil, W. Erhardt, November-December 1917.

Crossodactylodes pinto Cochran

Proc. Biol. Soc. Washington vol. 51, p. 42, 1938.

PARATYPES (3): 102607-8, 102611, Macaé, Rio de Janeiro, Brazil, MP. 102609 was exchanged to MCZ in 1944; 102610 to LC, 1944. The type assigned to temporary USNM number 102606 has been returned to DZ.

Crossodactylus dispar Lutz

Compt. Rend. Soc. Biol. Paris, vol. 93, p. 138, 1925.

COTYPES (3): 96738-40, Bonito, Serra da Bocaina, boundary of São Paulo and Rio de Janeiro, Brazil, A. Lutz, Jan. 20, 1925.

Cyclorhamphus boulengeri Lutz

Mem. Inst. Oswaldo Cruz, vol. 22, pt. 1, pp. 11, 23, pl. 3, figs. 1-2, 1929.

=*Cycloramphus neglectus* Lutz

COTYPES (3): 97179-81, São Bento, Santa Catarina, Brazil, A. Lutz, 1929.

Cyclorhamphus granulatus Lutz

Mem. Inst. Oswaldo Cruz, vol. 22, pp. 13, 24, pl. 3, figs. 3-4, 1929.

=*Cycloramphus granulatus* Lutz

COTYPE: 96742, Bonito, Serra da Bocaina, boundary of São Paulo and Rio de Janeiro, Brazil, A. Lutz, February 1928.

Cystignathus labialis Cope

Proc. Amer. Philos. Soc., vol. 17 (1877), p. 90, 1878.

=*Leptodactylus labialis* (Cope)

HOLOTYPE: 31302, probably Tehuantepec, Oaxaca, Mexico, F. Sumichrast.

PARATYPES (5): 31300-1, 31303-5, same data.

Cystignathus microtis Cope

Proc. Amer. Philos. Soc., vol. 13, p. 265, 1879.

=*Leptodactylus melanonotus* (Hallowell)

HOLOTYPE: 9906, Guanajuato, Mexico, A. Dugès, 1877.

PARATYPES (2): 9908-9, same data.

Cystignathus perlaevis Cope

Proc. Amer. Philos. Soc., vol. 13, p. 269, 1879.

=*Leptodactylus melanonotus* (Hallowell)

HOLOTYPE: 10041, from a well near Tapana, Tehuantepec, Oaxaca, Mexico, F. Sumichrast.

Cystignathus poecilochilus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 156, 1862.

=*Leptodactylus poecilochilus* (Cope)

HOLOTYPE: 4347a, near Turbo, New Granada (Colombia), A. Schott.

Diaglena reticulata Taylor

Univ. Kansas Sci. Bull., vol. 28, p. 60, figs. 1-1c, pl. 5, fig. 1, 1942.

HOLOTYPE: 115500, Cerro Arenal, 30 km. northwest of Tehuantepec, Oaxaca, Mexico, T. MacDougall, Jan. 2, 1940.

***Eleutherodactylus abbotti* Cochran**

Proc. Biol. Soc. Washington, vol. 36, p. 93, 1923.

HOLOTYPE: 65055, Laguna, Samaná Peninsula, Dominican Republic, W. L. Abbott, May 11–15, 1922.

PARATYPES (2): 65056–7, same data.

***Eleutherodactylus albipes* Barbour and Shreve**

Bull. Mus. Comp. Zool., vol. 80, p. 383, 1937.

PARATYPE: 103376, Pico Turquino, Oriente, Cuba, 5,400–6,000 ft., P. J. Darlington, June 1936.

***Eleutherodactylus alticola* Lynn**

Herpetologica, vol. 1, p. 89, pl. 8, figs. 2a–d, 1937.

HOLOTYPE: 102524, Blue Mountain Peak, St. Thomas Parish, Jamaica, W. G. Lynn, July 14, 1936.

PARATYPES (20): 102525, 102527–39, 102541, 102543–45, 102547, Blue Mountain Peak, St. Thomas Parish, Jamaica, W. G. Lynn, July 11–14, 1926. 102548, Sir John Peak, St. Thomas Parish, Jamaica, W. G. Lynn, June 7, 1936. 102540 was exchanged to MZUM in 1944; 102542 to MCZ, 1937; 102526 and 102546 were exchanged to R in 1953.

***Eleutherodactylus andrewsi* Lynn**

Herpetologica, vol. 1, p. 88, pl. 8, figs. 1a–1b, 1937.

HOLOTYPE: 102515, ♂, Chester Vale, St. Andrew Parish, Jamaica, H. N. Lynn, June 21, 1936.

PARATYPES (6): 102516–7, Chester Vale, St. Andrew Parish, Jamaica, W. G. Lynn, July 2, 5, 1936. 102519–20, Clydesdale, St. Andrew Parish, Jamaica, W. G. Lynn, June 13, 20, 1936. 102522, Catherine's Peak, St. Andrew Parish, Jamaica, W. G. Lynn, June 19, 1936. 102523, Morce's Cap, St. Andrew Parish, Jamaica, W. G. Lynn, June 21, 1936. 102518 was exchanged to MCZ in 1937; 102521 to MZUM, 1944.

***Eleutherodactylus atkinsi orientalis* Barbour and Shreve**

Bull. Mus. Comp. Zool., vol. 80, p. 382, pl. 3, fig. 2, 1937.

PARATYPES (2): 139346–7 (formerly MCZ 22073–4), El Yunque de Baracoa, Oriente, Cuba, P. J. Darlington, July 13–14, 1936.

***Eleutherodactylus audanti* Cochran**

Occ. Pap. Boston Soc. Nat. Hist., vol. 8, p. 164, 1934.

PARATYPES (7): 72595–7, Morne Cabaio, Massif de la Selle, Haiti, A. Wetmore, Apr. 10, 1927. 85009, Morne La Selle Range, Haiti, 7,000 ft., S. S. Cook, 1932. 95111–3, Peak La Selle, Haiti, A. Audant.

***Eleutherodactylus avocalis* Taylor and Smith**

Proc. U.S. Nat. Mus., vol. 95, p. 580, 1945.

HOLOTYPE: 116885, ♀, Tres Cruces, near Tehuantepec, Oaxaca, Mexico, Dr. and Mrs. H. M. Smith, March 1940.



PARATYPES (10): 116876-82, Tres Cruces, near Tehuantepec, Oaxaca, Mexico, H. M. Smith, Jan. 14, 1940. 116886, Tehuantepec, Oaxaca, Mexico, H. M. Smith, March 1940. 116887-8, Cerro Arenal, Oaxaca, Mexico, H. M. Smith, Jan. 19, Jan. 2, 1940. 116883-4 were exchanged to MCZ in 1957.

Eleutherodactylus bakeri Cochran

Proc. Boston Soc. Nat. Hist., vol. 40, p. 369, 1935.

PARATYPES (4): 95418-21, Mount La Hotte, La Selle Range, Haiti, 5,000-8,000 ft., P. J. Darlington, October 1934.

Eleutherodactylus beebei Cochran

Zoologica, New York, vol. 41, p. 11, text-fig. 1, 1956.

HOLOTYPE: 129526, Kartabo, British Guiana, W. Beebe, June 22, 1924.

Eleutherodactylus binghami Stejneger

Proc. U.S. Nat. Mus., vol. 45, p. 542, 1913.

=*Gastrotheca boliviana* (Steindachner)

HOLOTYPE: 49559, Cuzco, Peru, 11,500 ft., Yale-Peruvian Expedition, July 9, 1911.

PARATYPE: 49560, same data.

Eleutherodactylus casparii Dunn

Occ. Pap. Boston Soc. Nat. Hist., vol. 5, p. 215, 1926.

=*Eleutherodactylus ricordii casparii* Dunn

PARATYPE: 118204 (formerly MCZ 11439), Mina Carlota, near Cumana-yagua, Cuba, E. R. Dunn, August 1925.

Eleutherodactylus cavernicola Lynn

Journ. Washington Acad. Sci., vol. 44, p. 400, figs., 1954.

HOLOTYPE: 134239, cave on Portland Ridge, Clarendon, Jamaica, W. G. Lynn, June 28, 1953.

PARATYPES (16): 135242-57, cave on Portland Ridge, Clarendon, Jamaica, W. G. Lynn, Oct. 24, 1952-July 28, 1953. 135240-1 were exchanged to MCZ in 1957.

Eleutherodactylus conspicuus Taylor and Smith

Proc. U.S. Nat. Mus., vol. 95, p. 567, fig. 60a, 1945.

HOLOTYPE: 116509, ♀, Piedras Negras, Petén, Guatemala, Dr. and Mrs. H. M. Smith, May 29, 1939.

PARATYPES (5): 116506-8, 116510-11, Piedras Negras, Petén, Guatemala, H. M. Smith, May 24-June 10, 1939.

Eleutherodactylus cubanus Barbour

Copeia No. 3, p. 179, 1942. (New name for *Eleutherodactylus parvus* Barbour and Shreve, 1937, not of Girard, 1853.)

PARATYPES (2): 139344-5 (formerly MCZ 21657-8), Pico Turquino, Oriente, Cuba, 1,500-4,000 ft., P. J. Darlington, June 1936.

Eleutherodactylus cunctator Dunn

Proc. U.S. Nat. Mus., vol. 74, No. 2770, p. 2, 1928.

=*Eleutherodactylus orcutti* Dunn

HOLOTYPE: 73865, ♀, Arntully, St. Thomas Parish, Jamaica, C. R. Orcutt, August 1927.

Eleutherodactylus darlingtoni Cochran

Proc. Boston Soc. Nat. Hist., vol. 40, p. 368, 1935.

PARATYPE: 95422, La Visite, La Selle Range, Haiti, 5,000–7,000 ft., P. J. Darlington, Sept. 16–23, 1923.

Eleutherodactylus dorsoconcolor Taylor

Univ. Kansas Sci. Bull., vol. 27, p. 152, pl. 10, 1941.

HOLOTYPE: 110619, ♀, Tequeyutepec, Veracruz, Mexico, H. M. Smith. Mar. 23, 1940.

PARATYPES (3): 110615, 110617–8, same data. 110616 was exchanged to MCZ in 1944.

Eleutherodactylus eileenae Dunn

Occ. Pap. Boston Soc. Nat. Hist., vol. 5, p. 212, 1926.

PARATYPE: 27862, San Diego de los Baños, Cuba, W. Palmer and J. H. Riley, Apr. 16, 1900.

Eleutherodactylus footei Stejneger

Proc. U.S. Nat. Mus., vol. 45, p. 543, 1913.

HOLOTYPE: 49562, Tincocochaca (probably Tincocochaca), Peru, 7,000 ft., Yale-Peruvian Expedition, Aug. 8, 1911.

PARATYPES (2): 49561, 49563, juv., Cuzco, Peru, 11,500 ft., Yale-Peruvian Expedition, July 9, 1911.

Eleutherodactylus fuscus Lynn

Copeia, No. 4, p. 235, fig. 1, 1943.

HOLOTYPE: 115976, ♂, Bush Mountain at base of Dolphin Head, Westmoreland Parish, Jamaica, W. G. Lynn, Aug. 12, 1941.

PARATYPES (2): 115977, ♂, same data. 115978, ♂, Quickstep, St. Elizabeth Parish, Jamaica, W. G. Lynn, Aug. 9, 1941.

Eleutherodactylus goldmani Noble

Proc. Biol. Soc. Washington, vol. 37, p. 67, 1924.

=*Eleutherodactylus gollmeri* (Peters)

HOLOTYPE: 54033, ♂, Cerro Brujo, Panama, E. A. Goldman, June 7, 1911.

PARATYPE: 54034, juv., same data.

Eleutherodactylus gossei Dunn

Proc. Boston Soc. Nat. Hist., vol. 38, p. 118, 1926.

PARATYPES (2): 118212-3 (formerly MCZ 10169), Balaclava, St. Elizabeth, Jamaica, 700 ft., A. E. Wright, Apr. 29, 1909.

Eleutherodactylus grabhami Dunn

Proc. Boston Soc. Nat. Hist., vol. 38, p. 121, 1926.

PARATYPE: 30700, Pratville District, Clarendon, Jamaica, C. B. Taylor, March 1902.

Eleutherodactylus greyi Dunn

Occ. Pap. Boston Soc. Nat. Hist., vol. 5, p. 213, 1926.

PARATYPES (2): 118205-6 (formerly MCZ 11510), Mina Carlota, near Cumanayagua, Cuba, E. R. Dunn, July 1925.

Eleutherodactylus intermedius Barbour and Shreve

Bull. Mus. Comp. Zool., vol. 80, p. 384, pl. 4, figs. 3-4, 1937.

PARATYPES (2): 139342-3 (formerly MCZ 21974), near Cueva del Aura, Pico Turquino, Oriente, Cuba, 1,500-4,000 ft., P. J. Darlington, June 1936.

Eleutherodactylus intermedius Cochran

U.S. Nat. Mus. Bull. 177, p. 70, fig. 22, 1941.

=*Eleutherodactylus haitianus* Barbour and Shreve, new name for *Eleutherodactylus intermedius* Cochran, preoccupied

HOLOTYPE: 107566, ♂, Loma Rucilla, Dominican Republic, 8,000-10,000 ft., P. J. Darlington, June 1938.

PARATYPES (17): 107567, 107569-74, Loma Rucilla, Dominican Republic, P. J. Darlington, June 1938. 107575-6, Loma Vieja, Dominican Republic, 6,000 ft., P. J. Darlington, August 1938. 107578-85, Valle Nuevo, southeast of Constanza, Dominican Republic, 6,000-8,000 ft., P. J. Darlington, August 1938. 107568 and 107577 were exchanged to MZUM in 1943.

Eleutherodactylus jugans Cochran

Journ. Washington Acad. Sci., vol. 27, p. 312, 1937. (New name for *Leptodactylus darlingtoni* Cochran, preoccupied.)

PARATYPES (8): 95423-7, La Visite, La Selle Range, Haiti, P. J. Darlington, Sept. 16-23, 1934. 95428-30, from Mount La Hotte, La Selle Range, Haiti, P. J. Darlington, October 1934, were described by Shreve as *Eleutherodactylus ventrilineatus*, in Proc. New England Zool. Club, vol. 15, p. 98, 1936.

Eleutherodactylus junori Dunn

Proc. Boston Soc. Nat. Hist., vol. 38, p. 120, 1926.

PARATYPE: 118214 (formerly MCZ 11351), Spaldings, Clarendon Parish, Jamaica, 2,900 ft., P. Junor, August 1925.

***Eleutherodactylus karlschmidti* Grant**

Copeia, No. 2, p. 55, 1931.

PARATYPES (2): 118647-8 (formerly MCZ 73432), 10 km. south of Canóvanas, Puerto Rico, C. Grant, Mar. 21, 1931.

***Eleutherodactylus lewisi* Lynn and Dent**

Herpetologica, vol. 2, p. 72, 1942.

=*Eleutherodactylus luteolus* (Gosse)

HOLOTYPE: 115435, ♀, near Bush Mountain at base of Dolphin Head, Westmoreland Parish, Jamaica, W. G. Lynn, Aug. 12, 1941.

PARATYPES (5): 115436-9, 115442, same data. 115440 was exchanged to MZUM in 1944; 115441 to MCZ, 1944.

***Eleutherodactylus lymani* Barbour and Noble**

Bull. Mus. Comp. Zool., vol. 63, p. 403, 1920.

PARATYPE: 118189 (formerly MCZ 5432), Bellavista, Peru, G. K. Noble, Sept. 23-26, 1916.

***Eleutherodactylus lynni* Goin and Cooper**

Occ. Pap. Mus. Inst. Jamaica, No. 4, p. 4, 1 pl., 1950.

HOLOTYPE: 127976, ♀, Sweetwater, near Horse Guards Road, Saint James Parish, Jamaica, C. J. Goin, July 5, 1948.

PARATYPE: 117735, John's Hall, Saint James Parish, Jamaica, W. G. Lynn, Aug. 11, 1941.

***Eleutherodactylus matudai* Taylor**

Univ. Kansas Sci. Bull., vol. 27, p. 154, pl. 11, 1941.

HOLOTYPE: 110626, ♀, Mount Ovando, Chiapas, Mexico, H. M. Smith, Apr. 16, 1940.

PARATYPES (9): 110620-3, 110625, 110627-30, same data. 110624 was exchanged to MCZ in 1944.

***Eleutherodactylus megalops* Ruthven**

Occ. Pap. Mus. Zool. Univ. Michigan, No. 39, p. 3, pl. 1, figs. 1-2, 1917.

PARATYPE: 64026, (formerly MZUM 48548), San Lorenzo, Santa Marta Mountains, Colombia, A. G. Ruthven, July 7, 1913.

***Eleutherodactylus milesi* Schmidt**

Field Mus. Nat. Hist., Zool. Ser., vol. 20, p. 18, 1933.

PARATYPE: 118202 (formerly MCZ 17436), mountains west of San Pedro, Honduras, K. P. Schmidt, Mar. 31, 1923.

***Eleutherodactylus nubicola* Dunn**

Proc. Boston Soc. Nat. Hist., vol. 38, p. 116, 1926.

PARATYPE: 118236 (formerly MCZ 9227), Bocas del Toro, Gutiérrez, Panama, E. R. Dunn and C. B. Duryea, 1923.

***Eleutherodactylus orcutti* Dunn**

Proc. U.S. Nat. Mus., vol. 74, No. 2767, p. 1, 1928.

HOLOTYPE: 73866, ♀, Arntully, Saint Thomas Parish, Jamaica, C. R. Orcutt, August 1927.

PARATYPES (3): 73867-9, same data. 73870 was exchanged to MCZ in 1944.

***Eleutherodactylus pantoni* Dunn**

Proc. Boston Soc. Nat. Hist., vol. 38, p. 119, 1926.

PARATYPES (3): 35861-2, Holly Mount, Mount Diablo, Jamaica, W. Harris, Aug. 18, 1905. 30699, Mandeville, Manchester, Jamaica, C. B. Taylor, May 1902.

***Eleutherodactylus pinarensis* Dunn**

Occ. Pap. Boston Soc. Nat. Hist., vol. 5, p. 213, 1926.

PARATYPES (19): 26651-2, 27860-1, San Diego de los Baños, Cuba, W. Palmer and J. H. Riley, Apr. 14-16, 1900. 27417-8, El Guamá, Pinar del Río, Cuba, W. Palmer and J. H. Riley, Mar. 8, 1900. 27635-7, Guanajay, Cuba, W. Palmer and J. H. Riley, June 11, 1900. 57862-3, Havana, Cuba, J. Hurter, Jan. 18, 1914, 59329-35, 59348, San Diego de los Baños, Cuba, W. Palmer, Jan. 31-Feb. 1, 1917. 26741 was exchanged to MZUM in 1944; 54399 to R, 1953.

***Eleutherodactylus poolei* Cochran**

Proc. Biol. Soc. Washington, vol. 51, p. 93, 1938.

HOLOTYPE: 73999, Citadel of King Christophe, Haiti, A. J. Poole, Feb. 14, 1928.

PARATYPES (4): 74000-1 74004-5, same data. 74002 was exchanged to MCZ in 1944; 74003 to MZUM, 1943.

***Eleutherodactylus richmondi* Stejneger**

Report of the U.S. National Museum . . . for the year ending June 30, 1902, p. 593, figs. 25-29, 1904.

HOLOTYPE: 26884, Catalina plantation, Puerto Rico, 890 ft., L. Stejneger, Feb. 22, 1900.

PARATYPES (7): 26956-62, Camp El Yunque, Puerto Rico, 2,863 ft., L. Stejneger, Feb. 24-26, 1900.

***Eleutherodactylus ricordii rogersi* Goin**

Amer. Mus. Novit., No. 1708, p. 1, fig. 1, 1955.

PARATYPES (4): 66250-3, Watling Island (Bahama Islands), H. and P. Bartsch, Aug. 26, 1923.

***Eleutherodactylus saltator* Taylor**

Proc. Biol. Soc. Washington, vol. 54, p. 89, 1941.

PARATYPE: 134261 (formerly UIMNH 21440), Omilteme, Guerrero, Mexico, E. H. Taylor, Aug. 2-4, 1940.

***Eleutherodactylus schmidti limbensis* Lynn**

Herpetologica, vol. 14, p. 157, 1958.

HOLOTYPE: 140166, 3 miles southwest of Limbé, Haiti, W. G. Lynn, Apr. 15, 1953.

PARATYPES (36): 140167-202, same data.

***Eleutherodactylus schmidti rucillensis* Cochran**

Proc. New England Zool. Club, vol. 18, p. 3, 1939.

PARATYPE: 107565, Loma Rucilla, Dominican Republic, 4,000-7,000 ft., P. J. Darlington, June 1938.

***Eleutherodactylus spatulatus* Smith**

Proc. Biol. Soc. Washington, vol. 52, p. 187, pl. 2, figs. 4-5, 1939.

HOLOTYPE: 116026, ♀, Cuautlapan, Veracruz, Mexico, H. M. Smith, Jan. 5, 1939.

PARATYPES (6): 116027-32, Cuautlapan, Veracruz, Mexico, H. M. Smith, Jan. 5, 1939 (116027-8), Jan. 16, 1939 (116029-32).

***Eleutherodactylus talamancae* Dunn**

Occ. Pap. Boston Soc. Nat. Hist., vol. 5, p. 385, 1931.

PARATYPE: 73222 (formerly MCZ 9784), Suretka, Costa Rica, E. R. Dunn, 1922.

***Eleutherodactylus thomasi trinidadensis* Schwartz**

Amer. Mus. Novit., No. 1926, p. 11, fig. 2, 1959.

PARATYPES (60): 137985-90, 10 miles west of Trinidad, Las Villas Province, Cuba, J. D. Hardy, Sept. 5-10, 1956.

***Eleutherodactylus turquinensis* Barbour and Shreve**

Bull. Mus. Comp. Zool., vol. 80, p. 380, 1937.

PARATYPE: 103377, Pico Turquino, near Cueva del Aura, Oriente, Cuba, 1,500-4,000 ft., P. J. Darlington, June 1936.

***Eleutherodactylus unicolor* Stejneger**

Report of the U.S. National Museum . . . for the year ending June 30, 1902, p. 597, figs. 35-39, 1904.

HOLOTYPE: 26963, camp at El Yunque Mountain, Puerto Rico, 2,978 ft., L. Stejneger, Feb. 26, 1900.

***Eleutherodactylus vocalis* Taylor**

Univ. Kansas Sci. Bull., vol. 26 (1939), p. 401, pl. 44, text-fig. 8, 1940.

PARATYPE: 137199, Hacienda El Sabino, Uruapan, Michoacán, Mexico, E. H. Taylor and H. M. Smith, July 25, 1936.

***Eleutherodactylus vulcani* Shannon and Werler**

Trans. Kansas Acad. Sci., vol. 58, p. 379, pl. 3, fig. 2, text-fig. 5, 1955.

PARATYPE: 137200, Volcán San Martín, lower slope, Veracruz, Mexico, 3,000-4,000 ft., J. E. Werler, Feb. 4-19, 1953.

Eleutherodactylus wetmorei Cochran

Proc. Biol. Soc. Washington, vol. 45, p. 191, 1932.

HOLOTYPE: 72617, Fonds-des-Nègres, Haiti, B. H. Swales and A. Wetmore, Apr. 5, 1927.

PARATYPES (11): 72618-19, same data. 60627-35, Moron, Haiti, W. L. Abbott, Dec. 20 and Dec. 24, 1917. 72620 was exchanged to MZUM in 1943; 60650-1 were exchanged to MCZ, 1957.

Elosia vomerina Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 423, 1853.

=*Crossodactylus gaudichaudii* Duméril and Bibron

COTYPES (2): 15481-2, Rio de Janeiro, Brazil, U.S. Exploring Expedition, 1837-38.

Engystoma areolata Strecker

Proc. Biol. Soc. Washington, vol. 22, p. 118, 1909.

=*Microhyla olivacea* Hallowell

HOLOTYPE: 38999, Guadalupe River bottom, Victoria County, Texas, J. D. Mitchell, January 1909.

PROBABLE PARATYPES (7): 42317-23, 8 miles below Victoria, Victoria County, Texas, J. D. Mitchell, January 1909. 42316 was exchanged to MCZ in 1928.

Engystoma pictiventre Cope

Proc. Amer. Philos. Soc., vol. 23, p. 272, 1886.

=*Microhyla pictiventris* (Cope)

HOLOTYPE: 14196, Nicaragua, J. F. Bransford, Aug. 7, 1885.

Engystoma texense Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 11 (1859), p. 169, 1860.

=*Microhyla olivacea* (Hallowell)

COTYPES (2): 2644 (2), Rio Seco, Medina County, Texas, J. Pope.

Engystoma ustum Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 18, p. 131, 1866.

=*Microhyla usta usta* (Cope)

HOLOTYPE: 24965, Guadalajara, Jalisco, Mexico, J. J. Major.

Engystoma variolosum Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 18, p. 131, 1866.

=*Hypopachus variolosus* (Cope)

HOLOTYPE: 6486, Arriba, Costa Rica, C. N. Riotte.

Eupemphix bolbodaetyla Lutz

Compt. Rend. Soc. Biol. Paris, vol. 93, p. 138, 1925.

=*Basanitia bolbodaetyla* (Lutz)

COTYPES (2) : 96542-3, Angra dos Reis, Rio de Janeiro, Brazil, A. Pugas, January 1925.

Eupemphix maculiventris Lutz

Compt. Rend. Soc. Biol. Paris, vol. 93, p. 138, 1925.

=*Physalaemus signiferus* (Girard)

COTYPES (6) : 96837-9, Alto da Serra, São Paulo, Brazil, A. Lutz, Feb. 12, 1923. 96840, 96842-3, Alto da Serra, São, Paulo, Brazil, A. Lutz. 96841 was exchanged to LC in 1944; 96844 to MZUM, 1944.

Eupemphix olfersioides Lutz

Compt. Rend. Soc. Biol. Paris, vol. 93, p. 138, 1925.

=*Phyllobates brunneus* (Cope)

COTYPES (3) : 96412, Niterói, Rio de Janeiro, Brazil, A. Lutz, 1923. 96539-40, Angra dos Reis, Rio de Janeiro, Brazil, A. Pugas, April 1924.

Eupemphix ruthveni Netting

Ann. Carnegie Mus., vol. 19, p. 167, pl. 7, fig. 1, 1930.

PARATYPE: 118717 (formerly MZUM 45579), Fundación, Colombia, A. G. Ruthven, Aug. 13, 1913.

Gastrotheca coronata Stejneger

Proc. U.S. Nat. Mus., vol. 41, p. 287, 1911.

=*Anotheca coronata* (Stejneger)

HOLOTYPE: 48279, ♂, Palomo, Valle de Orosi, Cartago, Costa Rica, 1,200 m., C. Picado, Apr. 19, 1911.

Gastrotheca monticola Barbour and Noble

Bull. Mus. Comp. Zool., vol. 63, p. 426, pl. 2, figs. 1-2, 1920.

PARATYPE: 118241 (formerly MCZ 5331), Quercotilla, Peru, G. K. Noble, October 1916.

Halophila heros Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 423, 1853.

=*Cornufer heros* (Girard)

HOLOTYPE: 14539, Sebukeya (Levuka in Wilkes' narrative), Fiji Islands, U.S. Exploring Expedition, 1840.

Halophila vitiensis Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 423, 1853.

=*Cornufer vitiensis* (Girard)

COTYPES (2) : 15475-6, Sebukeya (Levuka in Wilkes' narrative), Fiji Islands, U.S. Exploring Expedition, 1840.

Helocaetes clarkii Baird

Proc. Acad. Nat. Sci. Philadelphia, vol. 7, p. 60, 1854.

=*Pseudacris clarkii* (Baird)

COTYPE: 3313, Galveston, Galveston County, Texas, M. Dean.

Helocaetes feriarum Baird

Proc. Acad. Nat. Sci. Philadelphia, vol. 7, p. 60, 1854.

=*Pseudacris nigrita feriarum* (Baird)

COTYPES (12): 3592 (12), Carlisle, Cumberland County, Pennsylvania, S. F. Baird. One cotype was exchanged to MZUM.

Hyla acuminata Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 354, 1862.

COTYPES (2): 5843 and 102700, Paraguay, T. J. Page.

Hyla affinis Baird

Proc. Acad. Nat. Sci. Philadelphia, vol. 7, p. 61, 1854. (Not of Spix, 1824.)

=*Hyla arenicolor* Cope

COTYPES (5): 11410 (5, formerly 3261), northern Sonora (Mexico), J. H. Clark. These specimens were renamed *Hyla arenicolor* by Cope, Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 6, p. 84, 1866, to replace *Hyla affinis* Baird, preoccupied by Spix, 1824.

Hyla albofrenata Lutz

Compt. Rend. Soc. Biol. Paris, vol. 90, p. 241, 1924.

COTYPES (22): 96228-34, Tijuca, Rio de Janeiro, Brazil, A. Lutz, November 1920. 96235-6, Tijuca, Rio de Janeiro, Brazil, A. Lutz, December 1922. 96320-2, Tijuca, Rio de Janeiro, Brazil, A. Lutz, December 1923. 96319, Sumaré, Rio de Janeiro, Brazil, A. Lutz, October 1923. 96339-45, Corcovado, Rio de Janeiro, Brazil, A. Lutz, Nov. 13, 1923. 96347, Sumaré and Corcovado, Rio de Janeiro, Brazil, A. Lutz, November 1923. 96478, Angra dos Reis, Rio de Janeiro, Brazil, A. Pugas, Apr. 26, 1924. 97348 was exchanged to MCZ in 1944.

Hyla albosignata Lutz and Lutz

Ann. Acad. Brasileira Sci., vol. 10, p. 185, figs. 6-7a, 1938.

PARATYPES (3): 96644, Bonito, Serra da Bocaina, boundary of São Paulo and Rio de Janeiro, Brazil, A. Lutz, Dec. 15-31, 1931. 96781-2, Serra da Cubatão, São Paulo, Brazil, A. Lutz, Jan. 25, 1924.

Hyla anceps Lutz

Compt. Rend. Soc. Biol. Paris, vol. 101, p. 943, 1929.

COTYPES (2): 96441-2, Estrêlla, Rio de Janeiro, Brazil, A. Lutz, Mar. 1, 1929.

Hyla andersonii Baird

Proc. Acad. Nat. Sci. Philadelphia, vol. 7, p. 60, 1854.

HOLOTYPE: 3600, Anderson, Anderson County, South Carolina, C. Paine.

Hyla angustilineata Taylor

Univ. Kansas Sci. Bull., vol. 35, p. 850, fig. 54, 1952.

HOLOTYPE: 75060, La Palma, Costa Rica, M. Valerio, Apr. 4, 1928.

PARATYPES (2): 75061, 75065, same data.

Hyla arboricola Taylor

Univ. Kansas Sci. Bull., vol. 27, p. 118, pl. 5, fig. 1, 1941.

PARATYPE: 134262 (formerly UIMNH 27021), Omilteme, Guerrero, Mexico, E. H. Taylor, Aug. 4-6, 1940.

Hyla arenicolor Cope

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 6, p. 84, 1866.

HOLOTYPE: 11410 (originally 3261), northern Sonora, Mexico, J. H. Clark.

PARATYPES (4): 11410b-e, same data. These specimens were originally named as cotypes of *Hyla affinis* by Baird in Proc. Acad. Nat. Sci. Philadelphia, vol. 7, p. 61, 1854.

Hyla aurantiaca surda Cochran

Herpetologica, vol. 8, p. 112, figs. E-F, 1953.

PARATYPE: 132912 (formerly MZUM 104116A), Curitiba, Paraná, Brazil, J. R. Bailey, 1941.

Hyla avivoca Viosca

Proc. Biol. Soc. Washington, vol. 41, p. 89, 1928.

HOLOTYPE: 75017, ♂, Mandeville, Saint Tammany Parish, Louisiana, P. Viosca, Jr., June 12, 1926. Ten specimens, 75018-27, from the same locality and received here with the type, may be considered as paratypes, although they were not so listed.

Hyla bipunctata branneri Cochran

Journ. Washington Acad. Sci., vol. 38, p. 316, 1948.

HOLOTYPE: 48861, Bonito, Pernambuco, Brazil, J. C. Branner.

PARATYPES (5): 48862-4, 125719-20, ♂, ♂ (formerly MCZ 2827), same data.

Hyla bistincta Cope

Proc. Amer. Philos. Soc., vol. 17 (1878), p. 87, 1877.

=*Hyla bistincta bistincta* Cope

HOLOTYPE: 32261, probably Veracruz, Mexico, F. Sumichrast.

Hyla bistincta labeculata Shannon

Proc. U.S. Nat. Mus., vol. 101, p. 470, figs. 92, 93d, 1951.

HOLOTYPE: 123689, San Lucas Camotlán, Oaxaca, Mexico, W. S. Miller, Apr. 29, 1946.

PARATYPE: 123690, tadpole, same data.

Hyla cadaverina Cope

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 6, p. 84, 1866. (New name for *Hyla nebulosa* Hallowell (not of Spix).)

= *Hyla regilla* Baird and Girard

COTYPE?: 3230, Tejon Pass (Los Angeles County), California, A. L. Heermann. This specimen is also the type of *Hyla nebulosa* Hallowell, Proc. Acad. Nat. Sci., Philadelphia, vol. 7, p. 96, 1854.

Hyla cárdenasi Taylor

Univ. Kansas Sci. Bull., vol. 25 (1938), p. 430, pl. 47, fig. 2, 1939.

HOLOTYPE: 84403, Puebla, Puebla, Mexico, H. Ruano, September 1919.

Hyla ceratophrys Stejneger

Proc. U.S. Nat. Mus., vol. 41, p. 286, 1911.

HOLOTYPE: 47705, Upper Pequiní River, Panama, A. H. Jennings, Mar. 29, 1909.

Hyla colymba Dunn

Occ. Pap. Boston Soc. Nat. Hist., vol. 5, p. 400, 1931.

PARATYPES (3): 118244, 1 juv. and 2 tadpoles, La Loma, Panama, E. R. Dunn, 1923.

Hyla crosopedospila Lutz

Compt. Rend. Soc. Biol. Paris, vol. 93, p. 211, 1925.

COTYPES (8): 96926-32, 96934, Campo Bello, Itatiaia, Rio de Janeiro, Brazil, A. Lutz, Dec. 18, 1923-Jan. 20, 1924. 96933 was exchanged to LC in 1944.

Hyla culex Dunn and Emlen

Proc. Acad. Nat. Sci. Philadelphia, vol. 84, p. 24, 1932.

= *Hyla staufferi* Cope

PARATYPE: 20262 (incorrectly given as 20267 in original description), Patuca, Honduras, H. W. Perry, May 14, 1891.

Hyla curta Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 18 (1866), p. 313, 1867.

= *Hyla regilla* Baird and Girard

COTYPES (17): 5293 (17), in a spring at Soria, 15 miles north of Cape San Lucas, Baja California, Mexico, J. Xantus, Aug. 10, 1859. One cotype was exchanged to MCZ in 1914.

Hyla cuspidata Lutz

Compt. Rend. Soc. Biol. Paris, vol. 93, p. 211, 1925.

COTYPES (5): 96139, Manguinhos, Rio de Janeiro, Brazil, A. Lutz, April 1925. 96147-9, Manguinhos, Rio de Janeiro, Brazil, A. Lutz, June 23, 1922. 96370, Rio de Janeiro, Brazil, A. Lutz, January 1923.

Hyla decipiens Lutz

Compt. Rend. Soc. Biol. Paris, vol. 93, p. 212, 1925.

COTYPE: 96194, Bom Sucesso, Rio de Janeiro, Brazil, A. Lutz, Apr. 3, 1925. A young specimen, 96150, from Manguinhos, Rio de Janeiro, Brazil, A. Lutz, is probably also a cotype.

Hyla dendrocarta Taylor

Proc. U.S. Nat. Mus., vol. 89, p. 45, pls. 2-3, 1940.

HOLOTYPE: 108679, Cuautlapan, Veracruz, Mexico, H. M. Smith, Jan 1, 1939.

PARATYPES (10): 108676-8, 108680-6, same data.

Hyla elaeochroa Cope

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8 (1875), p. 105, pl. 26, fig. 3, 1876.

LECTOTYPE: 30689, east foot of mountains near Sipurio, Costa Rica, W. M. Gabb.

PARATYPES (2): 30688,, 30690, same data.

Hyla elongata Lutz

Compt. Rend. Soc. Biol. Paris, vol. 93, p. 139, 1925.

COTYPES (8): 96861, 96863, Campo de Aviação, São Paulo, Brazil, B. Lutz, January 1924. 96957-62, Belo Horizonte, Minas Gerais, Brazil, A. Lutz and J. Venancio, November-December 1924. 96862 was exchanged to LC in 1944.

Hyla euthysanota Kellogg

Proc. Biol. Soc. Washington, vol. 41, p. 123, 1928.

HOLOTYPE: 73296, Los Esesmiles, Chalatenango, El Salvador, 6,400 ft., R. A. Stirton, Mar. 6, 1927.

Hyla evelynae Schmidt

Field Mus. Nat. Hist., Zool. Ser., vol. 29, p. 156, fig. 21, 1944.

PARATYPES (2): 101463-4 (formerly FMNH 9560, 9565), 15 miles north-east of San Carlos, Uruguay, K. P. Schmidt, Oct. 21, 1926.

Hyla evittata Miller

Proc. Biol. Soc. Washington, vol. 13, p. 76, 1899.

=*Hyla cinerea evittata* Miller

HOLOTYPE: 26291, ♂, Four Mile Run, Alexandria County (now Arlington County), Virginia, G. S. Miller, Jr., and E. A. Preble, July 15, 1899.

Hyla eximia Baird

Proc. Acad. Nat. Sci. Philadelphia, vol. 7, p. 61, 1854.

=*Hyla eximia eximia* Baird

PROBABLE COTYPES (2): 3248 (2), City of Mexico, Distrito Federal, Mexico, W. Rich, 1853.

Hyla flavoguttata Lutz and Lutz

Ann. Acad. Brasileira Sci., vol. 11, p. 75, 1939.

=*Hyla strigilata flavoguttata* Lutz and Lutz

PARATYPE: 121636, Petrópolis, Rio de Janeiro, Brazil, B. Lutz, July 1, 1938.

Hyla fuscomarginata Lutz

Compt. Rend. Soc. Biol. Paris, vol. 93, p. 138, 1925.

COTYPE: 96964, Belo Horizonte, Minas Gerais, Brazil, A. Lutz, 1924.

Hyla fuscovaria Lutz

Compt. Rend. Soc. Biol. Paris, vol. 93, p. 212, 1925.

COTYPES (9): 96988, 96992-4, Agua Limpa, near Ouro Preto, Minas Gerais, Brazil, A. Lutz, 1922 (96988), Oct. 22, 1921 (96992-4). 97001-5, Lassance, Minas Gerais, Brazil, A. Lutz, July 14, 1921 (97001), Mar. 6, 1922 (97002-5).

Hyla gabbii Cope

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8 (1875), p. 103, 1876.

COTYPES (2): 30658-9, Costa Rica (near Sipurio), W. M. Gabb.

Hyla goughi baileyi Cochran

Herpetologica, vol. 8, p. 114, figs. g-h, 1953.

PARATYPES (4): 132913-6 (formerly MZUM 104119c, 104127a and c, and 104131d), 8 km. north of Juquiá, São Paulo, Brazil, J. R. Bailey, Feb. 25-28, 1941.

Hyla gracilipes Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 17, p. 194, 1865.

=*Hyla eximia eximia* Baird

LECTOTYPE: 15321, Mirador, Veracruz, Mexico, C. Sartorius.

PARATYPES (2): 15319-20, same data.

Hyla gratiosa LeConte

Proc. Acad. Nat. Sci. Philadelphia, vol. 8, p. 146, pl. 6, 1856.

COTYPES (6): 5901 (6), Riceboro, Liberty County, Georgia, J. L. LeConte.

Hyla hallowelli Thompson

Herpetological Notices, No. 2, p. 2, June 28, 1912 (privately published, San Francisco, California).

PARATYPE: 64206, Kikaiga, Loo Choo (Ryukyu) Islands, V. Kuhne, Apr. 30, 1910.

Hyla loquax Gaige and Stuart

Occ. Pap. Mus. Zool. Univ. Michigan, No. 281, p. 1, 1934.

PARATYPE: 94100, 2 miles south of Libertad, Petén, Guatemala, L. C. Stuart, Apr. 28, 1933.

Hyla marianae Dunn

Proc. Boston Soc. Nat. Hist., vol. 38, p. 129, pl. 1, fig. 3, pl. 2, fig. 3, 1926.

PARATYPES (2): 35860, 36668, Mount Diablo, Holly Mount, St. Catherine Parish, Jamaica, 2,700 ft., W. Harris, Aug. 18, 1905.

Hyla megapodia Miranda-Ribeiro

O Campo, Rio de Janeiro, p. 67, May 1937.

=*Hyla raniceps* (Cope)

PARATYPE: 102687, Porto Esperidião, Rio Jaurú, Mato Grosso, Brazil, A. de Miranda-Ribeiro, September 1908. A figure without a description of this specimen is given in Miranda-Ribeiro, Arch. Mus. Nac. Rio de Janeiro, vol. 27, pl. 5, fig. 5, 1926.

Hyla melanomma Taylor

Univ. Kansas Sci. Bull., vol. 26 (1939), p. 508, pl. 57, figs. 1-1b, 1940.

PARATYPE: 134263 (formerly UIMNH 27011), Chilapa Road, 7 miles east of Chilpancingo, Guerrero, Mexico, H. M. Smith and E. H. Taylor, Aug. 20, 1939.

Hyla microcephala martini Smith

Herpetologica, vol. 7, p. 187, 1951.

PARATYPES (2): 134264-5, Encarnación, Campeche, Mexico, H. M. Smith, Oct. 11-12, 1936.

Hyla microcephala sartori Smith

Herpetologica, vol. 7, p. 186, 1951.

PARATYPE: 134266 (formerly UIMNH 20941), El Limoncita near La Venta, Guerrero, Mexico, E. H. Taylor, July 1, 1938.

Hyla microeximia Maslin

Herpetologica, vol. 13, p. 81, fig. 1, 1957.

HOLOTYPE: 139246 (formerly UCM 8379), ♂, 3 miles northwest of Jacotepec, near Lake Chapala, Jalisco, Mexico, T. P. Maslin, June 20, 1956.

PARATYPE: 139247 (formerly UCM 8399), ♂, same data.

Hyla milleri Shannon

Proc. U.S. Nat. Mus., vol. 101, p. 473, figs. 33a-c, 1951.

HOLOTYPE: 123700, ♂, San Lucas Camotlán, Oaxaca, Mexico, W. S. Miller, May 7, 1946.

PARATYPE: 123699, ♂, same data.

Hyla multilineata Lutz and Lutz

Ann. Acad. Brasileira Sci., vol. 11, p. 72, 1939.

=*Hyla bischoffi multilineata* Lutz and Lutz

PARATYPES (11): 96784-94, Alto da Serra, Serra de Cubatão, São Paulo, Brazil, A. Lutz, Feb. 27, 1922 (96784), February 1923 (96785-94).

Hyla muricolor Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 359, 1862.

=*Smilisca baudinii baudinii* (Duméril and Bibron)

HOLOTYPE?: 25097 (probably 4818), Mirador, Veracruz, Mexico, C. Sartorius.

PARATYPE?: 25098, same data.

Hyla nasica Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 354, 1862.

COTYPES (2): 5835, 32371, Paraguay, T. J. Page.

Hyla nebulosa Hallowell

Proc. Acad. Nat. Sci. Philadelphia, vol. 7, p. 96, 1854. (Not of Spix, 1824.)

=*Hyla regilla* Baird and Girard

COTYPE?: 3230, Tejon Pass (Los Angeles County), California, A. L. Heermann. This specimen is also the type of *Hyla cadaverina* Cope, Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 6, p. 84, 1866.

Hyla nigripes Cope

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8 (1875), p. 104, pl. 23, fig. 7, 1876.

COTYPES (2): 30685-6, Pico Blanco, Costa Rica, 5,000-7,000 ft., W. M. Gabb.

Hyla (Sphoehyla) orophila Lutz and Lutz

Ann. Acad. Brasileira Sci., vol. 10, p. 178, 1938.

=*Hyla aurantiaca orophila* Lutz and Lutz

PARATYPES (5): 96692-6, Bonito, Serra da Bocaina, boundary of São Paulo and Rio de Janeiro, Brazil, A. Lutz, Jan. 16-30, 1925.

Hyla pachyderma Taylor

Univ. Kansas Sci. Bull., vol. 28, p. 308, pl. 27, figs. 1-3, 1942.

HOLOTYPE: 115029, Pan de Olla, Veracruz, Mexico, H. M. Smith, Mar. 22, 1940.

PARATYPES (3): 115026-8, same data.

Hyla pallens Lutz

Compt. Rend. Soc. Biol. Paris, vol. 93, p. 212, 1925.

=*Hyla minuta* Peters

COTYPES (3): 96197-9, Bom Successo, Rio de Janeiro, A. Lutz, Sept. 2, 1923 (96197-8), Aug. 6, 1921 (96199).

Hyla parkeri Gaige

Occ. Pap. Mus. Zool. Univ. Michigan, No. 207, p. 1, 1929.

PARATYPES (4): 84359-60, 101440-1, Buenavista, Department of Santa Cruz, Bolivia, J. Steinbach.

Hyla phaeota Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 358, 1862.

=*Hyla phaeota phaeota* Cope

HOLOTYPE: 39899 (formerly 4347), Turbo (Colombia), A. Schott.

Hyla phaeota cyanosticta Smith

Herpetologica, vol. 8, p. 150, 1952.

HOLOTYPE: 111147, ♂, Piedras Negras, Petén, Guatemala, H. M. Smith, May 28, 1939.

PARATYPES (8): 35907, Semococh, above Panzós, Alta Verapaz, Guatemala, G. P. Goll, Feb. 19, 1905. 111139-41, 111143-6, Piedras Negras, Petén, Guatemala, H. M. Smith, May 28-29, 1939.

Hyla phlebodes Stejneger

Proc. U.S. Nat. Mus., vol. 30, p. 817, 1906.

HOLOTYPE: 29970, ♀, San Carlos, Costa Rica, C. Burgdorf and P. Schild.

Hyla pickeli Lutz and Lutz

Ann. Acad. Brasileira Sci., vol. 10, p. 189, 1938.

PARATYPES (8): 97067-8, 97071-2, Tapera, Pernambuco, Brazil, P. B. Pickel, January 1928. 97042-3, 97045-6, Natal, Rio Grande do Norte, Brazil, P. B. Pickel, July 1928.

Hyla (Sphoenohyla) planicola Lutz and Lutz

Ann. Acad. Brasileira Sci., vol. 10, p. 182, figs. 3-5, 1938.

=*Hyla aurantiaca planicola* Lutz and Lutz

PARATYPES (9): 97563-8, Km. 40, road between Rio de Janeiro and São Paulo, near Rio de Janeiro, Brazil, A. Lutz, D. M. Cochran, and J. Venancio,

Feb. 20, 1935. 97619-21, Recreio dos Bandeirantes, Rio de Janeiro, Brazil, B. Lutz, D. M. Cochran, and J. Venancio, Feb. 9-10, 1935. 97569 was exchanged to AMNH in 1948.

Hyla puma Cope

Proc. Amer. Philos. Soc., vol. 22, p. 183, 1885.

=*Hyla molitor* Schmidt

HOLOTYPE: 13735 (given incorrectly as 18735 in original description), Nicaragua, J. F. Moser, Dec. 4, 1883.

Hyla punctariola moesta Cope

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8 (1875), p. 106, 1876.

=*Hyla moesta* Cope

PROBABLE HOLOTYPE: 30660, Pico Blanco, Costa Rica, W. M. Gabb.

Hyla punctariola monticola Cope

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8 (1875), p. 106, 1876.

=*Hyla monticola* Cope

HOLOTYPE: 30661, Pico Blanco, Costa Rica, W. M. Gabb.

Hyla punctariola pictipes Cope

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8 (1875), p. 106, 1876.

=*Hyla pictipes* Cope

HOLOTYPE: 30652, Cordilleras, Pico Blanco, Costa Rica, 5,000-7,000 ft., W. M. Gabb.

Hyla punctata Hallowell

Proc. Acad. Nat. Sci. Philadelphia, vol. 7, p. 193, 1854. (Not of Schneider, 1799, or of Wied, 1824.)

=*Hylambates aubryi* Günther

HOLOTYPE: 32369, Gabun (French West Equatorial Africa), P. du Chaillu.

Hyla quinquevittata Cope

Proc. Amer. Philos. Soc., vol. 23, p. 273, 1886.

=*Hyla elaeochroa* Cope

HOLOTYPE: 14187, Nicaragua, J. F. Bransford, Aug. 7, 1885.

Hyla regilla Baird and Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 174, 1852.

COTYPES (2): 9182, Puget Sound (Washington), C. J. Pickering. 15409, Sacramento River (restricted to Sacramento County, California), U.S. Exploring Expedition, 1841. One cotype was exchanged to MCZ in 1879.

Hyla regilla laticeps Cope

U.S. Nat. Mus. Bull. 34, p. 359, 1889.

=*Hyla regilla* Baird and Girard

COTYPES (12): 5308 (12), Cape San Lucas, Baja California, Mexico, J. Xantus. One cotype was exchanged to MCZ in 1915.

***Hyla rickardsi* Taylor**

Univ. Kansas Sci. Bull., vol. 25 (1938), p. 335, pl. 41, figs. 1-8, 1939.

PARATYPE: 139722 (formerly UIMNH 39423), Potrero Viejo, Veracruz, Mexico, Mr. and Mrs. D. M. Forbes and E. H. Taylor, Aug. 29, 1936.

***Hyla robertmertensi* Taylor**

Proc. Biol. Soc. Washington, vol. 50, p. 43, pl. 2, figs. 3-7, 1937.

PARATYPE: 134267 (formerly UIMNH 26991), near Asunción, Chiapas, Mexico, E. H. Taylor and H. M. Smith, Sept. 1, 1935.

***Hyla robertsororum* Taylor**

Univ. Kansas Sci. Bull., vol. 26 (1939), p. 393, figs. 5-6, 1940.

PARATYPE: 134268 (formerly UIMNH 27037), El Chico National Park, Hidalgo, Mexico, H. and R. Roberts, Aug. 7, 1938.

***Hyla rozellae* Taylor**

Univ. Kansas Sci. Bull. vol. 28, p. 78, pl. 9, figs. 1-1c, 1942.

HOLOTYPE: 115039, Salto de Agua, Chiapas, Mexico, H. M. Smith, Apr. 18, 1940.

PARATYPES (25): 115030-8, Las Nubes on Mount Ovando, Chiapas, Mexico, H. M. Smith, Apr. 15, 1940. 115040-51, Salto de Agua, Chiapas, Mexico, H. M. Smith, Apr. 18, 1940. 115052-5, Finca Juárez, 28 km. northeast of Escuintla, Chiapas, Mexico, H. M. Smith, May 8, 1940.

***Hyla sanborni* Schmidt**

Field Mus. Nat. Hist., Zool. Ser., vol. 29, p. 155, fig. 20, 1944.

PARATYPES (3): 101457-8 (formerly FMNH 9582-3), San Carlos, Uruguay, K. P. Schmidt, Oct. 21, 1926. 101459 (formerly FMNH 10345), 5 miles north of San Vicente de Castillos, Department of Rocha, Uruguay, K. P. Schmidt, Nov. 9, 1926.

***Hyla similis* Cochran**

Journ. Washington Acad. Sci., vol. 42, p. 50, 1952.

HOLOTYPE: 97317, ♂, Manguinhos, Rio de Janeiro, Brazil, J. Venancio, Feb. 25, 1935.

PARATYPES (51): 81119-21, Rio de Janeiro, Brazil, M. M. Metcalf, 1930. 96144-6, Manguinhos, Rio de Janeiro, Brazil, A. Lutz, January 1922. 96203, Bom Sucesso, Rio de Janeiro, Brazil, A. Lutz, Oct. 25, 1928. 96213, Amorim, Rio de Janeiro, Brazil, A. Lutz, January 1926. 97312-6, 97318-52, 97374-6, Manguinhos, Rio de Janeiro, Brazil, J. Venancio, Feb. 15, 1935.

Hyla smaragdina Taylor

Copeia, No. 1, p. 18, fig. 1, 1940.

PARATYPE: 134269 (formerly UIMNH 27001), 4 miles east of Cojumatlán, Michoacán, Mexico, H. M. Smith and E. H. Taylor, 1939.

Hyla squalirostris Lutz

Compt. Rend. Soc. Biol. Paris, vol. 93, p. 212, 1925.

COTYPES (2): 96719-20, Bonito, Serra da Bocaina, boundary of São Paulo and Rio de Janeiro, Brazil, A. Lutz, Jan. 16-30, 1925.

Hyla staufferi Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 17, p. 195, 1865.

HOLOTYPE: 15317, Orizaba, Veracruz, Mexico, F. Sumichrast.

Hyla uranochroa Cope

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8 (1875), p. 103, pl. 27, fig. 4, 1876.

HOLOTYPE: 30651, Pico Blanco (near Sipurio), Costa Rica, W. M. Gabb.

Hyla vanvlietii Baird

Proc. Acad. Nat. Sci. Philadelphia, vol. 7, p. 61, 1854.

=*Smilisca baudinii baudinii* (Duméril and Bibron)

COTYPE?: 3256, juv., Brownsville, Cameron County, Texas, S. Van Vliet.

Hyla variabilis Boulenger

Ann. Mag. Nat. Hist., ser. 6, vol. 17, p. 20, 1896.

COTYPE: 71115 (formerly MCZ 194), Cali, Cauca Valley, Colombia, 3,200 ft., W. F. H. Rosenberg.

Hyla versicolor phaeocrypta Cope

U.S. Nat. Mus. Bull. 34, p. 375, 1889.

=*Hyla versicolor versicolor* LeConte

HOLOTYPE: 12074, Mount Carmel, Wabash County, Illinois, L. M. Turner.

Hyla versicolor sandersi Smith and Brown

Proc. Biol. Soc. Washington, vol. 60, p. 48, 1947.

HOLOTYPE: 123978, 8 miles southwest of Somerset, in Atascosa County, Texas, A. J. Kirn, Apr. 27, 1946.

Hyla wrightorum Taylor

Univ. Kansas Sci. Bull., vol. 25 (1938), p. 436, pl. 47, fig. 1, 1939.

=*Hyla eximia wrightorum* Taylor

PARATYPES (8): 9338 (2), Santa Fe, Santa Fe County, New Mexico, H. W. Henshaw, June 18, 1874. 26605-9, Meadow Valley, Chihuahua, Mexico, H. H. and C. S. Brimley, July 18, 1898 (26605-8), June 29, 1898 (26609). 118643, David Lee Lake, southwest of Luna, Catron County, New Mexico, J. J. Cantrell, Aug. 3, 1935.

Hyla zeteki Gaige

Occ. Pap. Mus. Zool. Univ. Michigan, No. 207, p. 4, 1929.

PARATYPE: 118672 (formerly MZUM 63881), Boquete, Chiriquí Province, Panama, H. T. Gaige, Mar. 3, 1927.

Hylambates cochranæ Loveridge

Proc. U.S. Nat. Mus., vol. 91, p. 125, 1941.

COTYPE: 109569, ♂, Bendaja, Liberia, W. M. Mann, May 14–27, 1940. 109570 was exchanged to MCZ in 1940.

Hyalplesia nigriventris Lutz

Compt. Rend. Soc. Biol. Paris, vol. 93, p. 139, 1925.

=*Eleutherodactylus guentheri* (Steindachner) ?

COTYPE: 96846, Alto da Serra, São Paulo, Brazil, A. Lutz.

Hylella platycephala Cope

Proc. Amer. Philos. Soc., vol. 18, p. 267, 1879.

=*Hylella sumichrasti* (Brocchi)

COTYPES (5): 10037 (5), Tapana, near Santa Efigenia, Tehuantepec, Oaxaca, Mexico, 2,000–3,000 ft., F. Sumichrast.

Hylodes cerasinus Cope

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8 (1875), p. 112, pl. 26, fig. 4, 1876.

=*Eleutherodactylus cerasinus* (Cope)

HOLOTYPE: 32572, Pico Blanco, eastern slope, Costa Rica, W. M. Gabb.

Hylodes cuneatus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 152, 1862.

=*Eleutherodactylus cuneatus* (Cope)

COTYPES (2): 5202 (2), eastern Cuba, C. Wright.

Hylodes nasutus Lutz

Compt. Rend. Soc. Biol. Paris, vol. 93, p. 213, 1925.

=*Eleutherodactylus nasutus* (Lutz)

COTYPES (2): 96468–9, Hotel Lemburger, Nova Friburgo, Rio de Janeiro, Brazil, A. Lutz and J. Venancio, Feb. 22, 1923. 96470 was exchanged to LC in 1944.

Hylodes polyptychus Cope

Proc. Amer. Philos. Soc., vol. 23, p. 276, 1886.

=*Microbatrachylus polyptychus* (Cope)

COTYPES (2): 14199 (2), Nicaragua, J. F. Bransford, Aug. 7, 1885.

Hymenochirus boettgeri camerunensis Perret and Mertens

Bull. Inst. François Afr. Noire, ser. A, vol. 19, p. 552, fig. 1, 1957.

PARATYPE: 140246 (formerly SNG 52302), Foulassi, Sangmélima, South Cameroon (French Cameroons), J. L. Perret, Feb. 12, 1954.

Hypopachus alboventer Taylor

Univ. Kansas Sci. Bull., vol. 26 (1939), p. 522, pls. 60 and 63, figs. 3-3a, 1940.

PARATYPE: 137203, 8 miles east of Cuernavaca, Morelos, Mexico, E. H. Taylor, June 20, 1938.

Hypopachus caprimimus Taylor

Univ. Kansas Sci. Bull., vol. 26 (1939), p. 526, pl. 61, pl. 63, figs. 1-1a, 1940.

PARATYPE: 134270 (formerly UIMNH 27274), Kilometer 327, Guerrero, Mexico, E. H. Taylor, July 21, 1936.

Hypopachus championi Stuart

Proc. Biol. Soc. Washington, vol. 53, p. 19, 1940.

PARATYPES (2): 118667-8 (formerly MZUM 84082), 1 km. south of San Gerónimo, Baja Verapaz, Guatemala, L. C. Stuart, July 6, 1938.

Hypopachus cuneus Cope

Proc. U.S. Nat. Mus., vol. 11, p. 395, 1839.

=*Hypopachus cuneus cuneus* Cope

COTYPE: 15676, San Diego, Duval County, Texas, W. Taylor.

Hypopachus maculatus Taylor

Univ. Kansas Sci. Bull., vol. 26 (1939), p. 524, pl. 62, figs. E-F, pl. 63, figs. 2-2a, 1940.

PARATYPE: 137204, near Asunción, Chiapas, Mexico, E. H. Taylor and H. M. Smith, Sept. 1, 1935.

Hypopachus ovis Taylor

Univ. Kansas Sci. Bull., vol. 26 (1939), p. 520, pl. 62, fig. B, 1940.

PARATYPE: 134271 (formerly UIMNH 27298), Tepic, Nayarit, Mexico, E. H. Taylor, July 31, 1934.

Hypopachus pearsei Ruthven

Proc. Biol. Soc. Washington, vol. 27, p. 77, 1914.

=*Elachistocleis ovalis* (Schneider)

PARATYPE: 51215, (formerly MZUM 45574), Fundación, Colombia, A. S. Pearse.

Hypsiboas miliarius Cope

Proc. Amer. Philos. Soc., vol. 23, p. 272, 1836.

=*Hyla miliaria* (Cope)

HOLOTYPE: 14193, Nicaragua, J. F. Bransford, Aug. 7, 1885.

Hypsiboas raniceps Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 353, 1862.

=*Hyla raniceps* (Cope)

COTYPES (8): 5403 (4), 5408, 12160, 12172 (2), Paraguay, T. J. Page.

***Ixalus japonicus* Hallowell**

Proc. Acad. Nat. Sci. Philadelphia, vol. 12 (1860), p. 501, 1861.

=*Rhacophorus japonicus* (Hallowell)

COTYPES (5): 7313 (5), Ousima (Amami, Ryukyu Islands), W. Stimpson, May 1855. One cotype was exchanged to MCZ in 1909.

***Kalophrynus stellatus* Stejneger**

Proc. U.S. Nat. Mus., vol. 33, p. 575, 1908.

=*Kalophrynus pleurostigma pleurostigma* Tschudi

HOLOTYPE: 37375, Basilan, Philippine Islands, E. A. Mearns, Feb. 13-22, 1906.

***Kaloula conjuncta stickeli* Inger**

Chicago Nat. Hist. Mus., Fieldiana: Zool., vol. 33, p. 443, 1954.

PARATYPE: 134422 (formerly CNHM 60788), Carigara, Leyte, Philippine Islands, W. H. Stickel, Nov. 22, 1944.

***Kaloula macroptica* Liu**

Journ. West China Border Res. Soc., ser. B, vol. 15, p. 31, pl. 1, figs. 6-7, pl. 2, figs. 8-9, 1945.

PARATYPES (2): 124574-5, ♂, ♂, Sichang, Sikang, China, C. C. Liu, 1942.

***Kaloula rugifera* Stejneger**

Occ. Pap. Boston Soc. Nat. Hist., vol. 5, p. 119, 1924.

HOLOTYPE: 65520, ♀, Kiating, Szechwan, China, D. C. Graham.

***Leptodactylus caliginosus* Girard**

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 422, 1853.

=*Leptodactylus ocellatus* (Linnaeus)

COTYPES (2): 7389 (2), Rio de Janeiro, Brazil, U.S. Exploring Expedition, 1839.

***Leptodactylus curtus* Barbour and Noble**

Bull. Mus. Comp. Zool. vol. 63, p. 405, 1920.

PARATYPE: 75990, Bellavista, Peru, G. K. Noble, 1916.

***Leptodactylus darlingtoni* Cochran**

Proc. Boston. Soc. Nat. Hist., vol. 40, p. 372, 1935.

=*Eleutherodactylus jugans* Cochran

PARATYPES (8): 95423-7, La Visite, La Selle Range, Haiti, P. J. Darlington, Sept. 16-23, 1935. 95428-30, Mount La Hotte, La Selle Range, Haiti, P. J. Darlington, October 1934. Paratypes of *Leptodactylus darlingtoni* Cochran (MCZ 19357-63) from Mount La Hotte were renamed *Leptodactylus ventrilineatus* by Shreve, Proc. New England Zool. Club, vol. 15, p. 98, 1936, but are now known as *Eleutherodactylus ventrilineatus* (Shreve).

Leptodactylus dominicensis Cochran

Journ. Washington Acad. Sci., vol. 13, p. 184, 1923.

HOLOTYPE: 65670, Las Cañitas, Dominican Republic, W. L. Abbott, Feb. 25, 1923.

PARATYPES (3): 65671, tadpole, Samaná and Laguna, Dominican Republic, W. L. Abbott, March 1923. 66675-6, 4 miles west of Jovéro, Dominican Republic, W. L. Abbott, Dec. 4, 1923.

Leptodactylus gaigeae Cochran

Proc. Biol. Soc. Washington, vol. 51, p. 41, 1938.

HOLOTYPE: 96759, ♂, Bonito, Serra da Bocaina, boundaries of São Paulo and Rio de Janeiro, Brazil, A. Lutz, Dec. 28, 1931.

PARATYPE: 96760, same data.

Leptodactylus insularum Barbour

Bull. Mus. Comp. Zool., vol. 46, p. 228, 1906.

=*Leptodactylus bolivianus* Boulenger

PARATYPES (3): 35603-4, San Miguel Island (Pearl Islands), Bay of Panama, Panama, W. W. Brown, Jr., 1900-1901. 52404 (formerly MCZ 2444), Saboga Island, Pearl Islands, Bay of Panama, Panama, W. W. Brown, Jr., May 1904.

Leptodactylus natalensis Lutz

Mem. Inst. Oswaldo Cruz, vol. 23, pp. 7, 26, pl. 1, figs. 7, 7a, pl. 3, figs. 1-2, 1930.

=*Leptodactylus podicipinus* (Cope)

COTYPE: 81130, Natal, Rio Grande do Norte, Brazil, A. Lutz, July 1925.

Leptodactylus occidentalis Taylor

Trans. Kansas Acad. Sci., vol. 39 (1936), p. 349, pl. 1, figs. 1-2, 7, 1937.

PARATYPE: 134272, (formerly UIMNH 26805), near Tepic, Nayarit, Mexico, E. H. Taylor, July 31, 1934.

Leptodactylus pentadactylus mattogrossensis Schmidt and Inger

Chicago Nat. Hist. Mus., Fieldiana: Zool., vol. 31, p. 444, 1951.

PARATYPE: 132738 (formerly FMNH 9192), 1 day's run below Descalvados, Mato Grosso, Brazil, K. P. Schmidt, 1926.

Leptodactylus serialis Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 421, 1853.

=*Leptodactylus ocellatus* (Linnaeus)

COTYPES (2): 7357, Rio de Janeiro, Brazil, U.S. Exploring Expedition, 1839. 15477, Rio de Janeiro, Brazil, J. Drayton, 1839.

Leptodactylus trivittatus Lutz

Mem. Inst. Oswaldo Cruz, vol. 19, pp. 151, 170, pl. 32, figs. 14-15, 1926.

=*Leptodactylus marmoratus* (Steindachner)

COTYPES (12): 96815-9, Alto da Serra, São Paulo, Brazil, A. Lutz, Feb. 27, 1922 (96815-6), January 1924 (96817-9). 96936, 96938-9, 96942, Montserrat, Campo Bello, Rio de Janeiro, Brazil, A. Lutz, February 1925. 96940-1, Campo Bello, Rio de Janeiro, Brazil, A. Lutz, November 1924. 96943, Campo Bello, Rio de Janeiro, Brazil, A. Lutz, 1925.

Leptopelis bequaerti Loveridge

Proc. U.S. Nat. Mus., vol. 91, p. 128, 1941.

PARATYPE: 109051, Si Mountain, Gibi, Liberia, W. M. Mann, 1940.

Limnomedusa schmidti Cochran and Goin

Copeia, No. 3, p. 208, fig. la-c, 1959.

HOLOTYPE: 140245, Leticia, Amazonas, Colombia, E. R. Allen, March 1956.

Lithodytes bransfordii Cope

Proc. Amer. Philos. Soc., vol. 23, p. 274, 1886.

=*Microbatrachylus bransfordii* (Cope)

COTYPES (11): 14200 (11), Nicaragua, J. F. Bransford, Aug. 7, 1885.

Lithodytes diastema Cope

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8 (1875), p. 155, 1876.

=*Eleutherodactylus diastema* (Cope)

COTYPES (2): 25170-1, Camp Mary Caretta, Panama, J. F. Bransford. E. R. Dunn considered 25171 an example of *Eleutherodactylus ockendeni* Boulenger.

Lithodytes gulosus Cope

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8 (1875), p. 112, 1876.

=*Eleutherodactylus rugosus* (Peters)?

HOLOTYPE: 32590, Pico Blanco, Costa Rica, 6,000 ft., W. M. Gabb.

Lithodytes lanciformis Cope

Proc. Amer. Philos. Soc., vol. 17 (1878), p. 88, 1877.

=*Eleutherodactylus gollmeri* (Peters)

COTYPES (2): 32324-5, west coast of Central America.

Lithodytes latrans Cope

U.S. Nat. Mus. Bull. 17, p. 25, 1880.

=*Eleutherodactylus latrans* (Cope)

COTYPES (7): 10058 (2), 10529 (2), 10751-3, Helotes, Bexar County, Texas, G. W. Marnock, 1878 (10529), Feb. 10, 1880 (10751-3).

Lithodytes megacephalus Cope

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8 (1875), p. 110, pl. 23, fig. 11, 1876.

=*Eleutherodactylus rugosus* (Peters)

HOLOTYPE: 32579, ♀, Pico Blanco, Costa Rica, 6,000 ft., W. M. Gabb.

Lithodytes melanostictus Cope

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8 (1875), p. 109, pl. 23, fig. 10, 1876.

=*Eleutherodactylus melanostictus* (Cope)

HOLOTYPE: 30608, Pico Blanco, Costa Rica, 7,000 ft., W. M. Gabb.

Lithodytes pelviculus Cope

Proc. Amer. Philos. Soc., vol. 17 (1878), p. 89, 1877.

=*Eleutherodactylus rugosus* (Peters)?

HOLOTYPE: 32326, west coast of Central America.

Lithodytes podiciferus Cope

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8 (1875), p. 107, pl. 23, fig. 9, 1876.

=*Eleutherodactylus podiciferus* (Cope)

COTYPES (12): 30662, 30665-75, Pico Blanco, Costa Rica, 5,000-7,000 ft., W. M. Gabb. Cope gave the following varieties in the original description: "Variety A, 30662, 30664-71; variety B, 30672-3; variety C, 30674; and variety D, 30675." 30663 was exchanged to MCZ in 1927; 30664 to MZUM, 1944.

Lithodytes ranoides Cope

Proc. Amer. Philos. Soc., vol. 23, p. 275, 1886.

=*Eleutherodactylus ranoides* (Cope)

COTYPES (4): 14179 (4), Nicaragua, J. F. Bransford.

Lithodytes rhodopis Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 18 (1866), p. 323, 1867.

=*Eleutherodactylus rhodopis* (Cope)

LECTOTYPE: 16558, vicinity of Orizaba, Veracruz, Mexico, F. Sumichrast.

PARATYPES (4): 16557, 16559-61, same data.

Liyla rugulosa Cope

Proc. Amer. Philos. Soc., vol. 11, p. 160, 1869.

=*Eleutherodactylus rugulosus* (Cope)

COTYPES (2): 29771-2, Isthmus of Tehuantepec, Pacific side, Oaxaca, Mexico, F. Sumichrast.

Malachylodes guttilatus Cope

Proc. Amer. Philos. Soc., vol. 18, p. 264, 1879.

=*Syrrophus guttilatus* (Cope)

HOLOTYPE: 9888, Guanajuato, Guanajuato, Mexico, A. Dugès, 1877.

Megophrys abbotti Cochran

Journ. Washington Acad. Sci., vol. 16, p. 446, 1926.

HOLOTYPE: 39097, Balik Papan Bay, eastern Borneo, W. L. Abbott, Feb. 6, 1909.

Megophrys minor Stejneger

Proc. Biol. Soc. Washington, vol. 39, p. 53, 1926.

HOLOTYPE: 68816, above Kuan-hsien, Szechwan, China, 3,000 ft., D. C. Graham, July 5, 1924.

Megophrys omeimontis Liu

Chicago Nat. Hist. Mus., Fieldiana: Zool. Mem., vol. 2, p. 191, figs. 39a-d, 1950.

PARATYPE: 124571, ♂, Mount Omei, Szechwan, China, C. C. Liu, July 26-28, 1940.

Microbatrachylus imitator Taylor

Univ. Kansas Sci. Bull., vol. 28, p. 70, pl. 6, figs. 1a-c, 1942.

HOLOTYPE: 115508, La Esperanza, Chiapas, Mexico, H. M. Smith, May 15, 1940.

Microbatrachylus montanus Taylor

Univ. Kansas Sci. Bull., vol. 28, p. 67, pl. 6, figs. 2a-c, 1942.

HOLOTYPE: 115507, Mount Ovando, Chiapas, Mexico, 6,000 ft., H. M. Smith, Apr. 16, 1940.

Microbatrachylus oaxacae Taylor

Univ. Kansas Sci. Bull., vol. 26, (1939), p. 504, 1940.

PARATYPE: 139723 (formerly UIMNH 16113), Cerro San Felipe, Oaxaca, Mexico, E. H. Taylor, Aug. 18-22, 1938.

Microhyla annamensis Smith

Journ. Nat. Hist. Soc. Siam, vol. 6, p. 47, pl. 5, fig. 2, 1923.

PARATYPES (3): 65837-9, Sui Kat, Langbian Plateau, southern Annam (Vietnam), 1,000 m., M. A. Smith, March 1918.

Microhyla grahami Stejneger

Occ. Pap. Boston Soc. Nat. Hist., vol. 5, p. 119, 1924.

=*Microhyla butleri* Boulenger

HOLOTYPE: 65936, Suifu, Szechwan, China, D. C. Graham, 1923.

PARATYPE: 65937, same data.

Microhyla malcolmi Cochran

Proc. Biol. Soc. Washington, vol. 40, p. 182, 1927.

=*Microhyla berdmorei* (Blyth)

HOLOTYPE: 72172, Pak Jong, Siam (Thailand), Hugh M. Smith, Nov. 15, 1925.

Microhyla okinavensis Stejneger

Proc. Biol. Soc. Washington, vol. 14, p. 189, 1901.

HOLOTYPE: 36553, ♀, Okinawa, Riu Kiu (Ryukyu) Islands, SCMT.

PARATYPES (20): 23906, Kunshiau, Okinawa, Riu Kiu Islands, I. Ijima. 34472-90, Ishigaki, Riu Kiu Islands, A. Owston, April-June 1899. 34491 was exchanged to MZUM in 1944.

Microhyla sowerbyi Stejneger

Occ. Pap. Boston Soc. Nat. Hist., vol. 5, p. 119, 1924.

=*Microhyla butleri* Boulenger

HOLOTYPE: 65309, near Yen-ping-fu, Fukien, China, A. de C. Sowerby.

Microhyla undulata Brown

Proc. Acad. Nat. Sci. Philadelphia, vol. 54, p. 186, 1902.

=*Microhyla okinavensis* Stejneger

COTYPE: 75081 (formerly WIM 5728b), Loo Choo (Ryukyu) Islands, W. H. Furness and H. M. Hiller, 1896.

Montorana ahli Vogt

Zool. Anz., vol. 60, p. 340, 1924.

=*Nanorana pleskei* Günther

PARATYPE: 72739, Dschiesongla, southwest of Tatsienlu, Sikang, China, 3,400 m., H. Weigold.

Paludicola frenata Cope

Proc. U.S. Nat. Mus., vol. 12, p. 143, 1889.

=*Pleurodema bibroni* Tschudi?

HOLOTYPE: 15129, Lota, Chile, U.S. Fish Commission *Albatross*, Feb. 17, 1888.

Pharyngodon petasatus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 17, p. 193, 1865.

=*Tripurion petasatus* (Cope)

HOLOTYPE: 12287, ♀, near Mérida, Yucatán, Mexico, A. Schott.

Philautus gryllus Smith

Proc. Zool. Soc. London, p. 231, pl. 3, figs. 2-5, 1924.

PARATYPES (3): 65834-6, Langbian Peak, south Annam (Vietnam), 2,000 m., M. A. Smith, March 1918.

Philautus hansenae Cochran

Proc. Biol. Soc. Washington, vol. 40, p. 181, 1927.

HOLOTYPE: 70109, ♂, Nong Khor, southeast Siam (Thailand), Hugh M. Smith, Oct. 4, 1925.

PARATYPES (4): 70110, 70112-3, Nong Khor, southeast Siam, Hugh M. Smith, Oct. 4-5, 1925. 70135, Ban Sadet, southeast Siam, Hugh M. Smith, May 30, 1925. 70115 was exchanged to MCZ in 1944; 70111 to MZUM, 1944; 70114 to WCUUM, 1947.

***Philautus nongkhorensis* Cochran**

Proc. Biol. Soc. Washington, vol. 40, p. 179, 1927.

HOLOTYPE: 70108, Nong Khor, southeast Siam (Thailand), Hugh M. Smith, Oct. 5, 1925.

PARATYPES (5): 70099-100, 70102-3, 70105, Nong Khor, southeast Siam, Hugh M. Smith, Oct. 4-5, 1925. 70106 was exchanged to MCZ in 1944; 70101 to MZUM, 1944; 70104 to WCUUM, 1947; 70107 to BMNH, 1928.

***Philautus woodi* Stejneger**

Proc. U.S. Nat. Mus., vol. 28, p. 346, 1905.

HOLOTYPE: 34781, Mount Apo, Mindanao, Philippine Islands, 6,000 ft., E. A. Mearns, June 30, 1904.

PARATYPE: 34782, Mount Apo, Mindanao, Philippine Islands, 6,000 ft., E. A. Mearns, July 9, 1904.

***Phrynixalus anulatus* Stejneger**

Proc. U.S. Nat. Mus., vol. 33, p. 573, 1908.

HOLOTYPE: 35399, Davao, Mindanao, Philippine Islands, E. A. Mearns.

PARATYPE: 35272, Todaya, Mount Apo, Mindanao, Philippine Islands, 4,000 ft., E. A. Mearns, July 2-10, 1904.

***Phyllobates auratus* Girard**

Proc. Acad. Nat. Sci. Philadelphia, vol. 7, p. 226, 1855.

=*Dendrobates auratus* (Girard)

HOLOTYPE: 10307, Taboga Island, Bay of Panama, Panama, C. B. Adams.

***Phyllobates cystignathoides* Cope**

Proc. Amer. Philos. Soc., vol. 17 (1878), p. 89, 1877.

=*Syrrhophus cystignathoides* (Cope)

COTYPES (7): 32402-4, 32406-9, Potrero, near Córdoba, Veracruz, Mexico, F. Sumichrast. 32405 was exchanged to MCZ in 1945.

***Phyllobates flotator* Dunn**

Occ. Pap. Boston Soc. Nat. Hist., vol. 5, p. 389, 1931.

PARATYPE: 50177, Cana, Panama, 1,800-2,000 ft., E. A. Goldman, Mar. 1-24, 1912.

***Phyllobates hylaeformis* Cope**

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8 (1875), p. 107, pl. 23, fig. 8, 1876.

=*Eleutherodactylus hylaeformis* (Cope)

HOLOTYPE: 30687, Pico Blanco, Costa Rica, 7,000 ft., W. M. Gabb.

***Phyllobates limbatus* Cope**

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 154, 1862.

=*Sminthillus limbatus* (Cope)

COTYPES (6): 5205 (6), eastern Cuba, C. Wright.

Phyllobates nubicola Dunn

Occ. Pap. Mus. Zool. Univ. Michigan, No. 151, p. 7, fig., 1924.

PARATYPES (2): 118674-5 (formerly MZUM 58286), Boquete, Chiriquí Province, Panama, H. T. Gaige, Mar. 10, 1923.

Phyllobates sylvatica Barbour and Noble

Bull. Mus. Comp. Zool., vol. 63, p. 396, 1920.

PARATYPE: 118239 (formerly MCZ 5357), Tabaconas, Peru, G. K. Noble, September 1916.

Phyllomedusa appendiculata Lutz

Compt. Rend. Soc. Biol. Paris, vol. 93, p. 139, 1925.

COTYPE: 97147, São Bento, Santa Catarina, Brazil, E. Nahderer, February 1924.

Phyllomedusa azurea Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 355, 1862.

=*Phyllomedusa hypochondrialis* (Daudin)

COTYPE: 5832, Paraguay, T. J. Page.

Phyllomedusa guttata Lutz

Compt. Rend. Soc. Biol. Paris, vol. 90, p. 241, 1924.

COTYPES (5): 96244-5, Tijuca, Rio de Janeiro, A. Lutz, November 1923. 96246, tadpoles, Tijuca, Rio de Janeiro, Brazil, A. Lutz, Feb. 2, 1920. 96317, Sumaré, Rio de Janeiro, Brazil, A. Lutz, Jan. 28, 1920. 96388, Paineiras, on the slope of Corcovado, Rio de Janeiro, Brazil, A. Lutz, November 1923.

Platymanthis aculeodactylus Brown

Bull. Mus. Comp. Zool., vol. 107, p. 46, pl. 5, fig. 1, 1952.

HOLOTYPE: 119769, ♀, Torakina, Bougainville Island, Solomon Islands, W. L. Necker, 1944.

PARATYPE: 119580, juvenile, Torakina, Bougainville Island, Solomon Islands, W. L. Necker, Aug. 2, 1944.

Plectrohyla sagorum Hartweg

Occ. Pap. Mus. Zool. Univ. Michigan, No. 437, p. 2, pl. 1, figs. 1-3, 1941.

PARATYPE: 118659 (formerly MZUM 87874), Mount Ovando, Soconusco, Chiapas, Mexico, N. Hartweg, Aug. 31, 1937.

Polypedates omeimontis Stejneger

Occ. Pap. Boston Soc. Nat. Hist., vol. 5, p. 120, 1924.

=*Rhacophorus omeimontis* (Stejneger)

HOLOTYPE: 66548, Shin-Kai-Si, Mount Omei, Szechwan, China, D. C. Graham, 1923.

Polypedates owstoni Stejneger

U.S. Nat. Mus. Bull. 58, p. 149, 1907.

=*Rhacophorus owstoni* (Stejneger)

HOLOTYPE: 34333, ♀, Ishigaki, Yaeyama Group, Loo Choo (Ryukyu) Islands, A. Owston, April-June 1899.

PARATYPES (15): 34330, 34332, 34334-5, 34337, 34339-48, same data. 34349 and 34331 were exchanged to MCZ in 1909 and 1944; 34336 and 34338 to WCUUM, 1947.

Polypedates viridis Hallowell

Proc. Acad. Nat. Sci. Philadelphia, vol. 12 (1860), p. 500, 1861.

=*Rhacophorus viridis* (Hallowell)

HOLOTYPE: 25397, Okinawa, Loo Choo (Ryukyu) Islands, W. Stimpson, December 1854.

Prostherapis boulengeri Barbour

Proc. Biol. Soc. Washington, vol. 22, p. 89, 1909. (New name for *Prostherapis femoralis* T. Barbour, 1905, preoccupied by *Prostherapis femoralis* Boulenger, 1883.)

=*Phyllobates boulengeri* (Barbour)

COTYPES (3): 52406, 118232-3 (formerly MCZ 2422), Gorgona Island, Colombia, W. W. Brown, Jr., June 17-July 2, 1904.

Prostherapis inguinalis Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 20, p. 137, 1868.

COTYPE: 4349, Truandó River, New Granada (Colombia), A. Schott.

Protopipa carvalhoi Miranda-Ribeiro

O Campo, Rio de Janeiro, p. 56, figs. 1-3, January 1937.

=*Hemipipa carvalhoi* (Miranda-Ribeiro)

PARATYPES (2): 102685-6, Poçoão, Pernambuco, Brazil, 1,000 m., A. L. de Carvalho, 1936.

Pseudacris streckeri illinoensis Smith

Bull. Chicago Acad. Sci., vol. 9, p. 190, pl. 1, 1951.

PARATYPE: 134273 (formerly UIMNH 24089), 3 miles east of Beardstown, Cass County, Illinois, D. and A. Smith, Mar. 6, 1951.

Pseudis bolbodactyla Lutz

Compt. Rend. Soc. Biol. Paris, vol. 93, p. 138, 1925.

COTYPE: 97022, Lassance, Minas Gerais, Brazil, A. Lutz.

Pseudophryne blanchardi Loveridge

Occ. Pap. Boston Soc. Nat. Hist., vol. 8, p. 91, 1933.

PARATYPE: 118771 (formerly MZUM 83367), Millgrove, Victoria, Australia, F. N. Blanchard, Apr. 1, 1928.

Ptychohyla adipoventris Taylor

Univ. Kansas Sci. Bull., vol. 30, p. 41, 1944.

PARATYPE: 114551, Agua del Obispo, Guerrero, Mexico, H. M. Smith, Aug. 31, 1939.

Rana areolata Baird and Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 173, 1852.

=*Rana areolata areolata* Baird and Girard

HOLOTYPE: 3304, Indianola, Calhoun County, Texas, J. H. Clark.

Rana areolata aesopus Cope

Proc. Amer. Philos. Soc., vol. 23, p. 517, 1886.

HOLOTYPE: 4743, Micanopy, Alachua County, Florida, T. H. Bean.

Rana aurora Baird and Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 174, 1852.

=*Rana aurora aurora* Baird and Girard

COTYPES (4): 11711 (4), Puget Sound (Washington), U.S. Exploring Expedition, May 1841.

Rana berlandieri Baird

Reptiles of the boundary, in Report U.S. and Mexican Bound. Surv., vol. 2, pt. 2, p. 27, pl. 36, figs. 7-10, 1859.

=*Rana pipiens berlandieri* Baird

COTYPES (10): 3293 (9), 131513, Brownsville, Cameron County, Texas, S. Van Vliet. Two cotypes from 3293 were exchanged to MCZ in 1879.

Rana boylei Baird

Proc. Acad. Nat. Sci. Philadelphia, vol. 7, p. 62, 1854.

=*Rana boylei boylei* Baird

COTYPES (2): 3370 (2), El Dorado County, California, C. C. Boyle.

Rana cantabrigensis evittata Cope

U.S. Nat. Mus. Bull. 34, p. 437, 1889.

=*Rana sylvatica cantabrigensis* Baird

COTYPES (24): 5169 (7), Puget Sound (Washington), U.S. Exploring Expedition. 5364 (3), Welthy R.? (probably Canada), R. W. Kennicott. 5365 (2), 5366 (3), 5922 (2), Moose Factory, British America (Canada), C. Drexler. 5929, British America? (Canada), R. W. Kennicott. 6505, Big Island, Great Slave Lake (Canada), J. Reid. 9385 (3), St. Catherine's, Canada, D. W. Beadle. 13727, Lake Aleknagik, north of Nushagak, Alaska, C. L. McKay, Sept. 27, 1883. 14495, Nulato River, Alaska, W. H. Dall.

Rana cantabrigensis latiremis Cope

Proc. Amer. Philos. Soc., vol. 23, p. 520, 1886.

=*Rana sylvatica cantabrigensis* Baird

COTYPES (4): 13723-6, Lake Aleknagik, north of Nushagak, Alaska, C. L. McKay, Sept. 27, 1883.

Rana capito LeConte

Proc. Acad. Nat. Sci. Philadelphia, vol. 7, p. 425, pl. 5, 1855.

=*Rana areolata capito* LeConte

HOLOTYPE: 5903, Riceboro, Liberty County, Georgia, J. L. LeConte.

Rana capito stertens Schwartz and Harrison

Proc. Biol. Soc. Washington, vol. 69, p. 136, 1956.

=*Rana areolata capito* LeConte

PARATYPES (2): 137279-80, 6 miles north of Cainhoy, Berkeley County, South Carolina, J. R. Harrison, J. A. Quinby, and A. Schwartz, Sept. 4, 1955.

Rana cascadae Slater

Herpetologica, vol. 1, p. 145, pl. 14, 1939.

=*Rana aurora cascadae* Slater

HOLOTYPE: 108689, Elysian Fields, Rainier National Park, Pierce County, Washington, J. R. Slater, June 19, 1938.

PARATYPES (2): 108690-1, same data.

Rana chaochiaensis Liu

Journ. West China Border Res. Soc., ser. B, vol. 16, p. 7, 1946.

PARATYPES (2): 124539-40, Chao-chiao, Sikang, China, C. C. Liu, May 7, 1942.

Rana draytonii Baird and Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 174, 1852.

=*Rana aurora draytonii* Baird and Girard

COTYPES (6): 11497 (6), San Francisco (San Francisco County), California, U.S. Exploring Expedition, 1841.

Rana fisheri Stejneger

North Amer. Fauna, No. 7, p. 227, pl. 3, figs. 5a-c, 1893.

=*Rana pipiens fisheri* Stejneger

HOLOTYPE: 18957, Vegas Valley, Clark County, Nevada, V. O. Bailey, Mar. 13, 1891.

PARATYPES (8): 18958-9, Vegas Valley, Clark County, Nevada, V. O. Bailey, Mar. 9-13, 1891. 18960-5, Las Vegas, Clark County, Nevada, E. W. Nelson, Mar. 9, 1891. 18966 was exchanged to MCZ in 1919.

Rana grisea milneana Loveridge

Bull. Mus. Comp. Zool., vol. 101, p. 414, 1948.

HOLOTYPE: 19173, ♀, Kwatto Branch Mission, Milne Bay, British New Guinea, 50 ft., J. F. Cassell, March 1944.

Rana grylio Stejneger

Proc. U.S. Nat. Mus., vol. 24, p. 212, fig., 1901.

HOLOTYPE: 27443, Bay Saint Louis, Hancock County, Mississippi, G. S. Miller, Jr., summer of 1900.

PARATYPES (10): 3688a-b, Pensacola, Escambia County, Florida, R. W. Jeffries. 27444-6, Bay Saint Louis, Hancock County, Mississippi, G. S. Miller, Jr., summer of 1900. 29007-11, Kissimmee, Osceola County, Florida, E. A. Mearns, May 4, 1902. 27447 was exchanged to B. B. Brandt.

Rana gryllus LeConte

Ann. Lyc. Nat. Hist. New York, vol. 1, p. 282, 1825.

=*Acris gryllus gryllus* (LeConte)

PROBABLE COTYPES (14): 3564 (7), 5909 (7), Georgia?, J. L. LeConte.

Rana jugans Stejneger

Proc. Biol. Soc. Washington, vol. 39, p. 53, 1926.

=*Staurois mantzorum* (David)

HOLOTYPE: 67819, near Wenchwan, about 100 km. northwest of Chengtu, Szechwan, China, D. C. Graham, Aug. 13, 1924.

Rana lecontei Baird and Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 301, 1853.

=*Rana aurora draytonii* Baird and Girard

COTYPES (2): 3362 (2), San Francisco (San Francisco County), California, J. L. LeConte, February 1850.

Rana longipes Hallowell

Report upon the reptiles collected on the survey, in Report U.S. Pacific R. R. Explor. and Surv., vol. 10, pt. 4, No. 1, p. 20, 1859.

=*Rana aurora draytonii* Baird and Girard

COTYPES (2): 3366, 3376, El Paso Creek, California, A. L. Heermann. These specimens are also cotypes of *Rana nigricans* Hallowell, 1854 (not of Agassiz, 1850).

Rana macrodon macrocephala Inger

Chicago Nat. Hist. Mus., Fieldiana: Zool., vol. 33, p. 287, 1954.

=*Rana magna macrocephala* Inger

PARATYPES (3): 39963, Maaguas River, Lagonoy Gulf, Philippine Islands, U.S. Fish Commission *Albatross*, June 17, 1909. 39968, Antipolo, Luzon, Philippine Islands, U.S. Fish Commission *Albatross*, Jan. 26, 1908. 57841, Luzon, Philippine Islands, J. Hurter, 1904.

Rana macrodon visayanus Inger

Chicago Nat. Hist. Mus., Fieldiana: Zool., vol. 33, p. 289, 1954.

=*Rana magna visayanus* Inger

PARATYPES (9): 78072-8, Iloilo, Island of Panay, Philippine Islands, H. C. Kellers, Apr. 10, 1929. 121597-8, 5 miles south of Tarragone, Leyte, Philippine Islands, D. J. Pletsch.

Rana magna Stejneger

Smithsonian Misc. Coll., vol. 52, p. 437, 1909.

=*Rana magna magna* Stejneger

HOLOTYPE: 35231, ♂, Mount Apo, Mindanao, Philippine Islands, 4,000-6,000 ft., E. A. Mearns, June-July 1904.

PARATYPES (17): 35232-6, 35238-40, 35242-50, same data. 35241 was exchanged to MCZ in 1910; 35237 to MHNB, 1909.

Rana marmorata Hallowell

Proc. Acad. Nat. Sci. Philadelphia, vol. 12 (1860), p. 500, 1861.

=*Rana nigromaculata* Hallowell

COTYPES (2): 12159 (2), ♀, Simoda, Hondo, Japan, Morrow.

Rana mascareniensis Duméril and Bibron

Épétologie générale ou histoire naturelle complète des reptiles, vol. 8, p. 350, 1841.

PROBABLE COTYPES (2): 10975 (2), Island of Réunion, Mascarene Islands (Indian Ocean), MHNP.

Rana mearnsi Stejneger

Proc. U.S. Nat. Mus., vol. 28, p. 343, 1905.

HOLOTYPE: 35258, Baganga River, Mindanao, Philippine Islands, 1,000-5,000 ft., E. A. Mearns, Apr. 28-May 7, 1904. Catalog states: "Originally preserved in blackberry brandy."

Rana megapoda Taylor

Univ. Kansas Sci. Bull., vol. 23, p. 310, pl. 28, 1942.

PARATYPES (9): 113998-114005, La Palma, Michoacán, Mexico, H. M. Smith, Mar. 23, 1939. 137197, near Chapala, Jalisco, Mexico, H. M. Smith, July 2, 1935.

Rana montezumae Baird

Proc. Acad. Nat. Sci. Philadelphia, vol. 7, p. 61, 1854.

COTYPES (15): 3344, 39383-96 (formerly 3344), City of Mexico, Distrito Federal, Mexico. 3344 is the lectotype. 39397 was exchanged to MCZ in 1909.

Rana montezumae concolor Baird

Proc. Acad. Nat. Sci. Philadelphia, vol. 7, p. 61, 1854.

=*Rana montezumae* Baird

LECTOTYPE: 81101 (formerly 10247), Guanajuato, Mexico, A. Dugès, 1879.

PARALECTOTYPES (3): 10247 (3), same data.

Rana nigricans Hallowell

Proc. Acad. Nat. Sci. Philadelphia, vol. 7, p. 96, 1854. (Not of Agassiz, 1850.)

=*Rana aurora draytonii* Baird and Girard

COTYPES (2): 3366, 3376, El Paso Creek, California, A. L. Heermann. These specimens are also the cotypes of *Rana longipes* Hallowell, in Report upon the reptiles collected on the survey, in Report U.S. Pacific R.R. Explor. and Surv., vol. 10, pt. 4, No. 1, p. 20, 1859. The catalog gives the locality for 3366 as "Pacific R.R. Survey," and for 3376 as simply "California."

Rana onca Cope

in Yarrow, Report upon the collections of batrachians and reptiles . . . 1871, 1872, 1873, and 1874, in Report U.S. Geogr. and Geol. Expl. and Surv. West of One Hundr. Mer., vol. 5, p. 523, pl. 25, figs. 1-3, 1875.

=*Rana pipiens brachycephala* Cope

COTYPE: 25331, ♀, Utah, H. C. Yarrow, 1872.

Rana pretiosa Baird and Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 378, 1853.

=*Rana pretiosa pretiosa* Baird and Girard

COTYPES (5): 11409 (5), Puget Sound (Washington), U.S. Exploring Expedition, 1841.

Rana sevosia Goin and Netting

Ann. Carnegie Mus., vol. 28, p. 137, pl. 12, figs. 1-2, 1940.

=*Rana areolata sevosia* Goin and Netting

PARATYPE: 109226 (formerly CM 18116), Van Cleave, Jackson County, Mississippi, CM, Oct. 21, 1930.

Rana sierramadrensis Taylor

Univ. Kansas Sci. Bull., vol. 25 (1938), p. 397, pl. 39, fig. 1, 1939.

PARATYPE: 139724 (formerly UIMNH 27054), near Agua del Obispo, between Rincón and Cajones, Guerrero, Mexico, E. H. Taylor, July 21, 1936.

Rana sinuata Baird

Proc. Acad. Nat. Sci. Philadelphia, vol. 7, p. 61, 1854.

=*Rana septentrionalis* Baird

COTYPES (5): 3420 (5), Sacketts Harbor, Garrison Creek, Jefferson County, New York, S. F. Baird, summer of 1850.

Rana sylvatica cherokiana Witschi

Proc. Iowa Acad. Sci., vol. 60, p. 764, figs. 1-4, 1953.

HOLOTYPE: 134417, ♂, Murphy, Cherokee County, North Carolina, J. C. Nicholls, Jr., Jan. 20, 1953.

PARATYPE: 134418, ♀, same data.

Rana tsushimensis Stejneger

U.S. Nat. Mus. Bull. 58, p. 116, figs. 94-98, 1907.

HOLOTYPE: 17519, ♀, Tsushima Island, Japan, P. L. Jouy, May 1885.

PARATYPE: 17520, ♂, same data.

Rana virescens brachycephala Cope

U.S. Nat. Mus. Bull. 34, p. 403, fig. 101, 1889.

=*Rana pipiens brachycephala* Cope

COTYPES (15): 3363 (15), Yellowstone River, Montana, F. V. Hayden.

Rana woodworthi Taylor

Philippine Journ. Sci., vol. 22, p. 519, pl. 1, figs. 1-2, 1923.

PARATYPE: 139725, Polillo, Philippine Islands, E. H. Taylor, 1920.

Rana yakani Taylor

Philippine Journ. Sci., vol. 21, p. 262, pl. 1, fig. 1, pl. 2, fig. 1, 1922.

=*Rana signata* Günther

PARATYPE: 139726, Basilan Island, Philippine Islands, E. H. Taylor, October 1920.

Ranoidea flavoviridis Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 422, 1853.

=*Hyla lesueurii* Duméril and Bibron

COTYPES (2): 15484-5, Wollongong, Illawara, New South Wales, Australia, U.S. Exploring Expedition, January 1840.

Ranoidea resplendens Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 422, 1853.

=*Hyla aurea* Lesson

HOLOTYPE: 15478, Wollongong, Illawara, New South Wales, Australia, U.S. Exploring Expedition, January 1840.

Rhacophorus baluensis Inger

Journ. Washington Acad. Sci., vol. 44, p. 250, 1954.

HOLOTYPE: 130215, ♀, Bundu Tuhan, Mount Kina Balu, North Borneo, 4,500 ft., D. H. Johnson, July 5, 1951.

PARATYPE: 130217, ♂, same data. 130216 was exchanged to CNHM in 1954.

Rhacophorus chenfui Liu

Journ. West China Border Res. Soc., ser. B, vol. 15, p. 35, pl. 3, figs. 13-15, 1945.

PARATYPE: 124564, ♂, Mount Omei, Szechwan, China, C. C. Liu.

Rhacophorus pleurostictus batangensis Vogt

Zool. Anz., vol. 60, p. 341, 1924.

=*Rhacophorus bambusicola* Barbour

PARATYPE: 72741, Batang, Sikang Province, China, H. Weigold.

Rana nigricans Hallowell

Proc. Acad. Nat. Sci. Philadelphia, vol. 7, p. 96, 1854. (Not of Agassiz, 1850.)

=*Rana aurora draytonii* Baird and Girard

COTYPES (2): 3366, 3376, El Paso Creek, California, A. L. Heermann. These specimens are also the cotypes of *Rana longipes* Hallowell, in Report upon the reptiles collected on the survey, in Report U.S. Pacific R.R. Explor. and Surv., vol. 10, pt. 4, No. 1, p. 20, 1859. The catalog gives the locality for 3366 as "Pacific R.R. Survey," and for 3376 as simply "California."

Rana onca Cope

in Yarrow, Report upon the collections of batrachians and reptiles . . . 1871, 1872, 1873, and 1874, in Report U.S. Geogr. and Geol. Expl. and Surv. West of One Hundr. Mer., vol. 5, p. 528, pl. 25, figs. 1-3, 1875.

=*Rana pipiens brachycephala* Cope

COTYPE: 25331, ♀, Utah, H. C. Yarrow, 1872.

Rana pretiosa Baird and Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 378, 1853.

=*Rana pretiosa pretiosa* Baird and Girard

COTYPES (5): 11409 (5), Puget Sound (Washington), U.S. Exploring Expedition, 1841.

Rana sevosia Goin and Netting

Ann. Carnegie Mus., vol. 28, p. 137, pl. 12, figs. 1-2, 1940.

=*Rana areolata sevosia* Goin and Netting

PARATYPE: 109226 (formerly CM 18116), Van Cleave, Jackson County, Mississippi, CM, Oct. 21, 1930.

Rana sierramadrensis Taylor

Univ. Kansas Sci. Bull., vol. 25 (1938), p. 397, pl. 39, fig. 1, 1939.

PARATYPE: 139724 (formerly UIMNH 27054), near Agua del Obispo, between Rincón and Cajones, Guerrero, Mexico, E. H. Taylor, July 21, 1936.

Rana sinuata Baird

Proc. Acad. Nat. Sci. Philadelphia, vol. 7, p. 61, 1854.

=*Rana septentrionalis* Baird

COTYPES (5): 3420 (5), Sacketts Harbor, Garrison Creek, Jefferson County, New York, S. F. Baird, summer of 1850.

Rana sylvatica cherokeiana Witschi

Proc. Iowa Acad. Sci., vol. 60, p. 764, figs. 1-4, 1953.

HOLOTYPE: 134417, ♂, Murphy, Cherokee County, North Carolina, J. C. Nicholls, Jr., Jan. 20, 1953.

PARATYPE: 134418, ♀, same data.

Rana tsushimensis Stejneger

U.S. Nat. Mus. Bull. 58, p. 116, figs. 94-98, 1907.

HOLOTYPE: 17519, ♀, Tsushima Island, Japan, P. L. Jouy, May 1885.

PARATYPE: 17520, ♂, same data.

Rana virescens brachycephala Cope

U.S. Nat. Mus. Bull. 34, p. 403, fig. 101, 1889.

=*Rana pipiens brachycephala* Cope

COTYPES (15): 3363 (15), Yellowstone River, Montana, F. V. Hayden.

Rana woodworthi Taylor

Philippine Journ. Sci., vol. 22, p. 519, pl. 1, figs. 1-2, 1923.

PARATYPE: 139725, Polillo, Philippine Islands, E. H. Taylor, 1920.

Rana yakani Taylor

Philippine Journ. Sci., vol. 21, p. 262, pl. 1, fig. 1, pl. 2, fig. 1, 1922.

=*Rana signata* Günther

PARATYPE: 139726, Basilan Island, Philippine Islands, E. H. Taylor, October 1920.

Ranoidea flavoviridis Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 422, 1853.

=*Hyla lesueurii* Duméril and Bibron

COTYPES (2): 15484-5, Wollongong, Illawara, New South Wales, Australia, U.S. Exploring Expedition, January 1840.

Ranoidea resplendens Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 422, 1853.

=*Hyla aurea* Lesson

HOLOTYPE: 15478, Wollongong, Illawara, New South Wales, Australia, U.S. Exploring Expedition, January 1840.

Rhacophorus baluensis Inger

Journ. Washington Acad. Sci., vol. 44, p. 250, 1954.

HOLOTYPE: 130215, ♀, Bundu Tuhan, Mount Kina Balu, North Borneo, 4,500 ft., D. H. Johnson, July 5, 1951.

PARATYPE: 130217, ♂, same data. 130216 was exchanged to CNHM in 1954.

Rhacophorus chenfui Liu

Journ. West China Border Res. Soc., ser. B, vol. 15, p. 35, pl. 3, figs. 13-15, 1945.

PARATYPE: 124564, ♂, Mount Omei, Szechwan, China, C. C. Liu.

Rhacophorus pleurostictus batangensis Vogt

Zool. Anz., vol. 60, p. 341, 1924.

=*Rhacophorus bambusicola* Barbour

PARATYPE: 72741, Batang, Sikang Province, China, H. Weigold.

Scaphiopus albus Garman

Proc. Amer. Assoc. Adv. Sci., vol. 25, p. 194, 1877.

=*Scaphiopus holbrookii albus* Garman

COTYPE: 52403 (formerly MCZ 1452), Key West, Monroe County, Florida, S. Garman, 1879.

Scaphiopus dugesi Brocchi

Bull. Soc. Philom. Paris, ser. 7, vol. 3, p. 23, 1879.

=*Scaphiopus hammondii multiplicatus* Cope

COTYPES (3): 16205-7, Silao de la Victoria, Guanajuato, Mexico, A. Dugès.

Scaphiopus multiplicatus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 15, p. 52, 1863.

=*Scaphiopus hammondii multiplicatus* Cope

HOLOTYPE: 3694, Valley of Mexico, Mexico, J. Potts.

Scaphiopus varius Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 15, p. 52, 1863.

=*Scaphiopus couchii* Baird

COTYPES (3): 5893 (3), Cape San Lucas, Baja California, Mexico, J. Xantus.

Scutiger pingi Liu

Journ. West China Border Res. Soc., ser. B., vol. 14, p. 35, figs. 1-5, 1943.

PARATYPE: 124567, Chaochiao, Sikang, China, C. C. Liu, 1942.

Scutiger schmidti Liu

Copeia, No. 2, p. 123, fig. 1, 1947.

PARATYPES (3): 124568-70, Mount Omei, Szechwan, China, C. C. Liu.

Scytopsis boulengeri Cope

U.S. Nat. Mus. Bull. 32, p. 12, 1887.

=*Hyla boulengeri* (Cope)

HOLOTYPE: 13974, Nicaragua, J. A. McNeil, Aug. 29, 1884.

Scytopsis hebes Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 354, 1862.

=*Phrynohyas hebes* (Cope)

HOLOTYPE: 5837, Paraguay, T. J. Page.

Spea stagnalis Cope

in Yarrow, Report upon the collections of batrachians and reptiles . . .

1871, 1872, 1873, and 1874, in Report U.S. Geogr. and Geol. Expl. and Surv. West of One Hundr. Mer., vol. 5, p. 525, pl. 25, figs. 6-8, 1875.

=*Scaphiopus hammondii hammondii* Baird

COTYPES (4): 8188 (2), 8563, 25335, Utah, H. C. Yarrow, 1872.

Sphoerohyla habra Goin

Caldasia, vol. 8, p. 18, 1957.

PARATYPE: 138859, Letitia, Amazonas Comissario, Colombia, E. R. Allen, 1956.

Staurois kangtingensis Liu

Chicago Nat. hist. Mus., Fieldiana: Zool. Mem., vol. 2, p. 349, figs. 89-a-b, 1950.

PARATYPE: 124559, ♂, Kangting, Sikang, China, C. C. Liu, 1942.

Staurois lifanensis Liu

Journ. West China Border Res. Soc., ser. B, vol. 15, p. 33, pl. 2, figs. 10-11, 1945.

PARATYPE: 124560, ♂, Lifan, Sikang, China, C. C. Liu, 1942.

Staurois loloensis Liu

Chicago Nat. Hist. Mus. Fieldiana: Zool. Mem., vol. 2, p. 353, 1950.

PARATYPES (3): 124561, ♀, Lo-lo-kou, Sikang, China, C. C. Liu, 1942. 79707, ♀, Yao-Gi (Yaochin), near Mupin, Szechwan, China, 8,000 ft., D. C. Graham, July 14, 1929. 93857, near Mupin, Szechwan, China, D. C. Graham, 1931.

Syrrhophus campi Stejneger

Proc. Biol. Soc. Washington, vol. 28, p. 131, 1915.

HOLOTYPE: 52290, Brownsville, Cameron County, Texas, R. D. Camp, Mar. 31, 1915.

PARATYPES (5): 52291, same data. 52304-7, Brownsville, Cameron County, Texas, R. D. Camp.

Syrrhophus interorbitalis Langebartel and Shannon

Herpetologica, vol. 12, p. 161, figs. 1-2, 1956.

PARATYPE: 139727 (formerly UIMNH 38098), 36 miles north of Mazatlán, Sinaloa, Mexico, D. A. Langebartel, Nov. 17, 1955.

Syrrhophus latodactylus Taylor

Univ. Kansas Sci. Bull., vol. 26 (1939), p. 397, fig. 7, pl. 43, 1940.

PARATYPE: 139728 (formerly UIMNH 13289), Sabinas Hidalgo, Nuevo León, Mexico, E. H. Taylor, June 17, 1936.

Syrrhophus leprus Cope

Proc. Amer. Philos. Soc., vol. 18, p. 268, 1879.

HOLOTYPE: 10040, Santa Efigenia, Tehuantepec, Oaxaca, Mexico, F. Sumichrast.

Syrrhophus modestus Taylor

Univ. Kansas Sci. Bull., vol. 28, p. 304, pl. 29, 1942.

PARATYPE: 139729 (formerly UIMNH 13301), Hacienda Paso del Río, Colima, Mexico, H. M. Smith, 1935.

Syrrhophus nebulosus Taylor

Univ. Kansas Sci. Bull., vol. 29, p. 353, pl. 27, figs. 3-5, 1943.

PARATYPE: 134274 (formerly UIMNH 13299), Tuxtla Gutiérrez, Chiapas, Mexico, H. M. Smith and E. H. Taylor, Sept. 4, 1935.

Syrrhophus pipilans Taylor

Proc. Biol. Soc. Washington, vol. 53, p. 95, pl. 1, 1940.

PARATYPE: 134275 (formerly UIMNH 13316), Agua del Obispo, Guerrero, Mexico, E. H. Taylor, June 24, 1938.

Syrrhophus rubrimaculatus Taylor and Smith

Proc. U.S. Nat. Mus., vol. 95, p. 583, 1945.

HOLOTYPE: 114070, ♂, La Esperanza, Chiapas, Mexico, Dr. and Mrs. H. M. Smith, May 13, 1940.

PARATYPES (19): 114054-69, 114071-2, La Esperanza, Chiapas, Mexico, H. M. Smith, Apr. 5-May 25, 1940. 114073, Rancho Las Gradadas, 4 km. northwest of La Esperanza, Chiapas, Mexico, H. M. Smith, May 20, 1940.

Syrrhophus smithi Taylor

Proc. U.S. Nat. Mus., vol. 89, p. 43, pl. 1, 1940.

HOLOTYPE: 108594, 15 miles west of Galeana, Nuevo León, Mexico, 5,200 ft. H. M. Smith, Oct. 13, 1939.

Telmatobius niger Barbour and Noble

Bull. Mus. Comp. Zool., vol. 63, p. 413, 1920.

PARATYPES (2): 38118-9, Palmyra Desert, Ecuador, 10,500 ft., T. Barbour, 1909.

Tetraprion jordani Stejneger and Test

Proc. U.S. Nat. Mus., vol. 14, p. 167, pl. 3, 1892.

HOLOTYPE: 12274 (formerly 6862), Guayaquil, Ecuador.

Thoropa lutzi Cochran

Proc. Biol. Soc. Washington, vol. 51, p. 41, 1938.

=*Eupsophus lutzi* (Cochran)

HOLOTYPE: 97622, ♂, Recreio dos Bandeirantes, Rio de Janeiro, Brazil, B. Lutz, D. Cochran, J. Venancio, Feb. 9-10, 1935.

PARATYPES (6): 96297-8, ♂, ♂, Tijuca, Rio de Janeiro, Brazil, A. Lutz, 96422-3, ♂ and tadpoles, Independencia, Petrópolis, Rio de Janeiro, Brazil, A. Lutz, 1926. 97328, Sumaré, Rio de Janeiro, Brazil, A. Lutz, Feb. 15, 1924. 97623, same data as holotype. 97327 was exchanged to LC in 1944.

Tomodactylus albolabris Taylor

Univ. Kansas Sci. Bull., vol. 29, p. 351, 1943.

PARATYPE: 139730 (formerly UIMNH 15974), Agua del Obispo, Guerrero, Mexico, E. H. Taylor, 1941.

Tomodactylus angustidigitorum Taylor

Univ. Kansas Sci. Bull., vol. 26 (1939), p. 494, pl. 55, figs. 1-1b, 1940.

PARATYPE: 137201, Zacapú, Michoacán, Mexico, E. H. Taylor, Sept. 6-7, 1938.

Tomodactylus macrotympanum Taylor

Univ. Kansas Sci. Bull., vol. 26 (1939), p. 496, pl. 55, figs. 2-2b, 1940.

PARATYPE: 137202, 18 miles south of Jacala, Hidalgo, Mexico, E. H. Taylor, July 2, 1936.

Trachycephalus scutigerus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 15, p. 45, 1863.

=*Hyla septentrionalis brunnea* (Gosse)

PARATYPE: 6267, Jamaica, C. B. Adams.

Trachycephalus wrightii Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 15, p. 45, 1863.

=*Hyla wrightii* (Cope)

HOLOTYPE: 5174, Guantánamo District, Cuba, C. Wright.

Zachaenus roseus Cope

Proc. U.S. Nat. Mus., vol. 12, p. 142, 1889.

HOLOTYPE: 15126, Port Otway, Patagonia, Argentina, L. A. Lee, Feb. 9, 1888. This specimen is now nearly disintegrated.

Zachaenus sawayae Cochran

Herpetologica, vol. 8, p. 111, figs. a-d, 1953.

HOLOTYPE: 125530, Banhado, Paraná, Brazil, R. Hertel, July 1945.

Class Reptilia

Order Squamata

Suborder Sauria

**Ablepharus gloriosus** Stejneger

Proc. U.S. Nat. Mus., vol. 16, p. 723, 1894.

=*Cryptoblepharus gloriosus* (Stejneger)

HOLOTYPE: 20463, Gloriosa Island, Madagascar, W. L. Abbott, January 1873.

PARATYPES (3): 20464-6, same data.

Agama agama turuensis Loveridge

Bull. Mus. Comp. Zool., vol. 72, p. 376, 1932.

=*Agama agama elgonis* Lönnberg

PARATYPES (3): 95597-9 (formerly MCZ 30709-10), Unyanganyi, Turu, Tanganyika, A. Loveridge, 1929.

Agama rueppelli occidentalis Parker

Journ. Linn. Soc. London, Zool., vol. 38, p. 225, 1932.

COTYPE: 66927, between Abyssinia and British East Africa (between Ethiopia and Kenya), E. A. Mearns, 1912.

Agama rueppelli septentrionalis Parker

Journ. Linn. Soc. London, Zool., vol. 38, p. 225, 1932.

COTYPES (2): 49056, British East Africa (probably Kenya), E. Heller. 66901, Mount Nyeri, south of Lake Rudolph, British East Africa (Kenya), E. A. Mearns, July 13, 1912.

Ameiva atrigularis Garman

Bull. Essex Inst., vol. 19, p. 2, 1887.

COTYPE: 120777, Trinidad, West Indies, C. S. Cazabon, 1879.

Ameiva birdorum Grant

Journ. Dept. Agr. Puerto Rico, vol. 16, p. 160, 1932.

PARATYPE: 120778, Diablo Island, opposite Fajardo, Puerto Rico, C. Grant, Mar. 11, 1932.

Ameiva chrysolema Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 20, p. 127, 1868.

=*Ameiva chrysolema chrysolema* Cope

COTYPES (2): 12140, 12142, within 25 miles of Port-au-Prince (given erroneously as from Gonave Island in original description), Haiti, A. C. Younglove, January-June 1866.

Ameiva chrysolema woodi Cochran

Occ. Pap. Boston Soc. Nat. Hist., vol. 8, p. 181, 1934.

PARATYPES (7): 59434, La Tortue Island, Haiti, W. L. Abbott, Jan. 31, 1917. 95127-32, La Tortue Island, Haiti, *Utowana* Expedition, Apr. 2, 1934.

Ameiva corvina Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 13, p. 312, 1861.

COTYPES (2): 52215-6 (formerly MCZ 3616), Sombrero Island?, West Indies, A. H. Riise?.

Ameiva eutropia Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 62, 1862.

=*Ameiva festiva festiva* (Lichtenstein and von Martens)

COTYPES (3): 4320 (3), Truando, New Granada (Colombia), A. Schott.

Ameiva panchlora Barbour

Proc. New England Zool. Club, vol. 7, p. 83, 1921.

=*Ameiva ameiva fuliginosa* (Cope)

HOLOTYPE: 13879, Old Providence Island, Colombia, U.S. Fish Commission *Albatross* Apr. 4-9, 1884.

PARATYPE: 63398, same data. 63397 was exchanged to MCZ in 1921.

***Ameiva plei exsul* Cope**

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 66, 1862.

=*Ameiva exsul* Cope

HOLOTYPE: 30696, Water Island (Virgin Islands), A. H. Riise.

***Ameiva polops* Cope**

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 66, 1862.

HOLOTYPE: 30695, St. Croix Island (Virgin Islands), A. H. Riise.

***Ameiva undulata amphigramma* Smith and Laufe**

Trans. Kansas Acad. Sci., vol. 48, p. 338, 1945.

PARATYPES (14): 133837-8, Atoyac, Veracruz, Mexico, H. M. Smith, Dec. 17, 1938. 133839, Orizaba, Veracruz, Mexico, H. M. Smith, 1938. 133840-8, Potrero Viejo, Veracruz, Mexico, H. M. Smith, 1938-39. 133920, Cosalapa, Oaxaca, Mexico, H. M. Smith, Jan. 11, 1939. 134298 (formerly UIMNH 27352), Tierra Colorado, Veracruz, Mexico, E. H. Taylor and H. M. Smith, July 15, 1932.

***Ameiva undulata dextra* Smith and Laufe**

Univ. Kansas Sci. Bull., vol. 31, p. 54, fig. 2c, 1946.

PARATYPES (5): 133849-53, 8 km. east of Coyuca, Guerrero, Mexico, Dr. and Mrs. H. M. Smith, Feb. 7, 1939.

***Ameiva undulata hartwegi* Smith**

Proc. Biol. Soc. Washington, vol. 53, p. 55, 1940.

HOLOTYPE: 103600, across Usumacinta River from Piedras Negras, Guatemala, in Chiapas, Mexico, H. M. Smith, June 18, 1939.

PARATYPES (11): 129274-83, 133854, Piedras Negras, Petén, Guatemala, H. M. Smith, May-June 1939.

***Ameiva undulata podarga* Smith and Laufe**

Univ. Kansas Sci. Bull., vol. 31, p. 40, figs. 1d, 2a, 1946.

PARATYPES (3): 106141-2, Hacienda La Clementina, 4 miles west of Forlón, Tamaulipas, Mexico, H. M. Smith, Nov. 19, 1938. 133859, Huichihuayán, San Luis Potosí, Mexico, H. M. Smith, Dec. 7, 1938.

***Ameiva undulata sinistra* Smith and Laufe**

Univ. Kansas Sci. Bull., vol. 31, p. 59, 1946.

PARATYPE: 134299 (formerly UIMNH 27343), Quesería, Colima, Mexico, H. M. Smith, June 18, 1935.

***Ameiva undulata stuarti* Smith**

Proc. Biol. Soc. Washington, vol. 53, p. 55, 1940.

HOLOTYPE: 103601, San Juanito, Palenque, Chiapas, Mexico, H. M. Smith, July 21, 1939.

PARATYPES (38): 133860, Río Chacama, Palenque, Chiapas, Mexico, H. M. Smith, Jan. 13, 1939. 133861-97, Palenque, Chiapas, Mexico, H. M. Smith, July 4-Aug. 2, 1939.

***Ameiva undulata thomasi* Smith and Laufe**

Univ. Kansas Sci. Bull., vol. 31, p. 47, pl. 1, fig. A, 1946.

PARATYPE: 137216, La Libertad, Chiapas, Mexico, H. D. Thomas, Sept. 1, 1937.

***Ameiva wetmorei* Stejneger**

Proc. Biol. Soc. Washington, vol. 26, p. 70, 1913.

=*Ameiva wetmorei wetmorei* Stejneger

HOLOTYPE: 49731, Guanica, above Río Loco, Puerto Rico, A. Wetmore, May 20, 1912.

***Ameiva wetmorei eleanorae* Grant and Roosevelt**

Journ. Dept. Agr. Puerto Rico, vol. 16, p. 48, 1932.

PARATYPES (2): 120785-6, islet south of Caja de Muertos Island, Puerto Rico, C. Grant, May 1932.

***Amiva gabbiana* Cope**

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8 (1875), p. 117, pl. 28, fig. 3, 1876.

=*Ameiva quadrilineata* (Hallowell)

COTYPES (3): 32614-6, Old Harbor, Costa Rica, W. M. Gabb.

***Amiva surinamenis tobaganus* Cope**

Proc. Amer. Philos. Soc., vol. 18, p. 276, 1879.

=*Ameiva tobagana* Cope

HOLOTYPE: 10113, Tobago Island, West Indies, F. A. Ober.

***Amphisbaena bakeri* Stejneger**

Report of the U.S. National Museum . . . for the year ending June 30, 1902, p. 681, fig. 133, 1904.

HOLOTYPE: 25541, Lares, Puerto Rico, A. B. Baker, Jan. 26, 1899.

PARATYPES (2): 25537, same data. 27458, Puerto Rico, C. E. Adams, 1900.

***Amphisbaena camura* Cope**

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 350, 1862.

COTYPES (2): 5860 (2), Paraguay, T. J. Page.

***Amphisbaena manni* Barbour**

Mem. Mus. Comp. Zool., vol. 44, p. 318, 1914.

PARATYPE: 67113 (formerly MCZ 8648), Cape Haitien, Haiti, W. M. Mann, 1913.

***Amphisbaena pericensis* Noble**

Ann. New York Acad. Sci., vol. 29, p. 141, 1921.

PARATYPE: 75970 (formerly MCZ 14786), Perico, Peru, G. K. Noble, September 1916.

***Amphisbaena townsendi* Stejneger**

Proc. U.S. Nat. Mus., vol. 41, p. 283, 1911.

HOLOTYPE: 47087, Piura, Peru, C. H. Townsend, Mar. 14, 1911.

***Anolis agassizii* Stejneger**

Bull. Mus. Comp. Zool., vol. 36, p. 161, pl., 1900.

=*Mariguana agassizii* (Stejneger)

HOLOTYPE: 22101, ♂, Malpelo Island, Colombia, C. H. Townsend, Mar. 5, 1891.

PARATYPES (2): 22103-4, ♀, ♀, same data. 22102 and 22105 were exchanged to MCZ in 1923.

***Anolis allisoni* Barbour**

Proc. New England Zool. Club, vol. 10, p. 58, 1928.

PARATYPE: 75859, Ruatán Island, Bay Islands, Honduras, T. Barbour, 1928.

***Anolis (Dracontura) alutaceus* Cope**

Proc. Acad. Nat. Sci. Philadelphia, vol. 13, p. 212, 1861.

=*Anolis alutaceus* Cope

COTYPES (3): 27485-7, Monte Verde, Cuba, C. Wright. 27488 was exchanged to MCZ in 1915.

***Anolis asper* Garman**

Bull. Essex Inst., vol. 19, p. 31, 1887.

=*Anolis ferreus* (Cope)

COTYPES (2): 39293-4 (formerly MCZ 6162), Marie Galante Island, West Indies, W. B. Richardson, 1886.

***Anolis aureolus* Cope**

Proc. Amer. Philos. Soc., vol. 22, p. 390, 1885.

COTYPES (5): 24850-1, Guatemala, H. Hague. 24854-6, Yucatán, Mexico, A. Schott. 24858 was exchanged to MCZ in 1915; 24857 to UIMNH, 1956.

***Anolis cepedii cinereus* Garman**

Bull. Essex Inst., vol. 19, p. 35, 1887.

=*Anolis roquet cinereus* Garman

COTYPE: 39298 (formerly MCZ 6182 part), St. George, Grenada Island, West Indies, S. Garman, 1879.

Anolis cepedii gentilis Garman

Bull. Essex Inst., vol. 19, p. 34, 1887.

= *Anolis roquet aeneus* Gray

COTYPE: 39295, Petite Martinique, Grenadines, West Indies, S. Garman, 1879.

Anolis compressicauda Smith and Kerster

Herpetologica, vol. 11, p. 193, figs. a-d, 1955.

PARATYPE: 138888 (formerly UIMNH 38032), El Ocote, Chiapas, Mexico, M. A. del Toro, Apr. 15, 1955.

Anolis (Gastrotropis) concolor Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 180, 1862.

= *Anolis concolor* Cope

COTYPES (3): 6055 (3), Nicaragua, C. Wright. One cotype was exchanged to MCZ in 1926.

Anolis conspersus Garman

Proc. Amer. Philos. Soc., vol. 24, p. 273, 1887.

COTYPE: 39292 (formerly MCZ 6021), Grand Cayman Island, West Indies, W. B. Richardson, 1887

Anolis conspersus lewisi Grant

The herpetology of the Cayman Islands, Bull. Inst. Jamaica Sci. Ser., No. 2 (1940), p. 23, 1941.

= *Anolis grahami lewisi* Grant

PARATYPES (3): 108027-9, east end, Grand Cayman Island, West Indies, C. B. Lewis, July 3, 1938.

Anolis cristatellus wileyi Grant

Journ. Dept. Agr. Puerto Rico, vol. 15, p. 220, 1931.

PARATYPES (2): 140281-2 (formerly MCZ untagged), Culebra Island, Puerto Rico, C. Grant, Apr. 15-26, 1931.

Anolis cupreus Hallowell

Proc. Acad. Nat. Sci. Philadelphia, vol. 12 (1860), p. 481, 1861.

COTYPES (11): 12211 (11), Nicaragua, C. Wright. Two cotypes were exchanged to MCZ in 1923; one cotype was exchanged to UIMNH in 1956.

Anolis (Dracontura) cyanopleurus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 13, p. 211, 1861.

= *Anolis cyanopleurus* Cope

COTYPES (3): 62068-70, Monte Verde, Cuba, C. Wright.

Anolis darlingtoni Cochran

Proc. New England Zool. Club, vol. 18, p. 1, 1939.

PARATYPES (2): 107556, ♂, Loma Vieja, Cordillera Central, Dominican Republic, 6,000 ft., P. J. Darlington, August 1938. 107557, Loma Rucilla and mountains north, Dominican Republic, P. J. Darlington, June 1938.

Anolis distichoides Rosén

Contributions to the fauna of the Bahamas, Lunds Univ. Årrskr. N. F., Afd. 2, Bd. 7, Nr. 5, p. 29, figs. 12a-d, 1911.

PARATYPES (2): 103709-10, Stanniard Creek, Andros Island, Bahama Islands, N. Rosén, January 1909.

Anolis distichus floridanus Smith and McCauley

Proc. Biol. Soc. Washington, vol. 61, p. 160, pl. 7, 1948.

HOLOTYPE: 127114, ♂, Brickell Park, Miami, Dade County, Florida, R. H. McCauley, Jr., Nov. 6, 1946.

Anolis dominicensis caudalis Cochran

Proc. Biol. Soc. Washington, vol. 45, p. 185, 1932.

=*Anolis distichus caudalis* Cochran

HOLOTYPE: 76801, En Café, Gonave Island, Haiti, A. J. Poole and W. M. Perrygo, March 1929.

PARATYPES (9): 76799-800, same data. 77080-2, Point-à-Raquette, Gonave Island, Haiti, W. J. Eyerdam, 1927. 80386, 80388-9, Anse à Galets, Gonave Island, Haiti, L. H. Parish and W. M. Perrygo, Mar. 23, 1930. 80390, Petite Gonave Island, Haiti, L. H. Parish and W. M. Perrygo, Mar. 19, 1930. 80387 was exchanged to MZUM in 1943.

Anolis dominicensis juliae Cochran

Occ. Pap. Boston Soc. Nat. Hist., vol. 8, p. 169, 1934.

=*Anolis distichus juliae* Cochran

PARATYPE: 95122, Isle Vache, Haiti, Utowana Expedition, Apr. 12, 1934.

Anolis dominicensis wetmorei Cochran

Proc. Biol. Soc. Washington, vol. 44, p. 89, 1931.

=*Anolis distichus wetmorei* Cochran

HOLOTYPE: 83381, Beata Island, Dominican Republic, A. Wetmore and F. C. Lincoln, May 13, 1931.

Anolis doris Barbour

Proc. Biol. Soc. Washington, vol. 33, p. 101, 1925.

=*Anolis cybotes doris* Barbour

PARATYPE: 69384 (formerly MCZ 13741), Gonave Island, Haiti, G. M. Allen, August 1919.

Anolis evermanni Stejneger

Report of the U.S. National Museum . . . for the year ending June 30, 1902, p. 647, figs. 102-104, 1904.

HOLOTYPE: 26855, ♂, Catalina Plantation, Puerto Rico, 890 ft., L. Stejneger, Feb. 21, 1900.

PARATYPES (29): 25721-2, 26085, 26906-7, El Yunque, Puerto Rico, U.S. Fishhawk, Feb. 19, 1899. 26853-4, 26861-3, 26866-9, Catalina Planta-

tion, Puerto Rico, L. Stejneger and C. W. Richmond, February–March 1900. 26899, between Catalina Plantation and Mameyes, Puerto Rico, L. Stejneger, Mar. 5, 1900. 27154–6, 58479 (formerly 27213), near Utuado, Puerto Rico, L. Stejneger, Apr. 7, 1900. 27237–8, 27256, 27259, 27273–4, 27278, 58480–1 (formerly 27239–40), Adjuntas, Puerto Rico, L. Stejneger, Apr. 13, 1900. 27279, between Adjuntas and Ponce, C. W. Richmond, Apr. 15, 1900. 25454 was exchanged to MCZ in 1912; 27153 to R, 1953.

Anolis fairchildi Barbour and Shreve

Proc. Boston Soc. Nat. Hist., vol. 40, p. 357, 1935.

HOLOTYPE: 81527, ♂, Cay Sal, Bahama Islands, P. Bartsch, June 17, 1930.

PARATYPES (3): 81525–6, same data. 81559, Cotton Cay, Cay Sal Group, Bahama Islands, P. Bartsch, June 23, 1930. 81558 was exchanged to MCZ in 1934.

Anolis greyi Barbour

Mem. Mus. Comp. Zool., vol. 44, p. 287, 1914.

COTYPES (3): 39516–8, Puerto Principe, Camagüey, Cuba, T. Barbour, March 1909.

Anolis griseus Garman

Bull. Essex Inst., vol. 19, p. 36, 1887.

= *Anolis richardii griseus* Garman

COTYPE: 120754 (formerly MCZ 6164), Kingston, St. Vincent, West Indies, S. Garman, Feb. 18, 1879.

Anolis hendersoni Cochran

Journ. Washington Acad. Sci., vol. 13, p. 225, 1923.

HOLOTYPE: 59210, Pétionville, Haiti, J. B. Henderson and P. Bartsch, Apr. 1, 1917.

Anolis leucophaeus mariguanae Cochran

Journ. Washington Acad. Sci., vol. 21, p. 40, 1931.

HOLOTYPE: 81346, Mariguana Cay, Bahama Islands, P. Bartsch, July 18, 1930.

PARATYPES (29): 81344–5, 81347–50, same data. 81351–5, 81357, 81359–72, Betsy Bay, Mariguana Cay, Bahama Islands, P. Bartsch, July 19–20, 1930. 81373–5, Booby Island, east of Mariguana Cay, Bahama Islands, P. Bartsch, July 21, 1930. 81356 and 81358 were exchanged to MCZ in 1934.

Anolis lineatopus coxi Grant

Jamaica To-day, p. 155, 1940.

PARATYPES (8): 108294–301, Portland Point, Jamaica, C. Grant, April 1937–January 1938.

Anolis lineatopus lynni Grant

Jamaica To-day, p. 155, 1940.

HOLOTYPE: 107902, near Chester Vale, St. Andrew, Jamaica, W. G. Lynn, 1936.

PARATYPE: 107903, same data.

Anolis lividus Garman

Bull. Essex Inst., vol. 19, p. 43, 1887.

=*Anolis bimaculatus lividus* Garman

COTYPES (2): 39303-4, Plymouth, Montserrat Island, West Indies, S. Garman, 1879.

Anolis luciae Garman

Bull. Essex Inst., vol. 19, p. 44, 1887.

=*Anolis trinitatis luciae* Garman

COTYPES (2): 39296-7, Punta Castries, Santa Lucia Island, West Indies, S. Garman, 1879.

Anolis matudai Smith

Herpetologica, vol. 12, p. 1, 1956.

PARATYPE: 137289, Mount Ovando, Chiapas, Mexico, Dr. and Mrs. H. M. Smith, Apr. 16, 1940. This specimen is not from Finca La Esperanza, as stated in original description (fide H. M. Smith).

Anolis mestrei Barbour and Ramsden

Proc. Biol. Soc. Washington, vol. 29, p. 19, 1916.

PARATYPES (4): 26731-3, San Diego de los Baños, Cuba, W. Palmer and J. H. Riley, Apr. 12, 1900. 27344, El Guamá, near city of Pinar del Río, Cuba, W. Palmer and J. H. Riley, Mar. 24, 1900.

Anolis microtus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 23, p. 214, pl. 24, fig. 2, 1871.

HOLOTYPE: 31282, San José, Costa Rica, C. H. Van Patten.

Anolis milleri Smith

in Smith and Taylor, U.S. Nat. Mus. Bull. 199, p. 64, 1950.

HOLOTYPE: 120957, Quetzaltepec, Oaxaca, Mexico, W. S. Miller, Apr. 17, 1944.

Anolis monensis Stejneger

Report of the U.S. National Museum . . . for the year ending June 30, 1902, p. 646, figs. 98-101, 1904.

HOLOTYPE: 29387, Mona Island, Puerto Rico, B. S. Bowdish, Aug. 19, 1901.

PARATYPES (12): 29376-86, 29388, Mona Island, Puerto Rico, B. S. Bowdish, Aug. 8-12, 1901. 29375 was exchanged to MCZ in 1912.

Anolis nannodes Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 16, p. 173, 1864.

=*Anolis laevis* (Wiegmann)

COTYPES (3): 12206 (3), Arriba, Costa Rica, C. N. Riotte.

Anolis nelsoni Barbour

Mem. Mus. Comp. Zool., vol. 44, p. 287, 1914.

PARATYPE: 75957, Swan Island, Honduras, G. Nelson, March 1912.

Anolis oxylophus Cope

Journ. Acad. Sci. Philadelphia, ser. 2, vol. 8 (1875), p. 123, pl. 24, fig. 4, pl. 28, fig. 5, 1876.

=*Anolis lionotus* Cope

COTYPES (2): 30556-7, Costa Rica, W. M. Gabb.

Anolis pachypus Cope

Journ. Acad. Sci. Philadelphia, ser. 2, vol. 8 (1875), p. 120, pl. 24, fig. 3, 1876.

HOLOTYPE: 30683, Pico Blanco, Limón, Costa Rica, W. M. Gabb.

Anolis pinchoti Cochran

Journ. Washington Acad. Sci., vol. 21, p. 354, 1931.

=*Anolis concolor* Cope

HOLOTYPE: 76945, ♂, Old Providence Island, Colombia, A. K. Fisher, Apr. 23, 1929.

PARATYPE: 76946, ♀, same data.

Anolis poncensis Stejneger

Report of the U.S. National Museum . . . for the year ending June 30, 1902, p. 665, figs. 117-120, 1904.

HOLOTYPE: 27294, ♂, 3 miles east of Ponce, Puerto Rico, C. W. Richmond, Apr. 16, 1900.

PARATYPES (5): 27289-93, 3 miles east of Ponce, Puerto Rico, L. Stejneger, Apr. 16, 1900.

Anolis purpurescens Cope

Sci. Bull., Acad. Nat. Sci. Philadelphia, No. 1, p. 7, 1899.

HOLOTYPE: 4321, Truandó River New Granada (Colombia), A. Schott.

Anolis quaggulus Cope

Proc. Amer. Philos. Soc., vol. 22, p. 391, 1885.

=*Anolis humilis humilis* Peters

HOLOTYPE: 24979, San Juan River, Nicaragua, R. W. Kennicott.

Anolis ruthveni Stuart

Occ. Pap. Mus. Zool. Univ. Michigan, No. 310, p. 1, 1935.

=*Anolis humilis uniformis* Cope

PARATYPES (6): 71866, 71869, 71892, 71908, 71914, 71915, El Petén, Guatemala, H. Malleis.

Anolis sabanus Garman

Bull. Essex Inst., vol. 19, p. 39, 1887.

=*Anolis bimaculatus sabanus* Garman

COTYPE: 39299, Saba Island, West Indies, F. Lagois, 1884.

Anolis sagrei mayensis Smith and Burger

Anal. Inst. Biol., Mexico, vol. 20 (1949), p. 407, 1950.

PARATYPE: 137211, Encarnación, Campeche, Mexico, H. M. Smith, Oct. 1-10, 1936.

Anolis stejnegeri Barbour

Copeia, No. 3, p. 88, 1931.

=*Anolis sagrei stejnegeri* Barbour

PARATYPE: 6002, Cape Punta (Florida).

Anolis subocularis Davis

Herpetologica, vol. 10, p. 3, 1954.

PARATYPES (93): 133725-816, Tierra Colorada, Guerrero, Mexico, H. M. Smith, Feb. 12-14, 1939. 133817, Agua del Obispo, Guerrero, Mexico, H. M. Smith, Aug. 31, 1939.

Anolis taylori Smith and Spieler

Copeia, No. 3, p. 165, 1945.

HOLOTYPE: 132358, ♂, 1 mile north of Acapulco, Guerrero, Mexico, E. H. Taylor, Sept. 3, 1939.

PARATYPES (5): 132359-63, 1 mile north of Acapulco, Guerrero, Mexico, E. H. Taylor and Dr. and Mrs. H. M. Smith, Sept. 3, 1939. 132364 was exchanged to MCZ in 1958.

Anolis townsendi Stejneger

Bull. Mus. Comp. Zool., vol. 36, p. 163, 1900.

HOLOTYPE: 22106, ♂, Cocos Island, Costa Rica, C. H. Townsend, Feb. 28, 1891.

PARATYPE: 22107, ♀, same data.

Anolis trinitatis extremus Garman

Bull. Essex Inst., vol. 19, p. 35, 1887.

=*Anolis roquet cinereus* Garman

COTYPES (2): 39290-1 (formerly, MCZ 2743), Bridgetown, Barbados Island, West Indies, L. Agassiz, 1872.

Anolis tropidonotus spilorhipis del Toro and Smith

Herpetologica, vol. 12, p. 9, 1956.

PARATYPE: 138889 (formerly UIMNH 37974), Villa Allende, Cerro Ombligo, Chiapas, Mexico, M. A. del Toro, May 5, 1954.

Anolis trossulus Garman

Bull. Essex Inst., vol. 19, p. 38, 1887.

=*Anolis richardii richardii* Duméril and Bibron

COTYPE: 39289, St. George, Grenada Island, West Indies, S. Garman, 1879.

Anolis uniformis Cope

Proc. Amer. Philos. Soc., vol. 22, p. 392, 1885.

=*Anolis humilis uniformis* Cope

COTYPES (23): 6774 (6), 24734-48, 24750, Guatemala, H. Hague. 24859, Yucatán, Mexico, A. Schott. 24749 was exchanged to MCZ in 1915.

Anolis vincentii Garman

Bull. Essex Inst., vol. 19, p. 46, 1887.

=*Anolis trinitatis vincentii* Garman

COTYPES (2): 39301-2, Kingston, St. Vincent, West Indies, S. Garman, 1879.

Anolis virgatus Garman

Bull. Essex Inst., vol. 19, p. 41, 1887.

=*Anolis gingivinus* Cope

COTYPE: 39300, St. Bart's Island, West Indies, F. Lagois, 1884.

Anota calidiarum Cope

Amer. Nat., vol. 30, p. 833, 1896.

=*Phrynosoma platyrhinos calidiarum* Cope

HOLOTYPE: 18444, ♀, Death Valley, California, Death Valley Expedition, 1891.

Aristelliger expectatus Cochran

Proc. Biol. Soc. Washington, vol. 46, p. 33, 1933.

=*Aristelligella expectata* (Cochran)

HOLOTYPE: 75908, ♂, Jacmel, Haiti, J. S. C. Boswell, 1928.

Aristelliger irregularis Cope

Proc. Amer. Philos. Soc., vol. 22, p. 387, 1885.

=*Aristelliger georgeensis* (Bocourt)

HOLOTYPE: 13903, Cozumel Island, off Yucatán, Mexico, U.S. Fish Commission *Albatross*, Jan. 23-29, 1885.

Audantia armouri Cochran

Occ. Pap. Boston Soc. Nat. Hist., vol. 8, p. 171, 1934.

PARATYPES (5): 72593-4, head of Rivière Chotard, Massif de La Selle, Haiti, 6,500 ft., B. H. Swales and A. Wetmore, Apr. 12, 1927. 72598-9, Morne Cabaio, Massif de La Selle, Haiti, 7,500 ft., B. H. Swales and A. Wetmore, Apr. 10, 1927. 85008, Morne La Selle, Haiti, 7,000 ft., S. S. Cook, 1932.

Audantia shrevei Cochran

Proc. New England Zool. Club, vol. 18, p. 2, 1939.

PARATYPES (5): 107558, ♂, Valle Nuevo, Cordillera Central, Dominican Republic, P. J. Darlington, August, 1938. 107559-61, Loma Pelona, Dominican Republic, P. J. Darlington, June 1938. 107562, Loma Rucilla, Dominican Republic, P. J. Darlington, June 1938.

Aulacoplax leptosoma Brown and Fehlmann

Occ. Pap. Nat. Hist. Mus. Stanford Univ., No. 6, p. 1, figs. 1-5, 1958.

PARATYPES (2): 140413-4 (formerly SU 20845-6), Koror Island, Palau Islands (West Caroline Islands, West Pacific), H. A. Fehlmann and Y. Sumang.

Bachia intermedia Noble

Ann. New York Acad. Sci., vol. 29, p. 142, 1921.

PARATYPE: 59927, Perico, Chinchipe River, northeastern Peru, G. K. Noble, 1916.

Bachia parkeri Ruthven

Proc. Boston Soc. Nat. Hist., vol. 38, p. 103, figs. 4-6, 1925.

PARATYPE: 65437, Río Tiquié, Colombia, G. McCreagh.

Barisia gadovii levigata Tihen

Univ. Kansas Sci. Bull., vol. 33, p. 231, pl. 2, figs. 3-5, 1949.

HOLOTYPE: 47212, ♂, Valley of Oaxaca, Oaxaca, Mexico, E. W. Nelson and E. A. Goldman, Sept. 14, 1894.

PARATYPE: 47855, ♂, mountains west of Oaxaca City, Oaxaca, Mexico, 9,000 ft., E. W. Nelson and E. A. Goldman, Sept. 14, 1894.

Barissia antauges Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 18, p. 132, 1866.

=*Barisia antauges* Cope

HOLOTYPE: 30221, Orizaba, Veracruz, Mexico, F. Sumichrast.

Barissia levicollis Stejneger

Proc. U.S. Nat. Mus., vol. 13, p. 184, 1890.

=*Barisia levicollis* Stejneger

HOLOTYPE: 9362, Mexican Boundary (probably Chihuahua), Mexico.

Basiliscus barbouri Ruthven

Proc. Biol. Soc. Washington, vol. 27, p. 9, pl. 1, figs. 1-2, 1914.

PARATYPE: 50671, San Lorenzo, Tamocal River, Santa Marta Mountains, Colombia, 400 ft., M. A. Carriker, Jr., July 7, 1913.

Basiliscus guttulatus Cope

Journ. Acad. Nat. Sci., Philadelphia, ser. 2, vol. 8 (1875), p. 156, 1876.

=*Basiliscus basiliscus* (Linnaeus)

HOLOTYPE: 25164, Buhio Soldado, Panama, J. F. Bransford. This specimen is now badly macerated.

Basiliscus (Cristasaura) nuchalis Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 181, 1862.

=*Basiliscus vittatus* Wiegmann

HOLOTYPE: 5845, near Greytown, Nicaragua, H. C. Caldwell.

Basiliscus plumifrons Cope

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8 (1875), p. 125, pl. 25, fig. 1, 1876.

COTYPES (5): 32622-6, Sipurio, Costa Rica, W. M. Gabb. 35627 was exchanged to UIMNH in 1956.

Cachryx defensor Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 18, p. 124, 1866.

=*Enyaliosaurus defensor* (Cope)

COTYPES (3): 12282 (3), Yucatán (restricted to Chichén Itzá), Mexico, A. Schott.

Cadea palirostrata Dickerson

Bull. Amer. Mus. Nat. Hist., vol. 35, p. 659, 1916.

PARATYPE: 54379 (formerly AMNH 2718), San Pedro, Isle of Pines, Cuba, C. S. Mead, winter of 1911-1912.

Callisaurus carmenensis Dickerson

Bull. Amer. Mus. Nat. Hist., vol. 41, p. 465, 1919.

=*Callisaurus draconoides carmenensis* Dickerson

HOLOTYPE: 64253, ♂, Carmen Island, Gulf of California, Baja California, Mexico, C. H. Townsend, Apr. 2, 1911.

Callisaurus crinitus Cope

Amer. Nat., vol. 30, p. 1049, 1896.

=*Callisaurus draconoides crinitus* Cope

HOLOTYPE: 14895, Ballenas Bay, Baja California, Mexico, U.S. Fish Commission *Albatross*, May 3, 1888.

Callisaurus inusitatus Dickerson

Bull. Amer. Mus. Nat. Hist., vol. 41, p. 465, 1919.

=*Callisaurus draconoides inusitatus* Dickerson

HOLOTYPE: 64256, ♂, Tiburón Island, Gulf of California, Sonora, Mexico, C. H. Townsend, Apr. 13, 1911.

Callisaurus plasticus Dickerson

Bull. Amer. Mus. Nat. Hist., vol. 41, p. 466, 1919.

=*Callisaurus draconoides carmenensis* Dickerson

HOLOTYPE: 64255, ♂, Agua Verde Bay, Baja California, Mexico, C. H. Townsend, Apr. 1, 1911.

Callisaurus rhodostictus Cope

Amer. Nat., vol. 30, p. 1049, 1896.

=*Callisaurus draconoides rhodostictus* Cope

COTYPES (8): 21504, El Rosario, Baja California, Mexico, A. W. Anthony, Apr. 24, 1894. 21505-11, San Fernando, Baja California, Mexico, A. W. Anthony, April-June, 1894. 21511 is the lectotype.

Callisaurus splendidus Dickerson

Bull. Amer. Mus. Nat. Hist., vol. 41, p. 467, 1919.

=*Callisaurus draconoides splendidus* Dickerson

HOLOTYPE: 64254, ♂, Angel de la Guarda Island, Gulf of California, Baja California, Mexico, C. H. Townsend, Apr. 10, 1911.

Callisaurus ventralis myurus Richardson

Proc. U.S. Nat. Mus., vol. 48, p. 408, 1915.

=*Callisaurus draconoides myurus* Richardson

HOLOTYPE: 51786, The Willows, Indian Agency, Pyramid Lake, Washoe County, Nevada, J. O. Snyder and C. H. Richardson, June 2, 1911.

Celestus badius Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 20, p. 126, 1868.

COTYPES (2): 25817-8, Navassa Island, West Indies, W. J. Rasin.

Celestus cruscus molesworthi Grant

Jamaica To-day, p. 157, 1940.

PARATYPES (2): 108158-9, 1 mile south of Buff Bay, Portland, Jamaica, C. Grant, May 2, 1939.

Celestus curtissi Grant

Copeia, No. 1, p. 68, 1 pl., 1951.

HOLOTYPE: 117337, Trou Forban, Haiti, A. Curtiss, Apr. 19, 1943.

PARATYPES (6): 117265, Trou Caïman, Haiti, A. Curtiss, Feb. 16, 1943. 117266-8, 117338, 129399, Trou Forban, Haiti, A. Curtiss, August 1942-April 1943.

Celestus darlingtoni Cochran

Proc. New England Zool. Club, vol. 18, p. 2, 1939.

PARATYPES (2): 107563-4, Valle Nuevo, Cordillera Central, Dominican Republic, 6,000-8,000 ft., P. J. Darlington, August 1938.

Celestus rozellae Smith

Proc. U.S. Nat. Mus., vol. 92, p. 372, 1942.

HOLOTYPE: 113526, Palenque, Chiapas, Mexico, R. B. Smith, July 7, 1939.

PARATYPES (2): 62992, Central America (from fruit ship at New Orleans, La.), H. E. Hubert. 113527, Piedras Negras, Petén, Guatemala, H. M. Smith, June 2, 1939.

Celestus rugosus Cope

Proc. Amer. Philos. Soc., vol. 18, p. 272, 1879.

HOLOTYPE: 10260, Puerto Plata, Dominican Republic, C. A. Fraser, 1878.

Celestus weinlandii Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 20, p. 125, 1868.

HOLOTYPE: 12145, within 25 miles of Port-au-Prince (given erroneously as Gonave Island in original description), Haiti, A. C. Younglove.

Chalcidolepis metallicus Cope

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8 (1875), p. 116, pl. 24, fig. 5, pl. 28, fig. 2, 1876.

=*Anadia metallica metallica* (Cope)

HOLOTYPE: 30568, Aguacate Mountains, Costa Rica, W. M. Gabb.

Chamaeleo abbotti Stejneger

Proc. U.S. Nat. Mus., vol. 14, p. 353, 1 fig., 1891.

=*Chamaeleo taitensis* Steindachner

HOLOTYPE: 16744, ♂, Kilimanjaro, East Africa (Tanganyika), 4,500 ft., W. L. Abbott, 1889.

PARATYPES (2): 16743, 16747, same data.

Chamaeleon ituriensis Schmidt

Bull. Amer. Mus. Nat. Hist., vol. 39, p. 589, text fig. 25, pl. 31, fig. 2, pl. 32, figs. 1-5, 1919.

=*Chamaeleo johnstonii* Boulenger

PARATYPE: 62015 (formerly AMNH 11461), Gamangui, Belgian Congo, H. Lang and J. P. Chapin Expedition, February 1910.

Chamaelinorops wetmorei Cochran

Proc. Biol. Soc. Washington, vol. 41, p. 45, 1928.

HOLOTYPE: 72630, Fonds-des-Nègres, Haiti, A. Wetmore, Apr. 4, 1927.

Cnemidophorus angusticeps Cope

Proc. Amer. Philos. Soc., vol. 17 (1878), p. 95, 1877.

=*Cnemidophorus sackii angusticeps* Cope

COTYPES (2): 24876-7, Yucatán, Mexico, A. Schott. 12284 (2) and 24879, same data, are probably also cotypes. 24878 was exchanged to MCZ in 1944.

Cnemidophorus bartolomas Dickerson

Bull. Amer. Mus. Nat. Hist., vol. 41, p. 476, 1919.

=*Cnemidophorus tigris multiscutatus* Cope

HOLOTYPE: 64448, San Bartolomé Bay, Baja California, Mexico, C. H. Townsend, March 11, 1913.

Cnemidophorus burti Taylor

Univ. Kansas Sci. Bull., vol. 24 (1936), p. 485, pl. 43, fig. 2, 1938.

PARATYPE: 138898 (formerly UIMNH 35973), 10 miles northeast of Guaymas, Sonora, Mexico, H. M. Smith and D. H. Dunkle, 1934.

Cnemidophorus celeripes Dickerson

Bull. Amer. Mus. Nat. Hist., vol. 41, p. 472, 1919.

=*Cnemidophorus tigris celeripes* Dickerson

HOLOTYPE: 64444, San José Island, Gulf of California, Baja California, Mexico, C. H. Townsend, 1911.

Cnemidophorus communis Cope

Proc. Amer. Philos. Soc., vol. 17 (1878), p. 95, 1877.

=*Cnemidophorus communis communis* Cope

COTYPES (114): 6779, Guatemala, H. Hague. 24941-60, Guadalajara?, Jalisco, Mexico, J. J. Major. 30240, Córdoba, Veracruz, Mexico, F. Sumichrast. 31398-420, 31506-27, 31530-62, 31604-17, Colima, Colima, Mexico, J. Xantus, March-October 1863. When these specimens were returned by Cope's estate after his death, they were marked as follows: Var. I, 31506-27, 31530-62, 31604-17; var. II, 24941-60, 30240. 31535, 31545, 31549, and 31561 were considered to be *Cnemidophorus deppii lineatissimus* Cope, fide R. G. Zweifel.

Cnemidophorus costatus Cope

Proc. Amer. Philos. Soc., vol. 17 (1878), p. 95, 1877.

=*Cnemidophorus sackii sackii* Wiegmann

HOLOTYPE: 31359, Mexico (restricted to Cuernavaca, Morelos), A. Boucard.

Cnemidophorus decemlineatus Hallowell

Proc. Acad. Nat. Sci. Philadelphia, vol. 12, p. 482, 1861.

=*Cnemidophorus deppii deppii* Wiegmann

COTYPES (11): 6058 (11), Nicaragua, C. Wright. One cotype was exchanged to MCZ in 1944.

Cnemidophorus disparilis Dickerson

Bull. Amer. Mus. Nat. Hist., vol. 41, p. 473, 1919.

=*Cnemidophorus tigris aethiops* Cope

HOLOTYPE: 64445, Tiburón Island, Gulf of California, Sonora, Mexico, C. H. Townsend, Apr. 12, 1911.

Cnemidophorus estebanensis Dickerson

Bull. Amer. Mus. Nat. Hist., vol. 41, p. 474, 1919.

=*Cnemidophorus tigris estebanensis* Dickerson

HOLOTYPE: 64446, San Estéban Island, Gulf of California, Sonora, Mexico, C. H. Townsend, Apr. 13, 1911.

Cnemidophorus gadovi Burger

Chicago Acad. Sci., Nat. Hist. Misc., No. 65, p. 2, 1950.

=*Cnemidophorus tigris aethiops* Cope

HOLOTYPE: 40042, Hermosillo, Sonora, Mexico, J. N. Rose, Mar. 5, 1910.

Cnemidophorus gigas Davis and Smith

Herpetologica, vol. 8, p. 98, 1952.

PARATYPE: 134300, Cuernavaca, Morelos, Mexico, H. M. Smith, Jan. 24, 1939.

Cnemidophorus gracilis Baird and Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 128, 1852.

=*Cnemidophorus tigris gracilis* Baird and Girard

HOLOTYPE: 3034, "Colorado Desert, California" (restricted to Yuma, Yuma County, Arizona), J. L. LeConte.

Cnemidophorus grahamii Baird and Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 128, 1852.

=*Cnemidophorus tessellatus* (Say)

COTYPES (2): 3046 (2), between San Antonio and El Paso del Norte (El Paso in the catalogue, but restricted to Fort Davis, Jeff Davis County, Texas), J. H. Clark.

Cnemidophorus gularis Baird and Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 128, 1852.

=*Cnemidophorus sackii gularis* Baird and Girard

COTYPES (13): 2989 (8), Indianola (Calhoun County), Texas, J. D. Graham. 3022 (5), Rio San Pedro of the Rio Grande, Texas (restricted to mouth of Devils River, Val Verde County), J. D. Graham. 3022a is the lectotype.

Cnemidophorus gularis gularis obsoletus Cope

Trans. Amer. Philos. Soc., vol. 17, p. 45, 1892.

=*Cnemidophorus sackii scalaris* Cope

COTYPES (3): 14231b, 14308d-e, Chihuahua, Chihuahua, Mexico, E. Wilkinson.

Cnemidophorus gularis gularis verus Cope

Trans. Amer. Philos. Soc., vol. 17, p. 45, 1892.

=*Cnemidophorus sackii scalaris* Cope

COTYPES (10): 14231a, 14241c-e, 14305b-d, 15308a-c, Chihuahua, Chihuahua, Mexico, E. Wilkinson.

Cnemidophorus gularis scalaris Cope

Trans. Amer. Philos. Soc., vol. 17, p. 47, pl. 10, fig. 10, 1892.

=*Cnemidophorus sackii scalaris* Cope

COTYPES (11): 8319 (4), between Chihuahua and Mexico, Mexico, J. Potts. 14302 (7), Chihuahua, Chihuahua, Mexico, E. Wilkinson. 14302a is the lectotype, and the type locality is restricted to Chihuahua, Chihuahua. One cotype from 8319 was exchanged to UIMNH in 1956.

Cnemidophorus gularis semifasciatus Cope

Trans. Amer. Philos. Soc., vol. 17, p. 49, pl. 11, fig. 12, 1892.

=*Cnemidophorus sackii semifasciatus* Cope

COTYPES (2): 9248 (formerly 3041), Agua Nueva, Coahuila, Mexico, D. N. Couch. 3033, Patos, Coahuila, Mexico, D. N. Couch.

Cnemidophorus gularis sericeus Cope

Trans. Amer. Philos. Soc., vol. 17, p. 48, pl. 11, fig. 11, 1892.

=*Cnemidophorus sackii gularis* Baird and Girard

HOLOTYPE: 15650, San Diego, Duval County, Texas, W. Taylor.

Cnemidophorus hyperythrus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 15, p. 103, 1863.

=*Cnemidophorus hyperythrus hyperythrus* Cope

COTYPES (24): 5299 (13), 69495-6, Cape San Lucas, Baja California, Mexico, J. Xantus, May-September, 1859. 69497, Rancho Viejo, 10 leagues north of San José, Baja California, Mexico, J. Xantus, December 1859. 69498, Santiago Sierras, between San José and La Paz, Baja California, Mexico, J. Xantus, November 1859. 69499-69505, Miraflores, 10 leagues north of San José, Baja California, Mexico, J. Xantus, December 1859.

Cnemidophorus immutabilis Cope

Proc. Amer. Philos. Soc., vol. 17 (1878), p. 93, 1877.

=*Cnemidophorus guttatus immutabilis* Cope

PROBABLE COTYPE: 30141, West Tehuantepec, Oaxaca, Mexico, F. Sumichrast.

Cnemidophorus inornatus Baird

Proc. Acad. Nat. Sci. Philadelphia, vol. 10 (1858), p. 255, 1859.

COTYPES (2): 3032 (2), Pesquería Grande, Nuevo León, Mexico, D. N. Couch.

Cnemidophorus labialis Stejneger

Proc. U.S. Nat. Mus., vol. 12, p. 643, 1890.

HOLOTYPE: 15596, Cedros Island (given as "Cerros Island" in the original description, but probably from San Quintín Bay), Baja California, Mexico, L. Belding.

PARATYPES (3): 15597-8, 15600, same data. 15599 was exchanged to MCZ.

Cnemidophorus lativittis Cope

Proc. Amer. Philos. Soc., vol. 17 (1878), p. 94, 1877.
 =*Cnemidophorus deppii deppii* Wiegmann
 HOLOTYPE: 30227, Juchitán, Tehuantepec, Oaxaca, Mexico, F. Sumichrast.

Cnemidophorus lineatissimus Cope

Proc. Amer. Philos. Soc., vol. 17 (1878), p. 94, 1877.
 =*Cnemidophorus deppii lineatissimus* Cope
 COTYPES (20): 24937-40, Guadalajara?, Jalisco, Mexico, J.J. Major.
 32299-314, Colima, Colima, Mexico, J. Xantus, July 1863.

Cnemidophorus marmoratus Baird and Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 128, 1852.
 =*Cnemidophorus tigris marmoratus* Baird and Girard
 COTYPES (2): 3024 (2), between San Antonio and El Paso, Texas (restricted to El Paso), J. H. Clark. 3024a is the lectotype.

Cnemidophorus martyr Stejneger

Proc. U.S. Nat. Mus., vol. 14, p. 407, 1891.
 =*Cnemidophorus tigris martyr* Stejneger
 HOLOTYPE: 15620, San Pedro Mártir Island, Gulf of California, Sonora, Mexico, E. Palmer.
 PARATYPE: 15621, same data.

Cnemidophorus maximus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 15, p. 104, 1863.
 HOLOTYPE: 5297, Cape San Lucas, Baja California, Mexico, J. Xantus, September 1859.

Cnemidophorus melanostethus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 15, p. 104, 1863.
 =*Cnemidophorus tigris gracilis* Baird and Girard
 COTYPES (2): 3067a (2, given incorrectly in original description as 3064), "Colorado River, California" (restricted to Yuma, Yuma County, Arizona), H. B. Möllhausen.

Cnemidophorus microlepidopus Cope

Proc. Amer. Philos. Soc., vol. 17 (1878), p. 93, 1877.
 =*Cnemidophorus guttatus immutabilis* Cope
 HOLOTYPE: 30187, west Tehuantepec, Oaxaca, Mexico (restricted to city of Tehuantepec), F. Sumichrast.

Cnemidophorus neomexicanus Lowe and Zweifel

Bull. Chicago Acad. Sci., vol. 9, p. 230, fig. 1c, pl. 1A, 1952.
 PARATYPES (2): 132467-8, McDonald Ranch Headquarters, Socorro County, New Mexico, C. H. Lowe, Jr., Aug. 18, Aug. 3, 1947.

Cnemidophorus octolineatus Baird

Proc. Acad. Nat. Sci. Philadelphia, vol. 10 (1858), p. 255, 1859.

=*Cnemidophorus inornatus* Baird

HOLOTYPE: 3009, Pesquería Grande, Nuevo León, Mexico, D. N. Couch.

Cnemidophorus perplexus Baird and Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 128, 1852.

LECTOTYPE: 3060, probably from western Texas. The original description states: "Collected . . . by Dr. William Gambel on his last journey to California."

Cnemidophorus praesignis Baird and Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 129, 1852.

=*Ameiva praesignis* (Baird and Girard)

COTYPE: 5519, Chagres, Panama, C. B. Adams. One cotype was exchanged to MZUM.

Cnemidophorus punctilinealis Dickerson

Bull. Amer. Mus. Nat. Hist., vol. 41, p. 475, 1919.

=*Cnemidophorus tigris aethiops* Cope

HOLOTYPE: 64447, ♂, Tiburón Island, Gulf of California, Sonora, Mexico, C. H. Townsend, Apr. 12, 1911.

PARATYPES (5): 64456-60, same data.

Cnemidophorus quadrilineatus Hallowell

Proc. Acad. Nat. Sci. Philadelphia, vol. 12 (1860), p. 483, 1861.

=*Ameiva quadrilineata* (Hallowell)

COTYPES (2): 6053 (2), Greytown, Nicaragua, C. Wright.

Cnemidophorus sacki huico Zweifel

Bull. Amer. Mus. Nat. Hist., vol. 117, p. 85, pl. 43, pl. 44, figs. 1, 4, 1959.

=*Cnemidophorus sackii huico* Zweifel

PARATYPE: 47671, Acaponeta, Tepic, Mexico, E. W. Nelson and E. A. Goldman, June 23, 1897.

Cnemidophorus sacki stictogrammus Burger

Chicago Acad. Sci., Nat. Hist. Misc. No. 65, p. 5, 1950.

=*Cnemidophorus sackii stictogrammus* Burger

HOLOTYPE: 132456, Yank Springs, 6 miles southeast of Ruby, Santa Cruz County, Arizona, M. M. Hensley and W. L. Burger, Aug. 17, 1948.

Cnemidophorus septemvittatus Cope

Trans. Amer. Philos. Soc., ser. 2, vol. 17, p. 40, 1892.

=*Cnemidophorus sackii semifasciatus* Cope

HOLOTYPE: 42141, "El Dorado County California" (stricted to Marfa, Presidio County, Texas).

Cnemidophorus sexlineatus oligoporus Hoffman

Journ. Washington Acad. Sci., vol. 47, p. 351, 1957. (Not of Smith, 1939.)

=*Cnemidophorus sexlineatus pauciporus* Hoffman

HOLOTYPE: 135232, ♂, Griffith, Alleghany County, Virginia, R. L. Hoffman, May 27, 1953. 135233, allotype, same data.

PARATYPES (6): 125589, Wilson Creek, 2 miles northeast of Clifton Forge, Alleghany County, Virginia, R. L. Hoffman, June 13, 1945. 125590, Potts Creek Valley, 2 miles southwest of Covington, Alleghany County, Virginia, R. L. Hoffman, July 19, 1947. 125591, 1 mile north of Eagle Rock, Botetourt County, Virginia, 950 ft., R. L. Hoffman, Aug. 24, 1944. 133046, Clifton Forge, Alleghany County, Virginia, R. L. Hoffman, Sept. 26, 1944. 135235-6, Roxbury Hollow, Alleghany County, Virginia, R. L. Hoffman, May 19, 1944.

Cnemidophorus tessellatus aethiops Cope

Report of the U.S. National Museum . . . for the year ending June 30, 1898, p. 582, 1900.

=*Cnemidophorus tigris aethiops* Cope

COTYPES (5): 64240-1, 64243-5, Hermosillo, Sonora, Mexico, O. P. Jenkins and B. W. Evermann, July 4-8, 1887. 64242 was exchanged to MCZ in 1944.

Cnemidophorus tessellatus multiscutatus Cope

Trans. Amer. Philos. Soc., vol. 17, p. 38, 1892.

=*Cnemidophorus tigris multiscutatus* Cope

COTYPES (4): 15160-3, Cedros Island (given as "Cerros Island" in original description), Baja California, Mexico, U.S. Fish Commission *Albatross*, May 5, 1888.

Cnemidophorus tessellatus rubidus Cope

Trans. Amer. Philos. Soc., ser. 2, vol. 17, p. 36, pl. 12, fig. F, 1892.

=*Cnemidophorus tigris rubidus* Cope

COTYPES (7): 15149-55, Santa Margarita Island, Baja California, Mexico, U.S. Fish Commission *Albatross*, May 2, 1888.

Cnemidophorus tigris Baird and Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 69, 1852.

=*Cnemidophorus tigris tigris* Baird and Girard

LECTOTYPE: 4103, valley of the Great Salt Lake, Utah, H. Stansbury.

PARATYPES (4): 3047 (4), same data.

Cnemidophorus undulatus Hallowell

Proc. Acad. Nat. Sci. Philadelphia, vol. 7, p. 94, 1854. (Not of Wiegmann, 1834.)

=*Cnemidophorus tigris gracilis* Baird and Girard

COTYPES (2): 3040 (2), Fort Miller, Fresno County, California (restricted to Imperial County, California, near Yuma, Arizona), A. L. Heermann.

Cnemidophorus unicolor Cope

Proc. Amer. Philos. Soc., vol. 17 (1878), p. 93, 1877.

=*Cnemidophorus guttatus guttatus* Wiegmann

HOLOTYPE: 30385, Juchitán, Oaxaca, Mexico, F. Sumichrast.

Cnemidophorus vandenburghi Dickerson

Bull. Amer. Mus. Nat. Hist., vol. 41, p. 477, 1919.

=*Cnemidophorus tigris rubidus* Cope

HOLOTYPE: 64449, Carmen Island, Gulf of California, Baja California, Mexico, C. H. Townsend, Apr. 2, 1911.

Cnemidophorus variolosus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 15, p. 39, 1863.

=*Cnemidophorus tigris variolosus* Cope

HOLOTYPE: 3066, Alamo de Parras, Coahuila, Mexico, D. N. Couch.

Coleonyx brevis Stejneger

North Amer. Fauna, No. 7, p. 163, 1893.

=*Coleonyx variegatus brevis* Stejneger

HOLOTYPE: 13627, Helotes, Bexar County, Texas, G. W. Marnock, Nov. 30, 1883.

PARATYPES (5): 42471-3, 50040-1 (all formerly 13627, which originally had 6 specimens), same data.

Coleonyx variegatus abbotti Klauber

Trans. San Diego Soc. Nat. Hist., vol. 10, p. 154, 1945.

PARATYPE: 62247, Calmallí, Baja California, Mexico, E. W. Nelson and E. A. Goldman, 1905.

Coleonyx variegatus bogerti Klauber

Trans. San Diego Soc. Nat. Hist., vol. 10, p. 176, 1945.

PARATYPES (4): 104256-9, Roosevelt Reservoir, Gila County, Arizona, E. L. Little, Jr., 1936-38.

Coleonyx variegatus peninsularis Klauber

Trans. San Diego Soc. Nat. Hist. vol. 10, p. 160, 1945.

PARATYPE: 67382, ♂, between Loreto and Comondú, Baja California, Mexico, W. M. Mann, February 1924.

Coleonyx variegatus sonoriensis Klauber

Trans. San Diego Soc. Nat. Hist., vol. 10, p. 162, 1945.

PARATYPES (2): 134301, San Carkis Bay, La Posa, Sonora, Mexico, E. H. Taylor, Aug. 8, 1934. 16808, Nogales, Arizona? (probably Nogales in Sonora, Mexico), P. L. Jouy.

Crotaphytus baileyi Stejneger

North Amer. Fauna, No. 3, p. 103, pl. 12, fig. 1, 1890.

=*Crotaphytus collaris baileyi* Stejneger

HOLOTYPE: 15821, Painted Desert, Little Colorado River, Arizona, C. H. Merriam and V. Bailey, Sept. 26, 1889.

PARATYPES (6): 15822-7, Painted Desert, Little Colorado River, Arizona, C. H. Merriam and V. Bailey, Sept. 21-26, 1889.

Crotaphytus copei Yarrow

Proc. U.S. Nat. Mus., vol. 5, p. 441, 1882.

=*Gambelia wislizenii wislizenii* (Baird and Girard)

HOLOTYPE: 12663, La Paz, Baja California, Mexico, L. Belding, 1882.

Crotaphytus dickersonae Schmidt

Bull. Amer. Mus. Nat. Hist., vol. 46, p. 638, fig. 2, 1922.

HOLOTYPE: 64451, ♀, Tiburón Island, Gulf of California, Sonora, Mexico, C. H. Townsend, Apr. 12, 1911.

Crotaphytus dorsalis Baird and Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 126, 1852.

=*Dipsosaurus dorsalis* (Baird and Girard)

HOLOTYPE: 2699, Colorado Desert, California (restricted to Winterhaven, Imperial County), J. L. LeConte.

Crotaphytus fasciatus Hallowell

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 207, 1852.

=*Gambelia wislizenii wislizenii* (Baird and Girard)

HOLOTYPE: 2736, Jornada del Muerte, Dona Ana County, New Mexico, S. W. Woodhouse.

Crotaphytus gambelii Baird and Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 126, 1852.

=*Gambelia wislizenii wislizenii* (Baird and Girard)

HOLOTYPE: 2722, California, W. Gambel.

Crotaphytus reticulatus Baird

Proc. Acad. Nat. Sci. Philadelphia, vol. 10, p. 253, 1858.

COTYPES (4): 2692 (3), Ringgold Barracks, Montague County, Texas, J. H. Clark. 2731, below Laredo, Webb County, Texas, A. Schott.

Crotaphytus silus Stejneger

North Amer. Fauna, No. 3, p. 105, 1890.

=*Gambelia wislizenii silus* (Stejneger)

HOLOTYPE: 11790A, Fresno, Fresno County, California, G. Eisen.

PARATYPES (8): 104578-85, same data.

Crotaphytus wislizenii Baird and Girard

in Stansbury, Report Expl. and Surv. Valley of the Great Salt Lake of Utah . . . , p. 340, pl. 3, 1852.

=*Gambelia wislizenii wislizenii* (Baird and Girard)

HOLOTYPE: 2770, Colorado, H. B. Möllhausen. The original description states: "Figured specimen caught near Santa Fe, New Mexico, by Dr. Wislizenus."

Ctenosaura bakeri Stejneger

Proc. U.S. Nat. Mus., vol. 23, p. 467, 1901.

=*Enyaliosaurus bakeri* (Stejneger)

HOLOTYPE: 26317, Utila Island, Honduras, J. E. Jarnigan, 1897.

PARATYPE: 25324, same data.

Ctenosaura brevirostris Cope

Proc. Amer. Philos. Soc., vol. 23, p. 268, 1886.

=*Ctenosaura pectinata* (Wiegmann)

COTYPES (2): 24708-9, Colima, Colima, Mexico, J. Xantus.

Ctenosaura clarkii Bailey

Proc. U.S. Nat. Mus., vol. 73, No. 2733, p. 44, pl. 27, 1928.

=*Enyaliosaurus clarkii* (Bailey)

PARATYPE: 21449 (given incorrectly as 21499 in original description), Mexico, Mexican Commission through F. Ferrari-Perez.

Ctenosaura completa Bocourt

Récherches zoologiques, in Mission Scientifique au Mexique et dans l'Amérique Centrale, pt. 3, p. 145, 1874.

=*Ctenosaura similis similis* (Gray)

COTYPE: 11003, Mexico, MHNP.

Ctenosaura conspicuosa Dickerson

Bull. Amer. Mus. Nat. Hist., vol. 41, p. 461, 1919.

=*Ctenosaura hemilopha* (Cope)

COTYPE: 64440, ♂, San Estéban Island, Gulf of California, Sonora, Mexico, C. H. Townsend, Apr. 13, 1911.

Ctenosaura insulana Dickerson

Bull. Amer. Mus. Nat. Hist., vol. 41, p. 462, 1919.

=*Ctenosaura hemilopha* (Cope)

HOLOTYPE: 64439, ♂, Cerralvo Island, Gulf of California, Baja California, Mexico, C. H. Townsend, Apr. 19, 1911.

Ctenosaura multispinis Cope

Proc. Amer. Philos. Soc., vol. 23, p. 267, 1886.

=*Ctenosaura acanthura* (Shaw)

COTYPE: 72737 (skin), ♂, Dondominguillo, Oaxaca, Mexico, F. Sumichrast, Aug. 10, 1868.

Ctenosaura palearis Stejneger

Proc. U.S. Nat. Mus., vol. 21, p. 381, 1898.

=*Enyaliosaurus palearis* (Stejneger)

HOLOTYPE: 22703, ♂, Gualán, Guatemala, K. I. P. McElroy.

PARATYPES (2): 22704, ♂, 24459, ♂, same data.

Ctenosaura parkeri Bailey

Proc. U.S. Nat. Mus., vol. 73, No. 2733, p. 29, pls. 14-15, 1928.

=*Ctenosaura pectinata* (Wiegmann)

HOLOTYPE: 18967, ♀, Barranca Ibarra, Jalisco, Mexico, P. L. Jouy, Apr. 22, 1892.

PARATYPE: 18970, same data.

Ctenosaura teres brachylopha Cope

Proc. Amer. Philos. Soc., vol. 23, p. 269, 1886.

=*Ctenosaura pectinata* (Wiegmann)

COTYPES (4): 7180-3, Mazatlán, Sinaloa, Mexico, F. Bischoff. Stuffed skins.

Cyclura carinata bartschi Cochran

Journ. Washington Acad. Sci., vol. 21, p. 39, 1931.

HOLOTYPE: 81212, ♂, Booby Cay, east of Mariguana Cay, Bahama Islands, P. Bartsch, July 21, 1930.

PARATYPES (4): 81213-5, 81217, same data. 81216 was exchanged to MCZ in 1934.

Cyclura (Ctenosaura) hemilopha Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 15, p. 105, 1863.

=*Ctenosaura hemilopha* (Cope)

COTYPES (5): 5295 (4), near Soria Ranch, Cape San Lucas, Baja California, Mexico, J. Xantus, September 1859. 69489, San Nicolás, between Cape San Lucas and La Paz, Baja California, Mexico, J. Xantus, September 1859.

Cyclura mattea Miller

Proc. U.S. Nat. Mus., vol. 54, p. 509, pl. 81, fig. 4, 1918.

HOLOTYPE: 59358, Magen's Bay, St. Thomas, Virgin Islands, T. de Booy.

PARATYPE: 59359, same data. Both holotype and paratype consist only of a left humerus of this apparently extinct form.

Cyclura nigerrima Cope

Proc. Amer. Philos. Soc., vol. 23, p. 264, 1886.

HOLOTYPE: 9974, Navassa Island, West Indies, G. A. Otis, July 1878.

Cyclura onchiopsis Cope

Amer. Nat., vol. 19, p. 1006, 1885.

=*Cyclura nigerrima* Cope

COTYPE: 9977, Navassa Island, West Indies, H. E. Klotz, July 1878. 12239, juvenile, from Navassa Island collected by W. J. Rasin, is probably also a cotype. One cotype from 9977 was exchanged to MCZ.

Cyclura rileyi Stejneger

Proc. Biol. Soc. Washington, vol. 16, p. 130, 1903.

HOLOTYPE: 31969, ♀, Watling Island, Bahama Islands, J. H. Riley, July 13, 1903.

PARATYPES (4): 31965-6, 31970-1, same data. 31964 was exchanged to MCZ in 1913; 31967 to UIMNH, 1956.

Cyclura stejnegeri Barbour and Noble

Bull. Mus. Comp. Zool., vol. 60, p. 163, pl. 12, figs. 1-2, 1916.

HOLOTYPE: 29367, Mona Island, Puerto Rico, B. S. Bowditch, August 1901.

PARATYPE: 29366, skin and humerus, same data. 29365 was exchanged to MCZ in 1915.

Daconura bivittata Hallowell

Proc. Acad. Nat. Sci. Philadelphia, vol. 12 (1860), p. 482, 1861.

=*Basiliscus vittatus* Wiegmann

HOLOTYPE: 6056, Nicaragua, C. Wright.

Dactyloperus insulensis Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 10, p. 197, 1858.

=*Peropus mutilatus* Gray

PROBABLE COTYPE: 21219, Sandwich Islands (Hawaiian Islands), U.S. Exploring Expedition, 1840-41. Dr. Stejneger left this note: "Found in collection without data, but in a bottle similar to the other exploring expedition specimens."

Deiropyx bartschi Cochran

Proc. Biol. Soc. Washington, vol. 41, p. 169, 1928.

HOLOTYPE: 75805, Baños San Vicente, Pinar del Río, Cuba, P. Bartsch, June 25, 1928.

PARATYPES (22): 51820, 51829-38, Viñales, Cuba, J. B. Henderson and P. Bartsch, May 30, 1914. 54417-8, Luis Lazo, Cuba, J. B. Henderson and P. Bartsch, 1916. 75797-804, 75806, Baños San Vicente, Pinar del Río, Cuba, P. Bartsch, June 25, 1928. 51825 was exchanged to MCZ in 1928; 51826 to BMNH, 1929; 51827 to R; 51828 to UIMNH, 1956.

Dicrodon holmbergi Schmidt

Chicago Nat. Hist. Mus., Fieldiana: Zoology, vol. 39, p. 66, figs. 10-11, 1957.

HOLOTYPE: 127823, Virú Valley, Peru, A. Holmberg, April 1948.

PARATYPES (2): 127824-5, same data. 127822 was exchanged to CNHM in 1957.

Diphalus fenestratus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 13, p. 76, 1861.

=*Amphisbaena fenestrata* (Cope)

COTYPE: 11715, St. John, Virgin Islands, A. H. Riise.

Diplodactylus inexpectatus Stejneger

Proc. U.S. Nat. Mus., vol. 16, p. 714, 1894.

HOLOTYPE: 20433, Ile Mahé, Seychelles Islands (Indian Ocean), W. L. Abbott, 1892.

Diplodactylus unctus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 15, p. 102, 1863.

=*Phyllodactylus unctus* (Cope)

COTYPE: 5304, Cape San Lucas, Baja California, Mexico, J. Xantus, March 1860.

Diploderma polygonatum Hallowell

Proc. Acad. Nat. Sci. Philadelphia, vol. 12 (1860), p. 490, 1861.

=*Japalura polygonata* (Hallowell)

HOLOTYPE: 12203, Amakarima, Riu Kiu (Ryukyu) Islands, J. Rodgers, April 1855.

Diploglossus chalybaeus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 18 (1866), p. 321, 1867.

=*Celestus enneagrammus* (Cope)

COTYPE: 6603, Mount Orizaba, Veracruz, Mexico, 4,000-6,000 ft., F. Sumichrast.

Diploglossus steindachneri Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 16, p. 179, 1864.

=*Celestus enneagrammus* (Cope)

HOLOTYPE: 6342, Orizaba, Veracruz, Mexico, F. Sumichrast.

Dipso-saurus dorsalis sonoriensis Allen

Occ. Pap. Mus. Zool. Univ. Michigan, No. 259, p. 4, 1933.

=*Dipsosaurus dorsalis sonoriensis* Allen

PARATYPE: 93206 (formerly MZUM 72128), 5 miles south of Hermosillo, Sonora, Mexico, M. J. Allen, July 1, 1932.

Doryura vulpecula Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 9, p. 197, 1857.

=*Hemidactylus garnotii* Duméril and Bibron

PROBABLE COTYPES (2): 11288 (2), Sandwich Islands? (Hawaii?), U.S. Exploring Expedition, 1840-41.

Draco mindanensis Stejneger

Proc. U.S. Nat. Mus., vol. 33, p. 677, 1908.

HOLOTYPE: 37388, Catagan, at base of Malindang Mountain, Mindanao, Philippine Islands, E. A. Mearns, May 11, 1906.

PARATYPE: 37387, same data.

Elgaria grandis Baird and Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 176, 1852.

=*Gerrhonotus multicaarinatus scincicaudus* Skilton

HOLOTYPE: 9057, Oregon, U.S. Exploring Expedition, 1841.

Elgaria nobilis Baird and Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 129, 1852.

=*Gerrhonotus kingii* Gray

COTYPES (2): 3076 (2), Copper Mines of the Gila (Santa Rita del Cobre), Fort Webster (Luna County), New Mexico, J. H. Clark.

Elgaria principis Baird and Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 175, 1852.

=*Gerrhonotus coeruleus principis* (Baird and Girard)

COTYPES (3): 3106, "Oregon" (restricted to Puget Sound, Washington), U.S. Exploring Expedition, 1841. 5134 (2), Puget Sound, C. J. Pickering.

Emoia nigrita Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 10, p. 197, 1858.

=*Emoia nigrita* Girard

HOLOTYPE: 5631, Navigator Islands, Polynesia, U.S. Exploring Expedition, 1839.

Emoia arnoensis Brown and Marshall

Copeia, No. 4, p. 201, 1953.

HOLOTYPE: 132131, ♀, Chittakinmatoroen Island, Arno Atoll, Marshall Islands, J. T. Marshall, Jr., Aug. 15, 1950.

PARATYPES (4): 132133, 132135, Eneman Island, Arno Atoll, Marshall Islands, J. T. Marshall, Jr., Aug. 25, 1950. 132134, East Tagelib Islet, Arno Atoll, Marshall Islands, J. T. Marshall, Jr., Aug. 26, 1950. 132138, Chittakinmatoroen Island, Arno Atoll, Marshall Islands, J. T. Marshall, Jr., Aug. 15, 1950.

Emoia boettgeri orientalis Brown and Marshall

Copeia, No. 4, p. 204, 1953.

HOLOTYPE: 132132, ♂, Ine Island, Arno Atoll, Marshall Islands, J. T. Marshall, Jr., July 30, 1950.

PARATYPES (4): 132137, 132140, same data. 132138, Dodo Island, Arno Atoll, Marshall Islands, J. T. Marshall, Jr., Aug. 26, 1950. 132139, Autore Island, Arno Atoll, Marshall Islands, J. T. Marshall, Jr., July 19, 1950.

Emoia frenata Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 187, 1862.

=*Mabuya frenata* (Cope)

COTYPE: 5855, Paraguay, T. J. Page.

Epaphelus sumichrasti Cope

Journ. Acad. Sci. Philadelphia, ser 2, vol. 8 (1875), p. 115, 1876.

=*Gymnophthalmus sumichrasti* (Cope)

COTYPES (2): 30245-6, near Ventose Bay, Tehuantepec, Oaxaca, Mexico, F. Sumichrast.

Eremias hoehneli Stejneger

Proc. U.S. Nat. Mus., vol. 16, p. 719, 1894.

HOLOTYPE: 20077, from the coast to Hameye, Tana River, East Africa (Kenya), W. A. Chanler, November 1892.

Eremias nitida garambensis Schmidt

Bull. Amer. Mus. Nat. Hist., vol. 39, p. 511, text-figs. 18-19, 1919.

PARATYPE: 62014 (formerly AMNH 10708), Garamba, Belgian Congo, Lang-Chapin Expedition, June 1912.

Eremias sextaeniata Stejneger

Proc. U.S. Nat. Mus., vol. 16, p. 718, 1894.

=*Eremias spekkii sextaeniata* Stejneger

HOLOTYPE: 20080, from the coast to Hameye, Tana River, British East Africa (Kenya), W. A. Chanler, November 1892.

PARATYPE: 20079, same data.

Euchirotos biporus Cope

Amer. Nat., vol. 28, p. 436, figs. 5a-c, 1894.

=*Bipes biporus* (Cope)

HOLOTYPE: 21324 (formerly 12599), La Paz, Baja California, Mexico, L. Belding, 1882.

PARATYPES (2): 8568, Cape San Lucas, Baja California, Mexico, T. H. Streets. 12599, La Paz, Cape San Lucas, Baja California, Mexico, L. Belding, 1882.

Eumeces brevilineatus Cope

U.S. Nat. Mus. Bull. 17, p. 18, 1880.

COTYPES (2): 10159 (2), Helotes Creek, Bexar County, Texas (given as "20 miles northwest of San Antonio" in original description), G. W. Marnock, May 1879. The lectotype is 10159a.

Eumeces brevirostris bilineatus Tanner

Great Basin Nat., vol. 18, p. 59, fig. 2, 1958.

PARATYPE: 64666, La Laguna, Jalisco, Mexico, E. W. Nelson and E. A. Goldman, Mar. 23, 1897.

Eumeces copei Taylor

Proc. Biol. Soc. Washington, vol. 46, p. 133, 2 figs., 1933.

PARATYPES (3): 7037, overland route, probably Mexico. 32291, Valley of Mexico and Toluca, Mexico, C. T. Hoege, 1884-85. 92547, 10 miles from Asunción, Mexico, Mexico, E. H. Taylor and H. M. Smith, Aug. 6, 1932.

Eumeces epipleurotis Cope

U.S. Nat. Mus. Bull. 17, p. 40, 1880.

=*Eumeces multivirgatus* Hallowell

COTYPES (2): 5263, northern boundary of Texas, J. H. Clark. 9219 (formerly 5411), Platte River, Fort Kearney, Buffalo County, Nebraska, C. S. McCarthy.

Eumeces indubitus Taylor

Univ. Kansas Sci. Bull., vol. 21 (1933), p. 258, pls. 24-25, 1934.

PARATYPE: 92548, ♂, near Asunción, Mexico, Mexico, E. H. Taylor and H. M. Smith, Aug. 6, 1932.

Eumeces lagunensis Van Denburgh

Proc. California Acad. Sci., ser. 2, vol. 5, p. 134, pl. 13, 1895.

NEOTYPE: 67398, on the trail between Loreto and Comondú, Lower California, Mexico, W. M. Mann, February 1924. This designation of neotype may be invalid since the designated specimen is not from the original type locality, given as San Francisquito, Sierra de la Laguna, Baja California, Mexico, fide H. M. Smith.

NEOPARATYPES (5): 67399-67403, same data.

Eumeces laticutatus okadae Stejneger

U.S. Nat. Mus. Bull. 58, p. 200, fig. 181, 1907.

HOLOTYPE: 23891, Miyakeshima, Idzu Seven Islands (Izu-shichito), Japan, N. Okada, May 3, 1887.

PARATYPES (5): 23892, 36534, same data. 23895-6, 36532, Niishima, Idzu Seven Islands, Japan, N. Okada, 1887. 36531 was exchanged to CAS; 36533 to MCZ, 1944.

Eumeces managuae Dunn

Proc. Biol. Soc. Washington, vol. 46, p. 67, 1933.

HOLOTYPE: 89474, ♂, Aviation Field, Managua, Nicaragua, J. H. Ivy, 1932.

Eumeces marginatus amamiensis Van Denburgh

Advance diagnoses of new reptiles and amphibians from the Loo Choo Islands and Formosa, p. 4, July 29, 1912 (published privately, San Francisco, California).

=*Eumeces oshimensis* Thompson

PARATYPE: 64210 (CAS 21547), Amami Oshima, Loo Choo (Ryukyu) Islands, V. Kuhne, May 1, 1910.

***Eumeces multilineatus* Tanner**

Great Basin Nat., vol. 17, p. 111, fig. 1a, 1957.

PARATYPE: 30833, Chihuahua, Mexico, H. H. and C. S. Brimley.

***Eumeces ochoterena* Taylor**

Proc. Biol. Soc. Washington, vol. 46, p. 129, 2 figs., 1933.

PARATYPE: 92549, between Rincón and Cajones, south of Chilpancingo, Guerrero, Mexico, E. H. Taylor and H. M. Smith, July 1, 1932.

***Eumeces parviauriculatus* Taylor**

Proc. Biol. Soc. Washington, vol. 46, p. 178, fig. 2, 1933.

HOLOTYPE: 47536, near Alamos, Sonora, Mexico, E. A. Goldman, Jan. 5, 1899.

***Eumeces parvulus* Taylor**

Proc. Biol. Soc. Washington, vol. 46, p. 175, fig. 1, 1933.

HOLOTYPE: 56903, Tepic, Nayarit, Mexico, J. Hurter (donor), Apr. 10, 1910.

***Eumeces pekinensis* Stejneger**

Occ. Pap. Boston Soc. Nat. Hist., vol. 5, p. 120, 1924.

HOLOTYPE: 60863, Hsin-Lung-Shan District, Chili Province, Imperial Hunting Grounds, North China, A. de C. Sowerby, August 1917.

PARATYPES (2): 60864-5, same data.

***Eumeces pluvialis* Cope**

U.S. Nat. Mus. Bull. 17, p. 19, 1880.

=*Eumeces anthracinus pluvialis* Cope

NEOTYPE: 75291, Mobile, Mobile County, Alabama, H. P. Löding.

***Eumeces quadrilineatus* Hallowell**

Report upon the reptiles collected on the survey, in Report U.S. Pacific R.R. Explor. and Surv., vol. 10, pt. 4, p. 10, pl. 9, figs. 3a-d, 1859.

(Not of Blyth, 1853.)

=*Eumeces skiltonianus skiltonianus* (Baird and Girard)

HOLOTYPE: 3168, near Mojave River in San Bernardino Valley, California (restricted to Mojave River), A. L. Heermann.

***Eumeces quadrivirgatus* Hallowell**

Proc. Acad. Nat. Sci. Philadelphia, vol. 12 (1860), p. 502, 1861.

=*Eumeces quadrilineatus* (Blyth)

HOLOTYPE: 7498, Hong Kong (off the mainland of China), D. Wright and W. Stimpson, May 4, 1854.

Eumeces septentrionalis pallidus Smith and Slater

Trans. Kansas Acad. Sci., vol. 52, p. 438, 1949.

PARATYPE: 58337, Paisano, Brewster County, Texas, J. Hurter, Apr. 30, 1905.

Eumeces skiltonianus brevipes Cope

Report of the U.S. National Museum . . . for the year ending June 30, 1898, p. 643, 1900.

=*Eumeces gilberti gilberti* Van Denburgh

HOLOTYPE: 12558, Fresno, Fresno County, California, G. Eisen.

Eumeces skiltonianus interparietalis Tanner

Great Basin Nat., vol. 17, p. 73, 1957.

PARATYPES (5): 23723, San Pedro Mountain, Baja California, Mexico, C. H. Townsend, 1889. 37686, Todos Santos Islands, Baja California, Mexico, E. W. Nelson and E. A. Goldman, May 21, 1905. 52412, North Island, Los Coronados Islands, Baja California, Mexico, A. E. Colburn, June 2, 1915. 64401, Los Coronados Islands, Baja California, Mexico, W. P. Taylor, Apr. 5, 1921. 75082, North Coronado Island, Baja California, Mexico, A. Wetmore, May 6, 1928.

Eumeces tunganus Stejneger

Journ. Washington Acad. Sci., vol. 14, p. 384, 1924.

HOLOTYPE: 66736, near Luting-Chiao, Szechwan, China, 5,000-6,000 ft., D. C. Graham, Aug. 9, 1923.

PARATYPE: 66737, same data.

Euphryne obesus Baird

Proc. Acad. Nat. Sci. Philadelphia, vol. 10 (1858), p. 253, 1859.

=*Sauromalus obesus obesus* (Baird)

COTYPE: 4172, Fort Yuma (Imperial County), California, A. Schott and G. G. Thomas.

Euprepis venustus Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 9 (1857), p. 195, 1858.

=*Mabuya delalandii* (Gray)

HOLOTYPE: 12205, San Jago, Cape Verde Islands, (off West Africa), U.S. Exploring Expedition, 1838.

Gaigeia dontomasi Smith

Proc. U.S. Nat. Mus., vol. 92, p. 374, 1942.

HOLOTYPE: 111473, Lachiguirí, Oaxaca, Mexico, 7,100 ft., T. MacDougall, Jan. 20, 1940.

PARATYPE: 111474, same data.

Gaigeia radula Smith

Proc. U.S. Nat. Mus., vol. 92, p. 376, 1942.

HOLOTYPE: 111472, ♀, San José Manteca, Oaxaca, Mexico, H. M. Smith, May 1940.

Gehyra vorax Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 9 (1857), p. 197, 1858.

COTYPES (3): 5698, Raraka Island, Tahiti, U.S. Exploring Expedition, 1840. 5699 (2), Fiji Islands, U.S. Exploring Expedition, 1840. One cotype from 5699 was exchanged to ANSP.

Gekko indicus Girard

in U.S. Exploring Expedition . . . : Herpetology, p. 290, pl. 16, figs. 9-16 (atlas), 1958.

=*Gekko gekko* (Linnaeus)

HOLOTYPE: 5681, Mangsi Island, Balabac Strait, Philippine Islands, U.S. Exploring Expedition, 1842.

Gerrhonotus alfaroi Stejneger

Proc. U.S. Nat. Mus., vol. 32, p. 505, 1907.

=*Barisia monticola* (Cope)

HOLOTYPE: 37342, Santa María, northeast of Laguna, Costa Rica, 2,000 m., J. F. Tristan, January 1907.

Gerrhonotus auritus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 20 (1868), p. 306, 1869.

=*Abronia aurita* (Cope)

HOLOTYPE: 6769, Vera Paz, Guatemala, H. Hague.

Gerrhonotus gramineus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 16, p. 179, 1864.

=*Abronia taeniata graminea* (Cope)

COTYPES (4): 6327 (4), Orizaba, Veracruz, Mexico, F. Sumichrast.

Gerrhonotus infernalis Baird

Proc. Acad. Nat. Sci. Philadelphia, vol. 10 (1858), p. 255, 1859.

=*Gerrhonotus liocephalus infernalis* Baird

HOLOTYPE: 3090, Devils River, Texas, C. B. R. Kennerly.

Gerrhonotus levicollis ciliaris Smith

Proc. U.S. Nat. Mus., vol. 92, p. 365, 1942.

=*Barisia imbricata ciliaris* (Smith)

HOLOTYPE: 47496, Sierra Guadalupe, Coahuila, Mexico, 9,500 ft., E. W. Nelson and E. A. Goldman, Apr. 28, 1902.

PARATYPES (8): 46723-4, Sierra Madre, Zacatecas, Mexico, 8,500 ft., E. W. Nelson and E. A. Goldman, Aug. 17, 1897. 46843, Indé, Durango, Mexico, 6,500 ft., E. W. Nelson and E. A. Goldman, Aug. 14, 1898. 47208-

10, mountains near Jesús María, San Luis Potosí, Mexico, E. W. Nelson and E. A. Goldman, Sept. 4, 1892. 47497-8, Sierra Guadalupe, Coahuila, Mexico, 9,500 ft., E. W. Nelson and E. A. Goldman, Apr. 28, 1902. 47207 was exchanged to UIMNH in 1956.

Gerrhonotus monticolus Cope

Proc. Amer. Philos. Soc., vol. 17 (1878), p. 97, 1877.

=*Barisia monticola* (Cope)

HOLOTYPE: 30591, Pico Blanco summit, 11,500 ft., Costa Rica, W. M. Gabb.

Gerrhonotus olivaceus Baird

Proc. Acad. Nat. Sci. Philadelphia, vol. 10 (1858), p. 255, 1859.

=*Barisia imbricata imbricata* (Wiegmann)

COTYPES (2): 3096 (2), Mexico (given as "near San Diego" in original description), A. Schott.

Gerrhonotus ophiurus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 18 (1866), p. 321, 1867.

=*Gerrhonotus liocephalus ophiurus* Cope

HOLOTYPE: 30206, Orizaba, Veracruz, Mexico, F. Sumichrast.

Gerrhonotus scincicauda palmeri Stejneger

North Amer. Fauna, No. 7, p. 196, 1893.

=*Gerrhonotus coerulea palmeri* (Stejneger)

HOLOTYPE: 18606, ♂, south fork, King's River, Fresno County, California, T. S. Palmer, 1891.

PARATYPES (9): 18607-9, ♀, ♀, ♂, east fork, Kaweah River, California, V. O. Bailey, Aug. 1-8, 1891. 18610-11, ♂, ♀, Soda Springs, north fork, Kern River, California, E. W. Nelson, Sept. 6, 1891. 18612, ♀, north fork, Kern River, California, E. W. Nelson, Sept. 15, 1891. 18613, ♂, Soda Springs, north fork, Kern River, California, V. O. Bailey, Aug. 15, 1891. 18614, ♀, Sequoia National Park, California, A. K. Fisher, Aug. 2, 1891. 18615, ♀, Mineral King, California, 8,000 ft., V. O. Bailey, Aug. 6, 1891.

Gerrhonotus webbii Baird

Proc. Acad. Nat. Sci. Philadelphia, vol. 10 (1858), p. 255, 1859.

=*Elgaria multicarinata webbii* (Baird)

HOLOTYPE: 3078, restricted to vicinity of San Diego, San Diego County, California (original description states: "San Diego to El Paso"), T. H. Webb.

Gonatodes atricucularis Noble

Ann. New York Acad. Sci., vol. 29, p. 135, 1921.

PARATYPES (10): 64040-44, Bellavista, Peru, H. C. Raven, 1916. 75941-5, Bellavista, Peru, G. K. Noble, 1916. 64039 was exchanged to UIMNH in 1956.

Gonatodes nigridius Smith

Sarawak Mus. Journ., vol. 3, p. 22, 1925.

PARATYPE: 76632, ♀, Lundu, Mount Gadin, North Borneo, M. A. Smith.

Gonocephalus abbotti Cochran

Proc. U.S. Nat. Mus., vol. 60, Art. 2421, p. 1, 1922.

HOLOTYPE: 24028, ♀, Trang, Lower Siam (Thailand), W. L. Abbott.

Heloderma suspectum Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 21, p. 5, 1869.

COTYPES (3): 2971 (3), Sierra de la Union, Sonora (Arizona), A. Schott.

Hemidactylus inornatus Hallowell

Proc. Acad. Nat. Sci. Philadelphia, vol. 12 (1860), p. 492, 1861.

=*Hemidactylus frenatus* Gray

COTYPES (3): 7429 (3), Loo Choo (Ryukyu Islands), W. Stimpson, November 1854.

Hemiphyllodactylus leucostictus Stejneger

Proc. U.S. Nat. Mus., vol. 21, p. 800, 1899.

HOLOTYPE: 23500, Kauai, Hawaii, V. Knudsen, May 1895.

PARATYPES (5): 21220, probably Hawaii, U.S. Exploring Expedition?, 1840?. 23459, Hawaii Island, Hawaii, H. W. Henshaw. 23485, 23499, Kauai, Hawaii, V. Knudsen, May 1895. 23509, Honolulu, Oahu, Hawaii, L. Stejneger, Nov. 17, 1896. 23460 was exchanged to R.

Heteronota pelagica Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 9 (1857), p. 197, 1858.

=*Gymnodactylus pelagicus* (Girard)

COTYPES (2): 5626 (2), Fiji Islands, U.S. Exploring Expedition, 1840.

Hispaniolus pratensis Cochran

Proc. Biol. Soc. Washington, vol. 41, p. 49, 1928.

HOLOTYPE: 69189, near San Michel, Haiti, G. S. Miller, Jr., Mar. 23, 1925.

PARATYPES (4): 69184-6, 59188, L'Atalaye Plantation, Haiti, G. S. Miller, Jr., Apr. 6, 1925. 69187 was exchanged to MCZ in 1959.

Holbrookia affinis Baird and Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 125, 1852.

=*Holbrookia texana texana* Troschel

COTYPES (3): 2662 (3), along Rio San Pedro of the Rio Grande del Norte, Texas, J. H. Clark.

Holbrookia dickersonae Schmidt

Amer. Mus. Novit., No. 22, p. 2, 1921.

=*Holbrookia maculata dickersonae* Schmidt

HOLOTYPE: 2668a, Castañuelas, Coahuila, Mexico, D. N. Couch.

PARATYPES (5): 2668b-d, same data. 2664 (2), Alamos de Parras, Coahuila, Mexico, D. N. Couch.

Holbrookia lacerata Cope

U.S. Nat. Mus. Bull. 17, p. 15, 1880.

=*Holbrookia lacerata lacerata* Cope

COTYPES (2): 10160 (2), Helotes, Bexar County, Texas, G. W. Marnock, May 1879.

Holbrookia lacerata subcaudalis Axtell

Bull. Chicago Acad. Sci., vol. 10, p. 174, pl. 2, figs. 1-2, 1956.

PARATYPES (3): 32899, near Cotulla, La Salle County, Texas, H. C. Oberholser, May 9, 1900. 46697, Sabinas, Coahuila, Mexico, E. W. Nelson and E. A. Goldman, May 27, 1902. 81983, Sinton, San Patricio County, Texas, J. D. Mitchell, Oct. 11, 1913. These specimens were used as paratypes, but were not so listed in original description.

Holbrookia maculata perspicua Axtell

Bull. Chicago Acad. Sci., vol. 10, p. 166, pl. 1, figs. 1-2, 1956.

PARATYPES (13): 4692 (3), Neosho Falls, Woodson County, Kansas, B. F. Goss. 22150-1, 10 miles from China Springs, Waco, McLennan County, Texas, H. H. and C. S. Brimley, June 8, 11, 1894. 32908-13, Henrietta, Clay County, Texas, H. C. Oberholser, July-August, 1900. 44981, Mayers, Comanche County, Oklahoma, J. H. Gaut, Apr. 28, 1904. 83123, 3 miles southwest of Dallas, Dallas County, Texas, C. E. Burt, 1930. These specimens were used as paratypes, but were not so listed in original description.

Holbrookia propinqua Baird and Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 126, 1852.

=*Holbrookia propinqua propinqua* Baird and Girard

COTYPES (7): 2671 (7), between Indianola and San Antonio, Texas (restricted to 9 miles southwest of Somerset, Atascosa County), J. H. Clark.

Holbrookia propinqua piperata Smith and Burger

Trans. Kansas Acad. Sci., vol. 53, p. 167, 1950.

PARATYPE: 137212, (formerly UIMNH 4049), ♀, Etiopia, 2 miles south of Tecolutla, Veracruz, Mexico, F. A. Shannon, W. L. Burger, and L. Firschein, Apr. 24, 1949.

Holbrookia pulchra Schmidt

Amer. Mus. Novit., No. 22, p. 1, 1921.

=*Holbrookia maculata pulchra* Schmidt

PARATYPES (20): 13973 (2), Tucson (Pima County), Arizona, E. W. Nelson, Aug. 18, 1884. 14749, Fort Huachuca (Cochise County), Arizona, H. C. Benson, Aug. 3, 1887. 5025 (3), Fort Buchanan, Arizona, B. J. D.

Irwin. 17228-33, Towne's Ranch, Nogales (Santa Cruz County), Arizona, P. L. Jouy, May 23, 1891. 19371-3, 19432, Mexican boundary, E. A. Mearns, 1892. 46980, Nogales, Sonora, Mexico, V. O. Bailey, Oct. 17, 1889. 47423-5, Mexico, U.S. Biological Survey. 17227 was exchanged to UIMNH in 1956.

Holbrookia thermophila Barbour

Proc. New England Zool. Club, vol. 7, p. 79, 1921.

=*Holbrookia maculata thermophila* Barbour

PARATYPE: 120762 (formerly MCZ 14289), San José de Guaymas, Sonora, Mexico, W. W. Brown, Jr., 1908.

Homalosaurus ventralis Hallowell

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 179, 1852.

=*Callisaurus draconoides ventralis* (Hallowell)

HOLOTYPE: 2670, "west of the Rio Grande, New Mexico" (restricted to Tucson, Pima County, Arizona), S. W. Woodhouse.

Hombroonia fasciolaris Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 9 (1857), p. 196, 1858.

=*Leiolopisma fasciolaris* (Girard)

HOLOTYPE: 5685, Bay of Islands, New Zealand, U.S. Exploring Expedition, March 1840.

Hombroonia undosa Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 9 (1857), p. 196, 1858.

=*Sphenomorphus undosus* (Girard)

HOLOTYPE: 5627, Bay of Islands, New Zealand, U.S. Exploring Expedition, March 1840.

Hoplodactylus pomarii Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 9 (1857), p. 197, 1858.

=*Hoplodactylus pacificus* (Gray)

HOLOTYPE: 5690, Bay of Islands, New Zealand, U.S. Exploring Expedition, March 1840.

Japalura mitsukurii Stejneger

Journ. Sci. Coll., Imp. Univ. Tokyo, vol. 12, p. 218, 1898.

HOLOTYPE: 36505, ♂, Botel Tobago Island, Formosa, T. Tada, May 1899.

PARATYPE: 36506, ♀, same data.

Japalura splendida Barbour and Dunn

Proc. New England Zool. Club, vol. 7, p. 18, 1919.

HOLOTYPE: 35522, gorge of Yangtze River near Ichang, Hupeh, China, E. Blackwelder, 1904.

PARATYPES (2): 35523, near Taningsien, east end of Chihsiting Pass, Szechwan, China, E. Blackwelder, 1904. 35524, Liang-ho, Chin-ling Mountains, Shensi, China, E. Blackwelder, 1904.

Laemanctus alticoronatus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 17, p. 192, 1865.

PROBABLE COTYPES (2): 12283 (2), near Mérida, Yucatán, Mexico, A. Schott.

Leiocephalus altavelensis Noble and Hassler

Amer. Mus. Novit., No. 652, p. 14, 1933.

=*Leiocephalus personatus altavelensis* Noble and Hassler

PARATYPE: 115362 (formerly AMNH 51069), Alta Vela Island, Dominican Republic, W. G. Hassler, Oct. 9-10, 1932.

Leiocephalus carinatus aquarius Schwartz and Ogren

Herpetologica, vol. 12, p. 100, 1956.

PARATYPES (7): 29586-7, Santiago de Cuba, Oriente Province, Cuba, B. S. Bowdish, Nov. 12, 7, 1901. 59202-3, Guantanamo Bay, Cuba, J. B. Henderson and P. Bartsch, Mar. 17, 1917. 81658, Macóla Hill, Guantanamo, Cuba, P. Bartsch, 1930. 81673, Río Puerco, Oriente Province, Cuba, P. Bartsch, 1930. 81687, Cabo Cruz, Oriente Province, Cuba, P. Bartsch, Aug. 31, 1930.

Leiocephalus carinatus armouri Barbour and Shreve

Proc. Boston Soc. Nat. Hist., vol. 40, p. 360, 1935.

PARATYPES (2): 120764-5, (formerly MCZ 38107-8), High Rock, Grand Bahama Island, Bahama Islands, T. Barbour, May 2, 1934.

Leiocephalus carinatus granti Rabb

Herpetologica, vol. 13, p. 109, 1957.

PARATYPES (14): 81742-9, Cayman Brac, West Indies, P. Bartsch, Sept. 8-10, 1930. 107966-71, Cayman Brac, West Indies, C. Grant, Mar. 31-Apr. 25, 1938.

Leiocephalus carinatus punctatus Cochran

Journ. Washington Acad. Sci., vol. 21, p. 39, 1931.

HOLOTYPE: 81560, Jamaica Wells, Acklin Island, Bahama Islands, P. Bartsch, July 6, 1930.

PARATYPES (22): 81481, Jamaica Bay, Acklin Island, Bahama Islands, P. Bartsch, July 5, 1930. 81482-9, hills near Cornucopia, Jamaica Wells, Acklin Island, Bahama Islands, P. Bartsch, July 7, 1930. 81492-6, Crooked Island, Bahama Islands, P. Bartsch, July 14, 1930. 81562-9, Castle Island, Bahama Islands, P. Bartsch, July 8, 1930.

***Leiocephalus inaguae* Cochran**

Journ. Washington Acad. Sci., vol. 21, p. 39, 1931.

HOLOTYPE: 81277, ♂, Man of War Bay, Great Inagua Island, Bahama Islands, P. Bartsch, Aug. 8, 1930.

PARATYPES (7): 81256, 81278, Carmichael Point, Great Inagua Island, Bahama Islands, P. Bartsch, Aug. 7, 1930. 81279, center of Ocean Bight Bay, Great Inagua Island, Bahama Islands, P. Bartsch, Aug. 7, 1930. 81280, northeast peninsula, Great Inagua Island, Bahama Islands, P. Bartsch, Aug. 6, 1930. 81281, northwest point, Great Inagua Island, Bahama Islands, P. Bartsch, Aug. 8, 1930. 81283-4, near Matthew Town, Great Inagua Island, Bahama Islands, P. Bartsch, Aug. 9, 1930. 81282 was exchanged to R in 1953.

***Leiocephalus macropus immaculatus* Hardy**

Journ. Washington Acad. Sci., vol. 48, p. 294, figs. 1-5, 1958.

HOLOTYPE: 138412, Ocuja, Oriente Province, Cuba, J. D. Hardy, Aug. 29-31, 1956.

PARATYPES (39): 138378-411, 138413-7, Ocuja, Oriente Province, Cuba, J. D. Hardy, Aug. 28-Sept. 1, 1956.

***Leiocephalus personatus aureus* Cochran**

Occ. Pap. Boston Soc. Nat. Hist., vol. 8, p. 175, 1934.

HOLOTYPE: 75909, ♂, Jacmel, Haiti, J. S. C. Boswell, 1928.

PARATYPES (6): 75910-1, 75913-5, Jacmel, Haiti, J. S. C. Boswell, 1928. 75916, Artibonite Valley, Haiti, J. S. C. Boswell, 1928. 75912 was exchanged to MZUM in 1943.

***Leiocephalus personatus louisae* Cochran**

Occ. Pap. Boston Soc. Nat. Hist., vol. 8, p. 177, 1934.

PARATYPE: 95117, Saona Island, Dominican Republic, T. Barbour, Apr. 8, 1934.

***Leiocephalus personatus lunatus* Cochran**

Occ. Pap. Boston Soc. Nat. Hist., vol. 8, p. 153, 1934.

PARATYPE: 94993, (formerly FMNH 152), Santo Domingo City, Dominican Republic, G. K. Cherrie, 1895.

***Leiocephalus personatus mentalis* Cochran**

Proc. Biol. Soc. Washington, vol. 45, p. 178, 1932.

HOLOTYPE: 65772, Jovéro, Dominican Republic, W. L. Abbott, Feb. 19, 1923.

PARATYPES (36): 10262-4, 10266 (4), 10269 (2), 10271, Puerto Plata, Dominican Republic, C. A. Fraser, 1878. 65770-1, 65775-9, 66708-9, 66712, Jovéro, Dominican Republic, W. L. Abbott, Nov. 30-Feb. 19, 1923. 66695, 66697-707, Dominican Republic, W. L. Abbott, 1923. 66710-1, Guarabo, Dominican Republic, W. L. Abbott, Nov. 21, 1923. 66764,

Samaná Peninsula, Dominican Republic, W. L. Abbott. 65774, Samaná and Laguna, Dominican Republic, W. L. Abbott, March 1923. 65773 was exchanged to AMNH in 1941; 66694 and 66696 were exchanged to MZUM in 1943.

***Leiocephalus personatus scalaris* Cochran**

Proc. Biol. Soc. Washington, vol. 45, p. 181, 1932.

HOLOTYPE: 74054, Cape Haïtien, Haiti, A. J. Poole, Mar. 3, 1928.

PARATYPES (23): 74047-51, 74053, 74055-6, 74058-63, 74065-70, 74072-4, same data. 74052 and 74064 were exchanged to MZUM in 1943; 74057 and 74071 to MCZ, 1944.

***Leiocephalus psammodromus* Barbour**

Copeia, No. 85, p. 73, 1920. (New name for *Leiocephalus arenarius* Barbour (not of Tschudi, 1845).)

PARATYPE: 81987, Bastion Cay, Turks Island, West Indies, L. A. Mowbray, June 1916.

***Leiocephalus vinculum* Cochran**

Proc. Biol. Soc. Washington, vol. 41, p. 54, 1928.

=*Leiocephalus personatus vinculum* Cochran

PARATYPE: 77061, Pointe-à-Raquettes, Gonave Island, Haiti, W. J. Eyerdam, Aug. 5, 1927.

***Leiocephalus virescens* Stejneger**

Proc. U.S. Nat. Mus., vol. 23, p. 471, 1901.

HOLOTYPE: 26758, Green Cay, Bahama Islands, U.S. Fish Commission *Albatross*, Apr. 12, 1886.

PARATYPES (13): 14568 (13), same data.

***Leiopisma barbouri* Stejneger**

Journ. Washington Acad. Sci., vol. 15, p. 150, 1925.

PARATYPES (2): 68723-4 (formerly MCZ 7261), Yunnan-Fu, Yunnan, China, 6,000 ft., J. Graham, 1906.

***Leiopisma caudaequinae* Smith**

Univ. Kansas Sci. Bull., vol. 34, p. 195, 1951.

PARATYPES (2): 137217-8, 10 miles west of Naranjo, San Luis Potosí, Mexico, E. H. Taylor.

***Leiopisma eunice* Cochran**

Proc. Biol. Soc. Washington, vol. 40, p. 187, 1927.

HOLOTYPE: 72180, Bang Suk, near Pak Jong, Siam (Thailand), Hugh M. Smith, Aug. 19, 1926.

PARATYPES (2): 70271-2, Pak Jong, Siam, Hugh M. Smith, May 18, 1925.

Leiolopisma forbesorum Taylor

Copeia, No. 1, p. 8, 1937.

=*Scincella gemmingeri forbesorum* (Taylor)

PARATYPE: 134307 (formerly UIMNH 27402), La Placita, Hidalgo, Mexico, 7,000 ft., E. H. Taylor, July 2-3, 1936.

Leiolopisma kohtaoensis Cochran

Proc. Biol. Soc. Washington, vol. 40, p. 188, 1927.

HOLOTYPE: 72284, Koh Tao, Gulf of Siam (Thailand), Hugh M. Smith, Dec. 31, 1926.

PARATYPES (2): 72282-3, same data.

Leiolopisma laterale boettgeri Van Denburgh

Advance diagnoses of new reptiles and amphibians from the Loo Choo Islands and Formosa, p. 7, July 29, 1912 (published privately, San Francisco, California).

PARATYPE: 64208 (CAS 21691), Ishigaki, Loo Choo (Ryukyu) Islands, V. Kuhne, June 2, 1910.

Leiolopisma pranensis Cochran

Proc. U.S. Nat. Mus., vol. 77, No. 2834, p. 18, 1930.

HOLOTYPE: 75591, Pran, Peninsular Siam (Thailand), Hugh M. Smith, May 25, 1928.

PARATYPE: 76850, Doi Angka, Peninsular Siam, Hugh M. Smith, Dec. 2, 1928.

Leiolopisma smithi Cochran

Copeia, No. 4, p. 238, fig. 1, 1941.

HOLOTYPE: 94536, Khun Tan Mountains, northern Siam (Thailand), Hugh M. Smith, Mar. 2, 1932.

Lepidoblepharis barbouri Noble

Ann. New York Acad. Sci., vol. 29, p. 133, 1921.

PARATYPES (6): 64035-7, Marañón Valley, Perico, northwestern Peru, H. C. Raven. 75952-4, Bellavista, Peru, G. K. Noble, 1916. 64038 was exchanged to UIMNH in 1956.

Lepidodactylus planicaudus Stejneger

Proc. U.S. Nat. Mus., vol. 28, p. 348, 1905.

HOLOTYPE: 34746, Mount Apo, Mindanao, Philippine Islands, 4,000-6,000 ft., E. A. Mearns, June-July, 1904.

Lepidophyma smithii oculator Smith

Proc. U.S. Nat. Mus., vol. 92, p. 378, 1942.

HOLOTYPE: 47133, Jalpan, Querétaro, Mexico, 2,000 ft., E. W. Nelson and E. A. Goldman, Aug. 30, 1896.

PARATYPES (2): 47134-5, same data.

Lepidophyma smithii tehuanae Smith

Proc. U.S. Nat. Mus., vol. 92, p. 377, 1942.

=*Lepidophyma flavimaculatum tehuanae* Smith

HOLOTYPE: 111488, Cerro Arenal, Oaxaca, Mexico, H. M. Smith, Jan. 19, 1940.

PARATYPES (9): 46687, Santa Efigenia, Oaxaca, Mexico, E. W. Nelson and E. A. Goldman, July 30, 1895. 46998-47000, near Tehuantepec, Oaxaca, Mexico, E. W. Nelson and E. A. Goldman, May 24, 1895. 48105, mountains near Santo Domingo, Oaxaca, Mexico, E. W. Nelson and E. A. Goldman, June 11, 1895. 111489, Tres Cruces, Oaxaca, Mexico, H. M. Smith, May 1940. 111490, El Limón, Oaxaca, Mexico, H. M. Smith, May 1940. 111491-2, La Concepción, Oaxaca, Mexico, H. M. Smith, May 1940. 46997 was exchanged to UIMNH in 1956.

Lepidophyma tuxtlae Werler and Shannon

Herpetologica, vol. 13, p. 119, fig. 1, 1957.

PARATYPE: 139731 (formerly UIMNH 40286), Rancho El Tular, 15 miles north of San Andrés Tuxtla, Veracruz, Mexico, D. A. Langebartel, Dec. 1, 1955.

Lepidosternon floridanum Baird

Proc. Acad. Nat. Sci. Philadelphia, vol. 10 (1858), p. 255, 1859.

=*Rhineura floridana* (Baird)

HOLOTYPE: 3202, Micanopy, Alachua County, Florida, J. B. Barratt.

Liocephalus eremitus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 20, p. 122, 1868.

=*Leiocephalus eremitus* Cope

HOLOTYPE: 12016, Navassa Island, West Indies, W. J. Rasin.

Liocephalus loxogrammus Cope

Proc. U.S. Nat. Mus., vol. 10, p. 437, 1887.

=*Leiocephalus loxogrammus* Cope

COTYPES (10): 14569 (10), Rum Cay, Bahama Islands, C. H. Townsend, March 1886. One cotype was exchanged to MCZ in 1915.

Liocephalus macropus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 184, 1862.

=*Leiocephalus macropus* Cope

COTYPES (11): 12254 (4), 25819-23, 25825-6, eastern Cuba (restricted to Monte Verde, near Guantanamo, Cuba), C. Wright. 25824 was exchanged to MCZ in 1915.

Liocephalus melanochlorus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 184, 1862.

=*Leiocephalus melanochlorus* Cope

COTYPES 52402 (formerly MCZ 3598), Jérémie, Haiti, D. F. Weinland.

Liocephalus raviceps Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 183, 1862.

=*Leiocephalus raviceps* Cope

COTYPES (8): 4162 (8), eastern Cuba, C. Wright. One cotype was exchanged to MCZ in 1915.

Liocephalus varius Garman

Proc. Amer. Philos. Soc., vol. 24, p. 274, 1887.

=*Leiocephalus varius* Garman

COTYPE: 52405 (formerly MCZ 6023), Grand Cayman Island, West Indies, W. B. Richardson.

Liolaemus hatcheri Stejneger

in Report Princeton Univ. Exped. to Patagonia, 1896-1899, vol. 3, pt. 1, Zoology, p. 218, 1909.

HOLOTYPE: 36912, plains of Patagonia, Santa Cruz Territory, Argentina, J. B. Hatcher, 1898.

PARATYPES (5): 36907-11, same data.

Liolaemus walkeri Shreve

Journ. Washington Acad. Sci., vol. 28, p. 404, 1938.

PARATYPES (2): 120768-9, Llocllapampa, Department of Junín, Peru, W. F. Walker, Jr., June 26-July 3, 1936.

Luperosaurus macgregori Stejneger

Proc. U.S. Nat. Mus., vol. 33, p. 545, 1907.

HOLOTYPE: 36191, Calayan Island, Philippine Islands, R. C. McGregor.

Lygodactylus capensis mossambica Loveridge

Proc. Zool. Soc. London, p. 135, 1920.

=*Lygodactylus grotei grotei* Sternfeld

COTYPES (5): 62831-5, ♀, ♀, ♀, ♂, Lumbo, Mozambique, A. Loveridge, July 11, 1918.

Lygodactylus manni Loveridge

Proc. U.S. Nat. Mus., vol. 72, No. 2720, p. 1, pl. 1, 1928.

HOLOTYPE: 72760, ♂, Saranda, Dodoma District, Tanganyika, A. Loveridge, July 15, 1926.

Lygosaurus pellopleurus Hallowell

Proc. Acad. Nat. Sci. Philadelphia, vol. 12 (1860), p. 496, 1861.

COTYPES (2): 42110, 42114, Amami-o-shima, Riu-Kiu (Ryukyu) Islands, W. Stimpson, 1855.

Lygosaurus sowerbyi Stejneger

Occ. Pap. Boston Soc. Nat. Hist., vol. 5, p. 120, 1924.

HOLOTYPE: 65375, Fukien, Futsin District, China, A. de C. Sowerby.

Lygosoma assatum ixbaac Stuart

Occ. Pap. Mus. Zool. Univ. Michigan, No. 421, p. 8, 1940.

=*Scincella cherriei ixbaac* (Stuart)

PARATYPE: 71379, San Miguel, El Petén, Guatemala, H. Malleis, Apr. 22, 1923.

Lygosoma cherriei stuarti Smith

Proc. Biol. Soc. Washington, vol. 54, p. 181, 1941.

=*Scincella cherriei stuarti* (Smith)

HOLOTYPE: 115174, Potrero Viejo, Veracruz, Mexico, H. M. Smith, Dec. 17, 1938.

PARATYPES (12): 115175-85, Potrero Viejo, Veracruz, Mexico, H. M. Smith, Dec. 13-26, 1938. 115187, Cuautlapan, Veracruz, Mexico, H. M. Smith, Jan. 3, 1939. 115186 was exchanged to UIMNH in 1956.

Lygosoma incertum Stuart

Occ. Pap. Mus. Zool. Univ. Michigan, No. 421, p. 10, 1940.

PARATYPE: 25224, vicinity of Guatemala, Guatemala, C. H. Van Patten.

Lygosoma kilimensis Stejneger

Proc. U.S. Nat. Mus., vol. 14, p. 405, 1891.

HOLOTYPE: 16749, Kilimanjaro, East Africa (Tanganyika), W. L. Abbott, 1889.

Lygosoma (Sphenomorphus) variegatum stickeli Loveridge

Bull. Mus. Comp. Zool., vol. 101, p. 345, 1948.

=*Sphenomorphus variegatus stickeli* Loveridge

PARATYPES (9): 119450-1, 119453, 119456-60, Gusiko, Australian New Guinea, W. H. Stickel, Apr. 5-30, 1944. 120106, Draeger Harbor, 5 miles east of Finschhafen, Australian New Guinea, G. H. Penn, Jr., 1944. 119454-5 were exchanged to MCZ in 1947.

Lysoptychus lateralis Cope

Proc. U.S. Nat. Mus., vol. 11, p. 397, pl. 36, figs. 1a-g, 1889.

=*Sceloporus couchii* Baird

HOLOTYPE: 14741, San Diego, Duval County, Texas, W. Taylor.

Mabuia alliacea Cope

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8 (1875), p. 115, pl. 28, fig. 1, 1876.

=*Mabuya mabouya mabouya* (Lacépède)

COTYPES (2): 30619-20, Costa Rica, W. M. Gabb.

Mabuia dorsivittata Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 350, 1862.

=*Mabuya dorsivittata* Cope

HOLOTYPE: 5405, Río Vermejo, Paraguay, T. J. Page.

Mabuia fulgida Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 185, 1862.

=*Mabuya spilonota* (Wiegmann)

COTYPES (6) : 5759 (6), Jamaica, C. B. Adams.

Mabuia lanceolata Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 187, 1862.

=*Mabuya lanceolata* Cope

COTYPES (2) : 6041 (2), Barbados (West Indies), T. N. Gill.

Mabuya bayonii keniensis Loveridge

Mus. Comp. Zool., Breviora, No. 59, p. 2, 1956.

PARATYPES (5) : 40710, west side of Mount Kenya, Kenya, J. A. Loring, Sept. 22, 1909. 40781, Wambugu, Kenya, J. A. Loring, Sept. 13, 1909. 40947-8, Sotik, Guaso Nyiro, Kenya, E. Heller, June 20, 1909. 42024, Sirgoit Lake?, Kenya, Smithsonian African Expedition, 1909.

Mabuya chanleri Stejneger

Proc. U.S. Nat. Mus., vol. 16, p. 721, 1894.

HOLOTYPE: 20104, Tana River, British East Africa (Kenya), W. A. Chanler, November 1892.

Mabuya deserticola Dunn

Proc. Acad. Nat. Sci. Philadelphia, vol. 37, p. 554, 1936.

=*Mabuya nigropalmata* Andersson

HOLOTYPE: 13826, Mollendo, Peru, W. H. Jones, Sept. 28, 1883.

Mabuya pergravis Barbour

Proc. New England Zool. Club, vol. 7, p. 85, 1921.

=*Mabuya mabouya pergravis* Barbour

HOLOTYPE: 13875, Old Providence Island (Colombia), U.S. Fish Commission *Albatross*, Apr. 4-9, 1884.

PARATYPES (2) : 63394-5, same data. 63396 was exchanged to MCZ in 1921.

Microlophus spinulosus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 351, 1862.

=*Tropidurus spinulosus* (Cope)

HOLOTYPE: 5956, Paraguay, T. J. Page.

Neoseps reynoldsi Stejneger

Proc. U.S. Nat. Mus., vol. 39, p. 34, figs. 1-6, 1910.

HOLOTYPE: 42147, near Spring Lake, Fruitland Park, Lake County, Florida, A. G. Reynolds, March 1910.

Oligosoma gemmingeri Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 16, p. 180, 1864.

=*Scincella gemmingeri gemmingeri* (Cope)

COTYPES (4): 6331 (4), Orizaba, Veracruz, Mexico, F. Sumichrast.

Ophisaurus attenuatus longicaudus McConkey

Amer. Midl. Nat., vol. 51, p. 151, pl. 1, center, bottom, 1954.

PARATYPES (22): 4169 (2), Knoxville, Knox County, Tennessee, J. B. Mitchell. 5038a, Georgia, W. L. Jones. 5322, Mississippi, B. F. Shumard. 8978, Florida. 10336, Clearwater, Pinellas County, Florida, S. T. Walker, July 1879. 12004a, Georgiana, Brevard County, Florida, W. Wittfield. 13383, Mount Mourne, Iredell County, North Carolina, J. A. D. Stephenson. 14724, Rockwood, Roane County, Tennessee, J. A. Sewell, Sept. 17, 1887. 18024, Greenville, Greenville County, South Carolina, J. C. Fitzgerald, 1891. 23972, New Smyrna, Volusia County, Florida, J. Y. Detwiler. 24343-4, Eustis, Lake County, Florida, H. J. Webber. 29618, Cleveland, Bradley County, Tennessee, J. D. Patton. 32530, Aiken, Aiken County, South Carolina, O. N. Cripps, October 1903. 38508, 3 miles west of Ashland, Hanover County, Virginia, F. B. Sheldon. 39043, Chuckatuck, Nansemond County, Virginia, J. S. Webb. 52212, forked tail only, Dunedin, Pinellas County, Florida, A. L. Holmes, February 1915. 61038, Gulfport, Pinellas County, Florida, A. G. Reynolds. 67596, Smithsonia, Lauderdale County, Alabama, R. R. Banks, September 1924. 69652, Eustis, Lake County, Florida, J. E. Brown, Dec. 10, 1922. 91413, Jedburg, Dorchester County, South Carolina, L. Brodie, May 29, 1932.

Ophisaurus ventralis compressus Cope

Report of the U.S. National Museum . . . for the year ending June 30, 1898, p. 501, fig. 90, 1900.

=*Ophisaurus compressus* Cope

HOLOTYPE: 21359, South Island, Georgetown County, South Carolina, E. P. Alexander, October 1893.

Oreosaurus anomalus Barbour and Noble

Proc. U.S. Nat. Mus., vol. 58, p. 614, 1921.

HOLOTYPE: 60704, ♂, Río San Miguel, San Fernando, Peru, 4,500 ft., E. Heller, Oct. 15, 1915.

Oreosaurus lacertus Stejneger

Proc. U.S. Nat. Mus. vol. 45, p. 546, 1913.

HOLOTYPE: 49551, Tincochchaca (Tinccoachaca), Peru, 7,000 ft., Yale-Peruvian Expedition, Aug. 10, 1911.

PARATYPES (2): 49549, Ollantaytambo, Peru, 9,000 ft., Yale-Peruvian Expedition, July 20, 1911. 49552, Tincochchaca (Tinccoachaca), Peru, 7,000 ft., Yale-Peruvian Expedition, Aug. 8, 1911.

Pantodactylus bivittatus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 15, p. 103, 1863.

=*Pantodactylus schreibersii* (Gray)

COTYPES (2) : 31279-80, Paysandú, Uruguay, H. W. Kennedy.

Phelsuma abbotti Stejneger

Proc. U.S. Nat. Mus., vol. 16, p. 716, 1894.

=*Phelsuma madagascariensis abbotti* Stejneger

HOLOTYPE: 20467, ♂, Aldabra Island (Seychelles Islands, Indian Ocean), W. L. Abbott, 1892.

PARATYPES (2) : 20468-9, same data.

Phoxophrys grahami Stejneger

Occ. Pap. Boston Soc. Nat. Hist., vol. 5, p. 120, 1924.

HOLOTYPE: 65500, Suifu, Szechwan, China, D. C. Graham.

Phrynocephalus erythrurus Zugmayer

Zool. Jahrb. Syst., vol. 27, p. 504, 1909.

PARATYPES (2) : 73854-5, West Tibet, China, E. Zugmayer, 1906.

Phrynosoma asio Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 16, p. 178, 1864.

PROBABLE COTYPES (2) : 32216-7, Colima, Mexico, J. Xantus.

Phrynosoma cerroense Stejneger

North Amer. Fauna, No. 7, p. 187, 1893.

HOLOTYPE: 11977, Cedros Island (given as "Cerros" Island in the original description), Baja California, Mexico, L. Belding.

Phrynosoma ditmarsii Stejneger

Proc. U.S. Nat. Mus., vol. 29, p. 565, 1906.

HOLOTYPE: 36022, ♂, State of Sonora, Mexico, near the boundary of Arizona, Eustace, 1897.

PARATYPE: 36013, ♀, same data.

Phrynosoma douglassii brachycercum Smith

Proc. U.S. Nat. Mus., vol. 92, p. 362, 1942.

HOLOTYPE: 23993, Durango, Mexico, E. Palmer, September 1896.

PARATYPES (3) : 23994-6, same data.

Phrynosoma douglassi pygmaea Yarrow

Proc. U. S. Nat. Mus., vol. 5, p. 443, 1882.

=*Phrynosoma douglassii douglassii* (Bell)

COTYPES (13) : 9199 (3), Fort Steilacoom, Puget Sound (Pierce County), Washington. 10918 (5), Fort Walla Walla (Walla Walla County), Wash-

ington, C. E. Bendire, June 1881. 11473 (2), Deschutes River, Oregon, H. W. Henshaw, Sept. 5, 1878. 11945 (3), Oregon, H. W. Henshaw, September 1878.

Phrynosoma goodei Stejneger

North Amer. Fauna, No. 7, p. 191, pl. 2, figs. 3a-c, 1893.

=*Phrynosoma platyrhinos goodei* Stejneger

HOLOTYPE: 8567A, west coast of Sonora, Mexico (restricted to Puerto Libertad), T. H. Streets.

Phrynosoma jamesi Schmidt

Bull. Amer. Mus. Nat. Hist., vol. 46, p. 668, pls. 55-56, 1922.

=*Phrynosoma coronatum jamesi* Schmidt

HOLOTYPE: 64450, ♀, San Bartolomé Bay, Baja California, Mexico, C. H. Townsend, Mar. 14, 1911.

Phrynosoma modestum Girard

in Stansbury, Report Expl. and Surv. Valley of the Great Salt Lake of Utah . . . , p. 361, pl. 6, figs. 4-8, 1852.

COTYPES (7): 164 (7), "between San Antonio and El Paso," Texas (restricted to Las Cruces, New Mexico), J. H. Clark. One cotype was exchanged to UIMNH in 1956.

Phrynosoma nelsoni Schmidt

Bull. Amer. Mus. Nat. Hist., vol. 46, p. 666, pl. 54, fig. 3, 1922.

=*Phrynosoma coronatum frontale* Van Denburgh

HOLOTYPE: 37585, ♀, San Quintín, Baja California, Mexico, E. W. Nelson and E. A. Goldman, August 15, 1905.

PARATYPE: 37584, ♂, same data.

Phrynosoma orbiculare durangoensis Horowitz

Amer. Midl. Nat., vol. 54, p. 211, 1955.

PARATYPES (3): 47469, El Salto, Durango, Mexico, E. W. Nelson and E. A. Goldman, July 20, 1895. 47872, near Bolaños, Jalisco, Mexico, E. W. Nelson and E. A. Goldman, Sept. 16, 1897. 47873, Sierra Madre, Zacatecas, Mexico, 8,000 ft., E. W. Nelson and E. A. Goldman, Aug. 16, 1897.

Phrynosoma planiceps Hallowell

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 178, 1852.

=*Phrynosoma cornutum* (Harlan)

COTYPE: 143, 90 miles below El Paso near the Rio Grande, Texas, S. W. Woodhouse.

Phrynosoma platyrhinos Girard

in Stansbury, Report Expl. and Surv. of the Valley of the Great Salt Lake of Utah . . . , p. 361, pl. 7, figs. 1-5, 1852.

=*Phrynosoma platyrhinos platyrhinos* Girard

COTYPES (3): 189 (3), Salt Lake Valley, Utah, H. Stansbury. One cotype exchanged to MCZ in 1879.

Phrynosoma regale Girard

in U.S. Exploring Expedition . . . 1838-42, vol. 20 (Herpetology), p. 406, 1858.

=*Phrynosoma solaris* Gray

COTYPES (2): 161 (2), Sierra de la Nariz, near Zuni, Sonora, Mexico, A. Schott.

Phyllodactylus homolepidurus Smith

Univ. Kansas Sci. Bull., vol. 22, p. 121, pl. 25, fig. 2, text-fig. 1a, 1935.

PARATYPE: 134302 (formerly UIMNH 27482), 5 miles southwest of Hermosillo, Sonora, Mexico, E. H. Taylor, July 8, 1934.

Phyllodactylus lanei Smith

Univ. Kansas Sci. Bull., vol. 22, p. 125, text-fig. 1b, pl. 25, fig. 3, 1935.

PARATYPE: 138887 (formerly UIMNH 19612), near Tierra Colorada, Guerrero, Mexico, E. H. Taylor and H. M. Smith, June 30, 1932.

Phyllodactylus leei Cope

Proc. U.S. Nat. Mus., vol. 12, p. 145, 1890.

HOLOTYPE: 14957, Chatham Island, Galápagos Islands (off the coast of Ecuador), U.S. Fish Commission *Albatross*, Apr. 4, 1888.

Phyllodactylus magister Noble

Occ. Pap. Boston Soc. Nat. Hist., vol. 5, p. 110, 1924.

PARATYPES (4): 75946-9, Bellavista, Peru, G. K. Noble, 1916. 75950 was exchanged to UIMNH in 1956.

Phyllodactylus magnus Taylor

Univ. Kansas Sci. Bull., vol. 28, p. 99, fig. 3, 1942.

PARATYPES (39): 115703-4, 115725-37, Cerro Arenal, Oaxaca, Mexico, H. M. Smith, Jan. 2, 19, 1940. 115706 (given in error as 115760 in original description), 115713-7, 115722, Tres Cruces, Oaxaca, Mexico, H. M. Smith, Jan. 14-15, 1940. 115707, Tehuantepec, Oaxaca, Mexico, H. M. Smith, Jan. 16, 1940. 115740-2, 115744-7, 115749-50, Cajón de Piedra, Oaxaca, Mexico, H. M. Smith, May 1940. 115751-6, Tonalá, Chiapas, Mexico, H. M. Smith, Jan. 26-30, 1940. 137210, Tierra Colorada, Guerrero, Mexico, 1939.

Phyllodactylus spatulatus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 176, 1862.

=*Phyllodactylus pulcher* Gray

COTYPES (3): 6040 (3), Barbados (West Indies), T. N. Gill.

Phyllodactylus wirshingi Kerster and Smith

Herpetologica, vol. 11, p. 229, fig. 1, pl. 1, 1955.

PARATYPE: 138899 (formerly UIMNH 37743), Caja de Muertos Islands, Puerto Rico, J. Wirshing, Mar. 10, 1953.

Platydactylus muralis Duméril and Bibron

Erpétologie générale ou histoire naturelle complète des reptiles, vol. 3, p. 319, 1836.

=*Tarentola mauritanica* (Linnaeus)

COTYPE: 10996, Nice, France, JPP.

Plestiodon anthracinus Baird

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 18, p. 294, 1849.

=*Eumeces anthracinus anthracinus* (Baird)

COTYPES (5): 3138 (5), Carlisle, Cumberland County, Pennsylvania, S. F. Baird.

Plestiodon egregius Baird

Proc. Acad. Nat. Sci. Philadelphia, vol. 10 (1858), p. 256, 1859.

=*Eumeces egregius* (Baird)

COTYPES (2): 3127 (2), Indian Key (Pinellas County), Florida, G. Würdemann.

Plestiodon inornatus Baird

Proc. Acad. Nat. Sci. Philadelphia, vol. 10 (1858), p. 256, 1859.

=*Eumeces multivirgatus* Hallowell

COTYPES (2): 3110 (2), sand hills of Platte, Nebraska, F. V. Hayden.

Plestiodon longirostris Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 13, p. 313, 1861.

=*Eumeces longirostris* (Cope)

COTYPES (5): 4737 (5), Bermuda, J. H. Darrell. One cotype was exchanged to UIMNH in 1956.

Plestiodon marginatus Hallowell

Proc. Acad. Nat. Sci. Philadelphia, vol. 12 (1860), p. 492, 1861.

=*Eumeces marginatus* (Hallowell)

LECTOTYPE: 11713, Okinawa (Ryukyu Islands), W. Stimpson, April 1855.

Plestiodon obsoletum Baird and Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 129, 1852.

=*Eumeces obsoletus* (Baird and Girard)

HOLOTYPE: 3133, Valley of the Rio San Pedro of the Rio Grande del Norte (now Devils River), Texas, J. H. Clark.



Plestiodon septentrionalis Baird

Proc. Acad. Nat. Sci. Philadelphia, vol. 10 (1858), p. 256, 1859.

=*Eumeces septentrionalis* (Baird)

COTYPES (3): 3156 (3), Fort Ripley (Crow Wing County), Minnesota, J. F. Head.

Plestiodon skiltonianum Baird and Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 69, 1852.

=*Eumeces skiltonianus skiltonianus* (Baird and Girard)

COTYPES (2): 3172a-b, Oregon (restricted to the Dalles, Wasco County), G. R. Geary. 3172a is the lectotype.

Plestiodon tetragrammus Baird

Proc. Acad. Nat. Sci. Philadelphia, vol. 10 (1858), p. 256, 1859.

=*Eumeces tetragrammus* (Baird)

COTYPES (10): 3124 (10), Matamoros, Tamaulipas, Mexico, D. N. Couch. The lectotype is 3124a. One cotype was exchanged to UIMNH in 1956.

Plica stejnegeri Burt and Burt

Proc. U.S. Nat. Mus., vol. 87, No. 2849, p. 19, 1930.

HOLOTYPE: 73505, Argentina, A. and A. Breyer.

Plistodon sumichrasti Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 18 (1866), p. 321, 1867.

=*Eumeces sumichrasti* (Cope)

HOLOTYPE: 6601, Potrero, Veracruz, Mexico ("Orizava" in original description), F. Sumichrast.

Prionodactylus marianus Ruthven

Occ. Pap. Mus. Zool. Univ. Michigan, No. 103, p. 1, 1921.

=*Euspondylus marianus* (Ruthven)

PARATYPE: 120789 (formerly MCZ 14660), San Pedro, near Medellín, Colombia, H. Nicéforo-María.

Proctoporus obesus Barbour and Noble

Proc. U.S. Nat. Mus., vol. 58, p. 616, 1921.

HOLOTYPE: 60748, ♂, Ñusta Hispana, Peru, E. C. Erdis, June 11, 1915.

Pterogasterus modestus Cope

Proc. Amer. Philos. Soc., vol. 17 (1878), p. 97, 1877.

=*Barisia modesta* (Cope)

COTYPES (3): 7084 (3), alpine regions, Orizaba, Veracruz, Mexico (given as "probably Guatemala" in original description, but restricted to Mount Orizaba), F. Sumichrast.

Riopa singha Taylor

Univ. Kansas Sci. Bull., vol. 33, p. 494, figs. 2a-b, 1950.

HOLOTYPE: 29411, Trincomalee, Ceylon, D. L. Karcher.

Saccodeira ornatissima Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 9 (1857), p. 198, 1858.

COTYPE: 5655, Yanga, Peru, U.S. Exploring Expedition, 1839.

Sator angustus Dickerson

Bull. Amer. Mus. Nat. Hist., vol. 41, p. 469, 1919.

HOLOTYPE: 64262, ♂, Santa Cruz Island, Gulf of California, Baja California, Mexico, C. H. Townsend, Apr. 17, 1911.

Sator grandaevus Dickerson

Bull. Amer. Mus. Nat. Hist., vol. 41, p. 469, 1919.

HOLOTYPE: 64261, ♂, Cerralvo Island, Gulf of California, Baja California, Mexico, C. H. Townsend, Apr. 19, 1911.

Sauromalus hispidus Stejneger

Proc. U.S. Nat. Mus., vol. 14, p. 409, 1891.

HOLOTYPE: 8563, Ángel de la Guarda Island, Gulf of California, Baja California, Mexico, T. H. Streets.

Sauromalus interbrachialis Dickerson

Bull. Amer. Mus. Nat. Hist., vol. 41, p. 463, 1919.

=*Sauromalus ater* Duméril

HOLOTYPE: 64443, La Paz, Baja California, Mexico, C. H. Townsend, Mar. 27, 1911. The locality is said to be incorrect by Shaw, Trans. San Diego Soc. Nat. Hist., vol. 10, pp. 274-5, 1945.

Sauromalus townsendi Dickerson

Bull. Amer. Mus. Nat. Hist., vol. 41, p. 464, 1919.

=*Sauromalus obesus townsendi* Dickerson

HOLOTYPE: 64442, ♂, Tiburón Island, Gulf of California, Sonora, Mexico, C. H. Townsend, Apr. 12, 1911.

Sauromalus varius Dickerson

Bull. Amer. Mus. Nat. Hist., vol. 41, p. 464, 1919.

HOLOTYPE: 64441, San Estéban Island, Gulf of California, Sonora, Mexico, C. H. Townsend, Apr. 13, 1911.

Scartiscus caducus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 182, 1862.

=*Leiocephalus caducus* (Cope)

HOLOTYPE: 5852, Paraguay, T. J. Page.

Sceloporus aeneus bicanthalis Smith

Occ. Pap. Mus. Zool. Univ. Michigan, No. 361, p. 6, 1937.

PARATYPES (2): 47280, Popocatepetl, Puebla, Mexico, E. W. Nelson, Feb. 27, 1893. 47395, Reyes, Oaxaca, Mexico, 10,200 ft., E. W. Nelson and E. A. Goldman, Oct. 22, 1894.

Sceloporus aeneus subniger Poglayen and Smith

Herpetologica, vol. 14, p. 13, 1958.

PARATYPES (25): 111563-87, 15 km. west of Toluca, Mexico, Mexico, H. M. Smith, Mar. 5, 1939.

Sceloporus boulengeri Stejneger

North Amer. Fauna, No. 7, p. 180, pl. 1, figs. 5a-c, 1893.

= *Sceloporus clarkii boulengeri* Stejneger

COTYPES (3): 14079 (3), Presidio, about 50 miles south of Mazatlán, Sinaloa, Mexico, A. Forrer, Apr. 3, 1885.

Sceloporus chrysostictus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 18, p. 125, 1866.

COTYPES (2): 24865-6, Yucatán, Mexico, A. Schott.

Sceloporus clarkii Baird and Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 127, 1852.

= *Sceloporus clarkii clarkii* Baird and Girard

LECTOTYPE: 2940, "Province of Sonora" (restricted to Santa Rita Mountains, Arizona), J. H. Clark. Two other specimens also listed as cotypes of *Sceloporus clarkii* are said to be examples of *Sceloporus magister magister* by Smith and Taylor, U.S. Nat. Mus. Bull. 199, p. 113, 1950.

Sceloporus cochranæ Smith

Proc. Biol. Soc. Washington, vol. 49, p. 87, pl. 2, fig. 1, pl. 3, 1936.

= *Sceloporus siniferus cupreus* Bocourt

HOLOTYPE: 47605, Mount Zempoaltepec, 8,000 ft., Oaxaca, Mexico, E. W. Nelson and E. A. Goldman, July 8, 1894.

Sceloporus couchii Baird

Proc. Acad. Nat. Sci. Philadelphia, vol. 10 (1858), p. 254, 1859.

COTYPES (9): 2739 (9), Santa Caterina, Nuevo León, Mexico, D. N. Couch, April 1853.

Sceloporus cozumelæ Jones

Occ. Pap. Mus. Zool. Univ. Michigan, No. 186, p. 1, 1927.

HOLOTYPE: 13904, Cozumel Island, Yucatán, Mexico, U.S. Fish Commission *Albatross*, Jan. 23-29, 1885.

PARATYPES (3): 17827-9, Cozumel Island, off Yucatán, Mexico, U.S. Fish Commission *Albatross*, 1885.

Sceloporus delicatissimus Hallowell

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 178, 1852.

= *Sceloporus variabilis marmoratus* Hallowell

HOLOTYPE: 16020, San Antonio, Bexar County, Texas, S. W. Woodhouse.

Sceloporus disparilis Stejneger

Proc. Biol. Soc. Washington, vol. 29, p. 228, 1916.

=*Sceloporus grammicus disparilis* Stejneger

HOLOTYPE: 33041, ♂, Lomita Ranch, 6 miles north of Hidalgo, Hidalgo County, Texas, W. Lloyd, June 17, 1891.

PARATYPES (4): 33042-5, Lomita Ranch, 6 miles north of Hidalgo, Hidalgo County, Texas, W. Lloyd, June 13-17, 1891.

Sceloporus edwardtaylori Smith

Herpetologica, vol. 1, p. 6, 1936.

PARATYPE: 138890 (formerly UIMNH 27256), near Totalpam, Oaxaca, Mexico, H. M. Smith, Aug. 7, 1935.

Sceloporus elongatus Stejneger

North Amer. Fauna, No. 3, p. 111, 1890.

=*Sceloporus undulatus elongatus* Stejneger

HOLOTYPE: 15858, ♂, Moa Ave, Painted Desert, Arizona, C. H. Merriam and V. O. Bailey, Sept. 23, 1889.

PARATYPES (3): 15856-7, ♀ and juvenile, Little Colorado River, Painted Desert, Arizona, C. H. Merriam and V. O. Bailey, Sept. 21, 26, 1889. 15859, ♀, Moa Ave, Painted Desert, Arizona, C. H. Merriam and V. O. Bailey, Sept. 23, 1889.

Sceloporus ferrariperezi Cope

Proc. Amer. Philos. Soc., vol. 22, p. 400, 1885.

=*Sceloporus torquatus torquatus* Wiegmann

COTYPES (4): 9874, 9876, 9878, 9880, Guanajuato, Mexico, A. Dugès, 1877. 9895 was exchanged to MCZ in 1944.

Sceloporus floridanus Baird

Proc. Acad. Nat. Sci. Philadelphia, vol. 10 (1853), p. 254, 1859.

=*Sceloporus undulatus undulatus* (Latreille)

HOLOTYPE: 2874, Pensacola, Escambia County, Florida, R. W. Jeffries.

Sceloporus formosus scitulus Smith

Proc. U.S. Nat. Mus., vol. 92, p. 352, 1942.

PARATYPES (4): 47738, Omilteme, Guerrero, Mexico, E. W. Nelson and E. A. Goldman, May 20, 1903. 111827-8, Omilteme, Guerrero, Mexico, R. C. and E. H. Taylor, Aug. 2-4, 1940. 134303, Omilteme, Guerrero, Mexico, E. H. Taylor, Aug. 4-6, 1940.

Sceloporus goldmani Smith

Occ. Pap. Mus. Zool. Univ. Michigan, No. 361, p. 5, 1937.

PARATYPE: 46869 (given incorrectly as 46839 in original description), Carneros, Coahuila, Mexico, 6,800 ft., E. W. Nelson and E. A. Goldman, Aug. 12, 1896.

Sceloporus graciosus Baird and Girard

in Stansbury, Report Expl. and Surv. Valley of the Great Salt Lake of Utah . . . , p. 346, pl. 5, figs. 1-3c, 1852.

=*Sceloporus graciosus graciosus* Baird and Girard

COTYPES (4): 2877 (4), Valley of the Great Salt Lake, Utah, H. Stansbury.

Sceloporus heterurus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 18 (1866), p. 322, 1867.

=*Sceloporus grammicus microlepidotus* Wiegmann

PROBABLE HOLOTYPE: 25064 (given incorrectly in the original description as 6539), Mirado, Veracruz, Mexico, C. Sartorius.

Sceloporus horridus albiventris Smith

Field Mus. Nat. Hist., Zool. Ser., vol. 26, p. 108, pl. 12, 1939.

PARATYPE: 137214, Mount San Juan near Tepic, Nayarit, Mexico, E. H. Taylor, July 29, 1934.

Sceloporus jarrovii Cope

in Yarrow, Report upon the collections of batrachians and reptiles . . . 1871, 1872, 1873, and 1874, in Report U.S. Geogr. and Geol. Expl. and Surv. West of One Hundr. Mer., vol. 5, p. 569, pl. 23, figs. 2-2d, 1875.

=*Sceloporus jarrovii jarrovii* Cope

COTYPES (2): 8495, 8611, southern Arizona, H. W. Henshaw, October 1873.

Sceloporus jarrovii immucronatus Smith

Copeia, No. 4 (1936), p. 223, 1937.

PARATYPE: 137215, 25 miles south of Jacala, Mexico, Mexico, E. H. Taylor and H. M. Smith, June 16, 1932.

Sceloporus jarrovii oberon Smith and Brown

Field Mus. Nat. Hist., Zool. Ser., vol. 24, p. 254, fig. 24, 1941.

HOLOTYPE: 105823, Arteaga, Coahuila, Mexico, H. M. and R. B. Smith, Nov. 16, 1938.

PARATYPES (139): 105606-744, Mount Zapaliname, Diamond Pass, near Saltillo, Coahuila, Mexico, 7,800 ft., H. M. Smith, Nov. 10-15, 1938.

Sceloporus jarrovii sugillatus Smith

Proc. U.S. Nat. Mus., vol. 92, p. 357, 1942.

HOLOTYPE: 112100, ♂, Lake No. 4, Zempoala, Morelos, Mexico, 10,000 ft., H. M. Smith, Feb. 22, 1939.

PARATYPES (39): 112072-99, 112101-11, Zempoala, Morelos, Mexico, H. M. Smith, Feb. 21-22, 1939.

Sceloporus licki Van Denburgh

Proc. California Acad. Sci., Ser. 2, vol. 5, p. 110, pl. 10, 1895.

—*Sceloporus orcutti licki* Van Denburgh

PARATYPE: 23749, Miraflores, Baja California, Mexico, G. Eisen and F. H. Vaslit, September 1894.

Sceloporus lineatulus Dickerson

Bull. Amer. Mus. Nat. Hist., vol. 41, p. 467, 1919.

—*Sceloporus magister lineatulus* Dickerson

HOLOTYPE: 64263, ♂, Santa Catalina Island, Gulf of California, Baja California, Mexico, C. H. Townsend, May 16, 1911.

Sceloporus lineolateralis Smith

Proc. Biol. Soc. Washington, vol. 49, p. 92, 1936.

PARATYPE: 138891 (formerly UIMNH 21322), 14 miles northeast of Pedriceña, Durango, Mexico, H. M. Smith and E. H. Taylor, Aug. 29, 1932.

Sceloporus longipes Baird

Proc. Acad. Nat. Sci., Philadelphia, vol. 10 (1858), p. 254, 1859.

—*Sceloporus occidentalis longipes* Baird

COTYPES (25): 4358 (25), Fort Tejon (Kern County), California, J. Xantus. 4358a is the lectotype. One cotype was exchanged to UIMNH in 1956; one to MZUM.

Sceloporus lundelli lundelli Smith

Field Mus. Nat. Hist., Zool. Ser., vol. 26, p. 66, pl. 4 (left), 1939.

PARATYPES (2): 46862, Jaina, Campeche, Mexico, E. W. Nelson and E. A. Goldman, June 15, 1900. 138892 (formerly UIMNH 27267), Balchacaj, Campeche, Mexico, H. M. Smith, Sept. 21–22, 1936.

Sceloporus macdougalli Smith and Bumzahem

Herpetologica, vol. 9, p. 186, fig. 1, pls. 1–2, 1953.

PARATYPE: 134304 (formerly UIMNH 34359), Rincón Bamba, Isthmus of Tehuantepec, Oaxaca, Mexico, T. MacDougall, Mar. 26–27, 1953.

Sceloporus magister Hallowell

Proc. Acad. Nat. Sci. Philadelphia, vol. 7, p. 93, 1854.

—*Sceloporus magister magister* Hallowell

HOLOTYPE: 2967, near Fort Yuma (Imperial County), California, A. L. Heermann.

Sceloporus magister bimaculosus Phelan and Brattstrom

Herpetologica, vol. 11, p. 9, 1955.

PARATYPES (10): 2964, between El Paso, Texas, and Janos, Chihuahua, Mexico, C. B. R. Kennerly, 1855. 22269, Las Cruces Agricultural College, Las Cruces, Dona Ana County, New Mexico, T. D. A. Cockerell. 22382,

Mesilla Valley, Dona Ana County, New Mexico, T. D. A. Cockerell. 32939-49, Boquillas, Brewster County, Texas, V. O. Bailey and H. C. Oberholser, May 23, 1901. 32942, Boquillas, Brewster County, Texas, H. C. Oberholser, May 28, 1901. 44592, Socorro Mountains, Socorro County, New Mexico, E. A. Goldman, Aug. 13, 1909. 56860-2, El Paso, El Paso County, Texas, J. Hurter, July 24-25, 1902.

Sceloporus magister uniformis Phelan and Brattstrom

Herpetologica, vol. 11, p. 7, 1955.

PARATYPES (35): 18096-101, Pahrana-gat Valley, Lincoln County, Nevada, C. H. Merriam and V. O. Bailey, May 23, 1891. 18102, ♀, Pahrana-gat Mountain, Lincoln County, Nevada, C. H. Merriam, May 26, 1891. 18103, ♂, Pah-rump Valley, Nye County, Nevada, C. H. Merriam, Apr. 29, 1891. 18104, ♂, Callville, Colorado River, Nevada, V. O. Bailey, May 4, 1891. 18105-6, ♂, ♂, Ash Meadows, Nye County, Nevada, E. W. Nelson and T. S. Palmer, Mar. 16, 20, 1891. 18107, ♂, Las Vegas Valley, Nevada, V. O. Bailey, May 1, 1891. 18108-9, ♂, ♂, Indian Springs Valley, Clark County, Nevada, C. H. Merriam and V. O. Bailey, May 29, 1891. 18111, ♂, Bunkerville, Clark County, Nevada, V. O. Bailey, May 8, 1891. 37659-60, Volcano Lake, Baja California, Mexico, E. W. Nelson and E. A. Goldman, Apr. 19-20, 1905. 37663, east base of Cocopah Mountains, Lower California, Mexico, E. W. Nelson, and E. A. Goldman, Apr. 11, 1905. 37664, San Felipe Bay, Lower California, Mexico, E. W. Nelson and E. A. Goldman, June 23, 1905. 44785, Pyramid Lake, Washoe County, Nevada, C. A. Keeler, June 1889. 44852, Mojave Valley, east of Morongo Valley, San Bernardino County, California, 3,100 ft., V. O. Bailey, Sept. 12, 1907. 45201-6, Pyramid Lake, Washoe County, Nevada, V. O. Bailey, June 23, 1893. 48691, Fallon, Churchill County, Nevada, V. O. Bailey, Sept. 6, 1911. 50808, 50810, near head of Pyramid Lake, Washoe County, Nevada, C. H. Richardson and J. O. Snyder. 50831-2, Pyramid Lake Indian Agency, Nevada, J. O. Snyder, May 31, 26, 1911. 51901, Fallon, Churchill County, Nevada, V. O. Bailey, Aug. 17, 1911. 87025, 4 miles west of Fallon, Churchill County, Nevada, C. E. Burt, Aug. 13, 1932. 118461, Fallon, Churchill County, Nevada, F. W. King, June 18, 1931.

Sceloporus malachiticus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 16, p. 178, 1864.

= *Sceloporus malachiticus malachiticus* Cope

COTYPE: 12186 (formerly 6492), near Arriba, Costa Rica, C. N. Riotte.

Sceloporus malachiticus internasalis Smith and Bumzahem

Herpetologica, vol. 11, p. 118, fig. 1, 1955.

PARATYPE: 135436, La Gloria, Oaxaca, Mexico, T. MacDougall, Mar. 12, 1953.

Sceloporus megalepidurus Smith

Trans. Kansas Acad. Sci., vol. 37, p. 272, pl. 9, figs. 7-8, pl. 10, fig. 13, 1934.

PARATYPE: 134305 (formerly UIMNH 27441), Pedregal, near Totalco, Veracruz, Mexico, E. H. Taylor, Sept. 2, 1936.

Sceloporus melanogaster Cope

Proc. Amer. Philos. Soc., vol. 22, p. 400, 1885.

=*Sceloporus torquatus melanogaster* Cope

HOLOTYPE: 9877, ♂, Mexico (restricted to Tupátaro, Guanajuato), A. Dugès, 1877.

Sceloporus melanorhinus calligaster Smith

Proc. U.S. Nat. Mus., vol. 92, p. 360, 1942.

HOLOTYPE: 112201, Acapulco, Guerrero, Mexico, H. M. Smith, Sept. 3, 1939.

PARATYPES (18): 24925-6, Guadalajara?, Jalisco, Mexico, J. J. Major 31496, Colima, Mexico, J. Xantus. 47731, Tamarindo, Guerrero, Mexico, E. W. Nelson and E. A. Goldman, Apr. 3, 1903. 47732, Acapulco, Guerrero, Mexico, E. W. Nelson and E. A. Goldman, Apr. 29, 1903. 51384-9, San Blas, Nayarit, Mexico, J. C. Thompson, Dec. 20, 25, 1913. 58159, Colima, Mexico, J. Hurter, July 14, 1902. 64667, San Blas, Nayarit, Mexico, E. W. Nelson and E. A. Goldman, June 11, 1897. 112186, near Coyuca, Guerrero, Mexico, H. M. Smith, Feb. 10, 1939. 112199-200, Acapulco, Guerrero, Mexico, H. M. Smith, Sept. 3, 1939. 112202-3, Apatzingán, Michoacán, Mexico, H. M. Smith, Mar. 16-17, 1939.

Sceloporus merriami Stejneger

Proc. Biol. Soc. Washington, vol. 17, p. 17, 1904.

=*Sceloporus merriami merriami* Stejneger

HOLOTYPE: 33039, ♂, East Painted Cave, near mouth of Pecos River, Texas, W. Lloyd, Sept. 2, 1890.

PARATYPES (7): 33033, ♂, Comstock, Val Verde County, Texas, M. Cary, July 26, 1902. 33034, ♀, Boquillas, Brewster County, Texas, V. Bailey, May 28, 1901. 33035, ♂, 55 miles northwest of Comstock, at mouth of Pecos River, Texas, M. Cary, Aug. 3, 1902. 33036-8, 33040, ♂, ♂, ♀, East Painted Cave, near mouth of Pecos River, Texas, W. Lloyd, Sept. 1-3, 1890.

Sceloporus merriami annulatus Smith

Proc. Biol. Soc. Washington, vol. 50, p. 83, 1937.

PARATYPE: 120770 (formerly MCZ 46524), Glenn Springs, Brewster County, Texas, H. T. Gaige.



Sceloporus mucronatus aureolus Smith

Proc. U.S. Nat. Mus., vol. 92, p. 356, 1942.

HOLOTYPE: 112232, ♂, 2 miles west of Acultzingo, Veracruz, Mexico, H. M. Smith, Jan. 17, 1939.

PARATYPES (26): 112233-42, 112244-58 from vicinity of Acultzingo, Veracruz, Mexico, H. M. Smith, January 1939 to August 1940. 112243, 8 km. east of Ozumbilla, Puebla, Mexico, H. M. Smith, Jan. 17, 1939.

Sceloporus nelsoni Cochran

Journ. Washington Acad. Sci., vol. 13, p. 185, 1923.

HOLOTYPE: 47676, Plomosas, Sinaloa, Mexico, E. W. Nelson and E. A. Goldman, July 18, 1897.

PARATYPES (8): 18979, Barranca Ibarra, Jalisco, Mexico, P. L. Jouy, Apr. 19, 1892. 46629, Culiacán, Sinaloa, Mexico, E. A. Goldman, Mar. 18, 1899. 47271, 47273-5, near Mazatlán, Sinaloa, Mexico, E. A. Goldman, Apr. 1-10, 1899. 47690-1, Rosario, Sinaloa, Mexico, E. W. Nelson and E. A. Goldman, July 21, 1897. 47272 was exchanged to MCZ in 1923; 47689 to MZUM, 1923.

Sceloporus ochoterena Smith

Trans. Kansas Acad. Sci., vol. 37, p. 269, pl. 9, figs. 6, 9; pl. 10, fig. 12, 1934.

PARATYPE: 138893 (formerly UIMNH 37949), 11 miles south of Puente de Ixtla, Guerrero, Mexico, H. M. Smith and E. H. Taylor, June 22, 1932.

Sceloporus oligoporus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 16, p. 177, 1864.

=*Sceloporus horridus oligoporus* Cope

COTYPES (6): 31386-90, 31393, Colima, Colima, Mexico, J. Xantus, March-July 1863. 31391 was exchanged to MCZ in 1944; 31392 to UIMNH, 1956.

Sceloporus orcutti Stejneger

North Amer. Fauna, No. 7, p. 181, pl. 1, figs. 4a-c, 1893.

=*Sceloporus orcutti orcutti* Stejneger

HOLOTYPE: 16330, Milquatay Valley, San Diego County, California, C. R. Orcutt, Jan. 5, 1890.

PARATYPES (7): 16328-9, 16332-6, same data. 16331 was exchanged to UIMNH in 1956.

Sceloporus ornatus Baird

Proc. Acad. Nat. Sci. Philadelphia, vol. 10 (1858), p. 254, 1859.

=*Sceloporus ornatus ornatus* Baird

HOLOTYPE: 2845, Patos, Coahuila, Mexico, D. N. Couch.

Sceloporus ornatus caeruleus Smith

Copeia, No. 4 (1936), p. 227, 1937.

PARATYPE: 138894 (formerly UIMNH 21453), 5 miles south of San Pedro, Coahuila, Mexico, H. M. Smith and D. H. Dunkle, June 29, 1934.

Sceloporus parvus Smith

Trans. Kansas Acad. Sci., vol. 37, p. 263, pl. 8, figs. 1, 3; pl. 10, fig. 10, 1934.

=*Sceloporus parvus parvus* Smith

PARATYPE: 138895 (formerly UIMNH 27419), near spring, Sabinas Hidalgo, Nuevo León, Mexico, H. M. Smith and E. H. Taylor, June 8, 1932.

Sceloporus poinsettii Baird and Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 126, 1852.

=*Sceloporus poinsettii poinsettii* Baird and Girard

COTYPES (5): 2948 (2), 131668, Rio San Pedro of the Rio Grande del Norte, Texas, J. H. Clark. 2952 (2), Province of Sonora, Mexico, J. H. Clark.

Sceloporus poinsetti macrolepis Smith and Chrapliwy

Herpetologica, vol. 13, p. 268, 1957.

=*Sceloporus poinsettii macrolepis* Smith and Chrapliwy

PARATYPES (6): 46844-5, Durango, Durango, Mexico, E. W. Nelson and E. A. Goldman, July 1, 1898. 47417, near Balleza, Chihuahua, Mexico, E. W. Nelson and E. A. Goldman, Sept. 23, 1899. 47419-21, near Guachochi, Sierra Madres, Chihuahua, Mexico, E. W. Nelson and E. A. Goldman, Sept. 28, 1898.

Sceloporus poinsetti polylepis Smith and Chrapliwy

Herpetologica, vol. 13, p. 269, 1957.

=*Sceloporus poinsettii polylepis* Smith and Chrapliwy

PARATYPES (15): 105486-92, near La Loma, Durango, Mexico, H. M. Smith, November 1-4, 1938. 105493-7, Pedriceña, Durango, Mexico, H. M. Smith, November 2-4, 1938. 105498-500, 14 miles north of Pedriceña, Durango, Mexico, H. M. Smith, November 4, 1938.

Sceloporus prezygus Smith

Proc. U.S. Nat. Mus., vol. 92, p. 354, 1942.

HOLOTYPE: 46861 (given incorrectly as 46881 in original description), Conjab, between San Bartolomé and Comitán, Chiapas, Mexico, 5,300 ft., E. W. Nelson and E. A. Goldman, May 2, 1894.

Sceloporus pyrocephalus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 16, p. 177, 1864.

COTYPES (37): 31428-32, 31434-54, 31456-65, 31495, near Colima, Colima, Mexico, J. Xantus, 1863. The lectotype is 31495. 31433, 31455 were exchanged to MCZ in 1923; 31427 was exchanged to UIMNH in 1956.

Sceloporus rufidorsum Yarrow

Proc. U.S. Nat. Mus., vol. 5, p. 442, 1882.

=*Sceloporus magister rufidorsum* Cope

HOLOTYPE: 11981, San Quintín Bay, Baja California, Mexico, L. Belding, 1882. This specimen is macerated; only the bones are left.

PARATYPES (5): 11971, Cedros Island, Baja California, Mexico (given as "Cerros Island" in the original description), L. Belding. 12667 (4), La Paz, Baja California, Mexico, L. Belding, 1882. One from 11971 was exchanged to MZUM in 1923.

Sceloporus scalaris slevini Smith

Occ. Pap. Mus. Zool. Univ. Michigan, No. 361, p. 3, 1937.

PARATYPES (3): 47064-5, Guadalupe y Calvo, Chihuahua, Mexico, E. W. Nelson and E. A. Goldman, Sept. 8, 1898. 58444, Chihuahua, Mexico, J. Hurter, May 1902.

Sceloporus scalaris unicanthalis Smith

Occ. Pap. Mus. Zool. Univ. Michigan, No. 361, p. 4, 1937.

PARATYPE: 47884, Atemajac, Jalisco, Mexico, E. W. Nelson, May 25, 1892.

Sceloporus serrifer Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 18, p. 124, 1866.

=*Sceloporus serrifer serrifer* Cope

HOLOTYPE: 24868, Yucatán, Mexico (restricted to Mérida, Yucatán), A. Schott.

PARATYPES (7): 10298, ♂, 24869-74, same data.

Sceloporus serrifer plioporus Smith

Field Mus. Nat. Hist., Zool. Ser., vol. 26, p. 212, pl. 18 (right), 1939.

PARATYPE: 138896 (formerly UIMNH 27362), 5 to 6 miles east of Jalapa, Veracruz, Mexico, E. H. Taylor, Sept. 6, 1936.

Sceloporus siniferus Cope

Proc. Amer. Philos. Soc., vol. 11, p. 159, 1869.

=*Sceloporus siniferus siniferus* Cope

COTYPES (18): 30453-6, 30458-71, Pacific side of Isthmus of Tehuantepec, Oaxaca, Mexico (restricted to Tehuantepec City, Oaxaca, F. Sumichrast. 30457 was exchanged to UIMNH in 1956.

Sceloporus smaragdinus Bocourt

Récherches zoologiques, in Mission scientifique au Mexique et dans l'Amérique Centrale, pt. 3, text and plates, p. 186, pl. 18, fig. 6, pl. 19, fig. 1, 1874.

=*Sceloporus malachiticus smaragdinus* Bocourt

COTYPE: 11000, "Mexico" (probably from Guatemala, in vicinity of Solola, Totonicapam, and Quetzaltenango, as given in original description), MHNP.

Sceloporus smaragdinus Cope

in Yarrow, Report upon the collections of batrachians and reptiles . . . 1871, 1872, 1873, and 1874, in Report U.S. Geogr. and Geol. Expl. and Surv. West of One Hundr. Mer., vol. 5, p. 572, pl. 24, figs. 2-2a, 1875. (Not of Bocourt, 1873.)

=*Sceloporus occidentalis longipes* Baird

COTYPES (4): 8612 (4), Beaver (Beaver County), Utah, and Dome Canyon, Nevada, H. C. Yarrow, 1872.

Sceloporus spinosus apicalis Smith and Smith

Proc. Biol. Soc. Washington, vol. 64, p. 101, 1951.

PARATYPE: 134306 (formerly UIMNH 10830), San Pedro Quiechapa, Oaxaca, Mexico, Dr. and Mrs. H. M. Smith, May 1940.

Sceloporus squamosus Bocourt

Récherches zoologiques, in Mission scientifique au Mexique et dans l'Amérique Centrale, pt. 3, text and plates, p. 212, pl. 18 bis, figs. 7-7c, pl. 19, fig. 3, 1874.

COTYPE: 10964, probably from the mouth of the Nagualate River, Guatemala.

Sceloporus stejnegeri Smith

Proc. U.S. Nat. Mus., vol. 92, p. 350, 1942.

HOLOTYPE: 112634, ♂, Tierra Colorada, Guerrero, Mexico, H. M. Smith, Aug. 31, 1939.

PARATYPES (14): 112635-48, Tierra Colorada, Guerrero, Mexico, H. M. Smith, Feb. 13-14, 1939.

Sceloporus taeniocnemis Cope

Proc. Amer. Philos. Soc., vol. 22, p. 399, 1885.

=*Sceloporus malachiticus taeniocnemis* Cope

HOLOTYPE: 24768, Guatemala (restricted to Cobán, Guatemala), H. Hague.

Sceloporus thayeri Baird and Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 127, 1852.

=*Sceloporus undulatus hyacinthinus* Green

COTYPES (3): 2887 (3), Indianola (Calhoun County), Texas, J. H. Clark.

Sceloporus torquatus cyanogenys Cope

Proc. Amer. Philos. Soc., vol. 22, p. 402, 1885.

=*Sceloporus cyanogenys* Cope

COTYPES (5): 31373-7, Monterrey, Nuevo León, Mexico, E. D. Cope.

Sceloporus torquatus minor Cope

Proc. Amer. Philos. Soc., vol. 22, p. 402, 1885.

=*Sceloporus jarrovi minor* Cope

COTYPES (2): 26166-7, Zacatecas, Mexico (restricted to Valparaiso Mountains, Zacatecas, Mexico), A. Dugès, 1884.

Sceloporus torquatus mucronatus Cope

Proc. Amer. Philos. Soc., vol. 22, p. 402, 1885.

=*Sceloporus mucronatus mucronatus* Cope

COTYPES (5): 25075-9, Mirador, Veracruz, Mexico, C. Sartorius. 25077 is the lectotype. 25074 was exchanged to MCZ in 1944.

Sceloporus tristichus Cope

in Yarrow, Report upon the collections of batrachians and reptiles . . . 1871, 1872, 1873, 1874, in Report U.S. Geogr. and Geol. Expl. and Surv. West of One Hundr. Mer., vol. 5, p. 571, 1875.

=*Sceloporus undulatus tristichus* Cope

HOLOTYPE: 8613, Taos (Taos County), New Mexico, W. G. Shedd, August 1874.

Sceloporus undulatus erythrocheilus Maslin

Herpetologica, vol. 12, p. 291, 1956.

HOLOTYPE: 137833, Purgatoire River, 19 miles east of Model, Las Animas County, Colorado, T. P. Maslin and H. A. Fehlmann, Sept. 17, 1949.

Sceloporus utiformis Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 16, p. 177, 1864.

COTYPES (2): 42089, near Colima, Colima, Mexico, J. Xantus, 1863. 42090, near Chacan River, Michoacán, Mexico, J. Xantus, May 1863.

Sceloporus vandenburgianus Cope

Amer. Nat., vol. 30, p. 834, 1896.

=*Sceloporus graciosus vandenburgianus* Cope

HOLOTYPE: 21931, ♂, Campbell's Ranch, Laguna, Imperial County, California, summit of coast range, E. A. Mearns, June 20, 1894.

Sceloporus variabilis olloporus Smith

Occ. Pap. Mus. Zool. Univ. Michigan, No. 358, p. 11, 1937.

PARATYPES (4): 37099-100, Ballena Bay, Nicoya, Peninsula, Costa Rica, A. Alfaro, Nov. 15, 1906. 37101-2, ♂, ♀, Circuelas, Costa Rica, 900 ft., A. Alfaro, Nov. 20, 1906.

Sceloporus variabilis smithi Hartweg and Oliver

Occ. Pap. Mus. Zool. Univ. Michigan, No. 356, p. 1, 1937.

PARATYPES (7): 30332-7, Tehuantepec, Oaxaca, Mexico, F. Sumichrast. 46994, Tehuantepec, Oaxaca, Mexico, E. W. Nelson and E. A. Goldman, May 24, 1895.

Sceloporus viviparus Cope

Proc. Amer. Philos. Soc., vol. 22, p. 398, 1885.

=*Sceloporus formosus formosus* Wiegmann

HOLOTYPE: 25073, Mirador, Veracruz, Mexico, C. Sartorius.

PARATYPES (4): 30209-12, Orizaba, Veracruz, Mexico, F. Sumichrast.

Sceloporus woodi Stejneger

Proc. Biol. Soc. Washington, vol. 31, p. 90, 1918.

HOLOTYPE: 48720, ♂, Auburndale, Polk County, Florida, N. R. Wood, 1912.

PARATYPES (21): 11997 (2), Georgiana, Brevard County, Florida, W. Wittfield. 48721, ♀, Auburndale, Polk County, Florida, N. R. Wood, 1912. 60829-39, 60841-7, Georgiana, Brevard County, Florida, W. Wittfield. 60840 was exchanged to UIMNH in 1956.

Sceloporus zosteromus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 15, p. 105, 1863.

=*Sceloporus magister zosteromus* Cope

COTYPES (18): 5298, Cape San Lucas, Baja California, Mexico, J. Xantus. 69472, Santiago, 15 leagues north of San José del Cabo, Baja California, Mexico, J. Xantus, December 1859. 69473-9, Miraflores, 10 leagues north of San José del Cabo, Baja California, Mexico, J. Xantus, December 1859. 69480, Santiago Sierras, between San José and La Paz, Baja California, Mexico, J. Xantus, November 1859. 69481-2, San José Mission, Baja California, Mexico, J. Xantus, May 15, 1859. 69483-7, Cape San Lucas, Baja California, Mexico, J. Xantus, May 1859-March 1860. 69488, Baja California, Mexico, J. Xantus. One from 5298 was exchanged to MCZ in 1944; 1 from 5298 to UIMNH, 1956.

Scincus lateralis Say

in Long, Account of an expedition from Pittsburgh to the Rocky Mountains . . . , vol. 2, p. 324, 1823.

=*Lygosoma lateralis* (Say)

HOLOTYPE: 3152, Mississippi River below Cape Girardeau, Cape Girardeau County, Missouri, S. W. Woodhouse.

Scincus philbyi Schmidt

Field Mus. Nat. Hist., Zool. Ser., vol. 24, p. 162, fig. 18, 1941.

PARATYPE: 134423, 'Anaiza, Saudi Arabia, H. St. J. B. Philby, June 11, 1938.

Scolecocaurus trinitatis Barbour

Mem. Mus. Comp. Zool., vol. 44, p. 316, 1914.

COTYPE: 120795 (formerly MCZ 8957), Caparo, Trinidad, West Indies, A. B. Carr, 1913.

Sphaerodactylus alopecus Cope

Proc. Acad. Nat. Sci., Philadelphia, vol. 13 (1861), p. 499, 1862.

=*Sphaerodactylus cinereus* Wagler

COTYPE: 52407 (formerly MCZ 3343), Grand' Anse River, near Jérémie, Haiti, D. F. Weinland.

Sphaerodactylus argus henriquesi Grant

Jamaica To-day, p. 134, 1940.

PARATYPE: 120710 (formerly MCZ 44976), Mandeville, Manchester, Jamaica, A. Henriques, June 1937–February 1938.

Sphaerodactylus bartschi Cochran

Smithsonian Misc. Coll., vol. 92, No. 3259, p. 5, 1934.

=*Sphaerodactylus argus bartschi* Cochran

HOLOTYPE: 81759, ♂, Little Cayman Island, West Indies, P. Bartsch, Sept. 13, 1930.

PARATYPES (3): 81757–8, 81760, Little Cayman Island, P. Bartsch, Sept. 12–13, 1930. 81761 was exchanged to MCZ in 1934.

Sphaerodactylus caicosensis Cochran

Smithsonian Misc. Coll., vol. 92, No. 3259, p. 7, 1934.

HOLOTYPE: 81443, ♀, South Caicos Island (Jamaica), P. Bartsch, July 29, 1930.

PARATYPE: 81447, ♀, Long Cay, off South Caicos Island, P. Bartsch, July 29, 1930.

Sphaerodactylus corticolus Garman

Bull. Essex Inst., vol. 20, p. 111, 1888.

COTYPE: 52408 (formerly MCZ 6219), Rum Cay, Bahama Islands, C. J. Maynard, January 1888.

Sphaerodactylus exsul Barbour

Mem. Mus. Comp. Zool., vol. 44, p. 264, 1914.

PARATYPE: 120711 (formerly MCZ 29975), Little Swan Island, Honduras, G. Nelson, March 1912.

Sphaerodactylus gaigeae Grant

Journ. Dept. Agr. Puerto Rico, vol. 16, p. 31, 1932.

COTYPES (2): 120712–3, 10 km. south of Canóvanas, Puerto Rico, C. Grant, May 1932.

Sphaerodactylus glaucus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 17, p. 192, 1865.

=*Sphaerodactylus glaucus glaucus* Cope

COTYPES (3): 6572, near Mérida, Yucatán, Mexico, A. Schott, February 1866. 62995-6, near Mérida, Yucatán, Mexico, A. Schott, Dec. 22, 1866. 62994 was exchanged to MCZ in 1920.

Sphaerodactylus grandisquamis Stejneger

Report of the U.S. National Museum . . . for the year ending June 30, 1902, p. 602, figs. 46-52, 1904.

=*Sphaerodactylus macrolepis* Günther

HOLOTYPE: 27007, Luquillo, Puerto Rico, L. Stejneger, Mar. 4, 1900.

PARATYPES (52): 25549-53, Isabel Island, Vieques Island, Puerto Rico, A. B. Baker, Feb. 7, 1899. 27008-16, 27018-28, 27030, 27032-45, Luquillo, Puerto Rico, L. Stejneger, March 1900. 27142-45, Vieques Island, Puerto Rico, L. Stejneger, Mar. 26, 1900. 27305-12, Ponce, Puerto Rico, L. Stejneger, Apr. 16-June 23, 1900. 27017, 27029, and 27031 were exchanged to MCZ in 1909.

Sphaerodactylus homolepis Cope

Proc. Amer. Philos. Soc., vol. 23, p. 277, 1886.

HOLOTYPE: 14207, Nicaragua, J. F. Bransford.

Sphaerodactylus klauberi Grant

Journ. Dept. Agr. Puerto Rico, vol. 15, p. 207, pl. 20, fig. 1, pl. 23, fig. 13, 1931.

PARATYPE: 120719, Rangers Cabin, El Junque, Puerto Rico, C. Grant, Jan. 11, 1931.

Sphaerodactylus mariguanae Cochran

Smithsonian Misc. Coll., vol. 92, No. 3259, p. 9, 1934.

HOLOTYPE: 81381, Booby Island, east of Mariguana Cay, Bahama Islands, P. Bartsch, July 21, 1930.

PARATYPES (6): 81376-8, 81380, 81382-3, same data. 81379 was exchanged to MCZ in 1934.

Sphaerodactylus nicholsi Grant

Journ. Dept. Agr. Puerto Rico, vol. 15, p. 204, pl. 23, fig. 11, 1931.

PARATYPES (7): 120731-7, Pazuera, Puerto Rico, C. Grant, July 30, 1932.

Sphaerodactylus pacificus Stejneger

Proc. Biol. Soc. Washington, vol. 16, p. 3, 1903.

HOLOTYPE: 31057, Cocos Island, Costa Rica, P. Biolley, January 1902.

PARATYPES (3): 31059-61, same data. 31058 was exchanged to MCZ in 1920.

Sphaerodactylus richardsoni gossei Grant

Copeia, No. 1, p. 10, fig. 2, 1939.

PARATYPES (2): 108406-7, near mouth of Roaring River, Jamaica, C. Grant, Mar. 20, 1938.

Sphaerodactylus sabanus Cochran

Proc. Biol. Soc. Washington, vol. 51, p. 148, 1938.

HOLOTYPE: 103985, Saba Island, West Indies, S. T. Danforth, June 19, 1937.

PARATYPES (16): 103986, 103988-93, 103995-104003, same data. 103987, 103994, and 104004 were exchanged to MCZ in 1938.

Sphaerodactylus samanensis Cochran

Proc. Biol. Soc. Washington, vol. 45, p. 183, 1932.

= *Sphaerodactylus difficilis* Barbour

HOLOTYPE: 74970, Boca del Infierno, Dominican Republic, G. S. Miller, Jr., Feb. 28, 1928.

PARATYPES (3): 74971-3, same data.

Sphaerodactylus stejnegeri Cochran

Copeia, No. 3, p. 89, 1931.

HOLOTYPE: 76640, San Michel, Haiti, A. J. Poole and W. M. Perrygo, Dec. 21-31, 1928.

PARATYPE: 60617, southwestern Haiti, W. L. Abbott.

Sphaerodactylus townsendi Grant

Journ. Dept. Agr. Puerto Rico, vol. 15, p. 208, pl. 23, fig. 12, 1931.

PARATYPES (4): 120738-41, Cape San Juan, northeast corner, Puerto Rico, C. Grant, Jan. 25, 1931.

Sphenomorphus atrigularis Stejneger

Proc. U.S. Nat. Mus., vol. 34, p. 199, figs. 1-2, 1908.

HOLOTYPE: 37386, Datu Ahib's place, base of Malingdang Mountains, Catagan, Misamis Province, Mindanao, Philippine Islands, E. A. Mearns, May 24, 1906.

Sphenomorphus deignani Taylor

Univ. Kansas Sci. Bull., vol. 33, p. 497, fig. 3a-b, 1950.

HOLOTYPE: 120326, Mount Ganoruva, Peradiniya, Kandy District, Ceylon, H. G. Deignan, Nov. 12, 1944.

Sphenomorphus helenae Cochran

Proc. Biol. Soc. Washington, vol. 40, p. 183, 1927.

HOLOTYPE: 67265, Nontaburi, Siam (Thailand), Hugh M. Smith, Sept. 2, 1923.

Sphenomorphus indicus formosensis Van Denburgh

Advance diagnoses of new reptiles and amphibians from the Loo Choo Islands and Formosa, p. 6, July 29, 1912 (published privately).

PARATYPE: 64209 (CAS 18687), Kanshirei, Formosa, V. Kuhne, Mar. 30, 1909.

Sphenomorphus melanopleurus Inger

Chicago Nat. Hist. Mus.: Fieldiana, Zool., vol. 39, p. 264, fig. 45, 1958.

PARATYPE: 139415, Sepik River, Marienberg, New Guinea, W. A. Weber and F. C. Wonder, May 1929.

Sphenomorphus multisquamatus Inger

Chicago Nat. Hist. Mus.: Fieldiana, Zool., vol. 39, p. 261, figs. 43 (right), 44, 1958.

PARATYPE: 130235, Bundu Tuhan, Mount Kinabalu, North Borneo, D. H. Johnson, July 1, 1951.

Sphenomorphus sabanus Inger

Chicago Nat. Hist. Mus., Fieldiana: Zool., vol. 39, p. 258, figs. 42, 43 (left), 1958.

PARATYPE: 139416, Deramakot, Kanabatangan District, North Borneo, R. F. Inger, Apr. 24, 1956.

Sphenomorphus steerei Stejneger

Proc. U.S. Nat. Mus., vol. 34, p. 201, figs. 3-4, 1908.

HOLOTYPE: 32658, Guimaras Island, Philippine Islands, J. B. Steere.

Sphenosoma hughii Cochran

Proc. Biol. Soc. Washington, vol. 40, p. 185, 1927.

HOLOTYPE: 72275, Koh Tao, Gulf of Siam (Thailand), Hugh M. Smith, Jan. 1, 1927.

PARATYPES (2): 72274, 72276, same data.

Stenocercus ervingi Stejneger

Proc. U.S. Nat. Mus., vol. 45, p. 545, 1913.

=*Stenocercus crassicaudatus* (Tschudi)

HOLOTYPE: 49550, Huadquiña, Peru, 5,000 ft., Yale-Peruvian Expedition, 1911.

Stenodactylus variegatus Baird

Proc. Acad. Nat. Sci. Philadelphia, vol. 10 (1858), p. 254, 1859.

=*Coleonyx variegatus variegatus* (Baird)

HOLOTYPE: 3217, Colorado Desert (restricted to Winterhaven, Imperial County, California), A. Schott.

Takydromus dorsalis Stejneger

Smithsonian Misc. Coll., vol. 47, No. 1487, p. 294, 1904.

=*Platyplacopus dorsalis* (Stejneger)

HOLOTYPE: 34162, Ishigaki, Yaeyama Group, Riu Kiu (Ryukyu) Islands, A. Owston, April-June, 1899.

PARATYPES (6): 34160-1, 34163-4, 34168-9, same data. 34165 was exchanged to BMNH in 1916; 34166-7 were exchanged to MCZ, 1944.

Takydromus intermedius Stejneger

Occ. Pap. Boston Soc. Nat. Hist., vol. 5, p. 120, 1924.

=*Platyplacopus intermedius* (Stejneger)

HOLOTYPE: 64437, Shin Kai Si Mountain, Szechwan, China, D. C. Graham, 1921.

PARATYPES (3): 64425, 65467, Mount Omei, Szechwan, China, D. C. Graham, 1921. 65933, Suifu, Szechwan, China, D. C. Graham, 1923. 63594 was exchanged to MCZ in 1944.

Takydromus khasiensis Boulenger

Mem. Asiatic Soc. Bengal, vol. 5, No. 6, p. 221, 1917.

PARATYPE: 84994, Loshiang, Yaoshan Mountains, Kwangsi, China, SYUM.

Takydromus tachydromoides oldi Walley

Herpetologica, vol. 14, p. 202, fig. 1, 1958.

HOLOTYPE: 129618, ♀, Taejon, Korea, picked up in an abandoned school by W. E. Old, January 1951.

PARATYPE: 129616, same data.

Tapaya brevirostris Girard

in U.S. Exploring Expedition . . . 1838-42, vol. 20 (Herpetology), p. 397, 1858.

=*Phrynosoma douglassii brevirostris* (Girard)

COTYPES (10): 208, 4592 (9), Pole Creek, Nebraska, S. W. Woodhouse, July 26, 1856.

Tapaya hernandesi Girard

in U.S. Exploring Expedition . . . 1838-42, vol. 20 (Herpetology), p. 395, 1858.

=*Phrynosoma douglassii hernandesi* (Girard)

COTYPES (2): 197, Sonora (restricted to Sante Fe, New Mexico), J. H. Clark. 198, Santa Fe (Santa Fe County, New Mexico), J. H. Clark. One cotype was exchanged to MHNP.

Tapaya ornatissima Girard

in U.S. Exploring Expedition . . . 1838-42, vol. 20 (Herpetology), p. 396, 1858.

=*Phrynosoma douglassii ornatissimum* (Girard)

COTYPES (2): 204 (2), Zuni Mountains, New Mexico, S. W. Woodhouse, 1851.

Tiaporus fuliginosus Cope

Proc. Amer. Philos. Soc., vol. 30, p. 132, 1 pl., 1892.

=*Ameiva ameiva fuliginosa* (Cope)

COTYPES (3): 14710, 32119-32120, Swan Island, Honduras, C. H. Townsend, May 20, 1887. 32118 was exchanged to MCZ in 1925.

Tiliqua dura Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 190, 1862.

=*Sphenosoma durum* (Cope)

HOLOTYPE: 5996, western Africa (probably Sierra Leone), S. M. E. Goheen.

Tropidophorus misaminius Stejneger

Proc. U.S. Nat. Mus., vol. 34, p. 202, figs. 5-6, 1908.

HOLOTYPE: 37383, Catagan, Misamis Province, Mindanao, Philippine Islands, 1,100 ft., E. A. Mearns, May 25, 1906.

PARATYPES (2): 37384-5, same data.

Tropidophorus partelloi Stejneger

Proc. U.S. Nat. Mus., vol. 39, p. 97, 1910.

HOLOTYPE: 39308, Mataling River Falls, Mindanao, Philippine Islands, 1,400 ft., J. M. T. Partello, Dec. 12, 1908.

Tropidurus abingdonensis Baur

Biol. Centralbl., vol. 10, p. 479, 1890.

=*Tropidurus pacificus* Steindachner

COTYPES (33): 14966-97, 15001, Abingdon Island, Galápagos Islands (Ecuador), U.S. Fish Commission *Albatross*, Apr. 16, 1888. 14999-15000 were exchanged to MCZ in 1926; 14998 was exchanged to UIMNH, 1956.

Tropidurus albemarlensis Baur

Biol. Centralbl., vol. 10, p. 478, 1890.

=*Tropidurus albemarlensis albemarlensis* Baur

COTYPES (9): 15003-9, 15011, 15013, Albemarle Island, Galápagos Islands (Ecuador), U.S. Fish Commission *Albatross*, Apr. 10, 1886. 15012 was exchanged to MCZ in 1926; 15010 to AMNH, 1930.

Tropidurus duncanensis Baur

Biol. Centralbl., vol. 10, p. 479, 1890.

COTYPES (4): 14941-4, Duncan Island, Galápagos Islands (Ecuador), U.S. Fish Commission, *Albatross*, Apr. 13, 1888.

Tropidurus hoodensis Baur

Biol. Centralbl., vol. 10, p. 478, 1890.

=*Tropidurus delanonis* Baur

COTYPES (16): 14926-30, Gardners Island, Galápagos Islands (Ecuador), U.S. Fish Commission *Albatross*, Apr. 7, 1888. 15014-8, 15020-5. Hood Island, Galápagos Islands, U.S. Fish Commission, *Albatross*, Apr. 7, 1888. 15026 was exchanged to MCZ in 1926.

Tropidurus indefatigabilis Baur

Biol. Centralbl., vol. 10, p. 478, 1890.

=*Tropidurus albemarlensis albemarlensis* Baur

COTYPES (36): 14897-915, 14917-24, James Island, Galápagos Islands (Ecuador), U.S. Fish Commission *Albatross*, Apr. 11, 1888. 14931-7, 14939-40, Indefatigable Island, Galápagos Islands, U.S. Fish Commission *Albatross*, Apr. 12, 1888. 14916 and 14938 were exchanged to MCZ in 1926.

Tropidurus lemniscatus Cope

Proc. U.S. Nat. Mus., vol. 12, p. 145, 1890.

=*Tropidurus bivittatus* (Peters)

COTYPES (16): 14946-55, 14959-64, Chatham Island, Galápagos Islands (Ecuador), U.S. Fish Commission *Albatross*, April 4, 1888. 14958 was exchanged to MCZ in 1928.

Uma notata Baird

Proc. Acad. Nat. Sci. Philadelphia, vol. 10 (1858), p. 253, 1859.

HOLOTYPE: 4124, "Mojave Desert, California" (restricted to Yuma, Yuma County, Arizona), A. L. Heerman.

Uma rufopunctata Cope

Amer. Nat., vol. 29, p. 939, 1895.

=*Uma notata notata* Baird

COTYPES (6): 21846-47, 21849-52, Monument 200, Yuma Desert (Yuma County), Arizona, E. A. Mearns, May 15-16, 1894. 21848 was exchanged to UIMNH in 1956.

Uma scoparia Cope

Amer. Nat., vol. 23, p. 435, figs. 3-4, 1894.

=*Uma notata scoparia* Cope

HOLOTYPE: 6063, Fort Buchanan?, Arizona? (probably from San Bernardino County, California), B. J. D. Irwin.

Urosaurus bicarinatus spinosus Bumzahem and Smith

Herpetologica, vol. 10, p. 181, fig. 1, 1954.

HOLOTYPE: 113341, ♂, Tonalá, Chiapas, Mexico, H. M. Smith, Jan. 26-30, 1940.

PARATYPE: 113342, ♀, same data.

Uta anonymorpha Mittleman

Herpetologica, vol. 2, p. 34, pl. 3, fig. 2, 1940.

=*Urosaurus bicarinatus anonymorphus* (Mittleman)

PARATYPES (3): 30095-7, Tehuantepec, Oaxaca, Mexico, F. Sumichrast.

Uta auriculata Cope

Proc. Boston Soc. Nat. Hist., vol. 14, p. 303, 1871.

=*Urosaurus auriculatus* (Cope)

HOLOTYPE: 46988, near Tehuantepec, Oaxaca, Mexico, E. W. Nelson and E. A. Goldman, May 27, 1895.

HOLOTYPE: 7027, skin, Socorro Island, Revillagigedo Islands, Colima, Mexico, A. J. Grayson.

Uta caerulea Smith

Univ. Kansas Sci. Bull., vol. 22, p. 172, pl. 26, fig. 3, 1935.

=*Urosaurus ornatus caeruleus* (Smith)

PARATYPE: 138897 (former UIMNH 22082), 20 miles south of Chihuahua, Chihuahua, Mexico, H. M. Smith and D. H. Dunkle, June 22, 1934.

Uta clarionensis Townsend

Proc. U.S. Nat. Mus., vol. 13, p. 143, 1890.

=*Urosaurus clarionensis* (Townsend)

HOLOTYPE: 15904, Clarión Island, Revillagigedo Islands, Colima, Mexico, C. H. Townsend, Mar. 4, 1889.

PARATYPES (4): 15905-8, same data.

Uta concinna Dickerson

Bull. Amer. Mus. Nat. Hist., vol. 41, p. 470, 1919.

HOLOTYPE: 64257, ♂, Cedros Island, Baja California, Mexico (given as "Cerros Island" in the original description), C. H. Townsend, Mar. 10, 1911.

Uta elegans Yarrow

Proc. U.S. Nat. Mus., vol. 5, p. 442, 1882.

=*Uta stansburiana elegans* Yarrow

COTYPES (13): 12666 (9), 12668 (4), La Paz, Baja California, Mexico, L. Belding, 1882. One from 12666 was exchanged to BMNH; one from 12666 to UIMNH, 1956.

Uta levis Stejneger

North Amer. Fauna, No. 3, p. 108, 1890.

HOLOTYPE: 11474, ♂, Tierra Amarilla, Rio Arriba County, New Mexico, 7,800 ft., E. D. Cope, Sept. 15, 1874.

PARATYPE: 8554, ♀, same data.

Uta mannophorus Dickerson

Bull. Amer. Mus. Nat. Hist., vol. 41, p. 470, 1919.

HOLOTYPE: 64260, ♂, Carmen Island, Gulf of California, Baja California, Mexico, C. H. Townsend, Apr. 5, 1911.

Uta mearnsi Stejneger

Proc. U.S. Nat. Mus., vol. 17, p. 589, 1894.

=*Streptosaurus mearnsi* (Stejneger)

HOLOTYPE: 21882, ♀, summit of Coast Range, U.S.-Mexican Boundary, California, E. A. Mearns, May 16, 1894.

PARATYPES (5, from the U.S.-Mexican Boundary at the summit of the Coast Range in California collected by E. A. Mearns): 21883, ♀, mountain spring, eastern slope, May 13, 1894. 21884-5, ♀, ♀, eastern base, May 11, 1894. 21886-7, ♂, ♂, mountain spring, eastern base, May 11, 9, 1894. 21888 was exchanged to UIMNH in 1956.

Uta nelsoni Schmidt

Amer. Mus. Novit., No. 22, p. 4, 1921.

=*Urosaurus bicarinatus nelsoni* (Schmidt)

HOLOTYPE: 46836, ♂, Cuicatlán, Oaxaca, Mexico, E. W. Nelson and E. A. Goldman, October 9, 1894.

Uta nigricauda Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 16, p. 176, 1864.

=*Urosaurus nigricaudus* (Cope)

COTYPES (31): 5307 (19), Cape San Lucas, Baja California, Mexico, J. Xantus. 69414-5, Miraflores, 10 leagues north of San José, Baja California, Mexico, J. Xantus, December 1859. 69416-8, Cape San Lucas, Baja California, Mexico, J. Xantus, December 1859. 69419-25, Santiago Sierras, between San José and La Paz, Baja California, Mexico, J. Xantus, November 1859.

Uta ornata Baird and Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 126, 1852.

=*Urosaurus ornatus ornatus* (part) (Baird and Girard)

COTYPES (2): 2750 (2), San Pedro, Texas (restricted to Rio San Pedro, Val Verde County, Texas), J. H. Clark.

Uta ornata linearis Baird

Reptiles of the boundary, in U.S. and Mexican Bound. Surv., vol. 2, pt. 2, p. 7, 1859.

=*Urosaurus ornatus linearis* (Baird)

NEOTYPE: 62077, Nogales, Sonora, Mexico, F. J. Dyer.

Uta ornata schmidti Mittleman

Herpetologica, vol. 2, p. 33, pl. 3, fig. 1, 1940.

=*Urosaurus ornatus schmidti* (Mittleman)

HOLOTYPE: 32929, ♂, Fort Davis, Jeff Davis County, Texas, Chisos Mountains, V. O. Bailey, June 3, 1901.

PARATYPES (15): 11957 (2), Fort Davis, Jeff Davis County, Texas, W. F. von Manteufel, July 1878. 30012, El Paso, El Paso County, Texas, H. H. and C. S. Brimley, Mar. 29, 1902. 32928-9, Fort Davis, Jeff Davis County, Texas, V. O. Bailey, Jan. 3-6, 1890. 32930-1, Davis Mountains, Jeff Davis County, Texas, V. O. Bailey, Jan. 3, 1890. 32932-3, Paisano, Presidio County, Texas, W. Lloyd, July 1890. 32934-5, Altuda, Brewster County, Texas, W. Lloyd, August 1890. 32936, Chisos Mountains, Brewster County, Texas, V. O. Bailey, June 6, 1901. 92897-8, 6 miles northwest of Alpine, Brewster County, Texas, L. Hoyle, June 17, 1933. 103646, Oak Creek, Chisos Mountains, Brewster County, Texas, T. F. Smith, June 1936.

Uta palmeri Stejneger

North Amer. Fauna, No. 3, p. 106, 1890.

HOLOTYPE: 16002, San Pietro Mártir Island, Gulf of California, Sonora, Mexico, E. Palmer.

PARATYPES (12): 16003-14, same data.

Uta parva Dickerson

Bull. Amer. Mus. Nat. Hist., vol. 41, p. 471, 1919.

=*Uta stansburiana hesperis* Richardson

HOLOTYPE: 64258, ♂, San Bartolomé Bay, Baja California, Mexico, C. H. Townsend, Mar. 13, 1911.

Uta squamata Dickerson

Bull. Amer. Mus. Nat. Hist., vol. 41, p. 471, 1919.

HOLOTYPE: 64259, ♂, Santa Catalina Island, Gulf of California, Baja California, Mexico, C. H. Townsend, May 16, 1911.

Uta stansburiana Baird and Girard

in Stansbury, Report Expl. and Surv. Valley of the Great Salt Lake of Utah . . ., p. 345, pl. 5, figs. 4-6, 1852.

=*Uta stansburiana stansburiana* Baird and Girard

COTYPES (4): 2753 (4), Valley of the Great Salt Lake, Utah, H. Stansbury.

Uta symmetrica Baird

Proc. Acad. Nat. Sci. Philadelphia, vol. 10, p. 253, 1858.

=*Urosaurus ornatus symmetricus* (Baird)

NEOTYPE: 2744A, Fort Yuma (Imperial County), California, G. W. Thomas, 1855.

Uta taylori Smith

Univ. Kansas Sci. Bull., vol. 22, p. 158, pl. 26, fig. 3, 1935.

PARATYPE: 134308 (formerly UIMNH 27454), La Posa, Sonora, Mexico, E. H. Taylor, July 13, 1934.

Uta thalassina Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 15, p. 104, 1863.

=*Petrosaurus thalassinus* (Cope)

HOLOTYPE: 5302, Cape San Lucas, Baja California, Mexico, J. Xantus.

Uta unica Mittleman

Journ. Washington Acad. Sci., vol. 31, p. 74, fig. 3, 1941.

=*Urosaurus unicus* (Mittleman)

HOLOTYPE: 14248, ♀, Chihuahua, Mexico, E. Wilkinson.

Verticaria beldingi Stejneger

Proc. U.S. Nat. Mus., vol. 17, p. 17, 1894.

=*Cnemidophorus hyperythrus beldingi* (Stejneger)

HOLOTYPE: 11980, Cerros (=Cedros) Island, Baja California, Mexico (probably from San Quintín Bay), L. Belding, 1878.

Verticaria caerulea Dickerson

Bull. Amer. Mus. Nat. Hist., vol. 41, p. 472, 1919.

=*Cnemidophorus hyperythrus caeruleus* (Dickerson)

HOLOTYPE: 64251, Carmen Island, Gulf of California, Baja California, Mexico, C. H. Townsend, Apr. 3, 1911.

Wetmorena haetiana Cochran

Proc. Biol. Soc. Washington, vol. 40, p. 91, 1927.

HOLOTYPE: 72600, Morne Cabaio, Massif de la Selle, Haiti, 7,500 ft., A. Wetmore, Apr. 10, 1927.

PARATYPES (2): 72601, 72604, same data. 72602 was exchanged to MZUM in 1943; 72603 to MCZ, 1927.

Xantusia arizonae Klauber

Trans. San Diego Soc. Nat. Hist., vol. 7, p. 3, pl. 1, figs. 1-2, 1931.

PARATYPE: 84587, 1 mile south of Yarnell, Yavapai County, Arizona, 4,940 ft., P. M. and L. M. Klauber, Aug. 21, 1931.

Xantusia henshawi Stejneger

Proc. U.S. Nat. Mus., vol. 16, p. 467, 1893.

HOLOTYPE: 20339, Witch Creek, San Diego County, California, 2,700 ft., H. W. Henshaw, May 1893.

Xantusia riversiana reticulata Smith

Journ. Washington Acad. Sci., vol. 36, p. 392, 1946.

=*Xantusia riversiana* Cope

HOLOTYPE: 124381, San Clemente Island, Orange County, California, C. W. Kern, summer of 1940.

Xantusia vigilis Baird

Proc. Acad. Nat. Sci. Philadelphia, vol. 10 (1858), p. 255, 1859.

COTYPES (3): 3063 (3), Fort Tejon (Kern County), California, J. Xantus.

Xiphosurus oculatus Cope

Proc. Amer. Philos. Soc., vol. 18, p. 274, 1879.

=*Anolis oculatus* (Cope)

COTYPES (10): 10139-48, Dominica, West Indies, F. A. Ober, July 1877.

Suborder Serpentes

Adelphicos veraepacis nigrilatus Smith

Proc. Rochester Acad. Sci., vol. 8, p. 182, figs. 1, 5, 1942.

PARATYPE: 137219, San Cristóbal, Chiapas, Mexico, H. D. Thomas, 1937.

Aepidia robusta Hallowell

Proc. Acad. Nat. Sci. Philadelphia, vol. 12 (1860), p. 488, 1861.

=*Elaphe oxycephala* (Boie)

HOLOTYPE: 7508, Gaspar Straits (off the south coast of Sumatra between Bangka and Billiton, Indonesia), J. Rodgers.

Agkistrodon blomhoffii brevicaudus Stejneger

U.S. Nat. Mus. Bull. 58, p. 463, 1907.

HOLOTYPE: 17507, ♂, Pusan, Korea, P. L. Jouy, 1885.

PARATYPES (3): 14612-3, ♂, ♀, Korea, N. M. Ferebee, 17508, ♂, Seoul, Korea, P. L. Jouy, 1883.

Agkistrodon mokeson pictigaster Gloyd and Conant

Bull. Chicago Acad. Sci., vol. 7, p. 156, fig. 10, 1943.

=*Agkistrodon contortrix pictigaster* (Gloyd and Conant)

PARATYPE: 103659, Pulliam Canyon, Chisos Mountains, Brewster County, Texas, 5,500 ft., T. F. Smith, July 4, 1936.

Aipysurus foliosquama Smith

Monograph of the sea-snakes (Hydrophidae), p. 22, figs. 11-12, 1926.

PARATYPE: 71050, ♀, Ashmore Reefs, Timor Sea, North Australia, M. A. Smith, 1925.

Alsophis angulifer ruttayi Grant

Herpetology of the Cayman Islands, Bull. Inst. Jamaica, Sci. Ser., No. 2, p. 50, pl. 3, figs. 5-6, 1941.

PARATYPES (2): 108010-11, Little Cayman, Jamaica, C. Grant, Apr. 8-12, 1938.

Alsophis funereus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 77, 1862.

=*Leimadophis funereus* (Cope)

PARATYPES (2): 5780, 12372 (formerly 5778), Jamaica, C. B. Adams.

Alsophis leucomelas danforthi Cochran

Proc. Biol. Soc. Washington, vol. 51, p. 153, 1938.

HOLOTYPE: 104237, ♂, Terre-de-Bas, Isles des Saintes, West Indies, S. T. Danforth, July 5, 1937.

Alsophis nicholsi richardi Grant

Journ. Agr. Univ. Puerto Rico, vol. 30, p. 124, 1949.

HOLOTYPE: 66522, ♂, St. Thomas, Virgin Islands, E. Sebastian, 1923.

PARATYPES (5): 12403a-b, St. Thomas, Virgin Islands, A. H. Riise. 66523-5, St. Thomas, Virgin Islands, E. Sebastian.

Alsophis sanctorum Barbour

Proc. Biol. Soc. Washington, vol. 28, p. 78, 1915.

=*Alsophis leucomelas sanctorum* Barbour

PARATYPE: 104497 (formerly MCZ 10690), Terre d'en Haut, Iles des Saintes, near Guadeloupe, West Indies, G. K. Noble, 1914.

Alsophis sibonius Cope

Proc. Amer. Philos. Soc., vol. 18, p. 275, 1879.

HOLOTYPE: 10138, Dominica, West Indies, F. A. Ober, July 1877.

Alsophis vudii raineyi Barbour and Shreve

Proc. Boston Soc. Nat. Hist., vol. 40, p. 363, 1935.

PARATYPE: 120808 (formerly MCZ 37940), Landrail Point, Crooked Island, Bahama Islands, T. Barbour, March 1924.

Amphiesma flavipunctatum Hallowell

Proc. Acad. Nat. Sci. Philadelphia, vol. 12 (1860), p. 503, 1861.

=*Natrix piscator flavipunctata* (Hallowell)

HOLOTYPE: 7387, ♂, Island of Hongkong (off the mainland of China), J. M. Brooke, May 1854.

Anomalepis aspinosus Taylor

Proc. New England Zool. Club, vol. 17, p. 92, 1939.

PARATYPE: 76295 (formerly MCZ 14784), Perico, Peru, G. K. Noble, September 1916.

Aporophis juliae Cope

Proc. Amer. Philos. Soc., vol. 18, p. 274, 1879.

=*Leimadophis juliae* (Cope)

HOLOTYPE: 10152, Dominica, West Indies, F. A. Ober, 1877.

Arizona elegans Kennicott

in Baird, Reptiles of the boundary, in Report U.S. and Mexican Bound.

Surv., vol. 2, pt. 2, p. 13, pl. 13, 1859.

=*Arizona elegans elegans* Kennicott

COTYPES (2): 1722, Lower Rio Grande, Texas, A. Schott. 4266, between Arkansas and Cimarron Rivers, Oklahoma, J. H. Clark.

Arizona elegans blanchardi Klauber

Trans. San Diego Soc. Nat. Hist., vol. 10, p. 328, 1946.

PARATYPE: 89239, near Hutchinson, Reno County, Kansas, W. T. Cole, September 1932.

Arizona elegans expolita Klauber

Trans. San Diego Soc. Nat. Hist., vol. 10, p. 340, 1946.

HOLOTYPE: 46374, ♂, Casas Grandes, Chihuahua, Mexico, E. W. Nelson and E. A. Goldman, May 27, 1899.

PARATYPE: 14298, Chihuahua, Mexico, E. Wilkinson, Oct. 10, 1885.

Arizona elegans noctivaga Klauber

Trans. San Diego Soc. Nat. Hist., vol. 10, p. 343, 1946.

PARATYPES (3): 8002, Camp Grant (Graham County), Arizona, E. Palmer. 37112, mesa south of Fort Mojave (Mohave County), Arizona, C. W. Adams, June 9, 1904. 44259, Phoenix (Maricopa County), Arizona, V. O. Bailey, May 21, 1889.

Arizona elegans occidentalis Blanchard

Occ. Pap. Mus. Zool. Univ. Michigan, No. 150, p. 1, 1924.

HOLOTYPE: 54372, La Jolla, San Diego County, California, J. C. Thompson, May 1916.

PARATYPES (10): 8408, southeastern Arizona, H. W. Henshaw, October 1873. 13839, San Diego, San Diego County, California, H. C. Orcutt. 14176, Mojave Desert, California, G. Eisen, July 29, 1885. 16848, Pacific Beach, San Diego County, California, C. R. Orcutt, January 1890. 17401, Alhambra, Los Angeles County, California, E. C. Thurber, Apr. 12, 1889. 29657, Yuma (Yuma County), Arizona, H. Brown, 1901. 37519, San Tomas, Lower California, Mexico, E. W. Nelson and E. A. Goldman, May 25, 1905. 37529, San Quintín, Marin County, California, E. W. Nelson and E. A. Goldman, Aug. 20, 1905. 44389, Fresno, Fresno County, California, C. P. Streater, July 4, 1893. 54371, La Jolla, San Diego County, California, J. C. Thompson, May 1916.

Arizona jani Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 12 (1860), p. 369, 1861.

==*Pituophis deppei jani* (Cope)

HOLOTYPE: 1522, Buena Vista, Coahuila, Mexico, D. N. Couch, 1853.

Arrhyton bivittatum Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 82, 1862.

==*Arrhyton vittatum* (Gundlach and Peters)

HOLOTYPE: 5784, Cuba, F. Poey.

Arrhyton fulvum Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 82, 1862.

=*Arrhyton taeniatum* Günther

HOLOTYPE: 12421, eastern Cuba.

Barbourina equatoriana Amaral

Journ. Washington Acad. Sci., vol. 14, p. 201, 1924.

=*Clelia clelia scytalina* (Cope)

HOLOTYPE: 62790, ♂, Guayaquil, Ecuador, F. W. Goding, April 1920.

Bascanion anthonyi Stejneger

Proc. U.S. Nat. Mus., vol. 23, p. 715, 1901.

=*Masticophis anthonyi* (Stejneger)

HOLOTYPE: 24390, Clarion Island, Revillagigedo Islands, Colima, off west coast of Mexico, A. W. Anthony.

PARATYPES (11): 24382-9, 24391-2, 24394, same data. 24393 was exchanged to CAS in 1921.

Bascanion flagellum frenatum Stejneger

North Amer. Fauna, No. 7, p. 208, 1893.

=*Masticophis flagellum piceus* (Cope)

HOLOTYPE: 16340, Mountain Spring, Colorado Desert, San Diego County, California, C. R. Orcutt, June 1890.

Bascanion foxii Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 96, 1853.

=*Coluber constrictor foxii* (Baird and Girard)

COTYPES (2): 4445, Grosse Ile, Wayne County, Michigan, C. Fox. 131711 (formerly 4446), Pittsburgh, Allegheny County, Pennsylvania, B. A. Fahnestock.

Bascanion suboculare Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 18 (1866), p. 319, 1867.

=*Masticophis mentovarius* (Duméril and Bibron)

COTYPES (2): 6753, 6762, between Cobán and Clusec, Guatemala, H. Hague.

Bascanion vetustus Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 97, 1853.

=*Coluber constrictor mormon* Baird and Girard

PARATYPES (3): 4418-9, 4454, Puget Sound (Washington), U.S. Exploring Expedition, 1841. One from 4419 was exchanged to ANSP.

Bascanium piceum Cope

Proc. U.S. Nat. Mus., vol. 14, p. 625, 1892.

=*Masticophis flagellum piceus* (Cope)

HOLOTYPE: 7891, Camp Grant (Graham County), Arizona, E. Palmer.

Bascanium semilineatum Cope

Proc. U.S. Nat. Mus., vol. 14, p. 626, 1892.

=*Masticophis bilineatus* Jan

COTYPES (2): 1981, Colorado River bottom, Arizona, A. Schott. 8434, Camp Grant (Graham County), Arizona, C. Rutter, Sept. 16, 1874.

Boiga dendrophila occidentalis Brongersma

Zool. Meded. Rijksmuseum Nat. Hist., vol. 17, p. 211, pl. 1, figs. 2, 14, pl. 2, figs. 5-7, text-figs. 34-38, 1934.

PARATYPES (5): 31667-8, ♂, ♀, Lofau, Nias Island, west coast of Sumatra, W. L. Abbott, Mar. 29-30, 1903. 35793-4, 36272, ♂, ♀, ♂, Nias Island, west coast of Sumatra, W. L. Abbott, Mar. 13, 14, 27, 1905.

Bothrops diporus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 347, 1862.

=*Bothrops neuwiedii diporus* Cope

HOLOTYPE: 12380 (formerly 5401), Vermejo River region (boundary of Paraguay and Argentina), La Plata Expedition.

Bothrops lansbergii venezuelensis Roze

Amer. Mus. Novit., No. 1934, p. 11, 1959.

PARATYPE: 129577, Caripito, Monagas, Venezuela, F. Pifano.

Bothrops leptura Amaral

Proc. New England Zool. Club, vol. 8, p. 102, 1923.

HOLOTYPE: 50110, Cana, Panama, 3,000 ft., E. A. Goldman, June 7, 1912.

Bothrops nummifer veraecrucis Burger

Bull. Chicago Acad. Sci., vol. 9, p. 65, 1950.

HOLOTYPE: 25045, Veracruz, Mexico (probably from Mirador, which was designated by Burger as the type locality), C. Sartorius.

PARATYPE: 6371, Mirador, Veracruz, Mexico, C. Sartorius.

Calamaria mearnsi Stejneger

Smithsonian, Misc. Coll., vol. 50, p. 30, 1907.

HOLOTYPE: 36991, Tangob, Mindanao, Philippine Islands, E. A. Mearns, June 10, 1906.

Calamaria raveni Cochran

Proc. Biol. Soc. Washington, vol. 36, p. 91, 1923.

HOLOTYPE: 51638, Sungai Merah, Borneo, H. C. Raven, April 1914.

Calamaria schmidtii Inger

Chicago Nat. Hist. Mus., Fieldiana: Zool., vol. 37, p. 197, 1955.

HOLOTYPE: 130240, Bundu Tuhan, Mount Kinabalu, North Borneo, D. H. Johnson, July 25, 1951.

Calamaria tenuis Baird and Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 176, 1852.

=*Contia tenuis* (Baird and Girard)

HOLOTYPE: 7289, Puget Sound (Washington), U.S. Exploring Expedition, 1841. This specimen is probably also a paratype of *Contia mitis* Baird and Girard.

Calamaria ventralis Cochran

Proc. Biol. Soc. Washington, vol. 36, p. 91, 1923.

=*Calamaria loyii ventralis* Cochran

HOLOTYPE: 51637, Sungai Merah, Borneo, H. C. Raven, April 1914.

Callirhinus patagoniensis Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 9 (1857), p. 182, 1858.

=*Philodryas schottii* (Schlegel)

COTYPE: 5536, coast of Patagonia near mouth of Río Negro (Argentina), U.S. Exploring Expedition, 1839. One of 5536 was exchanged to ANSP.

Callophis hughii Cochran

Proc. Biol. Soc. Washington, vol. 40, p. 190, 1927.

HOLOTYPE: 72307, Koh Tao, Gulf of Siam (Thailand), Hugh M. Smith, Jan. 1, 1927.

Cantoria violacea Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 9 (1857), p. 182, 1858.

=*Cantoria elapiformis* (Peters)

HOLOTYPE: 5523, neighborhood of Singapore (Malaya), U.S. Exploring Expedition, 1842.

Catostoma nasale Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 20, p. 131, 3 figs. 1868.

=*Geophis nasalis* (Cope)

PROBABLE COTYPE: 12425a, Guatemala, C. H. Van Patten.

Catostoma psephotum Cope

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 18, p. 146, 1875.

=*Ninia psephota* (Cope)

HOLOTYPE: 61971, Pico Blanco, Costa Rica, 5,000–7,000 ft., W. M. Gabb.

Caudisona basilisca Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 16, p. 166, 1864.

=*Crotalus basiliscus basiliscus* (Cope)

HOLOTYPE: 53586, skin, near Colima, Colima, Mexico, J. Xantus.

Caudisona enyo Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 13, p. 293, 1861.

=*Crotalus enyo* (Cope)

COTYPES (2): 5291 (2), Cape San Lucas, Baja California, Mexico, J. Xantus, May-June 1859.

Caudisona pyrrha Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 18 (1866), p. 308, 1867.

=*Crotalus mitchellii pyrrhus* (Cope)

HOLOTYPE: 6606, Cañon Prieto, Fort Whipple (Yavapai County), Arizona, E. Coues, 1870.

Causus nasalis Stejneger

Proc. U.S. Nat. Mus., vol. 16, p. 735, 1894.

=*Causus resimus* (Peters)

HOLOTYPE: 16055, ♂, Cunga, Angola, W. H. Brown.

PARATYPES (2): 16074, Cunga, Angola, W. H. Brown, Dec. 25, 1889. 20088, Tana River, East Africa (Kenya), W. A. Chanler, November 1892. 20089 was exchanged to MCZ.

Celuta helenae Kennicott

Proc. Acad. Nat. Sci. Philadelphia, vol. 11, p. 100, 1859.

=*Carphophis amoenus helenae* (Kennicott)

COTYPES (4): 2183 (3), 131708, southern Illinois, R. W. Kennicott. One of 2183 was exchanged to MZUM.

Charina brachyops Cope

Proc. U.S. Nat. Mus., vol. 11, p. 88, pl. 36, figs. 2a-f, 1888.

=*Charina bottae bottae* Blainville

HOLOTYPE: 15524, Point Reyes, Marin County, California, C. H. Merriam and C. A. Allen.

Chilomeniscus ehippicus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 19, p. 85, 1867.

=*Chilomeniscus cinctus* Cope?

HOLOTYPE: 8897, juv., Owens Valley, California, G. H. Horn.

Chilomeniscus stramineus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 12, p. 339, 1860.

=*Chilomeniscus stramineus stramineus* Cope

COTYPES (4): 4674 (2), Cape San Lucas, Baja California, Mexico, J. Xantus. 6495 (2), Cape San Lucas, Baja California, Mexico, J. Xantus, December 1859.

Chilomeniscus stramineus fasciatus Cope

Proc. U.S. Nat. Mus., vol. 14, p. 595, 1892.

=*Chilomeniscus cinctus* Cope

COTYPE: 12630 (2), La Paz, Baja California, Mexico, L. Belding, February 1882.

Chilopoma rufipunctatum Cope

in Yarrow, Report upon the collections of batrachians and reptiles . . .
1871, 1872, 1873, 1874, in Report U.S. Geogr. and Geol. Expl. and
Surv. West of One Hundr. Mer., vol. 5, p. 544, 1875.

=*Thamnophis rufipunctatus* (Cope)

HOLOTYPE: 8600 (formerly 1097), southern Arizona, H. W. Henshaw,
September 1874.

Chionactis diasii Cope

in Ferrari-Perez, Proc. U.S. Nat. Mus., vol. 9, p. 188, 1886.

=*Toluca lineata varians* (Jan)

PARATYPE: 31362, Puebla, Mexico, Mexican Geographical and Exploring
Commission.

Churchillia bellona Baird and Girard

in Stansbury, Report Expl. and Surv. Valley of the Great Salt Lake
of Utah . . ., p. 350, 1852.

=*Pituophis catenifer sayi* (Schlegel)

HOLOTYPE: 1519, San Antonio, west of Rio Grande (given as "Presidio
del Norte" in the original description), Chihuahua, Mexico, S. Churchill,
1846.

Clelia baileyi Smith

Proc. U.S. Nat. Mus., vol. 92, p. 391, 1942.

=*Oxyrhopus baileyi* (Smith)

HOLOTYPE: 111261, ♂ (stated to be a female in the original description),
Potrero Viejo, Veracruz, Mexico, D. M. Forbes, Jan. 11, 1939.

Clelia clelia immaculata Smith

Proc. U.S. Nat. Mus., vol. 92, p. 394, 1942.

HOLOTYPE: 24966, Guadalajara, Jalisco, Mexico, J. J. Major. This local-
ity was questioned by R. G. Zweifel.

Colorhogia redimita Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 81, 1862.

=*Arrhyton taeniatum* Günther

HOLOTYPE: 29769 (formerly 5747), eastern Cuba, C. Wright.

Coluber bairdi Yarrow

in Cope, U.S. Nat. Mus. Bull. 17, p. 41, 1880.

=*Elaphe obsoleta bairdi* (Yarrow)

HOLOTYPE: 10403, near Fort Davis, Jeff Davis County, Texas, W. F.
von Manteufel, 1878.

Coluber constrictor paludicola Auffenberg and Babbitt

Copeia, No. 1, p. 44, 1953.

HOLOTYPE: 131900, 3 miles west of Kendall, Dade County, Florida, L. H. Babbitt, Mar. 23, 1950.

PARATYPES (5): 28912, 30941, 32091, Lemon City, Dade County, Florida, J. E. Brown. 85304, 5 miles below Paradise Key, Dade County, Florida, A. Pfeuger, March 1932. 85305, Miami, Dade County, Florida, M. K. Brady, Apr. 4, 1932.

Coluber erythrogaster Forster

in Bossu, Travels through that part of America formerly called Louisiana, vol. 1, p. 364, 1771.

=*Natrix erythrogaster* (Forster)

NEOTYPE: 126890, near Parker's Ferry, Edisto River Swamp, Charleston County, South Carolina, R. Conant, Mar. 19, 1948.

Coluber flavirufus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 18 (1866), p. 319, 1867.

=*Elaphe flavirufa flavirufa* (Cope)

HOLOTYPE: 6566, Isamal, Yucatán, Mexico, A. Schott, 1865.

PARATYPE: 6626, Tabasco, Mexico, C. H. Berendt.

Coluber guttatus sellatus Cope

Proc. U.S. Nat. Mus., vol. 11, p. 387, 1889.

=*Elaphe guttata guttata* Linnaeus

COTYPES (2): 5507, Palatka, Putnam County, Florida, T. Glover. 9692, Arlington, Duval County, Florida, G. B. Goode.

Coluber (Masticophis) mentovarius centralis Roze

Herpetologica, vol. 9, p. 117, 1953.

HOLOTYPE: 115107, Maicao, Guajira, Colombia, A. Wetmore and M. A. Carriker, Jr., Apr. 16, 1941. This specimen was taken from the crop of a *Buteo albicaudatus colonus*.

Coluber mormon Baird and Girard

in Stansbury, Report Expl. and Surv. Valley of the Great Salt Lake of Utah . . . , Appendix C, p. 351, 1852.

=*Coluber constrictor mormon* Baird and Girard

HOLOTYPE: 2012, Great Salt Lake, Utah, H. Stansbury.

Coluber mutabilis Cope

Proc. Amer. Philos. Soc., vol. 22, p. 175, 1885.

=*Elaphe triaspis mutabilis* (Cope)

HOLOTYPE: 6745, Vera Paz, Guatemala, H. Hague.

PARATYPES (5): 9777, plateau of Costa Rica, J. C. Zeledon. 24729-30, 61066, Vera Paz, Guatemala, H. Hague. 30114, Guechecova, central part of Isthmus of Tehuantepec, Oaxaca, Mexico, F. Sumichrast.

Coluber obsoletus lemniscatus Cope

Proc. U.S. Nat. Mus., vol. 11, p. 386, 1889.

=*Elaphe obsoleta obsoleta* (Say)

COTYPES (2): 4710, skin, Whitfield County, Georgia. 8798, Augusta, Richmond County, Georgia, W. Phillips.

Coluber rhinomegas Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 12, p. 255, 1860. (Substitute name for *Scotophis calligaster* (not of Harlan), Kennicott, Proc. Acad. Nat. Sci. Philadelphia, vol. 11, p. 99, 1859.)

=*Elaphe guttata emoryi* (Baird and Girard)

HOLOTYPE: 2259, Hyatt (Anderson County), Kansas, S. Army.

Coluber rosaceus Cope

Proc. U.S. Nat. Mus., vol. 11, p. 388, pl. 36, figs. 3a-f, 1889.

=*Elaphe guttata rosacea* (Cope)

HOLOTYPE: 14418, Key West, Monroe County, Florida, H. Hemphill, June 20, 1885.

Coluber triaspis Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 18, p. 128, 1866.

=*Elaphe triaspis triaspis* (Cope)

HOLOTYPE: 24903, Belize, British Honduras, D. B. Parsons.

Coniophanes bipunctatus biseriatus Smith

Proc. Biol. Soc. Washington, vol. 53, p. 59, 1940.

HOLOTYPE: 108595, San Juanito, Palenque, Chiapas, Mexico, H. M. Smith, July 21, 1939.

PARATYPES (4): 109716-9, embryos, San Juanito, Palenque, Chiapas, Mexico, H. M. Smith, July-August 1939.

Coniophanes piceivittis Cope

Proc. Amer. Philos. Soc., vol. 11, p. 149, 1869.

=*Coniophanes piceivittis piceivittis* Cope

COTYPES (2): 30264-5, Chihuitán, Tehuantepec, Oaxaca, Mexico, F. Sumichrast.

Coniophanes proterops Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 12, p. 249, 1860.

=*Coniophanes fissidens proterops* Cope

COTYPE: 5285, Jalapa, Veracruz, Mexico, R. M. de Oca. One cotype was exchanged to ANSP.

Conophis concolor Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 18, p. 318, 1866.

=*Conophis lineatus concolor* Cope

COTYPES (2): 12368 (2), Yucatán, Mexico, A. Schott.

Conophis lineatus similis Smith

Journ. Washington Acad. Sci., vol. 31, p. 123, 1941. (Not of Bocourt, 1886.)

=*Conophis lineatus dumni* Smith

HOLOTYPE: 79963, ♀, Managua, Nicaragua, H. C. Kellers.

PARATYPES (3): 25237, Nicaragua, E. Flint. 37758, Esparta, Costa Rica, A. Alfaro, January 1906. 79964, Managua, Nicaragua, H. C. Kellers.

Conophis pulcher Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 20 (1868), p. 308, 1869.

=*Conophis pulcher pulcher* Cope

COTYPES (3): 6751 (2), 6803, Petén, Vera Paz, Guatemala, H. Hague.

Conophis pulcher plagosus Smith

Journ. Washington Acad. Sci., vol. 31, p. 121, 1941.

=*Conophis pulcher similis* Bocourt

HOLOTYPE: 109707, ♀, Tonalá, Chiapas, Mexico, H. M. Smith, Jan. 26-30, 1940.

Conophis sumichrastii Cope

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8 (1875), p. 137, 1876.

=*Conophis vittatus vittatus* Peters

COTYPES (2): 29123, Guadalajara?, Jalisco, Mexico, J. J. Major. 30258, western Tehuntepec, Oaxaca, Mexico (restricted to Guadalajara), F. Sumichrast. 29123 is the lectotype.

Conophis sumichrastii viduus Cope

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8 (1875), p. 137, 1876.

=*Conophis vittatus vittatus* Peters

HOLOTYPE: 30259, Tehuantepec, Oaxaca, Mexico, F. Sumichrast.

Conopsis biserialis Taylor and Smith

Univ. Kansas Sci. Bull., vol. 28, p. 33, figs. 2-3, 11, pl. 31, fig. 1, pl. 37, fig. 9, 1942.

PARATYPE: 137220, Kilometer 61 near Cuernavaca, Morelos, Mexico, E. H. Taylor and H. M. Smith, July 11, 1936.

Constrictor constrictor sigma Smith

Proc. U.S. Nat. Mus., vol. 93, p. 411, 1943.

HOLOTYPE: 46484, skin, María Madre Island, Tres Mariás Islands, Nayarit, Mexico, E. W. Nelson and E. A. Goldman, May 12, 1897.

PARATYPE: 24672, María Madre Island, Tres Mariás Islands, Nayarit, Mexico, E. W. Nelson, May 12, 1897.

Contia calligaster Cope

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8 (1875), p. 146, pl. 28, fig. 12, 1876.

—*Rhadinaea calligaster* (Cope)

COTYPES (2): 30607, 30679, Pico Blanco, Costa Rica, W. M. Gabb.

Contia isozona Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 18 (1866), p. 304, 1867.

—*Sonora semiannulata isozona* (Cope)

HOLOTYPE: 11417, Fort Whipple (Yavapai County), Arizona, E. Coues, 1865.

Contia mitis Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 110, 1853.

—*Contia tenuis* (Baird and Girard)

HOLOTYPE: 2034, San Jose, Santa Clara County, California, J. L. LeConte.

PARATYPE: 8075, Oregon, A. J. Skilton. 7289 from California, collected by the U.S. Exploring Expedition in 1841, the type of *Calamaria tenuis* Baird and Girard, is a probable paratype.

Contia pachyura Cope

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8 (1875), p. 175, 1876.

—*Rhadinaea pachyura pachyura* (Cope)

HOLOTYPE: 30618, Sipurio, Costa Rica, W. M. Cabb.

Coryphodon pantherinus Duméril and Bibron

Erpétologie générale ou histoire naturelle complète des reptiles, vol. 7, p. 181, 1854.

—*Dryadophis bifossatus bifossatus* (Raddi)

COTYPE: 391, Brazil, received from JPP.

Crotalophorus edwardsii Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 15, 1853.

—*Sistrurus catenatus edwardsii* (Baird and Girard)

COTYPES (2): 506, Sonora, Mexico, J. D. Graham. 509, Tamaulipas, Mexico, L. A. Edwards.

Crotalus adamanteus ruber Cope

Proc. U.S. Nat. Mus., vol. 14, p. 690, 1892.

—*Crotalus ruber ruber* Cope

HOLOTYPE: 9209, locality not given (restricted to the vicinity of San Diego, San Diego County, California).

Crotalus atrox Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 5, 1853.

HOLOTYPE: 7761, Indianola (Calhoun County), Texas, J. D. Graham.

PARATYPES (2): 255, skin, 7760, same data.

Crotalus atrox elegans Schmidt

Bull. Amer. Mus. Nat. Hist., vol. 46, p. 699, 1922.

=*Crotalus ruber ruber* Cope

HOLOTYPE: 64452, ♂, Angel de la Guarda Island, Gulf of California, Baja California, Mexico, C. H. Townsend, Apr. 10, 1911.

Crotalus basiliscus oaxacus Gloyd

Chicago Acad. Sci. Nat. Hist. Misc., No. 17, p. 1, figs. 1-3, 1948.

HOLOTYPE: 46467, ♀, Cerro San Felipe, near Oaxaca, Oaxaca, Mexico, 8,500 ft., E. W. Nelson and E. A. Goldman, June 20, 1894.

PARATYPE: 46468, ♀, San Pablo Yanganiza, Oaxaca, Mexico, 4,800 ft., E. W. Nelson and E. A. Goldman, July 2, 1894.

Crotalus cerastes Hallowell

Proc. Acad. Nat. Sci. Philadelphia, vol. 7, p. 95, 1854.

=*Crotalus cerastes cerastes* Hallowell

PROBABLE PARATYPE: 131801 (formerly 480), California, A. L. Heermann.

Crotalus cerastes cercobombus Savage and Cliff

Chicago Acad. Sci. Nat. Hist. Misc., No. 119, p. 2, 1953.

PARATYPE: 134309 (formerly UIMNH 5633), 21.6 miles south of Ajo, Pima County, Arizona, M. M. Hensley, June 26, 1949.

Crotalus confluentus abyssus Klauber

Trans. San Diego Soc. Nat. Hist., vol. 6, p. 114, pl. 11, fig. 1, 1930.

=*Crotalus viridis abyssus* Klauber

PARATYPES (2): 32725, skin, Grand Canyon, Arizona, H. S. Barber, May 10, 1904. 59747, Shinumo Creek, 3,000 ft., Grand Canyon, Arizona, E. A. Goldman, Sept. 10, 1916. These specimens were not listed by number in the original description, but Klauber considers them paratypes.

Crotalus confluentus lutosus Klauber

Trans. San Diego Soc. Nat. Hist., vol. 6, p. 100, pl. 10, fig. 1, 1930.

=*Crotalus viridis lutosus* Klauber

PARATYPE: 80069, 10 miles northwest of Abraham, Millard County, Utah, 4,650 ft., C. B. Perkins, May 12, 1929.

Crotalus confluentus stephensi Klauber

Trans. San Diego Soc. Nat. Hist., vol. 6, p. 108, 1930.

=*Crotalus mitchellii stephensi* Klauber

PARATYPES (13): 18661, Vegas Valley, Nevada, C. H. Merriam, May 1, 1891. 18662, Vegas Wash, Nevada, V. O. Bailey, May 2, 1891. 18663, juv., Indian Spring Valley, Nevada, C. H. Merriam, May 29, 1891. 18664, Grapevine Mountains above Salt Wells, Nevada, 3,000 ft., E. W. Nelson, June 6, 1891. 18665, Slate Range near summit of pass between Panamint Valley and Borax Flats, California, 3,100 ft., F. Stephens, Apr. 21, 1891. 18667, Johnson Canyon, Panamint Mountains, California, 5,000 ft., A. K. Fisher, Mar. 30, 1891. 18668-9, Emigrant Canyon, Panamint Mountains, California, 4,600 ft., F. Stephens, Apr. 16, 1891. 18670-1, Shepherd Canyon, Argus Range, California, A. K. Fisher, Apr. 29, 27, 1891. 18672-3, Coso Valley near Maturango Spring, California, T. S. Palmer, May 11, 1891. 18674, ♀, Independence Creek, Owen's Valley, California, 6,500 ft., F. Stephens, June 20, 1891.

***Crotalus durissus culminatus* Klauber**

Zool. Soc. San Diego Bull. 26, p. 65, 1952.

PARATYPE: 110610, skin, Xochicalco, Morelos, Mexico, H. M. Smith, Feb. 24, 1939.

***Crotalus durissus tzabcan* Klauber**

Zool. Soc. San Diego Bull. 26, p. 71, 1952.

PARATYPES (3): 6557 (2), Isamal, Mérida, Yucatán, Mexico, A. Schott. 46570, Chichén Itzá, Yucatán, E. W. Nelson and E. A. Goldman, February 1901.

***Crotalus gloydii lautus* Smith**

Univ. Kansas Sci. Bull., vol. 31, p. 75, figs. 1, 5, 9, 1946.

=*Crotalus intermedius intermedius* Troschel

HOLOTYPE: 110598, 1 km. east of El Limón, Totalco, Veracruz, Mexico, D. M. Forbes, Mar. 21, 1940.

***Crotalus goldmani* Schmidt**

Bull. Amer. Mus. Nat. Hist., vol. 46, p. 701, 1922.

=*Crotalus mitchellii pyrrhus* (Cope)

HOLOTYPE: 37573, ♂, skin, El Piñón, San Pedro Mártir Mountains, Baja California, Mexico, 5,300 ft., E. W. Nelson and E. A. Goldman, July 9, 1905.

***Crotalus jimenezii* Dugès**

Naturaleza, Mexico, vol. 4, p. 23, pl. 1, figs. 18-20, 1877.

=*Crotalus polystictus* Cope

COTYPE: 46508, juv., Guanajuato, Mexico, A. Dugès. This specimen was given to E. W. Nelson and E. A. Goldman by A. Dugès in 1896 as a "duplicate type" of *Crotalus jimenezii*.

Crotalus lecontei Hallowell

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 180, 1852.

=*Crotalus viridis viridis* (Rafinesque)

PARATYPE: 4224, juv., San Antonio, Bexar County, Texas, S. W. Woodhouse, March 1851.

Crotalus lucifer Baird and Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 177, 1852.

=*Crotalus viridis oreganus* Holbrook

COTYPES (4): 370-1, California, U.S. Exploring Expedition, 1841. 7762 (2), skins, Oregon, U.S. Exploring Expedition, 1841.

Crotalus molossus Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 10, 1853.

=*Crotalus molossus molossus* Baird and Girard

HOLOTYPE: 485, Fort Webster, Santa Rita del Cobre (Grant County), New Mexico, J. H. Clark.

Crotalus molossus estebanensis Klauber

Trans. San Diego Soc. Nat. Hist., vol. 11, p. 104, pl. 6, fig. 2, 1949.

PARATYPE: 64586, San Estéban Island, Gulf of California, Sonora, Mexico, C. H. Townsend, April 14, 1911.

Crotalus ornatus Hallowell

Proc. Acad. Nat. Sci. Philadelphia, vol. 7, p. 192, 1854.

=*Crotalus molossus molossus* Baird and Girard

HOLOTYPE: 486, skin, between El Paso and San Antonio, Pecos River, Texas, A. L. Heermann.

Crotalus rarus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 17, p. 191, 1865.

=*Sistrurus rarus* (Cope)

COTYPES (2): 25050-1, south tableland, Veracruz, Mexico, C. Sartorius.

Crotalus stejnegeri Dunn

Proc. Biol. Soc. Washington, vol. 32, p. 214, pl. 29, 1919.

HOLOTYPE: 46586, Plomosas, Sinaloa, Mexico, 2,500-3,000 ft., E. W. Nelson and E. A. Goldman, July 19, 1897.

PARATYPE: 46460, same data.

Crotalus tigris Kennicott

in Baird, Reptiles of the boundary, in Report U.S. and Mexican Bound.

Surv., vol. 2, pt. 2, p. 14, pl. 4, 1859.

HOLOTYPE: 471, Sierra Verde and Pozo Verde, Arizona, A. Schott.

PARATYPE: 472, same data.



***Crotalus triseriatus miquihuanus* Gloyd**

Chicago Acad. Sci. Spec. Publ. No. 4, p. 102, pl. 10, 1940.

=*Crotalus pricei miquihuanus* Gloyd

PARATYPE: 46402, ♂, Miquihuana, Tamaulipas, Mexico, 8,500 ft., E. W. Nelson and E. A. Goldman, June 14, 1898.

***Crotalus viridis caliginis* Klauber**

Trans. San Diego Soc. Nat. Hist., vol. 11, p. 90, pl. 5, fig. 2, 1949.

PARATYPE: 8564, Los Coronados Islands, Baja California, Mexico, T. H. Streets.

***Crotalus viridis nuntius* Klauber**

Trans. San Diego Soc. Nat. Hist., vol. 8, p. 78, pl. 8, fig. 1, map 1, 1935.

PARATYPES (5): 4251, San Francisco Mountains, Coconino County, Arizona, H. B. Möllhausen. 60240, 1 mile north of Winona, Coconino County, Arizona, 6,400 ft., C. M. Hoy, Aug. 11, 1917. 60241, 12 miles from mouth of Moencopie Wash, Coconino County, Arizona, 4,500 ft., C. M. Hoy, Aug. 3, 1917. 60244, Chin Lee, Apache County, Arizona, C. M. Hoy, Aug. 25, 1917. 73759, juv., Kayenta, Navajo County, Arizona, G. G. Cantwell, June 14, 1927.

***Crotalus willardi meridionalis* Klauber**

Trans. San Diego Soc. Nat. Hist., vol. 11, p. 131, 1949.

PARATYPE: 46332, Sierra Madre, Zacatecas, Mexico, E. W. Nelson and E. A. Goldman, Aug. 17, 1897.

***Crotalus willardi silus* Klauber**

Trans. San Diego Soc. Nat. Hist., vol. 11, p. 128, 1949.

PARATYPES (9): 26593, canyon, head of Río Pedras Verdes, Chihuahua, Mexico, 6900 ft., H. H. and C. S. Brimley, Aug. 31, 1898. 42496-7, headless skins, 105 km. south of Nueva Casas Grandes, District of Guerrero, Chihuahua, Mexico, E. C. Erdis. 42709, summit of Sierra Madres, Chihuahua, Mexico, 2,400 m., E. C. Erdis, June 1910. 46322-6, near Colonia Garcia, Chihuahua, Mexico, E. W. Nelson and E. A. Goldman, July 3, 1899.

***Dendroaspis jamesoni kaimosae* Loveridge**

Proc. Biol. Soc. Washington, vol. 49, p. 64, 1936.

=*Dendroaspis jamesoni kaimosae* Loveridge

PARATYPE: 49014, Kakamega (probably Kaimosi), Kenya, E. Heller, 1912.

***Dendrophidion vinitor* Smith**

Proc. Biol. Soc. Washington, vol. 54, p. 74, 1941.

HOLOTYPE: 110662, ♀, Piedras Negras, Petén, Guatemala, H. M. Smith, May 21, 1939.

PARATYPES (4): 7099, Tierra Caliente, Mexico, F. Sumichrast. 14215, 14220, Nicaragua, J. F. Bransford. 46589, Teapa, Tabasco, Mexico, E. W. Nelson and E. A. Goldman, Apr. 5, 1900.

Dendrophidium chloroticum Cope

Proc. Amer. Philos. Soc., vol. 23, p. 278, 1886.
 =*Drymobius chloroticus* (Cope)
 HOLOTYPE: 6755, Cobán, Guatemala, H. Hague.

Dendrophidium melanotropis Cope

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8 (1875), p. 134, pl. 26, fig. 1, 1876.
 =*Drymobius melanotropis* (Cope)
 HOLOTYPE: 32597, Costa Rica, W. M. Gabb.

Dendrophis prasinus Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 9 (1857), p. 181, 1858.
 =*Ahaetulla punctulata* (Gray)
 HOLOTYPE: 5535, New Holland (Sydney, Australia), U.S. Exploring Expedition, 1839.

Diadophis amabilis Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 113, 1853.
 =*Diadophis amabilis amabilis* Baird and Girard
 COTYPE: 2061, San Jose, Santa Clara County, California, J. L. LeConte.

Diadophis arnyi Kennicott

Proc. Acad. Nat. Sci. Philadelphia, vol. 11, p. 99, 1859.
 =*Diadophis punctatus arnyi* Kennicott
 HOLOTYPE: 1968, Hyatt (Anderson County), Kansas, S. Army.

Diadophis regalis Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 115, 1853.
 =*Diadophis regalis regalis* Baird and Girard
 HOLOTYPE: 2062, Sonora, Mexico, J. H. Clark.

Diadophis regalis arizonae Blanchard

Occ. Pap. Mus. Zool. Univ. Michigan, No. 142, p. 2, 1923.
 =*Diadophis regalis regalis* Baird and Girard
 HOLOTYPE: 62568, ♂, Sabino Canyon, Pima County, Arizona, G. H. Hofer, Sept. 15, 1919.

Diadophis regalis blanchardi Schmidt and Smith

Field Mus. Nat. Hist., Zool. Ser., vol. 29, p. 89, 1944.
 HOLOTYPE: 103641, Basin, Brewster County, Texas, Chisos Mountains, 5,200 ft., T. F. Smith, Aug. 7, 1936.

Diadophis texensis Kennicott

Proc. Acad. Nat. Sci. Philadelphia, vol. 12 (1860), p. 328, 1861.

=*Diadophis punctatus stictogenys* Cope

COTYPES (3): 1897, New Orleans (Louisiana) to Galveston, Texas, E. B. Andrews. 2076, head of Trinity River, Texas, J. Pope. 2155, Monticello, Lawrence County, Mississippi, H. Teunison.

Dinodon rufozonatum walli Stejneger

U.S. Nat. Mus. Bull. 58, p. 364, figs. 307-309, 1907.

HOLOTYPE: 34007, ♂, Ishigakishima, Yaeyama Group (Ryukyu Islands), A. Owston, June 1899.

PARATYPES (2): 34006, 34008, ♂, ♀, same data.

Dipsas gemmistrata latistrata Cope

U.S. Nat. Mus. Bull. 32, p. 68, 1887.

=*Imantodes latistratus* (Cope)

COTYPE: 24963, Guadalajara, Jalisco, Mexico, J. J. Major. This locality has been questioned by R. G. Zweifel.

Dipsas septentrionalis Kennicott

in Baird, Reptiles of the boundary, in Report U.S. and Mexican Bound. Surv., vol. 2, pt. 2, 1859.

=*Leptodeira annulata septentrionalis* (Kennicott)

COTYPES (4): 4267 (2), 4273, Matamoras, Tamaulipas, Mexico, D. N. Couch. 131739 (formerly 2288), Brownsville, Cameron County, Texas, S. Van Vliet.

Dipsas smithi Duméril and Bibron

Erpétologie générale ou histoire naturelle complète des reptiles, vol. 7, pt. 2, p. 1162, 1854.

=*Amplorhinus multimaculatus* Smith

COTYPES (2): 6217, 61227 (formerly 6216), Cape of Good Hope (South Africa), JPP.

Drepanodon attenuatus Barbour and Noble

Proc. U.S. Nat. Mus., vol. 58, p. 619, 1921.

HOLOTYPE: 60739, San Fernando, near Río Cosireni, Peru, 3,000 ft., E. Heller, August 1915. This specimen was taken from the alimentary tract of *Erythrolamprus aesculapii*.

Dromicus ignitus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 23, p. 201, 1871.

=*Rhadinaea decorata* (Günther)

COTYPES (2): 24501-2, Atlantic side, Isthmus of Darién (Panama-Colombia), T. O. Selfridge.

Dromicus leucomelas Duméril and Bibron

Erpétologie générale ou histoire naturelle complète des reptiles, vol. 7, pl. 1, p. 666, 1854.

=*Alsophis leucomelas* (Duméril and Bibron)

COTYPE: 416, Guadeloupe, West Indies, JPP.

Dromicus parvifrons rosamondae Cochran

Occ. Pap. Boston Soc. Nat. Hist., vol. 8, p. 186, 1934.

=*Leimadophis parvifrons rosamondae* (Cochran)

PARATYPES (3): 80919, Ile Vache, Haiti, C. R. Orcutt, June 5, 1929. 84292, Ile Vache, Haiti, L. H. Parish and W. M. Perrygo, 1930. 95116, Ile Vache, Haiti, Utowana Expedition, Apr. 12, 1934.

Dryadophis melanolomus stuarti Smith

Proc. U. S. Nat. Mus., vol. 93, p. 418, 1943.

HOLOTYPE: 110924, near Acapulco, Guerrero, Mexico, R. B. Smith, Sept. 3, 1939.

PARATYPE: 56283, Colima, Colima, Mexico, J. Hurter, July 2, 1902.

Dryadophis melanolomus tehuanae Smith

Proc. U. S. Nat. Mus., vol. 93, p. 420, 1943.

HOLOTYPE: 110917, Cerro Guengola, Oaxaca, Mexico, H. M. Smith, May 1940.

PARATYPES (13): 110909-13, 110915-6, Tres Cruces, Oaxaca, Mexico, H. M. Smith, January-March 1940. 110918-23, La Concepción, Oaxaca, Mexico, H. M. Smith, May 1940. 110914 was exchanged to MCZ in 1944.

Drymarchon corais rubidus Smith

Journ. Washington Acad. Sci., vol. 31, p. 474, 1941.

HOLOTYPE: 46430, ♀, Rosario, Sinaloa, Mexico, E. W. Nelson and E. A. Goldman, July 6, 1897.

PARATYPES (15): 24683, María Madre Island, Tres Mariás Islands, Gulf of California, Nayarit, Mexico, E. W. Nelson, May 18, 1897. 30425, Tehuantepec, Oaxaca, Mexico, F. Sumichrast. 46538, Acapulco, Guerrero, Mexico, E. W. Nelson and E. A. Goldman, Jan. 18, 1895. 46588, San Sebastián, Jalisco, Mexico, 3,900 ft., E. W. Nelson and E. A. Goldman, Mar. 17, 1897. 61948-9, Colima, Mexico, J. Xantus. 110877, San Diego, near Tehuacán, Puebla, Mexico, H. M. Smith, Aug. 27, 1939. 110878-9, Cerro Arenal, Oaxaca, Mexico, H. M. Smith, Jan. 10, 1940. 110880, Tres Cruces, Oaxaca, Mexico, H. M. Smith, Jan. 15, 1940. 110881-4, Tehuantepec, Oaxaca, Mexico, H. M. Smith, January-March 1940. 110885, Ixtepec, Oaxaca, Mexico, H. M. Smith, Jan. 22, 1940.

Drymarchon corais unicolor Smith

Journ. Washington Acad. Sci., vol. 31, p. 470, 1941.

HOLOTYPE: 110865, ♀, La Esperanza, Chiapas, Mexico, H. M. Smith, May 18, 1940.

PARATYPES (6): 6757a-b, Guatemala, H. Hague. 12687, Escuintla, Guatemala, H. J. Stuart, June 1882. 30424, ♀, Tehuantepec, Oaxaca, Mexico, F. Sumichrast. 46464, Huehuetán, Chiapas, Mexico, E. W. Nelson and E. A. Goldman, Feb. 26, 1896. 79960, Managua, Nicaragua, H. C. Kellers.

Drymobius aurigulus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 13, p. 301, 1861.

=*Masticophis aurigulus* (Cope)

HOLOTYPE: 5793, Cape San Lucas, Baja California, Mexico, J. Xantus.

Drymobius margaritiferus fistulosus Smith

Proc. U.S. Nat. Mus., vol. 92, p. 383, 1942.

HOLOTYPE: 51480, ♀, Miramar, Nayarit, Mexico, J. C. Thompson, 1913.

PARATYPES (7): 30484, Tehuantepec, Oaxaca, Mexico, F. Sumichrast. 31480-3, Colima, Mexico, J. Xantus. 56163, Colima, Mexico, J. Hurter, Aug. 5, 1902. 46545, Puente de Ixtla, Morelos, Mexico, E. W. Nelson and E. A. Goldman, June 6, 1903.

Dryophis brevirostris Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 12 (1860), p. 555, 1861.

=*Oxybelis brevirostris* (Cope)

HOLOTYPE: 31349, Veraguas, Panama, R. W. Mitchell.

Elaphe flavirufa matudai Smith

Copeia, No. 1, p. 132, 1941.

HOLOTYPE: 110303, ♀, Salto de Agua, Mount Ovando, Chiapas, Mexico, 1,200 ft., H. M. Smith, May 19, 1940.

Elaphe flavirufa phaescens Dowling

Occ. Pap. Mus. Zool. Univ. Michigan, No. 540, p. 7, fig. 2, 1952.

=*Elaphe triaspis phaescens* Dowling

PARATYPE: 46578, Chichén-Itzá, Yucatán, Mexico, E. W. Nelson and E. A. Goldman, February 1901.

Elaphe perlacea Stejneger

Proc. Biol. Soc. Washington, vol. 42, p. 129, 1929.

HOLOTYPE: 76257, ♂, Yachow, Szechwan, China, D. C. Graham, June 1928.

Elaphe quadrivittata deckerti Brady

Proc. Biol. Soc. Washington, vol. 45, p. 5, 1932.

=*Elaphe obsoleta quadrivittata* Holbrook

HOLOTYPE: 84295, ♂, Lower Matecumbe Key, Dade County, Florida, R. F. Deckert, June 1931.

Elapomorphus suspectus Amaral

Journ. Washington Acad. Sci., vol. 14, p. 202, 1924.

HOLOTYPE: 48939, ♂, Pilar, near Córdoba, Argentina, C. C. Craft.

Elaps distans Kennicott

Proc. Acad. Nat. Sci. Philadelphia, vol. 12, p. 338, 1860.

=*Micrurus distans distans* (Kennicott)

HOLOTYPE: 1144, Batosegatchic, Chihuahua, Mexico, A. Schott.

Elaps ehippifer Cope

Proc. Amer. Philos. Soc., vol. 23, p. 281, 1886.

=*Micrurus ehippifer* (Cope)

HOLOTYPE: 30085, Pacific side, Isthmus of Tehuantepec, Oaxaca, Mexico, F. Sumichrast.

Elaps euryxanthus Kennicott

Proc. Acad. Nat. Sci. Philadelphia, vol. 12, p. 337, 1860.

=*Micruroides euryxanthus* (Kennicott)

HOLOTYPE: 1122, Sonora, Mexico, T. H. Webb.

Elaps melanocephalus Hallowell

Proc. Acad. Nat. Sci. Philadelphia, vol. 12 (1860), p. 485, 1861.

=*Micrurus nigrocinctus nigrocinctus* (Girard)

HOLOTYPE: 7331, Nicaragua, C. Wright.

Elaps nigrocinctus Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 7, p. 226, 1854.

=*Micrurus nigrocinctus nigrocinctus* (Girard)

COTYPES (2): 7347 (2), Taboga Island, Bay of Panama (Panama), J. M. Gilliss.

Elaps tenere Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 22, 1853.

=*Micrurus fulvius tenere* (Baird and Girard)

COTYPE: 1119, Rio San Pedro of the Rio Grande, Texas, J. D. Graham.

Elaps tristis Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 23, 1853.

=*Micrurus fulvius tenere* (Baird and Girard)

COTYPES (2): 1123, Rio Grande west of San Antonio, Bexar County, Texas, S. Churchill. 1124, Kemper County, Mississippi, D. C. Lloyd.

Emydocephalus ijimae Stejneger

Journ. Sci. Coll. Imp. Univ. Tokyo, vol. 12, p. 223, 1898.

PARATYPE: 36517, Botel Tobago Island (Formosa), SCMT, 1888.

Epicrates crassus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 349, 1862.

=*Epicrates cenchria crassus* Cope

HOLOTYPE: 12413 (formerly 5409), Cadosa, Paraná River, Paraguay, C. Wood, La Plata Expedition.

Epicrates subflavus Stejneger

Proc. U.S. Nat. Mus., vol. 23, p. 469, 1901.

HOLOTYPE: 14507, Jamaica, ZSP, Nov. 25, 1885.

Erythrolamprus mimus micrurus Dunn and Bailey

Bull. Mus. Comp. Zool., vol. 86, p. 12, 1939.

PARATYPES (2): 11136, Isthmus of Darién (Panama), R. W. Shufeldt, 1870. 72353, San Juan River, Chocó Province, Colombia, E. J. Pampana.

Eudryas amarali Stuart

Copeia, No. 1, p. 7, 1938.

=*Dryadophis amarali* (Stuart)

HOLOTYPE: 22534, ♀, Margarita Island, Venezuela, W. Robinson, July 13, 1895.

PARATYPE: 10137, Tobago Island, West Indies, F. A. Ober, 1878.

Eurypholis semicarinatus Hallowell

Proc. Acad. Nat. Sci. Philadelphia, vol. 12 (1860), p. 493, 1861.

=*Liopeltis semicarinata* (Hallowell)

COTYPES (2): 7316 (2), ♂, ♀, near Napa, Loo Choo (Ryukyu Islands), C. Wright, November 1854.

Eutaenia angustirostris Kennicott

Proc. Acad. Nat. Sci. Philadelphia, vol. 12, p. 332, 1860.

=*Thamnophis angustirostris* (Kennicott)

HOLOTYPE: 959, Alamo de Parras, Coahuila, Mexico, D. N. Couch.

Eutaenia chrysocephala Cope

Proc. Amer. Philos. Soc., vol. 22, p. 173, 1885.

=*Thamnophis chrysocephalus* (Cope)

HOLOTYPE: 30494, Orizaba, Veracruz, Mexico, F. Sumichrast.

Eutaenia couchii Kennicott

in Baird, Report on reptiles collected on the survey, in Report U.S. Pacific R.R. Explor. and Surv., vol. 10, pt. 4, No. 4, p. 10, 1859.

=*Thamnophis elegans couchii* (Kennicott)

HOLOTYPE: 866, ♀, bank of Pitt River, California, (restricted to near mouth of Hat Creek), J. S. Newberry.

Eutaenia cyrtopsis Kennicott

Proc. Acad. Nat. Sci. Philadelphia, vol. 12, p. 333, 1860.

=*Thamnophis cyrtopsis cyrtopsis* (Kennicott)

HOLOTYPE: 8067 (formerly 930), Rinconada, Coahuila, Mexico, D. N. Couch, 1853.

PARATYPE: 8066, Durango, Mexico, probably collected by D. N. Couch.

Eutaenia cyrtopsis ocellata Cope

U.S. Nat. Mus. Bull 17, p. 22, 1880.

=*Thamnophis cyrtopsis cyrtopsis* (Kennicott)

COTYPES (2): 10528 (2), Helotes, Bexar County, Texas, G. W. Marnock.

Eutaenia elegans brunnea Cope

Proc. U.S. Nat. Mus., vol. 14, p. 654, 1892.

=*Thamnophis elegans biscutatus* (Cope)

HOLOTYPE: 10849, Camp Bidwell, Modoc County, California, H. W. Henshaw, July 1878.

PARATYPE: 10848, same data.

Eutaenia elegans lineolata Cope

Proc. U.S. Nat. Mus., vol. 14, p. 655, 1892.

=*Thamnophis elegans vagrans* (Baird and Girard)

HOLOTYPE: 8587, southern California, H. W. Henshaw, August 1875.

PARATYPES (21): 8579, Lake Tahoe, Nevada, H. W. Henshaw, September 1876. 8580 (2), southern California, H. W. Henshaw, 1875. 9565, eastern California, H. W. Henshaw. 10811, southern Oregon, H. W. Henshaw, July 1878. 10840, Fort Klamath (Klamath County), Oregon, H. W. Henshaw, August 29, 1878. 10843-44, Oregon, H. W. Henshaw, September 1878. 10845-46, Camp Warner, Oregon, H. W. Henshaw, Aug. 10, 1878. 10847, 10850, Camp Bidwell (Butte County), California, H. W. Henshaw, July 1878. 10911 (7), Fort Walla Walla (Walla Walla County), Washington, C. E. Bendire, June 1881. 11805, Baird, Shasta County, California, S. Stone. 12564, Fresno, Fresno County, California, G. Eisen.

Eutaenia flavilabris Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 18 (1866), p. 306, 1867.

=*Thamnophis eques eques* (Reuss)

COTYPES (2): 24991-2, tableland or southern mountains, Mexico, C. Sartorius.

Eutaenia hammondii Kennicott

Proc. Acad. Nat. Sci. Philadelphia, vol. 12, p. 332, 1860.

=*Thamnophis elegans hammondii* (Kennicott)

COTYPES (3): 894, ♂, San Diego, San Diego County, California, J. F. Hammond. 5496 (2), Fort Tejon (Kern County), California, J. Xantus.

Eutaenia henshawi Yarrow

Proc. U.S. Nat. Mus., vol. 6, p. 152, 1883.

=*Thamnophis elegans vagrans* (Baird and Girard)

HOLOTYPE: 10912, Fort Walla Walla (Walla Walla County), Washington, C. E. Bendire, June 1881.

Eutaenia infernalis vidua Cope

Proc. U.S. Nat. Mus., vol. 14, p. 658, 1892.

=*Thamnophis elegans atratus* (Kennicott)

COTYPES (2): 970 (2), San Francisco, San Francisco County, California, R. D. Cutts. These specimens are also cotypes of *Eutainia atrata* Kennicott, in Cooper, Report upon the reptiles collected on the survey, in Reports U.S. Expl. and Surv. . . . Mississippi River to the Pacific Ocean . . . 1853-5, vol. 12, pt. III, No. 4, p. 296, 1860.

Eutaenia insigniarum Cope

Proc. Amer. Philos. Soc., vol. 22, p. 172, 1885.

=*Thamnophis eques eques* (Reuss)

HOLOTYPE: 32163, castle of Chapultepec, Mexico, Mexico, E. D. Cope, 1884.

PARATYPE: 32164, Chapultepec, Mexico, A. L. Herrera, 1884.

Eutaenia macrostemma Kennicott

Proc. Acad. Nat. Sci. Philadelphia, vol. 12, p. 331, 1860.

=*Thamnophis eques eques* (Reuss)

COTYPES (2): 7247, 131775 (formerly 7247), City of Mexico, Distrito Federal, Mexico, W. Rich.

Eutaenia megalops Kennicott

Proc. Acad. Nat. Sci. Philadelphia, vol. 12, p. 330, 1860.

=*Thamnophis eques megalops* (Kennicott)

COTYPE: 965, "Tucson and Santa Magdalena" (restricted to Tucson, Pima County, Arizona), A. Schott.

Eutaenia pulchrilatus Cope

Proc. Amer. Philos. Soc., vol. 22, p. 174, 1885.

=*Thamnophis eques eques* (Reuss)

HOLOTYPE: 9899, Guanajuato, Mexico, A. Dugès.

Eutaenia radix twiningi Coues and Yarrow

Bull. U.S. Geol. Survey Terr., vol. 4, p. 279, 1878.

=*Thamnophis radix haydenii* (Kennicott)

HOLOTYPE: 9537, Two Forks of Milk River (Montana), E. Coues, July 15, 1874.

PARATYPES (4): 9538, Mouse River, North Dakota, E. Coues, Aug. 20, 1873. 9539, Turtle Mountain, North Dakota, E. Coues, July 22, 1873. 9540-1, near Frenchman's River, Montana, E. Coues, July 12, 8, 1874.

Eutaenia sirtalis semifasciata Cope

Proc. U.S. Nat. Mus., vol. 14, p. 662, 1892.

=*Thamnophis sirtalis sirtalis* (Linnaeus)

COTYPES (39): 8070 (30), Des Plaines, Cook County, Illinois, R. W. Kennicott. 1051 (7), Wisconsin, A. C. Berry. 1018 (2), Northfield, Cook County, Illinois, R. W. Kennicott.

Eutaenia sirtalis tetrataenia Cope

in Yarrow, Report upon the collections of batrachians and reptiles . . . 1871, 1872, 1873, and 1874, in Report U.S. Geogr. and Geol. Expl. and Surv. West of the One Hundr. Mer., vol. 5, p. 546, 1875.

=*Thamnophis sirtalis tetrataenia* (Cope)

COTYPES (2): 21383-4, locality not given (restricted to San Francisco Peninsula, California). The lectotype is 21384.

Eutaenia sumichrasti Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 18 (1866), p. 306, 1867.

=*Thamnophis sumichrasti sumichrasti* (Cope)

COTYPES (2): 26501-2, Orizaba, Veracruz, Mexico, F. Sumichrast.

Eutaenia vagrans plutonia Yarrow

Proc. U.S. Nat. Mus., vol. 6, p. 152, 1883.

=*Thamnophis elegans vagrans* (Baird and Girard)

HOLOTYPE: 8171, Arizona (restricted to northwestern Arizona), F. Bischoff, 1871.

Eutainia atrata Kennicott

in Cooper, Report upon the reptiles collected on the survey, in Reports U.S. Expl. and Surv. . . . Mississippi River to the Pacific Ocean . . . 1853-5, vol. 12, pt. III, No. 4, p. 296, 1860.

=*Thamnophis elegans atratus* (Kennicott)

COTYPES (2): 970 (2), San Francisco, San Francisco County, California, R. D. Cutts. These specimens are also cotypes of *Eutaenia infernalis vidua* Cope, Proc. U.S. Nat. Mus., vol. 14, p. 658, 1892. One cotype was exchanged to ANSP.

Eutainia elegans Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 34, 1853.

=*Thamnophis elegans elegans* (Baird and Girard)

HOLOTYPE: 882, ♂, El Dorado County, California, C. C. Boyle.

Eutainia faireyi Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 25, 1853.

=*Thamnophis sauritus proximus* (Say)

COTYPE: 752, Prairie Mer Rouge, Morehouse Parish, Louisiana, J. Fairie. One cotype was exchanged to MCZ; one to TCM.

Eutainia haydenii Kennicott

in Cooper, Report upon the reptiles collected on the survey, in Reports U.S. Expl. and Surv. . . . Mississippi River to the Pacific Ocean . . . 1853-5, vol. 12, pt. III, No. 4, p. 298, pl. 14, 1860.

=*Thamnophis radix haydenii* (Kennicott)

HOLOTYPE: 707, Fort Pierre (Stanley County), South Dakota, J. Evans. Since the type was in poor condition, a neotype, 128137, and a paraneotype, 128138, were designated, both from Whitlock Crossing, Davey County, South Dakota, collected by A. M. Jackley et al. on Oct. 6, 1947.

Eutainia leptcephala Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 29, 1853.

=*Thamnophis ordinoides* (Baird and Girard)

COTYPES (3): 961a-c, Puget Sound (Washington), U.S. Exploring Expedition, 1841.

Eutainia marciana Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 36, 1853.

=*Thamnophis marcianus* (Baird and Girard)

HOLOTYPE: 844, Indian Territory between Camp 5 and Red River (Tillman County near Hollister, Oklahoma, fide H. A. Dundee), R. B. Marcy, May 20, 1852.

PROBABLE PARATYPE: 1417, Indianola (Calhoun County), Texas, J. D. Graham.

Eutainia phenax Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 20, p. 134, 1868.

=*Thamnophis phenax phenax* (Cope)

COTYPES (2): 30498-9, Córdoba, Veracruz, Mexico, F. Sumichrast.

Eutainia pickeringi Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 27, 1853.

=*Thamnophis sirtalis pickeringi* (Baird and Girard)

HOLOTYPE: 936a, Puget Sound (Washington), U.S. Exploring Expedition, 1841.

PARATYPE: 936b, same data.

Eutainia radix Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 34, 1853.

=*Thamnophis radix radix* (Baird and Girard)

HOLOTYPE: 719, "Racine, Wisconsin" (restricted to vicinity of Chicago, Cook County, Illinois), P. R. Hoy.

Eutainia rutilorus Cope

Proc. Amer. Philos. Soc., vol. 22, p. 388, 1885.

=*Thamnophis sauritus chalconus* (Cope)

HOLOTYPE: 13906, Cozumel Island, Yucatán, Mexico, U.S. Fish Commission *Albatross*, Jan. 23-29, 1885.

Eutainia sirtalis obscura Cope

in Yarrow, Report upon the collections of batrachians and reptiles . . . 1871, 1872, 1873, and 1874, in Report U.S. Geogr. and Geol. Expl. and Surv. West of the One Hundr. Mer., vol. 5, p. 546, 1875.

=*Thamnophis sirtalis sirtalis* (Linnaeus)

COTYPES (5): 974 (5), Westport, Essex County, New York, S. F. Baird.

Eutainia sirtalis trilineata Cope

Proc. U.S. Nat. Mus., vol. 14, p. 665, 1892.

=*Thamnophis sirtalis trilineata* (Cope)

COTYPES (6): 5275, Port Townsend (Jefferson County), Washington, J. L. White. 5493 (5), Bitter Root Valley, west of Fort Benton, Montana, J. Mullan. The lectotype is 5275.

Eutainia vagrans Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 35, 1853.

=*Thamnophis elegans vagrans* (Baird and Girard)

COTYPES (2): 907, Humboldt River, California, J. S. Bowman. 908, ♀, "California" (restricted to Utah), W. Gambel. 907 is now considered to be *Thamnophis elegans elegans* (Baird and Girard).

Ficimia publia Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 18, p. 126, 1866.

=*Ficimia publia publia* Cope

COTYPES (2): 16427-8, Yucatán, Mexico, A. Schott, July 28, 1865.

Ficimia publia taylori Smith

Journ. Washington Acad. Sci., vol. 37, p. 411, 1947.

HOLOTYPE: 123491, ♂, Río Chiquito at San Lorenzo, Veracruz, Mexico, M. W. Stirling, February-April 1946.

PARATYPE: 123492, Río Chiquito at San Lorenzo, Veracruz, Mexico, M. W. Stirling, February-April 1946.

Geagras redimitus Cope

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8 (1875), p. 141, 1876.

HOLOTYPE: 30115, Tehuantepec, west side, Oaxaca, Mexico, F. Sumichrast.

Geophis anocularis Dunn

Proc. Biol. Soc. Washington, vol. 33, p. 127, 1920.

HOLOTYPE: 46556, ♂, Totontepec, Oaxaca, Mexico, E. W. Nelson and E. A. Goldman, July 26, 1894.

Geophis blanchardi Taylor and Smith

Univ. Kansas Sci. Bull., vol. 25 (1938), p. 245, fig. 2, 1939.

PARATYPE: 139732 (formerly UIMNH 27382), 2 miles southwest of Acultzingo, Veracruz, Mexico, E. H. Taylor, July 19, 1938.

Geophis cancellatus Smith

Smithsonian Misc. Coll., vol. 99, No. 19, p. 1, 1941.

HOLOTYPE: 46440, ♀, Chicharras, Chiapas, Mexico, 2,500-3,000 ft., E. W. Nelson and E. A. Goldman, February 1896.

PARATYPE: 46441, probably same data.

Gyalopion canum Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 12, p. 243, 1860.

=*Ficimia cana* (Cope)

HOLOTYPE: 5284 (formerly 4675), Fort Buchanan, Arizona, B. J. D. Irwin.

Helicops leprieurii Duméril and Bibron

Erpétologie générale ou histoire naturelle complète des reptiles, vol. 7, pt. 1, p. 750, pl. 68, 1854.

=*Helicops leopardinus* (Schlegel)

COTYPE: 394, Cayenne, French Guiana, JPP.

Herpetodryas bernieri Duméril and Bibron

Erpétologie générale ou histoire naturelle complète des reptiles, vol. 7, pt. 1, p. 211, pl. 66, figs. 1-4, 1854.

=*Dromicodryas bernieri* (Duméril and Bibron)

COTYPE: 6221, Madagascar, JPP.

Herpetodryas chloris Hallowell

Proc. Acad. Nat. Sci. Philadelphia, vol. 12 (1860), p. 503, 1861.

=*Eurypholis major* (Günther)

HOLOTYPE: 7335 ♂, Hongkong Island, (off the mainland of China), J. M. Brooke, May 1854.

Herpetodryas melas Cope

Proc. Amer. Philos. Soc., vol. 23, p. 278, 1886.

=*Chironius melas* (Cope)

HOLOTYPE: 14219, Nicaragua, J. F. Bransford.

Heterodon atmodes Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 57, 1853.

=*Heterodon platyrhinos* Latreille

COTYPES (4): 1158, Georgia, C. B. Adams. 1152-3, Charleston, Charleston County, South Carolina, S. B. Barker. 1199, Charleston, Charleston County, South Carolina, C. Girard.

Heterodon browni Stejneger

Proc. Biol. Soc. Washington, vol. 16, p. 123, 1903.

=*Heterodon platyrhinos* Latreille

HOLOTYPE: 32089, ♀, Lemon City, Dade County, Florida, E. J. Brown, 1903.

PARATYPES (2): 30925-6, ♂, ♂, Lemon City, Dade County, Florida, E. J. Brown, May 17, 1901.

Heterodon cognatus Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 54, 1853.

=*Heterodon nasicus nasicus* Baird and Girard

COTYPE: 1271, skin, Indianola (Calhoun County), Texas, J. D. Graham.

Heterodon dorbignyi Duméril and Bibron

Erpétologie générale ou histoire naturelle complète des reptiles, vol. 7, pt. 1, p. 772, 1854.

=*Lystrophis dorbignyi* (Duméril and Bibron)

COTYPE: 420, Brazil, JPP.

Heterodon kennerlyi Kennicott

Proc. Acad. Nat. Sci. Philadelphia, vol. 12, p. 336, 1860.

=*Heterodon nasicus kennerlyi* (Kennicott)

COTYPES (3): 1282 (2), "Sonora" (restricted to Lower Rio Grande, Texas, and Sonora, Mexico), D. N. Couch. 7290, Lower Rio Grande, Texas, A. Schott.

Heterodon nasicus gloydi Edgren

Chicago Acad. Sci. Nat. Hist. Misc., No. 112, p. 3, 1952.

HOLOTYPE: 5083a, ♀, Wheelock, Robertson County, Texas, F. Kellogg.

PARATYPES (2): 5083b-c, ♂, ♂, same data.

Himantodes leucomelas Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 13, p. 296, 1861.

=*Imantodes cenchoa leucomelas* Cope

COTYPES (2): 25035-6, Mirador, Veracruz, Mexico, C. Sartorius.

Himantodes tenuissimus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 18 (1866), p. 317, 1867.

=*Imantodes tenuissimus* Cope

HOLOTYPE: 6563, Merida, Yucatán, Mexico, A. Schott, Jan. 7, 1866.

Hydrophis ornatus maresinensis Mittleman

Proc. Biol. Soc. Washington, vol. 60, p. 3, 1947.

HOLOTYPE: 33933, ♀, Ishigaki Shima, Ryu Kyu (Ryukyu) Archipelago, A. Owston, April-May, 1899.

PARATYPES (2); 33935-6, ♂, ♂, same data. 33934 was given to TUM in 1950.

Hydrops lubricus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 23, p. 217, 1871.

=*Coniophanes quinquevittatus* Duméril and Bibron

HOLOTYPE: 61182, Coatzacoalcos River, Veracruz, Mexico, T. H. Streets.

Hydrops trianguligaris venezuelensis Roze

Acta Biol. Venezuelica, Univ. Centr. Venezuela, vol. 2, p. 78, fig. 13c, 1957.

PARATYPE: 56235, ♂, Ciudad Bolívar, Estado Bolívar, Venezuela, J. Hurter, 1895.

Hypsiglena chlorophaea Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 12, p. 247, 1860.

=*Hypsiglena torquata ochrorhyncha* (Cope)

COTYPE: 4676, Fort Buchanan, Arizona, B. J. D. Irwin.

Hypsiglena ochrorhynchus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 12, p. 246, 1860.

=*Hypsiglena torquata ochrorhyncha* Cope

COTYPES (9): 5283 (4), 9216, 69538-40, 131733, Cape San Lucas, Baja California, Mexico, J. Xantus, December 1859. 9216 was "captured under the cabin's poop of a wrecked vessel," according to Xantus' field notes; the rest were "captured under dead palm leaves." One cotype from 5283 was exchanged to MCZ.

Hypsiglena o[chrorhyncha] deserticola Tanner

Great Basin Nat., vol. 5, p. 59, pl. 2, fig. 3, 1944.

=*Hypsiglena torquata deserticola* Tanner

PARATYPE: 125552 (formerly BYU 2198), 36.4 miles northwest of Chimney Rock Pass, Utah County, Utah, W. W. Tanner, Aug. 11, 1939.

Hypsiglena texana Stejneger

North Amer. Fauna, No. 7, p. 205, 1893.

=*Hypsiglena torquata texana* Stejneger

HOLOTYPE: 1782, between Laredo (Webb County) and Camargo, Texas, A. Schott.

Hypsiglena torquata baueri Zweifel

Amer. Mus. Novit., No. 1895, p. 12, fig. 1a-c, 1958.

PARATYPE: 59371, ♀, northeast side of Cedros Island, Baja California, J. C. Thompson, Oct. 8, 1916.

***Ialtris parishi* Cochran**

Proc. Biol. Soc. Washington, vol. 45, p. 189, 1932.

HOLOTYPE: 80773, 10 miles east of Baradères, Haiti, L. H. Parish and W. M. Perrygo, Apr. 7, 1930.

***Imantodes splendidus oliveri* Smith**

Proc. U.S. Nat. Mus., vol. 92, p. 388, 1942.

PARATYPES (12): 12089, Mexico, C. Sartorius. 12443, 30164-6, 30178-9, Tehuantepec, Oaxaca, Mexico, F. Sumichrast. 30386-8, Juchitán, Oaxaca, Mexico, F. Sumichrast. 32171, Mexico, E. D. Cope. 110528, Tonalá, Chiapas, Mexico, H. M. Smith, Jan. 26-30, 1940.

***Lachesis barbouri* Dunn**

Proc. Biol. Soc. Washington, vol. 32, p. 213, 1919.

=*Bothrops barbouri* (Dunn)

HOLOTYPE: 46347, Omilteme, Guerrero, Mexico, 7,000 ft., E. W. Nelson and E. A. Goldman, May 20, 1903.

***Lachesis insularis* Amaral**

Anexos Mem. Inst. Butantan, vol. 1, p. 62, pls. 3-4, 1921.

=*Bothrops insularis* (Amaral)

PARATYPES (2): 66490-1, Ilha de Queimada Grande, São Paulo, Brazil, A. do Amaral.

***Lachesis stenophrys* Cope**

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8, p. 152, 1875.

=*Lachesis muta stenophrys* Cope

HOLOTYPE: 32479, Sipurio, Costa Rica, W. M. Gabb.

***Lampropeltis amaura* Cope**

Proc. Acad. Nat. Sci. Philadelphia, vol. 12, p. 258, 1860.

=*Lampropeltis triangulum amaura* Cope

HOLOTYPE: 5282, Mississippi? (restricted to New Orleans, Orleans Parish, Louisiana).

***Lampropeltis annulata* Kennicott**

Proc. Acad. Nat. Sci. Philadelphia, vol. 12, p. 329, 1860.

=*Lampropeltis triangulum annulata* Kennicott

PARATYPES (2): 425, Mexico, JPP. 1855, Brownsville, Cameron County, Texas. The holotype, originally USNM 4293, from Matamoras, Tamaulipas, Mexico, collected by J. L. Berlandier, is now ANSP 3613.

***Lampropeltis boylii conjuncta* Cope**

Proc. Acad. Nat. Sci. Philadelphia, vol. 13, p. 301, 1861.

=*Lampropeltis getulus conjuncta* Cope

COTYPES (4): 5288 (3), 131735, Cape San Lucas, Baja California, Mexico, J. Xantus.

Lampropeltis elapsoides virginiana Blanchard

Occ. Pap. Mus. Zool. Univ. Michigan, No. 81, p. 2, 1920.

=*Lampropeltis triangulum temporalis* (Cope)

HOLOTYPE: 21163, Raleigh, Wake County, North Carolina, H. H. and C. S. Brimley, June 16, 1893.

PARATYPES (4): 4466, ♀, Appomattox County, Virginia, C. W. Statham. 25321, ♂, Alberene, Albemarle County, Virginia, R. E. Shaw, July 1898. 26181, ♂, Cuscowilla, Mecklenburg County, Virginia, W. Townes. 56197, Raleigh, Wake County, North Carolina, J. Hurter, May 18, 1878.

Lampropeltis getulus floridanus Blanchard

Occ. Pap. Mus. Zool. Univ. Michigan, No. 70, p. 1, pl. 1, fig. 1, 1919.

=*Lampropeltis getulus floridana* Blanchard

HOLOTYPE: 22368, ♀, Orange Hammock, De Soto County, Florida, W. Palmer, March 1895.

PARATYPES (6): 5509, Charlotte Harbor, Charlotte County, Florida, E. B. Baker. 22367, Orange Hammock, De Soto County, Florida, W. Palmer. 36478-9, Florida, J. H. Egbert. 36564, skin, Lemon City, Dade County, Florida, E. J. Brown, May 10, 1904. 61276, Florida.

Lampropeltis getulus yumensis Blanchard

Occ. Pap. Mus. Zool. Univ. Michigan, No. 70, p. 6, pl. 1, fig. 2, 1919.

HOLOTYPE: 61318, ♂, 27 miles west of Indian Oasis, Pima County, Arizona, A. B. Howell, July 7, 1918.

PARATYPES (7): 4284, Fort Yuma (Imperial County), California, G. H. Thomas, 1855. 21720, ♂, San Domingo, Sonora, Mexico, E. A. Mearns, Jan. 16, 1894. 37532, Volcano Lake, Baja California, Mexico, E. W. Nelson and E. A. Goldman, Apr. 19, 1905. 37534, Yuma (Yuma County), Arizona, H. Brown, 1905. 44530, Yuma, Arizona, V. O. Bailey, Apr. 24, 1889. 51761, Ash Creek, Graham County, Arizona, E. G. Holt, June 23, 1914. 54717, Calva, Graham County, Arizona, W. P. Taylor, June 8, 1916. 15969 was exchanged to MCZ in 1928.

Lampropeltis multistriata Kennicott

Proc. Acad. Nat. Sci. Philadelphia, vol. 12, p. 328, 1860.

=*Lampropeltis triangulum gentilis* (Baird and Girard)

HOLOTYPE: 1842, Fort Lookout (Lyman County), South Dakota, F. V. Hayden.

Lampropeltis pyromelana woodini Tanner

Great Basin Nat., vol. 13, p. 54, 1953.

PARATYPE: 22195, Fort Huachuca (Cochise County), Arizona, A. K. Fisher, May 18, 1892.

Lampropeltis pyrrhomelaena celaenops Stejneger

Proc. U.S. Nat. Mus., vol. 25, p. 153, 1902.

=*Lampropeltis triangulum celaenops* Stejneger

HOLOTYPE: 22375, Mesilla Valley (Dona Ana County), New Mexico, H. B. Lane.

Lampropeltis ruthveni Blanchard

Occ. Pap. Mus. Zool. Univ. Michigan, No. 81, p. 8, pl. 1, fig. 2, 1920.

HOLOTYPE: 46558, Pátzcuaro, Michoacan, Mexico, E. W. Nelson, Aug. 2, 1892.

Lampropeltis triangulum nelsoni Blanchard

Occ. Pap. Mus. Zool. Univ. Michigan, No. 81, p. 6, pl. 1, fig. 1, 1920.

HOLOTYPE: 46552, Acámbaro, Guanajuato, Mexico, E. W. Nelson, October 1892.

PARATYPES (9): 12680 ♂, Guanajuato, Mexico, A. Dugès, 1881. 24684, María Madre Island, Tres Mariás Islands, Nayarit, Mexico, E. W. Nelson, May 16, 1897. 24967-8, Guadalajara, Jalisco, Mexico, J. J. Major. 30504-5, ♂, Mexican plateau, Mexico, F. Sumichrast. 31491, Neshpa River, Sierra Madre, Michoacan, Mexico, J. Xantus, April 1863. 31492-3, Colima, Mexico, J. Xantus, July 1863. 31494 was exchanged to MCZ in 1928. 24684 is also a paratype of *Lampropeltis triangulum schmidti* Stuart.

Lampropeltis triangulum schmidti Stuart

Occ. Pap. Mus. Zool. Univ. Michigan, No. 323, p. 2, 1935.

PARATYPE: 24684, María Madre Island, Tres Mariás Islands, Nayarit, Mexico, E. W. Nelson, May 16, 1897. This specimen is also a paratype of *Lampropeltis triangulum nelsoni* Blanchard.

Lampropeltis zonata parvirubra Zweifel

Copeia, No. 3, p. 160, pl. 1, fig. C., 1952.

PARATYPES (8): 31176-7, San Bernardino Mountains, California, C. C. Zeus. 44423, San Jacinto Mountains, California, V. O. Bailey, Aug. 12, 1907. 50677, San Bernardino County, California, 4,000 ft., E. S. Shepherd. 54987, Los Angeles County, California, E. J. Brown, 1916. 75184-5, Banning, Riverside County, California, G. Jessup and M. Morris, June 17, 1924, July 22, 1926. 84038, Tujunga Canyon, Los Angeles County, California.

Lampropeltis zonata pulchra Zweifel

Copeia, No. 3, p. 162, 1952.

PARATYPE: 13889, San Diego, San Diego County, California, C. R. Orcutt, June 16, 1884.

Lamprosoma annulatum Baird

Reptiles of the boundary, *in* Report U.S. and Mexican Bound. Surv., vol. 2, pt. 2, p. 22, pl. 21, fig. 1, 1859.

=*Chionactis occipitalis annulata* (Baird)

COTYPES (2): 2105-6, Colorado Desert, California, A. Schott.

Lamprosoma episcopum Kennicott

in Baird, Reptiles of the boundary, *in* Report U.S. and Mexican Bound. Surv., vol. 2, pt. 2, p. 22, pl. 8, fig. 2, 1859.

=*Sonora episcopa episcopa* (Kennicott)

LECTOTYPE: 2042, Eagle Pass, Maverick County, Texas, A. Schott.

Leimadophis alleni Dunn

Proc. New England Zool. Club, vol. 7, p. 40, 1920.

=*Leimadophis parvifrons alleni* Dunn

PARATYPES 10170, ♀, Gonaves Island, Haiti, J. J. Brown.

Leimadophis nebulatus Barbour

Ann. Carnegie Mus., vol. 10, p. 305, pl. 28, figs. 1-2, 1916.

=*Leimadophis andreae nebulatus* Barbour

PARATYPE: 120824 (formerly MCZ 11158), Los Indios, Isle of Pines, Cuba, G. A. Link.

Leimadophis parvifrons lincolni Cochran

Proc. Biol. Soc. Washington, vol. 44, p. 91, 1931.

HOLOTYPE: 83890, Beata Island, Dominican Republic, A. Wetmore and F. C. Lincoln, May 15, 1931.

Leimadophis parvifrons niger Dunn

Proc. New England Zool. Club, vol. 7, p. 39, 1920.

PARATYPES (7): 55026-32, Samaná Peninsula, Dominican Republic, W. L. Abbott, 1916.

Leimadophis stahli Stejneger

Report of the U.S. National Museum . . . for the year ending June 30, 1902, p. 695, fig. 161, 1904.

HOLOTYPE: 27323, ♀, Bayamón, Puerto Rico, A. Stahl, 1900.

PARATYPES (5): 7109, ♂, Puerto Rico, G. Latimer. 25529, ♂, Adjuntas, Puerto Rico, A. B. Baker, Feb. 3, 1899. 25724, ♂, El Yunque, eastern slope, Puerto Rico, U.S. Fish Commission, Feb. 19, 1899. 26883, ♀, Catalina Plantation, Puerto Rico, L. Stejneger, Mar. 2, 1900. 27768, ♀, Humacao, Puerto Rico, L. M. McCormick, 1900. 25616 was exchanged to R in 1953.

Leimadophis tortuganus Dunn

Proc. New England Zool. Club, vol. 7, p. 40, 1920.

=*Leimadophis parvifrons tortuganus* Dunn

HOLOTYPE: 59440, ♀, Tortuga Island, Haiti, W. L. Abbott, Feb. 7, 1917.

PARATYPE: 59439, ♀, Tortuga Island, Haiti, W. L. Abbott, Feb. 4, 1917.

Lepidocephalus fasciatus Hallowell

Proc. Acad. Nat. Sci. Philadelphia, vol. 12 (1860), p. 498, 1861.

=*Dinodon semicarinatum* (Cope)

COTYPES (2): 7354 (2), Japan and Loo Choo (Ryukyu Islands), Perry Expedition.

Leptodeira annulata pulchriceps Duellman

Bull. Amer. Mus. Nat. Hist., vol. 114, Art. 1, p. 51, 1958.

PARATYPES (2): 78335-6, Pôrto Esperanza, Mato Grosso, Brazil, IB.

Leptodeira annulata taylori Smith

Proc. Biol. Soc. Washington, vol. 54, p. 115, 1941.

=*Leptodeira septentrionalis polysticta* Günther

HOLOTYPE: 30208, Orizaba, Veracruz, Mexico, F. Sumichrast.

PARATYPES (10): 7088, 30207, 30508, Orizaba, Veracruz, Mexico, F. Sumichrast. 12113 (4), Mirador, Veracruz, Mexico, C. Sartorius. 65154, hills west of Veracruz, Veracruz, Mexico, C. Sartorius. 111224, Agua del Obispo, Guerrero, Mexico, H. M. Smith, Feb. 4, 1939. 111257, Cerro Gordo, Veracruz, Mexico, H. M. Smith, Mar. 24, 1940.

Leptodeira ephippiata Smith and Tanner

Copeia, No. 3, p. 131, 1944.

=*Leptodeira splendida ephippiata* Smith and Tanner

PARATYPE: 46459, ♀, Plomosas, Sinaloa, Mexico, 2,500-3,000 ft., E. W. Nelson and E. A. Goldman, July 19, 1897.

Leptodeira yucatanensis malleisi Dunn and Stuart

Occ. Pap. Mus. Zool. Univ. Michigan, No. 313, p. 1, 1935.

PARATYPES (8): 24901, Belize, British Honduras, D. B. Parsons. 24908, Belize, British Honduras. 26354, Belize, British Honduras, W. Bennett. 56010-2, Belize, British Honduras, J. Hurter, May 21, 1914. 71367, 71369, Libertad, Petén, Guatemala, H. Malleis, Sept. 12, Oct. 8, 1923.

Leptodira mystacina Cope

Proc. Amer. Philos. Soc., vol. 11 (1869), p. 151, 1870.

=*Leptodeira mystacina* Cope

COTYPES (2): 30339-40, western region near Isthmus of Tehuantepec, Oaxaca, Mexico, F. Sumichrast.

Leptodira pacifica Cope

Proc. Acad. Nat. Sci. Philadelphia, vol 20 (1868), p. 310, 1869.

=*Leptodeira punctata* (Peters)

HOLOTYPE: 6833, Mazatlán, Sinaloa, Mexico, F. Bischoff.

Leptodira personata Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 20 (1868), p. 310, 1869.

=*Leptodeira maculata* (Hallowell)

HOLOTYPE: 6836, Mazatlán, Sinaloa, Mexico, F. Bischoff.

Leptognathus argus Cope

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8 (1875), p. 130, pl. 27, fig. 1, pl. 28, fig. 7, 1876.

=*Dipsas argus* (Cope)

HOLOTYPE: 30656, Sipurio, Costa Rica, W. M. Gabb.

Leptognathus pictiventris Cope

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8 (1875), p. 130, pl. 28, fig. 8, 1876.

=*Dipsas pictiventris* (Cope)

HOLOTYPE: 30657, head and tail only, eastern Costa Rica, W. M. Gabb.

Leptognathus stratissima Cope

Proc. Amer. Philos. Soc., vol. 23, p. 280, 1886.

=*Imantodes elegans* Jan

HOLOTYPE: 14121, Panama, G. W. Nelson, May 9, 1885.

Leptognathus turgida Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 20, p. 136, 1868.

=*Sibynomorphus turgidus* (Cope)

COTYPES (2): 5815 (2), northern part of the Paraguay River (boundary of Brazil and Paraguay), T. J. Page.

Leptophis aeruginosus Cope

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8 (1875), p. 132, pl. 28, fig. 9, 1876.

HOLOTYPE: 30684, Costa Rica, W. M. Gabb. Identification of this specimen is doubtful because it is in a poor state of preservation.

Leptophis coeruleodorsus Oliver

Occ. Pap. Mus. Zool. Univ. Michigan, No. 462, p. 4, 1942.

=*Leptophis ahaetulla coeruleodorsa* Oliver

PARATYPES (9): 5587, Trinidad, West Indies, T. N. Gill. 15235, Trinidad, West Indies, H. Caracciola. 17746, Trinidad, West Indies, G. T. Wayman,

1891. 27821, Macuto, Venezuela, M. W. Lyon and W. Robinson, Aug. 8,
1900. 27831, Río Chico, Venezuela, L. Goldschmidt. 59931-3, Trinidad,
West Indies, H. D. Baker. 60598, Trinidad, West Indies, Mrs. J. B. Rorer.

Leptophis copei Oliver

Occ. Pap. Mus. Zool. Univ. Michigan, No. 462, p. 7, 1942.

=*Leptophis ahaetulla copei* Oliver

HOLOTYPE: 83564, ♂, Salto da Huá, Brazil-Venezuelan Boundary, E. G. Holt, Nov. 20, 1930.

PARATYPES (2): 83570, Salto da Huá, Brazil-Venezuelan Boundary, E. G. Holt, Nov. 25, 1930. 83617, San Antonio, Upper Orinoco River, Venezuela, E. G. Holt, Mar. 8, 1931.

Leptophis majalis Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 107, 1853.

=*Ophiodrys aestivus* (Linnaeus)

COTYPES (2): 1436, 1436a, Indianola (Calhoun County), Texas, J. D. Graham.

Leptophis mexicanus yucatanensis Oliver

Occ. Pap. Mus. Zool. Univ. Michigan, No. 462, p. 10, 1942.

PARATYPES (3): 24884-5, Yucatán, Mexico, A. Schott. 46567, Chichén-Itzá, Yucatán, Mexico, E. W. Nelson and E. A. Goldman, February 1901.

Leptophis nebulosus Oliver

Occ. Pap. Mus. Zool. Univ. Michigan, No. 462, p. 12, fig. 4, 1942.

PARATYPES (3): 19568, Greytown, Río San Juan, Nicaragua, C. W. Richmond, March 1892. 20270, Patuca, Honduras, H. W. Perry, July 24, 1891. 20683, Nicaragua, F. P. Davis.

Leptophis saturatus Cope

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8 (1875), p. 133, pl. 28, fig. 10, 1876.

=*Thalerophis depressirostris* (Cope)

HOLOTYPE: 32563, Sipurio, Costa Rica, W. M. Gabb.

Leptophis taeniata Hallowell

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 181, 1852.

=*Masticophis taeniatus taeniatus* (Hallowell)

HOLOTYPE: 2110, New Mexico, west of the Rio Grande, S. W. Woodhouse.

Leptophis trifrenatus Hallowell

Proc. Acad. Nat. Sci. Philadelphia, vol. 12 (1860), p. 503, 1861.

=*Ptyas mucosus* (Linnaeus)

HOLOTYPE: 7510, Island of Hongkong (off the mainland of China), J. M. Brooke, April 1854.

Leptotyphlops ater Taylor

Univ. Kansas Sci. Bull., vol. 26 (1939), p. 536, fig. 4, 1940.
 =*Leptotyphlops albifrons ater* Taylor
 HOLOTYPE: 79957, Managua, Nicaragua, H. C. Kellers.

Leptotyphlops bakewelli Oliver

Occ. Pap. Mus. Zool. Univ. Michigan, No. 360, p. 16, fig. 1a, 1937.
 =*Leptotyphlops phenops bakewelli* Oliver
 PARATYPE: 46340, La Salada, Michoacán, Mexico, E. W. Nelson and
 E. A. Goldman, Mar. 23, 1903.

Leptotyphlops humilis cahuilae Klauber

Trans. San Diego Soc. Nat. Hist., vol. 6, p. 339, 1931.
 PARATYPES (3): 15943, Fort Yuma (Imperial County, California), U.S.
 Fish Commission, 1889. 26289, Fort Yuma (Imperial County, California),
 H. Brown. 37114, Fort Mohave (Mohave County), Arizona, C. W. Adams,
 June 11, 1904.

Leptotyphlops humilis segregus Klauber

Trans. San Diego Soc. Nat. Hist., vol. 9, p. 67, 1939.
 HOLOTYPE: 103670, Chalk Draw, Brewster County, Texas, 3,000 ft., T. F.
 Smith, Aug. 11, 1936.

Leptotyphlops humilis slevini Klauber

Trans. San Diego Soc. Nat. Hist., vol. 6, p. 338, 1931.
 PARATYPES (4): 5292a-b, Cape San Lucas, Baja California, Mexico, J.
 Xantus. 12601, La Paz, Baja California, Mexico, L. Belding, February
 1882. 64580, Cape San Lucas, Baja California, Mexico, C. H. Townsend,
 Mar. 23, 1911.

Leptotyphlops magnamaculata Taylor

Univ. Kansas Sci. Bull., vol. 26 (1939), p. 532, fig. 1, 1940.
 =*Leptotyphlops albifrons magnamaculata* Taylor
 HOLOTYPE: 54760, Utila Island, Honduras, F. J. Dyer, Apr. 9, 1916. The
 catalog states: "Taken while crossing path in banana plantation in dry
 ground."

Leptotyphlops nasalis Taylor

Univ. Kansas Sci. Bull., vol. 26 (1939), p. 535, fig. 3, 1940.
 =*Leptotyphlops albifrons ater* Taylor
 HOLOTYPE: 16134, Managua, Nicaragua, Government of Nicaragua.

Leptotyphlops rufidorsum Taylor

Univ. Kansas Sci. Bull., vol. 26 (1939), p. 533, fig. 2, 1940.
 HOLOTYPE: 49993, Rímac Valley, Lima, Peru, 5,300 ft., C. H. Townsend,
 Apr. 16, 1913.

Leptotyphlops striatula Smith and Laufe

Proc. Biol. Soc. Washington, vol. 58, p. 29, figs. A-B, pl. 5, 1945.

HOLOTYPE: 98889, Yamachi, South Yungas, Bolivia, 3,000 m., M. Cárdenas, 1935.

Liasis amethystinus kinghorni Stull

Occ. Pap. Mus. Zool. Univ. Michigan, No. 267, p. 3, 1933.

PARATYPE: 11035, Australia, AM.

Lichanura orcutti Stejneger

West Amer. Sci., vol. 6, p. 83, 1889.

=*Lichanura roseofusca* (Cope)

HOLOTYPE: 15503, Colorado Desert, San Diego County, California, C. R. Orcutt, April 1889.

Lichanura roseofusca gracia Klauber

Trans. San Diego Soc. Nat. Hist., vol. 6, p. 307, pl. 21, fig. 1, 1931.

PARATYPES (5): 20643, 20953, Harquahala Mountains, Arizona, H. Brown, 1893. 29698, Phoenix (Maricopa County), Arizona, O. L. Geer, September 1901. 44317, Providence Mountains, California, F. Stephens, June 3, 1902. 60238, Harquahala Mountains, Arizona, 3,000 ft., E. A. Goldman, Oct. 14, 1917. While these specimens were not listed by number in the original description, Mr. Klauber considers them paratypes.

Lichanura simplex Stejneger

Proc. U.S. Nat. Mus., vol. 12, p. 97, fig. 2, 1890.

=*Lichanura roseofusca* Cope

HOLOTYPE: 13810, San Diego, San Diego County, California, R. Smith, Mar. 5, 1884.

Lichanura trivirgata Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 13, p. 304, 1861.

HOLOTYPE: 15502 (formerly 5023), Cape San Lucas, Baja California, Mexico, J. Xantus, September 1859.

Liodytes alleni lineapiatus Auffenberg

Herpetologica, vol. 6, p. 13, figs. 1-3, 1950.

PARATYPE: 129439, Ingraham Highway, 5 miles east of Paradise Key, Dade County, Florida, RPSP.

Lioninia vermiformis Hallowell

Proc. Acad. Nat. Sci. Philadelphia, vol. 12 (1860), p. 484, 1861.

=*Tantilla vermiformis* (Hallowell)

COTYPES (4): 32338-41 (formerly 5792), Nicaragua, C. Wright.

Liopeltis libertatis Barbour

Proc. Biol. Soc. Washington, vol. 23, p. 169, 1910.

=*Ptyas korros korros* (Schlegel)

HOLOTYPE: 42932, Buitenzorg, Java, O. Bryant, July 4, 1909.

Liophis perfuscus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 77, 1862.

=*Leimadophis perfuscus* (Cope)

COTYPES (2): 6044 (2), Barbados, West Indies, T. N. Gill.

Liophis subfasciatus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 77, 1862.

=*Leimadophis poecilogyrus* (Wied)

HOLOTYPE: 5809, Paraguay, T. J. Page.

Lycodon osmanhilli Taylor

Univ. Kansas Sci. Bull., vol. 33, p. 562, pl. 20, 1950.

PARATYPE: 19215, Ceylon, BMNH.

Lycodon subcinctus sealei Leviton

Philippine Journ. Sci., vol. 84, p. 195, 1955.

PARATYPES (2): 51636, Batu Lepo, Sungai, Borneo, H. C. Raven, Nov. 18, 1913. 130246, Bundu Tuhan, Mount Kinabalu, Borneo, D. H. Johnson, July 3, 1951.

Lycognathus lyrophanes Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 12, p. 343, 1860.

=*Trimorphodon lyrophanes* (Cope)

COTYPES (2): 4680 (2), Cape San Lucas, Baja California, Mexico, J. Xantus, May 1859. One cotype was exchanged to ANSP.

Lygophis flavifrenatus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 80, 1862.

=*Aporophis flavifrenatus* (Cope)

COTYPE: 5397, Buenos Aires, Argentine, T. J. Page.

Lytorhynchus sinai Schmidt and Marx

Chicago Nat. Hist. Mus., Fieldiana: Zool., vol. 39, p. 30, fig. 4, 1956.

HOLOTYPE: 134989, ♀, Wadi Feiran, Sinai Peninsula, R. E. Kuntz.

Masticophis flagellum lineatulus Smith

Journ. Washington Acad. Sci., vol. 31, p. 394, 1941.

HOLOTYPE: 105292, ♀, 11 miles north (original description states "south") of San Buena Ventura, Chihuahua, Mexico, H. M. Smith, Oct. 22, 1938.

PARATYPES (13): 1988, ♂, Alamo de Parras, Coahuila, Mexico, D. N. Couch. 1989, Cobre Mines (Grant County), New Mexico, H. B. Möllhausen. 4388, ♂, Laguna, Valencia County, New Mexico, H. B. Möllhausen. 8429, ♀, New Mexico, O. Loew, 1874. 12676, ♀, Guanajuato Mexico, A. Dugès, 1881. 14279, 14283, ♀, ♀, Chihuahua, Mexico, E. Wilkinson, Oct. 10, 1885. 22142, ♂, Las Cruces, Dona Ana County, New Mexico, T. D. A. Cockerell. 26151, Mexico, A. Dugès, 1884. 46355, ♂, Guanaceví, Durango, Mexico, E. W. Nelson and E. A. Goldman, Aug. 16, 1898. 102240, ♂, Jornada Experimental Range, Dona Ana County, New Mexico, E. T. Ares and E. L. Little, Jr., July 19, 1934. 104675-6, ♂, ♂, Río Santa María, near Progreso, Chihuahua, Mexico, H. M. Smith, Oct. 13, 1938.

Masticophis flagellum variolosus Smith

Proc. U.S. Nat. Mus., vol. 93, p. 448, 1943.

HOLOTYPE: 24681, ♂, María Madre Island, Tres Mariás Islands, Nayarit, Mexico, E. W. Nelson, May 12, 1897.

PARATYPES (2): 24680, 24682, ♀, ♀, María Magdalena Island, Tres Mariás Islands, Nayarit, Mexico, E. W. Nelson, May 12-16, 1897.

Masticophis melanolomus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 20, p. 134, 1868.

=*Dryadophis melanolomus melanolomus* (Cope)

HOLOTYPE: 24985, Yucatán, Mexico, A. Schott.

Masticophis ornatus Baird and Girard

Serpents, pt. 1, in Catalogue of North American reptiles in the Museum of the Smithsonian Institution, p. 102, 1853.

=*Masticophis taeniatus ornatus* Baird and Girard

COTYPE: 1971, skin, between San Antonio and El Paso, Texas (restricted to Fort Davis, Jeff Davis County), J. D. Graham. One cotype was exchanged to ANSP.

Masticophis ruthveni Ortenburger

Occ. Pap. Mus. Zool. Univ. Michigan, No. 139, p. 3, pls. 1-3, 1923.

PARATYPES (2): 37546, ♂, Soto la Marina, Tamaulipas, Mexico, E. W. Nelson and E. A. Goldman, Mar. 2, 1902. 64681, Tamaulipas, Mexico, 50 miles south of Brownsville, Texas, F. M. Woodruff.

Masticophis schotti Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 160, 1853.

=*Masticophis taeniatus schotti* Baird and Girard

HOLOTYPE: 1972, Eagle Pass, Maverick County, Texas, A. Schott.

Masticophis taeniatus australis Smith

Journ. Washington Acad. Sci., vol. 31, p. 390, 1941.

HOLOTYPE: 10240, ♀, Guanajuato, Mexico, A. Dugès, 1879.

PARATYPE: 111312, Tacícuaro, Michoacán, Mexico, H. M. Smith, 1939.

Megalops maculatus Hallowell

Proc. Acad. Nat. Sci. Philadelphia, vol. 12 (1860), p. 488, 1861.

= *Leptodeira maculata* (Hallowell)

HOLOTYPE: 7367, Tahiti (but probably from Nicaragua), C. B. Adams.

Mesopeltis longifrenis Stejneger

Proc. U.S. Nat. Mus., vol. 36, p. 457, 1909.

= *Sibynomorphus longifrenis* (Stejneger)

HOLOTYPE: 38750, Bocas del Toro, Panama, F. E. Read.

Mesopeltis sanniolus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 18 (1866), p. 318, 1867.

= *Dipsas sanniolus* (Cope)

HOLOTYPE: 6564, Yucatán, Mexico, A. Schott.

PROBABLE PARATYPE: 24888, same data.

Micrurus affinis mayensis Schmidt

Field Mus. Nat. Hist., Zool. Ser., vol. 20, p. 37, 1933.

PARATYPE: 46562, Chichén-Itzá, Yucatán, Mexico, E. W. Nelson and E. A. Goldman, February 1901.

Micrurus affinis stantoni Schmidt

Field Mus. Nat. Hist., Zool. Ser., vol. 20, p. 36, 1933.

PARATYPES (3): 16398, Mexico, MNHP. 56611, Belize, British Honduras, J. Hurter, May 20, 1914. 71784, Petén, Guatemala, H. Malleis.

Micrurus clarki Schmidt

Field Mus. Nat. Hist., Zool. Ser., vol. 20, p. 211, 1936.

PARATYPE: 14062, Costa Rica, J. C. Zeledon, Oct. 4, 1884.

Micrurus diastema proximans Smith and Chapliw

Herpetologica, vol. 13, p. 270, 1957.

PARATYPE: 67374, Magdalena, Jalisco, Mexico, C. C. Elliott.

Micrurus latifasciatus Schmidt

Field Mus. Nat. Hist., Zool. Ser., vol. 20, p. 35, 1933.

= *Micrurus fulvius* (Linnaeus)

PARATYPE: 20637, Guatemala, W. C. Shannon and C. A. Hedekin.

Micrurus nigrocinctus alleni Schmidt

Field Mus. Nat. Hist., Zool. Ser., vol. 20, p. 209, 1936.

PARATYPES (2): 15306, Cape Gracias, Nicaragua, C. Carlos. 20688, Nicaragua, F. P. Davis.

Micrurus nigrocinctus mosquitensis Schmidt

Field Mus. Nat. Hist., Zool. Ser., vol. 20, p. 33, 1933.

PARATYPES (7): 9785, Costa Rica, J. C. Zeledon, 1878. 19553, Greytown, Nicaragua, C. W. Richmond, Feb. 6, 1892. 30624-5, 30627-8, Costa Rica,

W. M. Gabb. 37345, Topaz Mine, 90 miles northwest of Bluefields, Nicaragua, W. F. Thornton.

Micrurus nigrocinctus ovandoensis Schmidt and Smith

Field Mus. Nat. Hist., Zool. Ser., vol. 29, p. 26, 1943.

HOLOTYPE: 111331, ♂, Salto de Agua, Mount Ovando, 1,200 ft., Chiapas, Mexico, H. M. Smith, May 19, 1940.

PARATYPE: 111332, La Esperanza, Chiapas, Mexico, 400 ft., H. M. Smith, May 25, 1940.

Micrurus olsoni Schmidt and Schmidt

Field Mus. Nat. Hist., Zool. Ser., vol. 12, p. 130, pl. 11, 1925.

PARATYPE: 38558, Peru, R. E. Coker.

Natrix compressicauda bivittata Cope

Proc. U.S. Nat. Mus., vol. 11, p. 392, pl. 56, figs. 4a-d, 1889.

=*Natrix sipedon compressicauda* (Kennicott)

COTYPES (2): 13659, 13692, Georgiana, Brevard County, Florida, W. Wittfield, 1883.

Natrix cyclopion floridana Goff

Occ. Pap. Mus. Zool. Univ. Michigan, No. 327, p. 1, 1936.

PARATYPES (6): 72590, Gainesville, Alachua County, Florida, O. C. Van Hyning, 1927. 83440, 84272, Miami, Dade County, Florida, M. K. Brady, 1930. 85268, Miami, Dade County, Florida, R. F. Deckert, Jan. 17, 1932. 85269, Paradise Key, Dade County, Florida, M. K. Brady, Jan. 27, 1932. 120836, Sebastian, Indian River County, Florida, G. Nelson, April 1935.

Natrix erythrogaster flavigaster Conant

Copeia, No. 1, p. 2, pl. 1, figs. 3a-c, 1949.

PARATYPE: 125718, Westwego, Jefferson Parish, Louisiana, E. B. Chamberlain, Oct. 24, 1943.

Natrix erythrogaster neglecta Conant

Copeia, No. 1, p. 5, pl. 1, figs. 1a-c, 1949.

PARATYPES (3): 30851-3, Olivet, Eaton County, Michigan, H. L. Clark, June 1902.

Natrix fasciata pictiventris Cope

Amer. Nat., vol. 29, p. 677, 1895.

=*Natrix sipedon pictiventris* Cope

COTYPES (11): 5473 (2), Palatka, Putnam County, Florida, T. Glover. 10449 (2), 11444, Gainesville, Alachua County, Florida, J. Bell. 10739, Clearwater, Pinellas County, Florida, S. T. Walker, July 14, 1879. 13641, 13698 (given erroneously as 13664 and 13693 in original description),

Georgiana, Brevard County, Florida, W. Wittfield. 13779, Punta Rassa (Lee County), Florida, C. W. Ward, Dec. 8, 1883. 19798, western Florida, J. A. Henshall, 1889. 19999, Lake Eustis (Lake County), Florida, T. Holm, Mar. 11, 1893.

***Natrix fasciata pleuralis* Cope**

Report of the U.S. National Museum . . . for the year ending June 30, 1898, p. 973, 1900.

=*Natrix sipedon pleuralis* Cope

HOLOTYPE: 1080, Summerville, Dorchester County, South Carolina, C. Girard.

PARATYPE: 8786, Augusta, Richmond County, Georgia, W. Phillips.

***Natrix harteri* Trapido**

Amer. Midl. Nat., vol. 25, p. 673, 1941.

HOLOTYPE: 110927, ♂, along Brazos River, north of Palo Pinto, Palo Pinto County, Texas, P. Harter, April 1940.

***Natrix olivacea uluguruensis* Loveridge**

Bull. Mus. Comp. Zool., vol. 79, p. 7, 1935.

PARATYPES (2): 120838-9 (formerly MCZ 23096-7), Nyange, Uluguru Mountains, Tanganyika, A. Loveridge, October 1926.

***Natrix rhombifera blanchardi* Clay**

Ann. Carnegie Mus., vol. 27, p. 252, pl. 25, figs. 1-2, 1938.

PARATYPE: 46533, Tlacotalpan, Veracruz, Mexico, E. W. Nelson and E. A. Goldman, May 18, 1894. This specimen is also a paratype of *Natrix rhombifera werleri* Conant, Chicago Acad. Sci. Nat. Hist. Misc., No. 126, p. 4, 1953.

***Natrix rhombifera werleri* Conant**

Chicago Acad. Sci. Nat. Hist. Misc., No. 126, p. 4, 1953.

=*Natrix rhombifera blanchardi* Clay

PARATYPES (2): 46533, Tlacotalpan, Veracruz, Mexico, E. W. Nelson and E. A. Goldman, May 18, 1894. 110511, Emiliano Zapata, Tabasco, Mexico, H. M. Smith, Aug. 9, 1939. 46533 is also a paratype of *Natrix rhombifera blanchardi* Clay, Ann. Carnegie Mus., vol. 27, p. 252, pl. 25, figs. 1-2, 1938.

***Natrix rigida deltae* Huheey**

Copeia, No. 4, p. 305, fig. 1b, 1959.

PARATYPE: 71180, Morgan City, Saint Mary Parish, Louisiana, A. and R. D. Svihla, June 1925.

***Natrix valida isabelleae* Conant**

Chicago Acad. Sci. Nat. Hist. Misc., No. 126, p. 7, 1953.

PARATYPE: 31384, Colima, Mexico, J. Xantus.

Nerodia agassizii Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 41, 1853.

=*Natrix sipedon sipedon* Linnaeus

HOLOTYPE: 1350, Lake Huron (Michigan), L. Agassiz.

Nerodia compressicauda Kennicott

Proc. Acad. Nat. Sci. Philadelphia, vol. 12, p. 335, 1860.

=*Natrix sipedon compressicauda* (Kennicott)

HOLOTYPE: 1348, south of Tampa Bay, Florida, D. Welsh.

Nerodia couchii Kennicott

Proc. Acad. Nat. Sci. Philadelphia, vol. 12, p. 335, 1860.

=*Natrix erythrogaster transversa* (Hallowell)COTYPES (2): 1314, Santa Catarina, Nuevo León, Mexico, D. N. Couch.
1319, skin, San Diego, Nuevo León, Mexico, D. N. Couch.**Nerodia holbrookii** Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 43, 1853.

=*Natrix rhombifera rhombifera* (Hallowell)

COTYPE: 1117, Prairie Mer Rouge, Morehouse Parish, Louisiana, J. Fairie. One cotype was exchanged to ANSP.

Nerodia woodhousii Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 42, 1853.

=*Natrix erythrogaster transversa* (Hallowell)COTYPES (6): 1308, Indianola (Calhoun County), Texas, J. D. Graham.
1327-8 (4), New Braunfels (Comal County), Texas, F. Lindheimer. 131742
(formerly 1321), Indianola to San Antonio (restricted to Indianola, Calhoun County), Texas, J. D. Graham.**Ninia diademata** Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 49, 1853.

=*Ninia diademata diademata* Baird and Girard

HOLOTYPE: 12122, Orizaba, Veracruz, Mexico, F. Sumichrast.

Ninia diademata nietoi Burger and Werler

Univ. Kansas Sci. Bull., vol. 36, p. 657, fig. 1, 1954.

PARATYPES (2): 35858, Finca Acleta, Alta Verapaz, Guatemala, W. R. Maxon and R. Hay, January 1905. 109807, Piedras Negras, Petén, Guatemala, H. M. Smith, June 20, 1939.

Ninia sebae immaculata Schmidt and Rand

Chicago Nat. Hist. Mus., Fieldiana: Zool., vol. 39, p. 81, 1957.

PARATYPES (15): 15207-8, Nicaragua, L. F. H. Birt. 15627-9, 15639, San Juan del Norte, Nicaragua, J. F. Le Baron. 19564, 19571, Greytown, Nicaragua, C. W. Richmond, Feb. 6, 9, 1892. 19890, Escondido River,

50 miles from Bluefields, Nicaragua, C. W. Richmond, Oct. 18, 1892. 25244, Nicaragua, E. Flint. 25319, Greytown, Nicaragua, E. P. Alexander, July 1898. 79971-2, Managua, Nicaragua, H. C. Kellers. 118041-2, San Carlos, Nicaragua, L. Marden, 1943.

***Ninia sebae tessellatus* Cope**

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8 (1875), p. 145, 1876.
 =*Ninia tessellata* Cope
 COTYPES (2): 32568-9, Costa Rica (Sipurio), W. M. Gabb.

***Nothopsis rugosus* Cope**

Proc. Acad. Nat. Sci. Philadelphia, vol. 23, p. 201, 1871.
 HOLOTYPE: 12427, Isthmus of Darién (Panama-Colombia), T. O. Selfridge.

***O[gmius] acutus* Cope**

in Ferrari-Perez, Proc. U.S. Nat. Mus., vol. 9, p. 189, 1886.
 =*Toluca lineata acuta* (Cope)
 HOLOTYPE: 30552, Tuchtán on the Pacific side of the Isthmus of Tehuantepec (but probably from Puebla), Mexico, F. Sumichrast.

***Oligodon annamensis* Leviton**

Journ. Washington Acad. Sci., vol. 43, p. 422, fig. 1, 1953.
 HOLOTYPE: 90408, ♀, Station Agricole, Blao, Haut Donai, Annam (Vietnam), E. Poilane, Mar. 11, 1933.

***Oligodon sub-griseum* Duméril and Bibron**

Erpétologie générale ou histoire naturelle complète des reptiles, vol. 7, pt. 1, p. 59, 1854.
 =*Oligodon subgriseum* Duméril and Bibron
 COTYPE: 6200, Pondicherry, India, L. T. Leschenault.

***Ophedrys vernalis blanchardi* Grobman**

Misc. Publ. Mus. Zool. Univ. Michigan, No. 50, p. 11, 1941.
 PARATYPES (6): 1476, ♀, road to Bridger's Pass (Carlson County), Wyoming, W. S. Wood, 8425, ♀, Abiquiu (Rio Arriba County), New Mexico, O. Loew, August 1874. 9095, ♂, Merino Valley, New Mexico, W. L. Carpenter, July 1875. 22377, ♀, Mesilla Valley, New Mexico, H. B. Lane, 25433, ♀, Ruidoso at Forks (Lincoln County), New Mexico, C. M. Barber, Aug. 18, 1898. 40199, ♀, Vallecito (La Plata County), Colorado, M. Cary, June 6, 1907.

***Ophiomorphus doliatus caesius* Cope**

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 348, 1862.
 =*Liophis poecilogyrus* (Wied)
 HOLOTYPE: 5393, Santa Fe, Paraguay, T. J. Page.

Ophibolus boylii Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 82, 1853.

=*Lampropeltis getulus californiae* (Blainville)

HOLOTYPE: 1698, El Dorado County, California, C. C. Boyle.

Ophibolus clericus Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 88, 1853.

=*Lampropeltis triangulum triangulum* (Lacépède)

COTYPE: 2380, Clarke County, Virginia, C. B. R. Kennerly.

Ophibolus doliatus collaris Cope

Proc. U.S. Nat. Mus., vol. 11, p. 383, 1889.

=*Lampropeltis triangulum triangulum* (Lacépède)

COTYPES (5): 2192, Roanoke County, Virginia, C. Shoemaker. 2433, Elmira, Stark County, Illinois, E. R. Boardman. 5449, Washington, D.C. 7849 (2) (formerly 2344), St. Louis, St. Louis County, Missouri, G. Engelmann. 7849 is listed under *Lampropeltis triangulum sypila* (Cope) by Blanchard, U.S. Nat. Mus. Bull. 114, p. 186, 1921.

Ophibolus doliatus parallelus Cope

Proc. U.S. Nat. Mus., vol. 11, p. 385, 1889.

=*Lampropeltis triangulum amaura* (Cope)

HOLOTYPE: 10544, Gainesville, Alachua County, Florida, J. Bell, March 1881.

Ophibolus doliatus sypilus Cope

Proc. U.S. Nat. Mus., vol. 11, p. 381, 1889.

=*Lampropeltis triangulum sypila* (Cope)

HOLOTYPE: 13380, Richland County, Illinois, R. Ridgway, 1883.

PARATYPES (10): 303, ♂, Independence, Jackson County, Missouri, J. G. Cooper, June 6, 1857. 1846 (2), Fort Towson, Choctaw County, Oklahoma, L. A. Edwards. 2296, Prairie Mer Rouge, Morehouse Parish, Louisiana, J. Fairie. 4291, Republican River, 95 miles west of Riley (in Clay County), Kansas, J. F. Hammond. 5188, Grand Coteau, Saint Landry Parish, Louisiana, J. W. Wallace. 12925, 13008, 13045, New Orleans, Orleans Parish, Louisiana, R. W. Shufeldt, 1883. 13361, ♂, Wheatland, Knox County, Indiana, R. Ridgway, 1883. 2296 is also a paratype of *Ophibolus gentilis* Baird and Girard.

Ophibolus evansii Kennicott

Proc. Acad. Nat. Sci. Philadelphia, vol. 11, p. 99, 1859.

=*Lampropeltis calligaster calligaster* Kennicott

COTYPES (4): 1593 (3), southern Illinois, R. Kennicott, 1855. 1702, Canadian River, Indian Territory (Oklahoma), H. B. Möllhausen. One cotype from 1593 was exchanged to TCM.

Ophibolus gentilis Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 90, 1853.

=*Lampropeltis triangulum gentilis* (Baird and Girard)

COTYPES (3): 1853, north fork of Red River near Sweetwater Creek, Wheeler County, Texas, R. B. Marcy, June 14, 1852. 2296, Prairie Mer Rouge, Morehouse Parish, Louisiana, J. Fairie. 131737 (formerly 1852), Red River, Oklahoma, G. B. McClellan. 2296 is also a paratype of *Ophibolus doliatus sypsilus* Cope.

Ophibolus getulus eiseni Yarrow

Proc. U.S. Nat. Mus., vol. 5, p. 439, 1882.

=*Lampropeltis getulus californiae* (Blainville)

HOLOTYPE: 11788, Fresno, Fresno County, California, G. Eisen, 1879.

PARATYPES (10): 11744, 11787 (8), 55100, same data.

Ophibolus getulus multicinctus Yarrow

Proc. U.S. Nat. Mus., vol. 5, p. 440, 1882.

=*Lampropeltis zonata multicincta* (Yarrow)

HOLOTYPE: 11753, Fresno, Fresno County, California, G. Eisen.

Ophibolus getulus niger Yarrow

Proc. U.S. Nat. Mus., vol. 5, p. 438, 1882.

=*Lampropeltis getulus niger* (Yarrow)

COTYPES (2): 12149 (2), Wheatland, Knox County, Indiana, R. Ridgway, 1881.

Ophibolus pyromelanus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 18 (1866), p. 305, 1867.

=*Lampropeltis pyromelana pyromelana* (Cope)

COTYPE: 7845, Fort Whipple (Yavapai County), Arizona, E. Palmer.

Ophibolus splendidus Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 83, 1853.

=*Lampropeltis getulus splendida* (Baird and Girard)

HOLOTYPE: 1726, skin, Sonora, Mexico, J. D. Graham.

PARATYPE: 4294, same data.

Opisthotropis multicincta Fan

Bull. Dept. Biol. Coll. Sci. Sun Yatsen Univ., No. 11, p. 82, figs. 5-6, 1931.

=*Opisthotropis balteata* (Cope)

PARATYPE: 85000, Loshiang, Kwangsi, China, SYUM through T. H. Fan.

Oreomorphus merremii semiaureus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 348, 1862.

=*Rhadinaea semiaurea* (Cope)

HOLOTYPE: 4665, Paraguay, T. J. Page.

Oxyrhopus fitzingeri frizzelli Schmidt and Walker

Field Mus. Nat. Hist., Zool. Ser., vol. 24, p. 314, 1943.

PARATYPE: 48552, ♂, Piura, Peru, C. H. Townsend, May 1, 1911.

Oxyrhopus trigeminus Duméril and Bibron

Erpétologie générale ou histoire naturelle complète des reptiles, vol. 7, pt. 2, p. 1013, 1854.

=*Clelia trigemina* (Duméril and Bibron)

COTYPE: 11137 (formerly 406), Bahia, Brazil, JPP.

Oxyrhopus doliatus semicinctus Cope

U.S. Nat. Mus. Bull. 32, p. 76, 1887.

=*Pseudoboa petola* (Linnaeus)

HOLOTYPE: 28900, Sipurio, Costa Rica, W. M. Gabb.

Pelophilus madagascariensis Duméril and Bibron

Erpétologie générale ou histoire naturelle complète des reptiles, vol. 6, p. 524, 1844.

=*Acrantophis madagascariensis madagascariensis* (Duméril and Bibron)

COTYPE: 10978, Madagascar, JPP.

Philodryas latirostris Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, pp. 73, 348, 1862.

=*Chlorosoma olfersii* (Lichtenstein)

COTYPE: 5811, Paraguay, T. J. Page.

Philothamnus irregularis battersbyi Loveridge

Bull. Mus. Comp. Zool., vol. 106, p. 190, 1951.

PARATYPES (31): 40766-7, 40953-5, 40984-5, 40988, 41512-3, 41978, Nairobi, Kenya, E. A. Mearns, Nov. 10-Dec. 4, 1909. 40956-8, 42048, 42063-4, 42256, Kenya, Smithsonian African Expedition. 40999 Juja Farm, Kenya, 5,000 ft., J. A. Loring, May 13, 1909. 41135, Fort Hall, Kenya, E. A. Mearns, Oct. 27, 1909. 41693, Juja Farm, Kenya, W. H. McMillan, 1908-09. 41701, southwest corner of Lake Naivasha, Kenya, J. A. Loring, July 12, 1909. 41970, Wambugu, Kenya, E. A. Mearns, July 24, 1909. 42935-6, Mount Kenya to Fort Hall, Kenya, Smithsonian African Expedition, Oct. 10, 1909. 48588, Mtoto Andei Station, Kenya, E. Heller. 49003, Kaimosi, Kenya, E. Heller, Jan. 27, 1912. 49007, 49012, 49021, Kenya, E. Heller. 120814, ♂, Amani, Usambare Mountains, Tanganyika, 3,000 ft., A. Loveridge, November 1926. 41136 was exchanged to MCZ. 41135 was shot in a tree; its stomach contained a clawed frog, *Xenopus laevis*?, now USNM 63066.

Phimothyra hexalepis Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 18 (1866), p. 304, 1867.

=*Salvadora hexalepis hexalepis* (Cope)

HOLOTYPE: 7894, Fort Whipple (Yavapai County), Arizona, E. Palmer.
The stomach contained a *Cnemidophorus sexlineatus*.

Phrynonax lyoni Stejneger

Proc. U.S. Nat. Mus., vol. 24, p. 185, 1901.

=*Pseustes poecilonotus polylepis* (Peters)

HOLOTYPE: 27826, Macuto, Venezuela, M. W. Lyon and W. Robinson,
Aug. 4, 1900.

Phyllorhynchus browni Stejneger

Proc. U.S. Nat. Mus., vol. 13, p. 152, 3 figs., 1890.

=*Phyllorhynchus browni browni* Stejneger

HOLOTYPE: 15719, Tucson (Pima County), Arizona, H. Brown.

Phyllorhynchus decurtatus perkinsi Klauber

Zool. Soc. San Diego Bull. 12, p. 11, fig. 2, 1935.

PARATYPES (2): 94953 (formerly SDSNH 20875), 7 miles east of The
Narrows, San Diego County, California, L. M. Klauber, July 23, 1933.
94954 (formerly SDSNH 22295), Dry Lake, San Diego County, California,
L. M. Klauber, May 5, 1934.

Phyllosira flavescens Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 14, p. 349, 1862.

=*Chironius fuscus* (Linnaeus)

HOLOTYPE: 5813, Paraguay, T. J. Page.

Pituophis annectens Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 72, 1853.

=*Pituophis catenifer annectens* Baird and Girard

COTYPES (3): 1839 (3, of which 2 are heads), San Diego County, Cali-
fornia, J. L. LeConte.

Pituophis catenifer bimaris Klauber

Trans. San Diego Soc. Nat. Hist., vol. 11, p. 7, 1946.

PARATYPES (2): 15157, Ballenas Bay, Baja California, Mexico U.S. Fish
Commission *Albatross*, May 3, 1888. 37536, El Alamo, Baja California,
Mexico, E. W. Nelson and E. A. Goldman, June 11, 1905. 37537, from
Margarita Island, Baja California, Mexico, E. W. Nelson and E. A. Gold-
man, Dec. 3, 1905, is a probable paratype.

Pituophis catenifer deserticola Stejneger

North Amer. Fauna, No. 7, p. 206, 1893.

LECTOTYPE: 18070, Beavercreek Mountains, Utah, C. H. Merriam, May 1,
1891.

PARATYPES (5): 18065, Jackass Spring, Panamint Mountains, Inyo
County, California, May 7, 1891. 18066, Surprise Canon, Panamint Moun-
tains, Inyo County, California, A. K. Fisher, Apr. 23, 1891. 18067, Shep-

herd Canon, Argus Range, Inyo County, California, A. K. Fisher, Apr. 26, 1891. 18068, Howes Meadows, 10 miles south of Owens Lake, Inyo County, California, 3,750 ft., F. Stephens, May 12, 1891. 18069, Lone Pine, Inyo County, California, A. K. Fisher, June 4, 1891.

***Pituophis catenifer fuliginatus* Klauber**

Trans. San Diego Soc. Nat. Hist., vol. 11, p. 14, 1946.

PARATYPES (2): 8565, San Martín Island, west coast of Baja California, Mexico, T. H. Streets. 24395, San Martín Island, Baja California, Mexico, A. W. Anthony, Mar. 12, 1897.

***Pituophis catenifer insulanus* Klauber**

Trans. San Diego Soc. Nat. Hist., vol. 11, p. 11, pl. 1, fig. 1, 1946.

PARATYPES (2): 14088, Cedros Island, west coast of Baja California, Mexico, C. H. Townsend, Apr. 4, 1885. 54764, Cedros Island, Baja California, Mexico, P. Wegeforth, July 1916.

***Pituophis lödingi* Blanchard**

Michigan Acad. Sci. Pap., vol. 4 (1924), p. 532, 1925.

=*Pituophis melanoleucus lödingi* Blanchard

PARATYPE: 62340, 14 miles southwest of Mobile, Mobile County, Alabama, H. P. Löding, June 1919.

***Pituophis mccllellani* Baird and Girard**

Smithsonian Misc. Coll., vol. 2, art. 5, p. 68, 1853.

=*Pituophis melanoleucus sayi* (Schlegel)

HOLOTYPE: 1540, Red River, Deaf Smith County, Texas, R. B. Marcy and G. B. McClellan, Jan. 28, 1852.

PARATYPE: 1541, same data.

***Pituophis melanoleucus ruthveni* Stull**

Occ. Pap. Mus. Zool. Univ. Michigan, No. 205, p. 1, 1929.

HOLOTYPE: 76278, ♂, Longleaf, Rapides Parish, Louisiana, W. D. Harris, Mar. 24, 1927.

***Pituophis mexicanus* Duméril and Bibron**

Erpétologie générale ou histoire naturelle complète des reptiles, vol. 7, pt. 1, p. 236, pl. 62, 1854.

=*Pituophis melanoleucus sayi* (Schlegel)

COTYPE: 1415, Mexico, JPP.

***Pituophis wilkesii* Baird and Girard**

Smithsonian Misc. Coll., vol. 2, art. 5, p. 71, 1953.

=*Pituophis melanoleucus catenifer* (Blainville)

COTYPE: 5471, Puget Sound (Washington), U.S. Exploring Expedition, 1841.

Pityophis haematois Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 12, p. 342, 1860.

=*Pituophis melanoleucus vertebralis* (Blainville)

COTYPES (2): 4682 (2), Cape San Lucas, Baja California, Mexico, J. Xantus.

Pliocercus bicolor Smith

Proc. Biol. Soc. Washington, vol. 54, p. 123, 1941.

HOLOTYPE: 25203, ♀, Tuxpan, Veracruz, Mexico, G. C. Lincecum.

Pliocercus dimidiatus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 17, p. 190, 1865.

HOLOTYPE: 6363, Arriba, Costa Rica, C. N. Riotte.

Pliocercus elapoides laticollaris Smith

Proc. Biol. Soc. Washington, vol. 54, p. 122, 1941.

HOLOTYPE: 110767, ♀, Tenosique, Tabasco, Mexico, H. M. Smith, June 30, 1939.

Pseudoboa robinsoni Stejneger

Proc. U.S. Nat. Mus., vol. 24, p. 190, figs. 25-27, 1901.

=*Pseudoboa newwiedii* (Duméril and Bibron)

HOLOTYPE: 22532, La Guaira, Venezuela, W. Robinson, June 21, 1895.

Python curtus brongersmai Stull

Occ. Pap. Boston Soc. Nat. Hist., vol. 8, p. 297, 1938.

PARATYPE: 53427, Kuala Lumpur, Malaya, T. C. Pasqual.

Rabdosoma crassicaudatum Duméril and Bibron

Erpétologie générale, ou histoire naturelle complète des reptiles, vol. 7, pt. 1, p. 103, 1854.

=*Atractus crassicaudatus* (Duméril and Bibron)

COTYPE: 11133 (formerly 387), New Granada (Panama-Colombia), JPP.

Regina clarkii Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 48, 1853.

=*Natrix sipedon clarkii* (Baird and Girard)

HOLOTYPE: 2264, Indianola (Calhoun County), Texas, J. D. Graham.

PARATYPES (3): 1392 (2), 44171, skins, same data.

Regina grahamii Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 47, 1853.

=*Natrix grahamii* (Baird and Girard)

HOLOTYPE: 1376, Rio Salado, 4 miles from San Antonio, Bexar County, Texas, J. D. Graham.

Regina kirtlandii Kennicott

Proc. Acad. Nat. Sci. Philadelphia, vol. 8, p. 95, 1856.

=*Natrix kirtlandii* (Kennicott)

COTYPES (3): 1514, 115115-6, West Northfield, Cook County, Illinois, R. W. Kennicott. One of 1514 was exchanged to MZUM. Conant, in 1943, considered the origin of 115116 to be doubtful.

Regina valida Kennicott

Proc. Acad. Nat. Sci. Philadelphia, vol. 12, p. 334, 1860.

=*Natrix valida valida* (Kennicott)

HOLOTYPE: 1309, Durango, Mexico, D. N. Couch.

Rena dulcis Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 142, 1853.

=*Leptotyphlops dulcis dulcis* (Baird and Girard)

HOLOTYPE: 7296, between San Pedro and Comanche Springs, Texas, J. D. Graham.

Rena humilis Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 143, 1853.

=*Leptotyphlops humilis humilis* (Baird and Girard)

HOLOTYPE: 2101, Vallicita, Calaveras County, California, J. L. LeConte.

Rhabdosoma guttulatatum Cope

Proc. Amer. Philos. Soc., vol. 22, p. 385, 1885.

=*Geophis chalybeus* (Wagler)

COTYPES (3): 25024-5, Mirador, Veracruz, Mexico, C. Sartorius. 30399, Veracruz, Mexico, C. Sartorius.

Rhadinaea albiceps Amaral

Journ. Washington Acad. Sci., vol. 14, p. 200, 1924.

=*Liophis albiceps* (Amaral)

HOLOTYPE: 22446, ♂, Ecuador?, M. B. Kerr, May 1895.

Rhadinaea crassa Smith

Proc. Biol. Soc. Washington, vol. 55, p. 190, pl. 3, figs. 4-5, 1942.

PARATYPE: 110366, Barranca de los Horcones, 10 km. south of Durango, Hidalgo, Mexico, H. M. Smith, Aug. 21, 1940.

Rhadinaea forbesi Smith

Proc. Biol. Soc. Washington, vol. 55, p. 188, fig. 3, 1942.

HOLOTYPE: 110365, ♂, Tequeyutepec, Veracruz, Mexico, H. M. Smith, Mar. 23, 1940.

PARATYPES (2): 29124, Guadalajara?, Mexico (probably near Orizaba, Veracruz), J. J. Major?. 110364, Tequeyutepec, Veracruz, Mexico, H. M. Smith, Mar. 23, 1940.

Rhadinaea fulviceps Cope

Proc. Amer. Philos. Soc., vol. 23, p. 279, 1886.

=*Rhadinaea pachyura fulviceps* Cope

HOLOTYPE: 61187 (formerly probably 14118), Panama, G. W. Nelson.

Rhadinaea fulvivittis Cope

Journ. Acad. Nat. Sci. Philadelphia (2), vol. 8 (1875), p. 139, 1876.

=*Rhadinaea vittata* (Jan)

HOLOTYPE: 7075, Orizaba (alpine section), Veracruz, Mexico, F. Sumichrast.

Rhadinaea hesperia Bailey

Occ. Pap. Mus. Zool. Univ. Michigan, No. 412, p. 8, pl. 2, fig. 3, 1940.

=*Rhadinaea hesperia hesperia* Bailey (in part)

PARATYPES (4): 15429-30, Guanajuato, Mexico, A. Dugès. 20166, Barranca, near Cuernavaca, Morelos, Mexico, P. L. Jouy, Aug. 1, 1892. 46456, Plomosas, Sinaloa, Mexico, E. W. Nelson and E. A. Goldman, July 19, 1897. 15430 and 46456 are also paratypes of *Rhadinaea hesperia hesperioides* Smith, 1942.

Rhadinaea hesperia hesperioides Smith

Proc. Biol. Soc. Washington, vol. 55, p. 186, fig. 1, 1942.

HOLOTYPE: 67373, ♂, Magdalena, Jalisco, Mexico, C. C. Elliott, June 23, 1924.

PARATYPES (2): 15430, Guanajuato (probably from the vicinity of Guadalajara), Mexico, A. Dugès. 46456, Plomosas, Sinaloa, Mexico, E. W. Nelson and E. A. Goldman, July 19, 1897. 15430 and 46456 are also paratypes of *Rhadinaea hesperia* Bailey.

Rhadinaea poecilopogon Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 15, p. 100, 1863.

=*Liophis poecilopogon* (Cope)

HOLOTYPE: 31278, Paysandú, Uruguay, H. W. Kennedy.

Rhadinaea quinquelineata Cope

Proc. Amer. Philos. Soc., vol. 23, p. 277, 1886.

LECTOTYPE: 31350, Teziutlán, Puebla, Mexico, S. Bernad.

Rhadinella schistosa Smith

Copeia, No. 1, p. 7, 1941.

PARATYPE: 109914, Cautlapan, Veracruz, Mexico, H. M. Smith, Aug. 7-14, 1940.

Rhamnophis aethiopissa elgonensis Loveridge

U.S. Nat. Mus. Bull. 151, p. 24, 1929.

PARATYPE: 49005, Kaimosi, Kenya, E. Heller, Jan. 27, 1912.

Rhegnops visoninus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 18, p. 128, 1866.

=*Adelphicos quadrivirgatus visoninus* (Cope)

HOLOTYPE: 24899, British Honduras (probably Belize), D. B. Parsons.

Salvadora bogerti Smith

Smithsonian Misc. Coll., vol. 99, No. 20, p. 2, figs. 1-2, 1941.

HOLOTYPE: 30296, ♀, Tehuantepec, Oaxaca, Mexico, F. Sumichrast.

Salvadora grahamiae Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 104, 1853.

HOLOTYPE: 2081, ♂, Sonora, Mexico, J. D. Graham.

Salvadora hexalepis celeris Smith

Smithsonian Misc. Coll., vol. 99, No. 20, p. 9, 1941.

HOLOTYPE: 40043, ♀, San Blas, Sinaloa, Mexico, J. N. Rose, Mar. 28, 1910.

Salvadora hexalepis klauberi Bogert

Amer. Mus. Novit., No. 1285, p. 2, figs. 1-4, 1945.

PARATYPES (8): 5347, Cape San Lucas, Baja California, Mexico, J. Xantus. 12637, 12637a-b, 12638, La Paz, Baja California, Mexico, L. Belding, 1882. 37517, 20 miles west of Santa Rosalía, Baja California, Mexico, E. W. Nelson and E. A. Goldman, Oct. 19, 1905. 37518, Todos Santos, Baja California, Mexico, E. W. Nelson and E. A. Goldman, Dec. 26, 1906. 64581, Miraflores, Baja California, Mexico, C. H. Townsend, May 9, 1911.

Salvadora hexalepis mojavensis Bogert

Amer. Mus. Novit., No. 1285, p. 6, figs. 5-9, 1945.

PARATYPES (6): 4470, Mojave Valley, California, H. B. Möllhausen. 9101, Cottonwood Canyon (Duchesne County), Utah, J. Kohlen, 1871. 18059, Shepherd Canyon, Argus Range, California, A. K. Fisher, Apr. 26, 1891. 18060, Maturango Spring, Argus Range (Inyo County), California, A. K. Fisher, May 2, 1891. 18061, Amargosa, California, T. S. Palmer, Mar. 16, 1891. 18062, Virgin River, Nevada, C. H. Merriam, May 8, 1891.

Salvadora intermedia Hartweg

Copeia, No. 4, p. 256, 1940.

=*Salvadora intermedia intermedia* Hartweg

PARATYPE: 109228, Chilpancingo, Guerrero, Mexico, H. M. Smith, 1939.

Salvadora lineata Schmidt

Field Mus. Nat. Hist., Zool. Ser., vol. 24, p. 148, fig. 15, 1940.

PARATYPES (3): 26441, Kinney County, Texas, E. A. Mearns. 55798, Waco, McLennan County, Texas, J. Hurter. 67363, San Antonio, Bexar County, Texas, C. S. East, 1924.

Scaphiodontophis albonuchalis Taylor and Smith

Univ. Kansas Sci. Bull., vol. 29, p. 323, fig. 9, pl. 23, fig. 1, pl. 24, pl. 25, figs. 1-2, 1943.

HOLOTYPE: 110413, ♀, La Esperanza, Chiapas, Mexico, H. M. Smith, Apr. 23, 1940.

PARATYPES (3): 6580, San Juan Bautista, Tabasco, Mexico, C. H. Berendt. 12087, Mexico, C. Sartorius. 12693, Escuintla, Guatemala, H. J. Stuart, June 1882.

Scaphiodontophis carpinctus Taylor and Smith

Univ. Kansas Sci. Bull., vol. 29, p. 315, pl. 22, fig. 1, text-fig. 6, 1943.

HOLOTYPE: 110411, ♂, Piedras Negras, Petén, Guatemala, H. H. Smith, June 3, 1939.

Scaphiodontophis nothus Taylor and Smith

Univ. Kansas Sci. Bull., vol. 29, p. 320, pl. 23, fig. 2, text-fig. 8, 1943.

HOLOTYPE: 110412, ♂, Potrero Viejo, Veracruz, Mexico, D. M. Forbes, Mar. 5, 1939.

Scolecophis fumiceps Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 12, p. 371, 1860.

=*Tantilla nigriceps fumiceps* (Cope)

HOLOTYPE: 110435, "probably Cuba" (restricted to San Antonio, Bexar County, Texas). This specimen was returned to MCZ in 1942.

Scolecophis scytalinus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 18 (1866), p. 320, 1867.

=*Clelia clelia clelia* (Daudin)

HOLOTYPE: 6581, San Juan Bautista, Tabasco, Mexico, C. H. Berendt.

Scoliophis atlanticus Davis, Bigelow, and Gray

Report of a committee of the Linnaean society of New England relative to a large marine animal, supposed to be a serpent seen near Cape Ann, Massachusetts, in August 1817, p. 41, pl. 1, 1817.

=*Coluber constrictor constrictor* Linnaeus

HOLOTYPE: 115617, Loblolly Cove, Cape Ann (Essex County), Massachusetts, 1817?. This specimen was reported to be the "progeny" of the Great Sea Serpent seen off the coast of Massachusetts in 1817.

Scotophis confinis Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 76, 1853.

=*Elaphe obsoleta obsoleta* (Say)

HOLOTYPE: 1563, Anderson, Anderson County, South Carolina, C. Paine.

Scotophis lindheimeri Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 74, 1853.

=*Elaphe obsoleta lindheimeri* (Baird and Girard)

HOLOTYPE: 1733, New Braunfels (Comal County), Texas, F. Lindheimer.
 PARATYPE: 1734, skin, Indianola (Calhoun County), Texas, J. D. Graham.

Scotophis vulpinus Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 75, 1853.

=*Elaphe vulpina vulpina* (Baird and Girard)

NEOTYPE: 9969, Racine, Racine County, Wisconsin, P. R. Hoy. A paratype, 1570, from Grosse Ile, Wayne County, Michigan, collected by C. Fox, is now identified as *Elaphe vulpina gloydi* Conant.

Sibon annulatum yucatanensis Cope

U.S. Nat. Mus. Bull. 32, p. 67, 1887.

=*Leptodeira yucatanensis yucatanensis* (Cope)

COTYPE: 24887 (formerly 6553), Yucatán, Mexico, A. Schott.

Sibynomorphus macrostomus Amaral

Journ. Washington Acad. Sci., vol. 14, p. 200, 1924.

=*Dipsas macrostomus* (Amaral)

HOLOTYPE: 14047, ♀, Ecuador, W. H. Jones, Dec. 18, 1884.

Simocephalus chanleri Stejneger

Proc. U.S. Nat. Mus., vol. 16, p. 726, 1894.

=*Mehelya chanleri chanleri* (Stejneger)

HOLOTYPE: 20126, Wange, Island of Manda, north of Lamu, East Africa (Kenya), G. Denhardt.

Sonora semiannulata Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 117, 1853.

=*Sonora semiannulata semiannulata* Baird and Girard

HOLOTYPE: 2109, "Sonora, Mexico" (restricted to Santa Rita Mountains, Arizona), J. D. Graham.

Spilotes chrysobronchus Cope

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8 (1875), p. 136, pl. 28, fig. 11, 1876.

=*Pseustes poecilonotus chrysobronchus* (Cope)

HOLOTYPE: 30623, coast region of Costa Rica, W. M. Gabb.

Spilotes erebennus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 12 (1860), p. 342, 1861.

=*Drymarchon corais erebennus* (Cope)

COTYPE: 1862, skin, Eagle Pass, Maverick County, Texas, A. Schott.

Spilotes melanurus Duméril and Bibron

Erpétologie générale, ou histoire naturelle complète des reptiles, vol. 7, pt. 1, p. 224, 1854.

=*Drymarchon corais melanurus* (Duméril and Bibron)

COTYPE: 1416, Mexico, JPP.

Spilotes pullatus auribundus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 13, p. 300, 1861.

=*Spilotes pullatus mexicanus* (Laurenti)

COTYPE (3): 25004-6, Mirador, Veracruz, Mexico, C. Sartorius.

Stenorhina degenhardtii apiata Cope

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8 (1875), p. 142, 1876.

HOLOTYPE: 70405, Tehuantepec, Oaxaca, Mexico, F. Sumichrast.

PARATYPE: 30090, Guichicovi, Tehuantepec, Oaxaca, Mexico, F. Sumichrast.

Stenorhina lactea Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 13, p. 303, 1861.

=*Stenorhina freminwillii lactea* Cope

HOLOTYPE: 4944, La Unión, Guatemala (but probably from La Unión, El Salvador), J. M. Dow.

Stenostoma phenops Cope

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8 (1875), p. 128, 1876.

=*Leptotyphlops phenops phenops* (Cope)

COTYPES (9): 6760, Cobán, Guatemala, H. Hague. 12444, 30289-95, Tehuantepec, Oaxaca, Mexico, F. Sumichrast.

Stilosoma extenuatum arenicola Highton

Bull. Florida State Mus., vol. 1, p. 86, 1956.

PARATYPE: 23333, Tarpon Springs, Pinellas County, Florida, C. S. and H. H. Brimley, January 1897.

Storeria dekayi temporalineata Trapido

Amer. Midl. Nat., vol. 31, p. 70, pl. 9, figs. 51-52, 1944.

HOLOTYPE: 32148, ♀, San Rafael, Jicaltepec, Veracruz, Mexico, about 100 ft., C. H. Townsend, July 2, 1896.

Storeria dekayi texana Trapido

Amer. Midl. Nat., vol. 31, p. 63, figs. 45-50, 1944.

PARATYPES (28): 2095, Brazos River, Texas, B. F. Shumard. 7279 (2), Matamoras, Tamaulipas, Mexico, D. N. Couch. 11820, 11823, Old Fort Cobb, Caddo County, Oklahoma, E. Palmer. 13090, New Orleans, Orleans Parish, Louisiana, R. W. Shufeldt, 1883. 14646, Waco, McLennan County, Texas, given by a Waco Indian chief, Nov. 10, 1886. 17699, New Braunfels, Comal County, Texas, B. W. Evermann, Dec. 3, 1891. 32802, Barnard Creek, west of Columbia (probably Columbus, Colorado County), Texas, W. Lloyd, Mar. 4, 1892. 42286, Victoria County, Texas, J. D. Mitchell, Sept. 24, 1904. 42290, Victoria, Victoria County, Texas, J. D. Mitchell, 1905-1916. 52281, 52284, ♀, Brownsville, Cameron County, Texas, R. D.

Camp, Mar. 31, 1915. 55912, Falls County, Texas, J. Hurter, June 12, 1906. 55914, Waco, McLennan County, Texas, J. Hurter, May 24, 1908. 65894, Winsted, McLeod County, Minnesota, H. F. Nachtrieb, May 1923. 71754, Putnam, Callahan County, Texas, A. J. Kirn, Apr. 30, 1926. 78611-8, Victoria, Victoria County, Texas, J. D. Mitchell, 1906-1910. 94153-4, 4 miles north of Lake City, Barber County, Kansas, C. E. Burt, Mar. 31, 1934. 103741, juv., Harlingen, Cameron County, Texas, N. O. Barry, Nov. 1, 1937.

***Storeria dekayi wrightorum* Trapido**

Amer. Midl. Nat., vol. 31, p. 57, figs. 38-44, 1944.

PARATYPES (74): 1945 (2), Pittsburgh, Allegheny County, Pennsylvania. 1958-59 (3), Racine, Racine County, Wisconsin. 2137, Lebanon, Boone County, Indiana, S. B. Davis. 2215, Port Huron, St. Clair County, Michigan, S. F. Baird. 2222, Pensacola, Escambia County, Florida, W. A. Hammond. 2226 (2), Monticello, Lawrence County, Mississippi, H. Teunison. 4798b, New Orleans, Orleans Parish, Louisiana, NOAS. 4814, southern Illinois, R. Kennicott. 6388, Chicago, Cook County, Illinois, R. Kennicott. 8770 (7), Belleville, St. Clair County, Illinois, A. Reuss. 8983 (3), Kinston, Lenoir County, North Carolina, J. W. Milner. 10211 (2), St. James Parish, Louisiana, O. de la Peichardiére, May 1879. 10653-5, 13358-9, Wheatland, Knox County, Indiana, R. Ridgway, 1881-1883. 10665-6, Monroe, Butler County, Ohio, R. T. Shepherd, Mar. 12, 1880. 10711, Nashville, Berrien County, Georgia, W. J. Taylor, Aug. 21, 1880. 12028 (18), Mount Carmel, Wabash County, Illinois, L. M. Turner. 12904, 12922, New Orleans, Orleans Parish, Louisiana, R. W. Shufeldt, 1883. 15577, Cherokee, Swain County, North Carolina, J. Mooney. 17967, ♀, Irvington, Marion County, Indiana, O. P. Hay. 26077, Wilmington, New Hanover County, North Carolina, W. P. Seal, May 9, 1899. 29606, Huguenot, Elbert County, Georgia, J. H. DuBose. 31666, Lobdell, West Baton Rouge Parish, Louisiana, A. Allison, June 1903. 33529, Lake Maxinkuckee, Jasper-Pulaski Counties, Indiana, B. W. Evermann, Oct. 8, 1900. 42542, Mobile County, Alabama, J. Hurter, May 26, 1910. 42589, Lake Maxinkuckee, Jasper-Pulaski Counties, Indiana, Gruff and Gard, Oct. 17, 1906. 44294, Marble Cave, Stone County, Missouri, C. H. Merriam, May 22, 1892. 55922, 55924, ♂, Little Rock, Carroll County, Arkansas, J. Hurter, May 7, 1913 (for 55922), Apr. 9, 1906 (55924). 55925, Mobile County, Alabama, J. Hurter, June 11, 1910. 55926-7, Mobile, Mobile County, Alabama, J. Hurter, 1907 (for 55926), Mar. 26, 1908 (for 55927). 55928, St. Louis County, Missouri, J. Hurter, Mar. 20, 1898. 55932, Stoddard County, Missouri, J. Hurter, Oct. 26, 1899. 55935, St. Francois County, Missouri, J. Hurter, 1894. 92062, island above Price Island, in Savannah River, Lincoln County, Georgia, C. E. Burt, July 19, 1933. 102589-90, Auburn, Lee County, Alabama, G. C. Moore.

Storeria tropica Cope

Proc. Amer. Philos. Soc., vol. 22, p. 175, 1885.

= *Storeria dekayi tropica* Cope

HOLOTYPE: 6759, Petén, Guatemala, H. Hague.

Symphimus leucostomus Cope

Proc. Amer. Philos. Soc., vol. 11, p. 150, 1869.

COTYPES (2): 30310, Chihuitán, Tehuantepec, Oaxaca, Mexico, F. Sumichrast. 30311, head and tail only, Oaxaca, Mexico, F. Sumichrast.

Sympholis lippiens Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 13 (1861), p. 524, 1862.

COTYPES (2): 31345-6, Guadalajara, Jalisco, Mexico, J. J. Major.

Tachymenis canilatus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 20, p. 104, 1868.

= *Chlorosoma elegans* (Tschudi)

HOLOTYPE: 12351, Guayaquil, Ecuador, Destruger and Reeve.

Taeniophis imperialis Kennicott

in Baird, Reptiles of the boundary, in Report U.S. and Mexican Bound. Surv., vol. 2, p. 23, pl. 19, fig. 1, 1859.

= *Coniophanes imperialis imperialis* (Kennicott)

HOLOTYPE: 2060, Brownsville (Cameron County), Texas (Matamoras, Tamaulipas, Mexico according to the catalog), S. Van Vliet.

Taeniophis tantillus Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 7, p. 227, 1854.

= *Leimadophis chamissonis* (Wiegmann)

COTYPE: 5515, near Santiago, Chile, J. M. Gilliss. One cotype was exchanged to ANSP.

Tantilla bimaculata Cope

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8 (1875), p. 143, 1876.

= *Tantilla calamarina* Cope

HOLOTYPE: 6834, Mazatlán, Sinaloa, Mexico, F. Bischoff.

Tantilla calamarina Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 13 (1866), p. 320, 1867.

HOLOTYPE: 6600, Guadalajara, Jalisco, Mexico, J. J. Major. This locality was questioned by R. G. Zweifel.

Tantilla canula Cope

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8 (1875), p. 144, 1876.

COTYPES (2): 24881-2, Yucatán, Mexico, A. Schott.

Tantilla coronata Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 131, 1853.

=*Tantilla coronata coronata* Baird and Girard

HOLOTYPE: 1875, Kemper County, Mississippi, D. C. Lloyd.

Tantilla eiseni Stejneger

Proc. U.S. Nat. Mus., vol. 18, p. 117, 1896.

=*Tantilla eiseni eiseni* Stejneger

HOLOTYPE: 11766a, Fresno, Fresno County, California, G. Eisen, 1879.

PARATYPES (5): 55387-91 (formerly 11766c-g), same data. 11766b was exchanged to MCZ in 1917.

Tantilla flavilineata Smith and Burger

Herpetologica, vol. 6, p. 117, 1950.

PARATYPE: 129524, 8 miles southeast of Nochixtlán, Oaxaca, Mexico, W. L. and J. Burger, Aug. 23, 1949.

Tantilla kirnia Blanchard

Field Mus. Nat. Hist., Zool. Ser., vol. 20, p. 373, 1938.

=*Tantilla nigriceps fumiceps* (Cope)

PARATYPES (2): 15669, San Diego, Duval County, Texas, W. J. Taylor. 22386, San Antonio, Bexar County, Texas, C. S. and H. H. Brimley.

Tantilla lintoni Smith

Proc. Biol. Soc. Washington, vol. 53, p. 61, fig. 1, 1940.

=*Tantillita lintoni* (Smith)

HOLOTYPE: 108603, Piedras Negras, Petén, Guatemala, H. M. Smith, May 24, 1939.

Tantilla miniator Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 15, p. 100, 1863.

=*Tantilla miniata* Cope

HOLOTYPE: 25031, Mirador, Veracruz, Mexico, C. Sartorius.

PROBABLE PARATYPES (2): 25032-3, same data.

Tantilla phrenitica Smith

Zoologica, New York, vol. 27, p. 39, 1942.

HOLOTYPE: 110381, ♀, Cuautlapan, Veracruz, Mexico, H. M. Smith, Aug. 7-14, 1940.

PARATYPES (7): 20835, Totontepec, Oaxaca, Mexico, M. Trujillo, 1887. 38134, Semocock, Guatemala, G. P. Goll, Apr. 8, 1905. 110379-80, Cuautlapan, Veracruz, Mexico, H. M. Smith, January-February 1940. 110383-5, Cuautlapan, Veracruz, Mexico, H. M. Smith, Aug. 7-14, 1940. 110382 was exchanged to MCZ.

Tantilla rubra Cope

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8 (1875), p. 144, 1876.
 HOLOTYPE: 26500, Tapana, Tehuantepec, Oaxaca, Mexico, F. Sumichrast.

Tantilla wilcoxi Stejneger

Proc. U.S. Nat. Mus., vol. 25, p. 156, 1902.

=*Tantilla wilcoxi wilcoxi* Stejneger

HOLOTYPE: 19674, Fort Huachuca (Cochise County), Arizona, T. E. Wilcox, 1892.

Tantilla wilcoxi rubricata Smith

Zoologica, New York, vol. 27, p. 40, 1942.

HOLOTYPE: 110399, ♂, 15 miles west ("southeast" in error in original description) of Galeana, Neuvo León, Mexico, H. M. Smith, Oct. 13, 1939.

PARATYPE: 110398, same data.

Thamnophis cyrtopsis cyclides Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 13, p. 299, 1861.

=*Thamnophis cyrtopsis cyrtopsis* (Kennicott)

HOLOTYPE: 5023, "Cape San Lucas" (probably from Fort Buchanan, Arizona), B. J. D. Irwin.

Thamnophis elegans nigrescens Johnson

Herpetologica, vol. 3, p. 161, 1947.

PARATYPE: 124628 (CPS 3027), Clover Creek Sanatorium, Tacoma, Pierce County, Washington, J. R. Slater, July 8, 1937.

Thamnophis melanogaster canescens Smith

Zoologica, New York, vol. 27, p. 117, 1942.

PARATYPES (3): 110799, La Palma, Michoacan, Mexico, H. M. Smith, Mar. 23, 1939. 110800, Tacúcuaro, Michoacán, Mexico, H. M. Smith, 1939. 134310, near Chapala, Jalisco, Mexico, H. M. Smith, July 2, 1935.

Thamnophis ordinoides errans Smith

Zoologica, New York, vol. 27, p. 112, 1942.

=*Thamnophis errans* Fitch

HOLOTYPE: 46336, ♀, Colonia Garcia, Sierra Madre Mountains at about 8,000 ft., Chihuahua, Mexico, E. W. Nelson and E. A. Goldman, July 1899.

PARATYPES (3): 46337-9, Colonia García, Chihuahua, Mexico, E. W. Nelson and E. A. Goldman, July 1899.

Thamnophis rozellae Smith

Proc. Biol. Soc. Washington, vol. 53, p. 56, 1940.

=*Thamnophis sumichrasti sumichrasti* Cope

HOLOTYPE: 108597, Palenque, Chiapas, Mexico, H. M. Smith, July 12, 1939.

PARATYPES (3): 46549, Monte Cristo, Tabasco, Mexico, E. W. Nelson and E. A. Goldman, May 25, 1900. 108598-9, Aguacate, Chiapas, Mexico, H. M. Smith, Aug. 8, 1939.

Thamnophis sumichrasti cerebrus Smith

Zoologica, New York, vol. 27, p. 111, 1942.

HOLOTYPE: 12734, ♀, Escuintla, Guatemala, H. J. Stuart, 1882.

PARATYPE: 12735, same data.

Thamnophis vicinus Smith

Zoologica, New York, vol. 27, p. 104, 1942.

PARATYPE: 137221, Morelia, Michoacan, Mexico, H. D. Thomas, 1938.

Thrasops praestans Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 20 (1868), p. 309, 1869.

=*Leptophis ahaetulla praestans* (Cope)

COTYPES (2): 6754 (2), near Petén, Guatemala, H. Hague.

Toluca amphisticta Smith and Laufe

Herpetologica, vol. 3, p. 5, 1945.

HOLOTYPE: 120950, ♂, Xuimygopk, between Ayutla and Cacalotepec, 8,650 ft., Oaxaca, Mexico, W. S. Miller, Mar. 15, 1944. Allotype: 120951, ♀, Kachky Walsky, near Cacalotepec, Oaxaca, Mexico, W. S. Miller, Mar. 15, 1944.

Toluca conica Taylor and Smith

Univ. Kansas Sci. Bull., vol. 28, p. 340, fig. 5, pl. 34, fig. 1, 1942.

PARATYPE: 46535, Guajamaloya, Oaxaca, Mexico, E. W. Nelson and E. A. Goldman, Aug. 6, 1894. This specimen was found dead on a road that ran through an oak forest at an altitude of 9,500 ft.

Toluca frontalis Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 16, p. 167, 1864.

=*Pseudoficimia frontalis* (Cope)

COTYPES (2): 31424-5, Colima, Mexico, J. Xantus.

Toluca lineata Kennicott

in Baird, Reptiles of the boundary, in Report U.S. and Mexican Bound. Surv., vol. 2, pt. 2, p. 23, pl. 21, fig. 2, 1859.

=*Toluca lineata lineata* Kennicott

HOLOTYPE: 2103, Valley of Mexico (Toluca on paper label in the jar), Mexico, J. Potts.

PARATYPE: 2104, Valley of Mexico, Mexico, J. Potts.

Toluca lineata wetmorei Smith

Proc. U.S. Nat. Mus., vol. 93, p. 489, 1943.

HOLOTYPE: 110727, ♂, Pan de Olla, Veracruz, Mexico, H. M. Smith, Mar. 22, 1940.

PARATYPES (19): 46405, 46442, Las Vigas, Veracruz, Mexico, E. W. Nelson, June 1893. 110726, 110728-43, Pan de Olla, Veracruz, Mexico, H. M. Smith, Mar. 22, 1940.

Tomodon nasutus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 16, p. 166, 1864.
 =*Manolepis putnami* (Jan)
 HOLOTYPE: 31478, Colima, Colima, Mexico, J. Xantus, July 1863.
 PARATYPE: 31479, same data.

Toxicophis pugnax Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 20, 1853.
 =*Ancistrodon piscivorus leucostoma* (Troost)
 HOLOTYPE: 61063 (formerly 4262), head only, Indianola (Calhoun County), Texas, J. D. Graham.

Tretanorhinus nigroluteus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 13, p. 298, 1861.
 =*Tretanorhinus nigroluteus nigroluteus* Cope
 HOLOTYPE: 5568, "Greytown, Nicaragua, Dr. Caldwell" (probably Aspinwall, Panama, J. Rowell), 1857.

Tretanorhinus variabilis binghami Schwartz and Ogren

Herpetologica, vol. 12, p. 105, 1956.
 PARATYPE: 136606, Finca Bucares, 22 km. southeast of Bueycito, Río Yao, Oriente Province, Cuba, L. H. Ogren and A. Schwartz, Dec. 26, 1954.

Trimeresurus flavoviridis tinkhami Gloyd

Bull. Chicago Acad. Sci., vol. 10, p. 123, figs. 1-3, 1955.
 HOLOTYPE: 137285, ♂, north slope of mountains south of Higa, in southeastern part of Kume Shima, Okinawa Group, Riu Kiu (Ryukyu) Islands, E. R. Tinkham, August 1945.
 PARATYPE: 137286, ♀, same data.

Trimeresurus garciai Smith

Proc. Biol. Soc. Washington, vol. 53, p. 62, 1940.
 =*Bothrops melanurus* (Müller)
 PARATYPE: 108602, Cacaloapam, Puebla, Mexico, H. M. Smith, Aug. 21, 1939.

Trimeresurus nummifer picadoi Dunn

Proc. Biol. Soc. Washington, vol. 52, p. 165, 1939.
 =*Bothrops picadoi* (Dunn)
 HOLOTYPE: 37753, ♀, La Palma, Costa Rica, 4,500 ft., E. Fernandez.

Trimeresurus stejnegeri Schmidt

Bull. Amer. Mus. Nat. Hist., vol. 54, p. 546, 1927.
 PARATYPES (2): 64022-3, Moh Kan Shan, Chekiang, China, C. H. Barlow.

Trimeresurus yucatanicus Smith

Zoologica, New York, vol. 26, p. 62, 1941.

=*Bothrops yucatanicus* (Smith)

HOLOTYPE: 46571, ♀, Chichén-Itzá, Yucatán, Mexico, E. W. Nelson and E. A. Goldman, February 1901.

Trimerorhinus tritaeniatus multisquamis Loveridge

Proc. Biol. Soc. Washington, vol. 45, p. 84, 1932.

PARATYPES (19): 40893, Juja Farm, Kenya, Smithsonian-African Expedition, 1908-1909. 40901-2, 41516-8, 41672-3, Lake Naivasha, Kenya, Smithsonian-African Expedition, July-August 1909. 40968, Nairobi, Kenya, H. Edgell, 1908-09. 40970, 40997, 41973, Guaso Nyiro, Kenya. E. Heller and J. A. Loring, June-July 1909. 41534, Athi Plains, Kenya. E. Heller, May 20, 1909. 42008, Kapiti Plains, Kenya, E. A. Mearns and E. Heller, Apr. 26, 1909. 42061, Kenya, Smithsonian-African Expedition 1909. 48584-6, Mtoto Andei Station, Uganda, E. Heller, 1911. 49080, Kaímosi, Kenya, E. Heller, Jan. 27, 1912. 40894, 40903, and 48587 were exchanged to MCZ in 1930.

Trimorphodon biscutatus quadruplex Smith

Proc. U.S. Nat. Mus., vol. 91, p. 157, 1941.

HOLOTYPE: 89476, ♀, Esteli, Nicaragua, J. H. Ivy, 1932.

Trimorphodon biscutatus semirutus Smith

Proc. U.S. Nat. Mus., vol. 93, p. 492, 1943.

HOLOTYPE: 110410, Acapulco, Guerrero, Mexico, H. M. Smith, Sept. 3, 1939.

Trimorphodon collaris Cope

Journ Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8 (1875), p. 131, 1876.

HOLOTYPE: 26499, Tuxpango, near Orizaba, Veracruz, Mexico, F. Sumichrast.

Trimorphodon fasciolata Smith

Proc. U.S. Nat. Mus., vol. 91, p. 160, 1941.

HOLOTYPE: 110400, ♂, Zarácua Falls, Uruapán, Michoacán, Mexico, H. M. Smith, Mar. 12, 1939.

Trimorphodon forbesi Smith

Proc. U.S. Nat. Mus., vol. 91, p. 163, 1941.

HOLOTYPE: 110402, ♂, near San Diego, south of Tehuacán, Puebla, Mexico, H. M. Smith, Dec. 17, 1939.

Trimorphodon lambda Cope

Proc. Amer. Philos. Soc., vol. 23, p. 286, 1886.

HOLOTYPE: 13487, Guaymas, Sonora, Mexico, H. F. Emeric, 1883.

Trimorphodon major Cope

Proc. Amer. Philos. Soc., vol. 11, p. 153, 1869.

=*Trimorphodon biscutatus biscutatus* (Duméril and Bibron)

COTYPES (3): 30427-9, near Tehuantepec, Oaxaca, Mexico, F. Sumichrast.

Trimorphodon tau Cope

Proc. Amer. Philos. Soc., vol. 11, p. 152, 1869.

HOLOTYPE: 30333, Quiotepec, between Tehuacán and Oaxaca, State of Oaxaca, Mexico, F. Sumichrast

Trimorphodon upsilon Cope

Proc. Amer. Philos. Soc., vol. 11, p. 151, 1869.

HOLOTYPE: 31358, Guadalajara, Jalisco, Mexico, J. J. Major.

Trimorphodon vilkinsonii Cope

Proc. Amer. Philos. Soc., vol. 23, p. 285, 1886.

HOLOTYPE: 14268, Chihuahua, Mexico, E. Wilkinson.

Tropidoclonium storerioides Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 17, p. 190, 1865.

=*Storeria storerioides* (Cope)

COTYPES (4): 24987-90, Mexican plateau between the eastern range and the Valley of Mexico, Mexico, C. Sartorius.

Tropidodipsas brevifacies Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 18, p. 127, 1866.

=*Dipsas brevifacies* (Cope)

HOLOTYPE: 24836, Yucatán, Mexico, A. Schott, Sept. 27, 1865.

Tropidodipsas lunulata Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 12 (1860), p. 517, 1861.

=*Pseustes poecilonotus poecilonotus* (Günther)

HOLOTYPE: 12398, Honduras, A. H. Riise?.

Tropidonotus bisectus Cope

Proc. U.S. Nat. Mus., vol. 10, p. 146, 1887.

=*Natrix erythrogaster transversa* (Hallowell)

HOLOTYPE: 14643, Fish Commission Armory Grounds, Washington, D.C., J. E. Brown, Nov. 2, 1886.

Tropidonotus celaeno Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 12, p. 341, 1860.

=*Natrix valida* (Kennicott)

COTYPES (8): 5281 (7), 131748, San José River, Cape San Lucas, Baja California, Mexico, J. Xantus, May 15, 1859. 5281a is the lectotype. One cotype from 5281 was exchanged to ANSP.

Tropidonotus compressicaudus walkeri Yarrow

Proc. U.S. Nat. Mus., vol. 6, p. 154, 1833.

=*Natrix sipedon compressicauda* (Yarrow)

HOLOTYPE: 10681, Clearwater, Pinellas County, Florida, S. T. Walker, October 1879.

Tropidonotus rhombifer Hallowell

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 177, 1852.

=*Natrix rhombifera rhombifera* (Hallowell)

COTYPE: 7253, Arkansas River, near north boundary of the Creek Nation, Indian Territory (near Keystone, Tulsa County, Oklahoma), S. W. Woodhouse.

Tropidonotus tephroleura Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 12, p. 341, 1860.

=*Natrix valida valida* (Kennicott)

COTYPES (6): 4683 (6), Cape San Lucas, Baja California, Mexico, J. Xantus.

Tropidonotus transversus Hallowell

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 177, 1852.

=*Natrix erythrogaster transversa* (Hallowell)

COTYPE: 1316, Arkansas River at Creek Boundary, Indian Territory (near Keystone, Tulsa County, Oklahoma), S. W. Woodhouse.

Tropidonotus ustus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 12, p. 340, 1860.

=*Natrix sipedon compressicauda* (Kennicott)

HOLOTYPE: 5286, Charlotte Harbor, Charlotte County, Florida, E. M. Baker.

Tropidophis maculatus jamaicensis Stull

Occ. Pap. Mus. Zool. Univ. Michigan, No. 195, p. 12, 1928.

PARATYPES (3): 42878-80, Montego Bay, Jamaica, E. A. Andrews, July-August 1910. These specimens are also listed as paratypes of *Tropidophis pardalis stejnegeri* Grant, in Lynn and Grant, Bull. Inst. Jamaica, Sci. Ser., No. 1, p. 120, 1940.

Tropidophis pardalis androsi Stull

in Lynn and Grant, Bull. Inst. Jamaica, Sci. Ser., No. 1, p. 119, 1940.

Occ. Pap. Mus. Zool. Univ. Michigan, No. 195, p. 34, pl. 2, fig. 5, 1928.

HOLOTYPE: 49471, ♂, Andros Island, West Indies, P. Bartsch, May 20, 1912, West Indies, P. Bartsch, 1921.

Tropidophis pardalis barbouri Bailey

Proc. New England Zool. Club, vol. 16, p. 49, 1937.

PARATYPES (2): 120804-5 (formerly MCZ 37905, 37924), Bannerman Town, South Eleuthera Island, Bahama Islands, T. Barbour, February 1934.

Tropidophis pardalis stejnegeri Grant

in Lynn and Grant, Bull. Inst. Jamaica, Sci. Ser., No. 1, p. 119, 1940.

PARATYPES (3): 42878-80, Montego Bay, Jamaica, E. A. Andrews, July-August 1910. These specimens are also listed as paratypes of *Tropidophis maculatus jamaicensis* Stull, Occ. Pap. Mus. Zool. Univ. Michigan, No. 195, p. 12, 1928.

Tropidophis wrighti Stull

Occ. Pap. Mus. Zool. Univ. Michigan, No. 195, p. 38, pl. 2, fig. 7, 1928.

HOLOTYPE: 12420, ♂, eastern Cuba.

Typhlops basimaculatus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 18 (1866), p. 320, 1867.

HOLOTYPE: 6602, Córdoba and Orizaba, Veracruz, Mexico, F. Sumichrast.

Typhlops mandensis Stejneger

Proc. U.S. Nat. Mus., vol. 16, p. 725, 1894.

HOLOTYPE: 20125, Wange, Island of Manda, north of Lamu, East Africa (Kenya), G. Denhardt.

Typhlops manni Loveridge

Proc. U.S. Nat. Mus., vol. 91, p. 118, 1941.

HOLOTYPE: 109634, Harbel, Liberia, W. M. Mann, Mar. 10-July 17, 1940.

Typhlops microstomus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 18, p. 125, 1866.

HOLOTYPE: 61064, Yucatán, Mexico, A. Schott.

PROBABLE PARATYPE: 6569, Mérida, Yucatán, Mexico, A. Schott, Apr. 20, 1866.

Typhlops praeocularis Stejneger

Proc. U.S. Nat. Mus., vol. 16, p. 709, 1894.

HOLOTYPE: 20799, Stanley Pool, Leopoldville, Congo Free State (Belgian Congo), J. H. Camp.

Typhlops rostellatus Stejneger

Report of the U.S. National Museum . . . for the year ending June 30, 1902, p. 686, figs. 145-147, 1904.

HOLOTYPE: 25463, Lares, Puerto Rico, A. B. Baker, Jan. 26, 1899.

PARATYPES (3): 25461-2, Lares, Puerto Rico, A. B. Baker, Jan. 26-30, 1899. 27460, Puerto Rico, C. E. Adams, 1900.

Typhlops stejnegeri Loveridge

Copeia, No. 3, p. 92, figs. 1-2, 1931.

HOLOTYPE: 23979, Luebo District, Belgian Congo, D. W. Snyder, 1897.

PARATYPE: 23980, same data.

Typhlops sulcatus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 20, p. 128, 1868.

HOLOTYPE: 12371, Navassa Island (West Indies), W. J. Rasin.

Ungalia bucculenta Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 20, p. 129, 1868.

=*Tropidophis pardalis bucculentus* (Cope)

COTYPES (3): 12377 (3), Navassa Island (West Indies), W. J. Rasin.

Ungalia cana Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 20, p. 129, 1868.

=*Tropidophis pardalis canus* (Cope)

COTYPES (6): 7111 (5), south side of Inagua Island, Bahama Islands, T. Blain. 26763, Inagua Island, Bahama Islands.

Ungaliophis panamensis Schmidt

Smithsonian Misc. Coll., vol. 89, No. 1, p. 12, 1933.

HOLOTYPE: 54059, Cerro Bruja, Panama, 2,000 ft., E. A. Goldman, June 6, 1911.

Ungualia haetiana Cope

Proc. Amer. Philos. Soc., vol. 18, p. 273, 1879.

=*Tropidophis maculatus haetianus* (Cope)

COTYPES (2): 10164, Port-au-Prince, Haiti, J. J. Brown. 10169, Île de la Gonâve, Haiti, J. J. Brown.

Uromacer scandax Dunn

Proc. New England Zool. Club, vol. 7, p. 43, 1920.

HOLOTYPE: 59438, Tortuga Island, Haiti, W. L. Abbott, Jan. 31, 1917.

Uromacer wetmorei Cochran

Proc. Biol. Soc. Washington, vol. 44, p. 91, 1931.

HOLOTYPE: 83891, Beata Island, Dominican Republic, A. Wetmore and F. C. Lincoln, May 13, 1931.

Urotheca williamsi Roze

Breviora, Mus. Comp. Zool., No. 88, p. 1, 1958.

PARATYPE: 121206, ♂, Colonia Tovar, 6,000 ft., Estado Aragua, Venezuela, E. G. Holt, Mar. 30, 1929.

Virginia valeriae Baird and Girard

Smithsonian Misc. Coll., vol. 2, art. 5, p. 127, 1853.

=*Haldea valeriae valeriae* (Baird and Girard)

HOLOTYPE: 1962, Kent County, Maryland, V. Blaney, 1847.

PARATYPE: 1876, Anderson, Anderson County, South Carolina, C. Paine.

Wenona isabella Baird and Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 176, 1852.

=*Charina bottae bottae* (Blainville)

HOLOTYPE: 7299, Puget Sound (Washington), U.S. Exploring Expedition, 1841.

Wenona plumbea Baird and Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 176, 1852.

=*Charina bottae bottae* (Blainville)

HOLOTYPE: 4492, Puget Sound (Washington), U.S. Exploring Expedition, 1841.

Xenodon ancorus Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 8, p. 182, 1857.

=*Holarchus ancorus* (Girard)

HOLOTYPE: 5521, Manila, Luzon, Philippine Islands, U.S. Exploring Expedition, 1842.

Xenodon mexicanus Smith

Proc. Biol. Soc. Washington, vol. 53, p. 57, 1940.

HOLOTYPE: 108596, Piedras Negras, Petén, Guatemala, H. M. Smith, June 12, 1939.

Xiphosoma annulatum Cope

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8 (1875), p. 129, pl. 28, fig. 6, 1876.

=*Boa annulata* (Cope)

HOLOTYPE: 32480, Costa Rica, W. M. Gabb.

Xiphosoma madagascariense Duméril and Bibron

Erpétologie générale ou histoire naturelle complète des reptiles, vol. 6, p. 549, 1844.

=*Sanzinia madagascariense* (Duméril and Bibron)

COTYPE: 10982, Madagascar, JPP.

Zamenis conirostris Cope

Amer. Nat., vol. 29, p. 679, 1895.

=*Coluber constrictor stejnegerianus* (Cope)

HOLOTYPE: 1768, Matamoros, Tamaulipas, Mexico, D. N. Couch.

Zamenis lateralis fuliginosus Cope

Amer. Nat., vol. 29, p. 679, 1895.

=*Masticophis flagellum piceus* (Cope)

COTYPES (2): 15135-6, Santa Margarita Island, Baja California, Mexico, U.S. Fish Commission *Albatross*, May 2, 1888.

Zamenis stejnegerianus Cope

Amer. Nat., vol. 29, p. 678, fig. 172, 1895.

=*Coluber constrictor stejnegerianus* (Cope)

HOLOTYPE: 17065, Cameron County, Texas, C. K. Worthen.

Zaocys oshimai Stejneger

Proc. U.S. Nat. Mus., vol. 66, No. 2562, p. 88, 1925.

HOLOTYPE: 52267, Urai, Formosa, F. Baker, Nov. 18, 1914.

Order Testudines

Amyda spinifera hartwegi Conant and Goin

Occ. Pap. Mus. Zool. Univ. Mich., No. 510, p. 1, pl. 1, 1948.

=*Trionyx spinifer hartwegi* (Conant and Goin)

PARATYPES (6): 55683, Greenwood County, Kansas, J. Hurter, July-August 1912. 91022, 5 miles north of Winfield, Cowley County, Kansas, L. Hoyle, May 28, 1933. 95301, 11 miles southeast of Winfield, Cowley County, Kansas, C. E. Burt, Aug. 31, 1934. 100529-30, Winfield, Cowley County, Kansas, C. E. Burt, Aug. 12, 1935. 123446, Topeka, Shawnee County, Kansas River, Kansas, C. E. Burt, June 26, 1945.

Aspidonectes asper Agassiz

Contributions to the natural history of the United States of America, vol. 1, p. 405, 1857.

=*Trionyx spinifer asper* (Agassiz)

COTYPES (2): 01086, skull, carapace, and parts of plastron, Washington, Adams County, Mississippi, B. L. C. Wailes. 012349, carapace, Lake Concordia (Concordia Parish), Louisiana, B. L. C. Wailes, 1851.

Aspidonectes emoryi Agassiz

Contributions to the natural history of the United States of America, vol. 1, p. 407, vol. 2, pl. 6, figs. 4-5, 1857.

=*Trionyx spinifer emoryi* (Agassiz)

COTYPE: 7855, Brownsville, Cameron County, Texas, C. B. R. Kennerly.

Caretta remivaga Hay

Proc. U.S. Nat. Mus., vol. 34, p. 194, pl. 10, pl. 11, fig. 5, 1908.

=*Lepidochelys olivacea* (Eschscholtz)

HOLOTYPE: 09973, skull, Ventosa Bay, Oaxaca, Mexico, F. Sumichrast.

Caretta rostrata Girard

in U.S. Exploring expedition . . . 1838-42, vol 20 (Herpetology),
p. 446, Atlas, pl. 30, figs. 8-13, 1858.

=*Eretmochelys imbricata squamata* Agassiz

PROBABLE COTYPES (2): 012387-8, carapaces, Fiji Islands, U.S. Exploring Expedition, 1840.

Chelonia formosa Girard

in U.S. Exploring Expedition . . . 1838-42, vol. 20 (Herpetology), p.
456, Atlas, pl. 31, figs. 1-4, 1858.

=*Chelonia mydas japonica* Thunberg

HOLOTYPE: 012386, carapace, Fiji Islands, U.S. Exploring Expedition, 1840.

Chelonia tenuis Girard

in U.S. Exploring Expedition . . . 1838-42, vol. 20 (Herpetology), p.
459, Atlas, pl. 31, fig. 8, 1858.

=*Eretmochelys imbricata bissa* Rüppell

COTYPE: 012390, Rosa Island (Tuamotu Islands, South Pacific), U.S. Exploring Expedition, 1840.

Chelopus funereus Cope

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8, p. 154, 1875.

=*Geoemyda funerea* (Cope)

COTYPES (4): 45900-1, 46134-5, Port Limón, Costa Rica, W. M. Gabb.

Chelopus gabbii Cope

Journ. Acad. Nat. Sci. Philadelphia, ser. 2, vol. 8, p. 153, 1875.

=*Geoemyda gabbii* (Cope)

HOLOTYPE: 45905, Costa Rica, W. M. Gabb.

Chelopus rubidus Cope

Proc. Amer. Philos. Soc., vol. 11, p. 148, 1869.

=*Geoemyda rubida rubida* (Cope)

COTYPES (3): 45612-4, Juchitán, Tehuantepec, Oaxaca, Mexico, F. Sumichrast.

Chelydra osceola Stejneger

Proc. Biol. Soc. Washington, vol. 31, p. 89, 1918.

=*Chelydra serpentina osceola* Stejneger

HOLOTYPE: 10369, ♂, Clearwater, Pinellas County, Florida, S. T. Walker, September 1879.

PARATYPES (7): 28761, shell and bones, Kissimmee River, Florida, E. A. Mearns, May 3, 1901. 29208, Alligator Bluff, Kissimmee River, Osceola County, Florida, E. A. Mearns, Apr. 24, 1901. 30013, ♀, skin, Alligator

Bluff, Kissimmee River, Osceola County, Florida, E. A. Mearns. 55317, shell, Vero Beach, Brevard County, Florida, I. M. Weills, Mar. 14, 1915. 60545-6, bones, ♂, ♀, Auburndale, Polk County, Florida, N. R. Wood, 1918. 60557, Auburndale, Polk County, Florida, N. R. Wood, Mar. 18, 1918.

Chrysemys dorsalis Agassiz

Contributions to the natural history of the United States of America, vol. 1, p. 440, 1857.

=*Chrysemys picta dorsalis* Agassiz

COTYPE: 21, Washington County, Mississippi, B. L. C. Wailes.

Chrysemys treleasei Hurter

Trans. Acad. Sci. St. Louis, vol. 20, p. 235, 1911.

=*Chrysemys picta bellii* Gray

COTYPES (3): 49427-8, St. Clair County, Illinois, J. Hurter, July 3, 1908, June 16, 1895. 49429, Madison County, Illinois, J. Hurter, July 9, 1905.

Cinosternum berendtianum Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 17, p. 189, 1865.

=*Kinosternon acutum* Gray

COTYPES (3): 6517, ♀, 106293-4, ♂, all shells, Tabasco, Mexico, C. H. Berendt, 1864.

Cinosternum brevigulare Cope

Proc. Amer. Philos. Soc., vol. 22, p. 389, 1885. (Not of Günther, May 1885.)

=*Kinosternon leucostomum* Duméril and Bibron

COTYPES (3): 19797, Old Harbor, Costa Rica, W. M. Gabb. 45582, ♀, 51165, ♂, Sipurio, Costa Rica, W. M. Gabb.

Cinosternum postinguinale Cope

U.S. Nat. Mus. Bull., No. 32, p. 23, 1887. (New name for *Cinosternum brevigulare* Cope, preoccupied.)

=*Kinosternon leucostomum* Duméril

COTYPES (3): 19797, Old Harbor, Costa Rica, W. M. Gabb. 45582, ♀, 51165, ♂, Sipurio, Costa Rica, W. M. Gabb.

Cistudo ornata Agassiz

Contributions to the natural history of the United States of America, vol. 1, p. 445, 1857.

=*Terrapene ornata ornata* (Agassiz)

COTYPES (4): 57, ♀, Yellowstone River, Montana, F. V. Hayden. 7541, ♂, southern boundary of Kansas, W. S. Wood and J. H. Clark. 7542, ♂, Illinois, R. W. Kennicott. 7547, ♀, Republican River, Kansas, W. S. Wood.

Cistudo triunguis Agassiz

Contributions to the natural history of the United States of America, vol. 1, p. 445, 1857.

=*Terrapene carolina triunguis* (Agassiz)

COTYPES (2): 86871-2 (formerly MCZ 1519), New Orleans, Orleans Parish, Louisiana, N. B. Benedict.

Claudius angustatus Cope

Proc. Acad. Nat. Sci., Philadelphia, vol. 17, p. 187, pl. 9, 1865.

COTYPES (2): 6518, 6525, Tabasco, Mexico, C. H. Berendt, 1863.

Claudius severus Cope

Proc. Acad. Nat. Sci. Philadelphia, vol. 24, p. 24, 1872.

=*Staurotypus salvinii* Gray

HOLOTYPE: 64005, shell, Santa Efigenia, west side of Isthmus of Tehuantepec, Oaxaca, Mexico, F. Sumichrast.

Clemmys nuchalis Dunn

Bull. Amer. Mus. Nat. Hist., vol. 37, p. 624, text fig. 6, pl. 61, 1917.

=*Clemmys muhlenbergii* Schoepff

PARATYPE: 62016, ♀, Brevard, Transylvania County, North Carolina, E. R. Dunn, July 12, 1916.

Deirochelys reticularia miaria Schwartz

Chicago Nat. Hist. Mus., Fieldiana: Zool., vol. 34, p. 486, figs. 107-8, 1956.

ALLOTYPE: 85145, ♀, White Rock Creek, near Dallas, Dallas County, Texas, C. E. Burt, Apr. 16, 1932.

Emys marmorata Baird and Girard

Proc. Acad. Nat. Sci. Philadelphia, vol. 6, p. 177, 1852.

=*Clemmys marmorata marmorata* (Baird and Girard)

COTYPES (5): 88, 7594-6, 131830 (formerly 7593), Puget Sound (Washington), U.S. Exploring Expedition, 1841.

Goniochelys minor Agassiz

Contributions to the natural history of the United States of America, vol. 1, p. 424, 1857.

=*Sternotherus minor minor* (Agassiz)

COTYPE: 7111 (formerly MCZ 1572), New Orleans, Orleans Parish, Louisiana, N. B. Benedict.

Gopherus flavomarginatus Legler

Univ. Kansas Publ., Mus. Nat. Hist., vol. 11, p. 337, fig. 1, pls. 7-8, 1959.

HOLOTYPE: 61253, 30 to 40 miles from Lerdo, Durango, Mexico, A. Busck.

PARATYPES (2): 60976, Durango, Mexico, E. Chaffee, May 1918. 61254, 30 to 40 miles from Lerdo, Durango, Mexico, A. Busck.

Graptemys pseudogeographica ouachitensis Cagle

Occ. Pap. Mus. Zool. Univ. Michigan, No. 546, p. 10, figs., 1953.

PARATYPE: 139733 (formerly UIMNH 26717), 4 miles east of Harrisonburg, Catahoula Parish, Louisiana, Tulane Univ. collecting party, June 10-11, 1950.

Graptemys pseudogeographica sabinensis Cagle

Occ. Pap. Mus. Zool. Univ. Mich., No. 546, p. 2, figs. 1, 3a, 3e, 3g, 1953.

PARATYPE: 134312 (formerly UIMNH 26722), 8 miles southwest of Negreet, Sabine Parish, Louisiana, Tulane Univ. collecting party, July 5-9, 1950.

Graptemys pseudogeographica versa Stejneger

Journ. Washington Acad. Sci., vol. 15, p. 463, 1925.

=*Graptemys versa* Stejneger

HOLOTYPE: 27473, Austin, Travis County, Texas, C. S. and H. H. Brimley, July 1900.

PARATYPES (7): 27474-80, same data.

Graptemys pulchra Baur

Amer. Nat., vol. 27, p. 675, 1893.

COTYPES (2): 8308, ♀, 029526, ♀, lake near Montgomery, Montgomery County, Alabama, T. H. Bean and L. Kumlien, July 1876.

Kinosternon abaxillare Baur

in Stejneger, Journ. Washington Acad. Sci., vol. 15, p. 462, 1925.

HOLOTYPE: 7518, ♂, shell, Tuxtla Gutiérrez, Chiapas, Mexico, C. H. Berendt.

PARATYPES (11): 7519-29, shells, same data.

Kinosternon bauri palmarum Stejneger

Journ. Washington Acad. Sci., vol. 15, p. 463, 1925.

HOLOTYPE: 61065, ♀, Paradise Key, Royal Palm State Park, Dade County, Florida, C. A. Mosier.

PARATYPES (2): 60597, shell, Royal Palm Hammock, Royal Palm State Park, Dade County, Florida, Apr. 6, 1918. 69559, Royal Palm State Park, Dade County, Florida, C. A. Mosier, Oct. 30, 1918.

Kinosternon cruentatum consors Stejneger

Proc. U.S. Nat. Mus., vol. 90, p. 458, 1941.

HOLOTYPE: 13912, ♂, Cozumel Island, Yucatán, Mexico, U.S. Fish Commission *Albatross*, Jan. 23-29, 1885.

PARATYPES (4): 6556, ♂, Progreso, Yucatán, Mexico, A. Schott. 13910-1, 13913, ♀, ♂, ♀, Cozumel Island, Yucatán, Mexico, U.S. Fish Commission *Albatross*, Jan. 23-29, 1885.

Kinosternon flavescens spooneri Smith

Bull. Chicago Acad. Sci., vol. 9, No. 10, p. 195, pl. 2, 1951.

PARATYPE: 83190, ♂, Meredosia, Morgan County, Illinois, D. H. Thompson, June 1927.

Kinosternon herrerae Stejneger

Journ. Washington Acad. Sci., vol. 15, p. 462, 1925.

HOLOTYPE: 61249, ♂, Xochimilco, Valley of Mexico, Mexico (restricted to La Laja, Veracruz), A. L. Herrera.

PARATYPES (3): 61250-2, ♀, ♀, ♀, same data.

Kinosternon louisianae Baur

Amer. Nat., vol. 27, p. 676, 1893.

=*Kinosternon subrubrum hippocrepis* Gray

HOLOTYPE: 15527, New Orleans, Orleans Parish, Louisiana, R. W. Shufeldt, 1833.

Kinosternon murrayi Glass and Hartweg

Copeia, No. 1, p. 50, fig. 1, 1951.

PARATYPE: 15860, Marfa, Presidio County, Texas, V. O. Bailey, Jan. 24, 1890.

Kinosternon panamensis Schmidt

Smithsonian Misc. Coll., vol. 106, No. 8, p. 5, 1946.

HOLOTYPE: 117369 (formerly MCZ 7996), Panama Railroad, Panama Canal Zone, T. Barbour, June 10, 1922.

PARATYPES (2): 7864, Panama. 117370 (formerly MCZ 7996), ♀, Panama Railroad, Panama Canal Zone, T. Barbour.

Malaclemmys littoralis Hay

Bull. [U.S.] Bureau of Fisheries, vol. 24, p. 18, pls. 8-9, pl. 12, figs. 2-3, 1904.

=*Malaclemmys terrapin littoralis* Hay

HOLOTYPE: 33913, Rockport, Aransas County, Texas, W. P. Hay.

Malaclemmys macrospilota Hay

Bull. [U.S.] Bureau of Fisheries, vol. 24, p. 16, pls. 6-7, pl. 11, fig. 1, 1904.

=*Malaclemys terrapin macrospilota* Hay

HOLOTYPE: 33917, ♀, Charlotte Harbor, Charlotte County, Florida, W. P. Hay, December 1902.

Malaclemmys terrapin tequesta Schwartz

Proc. Biol. Soc. Washington, vol. 68, p. 158, 1955.

PARATYPE: 37020, New Smyrna, Volusia County, Florida, B. W. Evermann, Dec. 3, 1905.

Malacoclemmys oculifera Baur

Science, ser. 1, vol. 16, p. 262, 1890.

=*Graptemys oculifera* (Baur)

HOLOTYPE: 15511, Mandeville, Saint Tammany Parish, Louisiana, G. Kohn.

PARATYPES (3): 15508-15510, same data. 15510 is a dry shell.

Ozotheca tristycha Agassiz

Contributions to the natural history of the United States of America, vol. 1, p. 425, vol. 2, pl. 5, figs. 20-22, 1857.

=*Sternotherus odoratus* Latreille

COTYPES (4): 69, 71-2, San Pedro near San Antonio (Bexar County), Texas, C. B. R. Kennerly. 7890, Medina River, Texas, C. B. R. Kennerly.

Platypeltis agassizii Baur

Amer. Nat., vol. 22, p. 1121, 1888.

=*Trionyx spinifer asper* Agassiz

PARATYPE: 92584, (formerly MCZ 1601), ♀, skull, Savannah, Chatham County, Georgia.

Platythyra flavescens Agassiz

Contributions to the natural history of the United States of America, vol. 1, p. 430, vol. 2, pl. 5, figs. 12-15, 1857.

=*Kinosternon flavescens flavescens* (Agassiz)

COTYPES (3): 50, 131823 (formerly 7867), ♂, Rio Blanco, Texas, C. B. R. Kennerly. 7892, Gila River, Camp Yuma (Yuma County), Arizona, R. O. Abbott.

Pseudemys alabamensis Baur

Proc. Amer. Philos. Soc., vol. 31, p. 224, 1893.

COTYPES (2): 20966-7, ♂, ♀, skins and skulls, Mobile Bay, Alabama, G. Kohn, May 7-11, 1885.

***Pseudemys decorata* Barbour and Carr**

Mem. Mus. Comp. Zool. vol. 54, No. 5, p. 409, pl. 9, figs. 3-4, 1940.

PARATYPES (5): 59093-6, Thomazeau, Haiti, J. B. Henderson and P. Bartsch, Apr. 2, 1917. 81091, 10 miles north of Port-au-Prince, W. M. Perrygo, 1930.

***Pseudemys elonae* Brimley**

Journ. Elisha Mitchell Sci. Soc., vol. 44, No. 1, p. 67, pl. 1, pl. 2, fig. 1, 1928.

=*Pseudemys floridana concinna* LeConte

HOLOTYPE: 79631, ♂, skin, Guilford County, not far from Elon College, North Carolina, D. W. Rumbold and J. F. Hall, November 1927.

***Pseudemys felis* Barbour**

Occ. Pap. Boston Soc. Nat. Hist., vol. 8, p. 205, 1935.

PARATYPE: 102447 (formerly MCZ 38388), Cat Island, Bahama Islands, W. G. Albury and C. S. Dolley, 1935.

***Pseudemys floridana peninsularis* Carr**

Copeia, No. 3, p. 105, 1938.

PARATYPES (2): 104389, ♀, Crystal Springs, Pasco County, Florida, L. Marchand, Feb. 20, 1938. 104390, ♂, near Umatilla, Lake County, Florida, A. F. Carr, Jr., Mar. 16, 1934.

***Pseudemys scripta gaigeae* Hartweg**

Occ. Pap. Mus. Zool. Univ. Mich., No. 397, p. 1, 1939.

PARATYPES (2): 60921, shell, Lerdo, Durango, Mexico, A. Busck, May 1918. 103706, skin and skull, Rio Nazos, Durango, Mexico, D. N. Couch.

***Pseudemys stejnegeri* Schmidt**

in New York Acad. Sci., Scientific Survey of Porto Rico and the Virgin Islands, vol. 10, pt. 1, p. 147, fig. 51, 1928.

HOLOTYPE: 25642, ♀, skull, shell, and feet, San Juan, Puerto Rico, U.S. Fish Commission *Fish Hawk*, Jan. 14, 1899.

PARATYPES (3): 25643-4, San Juan, Puerto Rico, U.S. Fish Commission *Fish Hawk*, Jan. 14, 1899. 25653, ♂, skin and skull, Caguas, Puerto Rico, U.S. Fish Commission *Fish Hawk*, Jan. 9, 1899.

***Pseudemys texana* Baur**

Proc. Amer. Philos. Soc., vol. 31, p. 223, 1893.

=*Pseudemys floridana texana* Baur

PARATYPES (2): 76, near Cadereita, Nuevo León, Mexico, D. N. Couch. 80, Pecos River, Guadalupe Mountains, Texas, T. J. Pope. A third specimen is a probable paratype: 7173, ♀, Old Fort Cobb (Caddo County), Oklahoma, E. Palmer, June 1, 1868.

***Pseudemys vioscana* Brimley**

Journ. Elisha Mitchell Sci. Soc., vol. 44, No. 1, p. 66, pl. 2, fig. 2, 1928.

=*Pseudemys floridana mobilensis* Holbrook

HOLOTYPE: 79632, ♂, skin, Lake des Allemands, Louisiana, P. Viosca, Jr., April 1927.

PARATYPES (2): 79633, 79634, skins, same data.

***Ptychemys hoyi* Agassiz**

Contributions to the natural history of the United States of America, vol. 1, p. 433, 1857.

=*Pseudemys floridana hoyi* (Agassiz)

NEOTYPE: 55516, ♀, skin, Newton County, Missouri, J. H. Black, 1899.

***Terrapene bauri* Taylor**

Proc. U.S. Nat. Mus., vol. 17, p. 576, fig. 3, 1895.

=*Terrapene carolina bauri* Taylor

HOLOTYPE: 8352, Florida, F. B. Meek, 1875.

***Terrapene goldmani* Stejneger**

Proc. Biol. Soc. Washington, vol. 46, p. 119, 1933.

=*Terrapene mexicana mexicana* Gray

HOLOTYPE: 46251, ♀, Chijol, San Luis Potosí, Mexico, E. W. Nelson and E. A. Goldman, May 11, 1898.

***Terrapene nelsoni* Stejneger**

Journ. Washington Acad. Sci., vol. 15, p. 463, 1925.

HOLOTYPE: 46252, ♂, Pedro Pablo, Nayarit, Mexico, 2,500 ft., E. W. Nelson and E. A. Goldman, Aug. 4, 1897.

***Testudo kleinmanni* Lortet**

Arch. Mus. Lyons, vol. 3, p. 188, 1883.

COTYPE: 10979, Egypt, JPP.

***Testudo mauritanica* Duméril and Bibron**

Erpétologie générale, ou histoire naturelle complète des reptiles, vol. 2, p. 44, 1835.

COTYPE: 10980, Algeria, JPP.

***Thalassochelys corticata* Girard**

in U.S. Exploring Expedition . . . 1838-42, vol. 20 (Herpetology), p. 451, pl. 29, 1858.

=*Caretta caretta caretta* Linnaeus

HOLOTYPE: 7778, skin, juv., Funchal, Madeira (Portugal), U.S. Exploring Expedition, 1838.

***Xerobates agassizii* Cooper**

Proc. California Acad. Sci., vol. 2, p. 120, 1863.

=*Gopherus agassizii* (Cooper)

COTYPE: 7888, juv., Utah Basin, Mojave River (catalog carries "Solado Valley, California"), J. G. Cooper, March 1861.

Xerobates berlandieri Agassiz

Contributions to the natural history of the United States of America, vol. 1, p. 447, vol. 2, pl. 3, figs. 17-19, 1857.

=*Gopherus berlandieri* (Agassiz)

COTYPES (2): 60 (2), Lower Rio Grande (restricted to Brownsville, Texas), Texas, J. L. Berlandier.

Order Loricata

Caiman sclerops apaporiensis Medem

Chicago Nat. Hist. Mus., Fieldiana: Zool., vol. 37, p. 340, figs. 75-76, 1955.

PARATYPE: 134499, (formerly CNHM 69822), ♀, skull, Laguna del Capitan Mario, Río Apapoporis, Intendencia Amazonas, Colombia, F. Medem, January 1952.

Index

- abaxillare, Kinosternon, 231
abbotti, Chamaeleo, 96
 Coleonyx variegatus, 103
 Eleutherodactylus, 41
 Gonocephalus, 116
 Megophrys, 67
 Phelsuma, 128
 Phelsuma madagascariensis, 128
abingdonensis, Tropicidurus, 151
Ablepharus gloriosus, 81
Abronia aurita, 114
 taeniata graminea, 114
abyssus, Crotalus confluentus, 169
 Crotalus viridis, 169
acanthura, Ctenosaura, 105
Acrantophis madagascariensis madagascariensis, 205
Acris gryllus gryllus, 74
 gryllus paludicola, 29
Acrocytes modesta, 29
aculeodactylus, Platymanthis, 70
acuminata, Hyla, 50
acuta, Toluca lineata, 202
acutum, Kinosternon, 229
acutus, O[gmius], 202
adamanteus ruber, Crotalus, 168
Adelphicos quadrivirgatus visoninus, 211
 veraepacis nigrilatus, 157
adipoventris, Ptychohyla, 72
adspersa, Corythomantis, 39
aduncus, Bufo, 31
aeneus, Anolis roquet, 86
 bicanthalis, Sceloporus, 133
 chermocki, Desmognathus, 12
 Desmognathus, 11
 Desmognathus aeneus, 11
 subniger, Sceloporus, 134
Aepidia robusta, 157
aeruginosus, Leptophis, 192
aesopus, Rana areolata, 72
aestivus, Opheodrys, 193
aethiopissa elgonensis, Rhamnophis, 210
aethiops, Cnemidophorus tessellatus, 102
 Cnemidophorus tigris, 97, 98, 101, 102
affinis, Holbrookia, 116
 Hyla, 50, 51
 mayensis, Micrurus, 198
 stantoni, Micrurus, 198
Agalychnis helenae, 29
Agama agama elongis, 81
 agama turuensis, 81
 rueppelli occidentalis, 82
 rueppelli septentrionalis, 82
agama elongis, Agama, 81
 turuensis, Agama, 81
agassizii, Anolis, 85
 Gopherus, 235
 Mariguana, 85
 Nerodia, 201
 Platypeltis, 233
 Xerobates, 235
agilis, Salamandra, 24
Agkistrodon blomhoffii brevicaudus, 157
 contortrix pictigaster, 157
 mokeson pictigaster, 157
ahaetulla coeruleodorsa, Leptophis, 192
 copei, Leptophis, 193
 praestans, Leptophis, 219
Ahaetulla punctulata, 173
ahli, Montorana, 68
Aipysurus foliosquama, 157
alabamensis, Necturus, 16
 Pseudemys, 233
albemarlensis, Tropicidurus, 151
 Tropicidurus albemarlensis, 151
albiceps, Liophis, 209
 Rhadinaea, 209
albifrons ater, Leptotyphlops, 194
 magnamaculata, Leptotyphlops, 194
albipes, Eleutherodactylus, 41
albiventris, Sceloporus horridus, 136
albofrenata, Hyla, 50
albolabris, Tomodactylus, 80
albonuchalis, Scaphiodontophis, 212
albosignata, Hyla, 50
albotunica, Cochranella, 38
alboventer, Hypopachus, 62
albus, Scaphiopus, 78
 Scaphiopus holbrookii, 78
alfaroi, Gerrhonotus, 114
alleganiensis bishopi, Cryptobranchus, 11
alleni, Leimadophis, 190
 Leimadophis parvifrons, 190
 lineapiatus, Liodytes, 195
 Micrurus nigrocinctus, 198
alliacea, Mabuia, 125

- allisoni, *Anolis*, 85
 allogus, *Anolis*, 89
 alopex, *Sphaerodactylus*, 146
 Alsophis angulifer ruttyi, 157
 funereus, 157
 leucomelas, 175
 leucomelas danforthi, 157
 leucomelas sanctorum, 158
 nicholsi richardi, 158
 sanctorum, 158
 sibonius, 158
 vudii raineyi, 158
 altavelensis, *Leiocephalus*, 119
 Leiocephalus personatus, 119
 alternans, *Hypogeophis*, 3
 alticola, *Eleutherodactylus*, 41
 alticoronatus, *Laemanctus*, 119
 Altirana parkeri, 29
 alutaceus, *Anolis*, 85
 Anolis (Dracontura), 85
 alvarius, *Bufo*, 31
 amabilis, *Diadophis*, 173
 Diadophis amabilis, 173
 amamiensis, *Eumeces marginatus*, 111
 amarali, *Dryadophis*, 178
 Eudryas, 178
 amaura, *Lampropeltis*, 187
 Lampropeltis triangulum, 187, 203
 Ambystoma annulatum, 4
 aterrimum, 4
 cingulatum, 4
 copeianum, 5
 decorticatam, 5
 lepturum, 5
 macroductyla, 5
 obscurum, 5
 paroticum, 5
 platineum, 5
 proserpine, 5
 tenebrosam, 6
 trisruptum, 6
 xiphias, 6
 Ambystoma annulatum, 4
 cingulatum bishopi, 6
 cingulatum cingulatum, 4, 5
 fluvinatum, 6
 gracile decorticatam, 5
 gracile gracile, 5, 25
 granulosum, 6
 jeffersonianum, 5, 24
 lacustris, 6
 mabeci, 6
 macroductylum macroductylum, 5, 7
 mavortia, 7
 nebulosum, 7
 Ambystoma—Continued
 ordinaria, 7
 ordinarium, 7
 rosaceum nigrum, 7
 rosaceum sonoraensis, 7
 stejnegeri, 7
 subsalsum, 7
 tigrinum diaboli, 7
 tigrinum mavortium, 5, 6, 7, 25
 tigrinum melanostictum, 8
 tigrinum nebulosum, 7
 tigrinum slateri, 8
 tigrinum stebbinsi, 8
 tigrinum tigrinum, 5, 6, 24
 Ameiva ameiva fuliginosa, 82, 150
 atrigularis, 82
 birdorum, 82
 chrysoleama, 82
 chrysoleama chrysoleama, 82
 chrysoleama woodi, 82
 corvina, 82
 eutropia, 82
 exsul, 83
 festiva festiva, 82
 panchhora, 82
 plei exsul, 83
 polops, 83
 praesignis, 101
 quadrilineata, 84, 101
 tobagana, 84
 undulata amphigramma, 83
 undulata dextra, 83
 undulata hartwegi, 83
 undulata podarga, 83
 undulata sinistra, 83
 undulata stuarti, 83
 undulata thomasi, 84
 wetmorei, 84
 wetmorei eleanorae, 84
 wetmorei wetmorei, 84
 ameiva fuliginosa, *Ameiva*, 82, 150
 amethystinus kinghorni, *Liasis*, 195
 Amiva gabbiana, 84
 surinamensis tobaganus, 84
 amoenus helenae, *Carphophis*, 163
 Amphibia, 3
 Amphiesma flavipunctatum, 158
 amphigramma, *Ameiva undulata*, 83
 Amphisbaena bakeri, 84
 camura, 84
 fenestrata, 108
 manni, 84
 pericensis, 85
 townsendi, 85

amphistieta, Toluca, 219
Amplorhinus multimaculatus, 174
Amyda spinifera hartwegi, 227
Anadia metallica metallica, 96
Anaides ferreus, 8
anceps, Hyla, 50
Ancistrodon piscivorus leucostoma, 220
ancorus, *Holarchus*, 226
 Xenodon, 226
andersonii, Hyla, 51
andreae nebulatus, *Leimadophis*, 190
andrewsi, *Eleutherodactylus*, 41
androsi, *Tropidophis pardalis*, 223
Aneides ferreus, 8
 lugubris lugubris, 19
angulifer ruttyi, *Alsophis*, 157
angustatus, *Claudius*, 230
angusticeps, *Cnemidophorus*, 96
 Cnemidophorus sackii, 96
angusticlavius, *Plethodon cinereus*, 19
 Plethodon dorsalis, 19
angustidigitum, *Tomodactylus*, 81
angustilineata, Hyla, 51
angustipes, *Bufo*, 31
angustirostris, *Eutaenia*, 178
 Thamnophis, 178
angustus, *Sator*, 133
annamensis, *Microhyla*, 67
 Oligodon, 202
annectens, *Pituophis*, 206
 Pituophis catenifer, 206
annulata, *Boa*, 226
 Chionactis occipitalis, 190
 Lampropeltis, 187
 Lampropeltis triangulum, 187
 pulchriceps, *Leptodeira*, 191
 septentrionalis, *Leptodeira*, 174
 taylori, *Leptodeira*, 191
annulatum, *Amblystoma*, 4
 Amblystoma, 4
 Lamprosoma, 190
 Xiphosoma, 226
 yucatanensis, *Sibon*, 213
annulatus, *Sceloporus merriami*, 139
anocularis, *Geophis*, 183
Anolis agassizii, 85
 allisoni, 85
 allogus, 89
 alutaceus, 85
 (*Dracontura*) *alutaceus*, 85
 asper, 85
 aureolus, 85
 bimaculatus lividus, 89
 bimaculatus sabanus, 91
 cepedii cinereus, 85

Anolis—Continued

cepedii gentilis, 86
 compressicauda, 86
 concolor, 86, 90
 (*Gastrotopis*) *concolor*, 86
 conspersus, 86
 conspersus lewisi, 86
 crisatellus wileyi, 86
 cupreus, 86
 cyanopleurus, 86
 (*Dracontura*) *cyanopleurus*, 86
 cybotes doris, 87
 darlingtoni, 86
 distichoides, 87
 distichus caudalis, 87
 distichus floridanus, 87
 distichus juliae, 87
 distichus wetmorei, 87
 dominicensis caudalis, 87
 dominicensis juliae, 87
 dominicensis wetmorei, 87
 doris, 87
 evermanni, 87
 fairchildi, 88
 ferreus, 85
 gingivinus, 92
 grahami lewisi, 86
 greyi, 88
 griseus, 88
 hendersoni, 88
 humilis humilis, 90
 humilis uniformis, 90, 92
 laeviventris, 90
 leucophaeus marignanae, 88
 lineatopus coxi, 88
 lineatopus lynni, 89
 lionotus, 90
 lividus, 89
 luciae, 89
 matudai, 89
 mestrei, 89
 microtus, 89
 milleri, 89
 monensis, 89
 nannodes, 90
 nelsoni, 90
 oculatus, 156
 oxylophus, 90
 pachypus, 90
 pinchoti, 90
 poncensis, 90
 purpureus, 90
 quaggulus, 90
 richardii griseus, 88

Anolis—Continued

- richardii richardii, 92
 roquet aeneus, 86
 roquet cinereus, 85, 91
 ruthveni, 90
 sabanus, 91
 sagrei mayensis, 91
 sagrei stejnegeri, 91
 stejnegeri, 91
 subocularis, 91
 taylori, 91
 townsendi, 91
 trinitatis extremus, 91
 trinitatis luciae, 89
 trinitatis vincentii, 92
 tripidonotus spilorhipis, 91
 trossulus, 92
 uniformis, 92
 vincentii, 92
 virgatus, 92
- Anomalepis aspinosus, 158
 anomalus, Oreosaurus, 127
 anonymorpha, Uta, 152
 anonymorphus, Urosaurus bicarinatus, 152
 Anota calidiarum, 92
 Anotheca coronata, 49
 antauges, Barisia, 93
 Barissia, 93
 anthonyi, Bascanion, 160
 Masticophis, 160
 anthracinus, Eumeces anthracinus, 131
 Plestiodon, 131
 anthracinus pluvialis, Eumeces, 112
 anulatus, Phrynixalus, 69
 apaporiensis, Caiman sclerops, 236
 Aparasphenodon brunoï, 39
 apiata, Stenorhina degenhardtii, 214
 apicalis, Sceloporus spinosus, 143
 Aporophis flavifrenatus, 196
 juliae, 158
 appendiculata, Phyllomedusa, 70
 aquarius, Leiocephalus carinatus, 119
 arborea, Bolitoglossa, 9
 Chiropterotriton, 9
 arboricola, Hyla, 51
 arenarius, Leiocephalus, 121
 arenicola, Stilosoma extenuatum, 214
 arenicolor, Hyla, 50, 51
 areolata aesopus, Rana, 72
 capito, Rana, 73
 Engystoma, 48
 Rana, 72
 Rana areolata, 72
 sevosa, Rana, 76
- argillaceus, Bufo, 31
 argus bartschi, Sphaerodactylus, 146
 Dipsas, 192
 henriquesi, Sphaerodactylus, 146
 Leptognathus, 192
 Aristelligella expectata, 92
 Aristelliger expectatus, 92
 georgeensis, 92
 irregularis, 92
 Arizona elegans, 158
 elegans blanchardi, 159
 elegans elegans, 158
 elegans expolita, 159
 elegans noctivaga, 159
 elegans occidentalis, 159
 jani, 159
 arizonae, Diadophis regalis, 173
 Xantusia, 156
 armouri, Audantia, 92
 Leiocephalus carinatus, 119
 arnoensis, Emoia, 109
 arnyi, Diadophis, 173
 Diadophis punctatus, 173
 Arrhyton bivittatum, 159
 fulvum, 160
 taeniatum, 160, 164
 vittatum, 159
 Ascaphus truei, 29
 truei californicus, 29
 truei montanus, 29
 truei truei, 29
 asio, Phrynosoma, 128
 asper, Anolis, 85
 Aspidonectes, 227
 Trionyx spinifer, 227, 233
 Aspidonectes asper, 227
 emoryi, 227
 aspinosus, Anomalepis, 158
 assatum ixbaac, Lygosoma, 125
 Atelopus cruciger vogli, 30
 elegans, 30
 gracilis, 30
 moreirae massarti, 30
 spurrelli certus, 30
 varius certus, 30
 varius glyphus, 30
 ater, Leptotyphlops, 194
 Leptotyphlops albifrons, 194
 Sauromalus, 133
 aterrimum, Amblystoma, 4
 atkinsi orientalis, Eleutherodactylus, 41
 atlanticus, Scoliophis, 212
 atmodes, Heterodon, 184
 Atractus crassicaudatus, 208

- atrata*, *Eutainia*, 180, 181
atratus, *Thamnophis elegans*, 180
atricucularis, *Gonatodes*, 115
atrigularis, *Ameiva*, 82
 Sphenomorphus, 148
atrox, *Crotalus*, 169
 elegans, *Crotalus*, 169
attenuatus, *Batrachoseps attentuatus*, 8
 Drepanodon, 174
 longicaudus, *Ophisaurus*, 127
aubryi, *Hylambates*, 58
audanti, *Eleutherodactylus*, 41
Audantia armouri, 92
 shrevei, 93
Aulacoplax leptosoma, 93
aurantiaca orophila, *Hyla*, 56
 planicola, *Hyla*, 57
 surda, *Hyla*, 51
auratus, *Dendrobates*, 69
 Phyllobates, 69
aurea, *Hyla*, 77
aureatagulus, *Desmognathus*, 11
aureolus, *Anolis*, 85
 Sceloporus mucronatus, 140
aureus, *Leiocephalus personatus*, 120
auribundus, *Spilotes pullatus*, 214
auriculata, *Uta*, 152
auriculatus, *Urosaurus*, 152
aurigulus, *Drymobius*, 176
 Masticophis, 176
aurita, *Abronia*, 114
auritus, *Bufo*, 31, 33
 Gerrhonotus, 114
aurora cascadae, *Rana*, 73
 draytonii, *Rana*, 73, 74, 76
 Rana, 72
 Rana aurora, 72
australis, *Masticophis taeniatus*, 197
avivoca, *Hyla*, 51
avocalis, *Eleutherodactylus*, 41
axanthus, *Pseudobranchius striatus*, 22
azurea, *Phyllomedusa*, 70
Bachia intermedia, 93
 parkeri, 93
badius, *Celestus*, 95
baileyi, *Clelia*, 164
 Crotaphytus, 104
 Crotaphytus collaris, 104
 Hyla goughi, 54
 Oxyrhopus, 164
bairdi, *Coluber*, 164
 Elaphe obsoleta, 164
bakeri, *Amphisbaena*, 84
 Ctenosaura, 105
 Eleutherodactylus, 42
 Enyaliosaurus, 105
bakewelli, *Leptotyphlops*, 194
 Leptotyphlops phenops, 194
balteata, *Opisthotropis*, 204
baluensis, *Rhacophorus*, 77
bambusicola, *Rhacophorus*, 77
barbouri, *Basiliscus*, 93
 Bothrops, 187
 Lachesis, 187
 Leiopisma, 121
 Lepidoblepharis, 122
 Tropidophis pardalis, 224
Barbourina equatoriana, 160
Barisia antauges, 93
 gadovii levigata, 93
 imbricata ciliaris, 114
 imbricata imbricata, 115
 levicollis, 93
 modesta, 132
 monticola, 114, 115
Barissia antauges, 93
 levicollis, 93
bartolomas, *Cnemidophorus*, 96
bartschi, *Cyclura carinata*, 106
 Deiroptyx, 107
 Sphaerodactylus, 146
 Sphaerodactylus argus, 146
Basanitia bolbodactyla, 48
Bascanion anthonyi, 160
 flagellum frenatum, 160
 foxi, 160
 suboculare, 160
 vetustus, 160
Basianium piceum, 160
 semilineatum, 161
basilica, *Caudisona*, 162
Basiliscus barbouri, 93
 basiliscus, 93
 guttulatus, 93
 (*Cristasaura*) *nuchalis*, 94
 plumifrons, 94
 vittatus, 94, 107
basiliscus, *Basiliscus*, 93
 Crotalus basiliscus, 162
 oaxacus, *Crotalus*, 169
basimaculatus, *Typhlops*, 224
batangensis, *Rhacophorus pleurostictus*, 77
Bathysiredon dumerilii, 24

- Batrachoseps attenuatus attenuatus*, 8
 catalinae, 8
 caudatus, 8
 leucopus, 8
 pacificus, 8, 15
 wrighti, 22
Batrachuperus karlschmidti, 8
 pinchonii, 24
 yenyanensis, 8
Batrachylodes trossulus, 31
battersbyi, *Philothamnus irregularis*, 205
baudinii, *Smilisca baudinii*, 56, 60
baueri, *Hypsiglena torquata*, 186
bauri palmarum, *Kinosternon*, 231
 Terrapene, 235
 Terrapene carolina, 235
bayonii keniensis, *Mabuia*, 126
beebei, *Eleutherodactylus*, 42
beldingi, *Bufo*, 31
 Cnemidophorus hyperythrus, 156
 Verticaria, 156
bellii, *Chrysemys picta*, 229
bellona, *Churchillia*, 164
bequaerti, *Leptopelis*, 65
berdmorei, *Microhyla*, 67
berendtianum, *Cinosternum*, 229
berlandieri, *Gopherus*, 236
 Rana, 72
 Rana pipiens, 72
 Xerobates, 236
bernieri, *Dromicodryas*, 184
 Herpetodryas, 184
beyeri, *Necturus*, 16
 Necturus maculosus, 16
bibroni, *Pleurodema*, 68
bicanthalis, *Sceloporus aeneus*, 133
bicarinatus anonymorphus, *Urosaurus*, 152
 nelsoni, *Urosaurus*, 154
 spinosus, *Urosaurus*, 152
bicolor, *Pliocercus*, 208
bifossatus, *Dryadophis bifossatus*, 168
bilineatus borealis, *Spelerpes*, 25
 Eumeces brevirostris, 110
 Masticophis, 161
bimaculata, *Tantilla*, 216
bimaculatus lividus, *Anolis*, 89
 sabanus, *Anolis*, 91
bimaculosus, *Sceloporus magister*, 137
himaris, *Pituophis catenifer*, 206
hinghami, *Eleutherodactylus*, 42
 Tretanorhinus variabilis, 220
Bipes biporus, 110
 Euchirotes, 110
bipunctata branneri, *Hyla*, 51
bipunctatus biseriatus, *Coniophanes*, 166
birdorum, *Ameiva*, 82
bischoffi multilineata, *Hyla*, 56
biscutatus quadruplex, *Trimorphodon*, 221
 semirutus, *Trimorphodon*, 221
 Thamnophis elegans, 179
 Trimorphodon biscutatus, 222
bisectus, *Tropidonotus*, 222
biserialis, *Conopsis*, 167
biseriatus, *Coniophanes bipunctatus*, 166
bishopi, *Ambystoma cingulatum*, 6
 Cryptobranchus, 11
 Cryptobranchus alleganiensis, 11
bislineata, *Eurycea bislineata*, 13, 25
 major, *Eurycea*, 13
 rivicola, *Eurycea*, 14
bissa, *Eretmochelys imbricata*, 228
bistincta, *Hyla*, 51
 Hyla bistincta, 51
 laberculata, *Hyla*, 52
bivittata, *Daconura*, 107
 Natrix compressicauda, 199
bivittatum, *Arrhyton*, 159
bivittatus, *Pantodactylus*, 128
 Tropidurus, 152
blanchardi, *Arizona elegans*, 159
 Diadophis regalis, 173
 Geophis, 184
 Natrix rhombifera, 200
 Opheodrys vernalis, 202
 Pseudophryne, 71
blasii, *Triton*, 27
blomhoffii brevicaudus, *Agkistrodon*, 157
Boa annulata, 226
boettgeri camerunensis, *Hymenochirus*, 61
 Leiopisma laterale, 122
 orientalis, *Emoia*, 109
bogerti, *Coleonyx variegatus*, 103
 Salvadora, 211
Boiga dendrophila occidentalis, 161
holbodactyla, *Basanitia*, 48
 Eupemphix, 48
 Pseudis, 71
Bolitoglossa arborea, 9
 borburata, 9
 cephalica rubrimembris, 9
 chondrostega, 9
 cochranae, 9
 dimidiata, 9
 flaviventris, 17
 galaenae, 9
 helmrichi, 18
 laevae, 10
 melanomolga, 10

- Bolitoglossa*—Continued
 moreleti, 10
 nigroflavescens, 10
 nigromaculata, 10
 occidentalis, 10
 odonnelli, 18
 striatula, 19
 terrestris, 11
 unguidentis, 11
 xolocalcae, 11
boliviana, *Gastrotheca*, 42
bolivianus, *Leptodactylus*, 64
Borborocoetes stejegeri, 31
borburata, *Bolitoglossa*, 9
borealis, *Spelerpes bilineatus*, 25
boreas, *Bufo*, 32
 Bufo boreas, 32, 35, 36
 exsul, *Bufo*, 33
 halophilus, *Bufo*, 34
 nelsoni, *Bufo*, 32
Bothrops barbouri, 187
 diporus, 161
 insularis, 187
 lansbergii venezuelensis, 161
 leptura, 161
 melanurus, 220
 neuwiedii diporus, 161
 nummifer veraecrucis, 161
 picadoi, 220
 yucatanicus, 221
bottae, *Charina bottae*, 163, 226
boulengeri, *Cyclorhamphus*, 40
 Hyla, 78
 Phyllobates, 71
 Prostherapis, 71
 Sceloporus, 134
 Sceloporus clarkii, 134
 Scytopsis, 78
boyllii conjuncta, *Lampropeltis*, 187
 Ophibolus, 203
 Rana, 72
 Rana boyllii, 72
brachycephala, *Rana pipiens*, 76, 77
 Rana virescens, 77
brachycercum, *Phrynosoma douglassii*, 128
brachylopha, *Ctenosaura teres*, 106
brachyops, *Charina*, 163
branneri, *Hyla bipunctata*, 51
bransfordii, *Lithodytes*, 65
 Microbatrachylus, 65
brevicaudus, *Agkistrodon blomhoffii*, 157
brevifacies, *Dipsas*, 222
 Tropidodipsas, 222
brevigulare, *Cinosternum*, 229
brevilineatus, *Eumeces*, 110
brevipes, *Eumeces skiltonianus*, 113
brevirostris bilineatus, *Eumeces*, 110
 Ctenosaurua, 105
 Dryophis, 176
 Oxybelis, 176
 Phrynosoma douglassii, 150
 Tapaya, 150
brevis, *Coleonyx*, 103
 Coleonyx variegatus, 103
brimleyorum, *Desmognathus*, 11
 Desmognathus fuscus, 11
bromeliacia, *Chiropterotriton*, 17
 Oedipus, 17
brongersmai, *Python curtus*, 208
browni, *Heterodon*, 185
 Phyllorhynchus, 206
 Phyllorhynchus browni, 206
brunnata, *Pseudoeurycea*, 23
brunnea, *Eutaenia elegans*, 179
 Hyla septentrionalis, 81
brunneus, *Phyllobates*, 49
brunoi, *Aparasphenodon*, 39
bucculenta, *Ungalia*, 225
bucculentus, *Tropidophis pardalis*, 225
Bufo aduncus, 31
 alvarius, 31
 angustipes, 31
 argillaceus, 31
 auritus, 31, 33
 beldingi, 31
 boreas, 32
 boreas boreas, 32, 35, 36
 boreas exsul, 33
 boreas halophilus, 34
 boreas nelsoni, 32
 canaliferus, 32
 coccifer, 32
 columbiensis, 32
 compactilis speciosus, 37
 coniferus, 32
 copei, 32
 debilis, 32
 debilis debilis, 32
 debilis insidior, 34
 debilis kelloggi, 35
 dorsalis, 33, 37
 dunni, 33
 exsul, 33
 fastidiosus, 39
 gabbi, 31, 33
 gemmifer, 33

Bufo—Continued

granulosus, 33, 36
güntheri, 33
haematiticus, 34
haematiticus lachrymans, 34
halophila, 34
hemiophrys, 34
houstonensis, 34
ictericus mertensi, 34
inca, 34
insidiator, 34
kelloggi, 35
lamentor, 35
lentiginosus pachycephalus, 35
longinasus, 35
lugubrosus, 35
marmoreus, 31
mazatlanensis nayaritensis, 36
melanochloris, 35
microscaphus, 35
minshanicus, 35
monksiae, 36
nayaritensis, 36
nebulifer, 33, 36
occidentalis, 36
paracnemis, 36
perplexus, 36
pictus, 36
politus, 36
punctatus, 31, 37
pygmaeus, 37
rubescens, 37
rufus, 37
smithi, 37
speciosus, 37
spinulosus, 35
tacanensis, 37
terrestris charlesmithi, 37
terrestris copei, 32
terrestris terrestris, 35
valliceps, 33, 36
woodhousei hemiophrys, 34
woodhousei woodhousei, 31, 33
woodhousii, 37
woodhousii velatus, 38
woodhousii woodhousii, 38
burti, *Cnemidophorus*, 97
butleri, *Microhyla*, 67, 68
Cachryx defensor, 94
cadaverina, *Hyla*, 52, 56
Cadea palirostrata, 94
caducus, *Leiocephalus*, 133
Scartiscus, 133

Caecilia isthmica, 3
ochrocephala, 3
tentaculata, 3
caerulea, *Uta*, 153
Verticaria, 156
caeruleus, *Cnemidophorus hyperythrus*, 156
Sceloporus ornatus, 141
Urosaurus ornatus, 153
caesius, *Opheomorphus doliatus*, 202
cahuilae, *Leptotyphlops humilis*, 194
caicosensis, *Sphaerodactylus*, 146
Caiman sclerops apaporiensis, 236
Calamaria loyii ventralis, 162
lumholtzii, 161
mearnsi, 161
raveni, 161
schmidti, 161
tenuis, 162, 168
ventralis, 162
calamarina, *Tantilla*, 216
calidiarum, *Anota*, 92
Phrynosoma platyrhinos, 92
californiae, *Lampropeltis getulus*, 203, 204
californicus, *Ascaphus truei*, 29
caliginis, *Crotalus viridis*, 172
caliginosus, *Leptodactylus*, 63
calligaster, *Contia*, 168
Lampropeltis calligaster, 203
Rhadinaea, 168
Sceloporus melanorhinus, 139
Scotophis, 166
Callirhinus patagoniensis, 162
Callisaurus carmenensis, 94
crinitus, 94
draconoides carmenensis, 94
draconoides crinitus, 94
draconoides inusitatus, 94
draconoides myurus, 95
draconoides rhodostictus, 95
draconoides splendidus, 95
draconoides ventralis, 118
inusitatus, 94
plasticus, 94
rhodostictus, 95
splendidus, 95
ventralis myurus, 95
Callophis hughi, 162
camerunensis, *Hymenochirus boettgeri*, 61
campi, *Syrhrophus*, 79
camura, *Amphisbaena*, 84
cana, *Ficimia*, 184
Ungalia, 225
canaliferus, *Bufo*, 32

- cancellatus, *Geophis*, 184
 canescens, *Thamnophis melanogaster*, 218
 canilatus, *Tachymenis*, 216
 cantabrigensis evittata, *Rana*, 72
 latiremis, *Rana*, 72
 Rana sylvatica, 72
 Cantoria elapiformis, 162
 violacea, 162
 canula, *Tantilla*, 216
 canum, *Gyalopion*, 184
 canus, *Tropidophis pardalis*, 225
 capensis mossambica, *Lygodactylus*, 124
 capito, *Rana*, 73
 Rana areolata, 73
 stertens, *Rana*, 73
 caprimimus, *Hypopachus*, 62
 cárdenasi, *Hyla*, 52
 caretta, *Caretta caretta*, 235
 Caretta caretta caretta, 235
 remivaga, 227
 rostrata, 228
 carinata bartschi, *Cyclura*, 106
 carinatus aquarius, *Leiocephalus*, 119
 armouri, *Leiocephalus*, 119
 granti, *Leiocephalus*, 119
 punctatus, *Leiocephalus*, 119
 carmenensis, *Callisaurus*, 94
 Callisaurus draconoides, 94
 carolina bauri, *Terrapene*, 235
 triunguis, *Terrapene*, 230
 carolinensis, *Desmognathus ochrophaea*, 12
 Desmognathus ochrophaeus, 11, 12
 Carphophis amoenus helenae, 163
 carpincinctus, *Scaphiodontophis*, 212
 carvalhoi, *Hemipipa*, 71
 Protopipa, 71
 cascadeae, *Rana*, 73
 Rana aurora, 73
 casparii, *Eleutherodactylus*, 42
 Eleutherodactylus ricordii, 42
 catalinae, *Batrachoseps*, 8
 catenatus edwardsii, *Sistrurus*, 168
 catenifer annectens, *Pituophis*, 206
 bimaris, *Pituophis*, 206
 deserticola, *Pituophis*, 206
 fuliginatus, *Pituophis*, 207
 insulanus, *Pituophis*, 207
 Pituophis melanoleucus, 207
 sayi, *Pituophis*, 164
 Catostoma nasale, 162
 psephotum, 162
 caudaequinae, *Leiopisma*, 121
 caudalis, *Anolis*, *distichus*, 87
 Anolis dominicensis, 87
 Caudata, 4
 caudatus, *Batrachoseps*, 8
 Caudisona basilisca, 162
 enyo, 162
 pyrrha, 163
 Causus nasalis, 163
 resimus, 163
 cavernicola, *Eleutherodactylus*, 42
 celaeno, *Tropidonotus*, 222
 celaenops, *Lampropeltis pyrrhomelaena*, 189
 Lampropeltis triangulum, 189
 celeripes, *Cnemidophorus*, 97
 Cnemidophorus tigris, 97
 celeris, *Salvadora hexalepis*, 211
 Celestus badius, 95
 cruscus molesworthi, 95
 curtissi, 95
 darlingtoni, 95
 enneagrammus, 108
 rozellae, 95
 rugosus, 96
 weinlandii, 96
 Celuta helenae, 163
 cenchoa leucomelas, *Imantodes*, 185
 cenchria crassus, *Epicrates*, 178
 centralis, *Coluber (Masticophis) mento-*
 varius, 165
 Centrolenella viridissima, 38
 cepedii cinereus, *Anolis*, 85
 gentilis, *Anolis*, 86
 cephalica, *Pseudoeurycea cephalica*, 26
 rubrimembris, *Bolitoglossa*, 9
 rubrimembris, *Pseudoeurycea*, 9
 cerasinus, *Eleutherodactylus*, 61
 Hylodes, 61
 cerastes cercobombus, *Crotalus*, 169
 Crotalus, 169
 Crotalus cerastes, 169
 Cerathyla panamensis, 38
 Ceratophrys fusceiventris, 38
 ceratophrys, *Hyla*, 52
 cercobombus, *Crotalus cerastes*, 169
 cerebrosus, *Thamnophis sumichrasti*, 219
 cerroense, *Phrynosoma*, 128
 certus, *Atelopus spurrelli*, 30
 Atelopus varius, 30
 chalceus, *Thamnophis sauritus*, 182
 Chalcidolepis metallicus, 96
 chalybaeus, *Diploglossus*, 108
 chalybeus, *Geophis*, 209
 Chamaeleo abbotti, 96
 johnstonii, 96
 taitensis, 96
 Chamaeleon ituriensis, 96

- Chamaelinorops wetmorei*, 96
chamissonis, *Leimadophis*, 216
championi, *Hypopachus*, 62
chanleri, *Mabuia*, 126
 Mehelya chanleri, 213
 Simocephalus, 213
chaochiaensis, *Rana*, 73
Charina bottae bottae, 163, 226
 brachyops, 163
charlesmithi, *Bufo terrestris*, 37
Chelonia formosa, 228
 mydas japonica, 228
 tenuis, 228
Chelopus funereus, 228
 gabbii, 228
 rubidus, 228
Chelydra osceola, 228
 serpentina osceola, 228
chenfui, *Rhacophorus*, 77
chermocki, *Desmognathus*, 12
 Desmognathus aeneus, 12
cherokiana, *Rana sylvatica*, 76
cherriei ixbaac, *Scincella*, 125
 stuarti, *Lygosoma*, 125
 stuarti, *Scincella*, 125
Chilomeniscus cinctus, 163
 ephippicus, 163
 stramineus, 163
 stramineus fasciatus, 163
 stramineus stramineus, 163
Chilopoma rufipunctatum, 164
Chionactis diasii, 164
 occipitalis annulata, 190
Chironius fuscus, 206
 melas, 184
Chiropterotriton arborea, 9
 bromeliacia, 17
 chondrostega, 9
 dimidiata, 9
 lavae, 10
 multidentata, 18
 nasalis, 18
 picadoi, 26
 terrestris, 11
 xolocalcae, 11
chloris, *Herpetodryas*, 184
chlorobryonis, *Plethodon glutinosus*, 19
chlorophaea, *Hypsigena*, 186
Chlorosoma elegans, 216
 olfersii, 205
chloroticum, *Dendrophidium*, 173
chloroticus, *Drymobius*, 173
chondrostega, *Bolitoglossa*, 9
 Chiropterotriton, 9
Chrysemys dorsalis, 229
 picta bellii, 229
 picta dorsalis, 229
 treleasei, 229
chrysobronchus, *Pseustes poecilonotus*, 213
 Spilotes, 213
chrysocephala, *Eutaenia*, 178
chrysocephalus, *Thamnophis*, 178
chrysoleama, *Ameiva*, 82
 chrysoleama, *Ameiva*, 82
 woodi, *Ameiva*, 82
chrysostictus, *Sceloporus*, 134
Churchillia bellona, 164
ciliaris, *Barisia imbricata*, 114
 Gerrhonotus levicollis, 114
cinctus, *Chilomeniscus*, 163
cinerea evittata, *Hyla*, 53
cinereus angusticlavius, *Plethodon*, 19
 Anolis cepedii, 85
 Anolis roquet, 85, 91
 dorsalis, *Plethodon*, 19
 Plethodon cinereus, 20, 24
 Sphaerodactylus, 146
cingulatum, *Amblystoma*, 4
 Ambystoma cingulatum, 4, 5
 bishopi, *Ambystoma*, 6
Cinosternum berendtianum, 229
 brevigulare, 229
 postinguinale, 229
Cistudo ornata, 229
 triunguis, 230
clarionensis, *Urosaurus*, 153
 Uta, 153
clarki, *Micrurus*, 198
clarkii boulengeri, *Sceloporus*, 134
 Ctenosaura, 105
 Enyaliosaurus, 105
 Helocaetes, 50
 Natrix sipedon, 208
 Pseudacris, 50
 Regina, 208
 Sceloporus, 134
 Sceloporus clarkii, 134
Claudius angustatus, 230
 severus, 230
Clelia baileyi, 164
 clelia clelia, 212
 clelia immaculata, 164
 clelia scytalina, 160
 trigemina, 205
clelia, *Clelia clelia*, 212
 immaculata, *Clelia*, 164
 scytalina, *Clelia*, 160

- clemsonae*, *Plethodon*, 19
Plethodon jordani, 19
Clemmys marmorata marmorata, 230
muhlenbergii, 230
nuchalis, 230
clericus, *Ophibolus*, 203
Cnemidophorus angusticeps, 96
bartolomas, 96
burti, 97
celeripes, 97
communis, 97
communis communis, 97
costatus, 97
decemlineatus, 97
deppii deppii, 97, 100
deppii lineatissimus, 97, 100
disparilis, 97
estebanensis, 97
gadovi, 98
gigas, 98
gracilis, 98
grahamii, 98
gularis, 98
gularis gularis obsoletus, 98
gularis gularis verus, 98
gularis scalaris, 98
gularis semifasciatus, 99
gularis sericeus, 99
guttatus guttatus, 103
guttatus immutabilis, 99, 100
hyperythrus, 99
hyperythrus heldingi, 156
hyperythrus caeruleus, 156
hyperythrus hyperythrus, 99
immutabilis, 99
inornatus, 99, 101
labialis, 99
lativittis, 100
lineatissimus, 100
marmoratus, 100
martyris, 100
maximus, 100
melanostethus, 100
microlepidopus, 100
neomexicanus, 100
octolineatus, 101
perplexus, 101
praesignis, 101
punctilinealis, 101
quadrilineatus, 101
sacki huico, 101
sackii angusticeps, 96
sackii gularis, 98, 99
sackii huico, 101
Cnemidophorus—Continued
sackii sackii, 97
sackii scalaris, 98
sackii semifasciatus, 99, 101
sackii stictogrammus, 101
septemvittatus, 101
sexlineatus, 206
sexlineatus oligoporus, 102
sexlineatus pauciporus, 102
tessellatus, 98
tessellatus aethiops, 102
tessellatus multiscutatus, 102
tessellatus rubidus, 102
tigris, 102
tigris aethiops, 97, 98, 101, 102
tigris celeripes, 97
tigris estebanensis, 97
tigris gracilis, 98, 100, 102
tigris marmoratus, 100
tigris martyris, 100
tigris multiscutatus, 96, 102
tigris rubidus, 102, 103
tigris tigris, 102
tigris variolosus, 103
undulatus, 102
unicolor, 103
vandenburghi, 103
variolosus, 103
coccifer, *Bufo*, 32
cochranae, *Bolitoglossa*, 9
Hylambates, 61
Pseudoeurycea, 9
Sceloporus, 134
Cochranella albotunica, 38
delicatissima, 38
divaricans, 38
dubia, 38
lutzorum, 39
petropolitana, 39
surda, 39
coeruleodorsa, *Leptophis ahaetulla*, 192
coeruleodorsus, *Leptophis*, 192
coeruleus palmeri, *Gerrhonotus*, 115
principis, *Gerrhonotus*, 109
cognatus, *Heterodon*, 185
Coleonyx brevis, 103
variegatus abbotti, 103
variegatus bogerti, 103
variegatus brevis, 103
variegatus peninsularis, 103
variegatus sonoriensis, 103
variegatus variegatus, 149
collaris baileyi, *Crotaphytus*, 104
Oedipina, 25

- collaris*—Continued
Ophibolus doliatus, 203
Spelerpes, 25
Trimorphodon, 221
Colorhogia redimita, 164
Coluber bairdi, 164
constrictor constrictor, 212
constrictor foxii, 160
constrictor mormon, 160, 165
constrictor paludicola, 165
constrictor stejnegerianus, 226, 227
erythrogaster, 165
flavirufus, 165
guttatus sellatus, 165
(Masticophis) *mentovarius centralis*, 165
mormon, 165
mutabilis, 165
obsoletus lemniscatus, 166
rhinomegas, 166
rosaceus, 166
triaspis, 166
columbiensis, *Bufo*, 32
colymba, *Hyla*, 52
communis, *Cnemidophorus*, 97
Cnemidophorus communis, 97
compactilis speciosus, *Bufo*, 37
completa, *Ctenosaura*, 105
compressicauda, *Anolis*, 86
bivittata, *Natrix*, 199
Natrix sipedon, 199, 201, 223
Nerodia, 201
compressicaudus walkeri, *Tropidonotus*, 223
compressus, *Ophisaurus*, 127
Ophisaurus ventralis, 127
concinna, *Pseudemys floridana*, 234
Uta, 153
concolor, *Anolis*, 86, 90
Anolis (Castrotropis), 86
Conophis, 167
Conophis lineatus, 167
Rana montezumae, 75
confinis, *Scotophis*, 212
confluentus abyssus, *Crotalus*, 169
lutosus, *Crotalus*, 169
stephensi, *Crotalus*, 169
conica, *Toluca*, 219
coniferus, *Bufo*, 32
Coniophanes bipunctatus biseriatus, 166
fissidens proterops, 166
imperialis imperialis, 216
piceivittis, 166
Coniophanes—Continued
piceivittis piceivittis, 166
proterops, 166
quinquevittatus, 186
conirostris, *Zamenis*, 226
conjuncta, *Lampropeltis boylii*, 187
Lampropeltis getulus, 187
stickeli, *Kaloula*, 63
Conophis concolor, 167
lineatus concolor, 167
lineatus dunni, 167
lineatus similis, 167
pulcher, 167
pulcher plagosus, 167
pulcher pulcher, 167
pulcher similis, 167
sumichrastii, 167
sumichrastii viduus, 167
vittatus vittatus, 167
Conopsis biserialis, 167
consors, *Kinosternon cruentatum*, 232
conspersus, *Anolis*, 86
Iewisi, *Anolis*, 86
conspicua, *Ctenosaura*, 105
conspicuus, *Eleutherodactylus*, 42
constrictor, *Coluber constrictor*, 212
foxii, *Coluber*, 160
mormon, *Coluber*, 160, 165
paludicola, *Coluber*, 165
sigma, *Constrictor*, 167
stejnegerianus, *Coluber*, 226, 227
Constrictor constrictor sigma, 167
Contia calligaster, 168
isozona, 168
mitis, 162, 168
pachyura, 168
tenuis, 162, 168
contortrix pictigaster, *Agkistrodon*, 157
copei, *Bufo*, 32
Bufo terrestris, 32
Crotaphytus, 104
Eumeces, 111
Leptophis, 193
Leptophis ahaetulla, 193
copeianum, *Amblystoma*, 5
corais erebennus, *Drymarchon*, 213
melanurus, *Drymarchon*, 213
rubidus, *Drymarchon*, 175
unicolor, *Drymarchon*, 176
Cornufer heros, 49
vitiensis, 49
worcesteri, 39
cornutum, *Phrynosoma*, 129

- coronata*, *Anotheca*, 49
 Gastrotheca, 49
 Tantilla, 217
 Tantilla coronata, 217
coronatum frontale, *Phrynosoma*, 129
 jamesi, *Phrynosoma*, 129
corticata, *Thalassochelys*, 235
corticulus, *Sphaerodactylus*, 146
corvina, *Ameiva*, 82
Coryphodon pantherinus, 168
Corythomantis adpersa, 39
costatus, *Cnemidophorus*, 97
couchii, *Eutaenia*, 178
 Nerodia, 201
 Scaphiopus, 78
 Sceloporus, 125, 134
 Thamnophis elegans, 178
coxi, *Anolis lineatopus*, 88
cozumelae, *Sceloporus*, 134
Cranopsis fastidiosus, 39
Craspedoglossa sanctae-catharinae, 39
 stejnegeri, 31
crassa, *Rhadinaea*, 209
crassicaudatum, *Rabdosome*, 208
crassicaudatus, *Atractus*, 208
 Stenocercus, 149
crassulus, *Plethodon*, 19
crassus, *Epicrates*, 178
 Epicrates cenchria, 178
crinitus, *Callisaurus*, 94
 Callisaurus draconoides, 94
crisatellus wileyi, *Anolis*, 86
croceator, *Ensatina*, 13
crospedospila, *Hyla*, 52
Crossodactylodes pintoi, 39
Crossodactylus dispar, 40
 gaudichaudii, 48
Crotalophorus edwardsii, 168
Crotalus adamanteus ruber, 168
 atrox, 169
 atrox elegans, 169
 basiliscus basiliscus, 162
 basiliscus oaxacus, 169
 cerastes, 169
 cerastes cerastes, 169
 cerastes cercobombus, 169
 confluentus abyssus, 169
 confluentus lutosus, 169
 confluentus stephensi, 169
 durissus culminatus, 170
 durissus tzabcan, 170
 enyo, 162
 gloydi lautus, 170
Crotalus—Continued
 goldmani, 170
 intermedius intermedius, 170
 jimenezii, 170
 lecontei, 171
 lucifer, 171
 mitchellii pyrrhus, 163, 170
 mitchellii stephensi, 169
 molossus, 171
 molossus estebanensis, 171
 molossus molossus, 171
 ornatus, 171
 polystictus, 170
 pricei miquihuanus, 172
 ravus, 171
 ruber ruber, 168, 169
 stejnegeri, 171
 tigris, 171
 triseriatus miquihuanus, 172
 viridis abyssus, 169
 viridis caliginis, 172
 viridis lutosus, 169
 viridis nuntius, 172
 viridis oreganus, 171
 viridis viridis, 171
 willardi meridionalis, 172
 willardi silus, 172
Crotaphytus baileyi, 104
 collaris baileyi, 104
 copei, 104
 dickersonae, 104
 dorsalis, 104
 fasciatus, 104
 gambelii, 104
 reticulatus, 104
 silus, 104
 wislizenii, 105
cruciger vogli, *Atelopus*, 30
cruentatus consors, *Kinosternon*, 232
crusculus molesworthi, *Celestus*, 95
Cryptoblepharus gloriosus, 81
Cryptobranchus alleganiensis bishopi, 11
 bishopi, 11
Ctenosaura acanthura, 105
 bakeri, 105
 brevirostris, 105
 clarkii, 105
 completa, 105
 conspicua, 105
 hemilopha, 105, 106
 insulana, 105
 multispinis, 105
 palearis, 106

- Ctenosaura—Continued
 parkeri, 106
 pectinata, 105, 106
 similis similis, 105
 teres brachylopha, 106
 cubanus, Eleutherodactylus, 42
 culex, Hyla, 52
 culminatus, Crotalus durissus, 170
 cunctator, Eleutherodactylus, 43
 cuneatus, Eleutherodactylus, 61
 Hylodes, 61
 cuneus, Hypopachus, 62
 Hypopachus cuneus, 62
 cupreus, Anolis, 86
 Sceloporus siniferus, 134
 curta, Hyla, 52
 curtissi, Celestus, 95
 curtus brongersmai, Python, 208
 Leptodactylus, 63
 cuspidata, Hyla, 53
 cyanogenys, Sceloporus, 143
 Sceloporus torquatus, 143
 cyanopleurus, Anolis, 86
 Anolis (Dracontura), 86
 cyanosticta, Hyla phacota, 57
 cybotes doris, Anolis, 87
 cyclides, Thamnophis cyrtopsis, 218
 cyclopion floridana, Natrix, 199
 Cycloramphus granulosus, 40
 neglectus, 40
 Cyclorhamphus boulengeri, 40
 granulosus, 40
 Cyclura carinata bartschi, 106
 (Ctenosaura) hemilopha, 106
 mattea, 106
 nigerrima, 106, 107
 onchiopsis, 107
 rileyi, 107
 stejnegeri, 107
 cyrtopsis cyclides, Thamnophis, 218
 Eutaenia, 179
 ocellata, Eutaenia, 179
 Thamnophis cyrtopsis, 179, 218
 cystignathoides, Phylllobates, 69
 Syrrophus, 69
 Cystignathus labialis, 40
 microtis, 40
 perlaevis, 40
 poecilochilus, 40
 Daconura bivittata, 107
 Dactyloperus insulensis, 107
 danforthi, Alsophis leucomelas, 157
 danieli dunni, Gyrinophilus, 15
 darlingtoni, Anolis, 86
 Celestus, 95
 Eleutherodactylus, 43
 Leptodactylus, 44, 63
 debilis, Bufo, 32
 Bufo debilis, 32
 insidiator, Bufo, 34
 kelloggi, Bufo, 35
 decemlineatus, Cnemidophorus, 97
 decipiens, Hyla, 53
 deekerti, Elaphe quadrivittata, 177
 decorata, Pseudemys, 234
 Rhadinaea, 174
 decorticatum, Amblystoma, 5
 Amblystoma gracile, 5
 decurtatus perkinsi, Phyllorhynchus, 206
 defensor, Cachryx, 94
 Enyaliosaurus, 94
 degenhardtii apiata, Stenorhina, 214
 deignani, Sphenomorphus, 148
 Deirochelys reticularia miaria, 230
 Deiroptyx bartschi, 107
 dekayi temporalineata, Storeria, 214
 texana, Storeria, 214
 tropica, Storeria, 216
 wrightorum, Storeria, 215
 delalandii, Mabuya, 113
 delanonis, Tropidurus, 151
 delicatissima, Cochranella, 28
 delicatissimus, Sceloporus, 134
 deltae, Natrix rigida, 200
 Dendraspis jamesoni kaimosae, 172
 Dendroaspis jamesoni kaimosae, 172
 Dendrobates auratus, 69
 Dendrophidion vinitor, 172
 Dendrophidium chloroticum, 173
 melanotropis, 173
 dendrophila occidentalis, Boiga, 161
 Dendrophis prasinus, 173
 dendrocarta, Hyla, 53
 deppei jani, Pituophis, 159
 deppii, Cnemidophorus deppii, 97, 100
 deppii lineatissimus, Cnemidophorus, 97,
 100
 depressirostris, Thalerophis, 193
 deserticola, Hypsiglena o[chrorhyncha],
 186
 Hypsiglena torquata, 186
 Mabuya, 126
 Pituophis catenifer, 206
 Desmognathus aeneus, 11
 aeneus aeneus, 11
 aeneus chermocki, 12

- Desmognathus—Continued
 aureatagulus, 11
 brimleyorum, 11
 chermocki, 12
 fuscus brimleyorum, 11
 fuscus imitator, 12
 fuscus welteri, 12
 monticola, 12
 monticola jeffersoni, 12
 monticola monticola, 12
 ochrophaea carolinensis, 12
 ochrophaeus carolinensis, 11, 12
 ocoee, 13
 wrighti, 13
 dextra, Ameiva undulata, 83
 diaboli, Ambystoma tigrinum, 7
 diademata nietoi, Ninia, 201
 Ninia, 201
 Ninia diademata, 201
 Diadophis amabilis, 173
 amabilis amabilis, 173
 arnyi, 173
 punctatus arnyi, 173
 punctatus stictogenys, 174
 regalis, 173
 regalis arizonae, 173
 regalis blanchardi, 173
 regalis regalis, 173
 texensis, 174
 Diaglena reticulata, 40
 diasii, Chionactis, 164
 diastema, Eleutherodactylus, 65
 Lithodytes, 65
 proximans, Micrurus, 198
 Dicamptodon ensatus, 4, 6
 dickersonae, Crotophytus, 104
 Holbrookia, 116
 Holbrookia maculata, 116
 Dicrodon holmbergi, 107
 Diemictylus viridescens dorsalis, 13
 viridescens evergladensis, 13
 viridescens perstriatus, 28
 viridescens piaropicola, 13
 Diemictylus viridescens vittatus, 13
 difficilis, Sphaerodactylus, 148
 dimidiata, Bolitoglossa, 9
 Chiropterotriton, 9
 dimidiatus, Pliocercus, 208
 Dinodon rufozonatum walli, 174
 semicarinatum, 191
 Diphalus fenestratus, 108
 Diplodactylus inexpectatus, 108
 unctus, 108
 Diploderma polygonatum, 108
 Diploglossus chalybaeus, 108
 steindachneri, 108
 diporus, Bothrops, 161
 Bothrops neuwiedii, 161
 Dipsas argus, 192
 brevifacies, 222
 gemmistrata latistrata, 174
 macrostomus, 213
 pictiventris, 192
 sanniolus, 198
 septentrionalis, 174
 smithi, 174
 Dipsosaurus dorsalis, 104
 dorsalis sonoriensis, 108
 dispar, Crossodactylus, 40
 disparilis, Cnemidophorus, 97
 Sceloporus, 135
 Sceloporus grammicus, 135
 distans, Elaps, 177
 Micrurus distans, 177
 distichoides, Anolis, 87
 distichus caudalis, Anolis, 87
 floridanus, Anolis, 87
 juliae, Anolis, 87
 wetmorei, Anolis, 87
 ditmarsii, Phrynosoma, 128
 divaricans, Cochranella, 38
 dixi, Plethodon, 19
 Plethodon wehrlei, 19
 doliatus caesius, Opheomorphus, 202
 collaris, Ophibolus, 203
 parallelus, Ophibolus, 203
 semicinctus, Oxyrrhopus, 205
 sypilus, Ophibolus, 203, 204
 dominicensis caudalis, Anolis, 87
 juliae, Anolis, 87
 Leptodactylus, 64
 wetmorei, Anolis, 87
 dontomasi, Gageia, 113
 dorbignyi, Heterodon, 185
 Lystrophis, 185
 doris, Anolis, 87
 Anolis cybotes, 87
 dorsalis angusticlavius, Plethodon, 19
 Bufo, 33, 37
 Chrysemys, 229
 Chrysemys picta, 229
 Crotophytus, 104
 Diemictylus viridescens, 13
 Dipsosaurus, 104
 Platyplacopus, 149
 Plethodon cinereus, 19
 Plethodon dorsalis, 19
 sonoriensis, Dipsosaurus, 108
 Takydromus, 149

- dorsivittata*, *Mabuia*, 125
 Mabuia, 125
dorsoconcolor, *Eleutherodactylus*, 43
Doryura vulpecula, 108
douglassi pygmaea, *Phrynosoma*, 128
douglassii brachycercum, *Phrynosoma*, 128
 brevirostris, *Phrynosoma*, 150
 hernandesi, *Phrynosoma*, 150
 ornatissimum, *Phrynosoma*, 150
 Phrynosoma douglassii, 128
Draco mindanensis, 109
draconoides carmenensis, *Callisaurus*, 94
 crinitus, *Callisaurus*, 94
 inuitatus, *Callisaurus*, 94
 myurus, *Callisaurus*, 95
 rhodostictus, *Callisaurus*, 95
 splendidus, *Callisaurus*, 95
 ventralis, *Callisaurus*, 118
draytonii, *Rana*, 73
 Rana aurora, 73, 74, 76
Drepanodon attenuatus, 174
Dromicodryas bernieri, 184
Dromicus ignitus, 174
 leucomelas, 175
 parvifrons rosamondae, 175
Dryadophis amarali, 178
 bifossatus bifossatus, 168
 melanolomus melanolomus, 197
 melanolomus stuarti, 175
 melanolomus tehuanae, 175
Drymarchon corais erebennus, 213
 corais melanurus, 213
 corais rubidus, 175
 corais unicolor, 176
Drymobius aurigulus, 176
 chloroticus, 173
 margaritiferus fistulosus, 176
 melanotropis, 173
Dryophis brevirostris, 176
dubia, *Cochranella*, 38
dubitus, *Thorius*, 27
dugesi, *Scaphiopus*, 78
dulcis, *Leptotyphlops dulcis*, 209
 Rena, 209
dumerilii, *Bathysiredon*, 24
 Siredon, 24
duncanensis, *Tropidurus*, 151
dunni, *Bufo*, 33
 Conophis lineatus, 167
 Gyrinophilus, 15
 Gyrinophilus danieli, 15
 Plethodon, 19
dura, *Tiliqua*, 151
durangoensis, *Phrynosoma orbiculare*, 129
durissus culminatus, *Crotalus*, 170
 tzabcan, *Crotalus*, 170
durum, *Sphenosoma*, 151
duryi, *Gyrinophilus porphyriticus*, 23
 Pseudotriton, 23
edwardsii, *Crotalophorus*, 168
 Sistrurus catenatus, 168
edwardtaylori, *Sceloporus*, 135
egregius, *Eumeces*, 131
 Plestiodon, 131
eileenae, *Eleutherodactylus*, 43
eiseni, *Ophibolus getulus*, 204
 Tantilla, 217
 Tantilla eiseni, 217
Elachistocleis ovalis, 62
elacochroa, *Hyla*, 53, 58
Elaphe flavirufa flavirufa, 165
 flavirufa matudai, 176
 flavirufa phaescens, 176
 guttata emoryi, 166
 guttata guttata, 165
 guttata rosacea, 166
 obsoleta bairdi, 164
 obsoleta lindheimeri, 212
 obsoleta obsoleta, 166, 212
 obsoleta quadrivittata, 177
 oxycephala, 157
 perlacea, 176
 quadrivittata deckerti, 177
 triaspis mutabilis, 165
 triaspis phaescens, 176
 triaspis triaspis, 166
 vulpina vulpina, 213
elapiformis, *Cantoria*, 162
elapoides laticollaris, *Pliocercus*, 208
Elapomorphus suspectus, 177
Elaps distans, 177
 ephippifer, 177
 euryxanthus, 177
 melanocephalus, 177
 nigrocinctus, 177
 teneris, 177
 tristis, 177
elapsoides virginiana, *Lampropeltis*, 188
eleanorae, *Ameiva wetmorei*, 84
elegans, *Arizona*, 158
 Arizona elegans, 158
 Atelopus, 30
 atratus, *Thamnophis*, 180
 biscutatus, *Thamnophis*, 179
 blanchardi, *Arizona*, 159
 brunnea, *Eutaenia*, 179
 Chlorosoma, 216
 couchii, *Thamnophis*, 178

elegans—Continued

- Crotalus atrox*, 169
- Eutainia, 181
- expolita, Arizona, 159
- hammondii, *Thamnophis*, 179
- Imantodes, 192
- lineolata, *Eutaenia*, 179
- nigrescens, *Thamnophis*, 218
- noctivaga, Arizona, 159
- occidentalis, Arizona, 159
- Thamnophis elegans*, 181, 183
- Uta, 153
- Uta stansburiana, 153
- vagrans, *Thamnophis*, 179, 180, 181, 183

Eleutherodactylus abbotti, 41

- albipes, 41
- alticola, 41
- andrewsi, 41
- atkinsi orientalis, 41
- audanti, 41
- avocalis, 41
- bakeri, 42
- beebei, 42
- binghami, 42
- casparii, 42
- cavernicola, 42
- cerasinus, 61
- conspicuus, 42
- cubanus, 42
- cunctator, 43
- cuneatus, 61
- darlingtoni, 43
- diastema, 65
- dorsoconcolor, 43
- eileenae, 43
- footei, 43
- fuscus, 43
- goldmani, 43
- gollmeri, 43, 65
- gossei, 44
- grabhami, 44
- greyi, 44
- guentheri, 61
- haitianus, 44
- hylaeformis, 69
- intermedius, 44
- jugans, 44, 63
- junori, 44
- karlschmidti, 45
- latrans, 65
- lewisi, 45
- luteolus, 45
- lymani, 45

Eleutherodactylus—Continued

- lynni, 45
 - matudai, 45
 - megalops, 45
 - melanostictus, 66
 - milesi, 45
 - nasutus, 61
 - nubicola, 45
 - orcutti, 43, 46
 - pantoni, 46
 - parvus, 42
 - pinarensis, 46
 - podiciferus, 66
 - poolei, 46
 - ranoides, 66
 - rhodopsis, 66
 - richmondi, 46
 - ricordii casparii, 42
 - ricordii rogersi, 46
 - rugosus, 65, 66
 - rugulosus, 66
 - saltator, 46
 - schmidti limbensis, 47
 - schmidti rucillensis, 47
 - spatulatus, 47
 - talamancae, 47
 - thomasi trinidadensis, 47
 - turquinensis, 47
 - unicolor, 47
 - ventrilineatus, 44, 63
 - vocalis, 47
 - vulcani, 47
 - wetmorei, 48
- Elgaria grandis*, 109
- multicarinata webbii, 115
 - nobilis, 109
 - principis, 109
- elgonensis*, *Rharnophis aethiopissa*, 210
- elonae*, *Pseudemys*, 234
- elongata*, *Hyla*, 53
- elongatus*, *Sceloporus*, 135
- Sceloporus undulatus*, 135
- elongis*, *Agama agama*, 81
- Elosia vomerina*, 48
- Emoa nigrita*, 109
- Emoia arnoensis*, 109
- boettgeri orientalis*, 109
 - frenata, 110
 - nigrita, 109
- emoryi*, *Aspidonectes*, 227
- Elaphe guttata*, 166
 - Trionyx spinifer*, 227
- Emydocephalus ijimae*, 178
- Emys marmorata*, 230

- Engystoma areolata*, 48
 pictiventre, 48
 texense, 48
 ustum, 48
 variolosum, 48
enneagrammus, *Celestus*, 108
Ensatina croceator, 13
 eschschooltzii klauberi, 13
 eschschooltzii oregonensis, 15
 eschschooltzii xanthoptica, 13
 klauberi, 13
ensatus, *Dicamptodon*, 4, 6
ensicauda, *Triton*, 27
ensicaudus, *Triturus*, 27, 28
ensicaudus popei, *Triturus*, 28
 Triturus ensicaudus, 27, 28
Enyaliosaurus bakeri, 105
 clarkii, 105
 defensor, 94
 palearis, 106
enyo, *Caudisona*, 162
 Crotalus, 162
Epaphelus sumichrasti, 110
ephippiata, *Leptodeira*, 191
 Leptodeira splendida, 191
ephippicus, *Chilomeniscus*, 163
ephippifer, *Elaps*, 177
 Micrurus, 177
Epicrates cenchria crassus, 178
 crassus, 178
 subflavus, 178
epipleurotis, *Eumeces*, 111
episcopa, *Sonora episcopa*, 190
episcopum, *Lamprosoma*, 190
equatoriana, *Barbourina*, 160
eques megalops, *Thamnophis*, 180
 Thamnophis eques, 179, 180
erebennus, *Drymarchon corais*, 213
 Spilotes, 213
Eremias hoehneli, 110
 nitida garambensis, 110
 sextaeniata, 110
 spekii sextaeniata, 110
eremitus, *Leiocephalus*, 123
 Liocephalus, 123
Eretmochelys imbricata bissa, 228
 imbricata squamata, 228
errans, *Thamnophis*, 218
 Thamnophis ordinoides, 218
ervingi, *Stenocercus*, 149
erythrocheilus, *Sceloporus undulatus*, 144
erythrogaster, *Coluber*, 165
 flavigaster, *Natrix*, 199
 Natrix, 165
erythrogaster—Continued
 neglecta, *Natrix*, 199
 transversa, *Natrix*, 201, 222, 223
Erythrolamprus mimus micrurus, 178
erythrurus, *Phrynocephalus*, 128
eschschooltzii klauberi, *Ensatina*, 13
 oregonensis, *Ensatina*, 15
 xanthoptica, *Ensatina*, 13
estebanensis, *Cnemidophorus*, 97
 Cnemidophorus tigris, 97
 Crotalus molossus, 171
Euchirotes biporus, 110
Eudryas amarali, 178
Eumeces anthracinus anthracinus, 131
 anthracinus pluvialis, 112
 brevilineatus, 110
 brevirostris bilineatus, 110
 copei, 111
 egregius, 131
 epipleurotis, 111
 gilberti gilberti, 113
 indubitus, 111
 lagunensis, 111
 laticutatus okadae, 111
 longirostris, 131
 managuae, 111
 marginatus, 131
 marginatus amamiensis, 111
 multilineatus, 112
 multivirgatus, 111, 131
 obsoletus, 131
 ochoterenae, 112
 oshimensis, 112
 parviauriculatus, 112
 parvulus, 112
 pekinensis, 112
 pluvialis, 112
 quadrilineatus, 112
 quadrivirgatus, 112
 septentrionalis, 132
 septentrionalis pallidus, 113
 skiltonianus brevipes, 113
 skiltonianus interparietalis, 113
 skiltonianus skiltonianus, 112, 132
 sumichrasti, 132
 tetragrammus, 132
 tunganus, 113
eunice, *Leiolopisma*, 121
Eupemphix bolbodactyla, 48
 maculiventris, 49
 olfersioides, 49
 ruthveni, 49
Euphyryne obesus, 113
Euprepis venustus, 113

- Eupsophus lutzi*, 80
Eurycea bislineata bislineata, 13, 25
 bislineata major, 13
 bislineata rivicola, 14
 griseogaster, 14
 latitans, 14
 longicauda pernix, 14
 multiplicata, 26
 multiplicata griseogaster, 14
 neotenes, 14
 tynercensis, 15
Eurypholis major, 184
 semicarinatus, 178
euryxanthus, *Elaps*, 177
 Micruroides, 177
Euspondylus marianus, 132
Eutaenia angustirostris, 178
 chrysocephala, 178
 couchii, 178
 cyrtopsis, 179
 cyrtopsis ocellata, 179
 elegans brunnea, 179
 elegans lineolata, 179
 flavilabris, 179
 hammondii, 179
 henshawi, 180
 infernalis vidua, 180, 181
 insigniarum, 180
 macrostemma, 180
 megalops, 180
 pulchrilatus, 180
 radix twiningi, 180
 sirtalis semifasciata, 181
 sirtalis tetrataenia, 181
 sumichrasti, 181
 vagrans plutonia, 181
Eutainia atrata, 180, 181
 elegans, 181
 faireyi, 181
 haydenii, 182
 leptocephala, 182
 marciana, 182
 phenax, 182
 pickeringi, 182
 radix, 182
 rutilorus, 182
 sirtalis obscura, 183
 sirtalis trilineata, 183
 vagrans, 183
euthysanota, *Hyla*, 53
eutropia, *Ameiva*, 82
evansii, *Ophibolus*, 203
evelynae, *Hyla*, 53
evergladensis, *Diemictylus viridescens*, 13
evermanni, *Anolis*, 87
evittata, *Hyla*, 53
 Hyla cinerea, 53
 Rana cantabrigensis, 72
eximia, *Hyla*, 54
 Hyla eximia, 54
 wrightorum, *Hyla*, 60
expectata, *Aristelligella*, 92
expectatus, *Aristelliger*, 92
expolita, *Arizona elegans*, 159
exsul, *Ameiva*, 83
 Ameiva plei, 83
 Bufo, 33
 Bufo boreas, 33
 Sphaerodactylus, 146
extenuatum arenicola, *Stilosoma*, 214
extremus, *Anolis trinitatis*, 91
fairchildi, *Anolis*, 88
faireyi, *Eutainia*, 181
fasciata pictiventris, *Natrix*, 199
 pleuralis, *Natrix*, 200
fasciatus, *Chilomeniscus stramineus*, 163
 Crotaphytus, 104
 Lepidocephalus, 191
fasciolaris, *Hombromia*, 118
 Leiolopisma, 118
fasciolata, *Trimorphodon*, 221
fastidiosus, *Bufo*, 39
 Cranopsis, 39
felis, *Pseudemys*, 234
femoralis, *Prostherapis*, 71
fenestrata, *Amphisbaena*, 108
fenestratus, *Diphalus*, 108
feriarum, *Helocaetes*, 50
 Pseudacris nigrita, 50
ferraripercezi, *Sceloporus*, 135
ferreus, *Anaides*, 8
 Anaides, 8
 Anolis, 85
festiva, *Ameiva festiva*, 82
Ficimia cana, 184
 publia, 183
 publia publia, 183
 publia taylori, 183
firscheini, *Pseudoeurycea*, 23
fisheri, *Rana*, 73
 Rana pipiens, 73
fissidens proterops, *Coniophanes*, 166
fistulosus, *Drymobius margaritiferus*, 176
fitzingeri frizzelli, *Oxyrhopus*, 205
flagellum frenatum, *Basconion*, 160
 lineatulus, *Masticophis*, 196
 piceus, *Masticophis*, 160, 227
 variolosus, *Masticophis*, 197

- flavescens*, *Kinosternon flavescens*, 233
 Phyllosira, 206
 Platythyra, 233
 spooneri, *Kinosternon*, 232
flavifrenatus, *Aporophis*, 196
 Lygophis, 196
flavigaster, *Natrix erythrogaster*, 199
flavilabris, *Eutaenia*, 179
flavilineata, *Tantilla*, 217
flavimaculatum tehuanae, *Lepidophyma*, 123
flavimembris, *Magnadigita*, 17
 Oedipus, 17
flavipunctata, *Natrix piscator*, 158
flavipunctatum, *Amphiesma*, 158
flavirufa, *Elaphe flavirufa*, 165
 matudai, *Elaphe*, 176
 phaescens, *Elaphe*, 176
flavirufus, *Coluber*, 165
flaviventris, *Bolitoglossa*, 17
 Oedipus, 17
flavoguttata, *Hyla*, 54
 Hyla strigilata, 54
flavomarginatus, *Gopherus*, 231
flavoviridis, *Ranoidea*, 77
 tinkhami, *Trimeresurus*, 220
floridana concinna, *Pseudemys*, 234
 hoyi, *Pseudemys*, 235
 Lampropeltis getulus, 188
 mobilensis, *Pseudemys*, 235
 Natrix cyclopion, 199
 peninsularis, *Pseudemys*, 234
 Rhineura, 123
 texana, *Pseudemys*, 234
floridanum, *Lepidosternon*, 123
floridanus, *Anolis distichus*, 87
 Lampropeltis getulus, 188
 Pseudotriton montanus, 23
 Sceloporus, 135
flotator, *Phyllobates*, 69
fluvinatum, *Ambystoma*, 6
foliosquama, *Aipysurus*, 157
footei, *Eleutherodactylus*, 43
forbesi, *Rhadinaea*, 209
 Trimorphodon, 221
forbesorum, *Leiopisma*, 122
 Scincella gemmingeri, 122
formosa, *Chelonia*, 228
formosensis, *Sphenomorphus indicus*, 148
formosus, *Sceloporus formosus*, 145
 scitululus, *Sceloporus*, 135
foxtii, *Bascanion*, 160
 Coluber constrictor, 160
franklini, *Magnadigita*, 17
 Oedipus, 17
freminvillii lactea, *Stenorhina*, 214
frenata, *Emoia*, 110
 Mabuya, 110
 Paludicola, 68
frenatum, *Bascanion flagellum*, 160
frenatus, *Hemidactylus*, 116
frizzelli, *Oxyrhopus fitzingeri*, 205
frontale, *Phrynosoma coronatum*, 129
frontalis, *Pseudoficimia*, 219
 Toluca, 219
fulgida, *Mabuia*, 126
fuliginatus, *Pituophis catenifer*, 207
fuliginosa, *Ameiva ameiva*, 82, 150
fuliginosus, *Tiaporus*, 150
 Zamenis lateralis, 227
fulviceps, *Rhadinaea*, 210
 Rhadinaea pachyura, 210
fulvius, *Micrurus*, 198
 teneris, *Micrurus*, 177
fulvivittis, *Rhadinaea*, 210
fulvum, *Arrhyton*, 160
funerea, *Geoemyda*, 228
funereus, *Alsophis*, 157
 Chelopus, 228
 Leimadophis, 157
fumiceps, *Scolecophis*, 212
 Tantilla nigriceps, 212, 217
fusciventris, *Ceratophrys*, 38
fuscocomarginata, *Hyla*, 54
fuscovaria, *Hyla*, 54
fuscus, *Chironius*, 206
 brimleyorum, *Desmognathus*, 11
 Eleutherodactylus, 43
 imitator, *Desmognathus*, 12
 welteri, *Desmognathus*, 12
gabbi, *Bufo*, 31, 33
gabbiana, *Amiva*, 84
gabpii, *Chelopus*, 228
 Geoemyda, 228
 Hyla, 54
gadovi, *Cnemidophorus*, 98
gadovii levigata, *Barisia*, 93
 Oedipus, 18
 Pseudoecurycea, 18
gaigeae, *Leptodactylus*, 64
 Pseudemys scripta, 234
 Sphaerodactylus, 146
Gaigeia dontomasi, 113
 radula, 114
galaenae, *Bolitoglossa*, 9
galeanae, *Pseudoecurycea*, 9
Gambelia wislizenii silus, 104
 wislizenii wislizenii, 104, 105
gambelii, *Crotaphytus*, 104

- garambensis, *Eremias nitida*, 110
 garciai, *Trimeresurus*, 220
 garnotii, *Hemidactylus*, 108
Gastrotheca boliviana, 42
 coronata, 49
 monticola, 49
 gaudichaudii, *Crossodactylus*, 48
 Geagras redimitus, 183
 Gehyra vorax, 114
 Gekko gekko, 114
 indicus, 114
 gemmifer, *Bufo*, 33
 gemmingeri forbesorum, *Scincella*, 122
 Oligosoma, 127
 Scincella gemmingeri, 127
 gemmistrata latistrata, *Dipsas*, 174
 gentilis, *Anolis cepedii*, 86
 Lampropeltis triangulum, 188, 204
 Ophibolus, 203, 204
 Geomyda funerea, 228
 gabbii, 228
 rubida rubida, 228
 Geophis anocularis, 183
 blanchardi, 184
 cancellatus, 184
 chalybeus, 209
 nasalis, 162
 georgeensis, *Aristelliger*, 92
 Gerrhonotus alfaroi, 114
 auritus, 114
 coerulea palmeri, 115
 coeruleus principis, 109
 gramineus, 114
 infernalis, 114
 kingii, 109
 levicollis ciliaris, 114
 liocephalus infernalis, 114
 liocephalus ophiurus, 115
 monticolus, 115
 multicarinatus scincicaudus, 109
 multicarinatus webbiai, 115
 olivaceus, 115
 ophiurus, 115
 scincicauda palmeri, 115
 webbiai, 115
 getulus californiae, *Lampropeltis*, 203, 204
 conjuncta, *Lampropeltis*, 187
 eiseni, *Ophibolus*, 204
 floridana, *Lampropeltis*, 188
 floridanus, *Lampropeltis*, 188
 multicinctus, *Ophibolus*, 204
 niger, *Lampropeltis*, 204
 niger, *Ophibolus*, 204
 splendida, *Lampropeltis*, 204
 yumensis, *Lampropeltis*, 188
 gibbicaudus, *Spelerpes*, 25, 26
 gigas, *Cnemidophorus*, 98
 gilberti, *Eumeces gilberti*, 113
 gingivinus, *Anolis*, 92
 glaucus, *Sphaerodactylus*, 146
 Sphaerodactylus glaucus, 146
 gloriosus, *Ablepharus*, 81
 Cryptoblepharus, 81
 gloydii, *Heterodon nasicus*, 185
 lautus, *Crotalus*, 170
 glutinosus chlorobryonis, *Plethodon*, 19
 glutinosus, *Plethodon glutinosus*, 19, 20
 glyphus, *Atelopus varius*, 30
 goebeli, *Oedipus*, 18
 Pseudoeurycea, 18
 goldmani, *Crotalus*, 170
 Eleutherodactylus, 43
 Sceloporus, 135
 Terrapene, 235
 gollmeri, *Eleutherodactylus*, 43, 65
 Gonatodes atricucularis, 115
 nigridius, 116
 Goniochelys minor, 230
 Gonocephalus abbotti, 116
 goodei, *Phrynosoma*, 129
 Phrynosoma platyrhinos, 129
 Gopherus agassizii, 235
 berlandieri, 236
 flavomarginatus, 231
 gossei, *Eleutherodactylus*, 44
 Sphaerodactylus richardsoni, 147
 goughi baileyi, *Hyla*, 54
 grabhami, *Eleutherodactylus*, 44
 gracia, *Lichanura roseofusca*, 195
 gracile, *Ambystoma gracile*, 5, 25
 decorticatum, *Ambystoma*, 5
 gracilipes, *Hyla*, 54
 gracilis, *Atelopus*, 30
 Cnemidophorus, 98
 Cnemidophorus tigris, 98, 100, 102
 Siredon, 25
 graciosus, *Sceloporus*, 136
 Sceloporus graciosus, 136
 vandenburgianus, *Sceloporus*, 144
 grahami lewisi, *Anolis*, 86
 Microhyla, 67
 Phoxophrys, 128
 grahamiae, *Salvadora*, 211
 grahamii, *Cnemidophorus*, 98
 Natrix, 208
 Regina, 208
 graminea, *Abronia taeniata*, 114
 gramineus, *Gerrhonotus*, 114
 grammicus disparilis, *Sceloporus*, 135

- grammicus microlepidotus*, *Sceloporus*, 136
grandaevus, *Sator*, 133
grandis, *Elgaria*, 109
grandisquamis, *Sphaerodactylus*, 147
granti, *Leiocephalus carinatus*, 119
granulata, *Salamandra*, 24
granulosum, *Ambystoma*, 6
granulosus, *Bufo*, 33, 36
 Cycloramphus, 40
 Cyclorhamphus, 40
Graptemys oculifera, 233
 pseudogeographica ouachitensis, 231
 pseudogeographica sabinensis, 231
 pseudogeographica versa, 231
 pulchra, 231
 versa, 231
gratiosa, *Hyla*, 54
greyi, *Anolis*, 88
 Eleutherodactylus, 44
grisea milneana, *Rana*, 73
griseogaster, *Eurycea*, 14
 Eurycea multiplicata, 14
griseus, *Anolis*, 88
 Anolis richardii, 88
grotei, *Lygodactylus grotei*, 124
gryllo, *Rana*, 74
gryllus, *Acris gryllus*, 74
 paludicola, *Acris*, 29
 Philautus, 68
 Rana, 74
guentheri, *Eleutherodactylus*, 61
gularis, *Cnemidophorus*, 98
 Cnemidophorus sackii, 98, 99
 gularis obsoletus, *Cnemidophorus*, 98
 gularis verus, *Cnemidophorus*, 98
 scalaris, *Cnemidophorus*, 98
 semifasciatus, *Cnemidophorus*, 99
 sericeus, *Cnemidophorus*, 99
gulosus, *Lithodytes*, 65
güntheri, *Bufo*, 33
guttata, *Elaphe guttata*, 165
 emoryi, *Elaphe*, 166
 Phyllomedusa, 70
 rosacea, *Elaphe*, 166
guttatus, *Cnemidophorus guttatus*, 103
 immutabilis, *Cnemidophorus*, 99, 100
 sellatus, *Coluber*, 165
guttulatus, *Malachyloides*, 66
 Syrrhophus, 66
guttulatum, *Rhabdosoma*, 209
guttulatus, *Basiliscus*, 93
Gyalopion canum, 184
Gymnodactylus pelagicus, 116
Gymnopia, 3
Gymnophthalmus sumichrasti, 110
Gymnopsis multiplicata multiplicata, 4
 multiplicata proxima, 4
 oligozona, 4
Gyrinophilus danielsi dunnii, 15
 dunnii, 15
 porphyriticus duryi, 23
 porphyriticus inagnoscus, 15
habra, *Sphoerohyla*, 79
haematiticus, *Bufo*, 34
 lachrymans, *Bufo*, 34
haematois, *Pityophis*, 208
haetiana, *Ungalia*, 225
 Wetmorena, 156
haetianus, *Tropidophis maculatus*, 225
haitianus, *Eleutherodactylus*, 44
Haldea valeriae valeriae, 226
hallowelli, *Hyla*, 55
halophila, *Bufo*, 34
Halophila heros, 49
 vitiensis, 49
halophilus, *Bufo boreas*, 34
hammondii, *Eutaenia*, 179
 multiplicatus, *Scaphiopus*, 78
 Scaphiopus hammondii, 78
 Thamnophis elegans, 179
hansena, *Philautus*, 68
harteri, *Natrix*, 200
hartwegi, *Ameiva undulata*, 83
 Amyda spinifera, 227
 Trionyx spinifer, 227
hatcheri, *Liolaemus*, 124
haydenii, *Eutania*, 182
 Thamnophis radix, 180, 182
hebes, *Phrynohyas*, 78
 Scytotis, 78
helenae, *Agalychnis*, 29
 Carphophis amoenus, 163
 Celuta, 163
 Sphenomorphus, 148
Helicops leopardinus, 184
 lepieurii, 184
helmrichi, *Bolitoglossa*, 18
 Oedipus, 18
Helocaetes clarkii, 50
 feriarum, 50
Heloderma suspectum, 116
Hemidactylum pacificum, 15
Hemidactylus frenatus, 116
 garnotii, 108
 inornatus, 116
hemilopha, *Ctenosaura*, 105, 106
 Cyclura (Ctenosaura), 106

- hemiophrys, *Bufo*, 34
 Bufo woodhousei, 34
Hemiphyllodactylus leucostictus, 116
Hemipipa carvalhoi, 71
hendersoni, *Anolis*, 88
henriquesi, *Sphaerodactylus argus*, 146
henshawi, *Eutaenia*, 180
 Xantusia, 156
Heredia oregonensis, 15
hernandesi, *Phrynosoma douglassii*, 150
 Tapaya, 150
heros, *Cornufer*, 49
 Halophila, 49
Herpetodryas bernieri, 184
 chloris, 184
 melas, 184
herrerai, *Kinosternon*, 232
hesperia hesperioides, *Rhadinaea*, 210
 Rhadinaea, 210
 Rhadinaea hesperia, 210
hesperioides, *Rhadinaea hesperia*, 210
hesperis, *Uta stansburiana*, 155
Heterodon atmodes, 184
 browni, 185
 cognatus, 185
 dorbignyi, 185
 kennerlyi, 185
 nasicus gloydi, 185
 nasicus kennerlyi, 185
 nasicus nasicus, 185
 platyrhinos, 184, 185
Heteronota pelagica, 116
heterurus, *Sceloporus*, 136
hexalepis celeris, *Salvadora*, 211
 klauberi, *Salvadora*, 211
 mojavensis, *Salvadora*, 211
 Phimothyra, 205
 Salvadora hexalepis, 205
Himantodes leucomelas, 185
 tenuissimus, 185
hippocrepis, *Kinosternon subrubrum*, 232
Hispaniolus pratensis, 116
hispidus, *Sauromalus*, 133
hoehneli, *Eremias*, 110
Holarchus ancorus, 226
Holbrookia affinis, 116
 dickersonae, 116
 lacerata, 117
 lacerata lacerata, 117
 lacerata subcaudalis, 117
 maculata dickersonae, 116
 maculata perspicua, 117
Gymnopiona, 3
Holbrookia—Continued
 maculata thermophila, 118
 propinqua, 117
 propinqua piperata, 117
 propinqua propinqua, 117
 pulchra, 117
 texana texana, 116
 thermophila, 118
holbrookii albus, *Scaphiopus*, 78
 Nerodia, 201
holmbergi, *Dicrodon*, 107
Homalosaurus ventralis, 118
Hombromia fasciolaris, 118
 undosa, 118
homolepidurus, *Phyllodactylus*, 130
homolepis, *Sphaerodactylus*, 147
hoodensis, *Tropidurus*, 151
Hoplodactylus pacificus, 118
 pomarii, 118
horribilis albiventris, *Sceloporus*, 136
horridus oligoporus, *Sceloporus*, 140
houstonensis, *Bufo*, 34
hoyi, *Pseudemys floridana*, 235
 Ptychemys, 235
hubrichti, *Plethodon*, 20
hughi, *Callophis*, 162
 Sphenosoma, 149
huico, *Cnemidophorus sacki*, 101
 Cnemidophorus sackii, 101
huldae, *Plethodon*, 20
humilis, *Anolis humilis*, 90
 cahuilae, *Leptotyphlops*, 194
 Leptotyphlops humilis, 209
 Rena, 209
 segregus, *Leptotyphlops*, 194
 slevini, *Leptotyphlops*, 194
 uniformis, *Anolis*, 90, 92
hyacinthinus, *Sceloporus undulatus*, 143
Hydrophis ornatus maresinensis, 185
Hydrops lubricus, 186
 trianguligaris venezuelensis, 186
Hyla acuminata, 50
 affinis, 50, 51
 albofrenata, 50
 albosignata, 50
 anceps, 50
 andersonii, 51
 angustilineata, 51
 arboricola, 51
 arenicolor, 50, 51
 aurantiaca orophila, 56
 aurantiaca planicola, 57
 aurantiaca surda, 51

Hyla—Continued

aurea, 77
 avivoca, 51
 bipunctata branneri, 51
 bischoffi multilineata, 56
 bistrincta, 51
 bistrincta bistrincta, 51
 bistrincta labeculata, 52
 boulengeri, 78
 cadaverina, 52, 56
 cárdenasi, 52
 ceratophrys, 52
 cinerea evittata, 53
 colymba, 52
 crospeospila, 52
 culex, 52
 curta, 52
 cuspidata, 53
 decipiens, 53
 dendroscarta, 53
 elaeochroa, 53, 58
 elongata, 53
 euthysanota, 53
 evelynae, 53
 evittata, 53
 eximia, 54
 eximia eximia, 54
 eximia wrightorum, 60
 flavoguttata, 54
 fuscomarginata, 54
 fuscovaria, 54
 gabbii, 54
 goughi baileyi, 54
 gracilipes, 54
 gratiosa, 54
 hallowelli, 55
 lesueurii, 77
 loquax, 55
 marianae, 55
 megapodia, 55
 melanomma, 55
 microcephala martini, 55
 microcephala sartori, 55
 microeximia, 55
 miliaria, 62
 milleri, 56
 minuta, 57
 moesta, 58
 molitor, 58
 monticola, 58
 multilineata, 56
 muricolor, 56
 nasica, 56
 nebulosa, 52, 56

Hyla—Continued

nigripes, 56
 (Sphoehyla) orophila, 56
 pachyderma, 57
 pallens, 57
 parkeri, 57
 phaeota, 57
 phaeota cyanosticta, 57
 phaeota phaeota, 57
 phlebodes, 57
 pickeli, 57
 pictipes, 58
 (Sphoehyla) planicola, 57
 puma, 58
 punctariola moesta, 58
 punctariola monticola, 58
 punctariola pictipes, 58
 punctata, 58
 quinquevittata, 58
 raniceps, 55, 62
 regilla, 52, 56, 58
 regilla laticeps, 58
 rickardsi, 59
 robertmertensi, 59
 robertsorum, 59
 rozellae, 59
 sanborni, 59
 septentrionalis brunnea, 81
 similis, 59
 smaragdina, 60
 squalirostris, 60
 staufferi, 52, 60
 strigilata flavoguttata, 54
 uranochroa, 60
 vanvlietii, 60
 variabilis, 60
 versicolor phaeocrypta, 60
 versicolor sandersi, 60
 versicolor versicolor, 60
 wrightii, 81
 wrightorum, 60
 zeteki, 61
 hylaeiformis, Eleutherodactylus, 69
 Phyllobates, 69
 Hylambates aubryi, 58
 cochranae, 61
 Hylaplesia nigriventris, 61
 Hylella platycephala, 61
 sumichrasti, 61
 Hylodes cerasinus, 61
 cuneatus, 61
 nasutus, 61
 polyptychus, 61
 Hymenochirus boettgeri camerunensis, 61

- Hynobius retardatus*, 15
 shihii, 15
 stejnegeri, 16
hyperythrus beldingi, *Cnemidophorus*, 156
 caeruleus, *Cnemidophorus*, 156
 Cnemidophorus, 99
 Cnemidophorus hyperythrus, 99
hypochondrialis, *Phyllomedusa*, 70
Hypogeophis alternans, 3
Hypopachus alboventer, 62
 caprimimus, 62
 championi, 62
 cuneus, 62
 cuneus cuneus, 62
 maculatus, 62
 ovis, 62
 pearsei, 62
 variolosus, 48
Hypsiboas miliarius, 62
 raniceps, 62
Hypsiglena chlorophaea, 186
 [ochrorhyncha] deserticola, 186
 ochrorhynchus, 186
 texana, 186
 torquata baueri, 186
 torquata deserticola, 186
 torquata ochrorhyncha, 186
 torquata texana, 186
Ialtris parishi, 187
Ichthyophis kotaensis, 3
 nigroflavus, 3
 paucidentulus, 3
 paucisulcus, 3
 sumatranus, 4
ictericus mertensi, *Bufo*, 34
idahoensis, *Plethodon*, 20
 Plethodon vandykei, 20
ignea, *Oedipina*, 17
ignitus, *Dromicus*, 174
ijimae, *Emydocephalus*, 178
illinoensis, *Pseudacris streckeri*, 71
Imantodes cenchoa leucomelas, 185
 elegans, 192
 latistratus, 174
 splendidus oliveri, 187
 tenuissimus, 185
imbricata, *Barisia imbricata*, 115
 bissa, *Eretmochelys*, 228
 ciliaris, *Barisia*, 114
 squamata, *Eretmochelys*, 228
imitator, *Desmognathus fuscus*, 12
 Microbatrachylus, 67
immaculata, *Clelia clelia*, 164
 Ninia sebae, 201
- immaculatus*, *Leiocephalus macropus*, 120
immucronatus, *Sceloporus jarrovi*, 136
immutabilis, *Cnemidophorus*, 99
 Cnemidophorus guttatus, 99, 100
imperialis, *Coniophanes imperialis*, 216
 Taeniophis, 216
inagnoscus, *Gyrinophilus porphyriticus*, 15
inaguae, *Leiocephalus*, 120
inca, *Bufo*, 34
incertum, *Lygosoma*, 125
indefatigabilis, *Tropidurus*, 151
indicus formosensis, *Sphenomorphus*, 148
 Gekko, 114
indubitus, *Eumeces*, 111
inexpectatus, *Diplodactylus*, 108
infernalis, *Gerrhonotus*, 114
 Gerrhonotus liocephalus, 114
 vidua, *Eutaenia*, 180, 181
inguinalis, *Prostherapis*, 71
inornata, *Cnemidophorus*, 101
inornatus, *Cnemidophorus*, 99
 Hemidactylus, 116
 Plestiodon, 131
insidior, *Bufo*, 34
 Bufo debilis, 34
insigniarum, *Eutaenia*, 180
insulana, *Ctenosaura*, 105
insulanus, *Pituophis catenifer*, 207
insularis, *Bothrops*, 187
 Lachesis, 187
insularum, *Leptodactylus*, 64
insulensis, *Dactyloperus*, 107
interbrachialis, *Sauromalus*, 133
intermedia, *Bachia*, 93
 nettingi, *Siren*, 25
 Salvadora, 211
 Salvadora intermedia, 211
 texana, *Siren*, 25
intermedius, *Crotalus intermedius*, 170
 Eleutherodactylus, 44
 Platyplacopus, 150
 Plethodon, 20
 Takydromus, 150
internasalis, *Sceloporus malachiticus*, 138
interorbitalis, *Syrrhophus*, 79
interparietalis, *Eumeces skiltonianus*, 113
inuitatus, *Callisaurus*, 94
 Callisaurus draconoides, 94
irregularis, *Aristelliger*, 92
 battersbyi, *Philothamnus*, 205
isabella, *Wenona*, 226
isabelleae, *Natrix valida*, 200
isozona, *Contia*, 168
 Sonora semiannulata, 168



- isthmica, *Caecilia*, 3
 ituriensis, *Chamaeleon*, 96
Ixalus japonicus, 63
ixbaac, *Lygosoma assatum*, 125
 Scincella cherriei, 125
jacksoni, *Plethodon*, 20
 Plethodon wehrlei, 20
jamaicensis, *Tropidophis maculatus*, 223,
 224
jamesi, *Phrynosoma*, 129
 Phrynosoma coronatum, 129
jamesoni kaimosae, *Dendraspis*, 172
 kaimosae, *Dendroaspis*, 172
jani, Arizona, 159
 Pituophis deppei, 159
Japalura mitsukurii, 118
 polygonata, 108
 splendida, 118
japonica, *Chelonia mydas*, 228
japonicus, *Ixalus*, 63
 Rhacophorus, 63
jarrovi *immucronatus*, *Sceloporus*, 136
 minor, *Sceloporus*, 143
 oberon, *Sceloporus*, 136
 Sceloporus, 136
 Sceloporus jarrovi, 136
 sugillatus, *Sceloporus*, 136
jeffersoni, *Desmognathus monticola*, 12
jeffersoniana, *Salamandra*, 24
jeffersonianum, *Ambystoma*, 5, 24
jimenezii, *Crotalus*, 170
johnstonii, *Chamaeleo*, 96
jordani clemsonae, *Plethodon*, 19
 metcalfi, *Plethodon*, 21
 shermani, *Plethodon*, 21
 Tetraprion, 80
jugans, *Eleutherodactylus*, 44, 63
 Rana, 74
juliae, *Anolis distichus*, 87
 Anolis dominicensis, 87
 Aporophis, 158
 Leimadophis, 158
junori, *Eleutherodactylus*, 44
kaimosae, *Dendraspis jamesoni*, 172
 Dendroaspis jamesoni, 172
Kalophrynus pleurostigma pleurostigma, 63
 stellatus, 63
Kaloula conjuncta stickeli, 63
 macroptica, 63
 rugifera, 63
kantingensis, *Staurois*, 79
karlschmidti, *Batrachuperus*, 8
 Eleutherodactylus, 45
kelloggi, *Bufo*, 35
 Bufo debilis, 35
keniensis, *Mabuya bayonii*, 126
kennerlyi, *Heterodon*, 185
 Heterodon nasicus, 185
kentuckyi, *Plethodon*, 20
khasiensis, *Takydromus*, 150
kilimensis, *Lygosoma*, 125
kinghorni, *Liasis amethystinus*, 195
kingii, *Gerrhonotus*, 109
Kinosternon abaxillare, 231
 acutum, 229
 bauri palmarum, 231
 cruentatum consors, 232
 flavescens flavescens, 233
 flavescens spooneri, 232
 herrerai, 232
 leucostomum, 229
 louisianae, 232
 murrayi, 232
 panamensis, 232
 subrubrum hippocrepis, 232
kirnia, *Tantilla*, 217
kirtlandii, *Natrix*, 209
 Regina, 209
klauberi, *Ensatina*, 13
 Ensatina eschscholtzii, 13
 Salvadora hexalepis, 211
 Sphaerodactylus, 147
kleinmanni, *Testudo*, 235
kohtaoensis, *Leiopisma*, 122
korros, *Ptyas korros*, 196
kotaensis, *Ichthyophis*, 3
laberculata, *Hyla bistincta*, 52
labialis, *Cnemidophorus*, 99
 Cystignathus, 40
 Leptodactylus, 40
lacerata, *Holbrookia*, 117
 Holbrookia lacerata, 117
 subcaudalis, *Holbrookia*, 117
lacertus, *Oreosaurus*, 127
Lachesis barbouri, 187
 insularis, 187
 muta stenophrys, 187
 stenophrys, 187
lachrymans, *Bufo haematiticus*, 34
lactea, *Stenorhina*, 214
 Stenorhina freminvillei, 214
lacustris, *Ambystoma*, 6
Laemanctus alticoronatus, 119
laevis, *Taricha*, 26
laeviventris, *Anolis*, 90
lagunensis, *Eumeces*, 111

- lambda*, *Trimorphodon*, 221
lamentor, *Bufo*, 35
Lampropeltis amaura, 187
 annulata, 187
 boylli conjuncta, 187
 calligaster calligaster, 203
 elapsoides virginiana, 188
 getulus californiae, 203, 204
 getulus conjuncta, 187
 getulus floridana, 188
 getulus floridanus, 188
 getulus niger, 204
 getulus splendida, 204
 getulus yumensis, 188
 multistriata, 188
 pyromelana pyromelana, 204
 pyromelana woodini, 188
 pyrrhomelaena celaenops, 189
 ruthveni, 189
 triangulum amaura, 187, 203
 triangulum annulata, 187
 triangulum celaenops, 189
 triangulum gentilis, 188, 204
 triangulum nelsoni, 189
 triangulum schmidti, 189
 triangulum sypila, 203
 triangulum temporalis, 188
 triangulum triangulum, 203
 zonata multicincta, 204
 zonata parvirubra, 189
 zonata pulchra, 189
Lamprosoma annulatum, 190
 episcopum, 190
lanceolata, *Mabuia*, 126
 Mabuia, 126
lanciformis, *Lithodytes*, 65
lanei, *Phyllodactylus*, 130
lansbergii venezuelensis, *Bothrops*, 161
larselli, *Plethodon vandykei*, 22
latastei, *Menobranthus lateralis*, 16
laterale boettgeri, *Leiopisma*, 122
lateralis fuliginosus, *Zamenis*, 227
 latastei, *Menobranthus*, 16
 Lygosoma, 145
 Lysoptychus, 125
 Scincus, 145
laticeps, *Hyla regilla*, 58
laticollaris, *Pliocercus elapoides*, 208
latifasciatus, *Micrurus*, 198
latiremisi, *Rana cantabrigensis*, 72
latirostris, *Philodryas*, 205
laticutatus okadae, *Eumeces*, 111
latistrata, *Dipsas gemmistrata*, 174
latistratus, *Imantodes*, 174
latitans, *Eurycea*, 14
lativittis, *Cnemidophorus*, 100
latodactylus, *Syrrhophus*, 79
latrans, *Eleutherodactylus*, 65
 Lithodytes, 65
lautus, *Crotalus gloydi*, 170
lavae, *Bolitoglossa*, 10
 Chiropterotriton, 10
lecontei, *Crotalus*, 171
 Rana, 74
leei, *Phyllodactylus*, 130
Leimadophis alleni, 190
 andreae nebulatus, 190
 chamissonis, 216
 funereus, 157
 juliae, 158
 nebulatus, 190
 parvifrons alleni, 190
 parvifrons lincolni, 190
 parvifrons niger, 190
 parvifrons rosamondae, 175
 parvifrons tortuganus, 191
 perfuscus, 196
 pocilogyrus, 196
 stahli, 190
 tortuganus, 191
Leiocephalus altavelensis, 119
 arenarius, 121
 caducus, 133
 carinatus aquarius, 119
 carinatus armouri, 119
 carinatus granti, 119
 carinatus punctatus, 119
 eremitus, 123
 inaguae, 120
 loxogrammus, 123
 macropus, 123
 macropus immaculatus, 120
 melanochlorus, 123
 personatus altavelensis, 119
 personatus aureus, 120
 personatus louisae, 120
 personatus lunatus, 120
 personatus mentalis, 120
 personatus scalaris, 121
 personatus vinculum, 121
 psammodromus, 121
 raviceps, 124
 varius, 124
 vinculum, 121
 virescens, 121
Leiopisma barbouri, 121
 caudaequinae, 121
 eunice, 121

Leiolopisma—Continued

- fasciolaris, 118
- forbesorum, 122
- kohtaoensis, 122
- laterale boettgeri, 122
- pranensis, 122
- smithi, 122
- lemniscatus, *Coluber* *obsoletus*, 166
- Tropidurus, 152
- lentiginosus pachycephalus, *Bufo*, 35
- leopardinus, *Helicops*, 184
- Lepidoblepharis barbouri, 122
- Lepidocephalus fasciatus, 191
- Lepidochelys olivacea, 227
- Lepidodactylus planicaudus, 122
- Lepidophyma flavimaculatum *tehuanae*, 123
 - smithii *oculor*, 122
 - smithii *tehuanae*, 123
 - tuxtlae, 123
- Lepidosternon floridanum, 123
- leprieurii, *Helicops*, 184
- leprosa, *Pseudoeurycea*, 25, 26
- leprosus, *Spelerpes*, 25, 26
- leprus, *Syrhophus*, 79
- leptocephala, *Eutainia*, 182
- Leptodactylus bolivianus, 64
 - caliginosus, 63
 - curtus, 63
 - darlingtoni, 44, 63
 - dominicensis, 64
 - gaigeae, 64
 - insularum, 64
 - labialis, 40
 - marmoratus, 65
 - melanonotus, 40
 - natalensis, 64
 - occidentalis, 64
 - ocellatus, 63, 64
 - pentadactylus mattogrossensis, 64
 - podicipinus, 64
 - poecilochilus, 40
 - serialis, 64
 - trivittatus, 65
 - ventrilineatus, 63
- Leptodeira annulata *pulchriceps*, 191
 - annulata *septentrionalis*, 174
 - annulata *taylori*, 191
 - ephippiata, 191
 - maculata, 192, 198
 - mystacina, 191
 - punctata, 192
 - septentrionalis *polysticta*, 191
 - splendida *ephippiata*, 191

Leptodeira—Continued

- yucatanensis *malleisi*, 191
- yucatanensis *yucatanensis*, 213
- Leptodira mystacina, 191
 - pacifica, 192
 - personata, 192
- Leptognathus argus, 192
 - pictiventris, 192
 - stratissima, 192
 - turgida, 192
- Leptopelis bequaerti, 65
- Leptopphis aeruginosus, 192
 - ahaetulla *coeruleodorsa*, 192
 - ahaetulla *copei*, 193
 - ahaetulla *praestans*, 219
 - coeruleodorsus, 192
 - copei, 193
 - majalis, 193
 - mexicanus *yucatanensis*, 193
 - nebulosus, 193
 - saturatus, 193
 - taeniata, 193
 - trifrenatus, 193
- leptosoma, *Aulacoplax*, 93
- Leptotyphlops albifrons *ater*, 194
 - albifrons *magnamaculata*, 194
 - ater, 194
 - bakewelli, 194
 - dulcis *dulcis*, 209
 - humilis *cahuilae*, 194
 - humilis *humilis*, 209
 - humilis *segregus*, 194
 - humilis *slevini*, 194
 - magnamaculata, 194
 - nasalis, 194
 - phenops *bakewelli*, 194
 - phenops *phenops*, 214
 - rufidorsum, 194
 - striatula, 195
- leptura, *Bothrops*, 161
- lepturum, *Amblystoma*, 5
- lermaensis, *Siredon*, 25
- lesueurii, *Hyla*, 77
- leucomelas, *Alsophis*, 175
 - danforthi, *Alsophis*, 157
 - Dromicus, 175
 - Himantodes, 185
 - Imantodes *cenchoa*, 185
 - sanctorum, *Alsophis*, 158
- leucophaeus *mariguanae*, *Anolis*, 88
- leucopus, *Batrachoseps*, 8
- leucostictus, *Hemiphyllodactylus*, 116
- leucostoma, *Ancistrodon piscivorus*, 220
- leucostomum, *Kinosternon*, 229

- leucostomus*, *Symphimus*, 216
levicollis, *Barisia*, 93
 Barissia, 93
 ciliaris, *Gerrhonotus*, 114
levigata, *Barisia gadovii*, 93
levis, *Uta*, 153
lewisi, *Anolis conspersus*, 86
 Anolis grahami, 86
 Eleutherodactylus, 45
 Necturus maculosus, 17
Liasis amethystinus kinghorni, 195
libertatis, *Liopeltis*, 196
Lichanura orcutti, 195
 roseofusca, 195
 roseofusca gracia, 195
 simplex, 195
 trivirgata, 195
lichenoides, *Siredon*, 25
licki, *Sceloporus*, 137
 Sceloporus orcutti, 137
lifanensis, *Stauroides*, 79
limbatus, *Phyllobates*, 69
 Sminthillus, 69
limbensis, *Eleutherodactylus schmidti*, 47
Limnomedusa schmidti, 65
lincolni, *Leimadophis parvifrons*, 190
lindheimeri, *Elaphe obsoleta*, 212
 Scotophis, 212
lineapiatus, *Liodytes alleni*, 195
linearis, *Urosaurus ornatus*, 154
 Uta ornata, 154
lineata acuta, *Toluca*, 202
 Salvadora, 211
 Toluca, 219
 Toluca lineata, 219
 varians, *Toluca*, 164
 wetmorei, *Toluca*, 219
lineatissimus, *Cnemidophorus*, 100
 Cnemidophorus deppi, 97
 Cnemidophorus deppii, 100
lineatopus coxi, *Anolis*, 83
 lynni, *Anolis*, 89
lineatulus, *Masticophis flagellum*, 196
 Sceloporus, 137
 Sceloporus magister, 137
lineatus concolor, *Conophis*, 167
 dunni, *Conophis*, 167
 similis, *Conophis*, 167
lineolata, *Eutaenia elegans*, 179
lineolateralis, *Sceloporus*, 137
lintoni, *Tantilla*, 217
 Tantillita, 217
Liocephalus eremitus, 123
 loxogrammus, 123
Liocephalus—Continued
 macropus, 123
 melanochlorus, 123
 raviceps, 124
 varius, 124
liocephalus infernalis, *Gerrhonotus*, 114
 ophiurus, *Gerrhonotus*, 115
Liodytes alleni lineapiatus, 195
Liolaemus hatcheri, 124
 walkeri, 124
Lioninia vermiformis, 195
lionotus, *Anolis*, 90
Liopeltis libertatis, 196
 semicarinata, 178
Liophis albiceps, 209
 perfuscus, 196
 poecilogyrus, 202
 poecilopogon, 210
 subfasciatus, 196
lippiens, *Sympholis*, 216
Lithodytes bransfordii, 65
 diastema, 65
 gulosus, 65
 lanciformis, 65
 latrans, 65
 megacephalus, 66
 melanostictus, 66
 pelviculus, 66
 podiciferus, 66
 ranoides, 66
 rhodopsis, 66
littoralis, *Malaclemmys*, 232
 Malaclemmys terrapin, 232
lividus, *Anolis*, 89
 Anolis bimaculatus, 89
Liyila rugulosa, 66
lödingi, *Necturus*, 17
 Pituophis, 207
 Pituophis melanoleucus, 207
loloensis, *Stauroides*, 79
longicauda pernix, *Eurycea*, 14
longicaudus, *Ophisaurus attenuatus*, 127
longifrenis, *Mesopeltis*, 198
 Sibynomorphus, 198
longinasus, *Bufo*, 35
longipes, *Rana*, 74, 76
 Sceloporus, 137
 Sceloporus occidentalis, 137, 143
longirostris, *Eumeces*, 131
 Plestiodon, 131
loquax, *Hyla*, 55
Loricata, 236
louisae, *Liocephalus personatus*, 120
louisiana, *Kinosternon*, 232

- louisianensis, *Necturus*, 17
 Necturus maculosus, 17
 loyii ventralis, *Calamaria*, 162
 loxogrammus, *Leiocephalus*, 123
 Leiocephalus, 123
 lubricus, *Hydrops*, 186
 luciae, *Anolis*, 89
 Anolis trinitatis, 89
 lucifer, *Crotalus*, 171
 lugubris, *Aneides lugubris*, 19
 lugubrosus, *Bufo*, 35
 lumboltzii, *Calamaria*, 161
 lunatus, *Leiocephalus personatus*, 120
 lundelli, *Sceloporus lundelli*, 137
 lunulata, *Tropidodipsas*, 222
 Luperosaurus macgregori, 124
 lurida, *Salamandra*, 24
 luteolus, *Eleutherodactylus*, 45
 lutosus, *Crotalus confluentus*, 169
 Crotalus viridis, 169
 lutzii, *Eupsophus*, 80
 Thoropa, 80
 lutzorum, *Cochranella*, 39
 Lycodon osmanhilli, 196
 subcinctus sealei, 196
 Lycognathus lyrophanes, 196
 Lygodactylus capensis mossambica, 124
 grotei grotei, 124
 manni, 124
 Lygophis flavifrenatus, 196
 Lygosaurus pellopleurus, 124
 sowerbyi, 124
 Lygosoma assatum ixbaac, 125
 cherriei stuarti, 125
 incertum, 125
 kilimensis, 125
 lateralis, 145
 (*Sphenomorphus*) *variegatum stickeli*,
 125
 lymani, *Eleutherodactylus*, 45
 lynni, *Anolis lineatopus*, 89
 Eleutherodactylus, 45
 lyoni, *Phrynonax*, 206
 lyrophanes, *Lycognathus*, 196
 Trimorphodon, 196
 Lyssoptychus lateralis, 125
 Lystrophis dorbignyi, 185
 Lytorhynchus sinai, 196
 mabecii, *Ambystoma*, 6
 mabouya, *Mabuya mabouya*, 125
 mabouya pergravis, *Mabuya*, 126
 Mabua alliacea, 125
 dorsivittata, 125
 fulgida, 126
 lanceolata, 126
 Mabuya bayonii keniensis, 126
 chanleri, 126
 delalandii, 113
 deserticola, 126
 dorsivittata, 125
 frenata, 110
 lanceolata, 126
 mabouya mabouya, 125
 mabouya pergravis, 126
 nigropalmata, 126
 pergravis, 126
 spilonota, 126
 macedougalli, *Sceloporus*, 137
 macgregori, *Luperosaurus*, 124
 macrocephala, *Rana macrodon*, 74
 Rana magna, 74
 macrodactyla, *Amblystoma*, 5
 macrodactylum, *Ambystoma macrodacty-*
 lum, 5, 7
 macrodon macrocephala, *Rana*, 74
 visayanus, *Rana*, 74
 macrolepis, *Sceloporus poinsetti*, 141
 Sceloporus poinsettii, 141
 Sphaerodactylus, 147
 macroptica, *Kaloula*, 63
 macropus immaculatus, *Leiocephalus*, 120
 Leiocephalus, 123
 Leiocephalus, 123
 macrospilota, *Malaclemmys*, 233
 Malaclemmys terrapin, 233
 macrostemma, *Eutaenia*, 180
 macrostomus, *Dipsas*, 213
 Sibynomorphus, 213
 macrotyimpanum, *Tomodactylus*, 81
 maculata dickersonae, *Holbrookia*, 116
 Leptodeira, 192, 198
 perspicua, *Holbrookia*, 117
 pulchra, *Holbrookia*, 117
 thermophila, *Holbrookia*, 118
 maculatus haetianus, *Tropidophis*, 225
 Hypopachus, 62
 jamaicensis, *Tropidophis*, 223, 224
 Megalops, 198
 maculiventris, *Eupemphix*, 49
 maculosus beyeri, *Necturus*, 16
 lewisi, *Necturus*, 17
 louisianensis, *Necturus*, 17
 Necturus maculosus, 16
 madagascariense, *Sanzinia*, 226
 Xiphosoma, 226
 madagascariensis abbotti, *Phelsuma*, 128
 Acrantophis madagascariensis, 205
 Pelophilus, 205

- magister himaculosus*, *Sceloporus*, 137
 lineatulus, *Sceloporus*, 137
 Phyllodactylus, 130
 rufidorsum, *Sceloporus*, 142
 Sceloporus, 137
 Sceloporus magister, 134, 137
 uniformis, *Sceloporus*, 138
 zosteromus, *Sceloporus*, 145
magna macrocephala, *Rana*, 74
 Rana, 75
 Rana magna, 75
 visayanus, *Rana*, 74
Magnadigita flavimembris, 17
 franklini, 17
 nigroflavescens, 10
magnamaculata, *Leptotyphlops*, 194
 Leptotyphlops albifrons, 194
magnus, *Phyllodactylus*, 130
majalis, *Leptopis*, 193
major, *Eurycea bislineata*, 13
 Eurypholis, 184
 Trimorphodon, 222
malachiticus, *Sceloporus*, 138
 Sceloporus malachiticus, 138
 internasalis, *Sceloporus*, 138
 smaragdinus, *Sceloporus*, 142
 taeniocnemis, *Sceloporus*, 143
Malachylodes guttilatus, 66
Malaclemmys littoralis, 232
 macrospilota, 233
 terrapin littoralis, 232
 terrapin macrospilota, 233
 terrapin tequesta, 233
Malacoclemmys oculifera, 233
malcolmi, *Microhyla*, 67
malleisi, *Leptodeira yucatanensis*, 191
managuae, *Eumeces*, 111
Manculus quadridigitatus, 16
 quadridigitatus paludicolus, 16
 quadridigitatus uvidus, 16
mandensis, *Typhlops*, 224
manni, *Amphisbaena*, 84
 Lygodactylus, 124
 Typhlops, 224
mannophorus, *Uta*, 153
Manolepis putnami, 220
mantzorum, *Stauroids*, 74
marciana, *Eutainia*, 182
marcianus, *Thamnophis*, 182
maresinensis, *Hydrophis ornatus*, 185
margaritiferus fistulosus, *Drymobius*, 176
marginatus amamiensis, *Eumeces*, 111
 Eumeces, 131
 Plestiodon, 131
marianae, *Hyla*, 55
marianus, *Euspondylus*, 132
 Prionodactylus, 132
Mariguana agassizii, 85
mariguanae, *Anolis leucophaeus*, 88
 Sphaerodactylus, 147
marmorata, *Clemmys marmorata*, 230
 Emys, 230
 Rana, 75
marmoratus, *Cnemidophorus*, 100
 Cnemidophorus tigris, 100
 Leptodactylus, 65
 Sceloporus variabilis, 134
 Triturus marmoratus, 27
marmoreus, *Bufo*, 31
martini, *Hyla microcephala*, 55
martyris, *Cnemidophorus*, 100
 Cnemidophorus tigris, 100
mascareniensis, *Rana*, 75
massarti, *Atelopus moreirae*, 30
Masticophis anthonyi, 160
 aurigulus, 176
 bilineatus, 161
 flagellum lineatulus, 196
 flagellum piceus, 160, 227
 flagellum variolosus, 197
 melanolomus, 197
 mentovarius, 160
 ornatus, 197
 ruthveni, 197
 schottii, 197
 taeniatus australis, 197
 taeniatus ornatus, 197
 taeniatus schottii, 197
 taeniatus taeniatus, 193
mattea, *Cyclura*, 106
matto grossensis, *Leptodactylus pentadactylus*, 64
matudai, *Anolis*, 89
 Elaphe flavirufa, 176
 Eleutherodactylus, 45
mauritanica, *Tarentola*, 131
 Testudo, 235
mavortia, *Ambystoma*, 7
 Ambystoma tigrinum, 5, 6, 7, 25
maxillabrochus, *Thorius*, 27
maximus, *Cnemidophorus*, 100
mayensis, *Anolis sagrei*, 91
 Micrurus affinis, 198
mazatlanensis nayaritensis, *Bufo*, 36
mcclellani, *Pituophis*, 207
mearnsi, *Calamaria*, 161
 Rana, 75

- mearnsi—Continued
 Streptosaurus, 153
 Uta, 153
- megacephalus, Lithodytes, 66
- megalepidurus, Sceloporus, 139
- megalops, Eleutherodactylus, 45
 Eutaenia, 180
 Thamnophis eques, 180
- Megalops maculatus, 198
- megapoda, Rana, 75
- megapodia, Hyla, 55
- Megophrys abbotti, 67
 minor, 67
 omeimontis, 67
- Mehelya chanleri chanleri, 213
- melanocephalus, Elaps, 177
- melanochloris, Bufo, 35
- melanochlorus, Leiocephalus, 123
 Liocephalus, 123
- melanogaster canescens, Thamnophis, 218
 Sceloporus, 139
 Sceloporus torquatus, 139
- melanolencus catenifer, Pituophis, 207
 lödingi, Pituophis, 207
 ruthveni, Pituophis, 207
 sayi, Pituophis, 207
 vertebralis, Pituophis, 208
- melanolomus, Dryadophis melanolomus, 197
 Masticophis, 197
 stuarti, Dryadophis, 175
 tehuanae, Dryadophis, 175
- melanomma, Hyla, 55
- melanomolga, Bolitoglossa, 10
 Pseudoeurycea, 10
- melanonotus, Leptodactylus, 40
- melanopleurus, Sphenomorphus, 149
- melanorhinus calligaster, Sceloporus, 139
- melanostethus, Cnemidophorus, 100
- melanostictum, Ambystoma tigrinum, 8
- melanostictus, Eleutherodactylus, 66
 Lithodytes, 66
- melanotropis, Dendrophidium, 173
 Drymobius, 173
- melanurus, Bothrops, 220
 Drymarchon corais, 213
 Spilotes, 213
- melas, Chironius, 184
 Herpetodryas, 184
- Menobanchus lateralis latastei, 16
 punctatus, 16
- mentalis, Leiocephalus personatus, 120
- mentovarius centralis, Coluber (Masticophis), 165
 Masticophis, 160
- meridionalis, Crotalus willardi, 172
- merremii semiaureus, Oreomorphus, 204
- merriami annulatus, Sceloporus, 139
 Sceloporus, 139
 Sceloporus merriami, 139
- mertensi, Bufo ictericus, 34
- Mesopeltis longifrenis, 198
 sanniolus, 198
- mestrei, Anolis, 89
- metcalfi, Plethodon, 21
 Plethodon jordani, 21
- metallica, Anadia metallica, 96
- metallicus, Chalcidolepis, 96
- mexicana, Terrapene mexicana, 235
- mexicanus, Pituophis, 207
 Spilotes pullatus, 214
 Xenodon, 226
 yucatanensis, Leptophis, 193
- miaria, Deirochelys reticularia, 230
- Microbatrachylus bransfordii, 65
 imitator, 67
 montanus, 67
 oaxacae, 67
 polyptychus, 61
- microcephala martini, Hyla, 55
 sartori, Hyla, 55
- microeximia, Hyla, 55
- Microhyla annamensis, 67
 berdmorei, 67
 butleri, 67, 68
 grahami, 67
 malcolmi, 67
 okinavensis, 68
 olivacea, 48
 pictiventris, 48
 sowerbyi, 68
 undulata, 68
 usta usta, 48
- microlepidopus, Cnemidophorus, 100
- microlepidotus, Sceloporus grammicus, 136
- Microlophus spinulosus, 126
- microscaphus, Bufo, 35
- microstomus, Typhlops, 224
- microtis, Cystignathus, 40
- microtus, Anolis, 89
- Micruroides euryxanthus, 177
- Micrurus affinis mayensis, 198
 affinis stantoni, 198
 clarki, 198
 diastema proximans, 198
 distans distans, 177
 ehippifer, 177
 fulvius, 198
 fulvius tenere, 177

- Micrurus—Continued
 latifasciatus, 198
 nigrocinctus alleni, 198
 nigrocinctus mosquitensis, 198
 nigrocinctus nigrocinctus, 177
 nigrocinctus ovandoensis, 199
 olssoni, 199
 micrurus, Erythrolamprus mimus, 178
 milesi, Eleutherodactylus, 45
 miliaria, Hyla, 62
 miliarius, Hysiboas, 62
 milleri, Anolis, 89
 Hyla, 56
 milneana, Rana grisea, 73
 minus micrurus, Erythrolamprus, 178
 mindanensis, Draco, 109
 miniata, Tantilla, 217
 miniator, Tantilla, 217
 minor, Goniocichelys, 230
 Megophrys, 67
 Sceloporus jarrovi, 143
 Sceloporus torquatus, 143
 Sternotherus minor, 230
 minshanicus, Bufo, 35
 minuta, Hyla, 57
 miquihuanus, Crotalus pricei, 172
 Crotalus triseriatus, 172
 misaminius, Tropidophorus, 151
 mitchellii pyrhus, Crotalus, 163, 170
 stephensi, Crotalus, 169
 mitis, Contia, 162, 168
 mitsukurii, Japalura, 118
 mobilensis, Pseudemys floridana, 235
 modesta, Acrodytes, 29
 Barisia, 132
 Phrynohyas, 29
 modestum, Phrynosoma, 129
 modestus, Pterogaster, 132
 Syrhophus, 79
 moesta, Hyla, 58
 Hyla punctariola, 58
 mojavensis, Salvadora hexalepis, 211
 mokeson pictigaster, Agkistrodon, 157
 molesworthi, Celestus cruscus, 95
 molitor, Hyla, 58
 molossus, Crotalus, 171
 Crotalus molossus, 171
 estebanensis, Crotalus, 171
 monensis, Anolis, 89
 monksiae, Bufo, 36
 montanus, Ascaphus truei, 29
 floridanus, Pseudotriton, 23
 Microbatrachylus, 67
 montanus—Continued
 Pseudotriton, 23
 Pseudotriton montanus, 23
 montezumae concolor, Rana, 75
 Rana, 75
 monticola, Barisia, 114, 115
 Desmognathus, 12
 Desmognathus monticola, 12
 Gastrotheca, 49
 Hyla, 58
 Hyla punctariola, 58
 jeffersoni, Desmognathus, 12
 monticolus, Gerrhonotus, 115
 Montorana ahli, 68
 moreirae massarti, Atelopus, 30
 moreleti, Bolitoglossa, 10
 mormon, Coluber, 165
 Coluber constrictor, 160, 165
 mosquitensis, Micrurus nigrocinctus, 198
 mossambica, Lygodactylus capensis, 124
 mucosus, Ptyas, 193
 mucronatus aureolus, Sceloporus, 140
 Sceloporus mucronatus, 144
 Sceloporus torquatus, 144
 mühlenbergii, Clemmys, 230
 multicarinatus scincicaudus, Gerrhonotus,
 109
 webbii, Elgaria, 115
 multicepta, Lampropeltis zonata, 204
 Opisthotropis, 204
 multiceptus, Ophibolus getulus, 204
 multidentata, Chiropterotriton, 18
 Oedipus, 18
 multilineata, Hyla, 56
 Hyla bischoffi, 56
 multilineatus, Eumeces, 112
 multimaculatus, Amplorhinus, 174
 multiplicata, Eurycea, 26
 griseogaster, Eurycea, 14
 Gymnopsis multiplicata, 4
 proxima, Gymnopsis, 4
 multiplicatus, Scaphiopus, 78
 Scaphiopus hammondii, 78
 Spelerpes, 26
 multiscutatus, Cnemidophorus tessellatus,
 102
 Cnemidophorus, tigris, 96, 102
 multispinis, Ctenosaura, 105
 multisquamatus, Sphenomorphus, 149
 multistriata, Lampropeltis, 188
 multivirgatus, Eumeces, 111, 131
 muralis, Platydictylus, 131
 muricolor, Hyla, 56

- murrayi*, Kinosternon, 232
muta stenophrys, Lachesis, 187
mutabilis, Coluber, 165
 Elaphe triaspis, 165
mutilatus, Peropus, 107
mydas japonica, Chelonia, 228
mystacina, Leptodeira, 191
 Leptodira, 191
myurus, Callisaurus draconoides, 95
 Callisaurus ventralis, 95
nannodes, Anolis, 90
Nanorana pleskei, 68
narisovalis, Thorius, 27
nasale, Catostoma, 162
nasalis, Causus, 163
 Chiropterotriton, 18
 Geophis, 162
 Leptotyphlops, 194
 Oedipus, 18
nasica, Hyla, 56
nasicus gloydi, Heterodon, 185
 Heterodon nasicus, 185
 kennerlyi, Heterodon, 185
nasutus, Hylodes, 61
 Eleutherodactylus, 61
 Tomodon, 220
natalensis, Leptodactylus, 64
Natrix compressicauda bivittata, 199
 cyclopion floridana, 199
 erythrogaster, 165
 erythrogaster flavigaster, 199
 erythrogaster neglecta, 199
 erythrogaster transversa, 201, 222, 223
 fasciata pictiventris, 199
 fasciata pleuralis, 200
 grahamii, 208
 harteri, 200
 kirtlandii, 209
 olivacea uluguruensis, 200
 piscator flavipunctata, 158
 rhombifera blanchardi, 200
 rhombifera rhombifera, 201, 223
 rhombifera werleri, 200
 rigida deltae, 200
 sipedon clarkii, 208
 sipedon compressicauda, 199, 201, 223
 sipedon pictiventris, 199
 sipedon pleuralis, 200
 sipedon sipedon, 201
 valida, 222
 valida isabelleae, 200
 valida valida, 209, 223
nyaritentis, Bufo, 36
 Bufo mazatlanensis, 36
 nebulatus, Leimadophis, 190
 Leimadophis andreae, 190
nebulifer, Bufo, 33, 36
nebulosa, Hyla, 52, 56
nebulosum, Ambystoma, 7
 Ambystoma tigrinum, 7
nebulosus, Leptophis, 193
 Syrrophus, 80
Necturus alabamensis, 16
 beyeri, 16
 lödingi, 17
 louisianensis, 17
 maculosus beyeri, 16
 maculosus lewisi, 17
 maculosus louisianensis, 17
 maculosus maculosus, 16
 punctatus, 16
neglecta, Natrix erythrogaster, 199
neglectus, Cycloramphus, 40
nelsoni, Anolis, 90
 Bufo boreas, 32
 Lampropeltis triangulum, 189
 Phrynosoma, 129
 Sceloporus, 140
 Terrapene, 235
 Urosaurus bicarinatus, 154
 Uta, 154
neomexicanus, Cnemidophorus, 100
 Plethodon, 21
Neoseps reynoldsi, 126
neotenes, Eurycea, 14
nerus, Typhlotriton, 28
Nerodia agassizii, 201
 compressicauda, 201
 couchii, 201
 holbrookii, 201
 woodhousii, 201
nettingi, Siren intermedia, 25
neuwiedii diporus, Bothrops, 161
 Pseudoboa, 208
nicholsi richardi, Alsophis, 158
 Sphaerodactylus, 147
nietoi, Ninia diademata, 201
niger, Lampropeltis getulus, 204
 Leimadophis parvifrons, 190
 Ophibolus getulus, 204
 Telmatobius, 80
nigerrima, Cyclura, 106, 107
nigrescens, Thamnophis elegans, 218
nigricans, Rana, 74, 76
nigricauda, Uta, 154
nigricaudus, Urosaurus, 154
nigriceps fumiceps, Tantilla, 212, 217

- nigridius*, *Gonatodes*, 116
nigrilatus, *Adelphicos veraepacis*, 157
nigripes, *Hyla*, 56
nigrita, *Emoa*, 109
 Emoia, 109
 feriarum, *Pseudacris*, 50
nigriventris, *Hylaplesia*, 61
nigrocinctus alleni, *Micrurus*, 198
 Elaps, 177
 Micrurus nigrocinctus, 177
 mosquitensis, *Micrurus*, 198
 ovandoensis, *Micrurus*, 199
nigroflavescens, *Bolitoglossa*, 10
 Magnadigita, 10
nigroflavus, *Ichthyophis*, 3
nigroluteus, *Tretanorhinus*, 220
 Tretanorhinus nigroluteus, 220
nigromaculata, *Bolitoglossa*, 10
 Pseudoeurycea, 10
 Rana, 75
nigropalmata, *Mabuya*, 126
nigrum, *Ambystoma rosaceum*, 7
Ninia diademata, 201
 diademata diademata, 201
 diademata nietoi, 201
 psephota, 162
 sebae immaculata, 201
 sebae tessellata, 202
 tessellata, 202
nitida garambensis, *Eremias*, 110
nobilis, *Elgaria*, 109
noctivaga, *Arizona elegans*, 159
nongkhorensis, *Philautus*, 69
notata scoparia, *Uma*, 152
 Uma, 152
 Uma notata, 152
Nothopsis rugosus, 202
nothus, *Scaphiodontophis*, 212
nubicola, *Eleutherodactylus*, 45
 Phyllobates, 70
nuchalis, *Basiliscus (Cristasaura)*, 94
 Clemmys, 230
nummifer picadoi, *Trimeresurus*, 220
 veraecrucis, *Bothrops*, 161
nuntius, *Crotalus viridis*, 172
oaxacae, *Microbatrachylus*, 67
oaxacus, *Crotalus basiliscus*, 169
oberon, *Sceloporus jarrovii*, 136
obesus, *Euphryne*, 113
 Proctoporus, 132
 Sauromalus obesus, 113
 townsendi, *Sauromalus*, 133
obscura, *Eutainia sirtalis*, 183
obscurum, *Amblystoma*, 5
obsoleta bairdi, *Elaphe*, 164
 Elaphe obsoleta, 166, 212
 lindheimeri, *Elaphe*, 212
 quadrivittata, *Elaphe*, 177
obsoletum, *Plestiodon*, 131
obsoletus, *Cnemidophorus gularis gularis*, 98
 Eumeces, 131
 lemniscatus, *Coluber*, 166
occidentalis, *Agama rueppelli*, 82
 Arizona elegans, 159
 Boiga dendrophila, 161
 Bolitoglossa, 10
 Bufo, 36
 Leptodactylus, 64
 longipes, *Sceloporus*, 137, 143
occipitalis annulata, *Chionactis*, 190
occulor, *Lepidophyma smithii*, 122
ocellata, *Eutaenia cyrtopsis*, 179
ocellatus, *Leptodactylus*, 63, 64
ochoterenaee, *Eumeces*, 112
 Sceloporus, 140
ochrocephala, *Caecilia*, 3
ochrophaeus carolinensis, *Desmognathus*, 11, 12
ochrorhyncha deserticola, *Hypsiglena*, 186
ochrorhyncha, *Hypsiglena torquata*, 186
ochrorhynchus, *Hypsiglena*, 186
ocoeae, *Desmognathus*, 13
octolineatus, *Cnemidophorus*, 101
oculatus, *Anolis*, 156
 Xiphosurus, 156
oculifera, *Graptomys*, 233
 Malacoclemmys, 233
odonnelli, *Bolitoglossa*, 18
 Oedipus, 18
odoratus, *Sternotherus*, 233
Oedipina collaris, 25
 ignea, 17
Oedipus bromeliacia, 17
 flavimembris, 17
 flaviventris, 17
 franklini, 17
 gadovii, 18
 goebeli, 18
 helmrichi, 18
 multidentata, 18
 nasalis, 18
 odonnellii, 18
 robertsi, 18
 smithi, 18
 striatulus, 19
 townsendi, 19

- Ogmia acutus*, 202
okadae, *Eumeces latiscutatus*, 111
okinavensis, *Microhyla*, 68
olfersii, *Chlorosoma*, 205
olfersioides, *Eupemphix*, 49
oldi, *Takydromus tachydromoides*, 150
Oligodon annamensis, 202
 subgriseum, 202
oligoporus, *Cnemidophorus sexlineatus*, 102
 Sceloporus, 140
 Sceloporus horridus, 140
Oligosoma gemmingeri, 127
oligozona, *Gymnopsis*, 4
oligozonus, *Siphonops*, 4
olivacea, *Lepidochelys*, 227
 Microhyla, 48
 uluguruensis, *Natrix*, 200
olivaceus, *Gerrhonotus*, 115
oliveri, *Imantodes splendidus*, 187
olporus, *Sceloporus variabilis*, 144
olssoni, *Micrurus*, 199
olympicus, *Ranodon*, 23
 Rhyacotriton, 23
omeimontis, *Megophrys*, 67
 Polypedates, 70
 Rhacophorus, 70
onca, *Rana*, 76
onchiopsis, *Cyclura*, 107
Opheodrys aestivus, 193
 vernalis blanchardi, 202
Opheomorphys doliatus caesius, 202
Ophibolus boylii, 203
 clericus, 203
 doliatus collaris, 203
 doliatus parallelus, 203
 doliatus sypilus, 203, 204
 evansii, 203
 gentilis, 203, 204
 getulus eiseni, 204
 getulus multincinctus, 204
 getulus niger, 204
 pyromelanus, 204
 splendidus, 204
Ophisaurus attenuatus longicaudus, 127
 compressus, 127
 ventralis compressus, 127
ophiurus, *Gerrhonotus*, 115
 Gerrhonotus liocephalus, 115
Opisthotropis balteata, 204
 multicincta, 204
orbiculare durangoensis, *Phrynosoma*, 129
orcutti, *Eleutherodactylus*, 43, 46
 Lichanura, 195
 licki, *Sceloporus*, 137
orcutti—Continued
 Sceloporus, 140
 Sceloporus orcutti, 140
ordinaria, *Ambystoma*, 7
ordinarium, *Ambystoma*, 7
ordinoides errans, *Thamnophis*, 218
 Thamnophis, 182
oreganus, *Crotalus viridis*, 171
oregonensis, *Ensatina eschscholtzii*, 15
 Heredia, 15
Oreomorphus merremii semiaureus, 204
Oreosaurus anomalus, 127
 lacertus, 127
orientalis, *Eleutherodactylus atkinsi*, 41
 Emoia boettgeri, 109
orizabensis, *Spelerpes*, 26
ornata, *Cistudo*, 229
 linearis, *Uta*, 154
 schmidti, *Uta*, 154
 Terrapene ornata, 229
 Uta, 154
ornatissima, *Saccodeira*, 133
 Tapaya, 150
ornatissimum, *Phrynosoma douglassii*, 150
ornatus caeruleus, *Sceloporus*, 141
 caeruleus, *Urosaurus*, 153
 Crotalus, 171
 linearis, *Urosaurus*, 154
 maresinensis, *Hydrophis*, 185
 Masticophis, 197
 Masticophis taeniatus, 197
 Sceloporus, 140
 Sceloporus ornatus, 140
 schmidti, *Urosaurus*, 154
 symmetricus, *Urosaurus*, 155
 Urosaurus ornatus, 154
orphila, *Hyla* (*Sphoerohyla*), 56
 Hyla aurantiaca, 56
osceola, *Chelydra*, 228
 Chelydra serpentina, 228
oshimai, *Zaocys*, 227
oshimensis, *Eumeces*, 112
osmanhilli, *Lycodon*, 196
ouachitae, *Plethodon*, 21
ouachitensis, *Graptemys pseudogeographica*, 231
ovalis, *Elachistocleis*, 62
ovandoensis, *Micrurus nigrocinctus*, 199
ovis, *Hypopachus*, 62
owstoni, *Polypedates*, 71
 Rhacophorus, 71
Oxybelis brevirostris, 176
oxycephala, *Elaphe*, 157
oxylophus, *Anolis*, 90

- Oxyrhopus baileyi*, 164
 fitzingeri frizzelli, 205
 trigeminus, 205
Oxyrrhopus doliatus semicinctus, 205
Ozotheca tristycha, 233
pachycephalus, *Bufo lentiginosus*, 35
pachyderma, *Hyla*, 57
pachypus, *Anolis*, 90
pachyura, *Contia*, 168
 fulviceps, *Rhadinaea*, 210
 Rhadinaea pachyura, 168
pacifica, *Leptodira*, 192
pacificum, *Hemidactylum*, 15
pacificus, *Batrachoseps*, 8, 15
 Hoplodactylus, 118
 Sphaerodactylus, 147
 Tropidurus, 151
palearis, *Ctenosaura*, 106
 Enyaliosaurus, 106
palirostrata, *Cadea*, 94
pallens, *Hyla*, 57
pallidus, *Eumeces septentrionalis*, 113
palmarum, *Kinosternon bauri*, 231
palmeri, *Gerrhonotus coeruleus*, 115
 Gerrhonotus scincicauda, 115
 Uta, 155
paludicola, *Acris gryllus*, 29
 Coluber constrictor, 165
Paludicola frenata, 68
paludicolus, *Manculus quadridigitatus*, 16
panamensis, *Cerathyla*, 38
 Kinosternon, 232
 Ungaliophis, 225
panchlora, *Ameiva*, 82
pantherinus, *Coryphodon*, 168
Pantodactylus bivittatus, 128
 schreibersii, 128
pantoni, *Eleutherodactylus*, 46
paracnemis, *Bufo*, 36
parallelus, *Ophibolus doliatus*, 203
pardalis androsi, *Tropidophis*, 223
 barbouri, *Tropidophis*, 224
 bucculentus, *Tropidophus*, 225
 canus, *Tropidophis*, 225
 stejnegeri, *Tropidophis*, 223, 224
parishi, *Ialtris*, 187
parkeri, *Altirana*, 29
 Bachia, 93
 Ctenosaura, 106
 Hyla, 57
paroticum, *Amblystoma*, 5
partelloi, *Tropidophorus*, 151
parva, *Uta*, 155
parviauriculatus, *Eumeces*, 112
parvifrons alleni, *Leimadophis*, 190
 lincolni, *Leimadophis*, 190
 niger, *Leimadophis*, 190
 rosamondae, *Dromicus*, 175
 rosamondae, *Leimadophis*, 175
 tortuganus, *Leimadophis*, 191
Parvimolge townsendi, 19
parvirubra, *Lampropeltis zonata*, 189
parvulus, *Eumeces*, 112
 Zachaeus, 38
parvus, *Eleutherodactylus*, 42
 Sceloporus parvus, 141
parvus, *Sceloporus*, 141
patagoniensis, *Callirhinus*, 162
paucidentulus, *Ichthyophis*, 3
pauciporus, *Cnemidophorus sexlineatus*, 102
paucisulcus, *Ichthyophis*, 3
pearsei, *Hypopachus*, 62
pectinata, *Ctenosaura*, 105, 106
pekinensis, *Eumeces*, 112
pelagica, *Heteronota*, 116
pelagicus, *Gymnodactylus*, 116
pelopleurus, *Lygosaurus*, 124
Pelophilus madagascariensis, 205
pelviculus, *Lithodytes*, 66
peninsularis, *Coleonyx variegatus*, 103
 Pseudemys floridana, 234
pennatulus, *Thorius*, 27
pentadactylus mattogrossensis, *Leptodactylus*, 64
perfuscus, *Leimadophis*, 196
 Liophis, 196
pergravis, *Mabuya*, 126
 Mabuya mabouya, 126
pericensis, *Amphisbaena*, 85
perkinsi, *Phyllorhynchus decurtatus*, 206
perlacea, *Elaphe*, 176
perlaevis, *Cystignathus*, 40
pernix, *Eurycea longicauda*, 14
Peropus mutilatus, 107
perplexus, *Bufo*, 36
 Cnemidophorus, 101
personata, *Leptodira*, 192
personatus altavelensis, *Leiocephalus*, 119
 aureus, *Leiocephalus*, 120
 louisae, *Leiocephalus*, 120
 lunatus, *Leiocephalus*, 120
 mentalis, *Leiocephalus*, 120
 scalaris, *Leiocephalus*, 121
 vinculum, *Leiocephalus*, 121
perspicua, *Holbrookia maculata*, 117
perstriatus, *Diemictylus viridescens*, 28
 Triturus, 28

- petasatus*, Pharyngodon, 68
 Tripriion, 68
petola, Pseudoboa, 205
petropolitana, Cochranella, 39
Petrosaurus thalassinus, 155
phaeocrypta, *Hyla versicolor*, 60
phaeota cyanosticta, *Hyla*, 57
 Hyla, 57
 Hyla phaeota, 57
phaescens, *Elaphe flavirufa*, 176
 Elaphe triaspis, 176
Pharyngodon petasatus, 68
Phelsuma abbotti, 128
 madagascariensis abbotti, 128
phenax, *Eutainia*, 182
 Thamnophis phenax, 182
phenops bakewelli, *Leptotyphlops*, 194
 Leptotyphlops phenops, 214
 Stenostoma, 214
Philautus gryllus, 68
 hansena, 68
 nongkhorensis, 69
 woodi, 69
philbyi, *Scincus*, 145
Philodryas latirostris, 205
 schottii, 162
Philothamnus irregularis battersbyi, 205
Phimothyrax hexalepis, 205
phlebodes, *Hyla*, 57
Phoxophrys grahami, 128
phrenitica, *Tantilla*, 217
Phrynixalus anulatus, 69
Phrynocephalus erythrurus, 128
Phrynohyas hebes, 78
 modesta, 29
Phrynonax lyoni, 206
Phrynosoma asio, 128
 cerroense, 128
 cornutum, 129
 coronatum frontale, 129
 coronatum jamesi, 129
 ditmarsii, 128
 douglasi pygmaea, 128
 douglasi brachycercum, 128
 douglasi douglasi, 128
 douglasi ornatissimum, 150
 douglasi brevisrostris, 150
 douglasi hernandesi, 150
 goodei, 129
 jamesi, 129
 modestum, 129
 nelsoni, 129
 orbiculare durangoensis, 129
Phrynosoma—Continued
 planiceps, 129
 platyrhinus, 129
 platyrhinus calidiarum, 92
 platyrhinus goodei, 129
 platyrhinus platyrhinus, 129
 regale, 130
 solaris, 130
Phyllobates auratus, 69
 boulengeri, 71
 brunneus, 49
 cystignathoides, 69
 flotator, 69
 hylaeformis, 69
 limbatus, 69
 nubicola, 70
 sylvatica, 70
Phyllodactylus homolepidurus, 130
 lanei, 130
 leei, 130
 magister, 130
 magnus, 130
 pulcher, 130
 spatulatus, 130
 unctus, 108
 wirshingi, 131
Phyllomedusa appendiculata, 70
 azurea, 70
 guttata, 70
 hypochondrialis, 70
Phyllorhynchus browni, 206
 browni browni, 206
 decurtatus perkinsi, 206
Phyllosira flavescens, 206
Physalaemus signiferus, 49
piaropicola, *Diemictylus viridescens*, 13
picadoi, *Bothrops*, 220
 Chiropterotriton, 26
 Spelerpes, 26
 Trimeresurus nummifer, 220
piceivittis, *Coniophanes*, 166
 Coniophanes piceivittis, 166
piceum, *Bascanium*, 160
piceus, *Masticophis flagellum*, 160, 227
pickeli, *Hyla*, 57
pickeringi, *Eutainia*, 182
 Thamnophis sirtalis, 182
picta bellii, *Chrysemys*, 229
 dorsalis, *Chrysemys*, 229
pictigaster, *Agkistrodon contortrix*, 157
 Agkistrodon mokeson, 157
pictipes, *Hyla*, 58
 Hyla punctariola, 58
pictiventre, *Engystoma*, 48

- pictiventris*, *Dipsas*, 192
 Leptognathus, 192
 Microhyla, 48
 Natrix fasciata, 199
 Natrix sipedon, 199
pictus, *Bufo*, 36
pinarensis, *Eleutherodactylus*, 46
pinchonii, *Batrachuperus*, 24
pinchoti, *Anolis*, 90
pingi, *Scutigera*, 78
pintoi, *Crossodactylodes*, 39
piperata, *Holbrookia propinqua*, 117
pipiens berlandieri, *Rana*, 72
 brachycephala, *Rana*, 76, 77
 fisheri, *Rana*, 73
pipilans, *Syrrophus*, 80
piscator flavipunctata, *Natrix*, 158
piscivorus leucostoma, *Ancistrodon*, 220
Pituophis annectens, 206
 catenifer annectens, 206
 catenifer bimarisi, 206
 catenifer deserticola, 206
 catenifer fuliginatus, 207
 catenifer insulanus, 207
 catenifer sayi, 164
 depei jani, 159
 lödingi, 207
 mcclellani, 207
 melanoleucus catenifer, 207
 melanoleucus lödingi, 207
 melanoleucus ruthveni, 207
 melanoleucus sayi, 207
 melanoleucus vertebralis, 208
 mexicanus, 207
 wilkesii, 207
Pityophis haematois, 208
plagosus, *Conophis pulcher*, 167
planicaudus, *Lepidodactylus*, 122
planiceps, *Phrynosoma*, 129
planicola, *Hyla* (*Sphoerohyla*), 57
 Hyla aurantiaca, 57
plasticus, *Callisaurus*, 94
platineum, *Amblystoma*, 5
platycephala, *Hylella*, 61
Platydictylus muralis, 131
Platymantis aculeodactylus, 70
Platypeltis agassizii, 233
Platyplacopus dorsalis, 149
 intermedius, 150
platyrhinus calidiarum, *Phrynosoma*, 92
 goodei, *Phrynosoma*, 129
 Heterodon, 184, 185
 platyrhinus, *Phrynosoma*, 129
Platythyra flavescens, 233
Plectrohyla sagorum, 70
plei exsul, *Ameiva*, 83
pleskei, *Nanorana*, 68
Plestiodon anthracinus, 131
 egregius, 131
 inornatus, 131
 longirostris, 131
 marginatus, 131
 obsoletum, 131
 septentrionalis, 132
 skiltonianum, 132
 tetragrammus, 132
Plethodon cinereus angusticlavius, 19
 cinereus cinereus, 20, 24
 cinereus dorsalis, 19
 clemsonae, 19
 crassulus, 19
 dixi, 19
 dorsalis angusticlavius, 19
 dorsalis dorsalis, 19
 dunni, 19
 glutinosus chlorobryonis, 19
 glutinosus glutinosus, 19, 20
 hubrichti, 20
 huldae, 20
 idahoensis, 20
 intermedius, 20
 jacksoni, 20
 jordani clemsonae, 19
 jordani metcalfi, 21
 jordani shermani, 21
 kentuckyi, 20
 metcalfi, 21
 neomexicanus, 21
 ouachitae, 21
 richmondi, 21
 richmondi popei, 21
 richmondi richmondi, 21
 shermani, 21
 vandykei idahoensis, 20
 vandykei larselli, 22
 wehrlei dixi, 19
 wehrlei jacksoni, 20
 welleri, 22
 welleri ventromaculatum, 22
 welleri welleri, 22
 yonahlossee, 22
Plethopsis wrighti, 22
pleuralis, *Natrix fasciata*, 200
 Natrix sipedon, 200
Pleurodema bibroni, 68
pleurostictus batangensis, *Rhacophorus*, 77

- pleurostigma, *Kalophrynus pleurostigma*, 63
- Plica stejnegeri*, 132
- Pliocerus bicolor*, 208
dimidiatus, 208
elapoides laticollaris, 208
- plioporus*, *Sceloporus serrifer*, 142
- Plistodon sumichrasti*, 132
- plumbea*, *Wenona*, 226
- plumifrons*, *Basiliscus*, 94
- plutonia*, *Eutaenia vagrans*, 181
- pluvialis*, *Eumeces*, 112
Eumeces anthracinus, 112
- podarga*, *Ameiva undulata*, 83
- podiciferus*, *Eleutherodactylus*, 66
Lithodytes, 66
- podicipinus*, *Leptodactylus*, 64
- poecilochilus*, *Cystignathus*, 40
Leptodactylus, 40
- pocilogyrus*, *Leimadophis*, 196
Liophis, 202
- pocilonotus chrysobronchus*, *Pseustes*, 213
polylepis, *Pseustes*, 206
Pseustes pocilonotus, 222
- pocilopogon*, *Liophis*, 210
Rhadinaea, 210
- poinsetti macrolepis*, *Sceloporus*, 141
polylepis, *Sceloporus*, 141
- poinsettii macrolepis*, *Sceloporus*, 141
polylepis, *Sceloporus*, 141
Sceloporus, 141
Sceloporus poinsettii, 141
- politus*, *Bufo*, 36
- polops*, *Ameiva*, 83
- polygonata*, *Japalura*, 108
- polygonatum*, *Diploderma*, 108
- polylepis*, *Pseustes pocilonotus*, 206
Sceloporus poinsettii, 141
Sceloporus poinsettii, 141
- Polypedates omeimontis*, 70
owstoni, 71
viridis, 71
- polyptychus*, *Hylodes*, 61
Microbatrachylus, 61
- polysticta*, *Leptodeira septentrionalis*, 191
- polystictus*, *Crotalus*, 170
- pomarii*, *Hoplodactylus*, 118
- poncensis*, *Anolis*, 90
- poolei*, *Eleutherodactylus*, 46
- popei*, *Plethodon richmondi*, 21
Triturus ensicaudus, 28
- porphyriticus duryi*, *Gyrinophilus*, 23
inagnoscus, *Gyrinophilus*, 15
- postinguinale*, *Cinosternum*, 229
- praeocularis*, *Typhlops*, 224
- praesignis*, *Ameiva*, 101
Cnemidophorus, 101
- praestans*, *Leptophis ahaetulla*, 219
Thrasops, 219
- pranensis*, *Leiopisma*, 122
- prasinus*, *Dendrophis*, 173
- pratensis*, *Hispaniolus*, 116
- pretiosa*, *Rana*, 76
Rana pretiosa, 76
- prezygus*, *Sceloporus*, 141
- pricei miquihuanus*, *Crotalus*, 172
- principis*, *Elgaria*, 109
Gerrhonotus coeruleus, 109
- Prionodactylus marianus*, 132
- Proctoporus obesus*, 132
- propinqua*, *Holbrookia*, 117
Holbrookia propinqua, 117
piperata, *Holbrookia*, 117
- proserpine*, *Amblystoma*, 5
- Prostherapis boulengeri*, 71
femoralis, 71
inguinalis, 71
- proterops*, *Coniophanes*, 166
Coniophanes fissidens, 166
- Protopipa carvalhoi*, 71
- proxima*, *Gymnopsis multiplicata*, 4
- proximans*, *Micrurus diastema*, 198
- proximus*, *Siphonops*, 4
Thamnophis sauritus, 181
- psammodromus*, *Leiocephalus*, 121
- psephota*, *Ninia*, 162
- psephotum*, *Catostoma*, 162
- Pseudacris clarkii*, 50
nigrita feriarum, 50
streckeri illinoensis, 71
- Pseudemys alabamensis*, 233
decorata, 234
elonae, 234
felis, 234
floridana concinna, 234
floridana hoyi, 235
floridana mobilensis, 235
floridana peninsularis, 234
floridana texana, 234
scripta gaigeae, 234
stejnegeri, 234
texana, 234
vioscana, 235
- Pseudis bolbodactyla*, 71
- Pseudoboa newwedii*, 208
robinsoni, 208
petola, 205

- Pseudobranchius striatus axanthus*, 22
striatus spheniscus, 22
Pseudoeurycea brunnata, 23
cephalica cephalica, 26
cephalica rubrimembris, 9
cochranae, 9
firscheini, 23
gadovii, 18
galeanae, 9
goebeli, 18
leprosa, 25, 26
melanomolga, 10
nigromaculata, 10
robertsi, 18
smithi, 18
unguidentis, 11
werleri, 23
Pseudoficimia frontalis, 219
pseudogeographica ouachitensis, *Graptemys*,
 231
sabinensis, *Graptemys*, 231
versa, *Graptemys*, 231
Pseudophryne blanchardi, 71
Pseudotriton duryi, 23
montanus, 23
montanus floridanus, 23
montanus montanus, 23
ruber schenki, 26
ruber sticticeps, 26
ruber vioscai, 23
Pseustes poecilonotus chrysobronchus, 213
poecilonotus poecilonotus, 222
poecilonotus polylepsis, 206
Pterogasterus modestus, 132
Ptyas korros korros, 196
mucosus, 193
Ptychemys hoyi, 235
Ptychohyla adipovertris, 72
publia, *Ficimia*, 183
Ficimia publia, 183
taylori, *Ficimia*, 183
pugnax, *Toxicophis*, 220
pulcher, *Conophis*, 167
Conophis pulcher, 167
Phyllodactylus, 130
plagosus, *Conophis*, 167
similis, *Conophis*, 167
pulchra, *Graptemys*, 231
Holbrookia, 117
Holbrookia maculata, 117
Lampropeltis zonata, 189
pulchriceps, *Leptodeira annulata*, 191
pulchrilatus, *Eutaenia*, 180
pullatus auribundus, *Spilotes*, 214
mexicanus, *Spilotes*, 214
pulmonaris, *Thorius*, 27
puma, *Hyla*, 58
punctariola moesta, *Hyla*, 58
monticola, *Hyla*, 58
pictipes, *Hyla*, 58
punctata, *Hyla*, 58
Leptodeira, 192
punctatus, *Bufo*, 31, 37
arnyi, *Diadophis*, 173
Leiocephalus carinatus, 119
Menobranchus, 16
Necturus, 16
stictogenys, *Diadophis*, 174
punctilinealis, *Cnemidophorus*, 101
punctulata, *Ahaetulla*, 173
purpurescens, *Anolis*, 90
putnami, *Manolepis*, 220
pygmaea, *Phrynosoma douglassi*, 128
pygmaeus, *Bufo*, 37
pyrocephalus, *Sceloporus*, 141
pyromelana, *Lampropeltis pyromelana*, 204
woodini, *Lampropeltis*, 188
pyromelanus, *Ophibolus*, 204
pyrrha, *Caudisona*, 163
pyrrhomelaena celaenops, *Lampropeltis*, 189
pyrrhus, *Crotalus mitchellii*, 163, 170
Python curtus brongersmai, 208
quadrigitatus, *Manculus*, 16
paludicolus, *Manculus*, 16
uvidus, *Manculus*, 16
quadrilineata, *Ameiva*, 84, 101
quadrilineatus, *Cnemidophorus*, 101
Eumeces, 112
quadrivirgatus, *Eumeces*, 112
visoninus, *Adelphicos*, 211
quadrivittata deckerti, *Elaphe*, 177
Elaphe obsoleta, 177
quadruplex, *Trimorphodon biscutatus*, 221
quagulus, *Anolis*, 90
quiquelineata, *Rhadinaea*, 210
quiquevittata, *Hyla*, 58
quiquevittatus, *Coniophanes*, 186
Rabdosoma crassicaudatum, 208
radix haydenii, *Thamnophis*, 180, 182
Eutaenia, 182
Thamnophis radix, 182
twiningi, *Eutaenia*, 180
radula, *Gaigeia*, 114
raineyi, *Alsophis vudii*, 158
Rana areolata, 72
aerolata aesopus, 72
areolata areolata, 72

Rana—Continued

- areolata capito*, 73
areolata sevosae, 76
aurora, 72
aurora aurora, 72
aurora cascadae, 73
aurora draytonii, 73, 74, 76
berlandieri, 72
boyllii, 72
boyllii boyllii, 72
cantabrigensis evittata, 72
cantabrigensis latiremisi, 72
capito, 73
capito stertens, 73
cascadae, 73
chaochiaoensis, 73
draytonii, 73
fisheri, 73
grisea milneana, 73
grylio, 74
gryllus, 74
jugans, 74
lecontei, 74
longipes, 74, 76
macrodon macrocephala, 74
macrodon visayanus, 74
magna, 75
magna macrocephala, 74
magna magna, 75
magna visayanus, 74
marmorata, 75
mascareniensis, 75
mearnsi, 75
megapoda, 75
montezumae, 75
montezumae concolor, 75
nigricans, 74, 76
nigromaculata, 75
onca, 76
pipiens berlandieri, 72
pipiens brachycephala, 76, 77
pipiens fisheri, 73
pretiosa, 76
pretiosa pretiosa, 76
septentrionalis, 76
sevosae, 76
sierramadrensis, 76
signata, 77
sinuata, 76
sylvatica cantabrigensis, 72
sylvatica cherokiana, 76
tsushimensis, 77
virescens brachycephala, 77

Rana—Continued

- woodworthi*, 77
yakani, 77
raniceps, Hyla, 55, 62
Hypsiboas, 62
Ranodon olympicus, 23
Ranoidea flavoviridis, 77
resplendens, 77
ranoides, Eleutherodactylus, 66
Lithodytes, 66
rathbuni, Typhlomolge, 28
raveni, Calamaria, 161
raviceps, Leiocephalus, 124
Leiocephalus, 124
ravus, Crotalus, 171
Sistrurus, 171
redimita, Colorhogia, 164
redimitus, Geagras, 183
regale, Phrynosoma, 130
regalis arizonae, Diadophis, 173
blanchardi, Diadophis, 173
Diadophis, 173
Diadophis regalis, 173
regilla, Hyla, 52, 56, 58
regilla laticeps, Hyla, 58
Regina clarkii, 208
grahamii, 208
kirtlandii, 209
valida, 209
remivaga, Caretta, 227
Rena dulcis, 209
humilis, 209
 Reptilia, 81
resimus, Causus, 163
resplendens, Rana, 77
retardatus, Hynobius, 15
reticularia miaria, Deirochelys, 230
reticulata, Diaglena, 40
Xantusia riversiana, 156
reticulatus, Crotaphytus, 104
reynoldsi, Neoseps, 126
Rhabdosoma guttatum, 209
Rhacophorus baluensis, 77
bambusicola, 77
chenfui, 77
japonicus, 63
omeimontis, 70
owstoni, 71
pleurostictus batangensis, 77
viridis, 71
Rhadinaea albiceps, 209
calligaster, 168
crassa, 209

- Rhadinaea—Continued
 decorata, 174
 forbesi, 209
 fulviceps, 210
 fulvivittis, 210
 hesperia, 210
 hesperia hesperia, 210
 hesperia hesperioides, 210
 pachyura fulviceps, 210
 pachyura pachyura, 168
 poecilopogon, 210
 quinquelineata, 210
 semiaurea, 204
 vittata, 210
- Rhadinella schistosa, 210
- Rhamnophis aethiopissa elgonensis, 210
- Rhegnops visoninus, 211
- Rhineura floridana, 123
- rhinomegas, Coluber, 166
- rhodopis, Eleutherodactylus, 66
 Lithodytes, 66
- rhodostictus, Callisaurus, 95
 Callisaurus draconoides, 95
- rhomبifer, Tropidonotus, 223
- rhomبifera blanchardi, Natrix, 200
 Natrix rhombifera, 201, 223
 werleri, Natrix, 200
- Rhyacosiredon, rivularis, 24
 zempoalaensis, 24
- Rhyacotriton olympicus, 23
- richardi, Alsophis richolsi, 158
- richardii, Anolis richardii, 92
 griseus, Anolis, 88
- richardsoni gossei, Sphaerodactylus, 147
- richmondi, Eleutherodactylus, 46
 Plethodon, 21
 Plethodon richmondi, 21
 popei, Plethodon, 21
- rickardsi, Hyla, 59
- ricordii casparii, Eleutherodactylus, 42
 rogersi, Eleutherodactylus, 46
- rigida deltae, Natrix, 200
- rileyi, Cyclura, 107
- Riopa singha, 132
- riversiana reticulata, Xantusia, 156
 Xantusia, 156
- rivicola, Eurycea bislineata, 14
- rivularis, Rhyacosiredon, 24
- robertmertensi, Hyla, 59
- robertsi, Oepidus, 18
 Pseudoeurycea, 18
- robertsorum, Hyla, 59
- robinsoni, Pseudoboa, 208
- robusta, Aepidia, 157
- rogersi, Eleutherodactylus ricordii, 46
- roquet aeneus, Anolis, 86
 cinereus, Anolis, 85, 91
- rosacea, Elaphe guttata, 166
- rosaceum nigrum, Ambystoma, 7
 sonoraensis, Ambystoma, 7
- rosaceus, Coluber, 166
- rosamondae, Dromicus parvifrons, 175
 Leimadophis parvifrons, 175
- roseofusca gracia, Lichanura, 195
 Lichanura, 195
- roseus, Zachaenus, 81
- rostellatus, Typhlops, 224
- rostrata, Caretta, 228
- rozellae, Celestus, 95
 Hyla, 59
 Thamnophis, 218
- ruber, Crotalus adamanteus, 168
 Crotalus ruber, 168, 169
 schenki, Pseudotriton, 26
 schenki, Spelerpes, 26
 sticticeps, Pseudotriton, 26
 sticticeps, Spelerpes, 26
 vioscae, Pseudotriton, 23
- rubescens, Bufo, 37
- rubida, Geoemyda rubida, 228
- rubidus, Chelopus, 228
 Cnemidophorus, 103
 Cnemidophorus tessellatus, 102
 Cnemidophorus tigris, 102
 Drymarchon corais, 175
- rubra, Tantilla, 218
- rubricata, Tantilla wilcoxi, 218
- rubrimaculatus, Syrrhophus, 80
- rubrimembris, Bolitoglossa cephalica, 9
 Pseudoeurycea cephalica, 9
- rucillensis, Eleutherodactylus schmidti, 47
- ruepelli occidentalis, Agama, 82
 septentrionalis, Agama, 82
- rufidorsum, Leptotyphlops, 194
 Sceloporus, 142
 Sceloporus magister, 142
- rufipunctatum, Chilopoma, 164
- rufipunctatus, Thamnophis, 164
- rufopunctata, Uma, 152
- rufozonatum walli, Dinodon, 174
- rufus, Bufo, 37
- rugifera, Kaloula, 63
- rugosus, Celestus, 96
 Eleutherodactylus, 65, 66
 Nothopsis, 202
- rugulosa, Liyla, 66
- rugulosus, Eleutherodactylus, 66

- ruthveni*, *Anolis*, 90
 Eupemphix, 49
 Lampropeltis, 189
 Masticophis, 197
 Pituophis melanoleucus, 207
rutilorus, *Eutainia*, 182
ruttyi, *Alsophis angulifer*, 157
sabanus, *Anolis*, 91
 Anolis bimaculatus, 91
 Sphaerodactylus, 148
 Sphenomorphus, 149
sabinensis, *Graptemys pseudogeographica*, 231
Saccodeira ornattissima, 133
sacki huico, *Cnemidophorus*, 101
sackii angusticeps, *Cnemidophorus*, 96
 Cnemidophorus sackii, 97
 gularis, *Cnemidophorus*, 98, 99
 huico, *Cnemidophorus*, 101
 scalaris, *Cnemidophorus*, 98
 semifasciatus, *Cnemidophorus*, 101
 stictogrammus, *Cnemidophorus*, 101
sagorum, *Plectrohyla*, 70
sagrei mayensis, *Anolis*, 91
 stejnegeri, *Anolis*, 91
Salamandra agilis, 24
 granulata, 24
 jeffersoniana, 24
 lurida, 24
Salamandrella sinensis, 24
Salientia, 29
saltator, *Eleutherodactylus*, 46
Salvadora bogerti, 211
 grahamiae, 211
 hexalepis celeris, 211
 hexalepis hexalepis, 205
 hexalepis klauberi, 211
 hexalepis mojavensis, 211
 intermedia, 211
 intermedia intermedia, 211
 lineata, 211
salvini, *Staurotypus*, 230
samanensis, *Sphaerodactylus*, 148
sanborni, *Hyla*, 59
sanctae-catharinae, *Craspedoglossa*, 39
sanctorum, *Alsophis*, 158
 Alsophis leucomelas, 158
sandersi, *Hyla versicolor*, 60
sanniolus, *Dipsas*, 193
 Mesopeltis, 198
Sanzinia madagascariense, 226
sartori, *Hyla microcephala*, 55
Sator angustus, 133
 grandaevus, 133
 saturatus, *Leptophis*, 193
Sauria, 81
sauritus chalceus, *Thamnophis*, 182
 proximus, *Thamnophis*, 181
Sauromalus ater, 133
 hispidus, 133
 interbrachialis, 133
 obesus obesus, 113
 obesus townsendi, 133
 townsendi, 133
 varius, 133
sawayae, *Zachaeus*, 81
sayi, *Pituophis catenifer*, 164
 Pituophis melanoleucus, 207
scalaris, *Cnemidophorus gularis*, 98
 Cnemidophorus sackii, 98
 Leiocephalus personatus, 121
 slevini, *Sceloporus*, 142
 unicanthalis, *Sceloporus*, 142
scandax, *Uromacer*, 225
Scaphiodontophis albonuchalis, 212
 carpicinctus, 212
 nothus, 212
Scaphiopus albus, 78
 couchii, 78
 dugesi, 78
 hammondii hammondii, 78
 hammondii multiplicatus, 78
 holbrookii albus, 78
 multiplicatus, 78
 varius, 78
Scartiscus caducus, 133
Sceloporus aeneus bicanthalis, 133
 aeneus subniger, 134
 boulengeri, 134
 chrysostrictus, 134
 clarkii, 134
 clarkii boulengeri, 134
 clarkii clarkii, 134
 cochranae, 134
 couchii, 125, 134
 cozumelae, 134
 cyanogenys, 143
 delicatissimus, 134
 disparilis, 135
 edwardtaylori, 135
 elongatus, 135
 ferraripezei, 135
 floridanus, 135
 formosus formosus, 145
 formosus scitulus, 135
 goldmani, 135
 graciosus, 136
 graciosus graciosus, 136

Sceloporus—Continued

graciosus vandenburgianus, 144
grammicus disparilis, 135
grammicus microlepidotus, 136
heterurus, 136
horridus albiventris, 136
horridus oligoporus, 140
jarrovii, 136
jarrovii immucronatus, 136
jarrovii jarrovii, 136
jarrovii minor, 143
jarrovii oberon, 136
jarrovii sugillatus, 136
licki, 137
lineatulus, 137
lineolateralis, 137
longipes, 137
lundelli lundelli, 137
macdougalli, 137
magister, 137
magister bimaculosus, 137
magister lineatulus, 137
magister magister, 134, 137
magister rufidorsum, 142
magister uniformis, 138
magister zosteromus, 145
malachiticus, 138
malachiticus internasalis, 138
malachiticus malachiticus, 138
malachiticus smaragdinus, 142
malachiticus taeniocnemis, 143
megalepidurus, 139
melanogaster, 139
melanorhinus calligaster, 139
merriami, 139
merriami annulatus, 139
merriami merriami, 139
mucronatus aureolus, 140
mucronatus mucronatus, 144
nelsoni, 140
occidentalis longipes, 137, 143
ochoterenae, 140
oligoporus, 140
orcutti, 140
orcutti licki, 137
orcutti orcutti, 140
ornatus, 140
ornatus caeruleus, 141
ornatus ornatus, 140
parvus, 141
parvus parvus, 141
poinsetti macrolepis, 141
poinsetti polylepis, 141
poinsettii, 141

Sceloporus—Continued

poinsettii macrolepis, 141
poinsettii poinsettii, 141
poinsettii polylepis, 141
prezygus, 141
pyrocephalus, 141
rufidorsum, 142
scalaris unicanthalis, 142
scalaris slevini, 142
serrifer, 142
serrifer plioporus, 142
serrifer serrifer, 142
siniferus, 142
siniferus cupreus, 134
siniferus siniferus, 142
smaragdinus, 142, 143
spinosus apicalis, 143
squamosus, 143
stejnegeri, 143
taeniocnemis, 143
thayeri, 143
torquatus cyanogenys, 143
torquatus melanogaster, 139
torquatus minor, 143
torquatus mucronatus, 144
torquatus torquatus, 135
tristichus, 144
undulatus elongatus, 135
undulatus erythrocheilus, 144
undulatus hyacinthinus, 143
undulatus tristichus, 144
undulatus undulatus, 135
uniformis, 144
vandenburgianus, 144
variabilis marmoratus, 134
variabilis olloporus, 144
variabilis smithi, 144
viviparus, 145
woodi, 145
zosteromus, 145
schenki, *Pseudotriton ruber*, 26
Spelerpes ruber, 26
schistosa, *Rhadinella*, 210
schmidti, *Calamaria*, 161
Lampropeltis triangulum, 189
limbensis, *Eleutherodactylus*, 47
Limnomedusa, 65
rucillensis, *Eleutherodactylus*, 47
Scutiger, 78
Thorius, 27
Urosaurus ornatus, 154
Uta ornata, 154

- schotti*, *Masticophis*, 197
 Masticophis taeniatus, 197
schottii, *Philodryas*, 162
schreibersii, *Pantodactylus*, 128
Scincella cherriei ixbaac, 125
 cherriei stuarti, 125
 gemmingeri forbesorum, 122
 gemmingeri gemmingeri, 127
scincicauda palmeri, *Gerrhonotus*, 115
scincicaudus, *Gerrhonotus multicarinatus*, 109
Scincus lateralis, 145
 philbyi, 145
scitulus, *Sceloporus formosus*, 135
scelopops apaporiensis, *Caiman*, 236
Scolecormorphus uluguruensis, 4
Scolecophis fumiceps, 212
 scytalinus, 212
Scolecosaurus trinitatis, 145
Scoliophis atlanticus, 212
scoparia, *Uma*, 152
 Uma notata, 152
Scotophis calligaster, 166
 confinis, 212
 lindheimeri, 212
 vulpinus, 213
scripta gaigeae, *Pseudemys*, 234
Scutigera pingi, 78
 schmidti, 78
scutigera, *Trachycephalus*, 81
scytalina, *Clelia clelia*, 160
scytalinus, *Scolecophis*, 212
Scytopsis boulengeri, 78
 hebes, 78
sealei, *Lycodon subcinctus*, 196
sebae immaculata, *Ninia*, 201
 tessellatus, *Ninia*, 202
segregus, *Leptotyphlops humilis*, 194
sellatus, *Coluber guttatus*, 165
semiannulata isozona, *Sonora*, 168
 Sonora, 213
 Sonora semiannulata, 213
semiaurea, *Rhadinaea*, 204
semiaureus, *Oreomorphus merremii*, 204
semicarinata, *Liopeltis*, 178
semicarinatum, *Dinodon*, 191
semicarinatus, *Eurypholis*, 178
semicinctus, *Oxyrrhopus doliatus*, 205
semifasciata, *Eutaenia sirtalis*, 181
semifasciatus, *Cnemidophorus gularis*, 99
 Cnemidophorus sackii, 99, 101
semilineatus, *Bascanium*, 161
semirutus, *Trimorphodon biscutatus*, 221
septemvittatus, *Cnemidophorus*, 101
 septentrionalis, *Agama rueppelli*, 82
 brunnea, *Hyla*, 81
 Dipsas, 174
 Eumeces, 132
 Leptodeira annulata, 174
 pallidus, *Eumeces*, 113
 Plestiodon, 132
 polysticta, *Leptodeira*, 191
 Rana, 76
 serialis, *Leptodactylus*, 64
 sericeus, *Cnemidophorus gularis*, 99
 Serpentes, 157
 serpentina osceola, *Chelydra*, 228
 serrifer pliopus, *Sceloporus*, 142
 Sceloporus, 142
 Sceloporus serrifer, 142
 severus, *Claudius*, 230
 sevosa, *Rana*, 76
 Rana areolata, 76
 sexlineatus, *Cnemidophorus*, 206
 oligoporus, *Cnemidophorus*, 102
 pauciporus, *Cnemidophorus*, 102
 sextaeniata, *Eremias*, 110
 Eremias spekii, 110
 shermani, *Plethodon*, 21
 Plethodon jordani, 21
 shihi, *Hynobius*, 15
 shrevei, *Audantia*, 93
 Sibon annulatum yucatanensis, 213
 sibonius, *Alsophis*, 158
 Sibynomorphus longifrenis, 198
 macrostomus, 213
 turgidus, 192
 sierramadrensis, *Rana*, 76
 sigma, *Constrictor constrictor*, 167
 signata, *Rana*, 77
 signiferus, *Physalaemus*, 49
 silus, *Crotalus willardi*, 172
 Crotaphytus, 104
 Gambelia wislizenii, 104
 similis, *Conophis lineatus*, 167
 Conophis pulcher, 167
 Ctenosaura similis, 105
 Hyla, 59
 Simocephalus chanleri, 213
 simplex, *Lichanura*, 195
 simus, *Siphonops*, 4
 sinai, *Lytorhynchus*, 196
 sinensis, *Salamandrella*, 24
 singha, *Riopa*, 132
 siniferus cupreus, *Sceloporus*, 134
 Sceloporus, 142
 Sceloporus siniferus, 142
 sinistra, *Ameiva undulata*, 83

- sinuata*, *Rana*, 76
sipedon clarkii, *Natrix*, 208
 compressicauda, *Natrix*, 199, 201, 223
 Natrix sipedon, 201
 pictiventris, *Natrix*, 199
 pleuralis, *Natrix*, 200
Siphonops oligozonus, 4
 proximus, 4
 simus, 4
Siredon dumerilii, 24
 gracilis, 25
 lermaensis, 25
 lichenoides, 25
Siren intermedia nettingi, 25
 intermedia texana, 25
sirtalis obscura, *Eutainia*, 183
 pickeringi, *Thamnophis*, 182
 semifasciata, *Eutaenia*, 181
 tetrataenia, *Eutaenia*, 181
 tetrataenia, *Thamnophis*, 181
 Thamnophis sirtalis, 181, 183
 trilineata, *Eutainia*, 183
 trilineata, *Thamnophis*, 183
Sistrurus catenatus edwardsii, 168
 ravus, 171
skiltonianum, *Plestiodon*, 132
skiltonianus brevipes, *Eumeces*, 113
 interparietalis, *Eumeces*, 113
 skiltonianus, *Eumeces*, 112, 132
slateri, *Ambystoma tigrinum*, 8
slevini, *Leptotyphlops humilis*, 194
 Sceloporus scalaris, 142
smaragdina, *Hyla*, 60
smaragdinus, *Sceloporus*, 142, 143
 Sceloporus malachiticus, 142
Smilisca baudinii baudinii, 56, 60
Sminthillus limbatus, 69
smithi, *Bufo*, 37
 Dipsas, 174
 Leiolopisma, 122
 Oedipus, 18
 Pseudoeurycea, 18
 Sceloporus variabilis, 144
 Syrhophus, 80
smithii occulor, *Lepidophyma*, 122
 tehuanae, *Lepidophyma*, 123
solaris, *Phrynosoma*, 130
Sonora episcopa episcopa, 190
 semiannulata, 213
 semiannulata isozona, 168
 semiannulata semiannulata, 213
sonoraensis, *Ambystoma rosaceum*, 7
sonoriensis, *Coleonyx variegatus*, 103
 Dipsosaurus dorsalis, 108
 sowerbyi, *Lygosaurus*, 124
 Microhyla, 68
spatulatus, *Eleutherodactylus*, 47
 Phyllodactylus, 130
Spea stagnalis, 78
speciosus, *Bufo*, 37
 Bufo compactilis, 37
spekii sextaeniata, *Eremias*, 110
spelaeus, *Typhlotriton*, 28
Spelerpes bilineatus borealis, 25
 collaris, 25
 gibbicaudus, 25, 26
 leprosus, 25, 26
 multiplicatus, 26
 orizabensis, 26
 picadoi, 26
 ruber schenki, 26
 ruber sticticeps, 26
Sphaerodactylus alopex, 146
 argus bartschi, 146
 argus henriquesi, 146
 bartschi, 146
 caicosensis, 146
 cinereus, 146
 corticulus, 146
 difficilis, 148
 exsul, 146
 gaigeae, 146
 glaucus, 146
 glaucus glaucus, 146
 grandisquamis, 147
 homolepis, 147
 klauberi, 147
 macrolepis, 147
 mariguanae, 147
 nicholsi, 147
 pacificus, 147
 richardsoni gossei, 147
 sabanus, 148
 samanensis, 148
 stejnegeri, 148
 townsendi, 148
spheniscus, *Pseudobranchius striatus*, 22
Sphenomorphus atrigularis, 148
 deignani, 148
 helena, 148
 indicus formosensis, 148
 melanopleurus, 149
 multisquamatus, 149
 sabanus, 149
 steerei, 149
 undosus, 118
 variegatus stickeli, 125

- Sphenosoma durum*, 151
 hughi, 149
Sphoenohyla habra, 79
spilonota, Mabuya, 126
spilorhipis, *Anolis tropidonotus*, 91
Spilotes chrysobronchus, 213
 erebennus, 213
 melanurus, 213
 pullatus auribundus, 214
 pullatus mexicanus, 214
spinifer asper, *Trionyx*, 227, 233
 emoryi, *Trionyx*, 227
 hartwegi, *Trionyx*, 227
spinifera hartwegi, *Amyda*, 227
spinosus apicalis, *Sceloporus*, 143
 Urosaurus bicarinatus, 152
spinulosus, *Bufo*, 35
 Microlophus, 126
 Tropidurus, 126
splendida ephippiata, *Leptodeira*, 191
 Japalura, 118
 Lampropeltis getulus, 204
splendidus, *Callisaurus*, 95
 Callisaurus draconoides, 95
 oliveri, *Imantodes*, 187
 Ophibolus, 204
spooneri, *Kinosternon flavescens*, 232
spurrelli certus, *Atelopus*, 30
squalirostris, *Hyla*, 60
Squamata, 81
squamata, *Eretmochelys imbricata*, 228
 Uta, 155
squamosus, *Sceloporus*, 143
stagnalis, *Spea*, 78
stahli, *Leimadophis*, 190
stansburiana elegans, *Uta*, 153
 hesperis, *Uta*, 155
 Uta, 155
 Uta stansburiana, 155
stantoni, *Micrurus affinus*, 198
staufferi, *Hyla*, 52, 60
Stauroids kangtingensis, 79
 lifanensis, 79
 loloensis, 79
 mantzorum, 74
Staurotypus salvinii, 230
stebbinsi, *Ambystoma tigrinum*, 8
steerei, *Sphenomorphus*, 149
steindachneri, *Diploglossus*, 108
stejnegeri, *Ambystoma*, 7
 Anolis, 91
 Anolis sagrei, 91
 Borborocoetes, 31
 Craspedoglossa, 31
 stejnegeri—Continued
 Crotalus, 171
 Cyclura, 107
 Hynobius, 16
 Plica, 132
 Pseudemys, 234
 Sceloporus, 143
 Sphaerodactylus, 148
 Trimeresurus, 220
 Tropidophis pardalis, 223, 224
 Typhlops, 225
stejnegerianus, *Coluber constrictor*, 226, 227
 Zamenis, 227
stellatus, *Kalophrynus*, 63
Stenocercus crassicaudatus, 149
 ervingi, 149
Stenodactylus variegatus, 149
stenophryys, *Lachesis*, 187
 Lachesis muta, 187
Stenorhina degenhardtii apiata, 214
 freminvillii lactra, 214
 lactea, 214
Stenostoma phenops, 214
stephensi, *Crotalus confluentus*, 169
 Crotalus mitchellii, 169
Sternotherus minor minor, 230
 odoratus, 233
stertens, *Rana capito*, 73
stickeli, *Kaloula conjuncta*, 63
 Lygosoma (Sphenomorphus) variegatum, 125
 Sphenomorphus variegatus, 125
sticticeps, *Pseudotriton ruber*, 26
 Spelerpes ruber, 26
stictogenys, *Diadophis punctatus*, 174
stictogrammus, *Cnemidophorus sackii*, 101
Stilosoma extenuatum arenicola, 214
Storeria dekayi temporalineata, 214
 dekayi texana, 214
 dekayi tropica, 216
 dekayi wrightorum, 215
 storerioides, 222
 tropica, 216
storerioides, *Storeria*, 222
 Tropidoclonium, 222
stramineus, *Chilomeniscus*, 163
 Chilomeniscus stramineus, 163
 fasciatus, *Chilomeniscus*, 163
stratissima, *Leptognathus*, 192
streckeri illinoensis, *Pseudacris*, 71
Streptosaurus mearnsi, 153
striatula, *Bolitoglossa*, 19
 Leptotyphlops, 195
striatulus, *Oedipus*, 19

- striatus axanthus*, *Pseudobranchus*, 22
 spheniscus, *Pseudobranchus*, 22
strigilata flavoguttata, *Hyla*, 54
stuarti, *Ameiva undulata*, 83
 Dryadophis melanolomus, 175
 Lygosoma cherriei, 125
 Scincella cherriei, 125
subcaudalis, *Holbrookia lacerata*, 117
subcinctus sealei, *Lycodon*, 196
subcristatus, *Triton*, 27, 28
subfasciatus, *Liophis*, 196
subflavus, *Epicrates*, 178
subgriseum, *Oligodon*, 202
subniger, *Sceloporus aeneus*, 134
suboculare, *Bascanion*, 160
subocularis, *Anolis*, 91
subrubrum hippocrepis, *Kinosternon*, 232
subsalsum, *Ambystoma*, 7
sugillatus, *Sceloporus jarrovii*, 136
sulcatus, *Typhlops*, 225
sumatranus, *Ichthyophis*, 4
sumichrasti cerebrosus, *Thamnophis*, 219
 Epaphelus, 110
 Eumeces, 132
 Eutaenia, 181
 Gymnophthalmus, 110
 Hylella, 61
 Plistodon, 132
 Thamnophis sumichrasti, 181, 218
sumichrastii, *Conophis*, 167
 viduus, *Conophis*, 167
surda, *Cochranella*, 39
 Hyla aurantiaca, 51
surinamensis tobaganus, *Amiva*, 84
suspectum, *Heloderma*, 116
suspectus, *Elapomorphus*, 177
sylvatica cantabrigensis, *Rana*, 72
 cherokiana, *Rana*, 76
 Phyllobates, 70
symmetrica, *Uta*, 155
symmetricus, *Urosaurus ornatus*, 155
Symphimus leucostomus, 216
Sympholis lippiens, 216
Syrhophus campi, 79
 cystignathoides, 69
 guttillatus, 66
 interorbitalis, 79
 latodactylus, 79
 leprus, 79
 modestus, 79
 nebulosus, 80
 pipilans, 80
 rubrimaculatus, 80
 smithi, 80
syspila, *Lampropeltis triangulum*, 203
syspilus, *Ophibolus doliatus*, 203, 204
tacanensis, *Bufo*, 37
tachydromoides oldi, *Takydromus*, 150
Tachymenis canilatus, 216
taeniata graminea, *Abronia*, 114
 Leptophis, 193
taeniatum, *Arrhyton*, 160, 164
taeniatus australis, *Masticophis*, 197
 Masticophis taeniatus, 193
 ornatus, *Masticophis*, 197
 schotti, *Masticophis*, 197
taeniocnemis, *Sceloporus*, 143
 Sceloporus malachiticus, 143
Taeniophis imperialis, 216
 tantillus, 216
taitensis, *Chamaeleo*, 96
Takydromus dorsalis, 149
 intermedius, 150
 khasiensis, 150
 tachydromoides oldi, 150
talamancae, *Eleutherodactylus*, 47
Tantilla bimaculata, 216
 calamarina, 216
 canula, 216
 coronata, 217
 coronata coronata, 217
 eiseni, 217
 eiseni eiseni, 217
 flavilineata, 217
 kirnia, 217
 lintoni, 217
 miniata, 217
 miniator, 217
 nigriceps fumiceps, 212, 217
 phrenitica, 217
 rubra, 218
 vermiformis, 195
 wilcoxi, 218
 wilcoxi rubricata, 218
 wilcoxi wilcoxi, 218
Tantillita lintoni, 217
tantillus, *Taeniophis*, 216
Tapaya brevirostris, 150
 hernandesi, 150
 ornatissima, 150
Tarentola mauritanica, 131
Taricha laevis, 26
 torosa torosa, 26
tau, *Trimorphodon*, 222
taylori, *Anolis*, 91
 Ficimia publia, 183
 Leptodeira annulata, 191
 Uta, 155



- tehuanæ, *Dryadophis melanolomus*, 175
 Lepidophyma flavimaculatum, 123
 Lepidophyma smithii, 123
Telmatobius niger, 80
 temporalineata, *Storeria dekayi*, 214
 temporalis, *Lampropeltis triangulum*, 188
 tenebrosum, *Amblystoma*, 6
 tenere, *Elaps*, 177
 Micrurus fulvius, 177
 tentaculata, *Cæcilia*, 3
 tenuis, *Calamaria*, 162, 168
 Chelonia, 228
 Contia, 162, 168
 tenuissimus, *Himantodes*, 185
 Imantodes, 185
 tephroleura, *Tropidonotus*, 223
 tequesta, *Malaclemmys terrapin*, 233
 teres brachylopha, *Ctenosaura*, 106
Terrapene bauri, 235
 carolina bauri, 235
 carolina triunguis, 230
 goldmani, 235
 mexicana mexicana, 235
 nelsoni, 235
 ornata ornata, 229
terrapin littoralis, *Malaclemmys*, 232
 macrospilota, *Malaclemmys*, 233
 tequesta, *Malaclemmys*, 233
 terrestris, *Bolitoglossa*, 11
 Bufo terrestris, 35
 charlesmithi, *Bufo*, 37
 Chiropterotriton, 11
 copei, *Bufo*, 32
 tessellata, *Ninia*, 202
 tessellatus aethiops, *Cnemidophorus*, 102
 Cnemidophorus, 98
 multiscutatus, *Cnemidophorus*, 102
 Ninia sebae, 202
 rubidus, *Cnemidophorus*, 102
 Testudines, 227
Testudo kleinmanni, 235
 mauritanica, 235
 tetragrammus, *Eumeces*, 132
 Plestiodon, 132
Tetraprion jordani, 80
 tetrataenia, *Eutaenia sirtalis*, 181
 Thamnophis sirtalis, 181
 texana, *Holbrookia texana*, 116
 Hypsiglena, 186
 Hypsiglena torquata, 186
 Pseudemys, 234
 Pseudemys floridana, 234
 Siren intermedia, 25
 Storeria dekayi, 214
 texense, *Engystoma*, 48
 texensis, *Diadophis*, 174
 thalassina, *Uta*, 155
 thalassinus, *Petrosaurus*, 155
Thalassochelys corticata, 235
Thalerophis depressirostris, 193
Thamnophis angustirostris, 178
 chrysocephalus, 178
 cyrtopsis cyclides, 218
 cyrtopsis cyrtopsis, 179, 218
 elegans atratus, 180
 elegans biscutatus, 179
 elegans couchii, 178
 elegans elegans, 181, 183
 elegans hammondii, 179
 elegans nigrescens, 218
 elegans vagrans, 179, 180, 181, 183
 eques eques, 179, 180
 eques megalops, 180
 errans, 218
 marcianus, 182
 melanogaster canescens, 218
 ordinoides, 182
 ordinoides errans, 218
 phenax phenax, 182
 radix haydenii, 180, 182
 radix radix, 182
 rozellæ, 218
 rufipunctatus, 164
 sauritus chalceus, 182
 sauritus proximus, 181
 sirtalis pickeringi, 182
 sirtalis sirtalis, 181, 183
 sirtalis tetrataenia, 181
 sirtalis trilineata, 183
 sumichrasti cerebrosus, 219
 sumichrasti sumichrasti, 181, 218
 vicinus, 219
thayeri, *Sceloporus*, 143
thermophila, *Holbrookia*, 118
 Holbrookia maculata, 118
thomasi, *Ameiva undulata*, 84
 trinidadensis, *Eleutherodactylus*, 47
Thorius dubitus, 27
 maxillabrochus, 27
 narisovalis, 27
 pennatulus, 27
 pulmonaris, 27
 schmidti, 27
Thoropa lutzi, 80
Thrasops praestans, 219
Tiaporus fuliginosus, 150
tigrinum, *Ambystoma tigrinum*, 5, 6, 24
 diaboli, *Ambystoma*, 7

- tigrinum—Continued
 mavortium, *Ambystoma*, 5, 6, 7, 25
 melanostictum, *Ambystoma*, 8
 nebulosum, *Ambystoma*, 7
 slateri, *Ambystoma*, 8
 stebbinsi, *Ambystoma*, 8
- tigris aethiops, *Cnemidophorus*, 97, 98, 101, 102
 celeripes, *Cnemidophorus*, 97
Cnemidophorus, 102
Cnemidophorus tigris, 102
Crotalus, 171
 estebanensis, *Cnemidophorus*, 97
 gracilis, *Cnemidophorus*, 98, 100, 102
 marmoratus, *Cnemidophorus*, 100
 martyris, *Cnemidophorus*, 100
 multiscutatus, *Cnemidophorus*, 96, 102
 rubidus, *Cnemidophorus*, 102, 103
 variolosus, *Cnemidophorus*, 103
- Tiliqua dura*, 151
- tinkhami, *Trimeresurus flavoviridis*, 220
- tobagana, *Ameiva*, 84
Amiva surinamensis, 84
- Toluca amphisticta*, 219
 conica, 219
 frontalis, 219
 lineata, 219
 lineata acuta, 202
 lineata lineata, 219
 lineata varians, 164
 lineata wetmorei, 219
- Tomodactylus albolabris*, 80
 angustidigitorum, 81
 macrotympanum, 81
- Tomodon nasutus*, 220
- torosa, *Taricha torosa*, 26
- torquata baueri, *Hypsiglena*, 186
 deserticola, *Hypsiglena*, 186
 ochrorhyncha, *Hypsiglena*, 186
 texana, *Hypsiglena*, 186
- torquatus cyanogenus, *Sceloporus*, 143
 melanogaster, *Sceloporus*, 139
 minor, *Sceloporus*, 143
 mucronatus, *Sceloporus*, 144
Sceloporus torquatus, 135
- tortuganus, *Leimadophis*, 191
Leimadophis parvifrons, 191
- townsendi, *Amphisbaena*, 85
Anolis, 91
Oedipus, 19
Parvimolge, 19
Sauromalus, 133
Sauromalus obesus, 133
Sphaerodactylus, 148
- Toxicophis pugnax*, 220
- Trachycephalus scutigerus*, 81
 wrightii, 81
- transversa, *Natrix erythrogaster*, 201, 222, 223
- transversus, *Tropidonotus*, 223
- treleasei, *Chrysemys*, 229
- Tretanorhinus nigroluteus*, 220
 nigroluteus nigroluteus, 220
 variabilis binghami, 220
- trianguligaris venezuelensis, *Hydrops*, 186
- triangulum amauro, *Lampropeltis*, 187, 203
 annulata, *Lampropeltis*, 187
 celaenops, *Lampropeltis*, 189
 gentilis, *Lampropeltis*, 188, 204
Lampropeltis triangulum, 203
 nelsoni, *Lampropeltis*, 189
 schmidti, *Lampropeltis*, 189
 syspila, *Lampropeltis*, 203
 temporalis, *Lampropeltis*, 188
- triaspis, *Coluber*, 166
Elaphe triaspis, 166
 mutabilis, *Elaphe*, 165
 phaescens, *Elaphe*, 176
- trifrenatus, *Leptophis*, 193
- trigemina, *Clelia*, 205
- trigeminus, *Oxyrhopus*, 205
- trilineata, *Eutainia sirtalis*, 183
Thamnophis sirtalis, 183
- Trimeresurus flavoviridis tinkhami*, 220
 garciai, 220
 nummifer picadoi, 220
 stejnegeri, 220
 yucatanicus, 221
- Trimerorhinus tritaeniatus multisquamis*, 221
- Trimorphodon biscutatus biscutatus*, 222
 biscutatus quadruplex, 221
 biscutatus semirutus, 221
 collaris, 221
 fasciolata, 221
 forbesi, 221
 lambda, 221
 lyrophanes, 196
 major, 222
 tau, 222
 upsilon, 222
 wilkinsonii, 222
- trinidadensis, *Eleutherodactylus thomasi*, 47
- trinitatis extremus, *Anolis*, 91
 luciae, *Anolis*, 89
Scolecosaurus, 145
 vincentii, *Anolis*, 92

- Trionyx spinifer asper*, 227, 233
 spinifer emoryi, 227
 spinifer hartwegi, 227
Tripriion petasatus, 68
triseriatus miquihuanus, *Crotalus*, 172
trisruptum, *Amblystoma*, 6
tristichus, *Sceloporus*, 144
 Sceloporus undulatus, 144
tristis, *Elaps*, 177
tristycha, *Ozotheca*, 233
tritaeniatus multisquamis, *Trimerorhinus*,
 221
Triton blasii, 27
 ensicauda, 27
 subcristatus, 27, 28
Triturus ensicaudus, 27
 ensicaudus ensicaudus, 28
 ensicaudus popei, 28
 marmoratus marmoratus, 27
 perstriatus, 28
triunguis, *Cistudo*, 230
 Terrapene carolina, 230
trivirgata, *Lichanura*, 195
trivittatus, *Leptodactylus*, 65
tropica, *Storeria* 216,
 Storeria dekayi, 216
Tropidoclonium storerioides, 222
Tropidodipsas brevifacies, 222
 lunulata, 222
Tropidonotus bisectus, 222
 celaeno, 222
 compressicaudus walkeri, 223
 rhombifer, 223
 tephropleura, 223
 transversus, 223
 ustus, 223
tropidonotus spilorhipis, *Anolis*, 91
Tropidophis maculatus haetianus, 225
 maculatus jamaicensis, 223, 224
 pardalis androsi, 223
 pardalis barbouri, 224
 pardalis bucculentus, 225
 pardalis canus, 225
 pardalis stejnegeri, 223, 224
 wrighti, 224
Tropidophorus misaminius, 151
 partelloi, 151
Tropidurus abingdonensis, 151
 albemarlensis, 151
 albemarlensis albemarlensis, 151
 bivittatus, 152
 delanonis, 151
 duncanensis, 151
 hoodensis, 151
Tropidurus—Continued
 indefatigabilis, 151
 lemniscatus, 152
 pacificus, 151
 spinulosus, 126
trossulus, *Anolis*, 92
 Batrachylodes, 31
truei, *Ascaphus*, 29
 Ascaphus truei, 29
 californicus, *Ascaphus*, 29
 montanus, *Ascaphus*, 29
tsushimensis, *Rana*, 77
tunganus, *Eumeces*, 113
turgida, *Leptognathus*, 192
turgidus, *Sibynomorphus*, 192
turquinensis, *Eleutherodactylus*, 47
turuensis, *Agama agama*, 81
tuxtlae, *Lepidophyma*, 123
twiningi, *Eutaenia radix*, 130
tynerensis, *Eurycea*, 15
Typhlomolge rathbuni, 28
Typhlops basimaculatus, 224
 mandensis, 224
 manni, 224
 microstomus, 224
 praecocularis, 224
 rostellatus, 224
 stejnegeri, 225
 sulcatus, 225
Typhlotriton nereus, 28
 spelaeus, 28
tzabcan, *Crotalus durissus*, 170
uluguruensis, *Natrix olivacea*, 200
 Scolecormorphus, 4
Uma notata, 152
 notata notata, 152
 notata scoparia, 152
 rufopunctata, 152
 scoparia, 152
unctus, *Diplodactylus*, 103
 Phyllodactylus, 103
undosa, *Hombromia*, 118
undosus, *Sphenomorphus*, 118
undulata amphigramma, *Ameiva*, 83
 dextra, *Ameiva*, 83
 hartwegi, *Ameiva*, 83
 Microhyla, 68
 podarga, *Ameiva*, 83
 sinistra, *Ameiva*, 83
 stuarti, *Ameiva*, 83
 thomasi, *Ameiva*, 84
undulatus, *Cnemidophorus*, 102
 elongatus, *Sceloporus*, 135
 erythrocheilus, *Sceloporus*, 144

- undulatus—Continued
 hyacinthinus, *Sceloporus*, 143
 Sceloporus undulatus, 135
 tristichus, *Sceloporus*, 144
Ungalia cana, 225
 bucculenta, 225
Ungaliophis panamensis, 225
Ungualia haetiana, 225
unguidentis, *Bolitoglossa*, 11
 Pseudoeurycea, 11
uniformis, *Anolis*, 92
 Anolis humilis, 90, 92
 Sceloporus magister, 138
unica, *Uta*, 155
unicanthalis, *Sceloporus scalaris*, 142
unicolor, *Cnemidophorus*, 103
 Drymarchon corais, 176
 Eleutherodactylus, 47
unicus, *Urosaurus*, 155
upsilon, *Trimorphodon*, 222
uranochroa, *Hyla*, 60
Uromacer scandax, 225
 wetmorei, 225
Urosaurus auriculatus, 152
 bicarinatus anonymorphus, 152
 bicarinatus nelsoni, 154
 bicarinatus spinosus, 152
 clarionensis, 153
 nigricaudus, 154
 ornatus caeruleus, 153
 ornatus linearis, 151
 ornatus ornatus, 154
 ornatus schmidti, 154
 ornatus symmetricus, 155
 unicus, 155
Urotheca williamsi, 225
usta, *Microhyla usta*, 48
ustum, *Engystoma*, 48
ustus, *Tropidonotus*, 223
Uta anonymorpha, 152
 auriculata, 152
 caerulea, 153
 clarionensis, 153
 concinna, 153
 elegans, 153
 levis, 153
 mannophorus, 153
 mearnsi, 153
 nelsoni, 154
 nigricauda, 154
 ornata, 154
 ornata linearis, 154
 ornata schmidti, 154
 palmeri, 155
Uta—Continued
 parva, 155
 squamata, 155
 stansburiana, 155
 stansburiana elegans, 153
 stansburiana hesperis, 155
 stansburiana stansburiana, 155
 symmetrica, 155
 taylori, 155
 thalassina, 155
 unica, 155
utiformis, *Sceloporus*, 144
uvidus, *Manculus quadridigitatus*, 16
vagrans, *Eutainia*, 183
 plutonia, *Eutaenia*, 181
 Thamnophis elegans, 179, 180, 181, 183
valeriae, *Haldea valeria*, 226
 Virginia, 226
valida isabelleae, *Natrix*, 200
 Natrix, 222
 Natrix valida, 209, 223
 Regina, 209
valliceps, *Bufo*, 33, 36
vandenburghi, *Cnemidophorus*, 103
vandenburghianus, *Sceloporus*, 144
 Sceloporus graciosus, 144
vandykei idahoensis, *Plethodon*, 20
 larselli, *Plethodon*, 22
vanvlietii, *Hyla*, 60
variabilis, *Hyla*, 60
 binghami, *Tretanorhinus*, 220
 marmoratus, *Sceloporus*, 134
 ollopopus, *Sceloporus*, 144
 smithi, *Sceloporus*, 144
varians, *Toluca lineata*, 164
variegatus abbotti, *Coleonyx*, 103
 bogerti, *Coleonyx*, 103
 brevis, *Coleonyx*, 103
 Coleonyx variegatus, 149
 peninsularis, *Coleonyx*, 103
 sonoriensis, *Coleonyx*, 103
 Stenodactylus, 149
 stickeli, *Lygosoma* (*Sphenomorphus*),
 125
 stickeli, *Sphenomorphus*, 125
variolosum, *Engystoma*, 48
variolosus, *Cnemidophorus*, 103
 Cnemidophorus tigris, 103
 Hypopachus, 48
 Masticophis flagellum, 197
varius certus, *Atelopus*, 30
 glyphus, *Atelopus*, 30
 Leiocephalus, 124
 Liocephalus, 124

- varius—Continued
 Sauromalus, 133
 Scaphiopus, 78
velatus, *Bufo woodhousii*, 38
venezuelensis, *Bothrops lansbergii*, 161
 Hydrops trianguligaris, 186
ventralis, *Calamaria*, 162
 Calamaria lovii, 162
 Callisaurus draconoides, 118
 compressus, *Ophisaurus*, 127
 Homalosaurus, 118
 myurus, *Callisaurus*, 95
ventrilineatus, *Eleutherodactylus*, 44, 63
 Leptodactylus, 63
ventromaculatum, *Plethodon welleri*, 22
venustus, *Euprepis*, 113
veraecrucis, *Bothrops nummifer*, 161
veraepacis nigrilatus, *Adelphicos*, 157
vermiformis, *Lioninia*, 195
 Tantilla, 195
vernalis blanchardi, *Opheodrys*, 202
versa, *Graptemys*, 231
 Graptemys pseudogeographica, 231
versicolor, *Hyla versicolor*, 60
 phaecrypta, *Hyla*, 60
 sandersi, *Hyla*, 60
vertebralis, *Pituophis melanoleucus*, 208
Verticaria heldingi, 156
 caerulea, 156
verus, *Cnemidophorus gularis gularis*, 98
vetustus, *Bascanion*, 160
vicinus, *Thamnophis*, 219
vidua, *Eutaenia infernalis*, 180, 181
viduus, *Conophis sumichrasti*, 167
vigilis, *Xantusia*, 156
vilkinsonii, *Trimorphodon*, 222
vincentii, *Anolis*, 92
 Anolis trinitatis, 92
vinculum, *Leiocephalus*, 121
 Leiocephalus personatus, 121
vinitor, *Dendrophidion*, 172
violacea, *Cantoria*, 162
vioscai, *Pseudotriton ruber*, 23
vioscana, *Pseudemys*, 235
virescens brachycephala, *Rana*, 77
 Leiocephalus, 121
virgatus, *Anolis*, 92
Virginia valeriae, 226
virginiana, *Lampropeltis elapsoides*, 188
viridescens dorsalis, *Diemictylus*, 13
 evergladensis, *Diemictylus*, 13
 perstriatus, *Diemictylus*, 28
 piaropicola, *Diemictylus*, 13
 vittatus, *Diemictylus*, 13
 viridis abyssus, *Crotalus*, 169
 caliginis, *Crotalus*, 172
 Crotalus viridis, 171
 lutosus, *Crotalus*, 169
 nuntius, *Crotalus*, 172
 oreganus, *Crotalus*, 171
 Polypedates, 71
 Rhacophorus, 71
viridissima, *Centrolenella*, 38
visayanus, *Rana macrodon*, 74
 Rana magna, 74
visoninus, *Adelphicos quadrivirgatus*, 211
 Rhegnops, 211
vitiensis, *Cornufer*, 49
 Halophila, 49
vittata, *Rhadinaea*, 210
vittatum, *Arrhyton*, 159
vittatus, *Basiliscus*, 94, 107
 Conophis vittatus, 167
 Diemictylus viridescens, 13
viviparus, *Sceloporus*, 145
vocalis, *Eleutherodactylus*, 47
vogli, *Atelopos cruciger*, 30
vomerina, *Elosia*, 48
vorax, *Gehyra*, 114
vudii raineyi, *Alsophis*, 158
vulcani, *Eleutherodactylus*, 47
vulpecula, *Doryura*, 108
vulpina, *Elaphe vulpina*, 213
vulpinus, *Scotophis*, 213
walkeri, *Liolaemus*, 124
 Tropidonotus compressicaudus, 223
walli, *Dinodon rufozonatum*, 174
webbii, *Elgaria multicarinata*, 115
 Gerrhonotus, 115
wehrlei dixi, *Plethodon*, 19
 jacksoni, *Plethodon*, 20
weinandii, *Celestus*, 96
welleri, *Plethodon*, 22
 Plethodon welleri, 22
 ventromaculatum, *Plethodon*, 22
welteri, *Desmognatus fuscus*, 12
Wenona isabella, 226
 plumbea, 226
werleri, *Natrix rhombifera*, 200
 Pseudoeurycea, 23
wetmorei, *Ameiva*, 84
 Ameiva wetmorei, 84
 Anolis distichus, 87
 Anolis dominicensis, 87
 Chamaelinorops, 96
 eleanorae, *Ameiva*, 84
 Eleutherodactylus, 48

- wetmorei—Continued
 Toluca lineata, 219
 Uromacer, 225
- Wetmorena haetiana, 156
- wilcoxi rubricata, Tantilla, 218
 Tantilla, 218
 Tantilla wilcoxi, 218
- wileyi, Anolis cristatellus, 86
- wilkesii, Pituophis, 207
- willardi meridionalis, Crotalus, 172
 silus, Crotalus, 172
- williamsi, Urotheca, 225
- wirshingi, Phyllodactylus, 131
- wislizenii, Crotaphytus, 105
 Gambelia wislizenii, 104, 105
 silus, Gambelia, 104
- woodhousei, Bufo woodhousei, 31, 33
 hemiophrys, Bufo, 34
- woodhousii, Bufo, 37
 Bufo woodhousii, 38
 Nerodia, 201
 velatus, Bufo, 38
- woodi, Ameiva chrysolema, 82
 Philautus, 69
 Sceloporus, 145
- woodini, Lampropeltis pyromelana, 188
- woodworthi, Rana, 77
- worcesteri, Cornufer, 39
- wrighti, Batrachoseps, 22
 Desmognathus, 13
 Plethopsis, 22
 Trepidophis, 224
- wrightii, Hyla, 81
 Trachycephalus, 81
- wrightorum, Storeria dekayi, 215
 Hyla, 60
 Hyla eximia, 60
- xanthoptica, Ensatina eschscholtzii, 13
- Xantusia arizonae, 156
 henshawi, 156
 riversiana, 156
 riversiana reticulata, 156
 vigilis, 156
- Xenodon ancorus, 226
 mexicanus, 226
- Xerobates agassizii, 235
 berlandieri, 236
- xiphias, Amblystoma, 6
- Xiphosoma annulatum, 226
 madagascariense, 226
- Xiphosurus oculus, 156
- xolocalcae, Bolitoglossa, 11
 Chiropterotriton, 11
- yakani, Rana, 77
- zenyuanensis, Batrachuperus, 8
- yonahlossee, Plethodon, 22
- yucatanensis, Leptophis mexicanus, 193
 Leptodeira yucatanensis, 213
 malleisi, Leptodeira, 191
 Sibon annulatum, 213
- yucatanicus, Bothrops, 221
 Trimeresurus, 221
- yumensis, Lampropeltis getulus, 188
- Zachaenus parvulus, 38
 roseus, 81
 sawayae, 81
- Zamenis conirostris, 226
 lateralis fuliginosus, 227
 stejnegerianus, 227
- Zaocys oshimai, 227
- zempoalaensis, Rhyacosiredon, 24
- zeteki, Hyla, 61
- zonata multicolor, Lampropeltis, 204
 parvirubra, Lampropeltis, 189
 pulchra, Lampropeltis, 189
- zosteromus, Sceloporus, 145
 Sceloporus magister, 145

